

National Park Service


U.S. Department of the Interior

Natural Resource Program Center



Classification of the Plant Communities of Craters of the Moon National Monument and Preserve, Idaho

Natural Resources Report NPS/UCBN/NRTR – 2008/096



ON THE COVER

Images of Craters of the Moon National Monument and Preserve (clockwise): view north looking over Craters of the Moon lava field, open limber pine woodlands on cinders cones; cushion buckwheat; moss on lava rock; Indian paintbrush; pahoehoe lava.
Photograph by: Northwest Management, Inc.

Classification of the Plant Communities of
Craters of the Moon National Monument and Preserve,
Idaho

Natural Resource Technical Report NPS/UCBN/NRTR – 2008/096

Steven K. Rust
Idaho Conservation Data Center
Department of Fish and Game
Boise, Idaho

Paige Wolken
Craters of the Moon National Monument and Preserve
National Park Service
Arco, Idaho

January 2008

U.S. Department of the Interior
National Park Service
Inventory and Monitoring Program
Upper Columbia Basin Network
Moscow, Idaho

The Natural Resource Publication series addresses natural resource topics that are of interest and applicability to a broad readership in the National Park Service and to others in the management of natural resources, including the scientific community, the public, and the NPS conservation and environmental constituencies. Manuscripts are peer-reviewed to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and is designed and published in a professional manner.

Natural Resource Reports are the designated medium for disseminating high priority, current natural resource management information with managerial application. The series targets a general, diverse audience, and may contain NPS policy considerations or address sensitive issues of management applicability. Examples of the diverse array of reports published in this series include vital signs monitoring plans; "how to" resource management papers; proceedings of resource management workshops or conferences; annual reports of resource programs or divisions of the Natural Resource Program Center; resource action plans; fact sheets; and regularly-published newsletters.

The Natural Resource Technical Reports series is used to disseminate the peer-reviewed results of scientific studies in the physical, biological, and social sciences for both the advancement of science and the achievement of the National Park Service's mission. The reports provide contributors with a forum for displaying comprehensive data that are often deleted from journals because of page limitations. Current examples of such reports include the results of research that addresses natural resource management issues; natural resource inventory and monitoring activities; resource assessment reports; scientific literature reviews; and peer reviewed proceedings of technical workshops, conferences, or symposia.

Views and conclusions in this report are those of the authors and do not necessarily reflect policies of the National Park Service. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the National Park Service.

Printed copies of reports in these series may be produced in a limited quantity and they are only available as long as the supply lasts. This report is also available from the Natural Resource Publications Management Web site (<http://www.nature.nps.gov/publications/NRPM>) on the Internet, or by sending a request to the address on the back cover.

Please cite this publication as:

Rust, S. K. and P. Wolken. 2008. Classification of the Plant Communities of Craters of the Moon National Monument and Preserve, Idaho. Natural Resource Technical Report, NPS/UCBN/NRTR – 2008/096. National Park Service, Fort Collins, Colorado.

NPS D-94, January 2008

Table of Contents

List of Figures.	iii
List of Tables.	iii
Abstract.	iv
Introduction.	1
Study Area.	1
Methods.	3
Results.	4
The Classification of Plant Communities.	4
Dichotomous Key to the Classification.	12
Literature Cited.	18

List of Figures

Figure 1. Study area location..	1
---	---

List of Tables

Table 1. Plant Community Classification.	6
--	---

Appendices

Appendix 1. Cross-walk to vegetation types..	24
Appendix 2. Summary of pertinent classification literature..	27
Appendix 3. Stand tables..	29

Abstract

Work on the vegetation map for Craters of the Moon National Monument and Preserve was initiated in May 2006. Craters of the Moon National Monument and Preserve lies on the Upper Snake River Plain in south central Idaho. The area encompasses sparse, vegetated lava and cinder cones; sagebrush steppe; grasslands; wetlands; and montane shrublands, woodlands, and forest. Three data sets were selected to augment field data collected specifically for the mapping project. Together with data collected in 2006, 1,241 vegetation samples were identified for plant community classification analyses. Initial plant community groups were identified using hierarchical cluster analysis. Plant community groups were successively partitioned to optimize within-group homogeneity, between-group difference, and the number of groups using indicator species analysis and multi-response permutation procedures. Decisions regarding the identification of plant associations within the data were informed by cross-walking candidate community groupings to the National Vegetation Classification. Classification and regression tree analysis (recursive partitioning) was employed in the development of dichotomous keys and the evaluation of cluster analysis results. Ninety-three plant communities are identified as occurring within the study area. One plant community is classified at the alliance level; the remainder are classified at the association level. Nearly 50% of the associations identified within the Monument and Preserve are currently listed in the National Vegetation Classification. Minor name-changes are proposed. The classification largely corresponds to vegetation types identified by Day and Wright (1985). Associations described but not recognized by the National Vegetation Classification include: (1) semi-natural vegetation (18 associations) and (2) plant communities (21 associations) that have not been described previously at the association level. Plant associations that should be considered for inclusion in the National Vegetation Classification are listed. A dichotomous key to the classification is presented. Summaries of stand composition are provided.

Introduction

The National Park Service Inventory and Monitoring Program Upper Columbia Basin Network (UCBN) is working in conjunction with the U.S. Geological Survey and the National Park Service Vegetation Mapping Program to complete vegetation maps for each of the national park units within the network. The National Park Service vegetation mapping protocol consists of the following major phases: preliminary collection and review of existing information; field data collection, management and analysis; photo-interpretation and mapping; map validation; and accuracy assessment (The Nature Conservancy and Environmental Systems Research Institute 1994a). Work on the vegetation map for Craters of the Moon National Monument and Preserve was initiated in May 2006. The initial phases of the project are reported by Northwest Management Inc. (2006). The objective of this document is to report on the field data analysis phase and provide a classification of the vegetation and a dichotomous key to the classification of plant communities.

Study Area

Craters of the Moon National Monument and Preserve lies on the Upper Snake River Plain in south central Idaho (Figure 1). The National Monument and Preserve encompasses vast areas of recent, relatively unweathered, flow basalt and cinder cones forming the Great Rift, Craters of the Moon Lava Field, and Wapi Lava Field (Kuntz et al. 1982 and 1988). Soils adjacent to the younger lava flows are predominately moderately deep to very deep eolian sand and loess forming loam, silt loam, sandy loam, and sandy soils (Johnson 1991, Jurs and Sands 2004). Elevation ranges from 1,615 m to 2,350 m (5,300 ft to 7,700 ft).

The climate is arid. Summers are hot and dry; winters are cold. In the northern portion of the Monument and Preserve average annual minimum and maximum temperatures are -1.0°C (30.2°F) and 13.0°C (55.4°F), respective; mean annual precipitation is 39.5cm (15.5in). In the south, hotter and drier conditions prevail. On the southern end of the Wapi Flow average minimum and maximum temperatures are (approximately) -1.0°C (30.2°F) and 15.4°C (59.7°F), respective; mean annual precipitation is 24.1cm (9.5in) (Western Regional Climate Center 2007).

Vegetation within the Monument and Preserve is diverse. The area encompasses sparse, vegetated lava and cinder cones; sagebrush steppe; grasslands; wetlands; and montane shrublands, woodlands, and forest. The study area encompasses several hundred kipukas (isolated areas of vegetation surrounded by more recent lava flows) and numerous parks. Many of these areas of relatively pristine native vegetation are protected from disturbances of exotic species invasion, livestock grazing, and recreational use. The Monument and Preserve commands a history of studies describing and conserving high quality, reference vegetation conditions (Tisdale et al. 1965; Caicco and Wellner 1983a, 1983b, 1983c). Day and Wright (1985) compiled an early vegetation map and describe plant communities of a portion of the study area. Other studies regarding the natural vegetation of the study area include Caicco

(1987), Day (1985), Egger (1941), Hironaka et al. (1983), Jurs and Sands (2004), Moseley (1989), Rust et al. (2001), Rust (1999) and USDI National Park Service and Bureau of Land Management (2005). USDI National Park Service and Bureau of Land Management (2005) provide a history of fire disturbance and describe livestock grazing practices within the area.




Figure 1. Study area location. The location of Craters of the Moon National Monument and Preserve (CRMO) is shown in relation to major geographical features. Inset shows the location of the study area within Idaho.

Methods

At the onset of the vegetation mapping project existing data were evaluated for potential use in the project (Northwest Management Inc. 2006, The Nature Conservancy 1996). Three data sets were selected to augment field data collected specifically for the mapping project: (1) Jurs and Sands (2004) data from Laidlaw Park, Little Park, and Paddelford Flat; (2) Rust (1999) data from Bureau of Land Management areas of critical environmental concern (Sand Kipuka and Big Juniper Kipuka); and (3) USDI Bureau of Land Management (2001) fuels inventory data collected within the Monument and Preserve. All four data sets, including the data collected in 2006, contributed 1,241 vegetation samples that were utilized for plant community classification analyses. Field methods employed in the collection of these data are described by Jurs and Sands (2004), Rust (1999), USDI Bureau of Land Management (2001), and Northwest Management Inc. (2006), respectively.

Analytical methods were employed in a manner consistent with the objective of identifying natural associations of existing vegetation (Jennings et al. 2006, Tart et al. 2005, The Nature Conservancy and Environmental Systems Research Institute 1994b). Multivariate statistical analyses were conducted using PC-ORD (McCune and Mefford 1999). Initial plant community groups were identified using hierarchical cluster analysis. Plant community groups were successively partitioned to optimize within-group homogeneity, between-group difference, and the number of groups. These 3 factors were evaluated using statistics generated by indicator species analysis (the mean p -value and number of significant ($p \leq 0.05$) indicator species) and multi-response permutation procedures (T , a measure of separation between groups; the chance-corrected within-group agreement (A), a measure of within-group homogeneity; and the compositional similarity of samples within a group (as measured by average relative Sorensen distance)). Environmental relations, within-group homogeneity, and between-group difference were evaluated graphically using indirect gradient analysis through detrended correspondence analysis (McCune and Grace 2002). Final decisions regarding the naming of plant associations were informed by cross-walking candidate community groupings to the National Vegetation Classification (NatureServe 2007). Classification and regression tree analysis (recursive partitioning) was employed in the development of dichotomous keys and the evaluation of classification results using the `rpart` package in R ® Development Core Team 2007).

The plant species nomenclature standard identified by the Monument and Preserve was maintained for the analysis to reflect local and regional taxonomic distinctions. Plant nomenclature of the Integrated Taxonomic Information System (ITIS) as reflected by the PLANTS Database (USDA, NRCS 2007) is employed in the naming of plant associations. This is the National Vegetation Classification standard and is necessary to discriminate differences among associations currently reported within the National Vegetation Classification.

Results

The Classification of Plant Communities

Ninety-three plant communities are identified as occurring within the study area. One plant community is classified at the alliance level; the remainder are classified at the association level. Table 1 provides a summary of the associations described for the Monument and Preserve.

Plant communities are named for the indicator (dominant or diagnostic) species of the vegetative strata present. The indicator species of the uppermost strata is listed first, followed by successively lower strata. Species that occur in the same strata (or are the same lifeform) are separated by a hyphen (-). Indicator species that occur in different strata (or are a different lifeform) are separated by a slash (/). Alliance names are concluded with the word “Alliance” to differentiate them from association names. Plant association names incorporate the physiognomic class in which the association is classified (e.g., Forest, Woodland, or Herbaceous) (FGDC 1997, 2006).

Forty-seven (nearly 50%) of the associations identified within the Monument and Preserve are currently listed in the National Vegetation Classification (as indicated by an assigned element code, Table 1) (NatureServe 2007). Physiognomic classes employed in names of plant associations currently listed in the National Vegetation Classification are used here, though the physiognomic class of the occurrence of any given association within the Monument and Preserve may not actually correspond to the assigned class.

More specific differences between the National Vegetation Classification and this classification of field observations from the Monument and Preserve are: (1) *Eriogonum ovalifolium* var. *depressum* Dwarf-Shrubland (as named within the National Vegetation Classification) occurs with <10% total vegetative cover. The association is referred to as *Eriogonum ovalifolium* var. *depressum* Sparse Vegetation here. (2) Day and Wright (1985) described their “Antelope Bitterbrush/Great Basin Wildrye” vegetation type as having substantial abundance of *Pseudoregneria spicata*, though *Leymus cinereus* was more conspicuous. This vegetation type is described as *Purshia tridentata* / *Pseudoregneria spicata* - *Leymus cinereus* Shrub Herbaceous Vegetation in the National Vegetation Classification (NatureServe 2007). Samples taken from stands mapped as the vegetation type, however have little *Pseudoregneria spicata* and are described here as *Purshia tridentata* / *Leymus cinereus* Shrub Herbaceous Vegetation. Coincidentally, these two associations are considered endemic to Idaho and are only known from the Monument and Preserve.

The classification largely corresponds to vegetation types identified by Day and Wright (1985). A cross walk between their classification and the associations identified here is provided in Appendix 1. Primary differences in the classifications are: (1) floristic associations are identified for “Low Density Lava Flows”, “Medium Density Lava Flows”, and “Riparian”; and (2) the classification presented here provides a finer resolution of *Pseudoregneria spicata*, and *Populus tremuloides* associations. Day and Wright’s (1985) “Antelope Bitterbrush” vegetation

type appears to be represented primarily by the *Purshia tridentata* / *Pseudoroegneria spicata* Shrub Herbaceous Vegetation association.

Many of the plant associations identified in this classification have been described previously by others. Reid et al. (2002) provide range-wide descriptions of many of the associations. A summary of previous classification work regarding vegetation that occurs within the study area is provided in Appendix 2. Two plant associations are listed in Table 1, but are not in the dichotomous key. These distinct associations are represented from only one sample, dissimilar to other associations within the study area, and are described in the literature or known to occur in the region at other locations. Twenty samples (~2% of available data) were not classified. These data represent (1) unique vegetation sampled in only one stand and (2) one group of samples in which key indicator species are unknown. These samples tentatively represent communities that are not described in the literature, but there is not sufficient data to verify an association.

Over one-half of the associations described in this classification are not recognized by the National Vegetation Classification or have not been described previously. These associations include: (1) semi-natural vegetation (18 associations, Table 1) that, due to the priority of conserving natural vegetation, has not been systematically reviewed nor incorporated into the National Vegetation Classification and (2) plant communities (21 associations, listed below) that have not been described previously at the association level. Distinctive plant associations that should be considered for inclusion in the National Vegetation Classification include:

Artemisia tripartita ssp. *tripartita* / *Achnatherum thurberianum* Shrubland
Artemisia tripartita ssp. *tripartita* / *Koeleria macrantha* Shrubland
Artemisia tripartita ssp. *tripartita* / *Pascopyrum smithii* Shrubland
Artemisia tripartita ssp. *tripartita* / *Poa secunda* Shrubland
Artemisia tridentata ssp. *vaseyana* - *Chamaebatiaria millefolium* / *Penstemon deustus* Shrubland
Artemisia tridentata ssp. *vaseyana* - *Purshia tridentata* / *Poa secunda* Shrubland
Artemisia tridentata ssp. *wyomingensis* / *Festuca idahoensis* Shrubland
Chamaebatiaria millefolium / *Penstemon deustus* Sparse Vegetation
Ericameria nana - *Holodiscus dumosus* / *Penstemon deustus* Shrubland
Ericameria nana / *Poa secunda* / *Penstemon deustus* Sparse Vegetation
Juniperus (osteosperma, scopulorum) / *Artemisia tridentata* ssp. *wyomingensis* -
Chamaebatiaria millefolium Wooded Shrubland
Juniperus osteosperma / *Artemisia tridentata* ssp. *wyomingensis* / *Hesperostipa comata*
Shrubland
Juniperus (scopulorum, osteosperma) / *Ericameria nana* Wooded Herbaceous Vegetation
Juniperus scopulorum / *Poa secunda* / *Penstemon deustus* Woodland
Philadelphus lewisii / *Penstemon deustus* Sparse Vegetation
Pinus flexilis / *Chamaebatiaria millefolium* / *Poa secunda* Sparse Vegetation
Poa secunda / *Leptodactylon pungens* Sparse Vegetation
Prunus virginiana / *Leymus cinereus* Shrubland
Purshia tridentata - *Ericameria nana* / *Penstemon deustus* Shrubland
Purshia tridentata - *Philadelphus lewisii* / *Poa secunda* Shrubland

Table 1. Plant Community Classification. Plant communities observed at Craters of the Moon National Monument and Preserve are listed alphabetically by National Vegetation Classification class with scientific name, common name, field code, element code (abbreviated as "elcode") and the number of stands sampled (N). One alliance is described (*Populus balsamifera* ssp. *trichocarpa* Temporarily Flooded Woodland Alliance); the remaining communities are plant associations.

Scientific Name	Common Name	Code	Elcode	N
Forest and Woodland				
<i>Juniperus (osteosperma, scopulorum) / Artemisia tridentata</i> ssp. <i>wyomingensis</i> - <i>Chamaebatiaria millefolium</i> Woodland	Juniper (Utah, Rocky Mountain) / Wyoming Big Sagebrush - Fernbush Woodland	JUOS-JUSC2/ARTRW8/CHMI2		7
<i>Juniperus scopulorum / Poa secunda / Penstemon deustus</i> Woodland	Rocky Mountain Juniper / Sandberg Bluegrass / Scabland Penstemon Woodland	JUSC2/POSE/PEDE4		14
<i>Pinus flexilis / Purshia tridentata</i> Woodland	Limber Pine / Antelope Bitterbrush Woodland	PIFL2/PUTR2	CEGL000814	16
<i>Populus balsamifera</i> ssp. <i>trichocarpa</i> Temporarily Flooded Woodland Alliance	Black Cottonwood Temporarily Flooded Woodland Alliance	POBAT	A.635	2
<i>Populus tremuloides</i> - <i>Pinus flexilis</i> Forest	Quaking Aspen - Limber Pine Forest	POTR5-PIFL2	CEGL000540	1
<i>Populus tremuloides / Artemisia tridentata</i> Forest	Quaking Aspen / Big Sagebrush Forest	POTR5/ARTR	CEGL000572	2
<i>Populus tremuloides / Calamagrostis rubescens</i> Forest	Quaking Aspen / Pinegrass Forest	POTR5/CARU	CEGL000575	6
<i>Populus tremuloides / Prunus virginiana</i> Forest	Quaking Aspen / Chokecherry Forest	POTR5/PRVI	CEGL000596	4
<i>Populus tremuloides / Symphoricarpos oreophilus / Calamagrostis rubescens</i> Forest	Quaking Aspen / Mountain Snowberry / Pinegrass Forest	POTR5/SYOR2/CARU	CEGL000612	8
<i>Populus tremuloides</i> / Tall Forbs Forest	Quaking Aspen / Tall Forbs Forest	POTR5/TALL FORBS	CEGL000618	1
<i>Pseudotsuga menziesii / Carex geyeri</i> Forest	Douglas-fir / Geyer's Sedge Forest	PSME/CAGE2	CEGL000430	2
<i>Pseudotsuga menziesii / Symphoricarpos oreophilus</i> Forest	Douglas-fir / Mountain Snowberry Forest	PSME/SYOR2	CEGL000462	8
Shrubland				
<i>Alnus incana / Mesic Forbs</i> Shrubland	Gray Alder / Mesic Forbs Shrubland	ALIN2/Mesic Forbs	CEGL001147	4
<i>Artemisia arbuscula</i> ssp. <i>arbuscula / Bromus tectorum</i> Semi-natural Shrubland	Low Sagebrush / Cheatgrass Semi-natural Shrubland	ARARA/BRTE		3
<i>Artemisia arbuscula</i> ssp. <i>arbuscula / Hesperostipa comata</i> Shrubland	Low Sagebrush / Needle and Thread Shrubland	ARARA/HECO26		3
<i>Artemisia arbuscula</i> ssp. <i>arbuscula / Poa secunda</i> Shrub Herbaceous Vegetation	Low Sagebrush / Sandberg Bluegrass Shrub Herbaceous Vegetation	ARARA/POSE	CEGL001411	8

Scientific Name	Common Name	Code	Elcode	N
<i>Artemisia arbuscula</i> ssp. <i>arbuscula</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation	Low Sagebrush / Bluebunch Wheatgrass Shrub Herbaceous Vegetation	ARARA/PSSP6	CEGL001412	7
<i>Artemisia arbuscula</i> ssp. <i>longiloba</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation	Early Low Sagebrush / Idaho Fescue Shrub Herbaceous Vegetation	ARARL/FEID	CEGL001522	3
<i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Festuca idahoensis</i> Shrubland	Basin Big Sagebrush / Idaho Fescue Shrubland	ARTRT/FEID	CEGL001014	1
<i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Leymus cinereus</i> Shrubland	Basin Big Sagebrush / Basin Wildrye Shrubland	ARTRT/LECI4	CEGL001016	5
<i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Poa secunda</i> Shrubland	Basin Big Sagebrush / Sandberg Bluegrass Shrubland	ARTRT/POSE	CEGL001008	6
<i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation	Basin Big Sagebrush / Bluebunch Wheatgrass Shrub Herbaceous Vegetation	ARTRT/PSSP6	CEGL001018	1
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> - <i>Chamaebatiaria millefolium</i> / <i>Penstemon deustus</i> Shrubland	Mountain Big Sagebrush - Fernbush / Scabland Penstemon Shrubland	ARTRV-CHMI2/PEDE4		5
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> - <i>Purshia tridentata</i> / <i>Poa secunda</i> Shrubland	Mountain Big Sagebrush - Antelope Bitterbrush / Sandberg Bluegrass Shrubland	ARTRV-PUTR2/POSE		5
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> - <i>Purshia tridentata</i> / <i>Pseudoroegneria spicata</i> Shrubland	Mountain Big Sagebrush - Antelope Bitterbrush / Bluebunch Wheatgrass Shrubland	ARTRV-PUTR2/PSSP6	CEGL001032	5
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> - <i>Symphoricarpos oreophilus</i> / <i>Pseudoroegneria spicata</i> Shrubland	Mountain Big Sagebrush - Mountain Snowberry / Bluebunch Wheatgrass Shrubland	ARTRV-SYOR2/PSSP6	CEGL001038	2
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Bromus tectorum</i> Semi-natural Shrubland	Mountain Big Sagebrush / Cheatgrass Semi-natural Shrubland	ARTRV/BRTE		24
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation	Mountain Big Sagebrush / Idaho Fescue Shrub Herbaceous Vegetation	ARTRV/FEID	CEGL001533	9
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Hesperostipa comata</i> Shrubland	Mountain Big Sagebrush / Needle and Thread Shrubland	ARTRV/HECO26	CEGL002931	4
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Leymus cinereus</i> Shrubland	Mountain Big Sagebrush / Basin Wildrye Shrubland	ARTRV/LECI4	CEGL001027	1
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Poa secunda</i> Shrubland	Mountain Big Sagebrush / Sandberg Bluegrass Shrubland	ARTRV/POSE	CEGL001029	15
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Pseudoroegneria spicata</i> Shrubland	Mountain Big Sagebrush / Bluebunch Wheatgrass Shrubland	ARTRV/PSSP6	CEGL001030	32
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Achnatherum hymenoides</i> Shrubland	Wyoming Big Sagebrush / Indian Ricegrass Shrubland	ARTRW8/ACHY	CEGL001046	5

Scientific Name	Common Name	Code	Elcode	N
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Achnatherum thurberianum</i> - <i>Bromus tectorum</i> Semi-natural Shrubland	Wyoming Big Sagebrush / Thurber's Needlegrass - Cheatgrass Semi-natural Shrubland	ARTRW8/ACTH7-BRTE		28
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Achnatherum thurberianum</i> Shrubland	Wyoming Big Sagebrush / Thurber's Needlegrass Shrubland	ARTRW8/ACTH7	CEGL001052	13
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Agropyron cristatum</i> Semi-natural Shrubland	Wyoming Big Sagebrush / Crested Wheatgrass Semi-natural Shrubland	ARTRW8/AGCR		5
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Bromus tectorum</i> Semi-natural Shrubland	Wyoming Big Sagebrush / Cheatgrass Semi-natural Shrubland	ARTRW8/BRTE		70
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Festuca idahoensis</i> Shrubland	Wyoming Big Sagebrush / Idaho Fescue Shrubland	ARTRW8/FEID		11
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Hesperostipa comata</i> Shrubland	Wyoming Big Sagebrush / Needle and Thread Shrubland	ARTRW8/HECO26	CEGL001051	33
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Pascopyrum smithii</i> Shrub Herbaceous Vegetation	Wyoming Big Sagebrush / Western Wheatgrass Shrub Herbaceous Vegetation	ARTRW8/PASM	CEGL001047	9
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Poa secunda</i> - <i>Bromus tectorum</i> Semi-natural Shrubland	Wyoming Big Sagebrush / Sandberg Bluegrass - Cheatgrass Semi-natural Shrubland	ARTRW8/POSE-BRTE		76
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Poa secunda</i> Shrubland	Wyoming Big Sagebrush / Sandberg Bluegrass Shrubland	ARTRW8/POSE	CEGL001049	76
<i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Pseudoroegneria spicata</i> Shrubland	Wyoming Big Sagebrush / Bluebunch Wheatgrass Shrubland	ARTRW8/PSSP6	CEGL001009	15
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Achnatherum thurberianum</i> Shrubland	Threetip Sagebrush / Thurber's Needlegrass Shrubland	ARTRT2/ACTH7		30
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Bromus tectorum</i> Semi-natural Shrubland	Threetip Sagebrush / Cheatgrass Semi-natural Shrubland	ARTRT2/BRTE		40
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation	Threetip Sagebrush / Idaho Fescue Shrub Herbaceous Vegetation	ARTRT2/FEID	CEGL001536	14
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Hesperostipa comata</i> Shrub Herbaceous Vegetation	Threetip Sagebrush / Needle and Thread Shrub Herbaceous Vegetation	ARTRT2/HECO26	CEGL001539	8
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Koeleria macrantha</i> Shrubland	Threetip Sagebrush / Prairie Junegrass Shrubland	ARTRT2/KOMA		9
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Leymus cinereus</i> Shrub Herbaceous Vegetation	Threetip Sagebrush / Basin Wildrye Shrub Herbaceous Vegetation	ARTRT2/LECI4	CEGL002994	1
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Pascopyrum smithii</i> - <i>Bromus tectorum</i> Semi-natural Shrubland	Threetip Sagebrush / Western Wheatgrass - Cheatgrass Semi-natural Shrubland	ARTRT2/PASM-BRTE		11

Scientific Name	Common Name	Code	Elcode	N
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Pascopyrum smithii</i> Shrubland	Threetip Sagebrush / Western Wheatgrass Shrubland	ARTRT2/PASM		19
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Poa secunda</i> - <i>Bromus tectorum</i> Semi-natural Shrubland	Threetip Sagebrush / Sandberg Bluegrass - Cheatgrass Semi-natural Shrubland	ARTRT2/POSE-BRTE		59
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Poa secunda</i> Shrubland	Threetip Sagebrush / Sandberg Bluegrass Shrubland	ARTRT2/POSE		17
<i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation	Threetip Sagebrush / Bluebunch Wheatgrass Shrub Herbaceous Vegetation	ARTRT2/PSSP6	CEGL001538	21
<i>Cercocarpus ledifolius</i> - <i>Purshia tridentata</i> / <i>Poa secunda</i> Shrubland	Curl-leaf Mountain Mahogany - Antelope Bitterbrush / Sandberg Bluegrass Shrubland	CELE3-PUTR2/POSE		2
<i>Chrysothamnus viscidiflorus</i> / <i>Agropyron cristatum</i> Shrubland	Yellow Rabbitbrush / Crested Wheatgrass Shrubland	CHVI8/AGCR		8
<i>Chrysothamnus viscidiflorus</i> / <i>Bromus tectorum</i> Semi-natural Shrubland	Yellow Rabbitbrush / Cheatgrass Semi-natural Shrubland	CHVI8/BRTE		39
<i>Chrysothamnus viscidiflorus</i> / <i>Hesperostipa comata</i> Shrubland	Yellow Rabbitbrush / Needle and Thread Shrubland	CHVI8/HECO26	CEGL002799	1
<i>Chrysothamnus viscidiflorus</i> / <i>Pascopyrum smithii</i> Shrubland	Yellow Rabbitbrush / Western Wheatgrass Shrubland	CHVI8/PASM		3
<i>Chrysothamnus viscidiflorus</i> / <i>Poa secunda</i> - <i>Bromus tectorum</i> Semi-natural Shrubland	Yellow Rabbitbrush / Sandberg Bluegrass - Cheatgrass Semi-natural Shrubland	CHVI8/POSE-BRTE		5
<i>Chrysothamnus viscidiflorus</i> / <i>Pseudoroegneria spicata</i> Shrubland	Yellow Rabbitbrush / Bluebunch Wheatgrass Shrubland	CHVI8/PSSP6		1
<i>Ericameria nana</i> - <i>Holodiscus dumosus</i> / <i>Penstemon deustus</i> Shrubland	Dwarf Goldenbush - Rockspirea / Scabland Penstemon Shrubland	ERNA7-HODU/PEDE4		11
<i>Juniperus osteosperma</i> / <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Hesperostipa comata</i> Wooded Shrubland	Utah Juniper / Wyoming Big Sagebrush / Needle and Thread Wooded Shrubland	JUOS/ARTRW8/HECO26		6
<i>Prunus virginiana</i> / <i>Leymus cinereus</i> Shrubland	Chokecherry / Basin Wildrye Shrubland	PRVI/LECI4		8
<i>Purshia tridentata</i> - <i>Ericameria nana</i> / <i>Penstemon deustus</i> Shrubland	Antelope Bitterbrush - Dwarf Goldenbush / Scabland Penstemon Shrubland	PUTR2-ERNA7/PEDE4		15
<i>Purshia tridentata</i> - <i>Philadelphus lewisii</i> / <i>Poa secunda</i> Shrubland	Antelope Bitterbrush - Lewis' Mock Orange / Sandberg Bluegrass Shrubland	PUTR2-PHLE4/POSE		4
<i>Purshia tridentata</i> / <i>Achnatherum thurberianum</i> Shrubland	Antelope Bitterbrush / Thurber's Needlegrass Shrubland	PUTR2/ACTH7		3
<i>Purshia tridentata</i> / <i>Leymus cinereus</i> Shrub Herbaceous Vegetation	Antelope Bitterbrush / Basin Wildrye Shrub Herbaceous Vegetation	PUTR2/LECI4	CEGL001497	4

Scientific Name	Common Name	Code	Elcode	N
<i>Purshia tridentata</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation	Antelope Bitterbrush / Bluebunch Wheatgrass Shrub Herbaceous Vegetation	PUTR2/PSSP6	CEGL001495	1
Herbaceous Vegetation				
<i>Achnatherum nelsonii</i> / <i>Lupinus argenteus</i> Herbaceous Vegetation	Columbia Needlegrass / Silvery Lupine Herbaceous Vegetation	ACNE9/LUAR3		1
<i>Achnatherum thurberianum</i> - <i>Bromus tectorum</i> Semi- natural Herbaceous Vegetation	Thurber's Needlegrass - Cheatgrass Semi-natural Herbaceous Vegetation	ACTH7-BRTE		9
<i>Agropyron cristatum</i> - <i>Bromus tectorum</i> Semi-natural Herbaceous Vegetation	Crested Wheatgrass - Cheatgrass Semi-natural Herbaceous Vegetation	AGCR-BRTE		38
<i>Bromus tectorum</i> Semi-natural Herbaceous Vegetation	Cheatgrass Semi-natural Herbaceous Vegetation	BRTE	CEGL003019	52
<i>Elymus lanceolatus</i> / <i>Lupinus argenteus</i> Herbaceous Vegetation	Streambank Wheatgrass / Silvery Lupine Herbaceous Vegetation	ELLA3/LUAR3		4
<i>Festuca idahoensis</i> - <i>Pseudoroegneria spicata</i> Herbaceous Vegetation	Idaho Fescue - Bluebunch Wheatgrass Herbaceous Vegetation	FEID-PSSP6	CEGL001624	3
<i>Hesperostipa comata</i> - <i>Poa secunda</i> Herbaceous Vegetation	Needle and Thread - Sandberg Bluegrass Herbaceous Vegetation	HECO26-POSE	CEGL001704	3
<i>Juncus balticus</i> Herbaceous Vegetation	Baltic Rush Herbaceous Vegetation	JUBA	CEGL001838	1
<i>Juniperus (scopulorum, osteosperma)</i> / <i>Ericameria nana</i> Wooded Herbaceous Vegetation	Juniper (Rocky Mountain, Utah) / Dwarf Goldenbush Wooded Herbaceous Vegetation	JUSC2-JUOS/ERNA7		14
<i>Leymus cinereus</i> Herbaceous Vegetation	Basin Wildrye Herbaceous Vegetation	LECI4	CEGL001479	4
<i>Pascopyrum smithii</i> Herbaceous Vegetation	Western Wheatgrass Herbaceous Vegetation	PASM	CEGL001577	2
<i>Poa secunda</i> - <i>Bromus tectorum</i> Semi-natural Herbaceous Vegetation	Sandberg Bluegrass - Cheatgrass Semi-natural Herbaceous Vegetation	POSE-BRTE		99
<i>Poa secunda</i> Herbaceous Vegetation	Sandberg Bluegrass Herbaceous Vegetation	POSE		8
<i>Pseudoroegneria spicata</i> - <i>Melica bulbosa</i> Herbaceous Vegetation	Bluebunch Wheatgrass - Oniongrass Herbaceous Vegetation	PSSP6-MEBU/ARAC2		2
<i>Pseudoroegneria spicata</i> - <i>Poa secunda</i> / <i>Balsamorhiza</i> <i>sagittata</i> Herbaceous Vegetation	Bluebunch Wheatgrass - Sandberg Bluegrass / Arrowleaf Balsamroot Herbaceous Vegetation	PSSP6-POSE/BASA3	CEGL001662	3
<i>Pseudoroegneria spicata</i> - <i>Poa secunda</i> Herbaceous Vegetation	Bluebunch Wheatgrass - Sandberg Bluegrass Herbaceous Vegetation	PSSP6-POSE	CEGL001677	11
<i>Pseudoroegneria spicata</i> / <i>Crepis acuminata</i> Herbaceous Vegetation	Bluebunch Wheatgrass / Tapertip Hawksbeard Herbaceous Vegetation	PSSP6/CRAC2		6

Scientific Name	Common Name	Code	Elcode	N
<i>Sisymbrium altissimum</i> / <i>Bromus tectorum</i> Semi-natural Herbaceous Vegetation	Tall Tumblemustard / Cheatgrass Semi-natural Herbaceous Vegetation	SIAL2/BRTE		24
Sparse Vegetation				
<i>Chamaebatiaria millefolium</i> / <i>Penstemon deustus</i> Sparse Vegetation	Fernbush / Scabland Penstemon Sparse Vegetation	CHMI2/PEDE4		5
<i>Ericameria nana</i> / <i>Poa secunda</i> / <i>Penstemon deustus</i> Sparse Vegetation	Dwarf Goldenbush / Sandberg Bluegrass / Scabland Penstemon Sparse Vegetation	ERNA7/POSE/PEDE4		11
<i>Eriogonum ovalifolium</i> var. <i>depressum</i> Sparse Vegetation	Cushion Buckwheat Sparse Vegetation	EROVD	CEGL001401	4
<i>Philadelphus lewisii</i> / <i>Penstemon deustus</i> Sparse Vegetation	Lewis' Mock Orange / Scabland Penstemon Sparse Vegetation	PHLE4/PEDE4		8
<i>Pinus flexilis</i> / <i>Chamaebatiaria millefolium</i> / <i>Poa secunda</i> Sparse Vegetation	Limber Pine / Fernbush / Sandberg Bluegrass Sparse Vegetation	PIFL2/CHMI2/POSE		4
<i>Poa secunda</i> / <i>Leptodactylon pungens</i> Sparse Vegetation	Sandberg Bluegrass / Granite Prickly Phlox Sparse Vegetation	POSE/LEPU		11

Dichotomous Key to the Classification

The field key to plant communities of the Monument and Preserve follows. The key is structured into physiognomic classes (or lifeform groups). These classes do not constraint the classification *per se*, rather they are employed to assist in applying the classification. In the field, different expressions of a given plant association may occur as different physiognomic classes. For example, *Ericameria nana* - *Holodiscus dumosus* / *Penstemon deustus* Shrubland occurs in stands that are classified as a range of different physiognomies: herbaceous, shrubland, and sparse vegetation (approximately 18%, 36%, and 45% of observations, respectively). Given this, several associations may be found through multiple pathways within the key.

Field Key to Plant Communities of Craters of the Moon National Monument and Preserve

Use of the Key

To use the field key most effectively, identify a representative, homogeneous stand of vegetation. Work through the entire hierarchy of the key, beginning with *Key to Lifeform Groups* to each sequential dichotomous lead. Estimate plant cover on an area of approximately 405m². If a satisfactory determination is not made in stands with low total cover consider adjusting diagnostic species cover break points downward. Assignment of individual species to lifeform follows USDA, NRCS (2007).

In the key the term *relative cover* is used. In this context relative cover refers to the proportional abundance of the given species (or group of species) with respect to the total abundance of the associated lifeform. The value is calculated by dividing the percent cover of the species (or group of species) under consideration by the total cover of the respective lifeform. For example, the relative cover of native grass species is calculated as follows: (sum of native grass species cover within the sample) / (total cover of all graminoids within the sample) = (relative native grass composition).

Key to Lifeform Groups

- 1a) Tree canopy cover (*Juniperus osteosperma*, *Juniperus scopulorum*, *Pinus flexilis*, *Populus balsamifera* ssp. *trichocarpa*, *Populus tremuloides*, or *Pseudotsuga menziesii* alone or combined) $\geq 10\%$, **Part A: Forest, Woodland, and Savannah**, lead 4a.
- 1b) Tree canopy cover $< 10\%$, lead 2a.
 - 2a) Shrub canopy cover $\geq 10\%$, **Part B: Shrublands and Shrub-Steppe**, lead 17a.
 - 2b) Shrub canopy cover $< 10\%$, lead 3a.
 - 3a) Vascular plant cover $\geq 10\%$, **Part C: Herbaceous Vegetation**, lead 81a.
 - 3b) Vascular plant cover $< 10\%$, **Part D: Sparse Vegetation**, lead 106a.

Part A: Forest, Woodland, and Savannah

- 4a) *Populus balsamifera* spp. *trichocarpa* $\geq 10\%$ cover, **POBAT**.
- 4b) *Populus balsamifera* spp. *trichocarpa* $< 10\%$ cover, lead 5a.
 - 5a) *Populus tremuloides* $> 10\%$ cover, lead 6a.
 - 6a) *Pinus flexilis* $\geq 5\%$ cover, **POTR5-PIFL2**.
 - 6b) *Pinus flexilis* $< 5\%$ cover, lead 7a.
 - 7a) *Acer glabrum*, *Amelanchier alnifolia*, *Prunus virginiana*, or *Salix scouleriana* (alone or in combination) $\geq 10\%$ cover (occasionally less abundant in sparse stands), **POTR5/PRVI**.
 - 7b) *Acer glabrum*, *Amelanchier alnifolia*, *Prunus virginiana*, or *Salix scouleriana* (alone or in combination) $< 10\%$ cover, lead 8a.
 - 8a) *Artemisia tridentata* ssp. *vaseyana* $\geq 10\%$ cover, **POTR5/ARTR**.

- 8b) *Artemisia tridentata* ssp. *vaseyana* <10% cover, lead 9a.
 - 9a) *Ribes aureum*, *Ribes cereum*, *Rosa woodsii*, or *Symphoricarpos oreophilus* (alone or in combination) \geq 10% cover (occasionally less abundant in sparse stands), **POTR5/SYOR2/CARU**.
 - 9a) *Ribes aureum*, *Ribes cereum*, *Rosa woodsii*, or *Symphoricarpos oreophilus* (alone or in combination) <10% cover, lead 10a.
 - 10a) *Agastache urticifolia*, *Delphinium occidentale*, *Hackelia floribunda*, *Heracleum maximum*, *Mertensia ciliata*, *Osmorhiza occidentalis*, *Senecio serra*, or *Valeriana occidentalis* (alone or in combination) \geq 10% cover (occasionally less abundant in sparse stands), **POTR5/TALL FORBS**.
 - 10b) *Calamagrostis rubescens* or *Carex geyeri* (alone or in combination) \geq 10% cover (occasionally less abundant in sparse stands), **POTR5/CARU**.
- 5b) *Populus tremuloides* \leq 10, lead 11a.
 - 11a) *Pinus flexilis* \geq 5% cover, **PIFL2/PUTR2**.
 - 11b) *Pinus flexilis* <5% cover, lead 12a.
 - 12a) *Pseudotsuga menziesii* \geq 5% cover, lead 13a.
 - 13a) *Carex geyeri* \geq 5% cover, **PSME/CAGE2**.
 - 13b) *Carex geyeri* <5% and *Prunus virginiana* or *Symphoricarpos oreophilus* (alone or in combination) \geq 5% cover (occasionally less abundant in sparse stands), **PSME/SYOR2**.
 - 12b) *Pseudotsuga menziesii* <5% cover and *Juniperus osteosperma* or *Juniperus scopulorum* is the principal tree species, lead 14a.
 - 14a) *Juniperus scopulorum* \geq 10% cover, **JUSC2/POSE/PEDE4**.
 - 14b) *Juniperus scopulorum* <10% cover, lead 15a.
 - 15a) *Chamaebatiaria millefolium* is present, **JUOS-JUSC2/ARTRW8/CHMI2**.
 - 15b) *Chamaebatiaria millefolium* is absent, lead 16a.
 - 16a) *Ericameria nana* is present, **JUSC2-JUOS/ERNA7**.
 - 16b) *Ericameria nana* is not present, **JUOS/ARTRW8/HECO26**.

Part B: Shrublands and Shrub-Steppe

- 17a) *Alnus incana* or *Prunus virginiana* (alone or in combination) \geq 10% cover, lead 18a.
 - 18a) *Alnus incana* \geq 10% cover, **ALIN2/Mesic Forbs**.
 - 18a) *Alnus incana* <10% cover, **PRVI/LECI4**.
- 17b) *Alnus incana* or *Prunus virginiana* (alone or in combination) <10% cover, lead 19a.
 - 19a) *Artemisia tripartita* spp. *tripartita* \geq 8 % cover, lead 20a.
 - 20a) Native grass composition <45% relative cover, **ARTRT2/BRTE**.
 - 20b) Native grass composition \geq 45% relative cover, lead 21a.
 - 21a) *Hesperostipa comata* \geq 18% of total graminoid cover, **ARTRT2/HECO26**.
 - 21b) *Hesperostipa comata* <18% relative graminoid cover, lead 22a.
 - 22a) Native grass composition \geq 74% relative cover, lead 23a.
 - 23a) *Pseudoroegneria spicata* \geq 30% relative graminoid cover, **ARTRT2/PSSP6**.
 - 23b) *Pseudoroegneria spicata* <30% relative graminoid cover, lead 24a.
 - 24a) *Poa secunda* \geq 42% relative graminoid cover, **ARTRT2/POSE**.
 - 24b) *Poa secunda* <42% relative graminoid cover, lead 25a.
 - 25a) *Pascopyrum smithii* <28% relative graminoid cover, lead 26a.
 - 26a) *Festuca idahoensis* \geq 15% relative cover¹, **ARTRT2/FEID**.
 - 26b) *Festuca idahoensis* <15% relative cover, lead 27a.
 - 27a) *Koeleria macrantha* <20% relative cover, **ARTRT2/ACTH7**.
 - 27b) *Koeleria macrantha* \geq 20% relative cover, **ARTRT2/KOMA**.
 - 25b) *Pascopyrum smithii* \geq 28% relative graminoid cover, **ARTRT2/PASM**.
 - 22b) Native grass composition <74% relative cover, lead 28a.
 - 28a) *Poa secunda* \geq 15% relative cover, **ARTRT2/POSE-BRTE**.

¹ Relative cover through lead 29 is in reference to relative graminoid cover.

- 54a) *Festuca idahoensis* $\geq 1\%$ cover, **ARTRV/FEID.**
- 54b) *Festuca idahoensis* $< 1\%$ cover, **ARTRV/POSE.**
- 53b) *Pseudoroegneria spicata* $\geq 2\%$ cover, **ARTRV/PSSP6.**
- 44b) *Artemisia tridentata* ssp. *vaseyana* < 3 , lead 55a.
- 55a) *Chrysothamnus viscidiflorus* $< 5\%$ cover, lead 56a.
- 56a) *Artemisia arbuscula* ssp. *arbuscula* $\geq 3\%$ cover, lead 57a.
- 57a) *Pseudoroegneria spicata* $\geq 2\%$ cover, **ARARA/PSSP6.**
- 57b) *Pseudoroegneria spicata* $< 2\%$ cover, lead 58a.
- 58a) *Hesperostipa comata* is present, **ARARA/HECO26.**
- 58b) *Hesperostipa comata* is not present, lead 59a.
- 59a) *Poa secunda* $\geq 2\%$ cover, **ARARA/POSE.**
- 59b) *Poa secunda* $< 2\%$ cover, **ARARA/BRTE.**
- 56b) *Artemisia arbuscula* ssp. *arbuscula* $< 3\%$ cover, lead 60a.
- 60a) *Artemisia arbuscula* ssp. *longiloba* $\geq 10\%$ cover, **ARARL/FEID.**
- 60b) *Artemisia arbuscula* ssp. *longiloba* $< 10\%$ cover, lead 61a.
- 61a) *Artemisia tridentata* ssp. *tridentata* $\geq 5\%$ cover, lead 62a.
- 62a) *Festuca idahoensis* $\geq 2\%$ cover, **ARTRT/FEID.**
- 62b) *Festuca idahoensis* $< 2\%$ cover, lead 63a.
- 63a) *Leymus cinereus* $\geq 2\%$ cover, **ARTRT/LECI4.**
- 63b) *Leymus cinereus* $\leq 2\%$ cover, lead 64a.
- 64a) *Pseudoroegneria spicata* $\geq 2\%$ cover, **ARTRT/PSSP6.**
- 64b) *Pseudoroegneria spicata* $< 2\%$ cover, **ARTRT/POSE.**
- 61b) *Artemisia tridentata* ssp. *tridentata* $< 5\%$ cover, lead 65a.
- 65a) *Artemisia tripartita* spp. *tripartita* $\geq 2\%$ cover, lead 20a (ARTRT2).
- 65b) *Artemisia tripartita* spp. *tripartita* $< 2\%$ cover, lead 66a.
- 66a) *Ericameria nana* $\geq 3\%$ cover, lead 67a.
- 67a) Tree cover $\geq 2\%$, lead 14a.
- 67b) Tree cover $< 2\%$, lead 107a.
- 66b) *Ericameria nana* $< 3\%$ cover, lead 68a.
- 68a) *Philadelphus lewisii* $> 2\%$ cover, **PHLE4/PEDE4.**
- 68b) *Philadelphus lewisii* $\leq 2\%$ cover, lead 69a.
- 69a) *Cercocarpus ledifolius* $\geq 5\%$ cover,
CELE3-PUTR2/POSE.
- 69b) *Cercocarpus ledifolius* $< 5\%$ cover, lead 70a.
- 70a) *Pinus flexilis* $\geq 1\%$ cover, **PIFL2/PUTR2.**
- 70b) *Pinus flexilis* $< 1\%$ cover, lead 71a.
- 71a) *Achnatherum thurberianum* $\geq 2\%$ cover,
PUTR2/ACTH7.
- 71b) *Achnatherum thurberianum* $< 2\%$ cover, lead 72a.
- 72a) *Leymus cinereus* present, **PUTR2/LECI4.**
- 72b) *Leymus cinereus* is absent, lead 73a.
- 73a) *Pseudoroegneria spicata* $\geq 2\%$ cover,
PUTR2/PSSP6.
- 73b) *Pseudoroegneria spicata* $< 2\%$ cover,
lead 74a.
- 74a) *Philadelphus lewisii* $\geq 1\%$ cover,
PUTR2-PHLE4/POSE.
- 74b) *Philadelphus lewisii* $< 1\%$ cover,
PUTR2-ERNA7/PEDE4.
- 55b) *Chrysothamnus viscidiflorus* $\geq 5\%$ cover, lead 75a.
- 75a) *Artemisia tripartita* spp. *tripartita* $\geq 3\%$ cover, lead 20a (ARTRT2).
- 75b) *Artemisia tripartita* spp. *tripartita* $< 3\%$ cover, lead 76a.
- 76a) Native grass composition $< 70\%$ relative cover, lead 77a.

- 77a) *Agropyron cristatum* or *Agropyron fragile* (alone or in combination) $\geq 2\%$ cover, **CHVI8/AGCR**.
- 77b) *Agropyron cristatum* or *Agropyron fragile* (alone or in combination) $< 2\%$ cover, **CHVI8/BRTE**.
- 76b) Native grass composition $\geq 70\%$ relative cover, lead 78a.
 - 78a) *Hesperostipa comata* $\geq 5\%$ cover, **CHVI8/HECO26**.
 - 78b) *Hesperostipa comata* $< 5\%$ cover, lead 79a.
 - 79a) *Pseudoroegneria spicata* $\geq 5\%$ cover, **CHVI8/PSSP6**.
 - 79b) *Pseudoroegneria spicata* $< 5\%$ cover, lead 80a.
 - 80a) *Achnatherum thurberianum* or *Pascopyrum smithii* (alone or in combination) $\geq 5\%$ cover, **CHVI8/PASM**.
 - 80b) *Achnatherum thurberianum* or *Pascopyrum smithii* (alone or in combination) $< 5\%$ cover, **CHVI8/POSE-BRTE**.

Part C: Herbaceous Vegetation

- 81a) Relative cover of trees, shrubs, and dwarf-shrubs (combined) $\geq 21\%$, lead 82a.
 - 82a) *Artemisia tridentata* spp. *wyomingensis* $< 3\%$ cover, lead 83a.
 - 83a) *Artemisia tridentata* spp. *wyomingensis* and *Chrysothamnus viscidiflorus*, combined, $< 6\%$ cover, lead 84a.
 - 84a) *Artemisia tripartita* spp. *tripartita* $\geq 3\%$ cover, lead 20a (ARTRT2).
 - 84b) *Artemisia tripartita* spp. *tripartita* $< 3\%$ cover, lead 85a.
 - 85a) *Artemisia tridentata* ssp. *vaseyana* $\geq 3\%$ cover, lead 44a (ARTRV).
 - 85b) *Artemisia tridentata* ssp. *vaseyana* $< 3\%$ cover, lead 86a.
 - 86a) *Artemisia arbuscula* ssp. *arbuscula* $\geq 4\%$ cover, lead 56a (ARARA).
 - 86b) *Artemisia arbuscula* ssp. *arbuscula* $< 4\%$ cover, lead 87a.
 - 87a) *Ericameria nana* is present, lead 88a.
 - 88a) Total vegetative cover $\geq 10\%$, lead 67a.
 - 88b) Total vegetative cover $< 10\%$, lead 107a.
 - 87b) *Ericameria nana* is absent, lead 90a.
 - 83b) *Artemisia tridentata* spp. *wyomingensis* and *Chrysothamnus viscidiflorus*, combined, $\geq 6\%$ cover, lead 75a (CHVI8).
 - 82b) *Artemisia tridentata* spp. *wyomingensis* $\geq 3\%$ cover, lead 89a.
 - 89a) *Artemisia tridentata* spp. *wyomingensis* and *Chrysothamnus viscidiflorus*, combined, $\geq 6\%$ cover, lead 31a (ARTRW8).
 - 89b) *Artemisia tridentata* spp. *wyomingensis* and *Chrysothamnus viscidiflorus*, combined, $< 6\%$ cover, lead 81b.
 - 81b) Relative cover of trees, shrubs, and dwarf-shrubs (combined) $< 21\%$, lead 90a.
 - 90a) *Agropyron cristatum* or *Agropyron fragile* (alone or in combination) $\geq 3\%$ cover, **AGCR-BRTE**.
 - 90b) *Agropyron cristatum* or *Agropyron fragile* (alone or in combination) $< 3\%$ cover, lead 91a.
 - 91a) Native grass composition $< 39\%$ relative cover, lead 92a.
 - 92a) *Sisymbrium altissimum* $\geq 3\%$ cover, **SIAL2/BRTE**.
 - 92b) *Sisymbrium altissimum* $< 3\%$ cover, lead 93a.
 - 93a) Combined cover of native perennial grass species (*Achnatherum hymenoides*, *Achnatherum thurberianum*, *Elymus elymoides*, *Elymus lanceolatus*, *Festuca idahoensis*, *Hesperostipa comata*, *Leymus cinereus*, *Pascopyrum smithii*, *Poa secunda*, or *Pseudoroegneria spicata*) $\geq 3\%$ cover, **POSE-BRTE**.
 - 93b) Combined cover of native perennial grass species $< 3\%$ cover, **BRTE**.
 - 91b) Native grass composition $\geq 39\%$ relative cover, lead 94a.
 - 94a) *Hesperostipa comata* $\geq 10\%$ cover, **HECO26-POSE**.
 - 94b) *Hesperostipa comata* $< 10\%$ cover, lead 95a.
 - 95a) *Bromus tectorum* $\geq 2\%$ cover, lead 96a.
 - 96a) *Achnatherum thurberianum* $\geq 5\%$ cover, **ACTH7-BRTE**.
 - 96b) *Achnatherum thurberianum* $< 5\%$ cover, **POSE-BRTE**.
 - 95b) *Bromus tectorum* $< 2\%$ cover, lead 97a.

- 97a) *Elymus lanceolatus* $\geq 3\%$ cover, **ELLA3/LUAR3**.
- 97b) *Elymus lanceolatus* $< 3\%$ cover, lead 98a.
 - 98a) *Festuca idahoensis* $\geq 2\%$ cover, **FEID-PSSP6**.
 - 98b) *Festuca idahoensis* $< 2\%$ cover, lead 99a.
 - 99a) *Leymus cinereus* $\geq 2\%$ and *Pseudoroegneria spicata* $< 1\%$ cover, **LECI4**.
 - 99b) *Leymus cinereus* $< 2\%$ cover, lead 100a.
 - 100a) *Pascopyrum smithii* $\geq 3\%$ cover, **PASM**.
 - 100b) *Pascopyrum smithii* $< 3\%$ cover, lead 101a.
 - 101a) *Pseudoroegneria spicata* $\geq 3\%$ cover, lead 102a.
 - 102a) *Melica bulbosa* $\geq 3\%$ cover, **PSSP6-MEBU/ARAC2**.
 - 102b) *Melica bulbosa* $< 3\%$ cover, lead 103a.
 - 103a) *Balsamorhiza sagittata* $\geq 5\%$ cover, **PSSP6-POSE/BASA3**.
 - 103b) *Balsamorhiza sagittata* $< 5\%$ cover, lead 104a.
 - 104a) *Poa secunda* $< 2\%$ cover, **PSSP6/CRAC2**.
 - 104b) *Poa secunda* $\geq 2\%$ cover, **PSSP6-POSE**.
 - 101b) *Pseudoroegneria spicata* $< 3\%$ cover, lead 105a.
 - 105a) *Leptodactylon pungens* is present, **POSE/LEPU**.
 - 105a) *Leptodactylon pungens* is not present, **POSE**.

Part D: Sparse Vegetation

- 106a) *Agropyron cristatum* $\geq 2\%$ cover, **AGCR**.
- 106b) *Agropyron cristatum* $< 2\%$ cover, lead 107a.
 - 107a) *Ericameria nana* present, lead 108a.
 - 108a) *Juniperus scopulorum* or *Juniperus osteosperma* (alone or in combination) $\geq 1\%$ cover, **JUSC2-JUOS/ERNA7**.
 - 108b) *Juniperus scopulorum* or *Juniperus osteosperma* (alone or in combination) $< 1\%$ cover, lead 109a.
 - 109a) *Holodiscus dumosus* $\geq 1\%$ cover, **ERNA7-HODU/PEDE4**.
 - 109b) *Holodiscus dumosus* $< 1\%$ cover, lead 110a.
 - 110a) *Purshia tridentata* $\geq 1\%$ cover, **PUTR2-ERNA7/PEDE4**.
 - 110b) *Purshia tridentata* $< 1\%$ cover, **ERNA7/POSE/PEDE4**.
 - 107b) *Ericameria nana* is not present, lead 111a.
 - 111a) *Philadelphus lewisii* is not present, lead 112a.
 - 112a) *Juniperus osteosperma* $\geq 1\%$ cover, **JUOS/ARTRW8/HECO26**.
 - 112b) *Juniperus osteosperma* $< 1\%$ cover, lead 113a.
 - 113a) *Purshia tridentata* is present, lead 114a.
 - 114a) *Leymus cinereus* is present, **PUTR2/LECI4**.
 - 114b) *Leymus cinereus* is not present, **PUTR2-ERNA7/PEDE4**.
 - 113b) *Purshia tridentata* is not present, lead 115a.
 - 115a) *Eriogonum ovalifolium* var. *depressum* is present, **EROVD**.
 - 115b) *Eriogonum ovalifolium* var. *depressum* is not present, lead 116a.
 - 116a) *Chamaebatiaria millefolium* is present, **CHMI2/PEDE4**.
 - 116b) *Chamaebatiaria millefolium* is not present, lead 117a.
 - 117a) *Ericameria nana* and *Holodiscus dumosus* each are present, **ERNA7/POSE/PEDE4**.
 - 117b) *Ericameria nana* and *Holodiscus dumosus* each are not present, **POSE/LEPU**.
 - 111b) *Philadelphus lewisii* is present, lead 118a.
 - 118a) *Purshia tridentata* $\geq 1\%$ cover, **PUTR2-PHLE3/POSE**.
 - 118b) *Purshia tridentata* $< 1\%$ cover, lead 119a.
 - 119a) *Pinus flexilis* $\geq 1\%$ cover, **PIFL2/CHMI2/POSE**.
 - 119b) *Pinus flexilis* $< 1\%$ cover, **PHLE4/PEDE4**.

Literature Cited

- Bowerman, T. S., J. Dorr, S. Leahy, K. Varga, and J. Warrick. 1997. Targhee National Forest ecological unit inventory. USDA Forest Service, Targhee National Forest, St. Anthony, ID. 789 pp.
- Caicco, S.L. 1987. National Natural Landmark Evaluation, Great Rift System (Idaho), Columbia Plateau Province, *Artemisia tridentata* (big sagebrush) Steppe Theme. Unpublished report prepared for U.S. Department of the Interior, National Park Service. Idaho Natural Heritage Program, Idaho Department of Fish and Game. 39 pp. plus appendices.
- Caicco, S.L. and C.A. Wellner. 1983a. Research Natural Area Recommendation for Big Juniper Kipuka. Unpublished report on file at U.S. Department of the Interior, Bureau of Land Management, Shoshone Field Office, Shoshone. 15 pp.
- Caicco, S.L. and C.A. Wellner. 1983b. Research Natural Area Recommendation for Brass Cap Kipuka. Unpublished report on file at U.S. Department of the Interior, Bureau of Land Management, Shoshone Field Office, Shoshone. 15 pp.
- Caicco, S.L. and C.A. Wellner. 1983c. Research Natural Area Recommendation for Sand Kipuka. Unpublished report on file at U.S. Department of the Interior, Bureau of Land Management, Shoshone Field Office, Shoshone. 14 pp.
- Cooper, S. V., K. E. Neiman, and D. W. Roberts. 1991. Forest habitat types of northern Idaho: a second approximation. USDA Forest Service General Technical Report INT-236. Intermountain Research Station, Ogden. 143 pp.
- Crowe, E. A., and R. R. Clausnitzer. 1995. Mid-montane wetlands classification of the Malheur, Umatilla, and Wallowa-Whitman National Forests. Wallowa-Whitman National Forest, Pacific Northwest Region, USDA Forest Service. 188 pp. plus appendices.
- Daubenmire, R. 1970. Steppe vegetation of Washington. Washington State University Technical Bulletin 62. Washington Agricultural Experiment Station, College of Agriculture, Pullman. 131 pp.
- Day, T. A. 1985. Plant association and soil factors in primary succession on cinder cones in Idaho. M.S. thesis, University of Idaho, Moscow. 62 pp.
- Day, T.A. and R.G. Wright. 1985. The vegetation types of Craters of the Moon National Monument. Bulletin Number 38, Forest, Wildlife and Range Experiment Station, University of Idaho, Moscow. 6 pp. plus map.
- Eggler, W. A. 1941. Primary succession on volcanic deposits in southern Idaho. *Ecological Monographs* 11(3): 277-298.

Forman, A. D., R. D. Blew, and J. R. Hafila. 2003. Sagebrush steppe ecosystem reserve plant community classifications. STOLLER-ESER-73. Environmental Surveillance, Education, Research Program, S. M. Stoller Corporation, Idaho Falls.

FGDC. 1997. Vegetation Classification Standard. FGDC-STD-005. Vegetation Subcommittee, Federal Geographic Data Committee, FGDC Secretariat, U.S. Geological Survey. Reston, VA. 58pp.

FGDC. 2006. National Vegetation Classification Standard Version 2 (submitted draft November 30, 2006). FGDC-STD-005 (Version 2). Vegetation Subcommittee, Federal Geographic Data Committee, FGDC Secretariat, U.S. Geological Survey. Reston, VA. 128pp.

French, N.R. and J.E. Mitchell. 1983. Long-term vegetation changes in permanent quadrats at the Idaho National Engineering Laboratory. Forestry, Wildlife, and Range Experiment Station Bulletin Number 36, University of Idaho, Moscow. 42 pp.

Hironaka, M., M.A. Fosberg, A.H. Winward. 1983. Sagebrush-grass Habitat Types of Southern Idaho. Forest, Wildlife, and Range Experiment Station. University of Idaho, Moscow, Idaho. Bulletin No. 35.

Jennings, M. D., D. Faber-Langendoen, R. K. Peet, O. L. Loucks, D. C. Glenn-Lewin, A. Damman, M. G. Barbour, R. Pfister, D. H. Grossman, D. Roberts, D. Tart, M. Walker, S. S. Talbot, J. Walker, G. S. Hartshorn, G. Waggoner, M. D. Abrams, A. Hill, M. Rejmanek. 2006. Description, documentation, and evaluation of associations and alliances within the U.S. National Vegetation Classification, Version 4.5. Ecological Society of America, Vegetation Classification Panel. Washington. 119p.

Johnson, C.G. Jr., and S.A. Simon. 1987. Plant associations of the Wallowa-Snake Province, Wallowa-Whitman National Forest. USDA Forest Service, Pacific Northwest Region, Wallowa-Whitman National Forest, R6-ECOL-TP-255A-86. 400 pp.

Johnson, M. E. 1991. Soil survey of Blaine County area, Idaho. USDA Soil Conservation Service, National Cooperative Soil Survey. U.S. Government Printing Office, Washington D.C. 368 pp. plus maps.

Jurs, L. P., and A. R. Sands. 2004. An inventory, assessment, and recommended management of sagebrush steppe vegetation in Laidlaw Park, Little Park, and Paddelford Flat, Craters of the Moon National Monument and Preserve, Idaho. Unpublished report prepared by Northwest Natural Resource Services and The Nature Conservancy for US Department of the Interior, Bureau of Land Management, Boise. 45 pp. plus appendices.

Kuntz, M. A., D. E. Champion, E. C. Spiker, R. H. Lefebvre, and L. A. McBroome. 1982. The Great Rift and evolution of the Craters of the Moon Lava Field, Idaho. In: Bonnicksen, B. and R. M. Breckenridge, editors. 1982. Cenozoic geology of Idaho. Idaho Bureau of Mines and Ecology, Bulletin No. 26. 725 pp.

Kuntz, M. A., D. E. Champion, R. H. Lefebvre, and H. R. Covington. 1988. Geologic map of the Craters of the Moon, Kings Bowl, and Wapi lava fields, and the Great Rift volcanic rift zone, south-central Idaho. USDI Geological Survey Miscellaneous Investigations Series Map I-1632.

Lauer, J.L. and J.M. Peek. 1976. Big game-livestock relationships on the bighorn sheep winter range, East Fork Salmon River. Forest, Wildlife, and Range Exp. Sta. Bull. No. 12. Univ. of Idaho, Moscow. 44 pp.

Manning, M. E., and W. G. Padgett. 1995. Riparian community type classification for the Humboldt and Toiyabe National Forests, Nevada and eastern California. USDA Forest Service, Intermountain Region Ecology and Classification Program, Ogden, UT. 274 pp.

McBride, R, N. R. French, A. H. Dahl, and J. E. Detmer. 1978. Vegetation types and surface of the Idaho National Engineering Laboratory Site. Environmental Sciences Branch, Radiological and Environmental Sciences Laboratory, U. S. Department of Energy, Idaho Falls.

McCune, B. and M. J. Mefford. 2006. PC-ORD. Multivariate Analysis of Ecological Data, Version 5.06. MjM Software, Gleneden Beach, Oregon, U.S.A.

McCune, B. and J. B. Grace. 2002. Analysis of Ecological Communities. With a contribution by Dean L. Urban. MjM Software, Gleneden Beach. 304 pp.

Moseley, R.K. 1989. National Natural Landmark Evaluation, Craters of the Moon National Monument, Idaho, Columbia Plateau Natural Region Low Sagebrush Theme, Low sagebrush/Idaho Fescue Subtheme. Unpublished Report provided by Idaho Natural Heritage Program, Idaho Department of Fish and Game. Boise, ID. pp. 34.

Mueggler, W. F. 1988. Aspen community types of the Intermountain Region. USDA Forest Service General Technical Report INT-250. Intermountain Research Station, Ogden. 135 pp.

Mueggler, W. F., and C. A. Harris. 1969. Some vegetation and soil characteristics of mountain grasslands in central Idaho. *Ecology* 50(4): 671-678.

NatureServe. 2007. NatureServe Explorer: An online encyclopedia of life [web application]. Version 6.2. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed on numerous dates; data current as of June 8, 2007).

Nelson, L. P., and M. E. Jensen. 1987. Sagebrush-grass community types of the Humboldt National Forest. Unpublished report prepared for USDA Forest Service, Humboldt National Forest. 80 pp. plus appendices.

Padgett, W. G., A. P. Youngblood, and A. H. Winward. 1989. Riparian community type classification of Utah and southeastern Idaho. USDA Forest Service R4-Ecol-89-01. Intermountain Region, Ogden, UT. 191 pp.

R Development Core Team. 2007. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna. Available online at <http://www.R-project.org>.

Reid, M., P. Comer, H. Barrett, S. Caicco, R. Crawford, C. Jean, G. Jones, J. Kagan, M. Karl, G. Kittel, P. Lyon, M. Manning, E. Peterson, R. Rosentreter, and S. Rust. 2002. International classification of ecological communities: Terrestrial vegetation of the United States. Sagebrush Vegetation of the Western United States. Final Report for the USGS Forest and Rangeland Ecosystem Science Center, Boise. NatureServe, Arlington. 154 pp. plus appendices.

Rust, S. K. 1999. Pinyon-juniper woodland classification and description in Research Natural Areas of southeastern Idaho. In: S. B. Monsen and R. Stevens (compilers). Proceedings: ecology and management of pinyon-juniper communities within the interior west. 1997 September 15-18, Provo, UT. Proc. RMRS-P-9. Ogden, UT: USDA Forest Service, Rocky Mountain Research Station. 411 pp.

Rust, S. K., C. J. Murphy, C. L. Coulter, A. G. Tackett, and G. B. Easdale. 2001. Inventory and Assessment of Upland Vegetation on the Pioneer Mountain Ranch. Unpublished report prepared in cooperation with U. S. Department of the Interior, Bureau of Land Management, Shoshone Field Office and The Nature Conservancy, Idaho Field Office for Lava Lake Land & Livestock, LLC, by Idaho Conservation Data Center, Department of Fish and Game, Boise. 83 pp.

Schlatterer, E. F. 1972. A preliminary description of plant communities found on the Sawtooth, White Cloud, Boulder, and Pioneer Mountains. USDA, Forest Service, Intermountain Region, Ogden, UT. 111 pp.

Steele, R., R. D. Pfister, R. A. Ryker, and J. A. Kittams. 1981. Forest habitat types of central Idaho. USDA Forest Service General Technical Report INT-114. Intermountain Forest and Range Experiment Station, Ogden, UT. 138 pp.

Steele, R., S. V. Cooper, D. M. Ondov, D. W. Roberts, and R. D. Pfister. 1983. Forest habitat types of eastern Idaho-western Wyoming. General Technical Report INT-144. USDA Forest Service, Intermountain Forest and Range Experiment Station, Ogden, UT. 122 pp.

Tart, D., C. Williams, J. DiBenedetto, E. Crowe, M. Girard, H. Gordon, K. Sleavin, M. Manning, J. Haglund, B. Short, and D. Wheeler. 2005. Section 2: Existing vegetation classification protocol. In: R. Brohman and L. Bryant, eds. Existing Vegetation Classification and Mapping Technical Guide. USDA Forest Service, Ecosystem Management Coordination Staff, Gen. Tech. Rep. WO-67. Washington. 34 pp.

The Nature Conservancy and Environmental Systems Research Institute. 1994a. Field Methods for Vegetation Mapping. Unpublished report prepared for US Department of the Interior, National Biological Survey and National Park Service by The Nature Conservancy and Environmental Systems Research Institute, Arlington.

The Nature Conservancy and Environmental Systems Research Institute. 1994b. Standardized National Vegetation Classification System. Unpublished report prepared for US Department of the Interior, National Biological Survey and National Park Service by The Nature Conservancy and Environmental Systems Research Institute, Arlington.

The Nature Conservancy. 1996. Methodology for assessing the utility of existing data for vegetation mapping. Unpublished report prepared for US Department of the Interior, Biological Resources Division and National Park Service by The Nature Conservancy, Arlington.

Tisdale, E. W. 1986. Canyon grasslands and associated shrublands of west-central Idaho and adjacent areas. Bulletin Number 40; Forest, Wildlife and Range Experiment Station; Moscow. 42 pp.

Tisdale, E. W., M. Hironaka, and M. A. Fosberg. 1965. An area of pristine vegetation in Craters of the Moon National Monument, Idaho. *Ecology* 46: 349-352.

Tuhy, J. S., and S. Jensen. 1982. Riparian classification for the Upper Salmon/Middle Fork Salmon River drainages, Idaho. White Horse Associates, Smithfield, UT. 183 pp.

US Department of the Interior, Bureau of Land Management. 2001. Shoshone Field Office fuels inventory line-intercept data. Unpublished fuels inventory data maintained by US Department of the Interior, Bureau of Land Management, Shoshone Field Office. Shoshone.

US Department of the Interior National Park Service and Bureau of Land Management. 2005. Craters of the Moon National Monument and Preserve Proposed Management Plan and Final Environmental Impact Statement. Document online at <http://parkplanning.nps.gov/>.

USDA, NRCS. 2007. The PLANTS Database (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

Western Regional Climate Center. 2007. Western US climate historical summaries: Idaho climate summaries. National Oceanic and Atmospheric Administration, United States Department of Commerce, Western Regional Climate Center, Reno. Online electronic data source : <http://www.wrcc.dri.edu/summary/climsmid.html>.

Winward, A. H. 1970. Taxonomic and ecological relationships of the big sagebrush complex in Idaho. Unpublished dissertation, University of Idaho, Moscow. 80 pp.

Northwest Management Inc. 2006. Report of the NPS Vegetation Mapping Project at Craters of the Moon National Monument. Unpublished report prepared for US Department of the Interior, Inventory and Monitoring Program, Upper Columbia Basin Network, Moscow, 18 pp.

Youngblood, A. P., W. G. Padgett, and A. H. Winward. 1985. Riparian community type classification of eastern Idaho - western Wyoming. USDA Forest Service R4-Ecol-85-01. Intermountain Region, Ogden, UT. 78 pp.

Appendix 1. Cross-walk to vegetation types. Vegetation types described by Day and Wright (1985) are cross-walked to plant associations identified in this classification. Vegetation types are listed in the order they appear in Day and Wright (1985) with the equivalent plant community common and scientific name.

| Vegetation Type (Day and Wright 1985) | Equivalent Common Name | Equivalent Scientific Name |
|--|--|---|
| Cinder Gardens | Cushion Buckwheat Sparse Vegetation | <i>Eriogonum ovalifolium</i> var. <i>depressum</i> Sparse Vegetation |
| Low Density Lava Flows and Medium Density Lava Flows | Mountain Big Sagebrush - Fernbush / Scabland Penstemon Shrubland | <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> - <i>Chamaebatiaria millefolium</i> / <i>Penstemon deustus</i> Shrubland |
| | Fernbush / Scabland Penstemon Sparse Vegetation | <i>Chamaebatiaria millefolium</i> / <i>Penstemon deustus</i> Sparse Vegetation |
| | Dwarf Goldenbush - Rockspirea / Scabland Penstemon Shrubland | <i>Ericameria nana</i> - <i>Holodiscus dumosus</i> / <i>Penstemon deustus</i> Shrubland |
| | Dwarf Goldenbush / Sandberg Bluegrass / Scabland Penstemon Sparse Vegetation | <i>Ericameria nana</i> / <i>Poa secunda</i> / <i>Penstemon deustus</i> Sparse Vegetation |
| | Juniper (Rocky Mountain, Utah) / Dwarf Goldenbush Herbaceous Vegetation | <i>Juniperus (scopulorum, osteosperma)</i> / <i>Ericameria nana</i> Herbaceous Vegetation |
| | Antelope Bitterbrush - Dwarf Goldenbush / Scabland Penstemon Shrubland | <i>Purshia tridentata</i> - <i>Ericameria nana</i> / <i>Penstemon deustus</i> Shrubland |
| | Antelope Bitterbrush - Lewis' Mock Orange / Sandberg Bluegrass Shrubland | <i>Purshia tridentata</i> - <i>Philadelphus lewisii</i> / <i>Poa secunda</i> Shrubland |
| Mountain Big Sagebrush / Bluebunch Wheatgrass | Mountain Big Sagebrush / Bluebunch Wheatgrass Shrubland | <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Pseudoroegneria spicata</i> Shrubland |
| Mountain Big Sagebrush / Sandberg Bluegrass | Mountain Big Sagebrush / Sandberg Bluegrass Shrubland | <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Poa secunda</i> Shrubland |
| Mountain Big Sagebrush / Needle Grass | Mountain Big Sagebrush / Needle and Thread Shrubland | <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Hesperostipa comata</i> Shrubland |
| Mountain Big Sagebrush / Idaho Fescue | Mountain Big Sagebrush / Idaho Fescue Shrub Herbaceous Vegetation | <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation |
| Big Sagebrush / Cheatgrass | not sampled | |
| Three-tip Sagebrush / Idaho Fescue | Threetip Sagebrush / Idaho Fescue Shrub Herbaceous Vegetation | <i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation |

| Vegetation Type (Day and Wright 1985) | Equivalent Common Name | Equivalent Scientific Name |
|---|--|---|
| Early Low Sagebrush / Idaho Fescue | Early Low Sagebrush / Idaho Fescue Shrub Herbaceous Vegetation | <i>Artemisia arbuscula</i> ssp. <i>longiloba</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation |
| Low Sagebrush / Sandberg Bluegrass | Low Sagebrush / Sandberg Bluegrass Shrub Herbaceous Vegetation | <i>Artemisia arbuscula</i> ssp. <i>arbuscula</i> / <i>Poa secunda</i> Shrub Herbaceous Vegetation |
| Low Sagebrush / Idaho Fescue | Low Sagebrush / Idaho Fescue Shrub Herbaceous Vegetation | <i>Artemisia arbuscula</i> ssp. <i>arbuscula</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation |
| Antelope Bitterbrush | Antelope Bitterbrush / Bluebunch Wheatgrass Shrub Herbaceous Vegetation | <i>Purshia tridentata</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation |
| Antelope Bitterbrush / Great Basin Wildrye | Antelope Bitterbrush / Basin Wildrye Shrub Herbaceous Vegetation | <i>Purshia tridentata</i> / <i>Leymus cinereus</i> Shrub Herbaceous Vegetation |
| Bluebunch Wheatgrass / Idaho Fescue | Idaho Fescue - Bluebunch Wheatgrass Herbaceous Vegetation | <i>Festuca idahoensis</i> - <i>Pseudoroegneria spicata</i> Herbaceous Vegetation |
| Bluebunch Wheatgrass / Sandberg Bluegrass | Bluebunch Wheatgrass - Sandberg Bluegrass / Arrowleaf Balsamroot Herbaceous Vegetation | <i>Pseudoroegneria spicata</i> - <i>Poa secunda</i> / <i>Balsamorhiza sagittata</i> Herbaceous Vegetation |
| | Bluebunch Wheatgrass - Sandberg Bluegrass Herbaceous Vegetation | <i>Pseudoroegneria spicata</i> - <i>Poa secunda</i> Herbaceous Vegetation |
| | Bluebunch Wheatgrass / Tapertip Hawksbeard Herbaceous Vegetation | <i>Pseudoroegneria spicata</i> / <i>Crepis acuminata</i> Herbaceous Vegetation |
| Great Basin Wildrye | Basin Wildrye Herbaceous Vegetation | <i>Leymus cinereus</i> Herbaceous Vegetation |
| Limber Pine / Antelope Bitterbrush (Low Total Cover) | Limber Pine / Fernbush / Sandberg Bluegrass Sparse Vegetation | <i>Pinus flexilis</i> / <i>Chamaebatiaria millefolium</i> / <i>Poa secunda</i> Sparse Vegetation |
| Limber Pine / Antelope Bitterbrush (High Total Cover) | Limber Pine / Antelope Bitterbrush Woodland | <i>Pinus flexilis</i> / <i>Purshia tridentata</i> Woodland |
| Limber Pine / Antelope Bitterbrush (High Density Limber Pine) | Limber Pine / Antelope Bitterbrush Woodland | <i>Pinus flexilis</i> / <i>Purshia tridentata</i> Woodland |
| Douglas-fir / Mountain Snowberry | Douglas-fir / Mountain Snowberry Forest | <i>Pseudotsuga menziesii</i> / <i>Symphoricarpos oreophilus</i> Forest |
| Upland Quaking Aspen | Quaking Aspen - Limber Pine Forest | <i>Populus tremuloides</i> - <i>Pinus flexilis</i> Forest |
| | Quaking Aspen / Big Sagebrush Forest | <i>Populus tremuloides</i> / <i>Artemisia tridentata</i> Forest |
| | Quaking Aspen / Pinegrass Forest | <i>Populus tremuloides</i> / <i>Calamagrostis rubescens</i> Forest |

| Vegetation Type (Day and Wright 1985) | Equivalent Common Name | Equivalent Scientific Name |
|---------------------------------------|--|---|
| | Quaking Aspen / Chokecherry Forest | <i>Populus tremuloides</i> / <i>Prunus virginiana</i> Forest |
| | Quaking Aspen / Mountain Snowberry / California Brome Forest | <i>Populus tremuloides</i> / <i>Symphoricarpos oreophilus</i> / <i>Bromus carinatus</i> Forest |
| | Quaking Aspen / Mountain Snowberry / Pinegrass Forest | <i>Populus tremuloides</i> / <i>Symphoricarpos oreophilus</i> / <i>Calamagrostis rubescens</i> Forest |
| | Quaking Aspen / Mountain Snowberry Forest | <i>Populus tremuloides</i> / <i>Symphoricarpos oreophilus</i> Forest |
| | Quaking Aspen / Tall Forbs Forest | <i>Populus tremuloides</i> / Tall Forbs Forest |
| Riparian | Gray Alder / Mesic Forbs Shrubland | <i>Alnus incana</i> / Mesic Forbs Shrubland |
| | Black Cottonwood Temporarily Flooded Woodland Alliance | <i>Populus balsamifera</i> ssp. <i>trichocarpa</i> Temporarily Flooded Woodland Alliance |

Appendix 2. Summary of pertinent classification literature. Local and regional literature regarding the classification of vegetation found within Craters of the Moon National Monument and Preserve is listed alphabetically by plant association within physiognomic class. The work of Day and Wright (1985) is treated in detail in Appendix 1 and is not listed here.

| Scientific Name | Reference |
|--|--|
| Forest and Woodland | |
| <i>Juniperus osteosperma</i> / <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Hesperostipa comata</i> | Forman et al. 2003 |
| <i>Populus tremuloides</i> - <i>Pinus flexilis</i> Forest | Mueggler 1988 |
| <i>Populus tremuloides</i> / <i>Artemisia tridentata</i> Forest | Mueggler 1988 |
| <i>Populus tremuloides</i> / <i>Calamagrostis rubescens</i> Forest | Mueggler 1988 |
| <i>Populus tremuloides</i> / <i>Symphoricarpos oreophilus</i> / <i>Calamagrostis rubescens</i> Forest | Mueggler 1988 |
| <i>Populus tremuloides</i> / Tall Forbs Forest | Bowerman et al. 1996, Mueggler 1988 |
| <i>Pseudotsuga menziesii</i> / <i>Carex geyeri</i> Forest | Cooper et al. 1991, Steele et al. 1981 |
| <i>Pseudotsuga menziesii</i> / <i>Symphoricarpos oreophilus</i> Forest | Bowerman et al. 1996, Johnson and Simon 1987, Steele et al. 1981, Steele et al. 1983 |
| Shrubland | |
| <i>Alnus incana</i> / <i>Mesic Forbs</i> Shrubland | Crowe and Clausnitzer 1995, Manning and Padgett 1995, Padgett et al. 1989 |
| <i>Artemisia arbuscula</i> ssp. <i>arbuscula</i> / <i>Poa secunda</i> Shrub Herbaceous Vegetation | Hironaka et al. 1983 |
| <i>Artemisia arbuscula</i> ssp. <i>arbuscula</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation | Bowerman et al. 1996, Hironaka et al. 1983, Nelson and Jensen 1987 |
| <i>Artemisia arbuscula</i> ssp. <i>longiloba</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation | Hironaka et al. 1983, Nelson and Jensen 1987 |
| <i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Festuca idahoensis</i> Shrubland | Hironaka et al. 1983, Nelson and Jensen 1987 |
| <i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Leymus cinereus</i> Shrubland | Hironaka et al. 1983 |
| <i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Poa secunda</i> Shrubland | Daubenmire 1970 |
| <i>Artemisia tridentata</i> ssp. <i>tridentata</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation | Bowerman et al. 1996, Daubenmire 1970, Hironaka et al. 1983, Nelson and Jensen 1987 |
| <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> - <i>Symphoricarpos oreophilus</i> / <i>Pseudoroegneria spicata</i> Shrubland | Bowerman et al. 1996, Hironaka et al. 1983, Nelson and Jensen 1987 |
| <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation | Bowerman et al. 1996, Hironaka et al. 1983, Johnson and Simon 1987, Winward 1970 |

| Scientific Name | Reference |
|---|---|
| <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Hesperostipa comata</i> Shrubland | Hironaka et al. 1983, Winward 1970 |
| <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Leymus cinereus</i> Shrubland | Hironaka et al. 1983, Nelson and Jensen 1987 |
| <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> / <i>Pseudoroegneria spicata</i> Shrubland | Bowerman et al. 1996, Hironaka et al. 1983, Lauer and Peek 1976, Nelson and Jensen 1987, Winward 1970 |
| <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Achnatherum hymenoides</i> | McBride et al. 1978 |
| <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Achnatherum thurberianum</i> Shrubland | Hironaka et al. 1983 |
| <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Hesperostipa comata</i> Shrubland | Hironaka et al. 1983, Winward 1970, McBride et al. 1978 |
| <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Poa secunda</i> Shrubland | Hironaka et al. 1983 |
| <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> / <i>Pseudoroegneria spicata</i> Shrubland | Hironaka et al. 1983, McBride et al. 1978 |
| <i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Festuca idahoensis</i> Shrub Herbaceous Vegetation | Daubenmire 1970, Hironaka et al. 1983 |
| <i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Hesperostipa comata</i> Shrub Herbaceous Vegetation | Daubenmire 1970, Hironaka et al. 1983 |
| <i>Artemisia tripartita</i> ssp. <i>tripartita</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation | Daubenmire 1970, Hironaka et al. 1983 |
| <i>Purshia tridentata</i> / <i>Pseudoroegneria spicata</i> Shrub Herbaceous Vegetation | Hironaka et al. 1983, Johnson and Simon 1987 |
| Herbaceous Vegetation | |
| <i>Festuca idahoensis</i> - <i>Pseudoroegneria spicata</i> Herbaceous Vegetation | Johnson and Simon 1987, Mueggler and Harris 1969, Tisdale 1986 |
| <i>Hesperostipa comata</i> - <i>Poa secunda</i> Herbaceous Vegetation | Daubenmire 1970 |
| <i>Juncus balticus</i> Herbaceous Vegetation | Crowe and Clausnitzer 1995, Manning and Padgett 1995, Padgett et al. 1989, Tuhy and Jensen 1982, Youngblood et al. 1985 |
| <i>Leymus cinereus</i> Herbaceous Vegetation | Johnson and Simon 1987 |
| <i>Pascopyrum smithii</i> Herbaceous Vegetation | French and Mitchell 1983 |
| <i>Pseudoroegneria spicata</i> - <i>Melica bulbosa</i> Herbaceous Vegetation | Mueggler and Harris 1969, Schlatterer 1972 |
| <i>Pseudoroegneria spicata</i> - <i>Poa secunda</i> Herbaceous Vegetation | Johnson and Simon 1987 |
| <i>Pseudoroegneria spicata</i> - <i>Poa secunda</i> / <i>Balsamorhiza sagittata</i> Herbaceous Vegetation | Mueggler and Harris 1969, Tisdale 1986 |

Appendix 3. Stand tables. The species compositions of plant associations that were sampled on 4 or more plots are summarized. The stand tables are organized alphabetically by physiognomic class: Part A: Forest, Woodland and Savanna; Part B: Shrublands and Shrub-Steppe; Part C: Herbaceous Vegetation; Part D: Sparse Vegetation. Species composition data is summarized as percent frequency followed by characteristic cover (the later shown in parentheses). For the purpose of comparison, species lists are selected by physiognomic class. One exception here are the deciduous shrublands, which have a unique species list and appear at the end of Part B. In some instances blank rows will appear as the summary for a given physiognomic class may span over several pages. Please refer to Table 1 for association sample sizes.

Part A: Forest, Woodland, and Savanna

| Species | JUOS-
JUSC2/
ARTRW8/
CHM12 | JUSC2/
POSE/
PEDE4 | PIFL2/
PUTR2 | POTR5/
CARU | POTR5/
PRVI | POTR5/
SYOR2/
CARU | PSME/
SYOR2 |
|--|-------------------------------------|--------------------------|-----------------|----------------|----------------|--------------------------|----------------|
| TREES | | | | | | | |
| Juniperus osteosperma | 100 (11.3) | 50 (4.2) | 13 (0.6) | - | - | - | - |
| Juniperus scopulorum | 86 (3.7) | 100 (20.4) | - | - | - | 13 (0.1) | - |
| Pinus flexilis | - | - | 100 (9.7) | 50 (0.4) | - | 38 (0.1) | 25 (0.2) |
| Populus tremuloides | - | - | 6 (10) | 100 (19.5) | 100 (18.8) | 100 (25.3) | - |
| Pseudotsuga menziesii | - | - | 6 (10) | - | - | - | 100 (24.1) |
| DWART-SHRUBS | | | | | | | |
| Ericameria nana | 86 (2.2) | 71 (0.5) | - | - | - | - | - |
| Eriogonum microthecum | - | 14 (0.1) | 19 (0.1) | - | - | - | - |
| SHRUBS | | | | | | | |
| Amelanchier alnifolia | - | - | 25 (0.1) | 17 (0.1) | 25 (0.1) | 25 (0.1) | - |
| Artemisia tridentata ssp. vaseyana | - | - | 63 (9.5) | 50 (0.4) | 25 (0.1) | 63 (0.3) | 38 (6.4) |
| Artemisia tridentata var. wyomingensis | 100 (6.3) | 43 (3.7) | 6 (2) | - | - | - | - |
| Chamaebatiaria millefolium | 100 (2.3) | 21 (1.4) | 19 (0.1) | - | - | - | - |
| Chrysothamnus viscidiflorus | 14 (5) | 7 (0.1) | 6 (1) | - | - | 25 (0.1) | - |
| Ericameria nauseosa ssp. consimilis var. oreophila | 71 (1) | 93 (1.1) | 88 (3) | 33 (0.1) | 25 (0.1) | 25 (0.1) | - |
| Holodiscus dumosus | 43 (1) | 29 (0.1) | - | - | - | - | - |
| Prunus virginiana | - | - | - | 67 (0.8) | 100 (19) | 63 (0.7) | 50 (12.8) |
| Purshia tridentata | 86 (0.7) | 79 (1.7) | 100 (8.9) | 17 (0.1) | - | 13 (0.1) | 13 (0.1) |
| Ribes aureum | - | 36 (0.6) | 25 (0.1) | - | 25 (0.1) | 13 (0.1) | 50 (0.1) |
| Ribes cereum var. pedicellare | - | - | 63 (1) | - | - | - | - |
| Salix boothii | - | - | - | 33 (0.1) | 25 (2) | 13 (2) | 13 (0.1) |
| Salix scouleriana | - | - | - | 33 (2.6) | - | - | 25 (1) |
| Symphoricarpos oreophilus | - | - | 6 (0.1) | 100 (2.4) | 75 (3.7) | 100 (11) | 50 (4.8) |
| GRAMINOIDS | | | | | | | |
| Achnatherum hymenoides | 29 (0.6) | 36 (0.5) | 50 (0.3) | - | - | - | - |
| Achnatherum nelsonii ssp. dorei | - | - | - | 50 (0.1) | 50 (0.1) | 50 (1.6) | 25 (0.1) |
| Bromus tectorum | 86 (1.5) | 100 (1.3) | 19 (0.1) | - | 25 (0.1) | - | 13 (3) |
| Calamagrostis rubescens | - | - | - | 100 (18.5) | 25 (1) | 75 (10.3) | - |
| Elymus elymoides | 14 (0.1) | 43 (0.3) | 56 (0.2) | - | - | - | - |
| Festuca idahoensis | - | - | 19 (0.7) | - | 25 (0.1) | 13 (0.1) | - |
| Hesperostipa comata ssp. comata | 14 (0.1) | 50 (0.9) | - | - | - | - | - |
| Koeleria macrantha | - | - | - | 17 (0.1) | 25 (0.1) | 63 (0.1) | - |
| Leymus cinereus | - | 7 (0.1) | 13 (0.6) | 17 (0.1) | 50 (0.6) | 38 (0.7) | - |
| Poa secunda | 86 (5) | 100 (1.3) | 88 (0.8) | - | - | 25 (0.1) | 38 (1.7) |
| Pseudoroegneria spicata | - | 14 (1) | 13 (1.1) | 17 (0.1) | 50 (1.5) | 13 (0.1) | 63 (1.4) |
| FORBS | | | | | | | |
| Achillea millefolium | - | - | - | 50 (0.1) | 25 (0.1) | 88 (0.4) | - |
| Agastache urticifolia | - | - | - | 17 (0.1) | 50 (0.1) | 25 (0.1) | 25 (0.1) |
| Agoseris glauca | - | - | 19 (0.1) | - | 25 (0.1) | - | - |
| Angelica pinnata | - | - | - | 33 (0.1) | 50 (0.1) | 13 (0.1) | 25 (0.1) |
| Antennaria microphylla | - | - | - | 17 (0.1) | - | - | 38 (0.1) |
| Asteraceae | - | - | - | 67 (0.2) | 50 (0.2) | 63 (0.1) | 25 (0.1) |
| Balsamorhiza sagittata | - | - | - | 17 (0.1) | 25 (0.1) | 50 (0.1) | 50 (5) |
| Chaenactis douglasii | 71 (0.5) | 36 (0.5) | 6 (0.1) | - | - | - | - |
| Cirsium canovirens | 57 (0.3) | 14 (0.1) | - | - | - | - | - |
| Collinsia parviflora | - | - | 13 (0.1) | - | 25 (0.1) | 13 (0.1) | 25 (0.1) |
| Crepis acuminata | - | - | 31 (0.1) | 17 (0.1) | 25 (0.1) | 50 (0.1) | 25 (0.1) |
| Delphinium andersonii | - | - | 81 (0.2) | - | - | - | - |
| Descurainia pinnata ssp. filipes | 57 (0.8) | 7 (0.1) | - | - | - | - | - |
| Eriogonum ovalifolium | - | - | 56 (0.1) | - | - | - | - |
| Eriogonum ovalifolium var. depressum | - | - | 31 (0.1) | - | - | - | - |
| Eriogonum umbellatum | - | 7 (0.1) | 81 (0.1) | - | - | - | - |
| Galium bifolium | - | - | - | - | 50 (0.1) | 13 (0.1) | 25 (0.6) |
| Gutierrezia sarothrae | - | 36 (0.8) | - | - | - | - | - |
| Hackelia floribunda | - | - | - | 33 (0.1) | 25 (0.1) | 38 (0.1) | 25 (0.1) |
| Lactuca serriola | 71 (0.8) | 57 (0.2) | - | 17 (0.1) | - | 13 (1) | - |
| Leptodactylon pungens | 86 (1.5) | 79 (0.5) | 56 (0.2) | - | - | - | - |
| Lithospermum ruderales | - | - | 13 (0.1) | - | 25 (0.1) | 25 (0.1) | - |
| Lupinus arbustus | - | - | - | - | 25 (0.1) | 25 (0.1) | - |
| Lupinus argenteus | - | - | - | 33 (0.1) | - | 25 (0.1) | - |
| Maianthemum racemosum ssp. amplexicaule | - | - | - | 17 (0.1) | 25 (0.1) | 25 (0.1) | 25 (0.1) |
| Maianthemum stellatum | - | - | - | 33 (0.1) | 50 (0.1) | 25 (0.1) | - |
| Opuntia polyacantha | 43 (0.4) | 43 (0.3) | - | - | - | - | - |
| Penstemon deustus | 86 (2.7) | 100 (0.7) | 38 (0.1) | - | - | - | 13 (0.1) |
| Phacelia hastata | - | 7 (0.1) | 81 (0.2) | - | 25 (1) | 13 (0.1) | - |
| Potentilla glandulosa | 43 (0.1) | 64 (0.2) | 19 (1) | 50 (0.1) | 25 (0.1) | 50 (0.1) | 25 (0.1) |
| Pteryxia terebinthina var. foeniculacea | - | - | 44 (0.4) | - | - | - | 13 (0.1) |
| Senecio integerrimus var. exaltatus | - | - | 31 (0.1) | 17 (0.1) | - | 13 (0.1) | - |
| Stephanomeria minor var. myrioclada | 100 (1) | 79 (0.3) | 6 (0.1) | - | - | - | - |
| Taraxacum officinale | - | - | - | 67 (0.1) | 50 (0.1) | 38 (0.1) | 25 (0.1) |
| Tragopogon dubius | 29 (0.1) | 14 (0.1) | - | 17 (0.1) | 25 (0.1) | 13 (0.1) | - |
| Unknown species | - | - | - | 33 (0.1) | 25 (0.2) | 13 (0.1) | 25 (0.1) |
| Viola nuttallii | - | - | 6 (0.1) | - | 25 (0.1) | 50 (0.1) | - |

Part B: Shrublands and Shrub-Steppe

| Species | ARARA/
POSE | ARARA/
PSSP6 | ARTRT/
LECI4 | ARTRT/
POSE | ARTRT2/
ACTH7 | ARTRT2/
BRTE | ARTRT2/
FEID |
|--|----------------|-----------------|-----------------|----------------|------------------|-----------------|-----------------|
| TREES | | | | | | | |
| Juniperus osteosperma | - | - | - | - | - | - | - |
| Pinus flexilis | - | - | - | - | - | - | - |
| DWARF-SHRUBS | | | | | | | |
| Artemisia arbuscula | 100 (12) | 100 (7.6) | - | - | - | 3 (0.1) | - |
| Ericameria nana | - | - | - | - | - | - | - |
| Eriogonum microthecum | 25 (0.1) | - | - | 17 (0.1) | - | - | - |
| Artemisia tridentata ssp. tridentata | - | - | 100 (12.4) | 100 (13.2) | - | 3 (0.1) | - |
| Artemisia tridentata ssp. vaseyana | - | 29 (2) | - | 17 (1) | - | - | 21 (0.1) |
| Artemisia tridentata var. wyomingensis | - | - | - | - | 50 (5.9) | 58 (2) | 43 (0.7) |
| Artemisia tripartita | - | - | 40 (2) | 33 (1.5) | 100 (21.5) | 100 (10.2) | 100 (25.1) |
| SHRUBS | | | | | | | |
| Chamaebatiaria millefolium | - | - | - | - | - | - | - |
| Chrysothamnus viscidiflorus | - | 29 (1) | 60 (2.7) | 33 (0.2) | 90 (4.8) | 80 (3.3) | 86 (1.5) |
| Ericameria nauseosa ssp. consimilis var. oreophila | 13 (0.1) | - | 20 (4) | 67 (0.8) | 37 (2.9) | - | 7 (2) |
| Holodiscus dumosus | - | - | - | - | - | - | - |
| Philadelphus lewisii | - | - | - | - | - | - | - |
| Purshia tridentata | - | 43 (1) | - | 67 (6.5) | 13 (2.3) | 8 (3) | 36 (0.6) |
| Ribes aureum | - | - | - | - | - | - | - |
| Symphoricarpos oreophilus | - | - | - | - | - | - | - |
| Tetradymia canescens | - | - | 20 (2) | - | 20 (1.5) | 5 (1) | 7 (4) |
| GRAMINOIDS | | | | | | | |
| Achnatherum hymenoides | - | - | - | 17 (0.1) | 13 (1.8) | 10 (1) | 14 (0.6) |
| Achnatherum thurberianum | - | - | 20 (1) | - | 100 (10.9) | 43 (1.5) | 43 (12.2) |
| Agropyron cristatum | - | - | - | - | - | 8 (5.7) | - |
| Agropyron fragile | - | - | - | - | - | - | - |
| Bromus japonicus | 38 (2.3) | - | - | - | - | 3 (0.1) | 7 (0.1) |
| Bromus tectorum | 50 (2.5) | 57 (2.5) | 80 (7) | 83 (7) | 87 (8) | 95 (7.3) | 57 (7.9) |
| Carex | - | - | - | - | 33 (4.9) | - | 7 (2) |
| Elymus elymoides | 38 (0.4) | 14 (1) | 20 (7) | 83 (0.1) | 67 (5.8) | 30 (1.3) | 29 (0.6) |
| Elymus lanceolatus | - | - | 40 (2.1) | 33 (0.2) | 23 (3.3) | - | 7 (2) |
| Festuca idahoensis | - | - | - | 17 (0.1) | 27 (3.6) | - | 100 (15.4) |
| Hesperostipa comata ssp. comata | - | - | 20 (0.1) | 17 (0.1) | 23 (2.9) | 13 (1.4) | 7 (1) |
| Koeleria macrantha | - | - | 20 (0.1) | 17 (0.1) | 40 (2.8) | - | 57 (1) |
| Leymus cinereus | - | - | 100 (7.2) | 67 (0.1) | 43 (1.8) | 8 (1.3) | 57 (0.4) |
| Melica bulbosa | - | 14 (1) | - | - | - | - | - |
| Pascopyrum smithii | - | - | 20 (29) | - | 63 (3.3) | 20 (2.1) | 50 (2.6) |
| Poa | - | 14 (1) | 20 (12) | - | 7 (2.5) | 43 (2.7) | 7 (0.1) |
| Poa ampla | - | - | - | - | 23 (2.6) | - | - |
| Poa nevadensis | - | - | - | - | 40 (2.3) | - | 7 (3) |
| Poa secunda | 100 (4.3) | 71 (2.4) | 80 (5.3) | 100 (1.4) | 100 (8.5) | 65 (2.3) | 79 (9.9) |
| Pseudoroegneria spicata | 38 (1) | 100 (2.6) | 60 (1) | 33 (0.2) | 77 (5) | 18 (1) | 71 (1.9) |
| FORBS | | | | | | | |
| Agoseris glauca | - | - | - | 33 (0.1) | 30 (2.2) | - | 21 (0.1) |
| Allium | - | - | 40 (0.6) | 17 (0.1) | 33 (1.6) | 5 (1) | 7 (0.1) |
| Allium acuminatum | - | - | 20 (0.1) | 50 (0.1) | - | 3 (0.1) | - |
| Alyssum desertorum | - | - | 20 (0.1) | 17 (0.2) | - | - | - |
| Amsinckia | - | - | - | - | - | 13 (1.6) | - |
| Arabis | - | - | - | - | 3 (1) | - | - |
| Arabis holboellii | 25 (0.1) | - | - | 33 (0.1) | - | - | - |
| Arenaria aculeata | 25 (1.6) | 14 (2) | - | - | - | 5 (1.5) | - |
| Aster | - | - | - | - | 10 (1.3) | 3 (1) | - |
| Astragalus | 13 (0.1) | - | 20 (1) | - | 33 (1.6) | - | 7 (0.1) |
| Astragalus purshii var. glareosus | 13 (0.1) | - | - | - | - | - | 7 (0.1) |
| Balsamorhiza sagittata | - | 43 (1) | 20 (10) | 17 (0.2) | 3 (1) | - | - |
| Calochortus | - | - | - | - | 3 (1) | - | - |
| Castilleja | - | - | - | - | 7 (1) | - | 7 (0.1) |
| Castilleja pallescens var. inverta | 25 (0.6) | - | - | 17 (0.2) | - | - | - |
| Ceratocephala testiculata | - | - | - | 17 (0.1) | - | - | - |
| Chaenactis douglasii | - | - | - | 33 (0.1) | - | - | - |
| Cirsium | - | - | - | - | - | - | - |
| Cirsium canovirens | - | - | - | - | - | - | - |
| Collinsia parviflora | 25 (0.6) | - | 40 (0.1) | 50 (0.1) | - | - | 14 (0.1) |
| Crepis acuminata | 25 (0.6) | - | 60 (0.7) | 83 (0.3) | - | 5 (1) | 21 (0.1) |
| Cryptantha | 13 (2) | 29 (1) | - | - | - | 5 (1.5) | - |
| Cryptantha spiculifera | - | - | - | 17 (0.1) | - | - | - |
| Delphinium andersonii | - | - | - | 83 (0.1) | - | 3 (0.1) | 21 (0.1) |
| Descurainia pinnata ssp. filipes | - | - | - | - | - | 3 (1) | - |
| Epilobium | - | - | - | - | 7 (2) | 15 (1) | - |
| Erigeron | - | - | 20 (1) | - | 3 (2) | - | - |
| Erigeron peregrinus | - | - | - | - | - | 5 (2) | - |
| Erigeron pumilus | - | - | 20 (0.1) | 17 (0.1) | - | - | 7 (0.1) |
| Eriogonum | - | 43 (2.3) | - | - | 13 (2.5) | 3 (1) | - |
| Eriogonum caespitosum | 38 (0.2) | - | - | - | - | - | 7 (0.1) |
| Eriogonum heracleoides | 13 (0.1) | - | - | - | - | - | - |
| Eriogonum ovalifolium | 13 (0.1) | - | 20 (0.1) | 33 (0.1) | - | - | - |
| Eriogonum umbellatum | - | - | - | 17 (0.1) | - | - | 7 (0.1) |
| Fabaceae | - | - | - | - | - | 13 (1.6) | - |
| Gayophytum | - | - | 20 (1) | - | 27 (2.4) | 55 (2.5) | 7 (0.1) |
| Gutierrezia sarothrae | - | - | - | - | - | - | - |

| | | | | | | | |
|--|----------|----------|----------|----------|----------|----------|----------|
| Gymnosteris nudicaulis | - | - | - | 33 (0.1) | - | - | - |
| Gymnosteris parvula | - | - | - | 50 (0.1) | - | - | - |
| Lactuca serriola | - | - | - | - | - | 3 (0.1) | - |
| Lappula occidentalis var. occidentalis | - | - | - | - | - | - | - |
| Leptodactylon pungens | - | - | 20 (0.1) | 33 (0.1) | 3 (9) | - | - |
| Lithophragma glabrum | - | - | 20 (0.1) | - | - | - | 7 (0.1) |
| Lithophragma tenellum | - | - | - | 17 (0.1) | - | - | 14 (0.1) |
| Lithospermum ruderales | - | - | - | 33 (0.1) | 3 (2) | 5 (1) | 7 (0.1) |
| Lomatium | - | - | 20 (1) | - | 53 (2.3) | - | - |
| Lomatium dissectum | - | - | - | 17 (0.1) | - | - | - |
| Lomatium foeniculaceum ssp. macdougallii | 13 (0.1) | - | 20 (0.1) | 33 (0.1) | - | - | 7 (0.1) |
| Lomatium nudicaule | - | - | - | - | - | 15 (1.7) | - |
| Lomatium triternatum | - | - | 40 (0.1) | 67 (0.1) | - | - | 43 (0.3) |
| Lupinus | - | 14 (2) | - | - | 10 (1.3) | - | 21 (0.1) |
| Machaeranthera canescens | - | - | - | - | - | - | - |
| Nothocalais troximoides | - | - | - | 33 (0.2) | - | - | - |
| Opuntia polyacantha | 13 (0.1) | - | 20 (0.1) | 17 (0.1) | 13 (1.5) | - | 7 (0.1) |
| Orthocarpus | - | - | 20 (3) | - | 27 (2.5) | 18 (1.1) | - |
| Packera cana | - | - | 20 (0.1) | 33 (0.1) | - | - | - |
| Penstemon | - | - | - | 17 (0.1) | 3 (1) | - | - |
| Penstemon cyaneus | - | - | 40 (0.1) | - | - | 3 (0.1) | 7 (0.1) |
| Penstemon deustus | - | 14 (1) | - | - | - | 5 (0.6) | - |
| Phacelia hastata | - | - | 20 (0.1) | 33 (0.1) | - | - | - |
| Phlox | - | 29 (1) | - | - | 3 (1) | 10 (1) | - |
| Phlox hoodii | 63 (1.4) | 29 (1.5) | 20 (0.1) | 33 (0.1) | 73 (4) | 23 (1.2) | 86 (2.6) |
| Phlox longifolia | 38 (2) | 43 (3) | 40 (5.1) | 17 (0.4) | 60 (4.2) | 23 (1.1) | 7 (1) |
| Potentilla glandulosa | - | - | - | - | - | - | - |
| Pteryxia terebinthina var. foeniculacea | 13 (0.1) | - | - | 17 (0.2) | - | - | - |
| Senecio integerrimus var. exaltatus | - | - | 20 (0.1) | 50 (0.1) | - | - | 21 (0.1) |
| Sisymbrium altissimum | - | - | 20 (0.1) | - | - | 15 (1.4) | - |
| Stephanomeria minor var. myrioclada | - | - | - | - | - | - | - |
| Tragopogon dubius | - | - | 20 (0.1) | 17 (0.1) | 13 (1.3) | 8 (0.7) | 7 (0.1) |
| Unknown forb | 13 (1) | - | 20 (1) | - | 17 (1) | 8 (0.7) | 7 (1) |
| Unknown species | - | - | - | 17 (0.1) | - | - | 14 (0.6) |
| Viola nuttallii | - | - | 20 (0.1) | - | - | - | - |

Part B: Shrublands and Shrub-Steppe (continued)

| Species | ARTRT2/
HECO26 | ARTRT2/
KOMA | ARTRT2/
PASM | ARTRT2/
PASM-
BRTÉ | ARTRT2/
POSE | ARTRT2/
POSE-
BRTÉ | ARTRT2/
PSSP6 |
|--|-------------------|-----------------|-----------------|--------------------------|-----------------|--------------------------|------------------|
| TREES | | | | | | | |
| Juniperus osteosperma | - | - | - | - | - | - | - |
| Pinus flexilis | - | - | - | - | - | - | - |
| DWARF-SHRUBS | | | | | | | |
| Artemisia arbuscula | - | - | - | - | - | - | - |
| Ericameria nana | - | - | - | - | - | - | 5 (0.1) |
| Eriogonum microthecum | 25 (0.1) | - | - | - | 6 (0.1) | - | 29 (0.4) |
| Artemisia tridentata ssp. tridentata | 13 (0.1) | - | - | - | - | 2 (3) | 10 (1.1) |
| Artemisia tridentata ssp. vaseyana | 13 (0.1) | - | - | - | 6 (5) | - | 19 (0.3) |
| Artemisia tridentata var. wyomingensis | 25 (1.5) | 22 (0.6) | 5 (1) | 18 (2.5) | 47 (3.9) | 49 (3.4) | 29 (2) |
| Artemisia tripartita | 100 (13.1) | 100 (29) | 100 (14.2) | 100 (11.1) | 100 (19.1) | 100 (17.4) | 100 (16.9) |
| SHRUBS | | | | | | | |
| Chamaebatiaria millefolium | - | - | - | - | - | - | 5 (5) |
| Chrysothamnus viscidiflorus | 88 (1.6) | 89 (4.1) | 84 (3.9) | 73 (4.5) | 82 (2.3) | 71 (3.9) | 62 (3.5) |
| Ericameria nauseosa ssp. consimilis var. oreophila | 13 (0.1) | 22 (0.6) | - | - | 24 (1.3) | 14 (1.8) | 29 (1.4) |
| Holodiscus dumosus | - | - | - | - | - | - | - |
| Philadelphus lewisii | - | - | - | - | - | - | - |
| Purshia tridentata | 25 (1.1) | 11 (0.1) | - | 9 (2) | 12 (0.6) | 12 (3.3) | 33 (1.3) |
| Ribes aureum | - | - | - | - | - | - | 10 (0.1) |
| Symphoricarpos oreophilus | - | - | - | - | - | - | 5 (3) |
| Tetradymia canescens | 25 (1.5) | 22 (2) | 21 (1) | 9 (1) | - | 10 (1.8) | 24 (2.4) |
| GRAMINOIDS | | | | | | | |
| Achnatherum hymenoides | 13 (2) | - | - | - | 18 (1.7) | 7 (1.3) | 24 (0.3) |
| Achnatherum thurberianum | 38 (2.7) | 44 (2.5) | 58 (3) | 82 (2.9) | 53 (3.2) | 59 (3.6) | 14 (5) |
| Agropyron cristatum | - | - | - | 9 (2) | 6 (1) | 2 (2) | - |
| Agropyron fragile | - | - | - | - | - | 5 (6) | - |
| Bromus japonicus | - | 11 (2) | - | - | - | - | - |
| Bromus tectorum | 63 (1.8) | 100 (4.5) | 32 (2) | 100 (2.7) | 71 (4.2) | 98 (8.8) | 33 (4.5) |
| Carex | 25 (3.5) | 33 (6.3) | 11 (2.5) | - | - | 5 (4.7) | 19 (2) |
| Elymus elymoides | 50 (1) | 33 (1) | 5 (1) | 27 (1.3) | 18 (3.7) | 32 (2.8) | 24 (2.2) |
| Elymus lanceolatus | - | - | - | - | 12 (3.1) | 3 (3) | 10 (1.1) |
| Festuca idahoensis | 13 (2) | 44 (2.5) | 5 (1) | - | 24 (10) | 19 (2.9) | 52 (3.5) |
| Hesperostipa comata ssp. comata | 100 (4.3) | 44 (2) | 16 (4) | - | 12 (2) | 12 (1.1) | 24 (3) |
| Koeleria macrantha | 25 (4.6) | 100 (12.8) | 21 (12.5) | - | 12 (2) | 5 (2.3) | 29 (6.2) |
| Leymus cinereus | 13 (0.1) | 44 (5) | 5 (1) | 18 (1) | 6 (2) | 8 (1.4) | 57 (1.5) |
| Melica bulbosa | - | - | - | - | - | - | - |
| Pascopyrum smithii | 25 (1.5) | 56 (6.6) | 100 (7.8) | 100 (2.5) | 29 (1.6) | 58 (4.5) | 19 (4.5) |
| Poa | 25 (3) | - | 32 (1.3) | 73 (2.6) | 12 (2.5) | 24 (2) | 5 (1) |
| Poa ampla | - | - | - | - | 6 (1) | 3 (1) | - |
| Poa nevadensis | 13 (2) | 44 (2) | 5 (3) | - | 12 (1) | 5 (1.7) | 5 (1) |
| Poa secunda | 75 (3.9) | 100 (2.9) | 74 (1.9) | 55 (1.2) | 100 (14.2) | 100 (9.8) | 86 (4.6) |
| Pseudoroegneria spicata | 63 (1.4) | 78 (3.3) | 63 (2.8) | 64 (2) | 59 (2.2) | 41 (1.6) | 100 (13.4) |
| FORBS | | | | | | | |
| Agoseris glauca | 13 (0.1) | - | - | - | - | 8 (2.2) | 5 (2) |
| Allium | 13 (1) | 56 (4.4) | 16 (3.7) | - | 12 (1.5) | 14 (1.4) | 24 (2.4) |
| Allium acuminatum | - | - | - | - | - | - | 5 (0.1) |
| Alyssum desertorum | - | 11 (0.1) | - | - | - | - | - |
| Amsinckia | 13 (1) | - | 11 (1) | - | 6 (1) | 3 (1.5) | - |
| Arabis | - | 11 (0.1) | - | - | 6 (1) | - | 5 (1) |
| Arabis holboellii | 13 (0.1) | - | - | - | - | - | - |
| Arenaria aculeata | 13 (0.1) | - | - | - | 6 (1) | 3 (1) | - |
| Aster | - | - | - | - | - | 2 (1) | - |
| Astragalus | - | 33 (2) | 16 (3.3) | - | 24 (0.6) | 10 (1.7) | 14 (0.7) |
| Astragalus purshii var. glareosus | - | - | - | - | 6 (0.1) | - | - |
| Balsamorhiza sagittata | 13 (0.1) | 11 (1) | - | - | 6 (0.1) | - | 19 (3.3) |
| Calochortus | - | - | - | - | - | 2 (1) | - |
| Castilleja | - | 11 (0.1) | - | - | - | 2 (1) | 5 (1) |
| Castilleja pallescens var. inverta | 13 (0.1) | - | - | - | - | - | - |
| Ceratocephala testiculata | - | - | - | - | - | - | - |
| Chaenactis douglasii | 13 (0.1) | - | - | - | - | - | 5 (0.1) |
| Cirsium | - | - | - | - | - | - | - |
| Cirsium canovirens | - | - | - | - | - | - | - |
| Collinsia parviflora | 13 (0.1) | 11 (0.1) | - | - | 6 (0.1) | - | 5 (0.1) |
| Crepis acuminata | 50 (0.6) | 11 (0.1) | - | - | 12 (0.6) | 5 (1.3) | 33 (0.2) |
| Cryptantha | - | - | - | - | 6 (0.1) | - | - |
| Cryptantha spiculifera | - | - | - | - | - | - | - |
| Delphinium andersonii | 13 (0.1) | 11 (0.1) | - | - | - | - | 19 (0.1) |
| Descurainia pinnata ssp. filipes | - | - | - | - | - | - | 5 (0.1) |
| Epilobium | - | - | 5 (1) | - | 6 (1) | 2 (1) | - |
| Erigeron | - | - | - | - | 6 (2) | 2 (1) | - |
| Erigeron peregrinus | 13 (1) | - | 5 (1) | - | - | 3 (1.5) | - |
| Erigeron pumilus | - | - | - | - | - | - | 10 (0.6) |
| Eriogonum | 13 (1) | 33 (1.3) | 16 (2) | - | 12 (1.5) | 5 (1) | 10 (1.5) |
| Eriogonum caespitosum | - | - | - | - | - | - | - |
| Eriogonum heracleoides | - | - | - | - | - | - | - |
| Eriogonum ovalifolium | 13 (0.1) | 11 (0.1) | - | - | 6 (0.1) | - | 33 (0.1) |
| Eriogonum umbellatum | 13 (0.1) | - | - | - | 6 (0.1) | - | 19 (0.1) |
| Fabaceae | - | - | 5 (1) | - | - | 3 (1) | 5 (0.1) |
| Gayophytum | 38 (2.3) | - | 58 (2.8) | 55 (1.8) | 47 (2.5) | 36 (2.7) | 10 (2) |

| | | | | | | | |
|--|----------|----------|----------|----------|----------|----------|----------|
| Gutierrezia sarothrae | - | - | - | - | - | - | - |
| Gymnosteris nudicaulis | 13 (0.1) | 11 (0.1) | - | - | 6 (0.1) | - | - |
| Gymnosteris parvula | 13 (0.1) | 11 (0.1) | - | - | 6 (0.1) | - | - |
| Lactuca serriola | - | - | - | - | - | - | 5 (0.1) |
| Lappula occidentalis var. occidentalis | - | - | - | - | - | - | - |
| Leptodactylon pungens | 25 (0.6) | 33 (2) | 5 (3) | - | 6 (1) | 2 (3) | 52 (1.9) |
| Lithophragma glabrum | - | - | - | - | - | - | - |
| Lithophragma tenellum | - | 11 (0.1) | - | - | - | - | - |
| Lithospermum ruderales | 13 (1) | 22 (0.1) | - | - | - | 2 (1) | 5 (0.1) |
| Lomatium | - | 67 (1.5) | 11 (2) | - | 18 (2.7) | 14 (2) | 19 (3.8) |
| Lomatium dissectum | - | - | - | - | 6 (0.1) | - | - |
| Lomatium foeniculaceum ssp. macdougallii | - | - | - | - | 6 (0.1) | - | 24 (0.1) |
| Lomatium nudicaule | 13 (1) | - | - | - | 6 (1) | - | - |
| Lomatium triternatum | 13 (0.1) | 11 (0.1) | - | - | 6 (0.1) | - | 5 (0.1) |
| Lupinus | 13 (0.1) | 22 (1) | - | - | 6 (0.1) | 3 (1) | 14 (0.7) |
| Machaeranthera canescens | - | - | - | - | - | 2 (1) | - |
| Nothocalais troximoides | 13 (0.1) | - | - | - | - | - | - |
| Opuntia polyacantha | - | - | - | - | 6 (0.1) | - | 19 (0.1) |
| Orthocarpus | 13 (2) | - | 47 (3.2) | 55 (2) | 29 (1.6) | 22 (1.8) | 10 (5.5) |
| Packera cana | - | - | - | - | - | - | 10 (0.6) |
| Penstemon | - | - | - | - | 6 (1) | 3 (1) | 14 (0.1) |
| Penstemon cyaneus | 13 (0.1) | 22 (0.6) | - | - | 6 (0.1) | - | 10 (0.1) |
| Penstemon deustus | - | - | - | - | - | - | 5 (0.1) |
| Phacelia hastata | 13 (0.1) | - | - | - | 6 (0.1) | - | 14 (0.1) |
| Phlox | 13 (3) | 11 (2) | 26 (2.4) | - | - | 5 (1) | 5 (3) |
| Phlox hoodii | 63 (0.8) | 78 (5.1) | 47 (2.7) | 45 (1.8) | 59 (2.9) | 54 (2.5) | 52 (1.9) |
| Phlox longifolia | 25 (0.6) | 56 (3.6) | 26 (3.6) | 9 (3) | 47 (2.1) | 39 (2.9) | 33 (3.3) |
| Potentilla glandulosa | - | - | - | - | - | - | 5 (1) |
| Pteryxia terebinthina var. foeniculacea | 13 (0.1) | - | - | - | 6 (0.1) | - | 5 (1) |
| Senecio integerrimus var. exaltatus | 13 (0.1) | - | - | - | - | - | 14 (0.1) |
| Sisymbrium altissimum | 13 (2) | - | 26 (1) | 27 (1) | - | 8 (2.4) | 5 (1) |
| Stephanomeria minor var. myrioclada | - | - | - | - | - | - | - |
| Tragopogon dubius | 13 (0.1) | - | 5 (1) | - | - | 5 (1) | 10 (0.6) |
| Unknown forb | 13 (1) | 11 (1) | - | - | 6 (2) | 14 (1.1) | 5 (2) |
| Unknown species | 13 (0.1) | - | - | - | 6 (0.1) | - | 10 (0.1) |
| Viola nuttallii | - | - | - | - | - | - | 14 (0.1) |

Part B: Shrublands and Shrub-Steppe (continued)

| Species | ARTRV-
CHMI2/
PEDE4 | ARTRV-
PUTR2/
POSE | ARTRV-
PUTR2/
PSSP6 | ARTRV/
BRTE | ARTRV/
FEID | ARTRV/
HECO26 | ARTRV/
POSE | ARTRV/
PSSP6 |
|--|---------------------------|--------------------------|---------------------------|----------------|----------------|------------------|----------------|-----------------|
| TREES | | | | | | | | |
| Juniperus osteosperma | - | - | - | - | - | - | - | - |
| Pinus flexilis | 40 (2.5) | - | - | 4 (0.1) | - | - | 7 (0.1) | - |
| DWARF-SHRUBS | | | | | | | | |
| Artemisia arbuscula | - | - | - | 17 (4.3) | - | - | 20 (4) | 9 (2.7) |
| Ericameria nana | 60 (1) | 20 (2) | 20 (0.1) | 8 (0.1) | - | 25 (0.1) | 33 (0.7) | 3 (0.1) |
| Eriogonum microthecum | - | 40 (0.6) | 40 (0.6) | 13 (0.1) | - | 50 (0.1) | 47 (0.1) | 3 (0.1) |
| Artemisia tridentata ssp. tridentata | 20 (1) | - | - | - | - | - | - | - |
| Artemisia tridentata ssp. vaseyana | 100 (10) | 100 (20) | 100 (18) | 100 (12.1) | 100 (11.8) | 100 (13.8) | 100 (12.6) | 100 (11.3) |
| Artemisia tridentata var. wyomingensis | - | - | - | - | 11 (2) | - | - | - |
| Artemisia tripartita | 20 (1) | - | - | 4 (6) | 11 (5) | - | - | 3 (5) |
| SHRUBS | | | | | | | | |
| Chamaebatiaria millefolium | 100 (4.8) | 20 (0.1) | 20 (0.1) | 17 (0.1) | 11 (0.1) | 75 (0.1) | 20 (0.1) | 6 (3) |
| Chrysothamnus viscidiflorus | - | 40 (1.1) | - | 25 (1.7) | 33 (2.3) | - | 13 (1.5) | 38 (1.1) |
| Ericameria nauseosa ssp. consimilis var. oreophila | 40 (1) | 40 (0.6) | 60 (0.4) | 21 (0.1) | - | 50 (0.6) | 53 (0.2) | 13 (0.3) |
| Holodiscus dumosus | - | - | - | - | - | - | - | - |
| Philadelphus lewisii | 20 (0.1) | - | - | - | - | - | - | - |
| Purshia tridentata | 100 (1.2) | 100 (8) | 100 (12) | 50 (3.2) | 22 (2.1) | 100 (1.3) | 60 (1.6) | 38 (0.8) |
| Ribes aureum | - | 20 (1) | 20 (0.1) | - | - | 25 (0.1) | - | - |
| Symphoricarpos oreophilus | - | 20 (0.1) | 20 (1) | 4 (2) | 22 (1) | - | - | 19 (1.2) |
| Tetradymia canescens | - | - | - | - | - | - | - | 3 (0.1) |
| GRAMINOIDS | | | | | | | | |
| Achnatherum hymenoides | 40 (0.6) | - | - | - | - | - | 13 (0.1) | 3 (1) |
| Achnatherum thurberianum | - | - | - | - | - | - | - | - |
| Agropyron cristatum | 20 (0.1) | - | - | - | - | - | - | 3 (1) |
| Agropyron fragile | - | - | - | - | - | - | - | - |
| Bromus japonicus | - | - | - | 25 (2.2) | - | - | 13 (0.6) | 9 (1.3) |
| Bromus tectorum | 40 (0.1) | 40 (0.6) | 40 (0.6) | 96 (7.8) | 11 (2) | 100 (0.3) | 33 (1.4) | 41 (3.4) |
| Carex | 40 (0.1) | - | - | - | - | - | - | 3 (0.1) |
| Elymus elymoides | 80 (0.1) | 40 (0.1) | 20 (0.1) | 25 (0.1) | 11 (0.1) | 75 (0.1) | 33 (0.1) | 16 (1.2) |
| Elymus lanceolatus | - | 40 (1.5) | 20 (0.1) | - | - | - | - | 3 (0.1) |
| Festuca idahoensis | 40 (1.1) | 20 (0.1) | 20 (1) | 4 (1) | 100 (2) | - | 7 (0.1) | 22 (1.7) |
| Hesperostipa comata ssp. comata | - | 20 (1) | 60 (0.4) | 21 (0.5) | - | 100 (1.3) | 7 (0.1) | - |
| Koeleria macrantha | - | 20 (0.1) | - | - | 11 (1) | - | - | 3 (1) |
| Leymus cinereus | 20 (0.1) | - | 20 (0.1) | 8 (0.6) | 22 (1.5) | - | 13 (0.6) | 22 (1) |
| Melica bulbosa | - | - | - | 17 (1.5) | 22 (1) | - | 13 (2) | 16 (2) |
| Pascopyrum smithii | - | - | - | - | - | - | - | - |
| Poa | - | - | 20 (2) | 8 (3) | 22 (1) | - | - | 6 (1) |
| Poa ampla | - | - | - | - | - | - | - | - |
| Poa nevadensis | - | - | - | - | - | - | - | - |
| Poa secunda | 100 (0.6) | 100 (2.8) | 60 (1.8) | 58 (2) | 56 (1.6) | 100 (2) | 93 (3.7) | 47 (3.2) |
| Pseudoroegneria spicata | 40 (0.6) | 40 (0.1) | 100 (2) | 33 (2.4) | 78 (1.6) | - | 27 (1.3) | 100 (5.8) |
| FORBS | | | | | | | | |
| Agoseris glauca | 20 (0.1) | - | - | 8 (0.1) | 11 (0.1) | - | 7 (0.1) | 3 (1) |
| Allium | - | - | - | 13 (0.1) | - | 25 (0.1) | 7 (0.1) | - |
| Allium acuminatum | - | 20 (0.1) | - | - | 11 (0.1) | - | 7 (0.1) | 3 (0.1) |
| Alyssum desertorum | - | - | - | - | - | - | - | - |
| Amsinckia | - | - | - | - | - | - | - | - |
| Arabis | - | - | - | - | - | - | - | 6 (1) |
| Arabis holboellii | - | - | - | 4 (0.1) | 11 (0.1) | 25 (0.1) | 7 (0.1) | 6 (0.1) |
| Arenaria aculeata | - | - | - | 4 (1) | 11 (1) | - | 7 (1) | 6 (1) |
| Aster | - | - | - | - | - | - | - | - |
| Astragalus | - | - | - | 8 (0.6) | - | - | - | 3 (0.1) |
| Astragalus purshii var. glareosus | - | - | - | - | - | - | - | 3 (0.1) |
| Balsamorhiza sagittata | - | - | - | 25 (2.7) | 33 (0.7) | - | 13 (2.5) | 44 (2.2) |
| Calochortus | - | - | - | 8 (0.6) | - | - | - | 3 (1) |
| Castilleja | - | - | - | - | - | - | - | - |
| Castilleja pallescens var. inverta | 20 (0.1) | - | - | - | - | 25 (0.1) | - | - |
| Ceratocephala testiculata | - | - | - | 4 (3) | - | - | - | 3 (1) |
| Chaenactis douglasii | 40 (0.1) | 20 (0.1) | 40 (0.1) | 4 (0.1) | - | - | - | - |
| Cirsium | 20 (0.1) | 20 (0.1) | 20 (0.1) | 13 (0.1) | 11 (0.1) | 50 (1.6) | - | - |
| Cirsium canovirens | 60 (0.1) | - | - | 8 (0.1) | - | - | 27 (0.1) | 3 (0.1) |
| Collinsia parviflora | - | 20 (0.1) | - | 13 (0.7) | 22 (1) | - | 13 (0.6) | 19 (1.2) |
| Crepis acuminata | - | 40 (0.1) | 60 (0.1) | - | 22 (1.5) | 25 (0.1) | 20 (0.1) | 25 (0.8) |
| Cryptantha | - | 20 (0.1) | 20 (0.1) | 8 (1) | - | - | - | 9 (0.4) |
| Cryptantha spiculifera | - | - | - | - | - | - | - | - |
| Delphinium andersonii | 40 (0.1) | 40 (0.1) | 40 (0.1) | 4 (0.1) | - | - | 27 (0.1) | 6 (0.1) |
| Descurainia pinnata ssp. filipes | - | - | - | - | 11 (1) | - | - | - |
| Epilobium | - | - | - | 13 (1.3) | - | - | - | - |
| Erigeron | - | - | - | 4 (0.1) | - | - | 13 (0.1) | - |
| Erigeron peregrinus | - | - | - | 4 (1) | - | - | - | - |
| Erigeron pumilus | - | - | - | 4 (0.1) | - | 25 (0.1) | 13 (0.1) | - |
| Eriogonum | - | 20 (0.1) | - | 25 (1) | 44 (6.3) | - | 33 (2.4) | 34 (1.8) |
| Eriogonum caespitosum | 20 (2) | - | 20 (1) | - | - | 25 (0.1) | - | 3 (0.1) |
| Eriogonum heracleoides | - | 40 (0.1) | 80 (1.3) | - | - | - | 7 (0.1) | 6 (1.1) |
| Eriogonum ovalifolium | 40 (0.1) | 60 (0.1) | 20 (0.1) | 8 (0.1) | - | 50 (0.1) | 40 (0.3) | 3 (0.2) |
| Eriogonum umbellatum | 40 (0.1) | 80 (0.1) | - | 4 (0.1) | - | - | 27 (0.1) | 9 (0.1) |
| Fabaceae | - | - | - | - | - | - | 7 (0.1) | - |
| Gayophytum | - | - | - | - | - | - | - | - |

| | | | | | | | | |
|--|-----------|----------|----------|----------|----------|----------|----------|----------|
| Gutierrezia sarothrae | - | - | - | - | - | - | - | - |
| Gymnosteris nudicaulis | - | 20 (0.1) | - | - | - | - | - | 3 (0.1) |
| Gymnosteris parvula | - | 20 (0.1) | - | - | - | - | - | 3 (0.1) |
| Lactuca serriola | - | - | - | - | - | 25 (0.1) | 7 (0.1) | 3 (1) |
| Lappula occidentalis var. occidentalis | - | - | - | 4 (0.1) | - | - | 7 (0.1) | 3 (0.1) |
| Leptodactylon pungens | 40 (0.1) | 60 (0.1) | 40 (2.1) | 17 (0.3) | 11 (1) | 75 (1) | 53 (0.5) | 6 (0.1) |
| Lithophragma glabrum | - | 20 (0.1) | 20 (0.1) | - | - | - | 13 (0.1) | - |
| Lithophragma tenellum | 60 (0.1) | 20 (0.1) | 20 (0.1) | 8 (0.1) | 11 (0.1) | 25 (0.1) | 13 (0.1) | - |
| Lithospermum ruderales | - | - | - | 8 (1) | - | - | 13 (0.6) | 9 (0.1) |
| Lomatium | - | - | - | 4 (0.1) | - | - | - | - |
| Lomatium dissectum | - | - | 20 (0.1) | 4 (0.1) | - | 25 (2) | 7 (0.1) | - |
| Lomatium foeniculaceum ssp. macdougallii | 20 (0.1) | - | - | 4 (0.1) | - | - | 13 (0.1) | 3 (0.1) |
| Lomatium nudicaule | - | - | - | 4 (1) | - | - | - | 3 (3) |
| Lomatium triternatum | - | 20 (0.1) | 40 (0.1) | 8 (0.1) | 11 (0.1) | - | 13 (0.1) | - |
| Lupinus | - | - | - | 21 (1.8) | 67 (4.9) | - | 20 (3.3) | 50 (3.1) |
| Machaeranthera canescens | - | - | - | - | - | - | - | - |
| Nothocalais troximoides | - | 20 (0.1) | - | 13 (0.1) | - | 50 (0.1) | - | 3 (0.1) |
| Opuntia polyacantha | 40 (0.1) | 20 (0.1) | 20 (0.1) | - | - | - | 27 (0.1) | - |
| Orthocarpus | - | - | - | - | - | - | - | - |
| Packera cana | - | 20 (0.1) | 20 (0.1) | - | 11 (0.1) | - | 13 (0.1) | - |
| Penstemon | - | - | 40 (0.1) | - | - | - | - | 3 (0.1) |
| Penstemon cyaneus | 20 (0.1) | - | 40 (0.1) | 8 (0.1) | - | - | 7 (0.1) | 3 (0.1) |
| Penstemon deustus | 100 (0.1) | 60 (0.4) | 80 (0.1) | 13 (0.1) | 44 (0.8) | 75 (0.1) | 47 (0.2) | 6 (1) |
| Phacelia hastata | 20 (0.1) | 20 (0.1) | 60 (0.1) | 4 (1) | - | 25 (0.1) | 20 (0.1) | 13 (0.1) |
| Phlox | - | - | - | 8 (1) | 11 (3) | - | - | 3 (1) |
| Phlox hoodii | - | 20 (0.1) | - | 8 (0.6) | 11 (3) | - | 27 (1.8) | 22 (2.4) |
| Phlox longifolia | - | 20 (0.1) | - | 13 (1) | 22 (5) | - | 7 (3) | 28 (1.8) |
| Potentilla glandulosa | 60 (0.1) | 20 (0.1) | 20 (0.1) | - | 11 (0.1) | - | 27 (0.1) | 3 (0.1) |
| Pteryxia terebinthina var. foeniculacea | 60 (0.7) | 20 (5) | - | 4 (0.1) | 11 (0.1) | - | 13 (0.1) | - |
| Senecio integerrimus var. exaltatus | 20 (0.1) | 20 (0.1) | 20 (0.1) | - | 11 (0.1) | - | 13 (0.1) | 9 (0.1) |
| Sisymbrium altissimum | - | - | 20 (0.1) | 4 (0.1) | 11 (0.1) | - | - | - |
| Stephanomeria minor var. myrioclada | - | - | - | - | - | - | 7 (0.1) | - |
| Tragopogon dubius | - | - | - | 8 (0.1) | 11 (0.1) | - | - | 3 (1) |
| Unknown forb | 20 (0.1) | - | - | 4 (0.1) | - | - | 7 (1) | 6 (1) |
| Unknown species | 20 (0.1) | - | - | - | - | - | 13 (0.6) | 9 (0.2) |
| Viola nuttallii | 40 (0.1) | 40 (0.1) | - | - | - | - | 7 (0.1) | 3 (0.1) |

Part B: Shrublands and Shrub-Steppe (continued)

| Species | ARTRW8/
ACHY | ARTRW8/
ACTH7 | ARTRW8/
ACTH7-
BRTE | ARTRW8/
AGCR | ARTRW8/
BRTE | ARTRW8/
FEID | ARTRW8/
HECO26 |
|--|-----------------|------------------|---------------------------|-----------------|-----------------|-----------------|-------------------|
| TREES | | | | | | | |
| Juniperus osteosperma | - | - | - | - | 1 (0.1) | 9 (2) | - |
| Pinus flexilis | - | - | - | - | - | - | - |
| DWARF-SHRUBS | | | | | | | |
| Artemisia arbuscula | - | - | - | - | 1 (0.1) | - | - |
| Ericameria nana | - | - | - | - | 9 (0.1) | 18 (0.6) | 12 (1) |
| Eriogonum microthecum | 20 (0.1) | - | - | - | 4 (0.1) | 36 (0.1) | 12 (0.6) |
| Artemisia tridentata ssp. tridentata | - | - | - | - | 1 (0.1) | - | - |
| Artemisia tridentata ssp. vaseyana | - | - | - | - | 6 (4.8) | 9 (5) | - |
| Artemisia tridentata var. wyomingensis | 100 (12.2) | 100 (8.4) | 100 (18.5) | 100 (14.4) | 99 (9.8) | 100 (14.5) | 100 (14.2) |
| Artemisia tripartita | 60 (3.3) | 31 (1.3) | 61 (5.5) | 40 (10) | 19 (3.2) | 18 (3.5) | 27 (3.9) |
| SHRUBS | | | | | | | |
| Chamaebatiaria millefolium | - | - | - | - | 4 (0.1) | 18 (0.1) | 3 (0.1) |
| Chrysothamnus viscidiflorus | 60 (0.7) | 77 (3.4) | 89 (5.9) | 60 (2.7) | 59 (2.4) | 27 (1.7) | 61 (3.8) |
| Ericameria nauseosa ssp. consimilis var. oreophila | - | - | 89 (4.2) | 80 (5.8) | 14 (0.6) | 27 (0.1) | 48 (1.7) |
| Holodiscus dumosus | - | - | - | - | - | - | 3 (1) |
| Philadelphus lewisii | - | - | - | - | - | - | - |
| Purshia tridentata | 40 (0.6) | 38 (2.2) | 25 (6.6) | - | 26 (0.7) | 55 (0.6) | 36 (1) |
| Ribes aureum | - | - | - | - | - | - | - |
| Symphoricarpos oreophilus | 20 (4) | - | - | - | - | - | - |
| Tetradymia canescens | - | 23 (2.3) | 14 (1.3) | - | - | - | 9 (1) |
| GRAMINOIDS | | | | | | | |
| Achnatherum hymenoides | 100 (1) | 8 (1) | 32 (1.4) | - | 10 (0.6) | 9 (0.1) | 15 (1.2) |
| Achnatherum thurberianum | 20 (2) | 100 (5.4) | 100 (9.2) | 60 (3.7) | 16 (1.5) | 9 (1) | 21 (3.4) |
| Agropyron cristatum | - | - | - | 100 (8.6) | 3 (5.5) | - | 3 (3) |
| Agropyron fragile | - | - | 4 (1) | 80 (16.5) | - | - | 6 (2) |
| Bromus japonicus | - | 8 (1) | - | - | 4 (2.7) | - | 3 (0.1) |
| Bromus tectorum | 60 (4) | 92 (4) | 100 (17.2) | 100 (10.2) | 99 (9.4) | 36 (2.8) | 91 (9.7) |
| Carex | - | 8 (1) | 11 (1.7) | - | 1 (1) | 9 (0.1) | - |
| Elymus elymoides | 40 (0.1) | 23 (1.3) | 96 (7.2) | 80 (3.3) | 41 (0.4) | 45 (2.1) | 64 (1.9) |
| Elymus lanceolatus | - | - | 36 (4) | 80 (2.8) | 1 (0.1) | - | 9 (8.7) |
| Festuca idahoensis | - | 8 (1) | 4 (1) | - | 6 (0.6) | 100 (5.6) | 9 (1.3) |
| Hesperostipa comata ssp. comata | 20 (0.1) | - | 29 (2.4) | 60 (6.7) | 19 (0.4) | - | 100 (5.2) |
| Koeleria macrantha | - | - | 7 (1.5) | - | 1 (0.1) | - | - |
| Leymus cinereus | - | 31 (1) | 57 (2.3) | - | 6 (0.3) | - | 12 (1.5) |
| Melica bulbosa | - | - | - | - | - | - | - |
| Pascopyrum smithii | - | 31 (1.5) | 71 (6.7) | 20 (2) | 10 (2.1) | - | 33 (3.8) |
| Poa | 20 (4) | 23 (1.3) | - | - | 13 (1.9) | 18 (0.6) | - |
| Poa ampla | - | - | 25 (1.1) | 20 (1) | - | - | - |
| Poa nevadensis | - | - | 29 (1.6) | 20 (1) | - | - | 12 (1.8) |
| Poa secunda | 80 (4.3) | 92 (4.1) | 100 (18.1) | 100 (10.4) | 81 (1.6) | 73 (5.9) | 94 (8.6) |
| Pseudoroegneria spicata | 20 (0.1) | 15 (1.5) | 61 (2.4) | 20 (1) | 14 (0.9) | 45 (1.8) | 39 (4) |
| FORBS | | | | | | | |
| Agoseris glauca | 20 (0.1) | - | 50 (1.6) | 20 (1) | 1 (0.1) | 9 (0.1) | 3 (1) |
| Allium | 60 (0.7) | 8 (2) | 54 (1.9) | 40 (1) | 9 (0.7) | 18 (2.1) | 15 (0.6) |
| Allium acuminatum | - | - | - | - | 11 (0.1) | 9 (0.1) | 9 (0.7) |
| Alyssum desertorum | - | - | - | - | 10 (1.5) | - | 3 (5) |
| Amsinckia | - | - | - | - | 3 (1) | - | - |
| Arabis | - | - | 14 (1) | - | - | - | 6 (1) |
| Arabis holboellii | - | - | - | - | 3 (0.1) | - | - |
| Arenaria aculeata | - | - | - | - | 1 (1) | 9 (2) | 6 (1) |
| Aster | - | - | 11 (1) | - | 1 (1) | 9 (0.1) | - |
| Astragalus | - | - | 54 (2.1) | 40 (2.5) | - | 27 (0.7) | 9 (1.7) |
| Astragalus purshii var. glareosus | 20 (0.1) | - | - | - | 3 (0.1) | 27 (0.1) | - |
| Balsamorhiza sagittata | 20 (2) | - | 14 (1.5) | - | 9 (1.2) | 18 (0.6) | 12 (5) |
| Calochortus | - | - | 4 (1) | - | 3 (0.1) | - | 3 (0.1) |
| Castilleja | - | - | 14 (1) | - | - | 18 (0.1) | 3 (0.1) |
| Castilleja pallescens var. inverta | 40 (0.1) | - | - | - | 1 (0.1) | 9 (0.1) | 3 (0.1) |
| Ceratocephala testiculata | - | - | - | - | 10 (0.7) | 9 (0.1) | - |
| Chaenactis douglasii | - | - | - | - | 10 (0.1) | - | 9 (0.1) |
| Cirsium | - | - | - | - | 1 (0.1) | - | 3 (0.1) |
| Cirsium canovirens | - | - | - | - | - | - | - |
| Collinsia parviflora | - | 8 (1) | - | - | 10 (0.9) | 27 (0.1) | 12 (1.3) |
| Crepis acuminata | 40 (0.1) | - | 11 (1.3) | - | 30 (0.6) | 45 (1.7) | 30 (0.8) |
| Cryptantha | - | - | - | - | - | - | - |
| Cryptantha spiculifera | - | - | - | - | 1 (0.1) | - | - |
| Delphinium andersonii | - | - | - | - | 10 (0.1) | 55 (0.3) | 9 (0.1) |
| Descurainia pinnata ssp. filipes | - | - | - | - | 3 (0.1) | - | - |
| Epilobium | - | - | - | - | 7 (1.4) | - | - |
| Erigeron | - | - | 25 (1.4) | 20 (1) | - | - | - |
| Erigeron peregrinus | 20 (1) | - | - | - | 4 (1) | - | 3 (1) |
| Erigeron pumilus | 20 (0.1) | - | - | - | 6 (0.1) | - | 6 (0.1) |
| Eriogonum | 20 (1) | 8 (1) | 18 (1.6) | 20 (1) | 1 (2) | 27 (3) | 9 (2) |
| Eriogonum caespitosum | - | - | - | - | - | 18 (0.1) | - |
| Eriogonum heracleoides | - | - | - | - | - | - | - |
| Eriogonum ovalifolium | 20 (0.1) | - | - | - | 4 (0.1) | 27 (0.1) | 6 (0.1) |
| Eriogonum umbellatum | - | - | - | - | 1 (0.1) | - | 9 (0.1) |
| Fabaceae | - | - | - | - | - | - | - |
| Gayophytum | 20 (6) | 31 (1.8) | - | - | 30 (1.8) | - | 12 (2.8) |

| | | | | | | | |
|--|----------|----------|----------|----------|----------|----------|----------|
| Gutierrezia sarothrae | - | - | - | - | 1 (0.1) | - | - |
| Gymnosteris nudicaulis | - | - | - | - | 4 (0.1) | 9 (0.1) | 3 (0.1) |
| Gymnosteris parvula | - | - | - | - | 4 (0.1) | 9 (0.1) | 3 (0.1) |
| Lactuca serriola | - | 8 (1) | - | - | 10 (0.4) | - | 3 (0.1) |
| Lappula occidentalis var. occidentalis | 20 (0.1) | - | - | - | 3 (0.1) | - | - |
| Leptodactylon pungens | 40 (2) | - | 14 (1.5) | - | 17 (0.8) | 45 (0.3) | 45 (0.8) |
| Lithophragma glabrum | 20 (0.1) | - | - | - | 6 (0.1) | - | - |
| Lithophragma tenellum | 20 (0.1) | - | - | - | 4 (0.1) | 9 (0.1) | 3 (0.1) |
| Lithospermum ruderales | - | - | 4 (2) | - | 9 (0.3) | - | - |
| Lomatium | - | - | 79 (3.5) | 80 (3.8) | - | 9 (15) | 12 (4) |
| Lomatium dissectum | - | - | - | - | 6 (2.8) | 9 (2) | 9 (3) |
| Lomatium foeniculaceum ssp. macdougallii | - | - | - | - | 3 (0.1) | 36 (0.1) | 9 (0.1) |
| Lomatium nudicaule | 20 (1) | 23 (1.3) | - | 20 (4) | 3 (1) | - | 3 (1) |
| Lomatium triternatum | 20 (0.1) | - | - | - | 10 (0.4) | 27 (0.1) | 9 (0.1) |
| Lupinus | - | - | 4 (1) | - | 1 (2) | 18 (4) | 3 (2) |
| Machaeranthera canescens | - | - | - | - | 6 (1) | - | 6 (1.5) |
| Nothocalais troximoides | - | - | - | - | 4 (0.1) | - | 12 (0.1) |
| Opuntia polyacantha | - | 8 (1) | 14 (1.5) | 20 (1) | 9 (0.1) | 18 (0.1) | 27 (0.4) |
| Orthocarpus | - | 38 (2.8) | - | - | 3 (1) | - | - |
| Packera cana | 20 (0.1) | - | - | - | 3 (0.1) | 9 (0.1) | 3 (0.1) |
| Penstemon | - | - | 7 (1) | - | - | 27 (0.4) | - |
| Penstemon cyaneus | - | - | - | - | 10 (0.1) | 9 (0.1) | 9 (0.4) |
| Penstemon deustus | - | 8 (1) | - | - | 6 (0.1) | 18 (0.1) | 15 (0.6) |
| Phacelia hastata | - | - | - | - | 3 (0.1) | 9 (0.1) | 3 (0.1) |
| Phlox | 20 (8) | - | - | - | - | 18 (1.5) | - |
| Phlox hoodii | 40 (0.6) | 15 (2) | 64 (3.9) | - | 26 (1) | 64 (0.8) | 30 (1.1) |
| Phlox longifolia | - | 23 (1) | 96 (4.9) | 80 (5.8) | 26 (0.9) | 18 (5.6) | 48 (4.1) |
| Potentilla glandulosa | - | - | - | - | 1 (0.1) | 9 (0.1) | 3 (1) |
| Pteryxia terebinthina var. foeniculacea | - | - | - | - | 4 (0.1) | 9 (0.1) | 3 (0.1) |
| Senecio integerrimus var. exaltatus | - | - | - | - | 4 (0.1) | 45 (0.3) | - |
| Sisymbrium altissimum | - | 8 (1) | - | - | 14 (0.5) | - | 3 (1) |
| Stephanomeria minor var. myrioclada | - | - | - | - | - | - | 12 (0.1) |
| Tragopogon dubius | 20 (0.1) | - | 50 (1.3) | 20 (1) | 13 (0.2) | 18 (0.1) | 18 (0.4) |
| Unknown forb | - | 8 (1) | 54 (1.7) | 20 (2) | 14 (0.7) | 9 (7) | 21 (1.4) |
| Unknown species | - | - | - | - | 7 (1.2) | 27 (0.1) | - |
| Viola nuttallii | - | - | - | - | - | 9 (0.1) | - |

Part B: Shrublands and Shrub-Steppe (continued)

| Species | ARTRW8/
PASM | ARTRW8/
POSE | ARTRW8/
POSE-
BRITE | ARTRW8/
PSSP6 | CHV18/
AGCR | CHV18/
BRITE | CHV18/
POSE-
BRITE |
|--|-----------------|-----------------|---------------------------|------------------|----------------|-----------------|--------------------------|
| TREES | | | | | | | |
| Juniperus osteosperma | - | 5 (2.8) | - | 7 (0.1) | - | - | - |
| Pinus flexilis | - | 5 (1.8) | - | 7 (0.1) | - | - | - |
| DWARF-SHRUBS | | | | | | | |
| Artemisia arbuscula | - | - | 1 (1) | - | - | - | - |
| Ericameria nana | - | 16 (0.5) | - | - | - | - | - |
| Eriogonum microthecum | - | 11 (0.2) | 1 (0.1) | 13 (0.1) | - | - | - |
| Artemisia tridentata ssp. tridentata | - | - | - | - | - | - | - |
| Artemisia tridentata ssp. vaseyana | - | 5 (0.6) | 1 (2) | - | 13 (0.1) | - | - |
| Artemisia tridentata var. wyomingensis | 100 (8.4) | 100 (10.8) | 100 (11.2) | 100 (15.7) | 63 (1.2) | 79 (2.1) | 40 (1.1) |
| Artemisia tripartita | 44 (2) | 17 (2.1) | 41 (3) | 20 (1.7) | 25 (2.5) | 18 (1.3) | 40 (1.5) |
| SHRUBS | | | | | | | |
| Chamaebatiaria millefolium | - | 13 (0.2) | 1 (1) | - | - | - | - |
| Chrysothamnus viscidiflorus | 78 (2.1) | 43 (1.8) | 87 (4) | 53 (3.3) | 100 (9.8) | 100 (8.3) | 100 (8.8) |
| Ericameria nauseosa ssp. consimilis var. oreophila | 22 (4.1) | 24 (0.9) | 30 (4.7) | 20 (0.7) | 50 (2.3) | 8 (3.7) | 20 (1) |
| Holodiscus dumosus | - | - | - | - | - | - | - |
| Philadelphus lewisii | - | - | - | - | - | - | - |
| Purshia tridentata | 11 (0.1) | 38 (1.4) | 16 (2.3) | 33 (1) | 13 (1) | 13 (2.2) | 20 (1) |
| Ribes aureum | - | 4 (0.1) | - | 7 (0.1) | - | - | - |
| Symphoricarpos oreophilus | - | - | - | 7 (1) | - | - | 20 (1) |
| Tetradymia canescens | - | - | 7 (2.8) | - | - | 5 (2.5) | - |
| GRAMINOIDS | | | | | | | |
| Achnatherum hymenoides | - | 1 (0.1) | 9 (2.6) | 7 (2) | - | 13 (2.4) | 20 (2) |
| Achnatherum thurberianum | 33 (1.7) | 14 (1.5) | 41 (2) | 20 (4) | - | 23 (2.7) | 80 (1.8) |
| Agropyron cristatum | - | 3 (2) | 4 (3.7) | - | 100 (5.6) | - | 20 (0.1) |
| Agropyron fragile | - | - | 4 (2.7) | 13 (3) | 38 (29.3) | 5 (8.5) | - |
| Bromus japonicus | - | 3 (1) | 4 (1) | - | - | 3 (4) | - |
| Bromus tectorum | 89 (3.6) | 87 (2.8) | 100 (13.3) | 60 (2.5) | 100 (7.6) | 100 (11.9) | 100 (2.8) |
| Carex | - | - | 4 (3) | - | 13 (4) | - | - |
| Elymus elymoides | 33 (1) | 51 (0.5) | 50 (4.6) | 47 (3.6) | 25 (0.6) | 36 (1.6) | 20 (1) |
| Elymus lanceolatus | - | 1 (0.1) | 11 (3.1) | 27 (1.1) | - | - | - |
| Festuca idahoensis | 11 (1) | 12 (0.1) | - | 27 (1.3) | 13 (0.1) | 5 (3) | 20 (4) |
| Hesperostipa comata ssp. comata | 11 (0.1) | 13 (0.1) | 7 (4.2) | 20 (0.1) | 13 (1) | 15 (1) | 40 (0.6) |
| Koeleria macrantha | - | - | 1 (1) | 7 (9) | - | 3 (10) | - |
| Leymus cinereus | 11 (1) | 4 (0.7) | 28 (2) | 33 (0.6) | 13 (1) | 18 (1.6) | 40 (3) |
| Melica bulbosa | - | 1 (1) | - | - | - | - | 20 (1) |
| Pascopyrum smithii | 100 (3.8) | 8 (1) | 45 (3.5) | 27 (2.3) | 25 (5) | 26 (4.5) | 40 (2) |
| Poa | 11 (2) | 7 (2.8) | 5 (2.3) | - | - | 21 (1.3) | - |
| Poa ampla | - | - | 4 (1) | 13 (4.5) | - | - | - |
| Poa nevadensis | 11 (1) | - | 11 (1.5) | 7 (2) | - | 5 (4) | - |
| Poa secunda | 89 (5.3) | 99 (4.5) | 96 (10.2) | 73 (3.9) | 100 (9.9) | 95 (4.2) | 100 (6.6) |
| Pseudoroegneria spicata | 11 (1) | 11 (1) | 29 (2.8) | 100 (8.3) | - | 21 (2.6) | 40 (3) |
| FORBS | | | | | | | |
| Agoseris glauca | 11 (2) | 9 (0.1) | 16 (1.7) | 20 (0.1) | 13 (0.1) | 3 (24) | - |
| Allium | - | 11 (0.8) | 8 (1) | 13 (1.5) | 13 (1) | 3 (1) | 20 (2) |
| Allium acuminatum | - | 17 (0.2) | - | 27 (0.1) | - | 3 (0.1) | - |
| Alyssum desertorum | - | 4 (3) | 1 (3) | 13 (0.1) | - | 5 (1.5) | 20 (2) |
| Amsinckia | - | 1 (1) | - | - | 13 (1) | 10 (1) | - |
| Arabis | - | 8 (0.1) | 3 (1) | 13 (0.6) | - | - | - |
| Arabis holboellii | - | 5 (0.1) | - | - | - | - | - |
| Arenaria aculeata | - | 7 (1.2) | 4 (1.3) | - | 13 (3) | 10 (1.3) | - |
| Aster | - | - | 3 (1) | 7 (0.1) | - | - | - |
| Astragalus | 11 (2) | - | 12 (1.1) | 20 (1) | 25 (1) | - | - |
| Astragalus purshii var. glareosus | - | - | - | 7 (0.1) | 13 (0.1) | 3 (0.1) | - |
| Balsamorhiza sagittata | 11 (2) | 7 (1.1) | 8 (0.9) | 33 (1.4) | - | - | 20 (0.1) |
| Calochortus | - | - | 1 (1) | - | - | - | - |
| Castilleja | - | - | 1 (1) | - | - | - | - |
| Castilleja pallescens var. inverta | - | 3 (0.1) | - | 13 (0.1) | - | - | - |
| Ceratocephala testiculata | - | 1 (2) | - | - | - | - | - |
| Chaenactis douglasii | - | 9 (0.1) | 1 (0.1) | 13 (0.1) | 13 (0.1) | 3 (0.1) | - |
| Cirsium | - | 3 (0.1) | - | 7 (0.1) | - | - | - |
| Cirsium canovirens | - | - | - | - | - | - | - |
| Collinsia parviflora | - | 11 (1.5) | 4 (1.7) | 7 (0.1) | 25 (2) | 8 (2.3) | - |
| Crepis acuminata | 11 (0.1) | 28 (0.2) | 9 (1.2) | 53 (0.2) | 13 (0.1) | 3 (0.1) | - |
| Cryptantha | - | 3 (0.1) | - | - | - | - | - |
| Cryptantha spiculifera | - | 3 (0.1) | - | - | - | - | - |
| Delphinium andersonii | - | 18 (0.1) | 1 (0.1) | 13 (0.1) | 13 (0.1) | - | - |
| Descurainia pinnata ssp. filipes | 11 (0.1) | 1 (0.1) | - | 7 (0.1) | - | - | 20 (0.1) |
| Epilobium | - | - | 5 (1.8) | 7 (1) | 13 (2) | 15 (1.7) | 20 (3) |
| Erigeron | - | - | 3 (1) | - | - | - | - |
| Erigeron peregrinus | 11 (1) | 5 (1.3) | 7 (1.2) | - | - | 10 (1) | - |
| Erigeron pumilus | - | 8 (0.1) | 3 (0.2) | 27 (0.1) | - | 3 (0.1) | - |
| Eriogonum | 11 (1) | 4 (0.4) | 7 (1.4) | 13 (1) | 13 (1) | 5 (1) | - |
| Eriogonum caespitosum | - | 5 (0.1) | - | - | - | - | - |
| Eriogonum heracleoides | - | - | - | - | - | - | - |
| Eriogonum ovalifolium | - | 12 (0.1) | 1 (0.2) | 13 (0.1) | - | - | - |
| Eriogonum umbellatum | - | 7 (0.1) | 1 (4) | 7 (0.1) | - | - | - |
| Fabaceae | 11 (1) | 3 (0.6) | 5 (1) | - | - | 3 (1) | - |
| Gayophytum | 67 (3.3) | 14 (2.2) | 26 (1.8) | - | 13 (2) | 51 (1.7) | 60 (1) |

| | | | | | | | |
|---|----------|----------|----------|----------|----------|----------|----------|
| Gutierrezia sarothrae | - | - | - | - | - | - | - |
| Gymnosteris nudicaulis | - | 4 (0.1) | 1 (0.1) | 7 (0.1) | - | - | - |
| Gymnosteris parvula | - | 4 (0.1) | 1 (0.1) | 7 (0.1) | - | - | - |
| Lactuca serriola | 11 (1) | 4 (0.7) | 1 (0.2) | 20 (0.1) | 13 (2) | 3 (0.1) | - |
| Lappula occidentalis var. occidentalis | - | 1 (0.1) | - | 7 (0.1) | - | - | 20 (0.1) |
| Leptodactylon pungens | - | 36 (0.4) | 5 (0.6) | 27 (0.1) | - | 3 (1) | - |
| Lithophragma glabrum | - | 8 (0.1) | 1 (0.1) | - | - | - | - |
| Lithophragma tenellum | - | 4 (0.7) | 1 (0.1) | 7 (0.1) | - | - | - |
| Lithospermum ruderales | - | 1 (0.1) | 4 (0.7) | 7 (1) | 13 (0.1) | 3 (0.1) | - |
| Lomatium | 11 (2) | - | 16 (3.5) | 20 (3.7) | - | - | - |
| Lomatium dissectum | 11 (0.1) | 3 (0.1) | 1 (0.1) | 13 (0.1) | - | - | - |
| Lomatium foeniculaceum ssp. macdougalii | - | 8 (0.1) | - | 7 (0.1) | - | - | - |
| Lomatium nudicaule | - | 11 (2) | 5 (1.5) | - | 13 (1) | 5 (1) | - |
| Lomatium triternatum | - | 16 (0.3) | 1 (0.1) | 27 (0.1) | 13 (1) | - | - |
| Lupinus | - | 1 (1) | - | 13 (1.5) | - | 3 (1) | - |
| Machaeranthera canescens | - | 1 (2) | 4 (1) | - | 13 (1) | 8 (1) | 20 (1) |
| Nothocalais troximoides | - | 8 (0.1) | 1 (0.1) | - | - | - | - |
| Opuntia polyacantha | 11 (1) | 14 (0.3) | 4 (1) | 20 (0.1) | - | 8 (0.7) | 20 (0.1) |
| Orthocarpus | 22 (1.5) | 3 (1) | 8 (2.3) | - | - | 10 (1.3) | 20 (1) |
| Packera cana | - | 16 (0.1) | - | - | - | - | - |
| Penstemon | - | - | 1 (1) | 7 (0.1) | - | 3 (0.1) | - |
| Penstemon cyaneus | - | 5 (0.1) | 1 (0.2) | 27 (0.1) | - | - | - |
| Penstemon deustus | - | 21 (0.2) | 1 (0.1) | - | - | - | 20 (1) |
| Phacelia hastata | - | 4 (0.1) | 1 (0.1) | 7 (0.1) | 13 (0.1) | - | 20 (2) |
| Phlox | - | 3 (1.5) | 4 (1.3) | - | 13 (1) | 5 (1) | - |
| Phlox hoodii | 33 (2.3) | 33 (1.1) | 43 (1.9) | 47 (1.8) | 13 (1) | 21 (1.5) | - |
| Phlox longifolia | 44 (2) | 17 (1) | 53 (2.7) | 60 (2.8) | 50 (2) | 18 (1.2) | - |
| Potentilla glandulosa | - | 12 (0.1) | - | - | - | - | - |
| Pteryxia terebinthina var. foeniculacea | - | 16 (0.5) | - | 7 (0.1) | - | 3 (0.1) | - |
| Senecio integerrimus var. exaltatus | - | 3 (0.1) | - | 13 (0.1) | - | - | - |
| Sisymbrium altissimum | 11 (0.1) | 5 (1.3) | 3 (1) | 13 (1.1) | 13 (7) | 10 (1.3) | 60 (1.7) |
| Stephanomeria minor var. myrioclada | - | 8 (0.1) | - | - | - | - | - |
| Tragopogon dubius | 22 (1.1) | 8 (0.3) | 20 (1.2) | 13 (0.1) | 13 (0.1) | 3 (0.1) | 20 (0.1) |
| Unknown forb | 22 (2.5) | 5 (0.8) | 25 (1.4) | - | 38 (1.4) | 13 (2) | - |
| Unknown species | 11 (0.1) | 5 (0.8) | 4 (2.7) | 27 (0.1) | 13 (8) | 3 (0.3) | - |
| Viola nuttallii | - | 1 (0.1) | 1 (0.1) | - | - | - | - |

Part B: Shrublands and Shrub-Steppe (continued)

| Species | ERNA7-
HODU/
PEDE4 | JUOS/
ARTRW8/
HECO26 | PUTR2-
ERNA7/
PEDE4 | PUTR2-
PHLE4/
POSE | PUTR2/
LECI4 |
|--|--------------------------|----------------------------|---------------------------|--------------------------|-----------------|
| TREES | | | | | |
| Juniperus osteosperma | 18 (1.1) | 100 (6.5) | 13 (0.1) | - | - |
| Pinus flexilis | - | - | 7 (0.1) | 50 (1.5) | - |
| DWARF-SHRUBS | | | | | |
| Artemisia arbuscula | - | - | - | - | - |
| Ericameria nana | 100 (8) | 50 (0.1) | 100 (0.8) | 50 (0.1) | 25 (0.1) |
| Eriogonum microthecum | 36 (0.6) | 17 (0.1) | 80 (0.9) | 50 (0.1) | - |
| Artemisia tridentata ssp. tridentata | - | - | - | - | - |
| Artemisia tridentata ssp. vaseyana | - | - | - | 25 (0.1) | 50 (0.1) |
| Artemisia tridentata var. wyomingensis | - | 100 (5.5) | 47 (2.6) | 25 (0.1) | - |
| Artemisia tripartita | - | - | - | - | - |
| SHRUBS | | | | | |
| Chamaebatiaria millefolium | 9 (0.1) | - | 13 (0.1) | 75 (2) | - |
| Chrysothamnus viscidiflorus | - | 50 (10.3) | - | - | - |
| Ericameria nauseosa ssp. consimilis var. oreophila | 36 (0.4) | 100 (1.8) | 60 (0.7) | 50 (1.1) | 25 (1) |
| Holodiscus dumosus | 100 (2) | 33 (0.1) | 40 (0.2) | - | - |
| Philadelphus lewisii | 27 (1.4) | - | 27 (1) | 100 (4.5) | - |
| Purshia tridentata | 27 (0.4) | 83 (0.6) | 100 (4.5) | 100 (1.3) | 100 (9.3) |
| Ribes aureum | 18 (0.1) | - | 13 (0.6) | 50 (0.1) | - |
| Symphoricarpos oreophilus | - | - | - | - | - |
| Tetradymia canescens | - | - | - | - | - |
| GRAMINOIDS | | | | | |
| Achnatherum hymenoides | 55 (0.7) | 50 (0.1) | 20 (0.4) | 25 (0.1) | 25 (0.1) |
| Achnatherum thurberianum | 27 (0.1) | 17 (1) | 13 (1.3) | - | - |
| Agropyron cristatum | - | - | - | - | - |
| Agropyron fragile | - | - | - | - | - |
| Bromus japonicus | - | - | - | - | - |
| Bromus tectorum | 27 (0.1) | 100 (21.9) | 93 (1.1) | 50 (0.1) | 50 (2) |
| Carex | - | - | - | - | - |
| Elymus elymoides | 82 (0.2) | 33 (0.6) | 80 (0.1) | 75 (0.1) | 25 (0.1) |
| Elymus lanceolatus | - | 17 (0.1) | - | - | - |
| Festuca idahoensis | - | - | - | - | - |
| Hesperostipa comata ssp. comata | 18 (0.1) | 67 (2.3) | 47 (0.8) | - | 25 (0.1) |
| Koeleria macrantha | - | - | - | - | - |
| Leymus cinereus | - | - | - | - | 100 (3.3) |
| Melica bulbosa | - | - | - | - | - |
| Pascopyrum smithii | - | - | - | - | - |
| Poa | - | - | - | - | - |
| Poa ampla | - | - | - | - | - |
| Poa nevadensis | - | - | - | - | - |
| Poa secunda | 91 (1.4) | 83 (0.7) | 100 (1.8) | 100 (0.6) | 50 (1.1) |
| Pseudoroegneria spicata | - | 33 (21) | - | - | - |
| FORBS | | | | | |
| Agoseris glauca | - | 17 (1) | - | - | 25 (0.1) |
| Allium | - | - | - | - | 25 (0.1) |
| Allium acuminatum | - | 50 (1) | - | - | - |
| Alyssum desertorum | - | 50 (0.1) | 7 (0.1) | - | - |
| Amsinckia | - | - | - | - | - |
| Arabis | - | - | - | 50 (0.1) | - |
| Arabis holboellii | - | - | 13 (0.1) | 25 (0.1) | - |
| Arenaria aculeata | - | - | - | - | - |
| Aster | - | - | - | - | - |
| Astragalus | - | - | - | - | - |
| Astragalus purshii var. glareosus | - | - | - | - | - |
| Balsamorhiza sagittata | - | 33 (5) | - | - | - |
| Calochortus | - | - | - | - | - |
| Castilleja | 18 (0.1) | - | - | - | - |
| Castilleja pallescens var. inverta | - | - | - | - | - |
| Ceratocephala testiculata | - | - | - | - | - |
| Chaenactis douglasii | 55 (0.3) | 33 (0.1) | 40 (0.1) | - | 25 (0.1) |
| Cirsium | 9 (0.1) | 17 (0.1) | 7 (0.1) | 25 (0.1) | - |
| Cirsium canovirens | 45 (0.2) | - | 13 (0.6) | - | - |
| Collinsia parviflora | - | - | - | - | 50 (0.1) |
| Crepis acuminata | - | 33 (0.6) | - | - | 50 (1.1) |
| Cryptantha | - | - | - | - | - |
| Cryptantha spiculifera | 45 (0.4) | - | 33 (0.1) | 25 (0.1) | - |
| Delphinium andersonii | - | - | - | - | 50 (0.1) |
| Descurainia pinnata ssp. filipes | 9 (0.1) | 50 (0.1) | 33 (0.1) | 25 (0.1) | - |
| Epilobium | - | - | - | - | - |
| Erigeron | - | - | - | - | - |
| Erigeron peregrinus | - | - | - | - | - |
| Erigeron pumilus | 73 (0.1) | - | - | - | - |
| Eriogonum | 36 (0.7) | - | 20 (0.1) | - | - |
| Eriogonum caespitosum | 55 (0.7) | - | 33 (0.1) | 25 (0.1) | - |
| Eriogonum heracleoides | 9 (0.1) | - | 7 (0.1) | - | 25 (0.1) |
| Eriogonum ovalifolium | 9 (0.1) | - | 13 (0.1) | 25 (0.1) | 25 (0.1) |
| Eriogonum umbellatum | - | - | 40 (0.1) | 50 (0.1) | 50 (0.1) |
| Fabaceae | - | - | - | - | - |
| Gayophytum | - | - | - | - | 25 (0.1) |

| | | | | | |
|--|-----------|----------|-----------|-----------|----------|
| Gutierrezia sarothrae | 27 (0.1) | 50 (0.1) | 20 (0.1) | - | - |
| Gymnosteris nudicaulis | - | - | - | - | - |
| Gymnosteris parvula | - | - | - | - | - |
| Lactuca serriola | 45 (0.3) | 83 (0.3) | 47 (0.1) | - | - |
| Lappula occidentalis var. occidentalis | - | 33 (0.1) | 13 (0.1) | - | - |
| Leptodactylon pungens | 91 (0.5) | 33 (0.1) | 87 (0.5) | 50 (0.1) | - |
| Lithophragma glabrum | - | - | - | - | - |
| Lithophragma tenellum | - | - | - | - | - |
| Lithospermum ruderales | - | 17 (2) | - | - | - |
| Lomatium | 9 (0.1) | - | - | - | - |
| Lomatium dissectum | - | 50 (2.3) | - | - | 50 (0.6) |
| Lomatium foeniculaceum ssp. macdougallii | - | - | - | - | - |
| Lomatium nudicaule | - | - | - | - | - |
| Lomatium triternatum | - | 17 (0.1) | 7 (0.1) | - | - |
| Lupinus | - | - | - | - | 25 (0.1) |
| Machaeranthera canescens | - | - | - | - | - |
| Nothocalais troximoides | - | - | - | - | - |
| Opuntia polyacantha | 9 (0.1) | 67 (1.3) | 13 (0.1) | 50 (0.1) | - |
| Orthocarpus | - | - | - | - | - |
| Packera cana | 36 (0.5) | 17 (0.1) | 27 (0.6) | 75 (0.1) | - |
| Penstemon | - | - | - | - | 25 (0.1) |
| Penstemon cyaneus | - | 33 (0.1) | - | - | 25 (0.1) |
| Penstemon deustus | 100 (0.7) | 50 (2.4) | 100 (0.6) | 100 (0.1) | - |
| Phacelia hastata | 36 (0.1) | - | 80 (0.1) | 25 (0.2) | 75 (0.1) |
| Phlox | - | - | - | - | - |
| Phlox hoodii | 9 (0.1) | 17 (0.1) | - | - | - |
| Phlox longifolia | - | 17 (2) | - | - | - |
| Potentilla glandulosa | 82 (0.5) | 50 (0.1) | 73 (0.1) | 100 (0.1) | - |
| Pteryxia terebinthina var. foeniculacea | - | - | 13 (0.1) | 75 (0.7) | - |
| Senecio integerrimus var. exaltatus | - | - | - | - | 25 (0.1) |
| Sisymbrium altissimum | 9 (0.1) | 67 (0.3) | 13 (0.1) | - | - |
| Stephanomeria minor var. myrioclada | 64 (0.1) | 17 (0.1) | 67 (0.1) | - | - |
| Tragopogon dubius | 18 (0.6) | 50 (0.1) | 40 (0.1) | - | - |
| Unknown forb | - | - | 7 (0.1) | - | - |
| Unknown species | - | 50 (0.1) | 7 (0.1) | 25 (0.1) | 25 (0.1) |
| Viola nuttallii | - | - | - | - | - |

Part B: Shrublands and Shrub-Steppe (continued)

| Species | ALIN2/
MesicForbs | PRVI/
LECI4 |
|--|----------------------|----------------|
| TREES | | |
| <i>Populus tremuloides</i> | 100 (1.8) | 25 (0.6) |
| SHRUBS | | |
| <i>Alnus incana</i> | 100 (27.5) | 13 (2) |
| <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> | 25 (0.1) | 25 (0.1) |
| <i>Prunus virginiana</i> | 75 (0.7) | 100 (55.6) |
| <i>Rosa woodsii</i> var. <i>ultramontana</i> | 50 (0.6) | 25 (0.1) |
| <i>Salix boothii</i> | 75 (0.7) | - |
| <i>Symphoricarpos oreophilus</i> | 50 (0.1) | 75 (3.1) |
| GRAMINOIDS | | |
| <i>Bromus tectorum</i> | - | 38 (3.4) |
| <i>Leymus cinereus</i> | 75 (1.7) | 88 (3.3) |
| <i>Poa pratensis</i> | 100 (0.1) | 13 (0.1) |
| FORBS | | |
| <i>Agastache urticifolia</i> | - | 38 (0.1) |
| <i>Angelica pinnata</i> | 100 (0.1) | 63 (1.6) |
| <i>Aquilegia formosa</i> | 50 (0.1) | 13 (0.1) |
| <i>Chamerion angustifolium</i> ssp. <i>angustifolium</i> | 75 (0.1) | 13 (0.1) |
| <i>Crepis acuminata</i> | - | 38 (0.1) |
| <i>Galium bifolium</i> | 100 (0.8) | 63 (1) |
| <i>Hackelia floribunda</i> | 50 (0.1) | 38 (0.1) |
| <i>Heracleum maximum</i> | 100 (12.5) | 38 (5) |
| <i>Hydrophyllum capitatum</i> var. <i>capitatum</i> | 100 (0.1) | 50 (0.1) |
| <i>Lactuca serriola</i> | 25 (0.1) | 38 (0.1) |
| <i>Maianthemum racemosum</i> ssp. <i>amplexicaule</i> | 75 (0.1) | 38 (0.1) |
| <i>Maianthemum stellatum</i> | 100 (0.1) | 38 (0.1) |
| <i>Osmorhiza berteroi</i> | 75 (0.1) | - |
| <i>Solidago canadensis</i> var. <i>salebrosa</i> | 100 (0.1) | 63 (0.3) |
| <i>Taraxacum officinale</i> | 50 (0.1) | 25 (0.1) |
| Unknown species | 75 (0.3) | 13 (0.2) |
| <i>Urtica dioica</i> ssp. <i>gracilis</i> | 100 (5.5) | 50 (3) |
| <i>Verbascum thapsus</i> | 25 (0.1) | 38 (0.1) |
| <i>Viola</i> | 25 (0.1) | 25 (0.1) |

Part C: Herbaceous Vegetation

| Species | ACTH7-
BRTE | BRTE | JUSC2-
JUOS/
ERNA7 | LECI4 | POSE | POSE-
BRTE | PSSP6-
POSE | PSSP6/
CRAC2 | SIAL2/
BRTE |
|--|----------------|------------|--------------------------|------------|----------|---------------|----------------|-----------------|----------------|
| TREES | | | | | | | | | |
| Juniperus osteosperma | - | - | 86 (3.8) | - | - | - | - | - | 4 (0.1) |
| Juniperus scopulorum | - | - | 93 (3.3) | - | - | - | - | - | - |
| DWARF-SHRUBS | | | | | | | | | |
| Artemisia arbuscula | - | 2 (2) | - | - | 13 (3) | 2 (1) | 9 (2) | - | - |
| Ericameria nana | - | 2 (0.1) | 100 (4.7) | - | - | - | - | - | - |
| Eriogonum microthecum | - | 6 (0.1) | 29 (1) | - | - | - | 27 (0.1) | 50 (0.1) | - |
| Monardella odoratissima | - | - | 29 (1.9) | - | - | - | - | - | - |
| SHRUBS | | | | | | | | | |
| Artemisia tridentata ssp. vaseyana | - | 2 (0.1) | - | 25 (0.1) | - | 4 (3.5) | 18 (2.6) | 33 (3) | - |
| Artemisia tridentata var. wyomingensis | 56 (2.8) | 38 (2.6) | 21 (3.7) | - | 75 (3.5) | 54 (2.6) | 45 (2.4) | 17 (0.1) | 25 (1.4) |
| Artemisia tripartita | - | 10 (1.8) | - | - | 25 (2) | 13 (1.7) | 27 (0.1) | 33 (0.1) | 4 (2) |
| Chrysothamnus viscidiflorus | 33 (2.7) | 42 (2.1) | - | - | 38 (1.7) | 62 (2.3) | 55 (1.5) | 17 (5) | 33 (1.5) |
| Ericameria nauseosa ssp. consimilis var. oreophila | - | 6 (0.7) | 71 (1) | - | - | 2 (1.5) | 36 (0.3) | 50 (0.4) | 4 (1) |
| Holodiscus dumosus | - | - | 57 (0.6) | - | - | - | - | - | - |
| Purshia tridentata | 11 (4) | 10 (0.9) | 57 (2.3) | - | 13 (2) | 4 (2.8) | 27 (0.1) | 17 (0.1) | 4 (0.1) |
| Ribes aureum | - | - | 29 (1.6) | - | - | - | - | 17 (0.1) | - |
| Tetradymia canescens | - | 2 (0.1) | - | - | - | 1 (2) | 27 (0.1) | 33 (3) | - |
| GRAMINOIDS | | | | | | | | | |
| Achnatherum hymenoides | - | 10 (0.7) | 50 (1.3) | - | - | 6 (1.4) | 18 (1.1) | 33 (0.1) | - |
| Achnatherum nelsonii ssp. dorei | - | 6 (1.3) | - | - | - | 1 (2) | - | - | 4 (1) |
| Achnatherum thurberianum | 100 (9.6) | - | 43 (0.7) | - | 25 (2.5) | 26 (2) | - | - | - |
| Agropyron cristatum | - | 17 (1.6) | - | - | - | 6 (1.3) | 9 (1) | - | 29 (1.7) |
| Aristida purpurea | - | - | 43 (1.8) | - | - | - | - | - | - |
| Bromus japonicus | 11 (2) | 2 (1) | - | - | 25 (3.5) | 6 (1.8) | - | - | - |
| Bromus tectorum | 100 (6.9) | 100 (17.5) | 57 (3.2) | 50 (1.1) | 50 (1.5) | 100 (10.8) | 55 (1.4) | 50 (0.7) | 100 (17.7) |
| Elymus elymoides | 11 (2) | 19 (0.5) | 57 (0.6) | - | 38 (1.3) | 27 (1.2) | 9 (0.1) | 17 (0.1) | 13 (1) |
| Elymus lanceolatus | - | 6 (0.4) | - | - | - | 1 (1) | - | - | 8 (0.1) |
| Festuca idahoensis | - | 2 (1) | - | - | - | - | 18 (0.1) | 17 (1) | - |
| Hesperostipa comata ssp. comata | - | 8 (0.6) | 43 (1) | - | - | 14 (2.2) | 18 (1.6) | 17 (0.1) | 4 (1) |
| Leymus cinereus | 11 (3) | 15 (0.8) | - | 100 (27.5) | 13 (1) | 17 (2) | 36 (2.6) | 33 (0.1) | 13 (1) |
| Pascopyrum smithii | 67 (1.7) | 4 (1) | - | - | 25 (1) | 17 (2.8) | 9 (2) | - | 4 (2) |
| Poa | 22 (1) | 6 (2) | - | - | 25 (1.5) | 14 (2) | - | 17 (1) | 8 (1.5) |
| Poa bulbosa | - | 2 (0.1) | - | 25 (0.1) | - | 5 (1.8) | - | - | 4 (10) |
| Poa secunda | 67 (3) | 73 (1.5) | 93 (2.7) | 25 (0.1) | 100 (11) | 94 (6.1) | 100 (2.4) | 83 (0.5) | 75 (6.3) |
| Pseudoroegneria spicata | 22 (1.5) | 19 (1) | 14 (1.1) | - | 50 (1.8) | 19 (4) | 100 (12.9) | 100 (9.5) | - |
| Thinopyrum intermedium | - | 10 (1.4) | - | - | - | 4 (5) | - | - | - |
| FORBS | | | | | | | | | |
| Agoseris glauca | - | 2 (0.1) | - | 25 (0.1) | - | 3 (0.7) | 18 (1.1) | - | 4 (0.1) |
| Allium | - | 2 (1) | - | - | 25 (1) | 6 (0.9) | - | - | 4 (1) |
| Allium acuminatum | - | 6 (0.1) | - | - | - | 4 (0.1) | 18 (0.6) | 17 (0.1) | - |
| Alyssum desertorum | 33 (1.3) | 15 (0.7) | - | - | - | 14 (1.8) | 9 (0.1) | - | 21 (2.8) |
| Amsinckia | - | 2 (1) | - | - | - | 5 (1.2) | - | - | 8 (1) |
| Arenaria aculeata | - | - | - | - | - | 5 (1.4) | - | - | 4 (1) |
| Aster | - | 4 (0.6) | - | - | - | 6 (1.2) | 9 (0.1) | - | 4 (2) |
| Astragalus | - | - | - | 25 (0.1) | - | 1 (0.1) | 9 (0.1) | 17 (0.1) | 4 (1) |
| Astragalus purshii var. glareosus | - | 2 (0.1) | - | - | - | 1 (0.1) | 9 (0.1) | 33 (0.1) | - |
| Balsamorhiza sagittata | - | 2 (9) | - | - | 25 (1.5) | 5 (3) | 18 (1.1) | 33 (0.1) | - |
| Ceratocephala testiculata | - | 8 (2.5) | - | - | 13 (1) | 5 (1) | - | - | 8 (1.5) |
| Chaenactis | - | - | - | - | - | 9 (1.3) | - | - | 17 (1.3) |
| Chaenactis douglasii | - | 4 (0.1) | 29 (0.1) | - | - | 1 (0.1) | - | 17 (0.1) | - |
| Cirsium canovirens | - | - | 50 (0.4) | - | - | - | - | - | - |
| Collinsia parviflora | 67 (3.3) | 12 (2.7) | - | - | - | 13 (2.3) | 9 (0.1) | - | 13 (1.7) |
| Crepis acuminata | - | 13 (0.2) | - | - | 13 (2) | 9 (0.9) | 73 (0.7) | 83 (0.5) | - |
| Cryptantha | - | 4 (0.6) | - | - | - | 4 (1) | - | 17 (0.1) | 13 (1.7) |
| Cryptantha spiculifera | - | - | 43 (0.2) | - | - | - | 9 (0.1) | - | - |
| Delphinium andersonii | - | 4 (0.1) | - | - | - | - | 55 (0.1) | 67 (0.1) | - |
| Descurainia pinnata ssp. filipes | - | - | 7 (0.1) | 50 (0.1) | - | 1 (0.1) | - | 17 (0.1) | 13 (1.7) |
| Epilobium | 11 (3) | 2 (3) | - | - | - | 16 (1.8) | - | 17 (1) | 4 (2) |
| Erigeron peregrinus | - | 4 (1) | - | - | - | 4 (1) | - | - | 4 (1) |
| Erigeron pumilus | - | 8 (0.1) | 14 (0.1) | - | - | 1 (0.1) | 9 (0.1) | 33 (0.1) | - |
| Eriogonum | - | - | 36 (0.9) | - | 25 (3) | 3 (1) | 9 (2) | - | - |
| Eriogonum caespitosum | - | - | 36 (0.6) | - | - | - | - | - | - |
| Eriogonum ovalifolium | - | 4 (0.1) | - | - | - | - | 27 (0.1) | 33 (0.1) | - |
| Fabaceae | - | - | - | - | - | 2 (1) | - | 17 (0.1) | 4 (0.1) |
| Fritillaria pudica | - | 8 (0.1) | - | - | - | 2 (1.5) | - | - | - |
| Gayophytum | 44 (2) | 10 (1.8) | - | - | 25 (3) | 26 (1.9) | 9 (5) | - | 17 (2.3) |
| Kochia scoparia | - | - | - | 25 (1) | - | 3 (0.7) | - | - | 17 (1.3) |
| Lactuca serriola | - | 19 (1) | 29 (0.1) | 25 (1) | - | 12 (1.7) | 9 (0.1) | - | 58 (2.3) |
| Leptodactylon pungens | - | 4 (0.1) | 86 (2) | - | - | 2 (0.1) | 36 (0.6) | 50 (1) | - |
| Lithospermum ruderales | - | 8 (0.6) | - | - | - | 4 (0.8) | 9 (1) | - | - |
| Lomatium dissectum | - | 2 (0.1) | - | - | - | 1 (10) | 18 (0.6) | - | - |
| Lomatium foeniculaceum ssp. macdougallii | - | 4 (0.1) | - | - | - | - | 36 (0.1) | 67 (0.1) | - |
| Lomatium nudicaule | - | - | - | - | 13 (4) | 10 (1.1) | 18 (1) | - | 8 (2) |
| Lomatium triternatum | - | 6 (0.4) | - | - | - | 1 (0.2) | 36 (0.1) | 67 (0.3) | - |
| Lupinus | - | 2 (1) | - | 25 (0.1) | 13 (1) | 3 (1) | 9 (9) | 50 (0.7) | - |
| Lupinus argenteus | - | - | - | - | - | 1 (1) | 36 (0.6) | - | - |
| Machaeranthera canescens | 11 (1) | 4 (1) | - | - | - | 9 (1.1) | - | - | 8 (1) |
| Mentzelia albicaulis | - | 2 (0.1) | - | 25 (0.1) | - | 1 (0.1) | - | - | 4 (0.1) |
| Nothocalais troximoides | - | 6 (0.1) | - | 50 (0.1) | - | - | - | - | - |

| | | | | | | | | | |
|---|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| Opuntia polyacantha | 11 (2) | 6 (0.1) | 29 (0.6) | - | - | 3 (0.7) | 9 (2) | 33 (0.1) | 4 (0.1) |
| Orthocarpus | - | - | - | - | - | 7 (1.1) | - | - | - |
| Packera cana | - | - | 36 (1.1) | - | - | 1 (0.1) | 9 (0.1) | - | - |
| Penstemon | - | 6 (0.1) | - | - | - | 1 (0.2) | - | 33 (0.1) | - |
| Penstemon cyaneus | - | - | - | - | - | 1 (0.1) | 36 (0.1) | 33 (1.1) | - |
| Penstemon deustus | - | 2 (0.1) | 93 (2.6) | 25 (0.1) | - | 2 (1) | - | 17 (0.1) | - |
| Phacelia hastata | - | 8 (0.8) | 43 (0.1) | - | - | 2 (0.6) | 36 (0.1) | 33 (0.6) | - |
| Phlox | - | - | 7 (0.1) | - | - | 4 (1) | - | - | - |
| Phlox hoodii | 33 (1.3) | 10 (0.1) | - | - | 25 (1.5) | 26 (1.6) | 36 (0.8) | 33 (0.1) | 8 (1) |
| Phlox longifolia | 44 (2) | 15 (0.8) | - | - | 13 (5) | 22 (1.7) | 18 (0.1) | 17 (3) | 17 (0.8) |
| Potentilla glandulosa | - | - | 93 (0.8) | - | - | - | - | - | - |
| Pteryxia terebinthina var. foeniculacea | - | 4 (0.1) | - | - | - | 1 (0.1) | 9 (0.1) | - | - |
| Salsola tragus | - | - | - | - | - | 5 (1.8) | - | - | 21 (2) |
| Sisymbrium altissimum | 22 (1.5) | 37 (1.3) | 7 (0.1) | 25 (0.1) | - | 24 (3.2) | 36 (0.3) | - | 100 (14) |
| Sisymbrium loeselii | - | 2 (0.1) | - | 25 (0.1) | - | 1 (0.1) | - | - | 4 (0.1) |
| Stephanomeria minor var. myrioclada | - | - | 79 (0.5) | - | - | - | - | - | - |
| Thlaspi | - | 4 (1.5) | - | - | - | 4 (2.8) | - | - | 4 (2) |
| Tragopogon dubius | 11 (1) | 31 (2.9) | 43 (0.1) | - | 13 (1) | 27 (2.7) | 18 (0.1) | 17 (0.1) | 71 (2.4) |
| Unknown forb | 22 (1) | 6 (0.7) | 21 (1) | - | - | 12 (1.4) | 27 (0.4) | - | 17 (1.5) |
| Unknown species | - | 17 (1.7) | - | 25 (0.2) | - | 7 (0.5) | 27 (0.1) | 17 (0.1) | 13 (1.7) |
| Viola nuttallii | - | 2 (0.1) | - | 75 (0.1) | - | - | 18 (0.1) | - | 4 (0.1) |

Part D: Sparse Vegetation

| Species | CHM12/
PEDE4 | ERNA7/
POSE/
PEDE4 | EROVD | PHLE4/
PEDE4 | PIFL2/
CHM12/
POSE | POSE/
LEPU |
|---|-----------------|--------------------------|-----------|-----------------|--------------------------|---------------|
| TREES | | | | | | |
| <i>Pinus flexilis</i> | - | - | 25 (7) | - | 100 (1.5) | - |
| Dwarf-Shrubs | | | | | | |
| <i>Ericameria nana</i> | - | 100 (1.5) | - | 25 (0.1) | - | 45 (0.3) |
| <i>Eriogonum microthecum</i> | - | 73 (1) | - | 25 (0.1) | - | 36 (1.6) |
| SHRUBS | | | | | | |
| <i>Artemisia tridentata</i> ssp. <i>tridentata</i> | 40 (0.1) | - | - | - | 25 (0.1) | 9 (2) |
| <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> | 20 (0.1) | - | - | - | 75 (1) | 18 (3) |
| <i>Artemisia tridentata</i> var. <i>wyomingensis</i> | 20 (0.1) | 27 (1) | - | - | - | 36 (3.8) |
| <i>Chamaebatiaria millefolium</i> | 100 (0.7) | - | - | 63 (0.3) | 100 (0.8) | 9 (0.1) |
| <i>Ericameria nauseosa</i> ssp. <i>consimilis</i> var. <i>oreophila</i> | - | 27 (0.7) | - | - | 25 (0.1) | 55 (0.1) |
| <i>Holodiscus dumosus</i> | - | 64 (0.1) | - | 13 (0.1) | 50 (0.1) | 9 (0.1) |
| <i>Philadelphus lewisii</i> | - | 27 (1.7) | - | 100 (0.6) | 100 (1.5) | 9 (0.1) |
| <i>Purshia tridentata</i> | - | 64 (0.7) | - | 13 (0.1) | 25 (0.1) | 18 (0.6) |
| <i>Ribes aureum</i> | - | 9 (0.1) | - | - | 75 (0.1) | - |
| GRAMINOIDS | | | | | | |
| <i>Achnatherum hymenoides</i> | - | 27 (0.1) | - | - | - | 18 (0.6) |
| <i>Bromus tectorum</i> | - | 45 (0.5) | - | - | - | 45 (1) |
| <i>Elymus elymoides</i> | 20 (0.1) | 91 (0.3) | - | 100 (0.1) | 25 (0.1) | 73 (0.8) |
| <i>Hesperostipa comata</i> ssp. <i>comata</i> | 40 (0.1) | 82 (0.8) | - | 13 (0.1) | - | 27 (0.4) |
| <i>Poa secunda</i> | 60 (0.1) | 100 (0.6) | - | 38 (0.1) | 100 (0.1) | 100 (4.8) |
| <i>Pseudoroegneria spicata</i> | - | - | - | 13 (0.1) | 75 (1.7) | 18 (2) |
| FORBS | | | | | | |
| <i>Arabis holboellii</i> | - | - | - | 25 (0.1) | - | 27 (0.1) |
| <i>Castilleja angustifolia</i> var. <i>angustifolia</i> | - | 27 (0.1) | - | 13 (0.1) | - | 18 (0.1) |
| <i>Chaenactis douglasii</i> | - | 27 (0.1) | 75 (0.1) | - | - | 27 (0.1) |
| <i>Cirsium</i> | - | 27 (0.1) | 25 (0.1) | 13 (0.1) | - | 18 (0.1) |
| <i>Cirsium canovirens</i> | - | 9 (0.1) | - | - | 25 (0.1) | 18 (0.1) |
| <i>Crepis acuminata</i> | - | - | - | - | 50 (0.1) | 9 (0.1) |
| <i>Dryopteris filix-mas</i> | - | 9 (0.1) | - | - | 75 (0.1) | - |
| <i>Eriogonum ovalifolium</i> | - | - | - | - | 25 (0.1) | 27 (0.1) |
| <i>Eriogonum ovalifolium</i> var. <i>depressum</i> | - | - | 100 (0.8) | - | - | - |
| <i>Eriogonum umbellatum</i> | - | 64 (0.1) | 25 (0.1) | 13 (0.1) | - | 27 (0.1) |
| <i>Gilia leptomeria</i> | 20 (0.1) | 9 (0.1) | - | 50 (0.1) | 100 (0.1) | 18 (0.1) |
| <i>Leptodactylon pungens</i> | - | 91 (0.8) | - | 25 (0.1) | 75 (0.1) | 82 (0.8) |
| <i>Lewisia rediviva</i> | - | - | 75 (0.1) | - | - | - |
| <i>Lithophragma tenellum</i> | - | - | - | - | 100 (0.1) | - |
| <i>Lomatium triternatum</i> | - | 27 (0.1) | - | - | - | 9 (0.1) |
| <i>Opuntia polyacantha</i> | - | 36 (0.1) | - | - | - | 55 (0.1) |
| <i>Packera cana</i> | 20 (0.1) | 27 (0.1) | - | 25 (0.1) | - | 36 (0.1) |
| <i>Penstemon deustus</i> | 60 (0.1) | 100 (0.3) | - | 100 (0.2) | 25 (0.1) | 64 (0.2) |
| <i>Phacelia hastata</i> | - | 91 (0.2) | 100 (0.3) | - | - | 36 (0.1) |
| <i>Phlox hoodii</i> | - | 9 (0.1) | - | 25 (0.2) | - | 18 (0.6) |
| <i>Potentilla glandulosa</i> | 20 (0.1) | 91 (0.6) | - | 88 (0.6) | 25 (0.1) | 27 (0.4) |
| <i>Pteryxia terebinthina</i> var. <i>foeniculacea</i> | - | 9 (0.1) | - | 75 (0.1) | 75 (0.1) | 36 (0.3) |
| <i>Stephanomeria minor</i> var. <i>myrioclada</i> | 20 (0.1) | 73 (0.1) | - | 13 (0.1) | - | 36 (0.1) |
| <i>Tragopogon dubius</i> | - | 18 (0.1) | - | - | - | 27 (0.1) |
| <i>Viola nuttallii</i> | - | - | - | - | 75 (0.1) | - |

National Park Service

U.S. Department of the Interior



Natural Resource Program Center

1201 Oak Ridge Drive, Suite 150
Fort Collins, CO 80525

www.nature.nps.gov

EXPERIENCE YOUR AMERICA™