

Conjunctival lymphoma

Category(ies): Cornea, External Disease

Contributor: [Jesse Vislisel, MD](#)

Photographer: Cindy Montague, CRA

Conjunctival lymphoma classically appears as a smooth, mobile, salmon-colored, fleshy mass. The lesions are most commonly unilateral. They cannot be differentiated from [benign lymphoid hyperplasia](#) based on clinical appearance alone, and thus require biopsy. This lesion was biopsied and found to be an extranodal marginal zone (MALT) B-cell lymphoma.

EyeRounds.org



 Enlarge

 Download

Image Permissions:



Ophthalmic Atlas Images by [EyeRounds.org, The University of Iowa](#) are licensed under a [Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License](#).

Related Articles



[Related Atlas Entry: Conjunctival lymphoid hyperplasia](#)

Address

University of Iowa
Roy J. and Lucille A. Carver College
of Medicine
Department of Ophthalmology and
Visual Sciences
200 Hawkins Drive
Iowa City, IA 52242

[Support Us](#)

Legal

Copyright © 2019 The University of
Iowa. All Rights Reserved
[Report an issue with this page](#)
[Web Privacy Policy](#) |
[Nondiscrimination Statement](#)

Related Links

[Cataract Surgery for Greenhorns](#)
[EyeTransillumination](#)
[Gonioscopy.org](#)
[Iowa Glaucoma Curriculum](#)
[Iowa Wet Lab](#)
[Patient Information](#)
[Stone Rounds](#)
[The Best Hits Bookshelf](#)

EyeRounds Social Media

Follow



Receive notification of new cases,
[sign up here](#)
[Contact Us](#)
[Submit a Suggestion](#)