

# A note on the abelianizations of finite-index subgroups of the mapping class group

Andrew Putman

November 3, 2018

## 1 Introduction

Let  $\Sigma_{g,b}^p$  be an oriented genus  $g$  surface with  $b$  boundary components and  $p$  punctures and let  $\text{Mod}(\Sigma_{g,b}^p)$  be its *mapping class group*, that is, the group of isotopy classes of orientation-preserving diffeomorphisms of  $\Sigma_{g,b}^p$  that fix the boundary components and punctures pointwise (we will omit  $b$  or  $p$  when they are zero). A long-standing conjecture of Ivanov (see [6] for a recent discussion) says that for  $g \geq 3$ , the group  $\text{Mod}(\Sigma_{g,b}^p)$  does not virtually surject onto  $\mathbb{Z}$ . In other words, if  $\Gamma$  is a finite-index subgroup of  $\text{Mod}(\Sigma_{g,b}^p)$ , then  $H_1(\Gamma; \mathbb{R}) = 0$ .

The goal of this note is to offer some evidence for this conjecture. If  $G$  is a group and  $g \in G$ , then we will denote by  $[g]_G$  the corresponding element of  $H_1(G; \mathbb{R})$ . Also, for a simple closed curve  $\gamma$  on  $\Sigma_{g,b}^p$ , we will denote by  $T_\gamma$  the corresponding right Dehn twist. Observe that if  $\Gamma$  is any finite-index subgroup of  $\text{Mod}(\Sigma_{g,b}^p)$ , then  $T_\gamma^n \in \Gamma$  for some  $n \geq 1$ . Our first result is the following.

**Theorem A** (Powers of twists vanish). *For some  $g \geq 3$ , let  $\Gamma < \text{Mod}(\Sigma_{g,b}^p)$  satisfy  $[\text{Mod}(\Sigma_{g,b}^p) : \Gamma] < \infty$  and let  $\gamma$  be a simple closed curve on  $\Sigma_{g,b}^p$ . Pick  $n \geq 1$  such that  $T_\gamma^n \in \Gamma$ . Then  $[T_\gamma^n]_\Gamma = 0$ .*

*Remark.* After this paper was written, Bridson informed us that in unpublished work, he had proven a result about mapping class group actions on  $\text{CAT}(0)$  spaces that implies Theorem A. Bridson's work will appear in [3].

We use this to verify Ivanov's conjecture for a class of examples. For a long time, the only positive evidence for Ivanov's conjecture was a result of Hain [5] that says that it holds for all finite-index subgroups containing the *Torelli group*  $\mathcal{I}_{g,b}^p$ , that is, the kernel of the action of  $\text{Mod}(\Sigma_{g,b}^p)$  on  $H_1(\Sigma_g; \mathbb{Z})$  induced by filling in all the punctures and boundary components. The group  $\mathcal{I}_{g,b}^p$  contains the *Johnson kernel*  $\mathcal{K}_{g,b}^p$ , which is the subgroup generated by Dehn twists about separating curves. A result of Johnson [7] says that  $\mathcal{K}_{g,b}^p$  is an infinite-index subgroup of  $\mathcal{I}_{g,b}^p$ .

For a subgroup  $\Gamma$  of  $\text{Mod}(\Sigma_{g,b}^p)$ , denote by  $K(\Gamma)$  the subgroup of  $\Gamma \cap \mathcal{K}_{g,b}^p$  generated by the set

$$\{T_\gamma^n \mid \gamma \text{ a separating curve, } n \in \mathbb{Z}, \text{ and } T_\gamma^n \in \Gamma\}.$$

If  $\mathcal{K}_{g,b}^p < \Gamma$ , then  $K(\Gamma) = \Gamma \cap \mathcal{K}_{g,b}^p$ , but the converse does not hold. Our second result is the following.

**Theorem B** (Subgroups containing large pieces of Johnson kernel). *For some  $g \geq 3$ , let  $\Gamma < \text{Mod}(\Sigma_{g,b}^p)$  satisfy  $[\text{Mod}(\Sigma_{g,b}^p) : \Gamma] < \infty$ . Assume that  $[\Gamma \cap \mathcal{K}_{g,b}^p : K(\Gamma)] < \infty$ . Then  $H_1(\Gamma; \mathbb{R}) = 0$ .*

As a corollary, we obtain the following result, which was recently proven by Boggi [2] via a difficult algebro-geometric argument under the assumption  $b = p = 0$ .

**Corollary C** (Subgroups containing Johnson kernel). *For some  $g \geq 3$ , let  $\Gamma < \text{Mod}(\Sigma_{g,b}^p)$  satisfy  $[\text{Mod}(\Sigma_{g,b}^p) : \Gamma] < \infty$ . Assume that  $\mathcal{K}_{g,b}^n < \Gamma$ . Then  $H_1(\Gamma; \mathbb{R}) = 0$ .*

*Remark.* McCarthy [11] proved that Ivanov’s conjecture fails in the case  $g = 2$ .

**Acknowledgments.** I wish to thank Martin Bridson, Benson Farb, Thomas Koberda, Dan Margalit, and Ben Wieland for useful comments and conversations. I also wish to thank Dongping Zhuang for showing me how to slightly weaken the hypotheses in my original version of Theorem B.

## 2 Notation and basic facts about group homology

If  $M$  is a  $G$ -module, then  $M_G$  will denote the *coinvariants* of the action, that is, the quotient of  $M$  by the submodule generated by the set  $\{x - g(x) \mid x \in M, g \in G\}$ . This appears in the 5-term exact sequence [4, Corollary VII.6.4], which asserts the following. If

$$1 \longrightarrow K \longrightarrow G \longrightarrow Q \longrightarrow 1$$

is a short exact sequence of groups, then for any ring  $R$ , there is an exact sequence

$$H_2(G; R) \longrightarrow H_2(Q; R) \longrightarrow (H_1(K; R))_Q \longrightarrow H_1(G; R) \longrightarrow H_1(Q; R) \longrightarrow 0.$$

If  $G_2 < G_1$  are groups satisfying  $[G_1 : G_2] < \infty$  and  $R$  is a ring, then for all  $k$  there exists a *transfer map* of the form  $t : H_k(G_1; R) \rightarrow H_k(G_2; R)$  (see, e.g., [4, Chapter III.9]). The key property of  $t$  (see [4, Proposition III.9.5]) is that if  $i : H_k(G_2; R) \rightarrow H_k(G_1; R)$  is the map induced by the inclusion, then  $i \circ t : H_k(G_1; R) \rightarrow H_k(G_1; R)$  is multiplication by  $[G_1 : G_2]$ . In particular, if  $R = \mathbb{R}$ , then we obtain a right inverse  $\frac{1}{[G_1 : G_2]}t$  to  $i$ . This yields the following standard lemma.

**Lemma 2.1.** *Let  $G_2 < G_1$  be groups satisfying  $[G_1 : G_2] < \infty$ . For all  $k$ , the map  $H_k(G_2; \mathbb{R}) \rightarrow H_k(G_1; \mathbb{R})$  is surjective.*

## 3 Proof of Theorem A

Let  $n \geq 1$  be the smallest integer such that  $T_\gamma^n \in \Gamma$ .

We first claim that there exists a subsurface  $S \hookrightarrow \Sigma_{g,b}^p$  whose genus is at least 2 with the following property. Let  $i : \text{Mod}(S) \rightarrow \text{Mod}(\Sigma_{g,b}^p)$  be the induced map (“extend by the identity”). Then there exists some boundary component  $\beta$  of  $S$  such that  $i(T_\beta) = T_\gamma$ . There are two cases. If  $\gamma$  is nonseparating, then let  $S$  be the complement of a regular neighborhood of  $\gamma$ . Observe that  $S \cong \Sigma_{g-1, b+2}^p$ , so the genus of  $S$  is at least 2. If instead  $\gamma$  is separating, then let  $S$  be the component of  $\Sigma_{g,b}^p$  cut along  $\gamma$  whose genus is maximal. Since  $g \geq 3$ , this subsurface must have genus at least 2. The claim follows.

Define  $\Gamma' = i^{-1}(\Gamma)$ . We have  $T_\beta^n \in \Gamma'$ , and it is enough to show that  $[T_\beta^n]_{\Gamma'} = 0$ . Let  $\bar{S}$  be the result of gluing a punctured disc to  $\beta$  and let  $\bar{\Gamma}'$  be the image of  $\Gamma'$  in  $\text{Mod}(\bar{S})$ . There is a diagram of central extensions

$$\begin{array}{ccccccccc} 1 & \longrightarrow & \mathbb{Z} & \longrightarrow & \Gamma' & \longrightarrow & \bar{\Gamma}' & \longrightarrow & 1 \\ & & \downarrow \times n & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \mathbb{Z} & \longrightarrow & \text{Mod}(S) & \longrightarrow & \text{Mod}(\bar{S}) & \longrightarrow & 1 \end{array}$$

with  $\mathbb{Z} < \text{Mod}(S)$  and  $\mathbb{Z} < \Gamma'$  generated by  $T_\beta$  and  $T_\beta^n$ , respectively. The last 4 terms of the corresponding diagram of 5-term exact sequences are

$$\begin{array}{ccccccccc} H_2(\bar{\Gamma}'; \mathbb{R}) & \xrightarrow{f_1} & \mathbb{R} & \longrightarrow & H_1(\Gamma'; \mathbb{R}) & \longrightarrow & H_1(\bar{\Gamma}'; \mathbb{R}) & \longrightarrow & 0 \\ \downarrow f_2 & & \downarrow \cong & & \downarrow & & \downarrow & & \\ H_2(\text{Mod}(\bar{S}); \mathbb{R}) & \xrightarrow{f_3} & \mathbb{R} & \longrightarrow & H_1(\text{Mod}(S); \mathbb{R}) & \longrightarrow & H_1(\text{Mod}(\bar{S}); \mathbb{R}) & \longrightarrow & 0 \end{array}$$

We remark that there are no nontrivial coinvariants in these sequences since our extensions are central. We must show that  $f_1$  is a surjection. Since  $S$  has genus at least 2, we have  $H_1(\text{Mod}(S); \mathbb{R}) = 0$  (see, e.g., [10]), so  $f_3$  is a surjection. Since  $[\text{Mod}(\bar{S}) : \bar{\Gamma}'] < \infty$ , Lemma 2.1 implies that  $f_2$  is a surjection, so  $f_1$  is a surjection, as desired.

## 4 Proof of Theorem B

### 4.1 Two facts about $\text{Sp}_{2g}(\mathbb{Z})$

We will need two standard facts about finite-index subgroups  $\Gamma$  of  $\text{Sp}_{2g}(\mathbb{Z})$ , both of which follow from the fact that  $\Gamma$  is a lattice in  $\text{Sp}_{2g}(\mathbb{R})$ .

For the first, since  $\text{Sp}_{2g}(\mathbb{R})$  is a connected simple Lie group with finite center and real rank  $g$ , the group  $\Gamma$  has Kazhdan's property (T) when  $g \geq 2$  (see, e.g., [13, Theorem 7.1.4]). One standard property of groups with property (T) is that they have no nontrivial homomorphisms to  $\mathbb{R}$  (see, e.g., [13, Theorem 7.1.7]). Combining these facts, we obtain the following theorem.

**Theorem 4.1.** *For some  $g \geq 2$ , let  $\Gamma < \text{Sp}_{2g}(\mathbb{Z})$  satisfy  $[\text{Sp}_{2g}(\mathbb{Z}) : \Gamma] < \infty$ . Then  $H_1(\Gamma; \mathbb{R}) = 0$ .*

For the second, since  $\text{Sp}_{2g}(\mathbb{R})$  is a connected noncompact simple real algebraic group, we can apply the Borel density theorem (see, e.g., [13, Theorem 3.2.5]) to deduce that  $\Gamma$  is Zariski dense in  $\text{Sp}_{2g}(\mathbb{R})$ . This implies that any finite dimensional nontrivial irreducible  $\text{Sp}_{2g}(\mathbb{R})$ -representation  $V$  must also be an irreducible  $\Gamma$ -representation; indeed, if  $V'$  was a nontrivial proper  $\Gamma$ -submodule of  $V$ , then the subgroup of  $\text{Sp}_{2g}(\mathbb{R})$  preserving  $V'$  would be a proper subvariety of  $\text{Sp}_{2g}(\mathbb{R})$  containing  $\Gamma$ . Recall that the ring of coinvariants  $V_\Gamma$  of  $V$  under  $\Gamma$  is the quotient  $V/K$ , where  $K = \langle x - g(x) \mid x \in V, g \in \Gamma \rangle$ . Since  $K \neq 0$ , we can apply Schur's lemma to deduce that  $K = V$ , i.e. that  $V_\Gamma = 0$ . We record this fact as the following theorem.

**Theorem 4.2.** *For some  $g \geq 1$ , let  $\Gamma < \text{Sp}_{2g}(\mathbb{Z})$  satisfy  $[\text{Sp}_{2g}(\mathbb{Z}) : \Gamma] < \infty$  and let  $V$  be a nontrivial irreducible  $\text{Sp}_{2g}(\mathbb{R})$ -representation. Then  $V_\Gamma = 0$ .*




Figure 1: a–f. Curves needed for proof of Lemma 4.3

## 4.2 Two preliminary lemmas

We will need two lemmas. The first is the following, which slightly generalizes a theorem of Johnson [8].

**Lemma 4.3.** *For  $g \geq 3$ , we have  $\mathcal{S}_{g,b}^p / \mathcal{K}_{g,b}^p \cong (\wedge^3 H) / H \oplus H^{b+p}$ , where  $H = H_1(\Sigma_g; \mathbb{Z})$ .*

*Proof.* Since  $\mathcal{K}_{g,b}^p$  contains all twists about boundary curves, we can assume that  $b = 0$ .

Building on work of Johnson [9], Hain [5] proved that

$$H_1(\mathcal{S}_g^p; \mathbb{R}) \cong (\wedge^3 H_{\mathbb{R}}) / H_{\mathbb{R}} \oplus H_{\mathbb{R}}^p,$$

where  $H_{\mathbb{R}} = H_1(\Sigma_g; \mathbb{R})$ . Also, Johnson [9, Lemma 2] proved that for  $x \in \mathcal{K}_g^p$ , we have  $[x]_{\mathcal{S}_g^p} = 0$  (Johnson only considered the case where  $p = 0$ , but his argument works in general). It follows that

$$H_1(\mathcal{S}_g^p / \mathcal{K}_g^p; \mathbb{R}) \cong (\wedge^3 H_{\mathbb{R}}) / H_{\mathbb{R}} \oplus H_{\mathbb{R}}^p. \quad (1)$$

We will prove the lemma by induction on  $p$ . The base case  $p = 0$  is a theorem of Johnson [8]. Assume now that  $p > 0$  and that the lemma is true for all smaller  $p$ . Fixing a puncture  $*$  of  $\Sigma_g^p$ , work of Birman [1] and Johnson [9] gives an exact sequence

$$1 \longrightarrow \pi_1(\Sigma_g^{p-1}, *) \longrightarrow \mathcal{S}_g^p \longrightarrow \mathcal{S}_g^{p-1} \longrightarrow 1,$$

where the map  $\mathcal{S}_g^p \rightarrow \mathcal{S}_g^{p-1}$  comes from “forgetting the puncture  $*$ ”. Quotienting out by  $\mathcal{K}_g^p$ , we obtain an exact sequence

$$1 \longrightarrow \pi_1(\Sigma_g^{p-1}, *) / (\pi_1(\Sigma_g^{p-1}, *) \cap \mathcal{K}_g^p) \longrightarrow \mathcal{S}_g^p / \mathcal{K}_g^p \longrightarrow \mathcal{S}_g^{p-1} / \mathcal{K}_g^{p-1} \longrightarrow 1.$$

By induction, we have

$$\mathcal{S}_g^{p-1} / \mathcal{K}_g^{p-1} \cong (\wedge^3 H) / H \oplus H^{p-1}.$$

Set  $A = \pi_1(\Sigma_g^{p-1}, *) / (\pi_1(\Sigma_g^{p-1}, *) \cap \mathcal{K}_g^p)$ . We will prove that  $A$  is a quotient of  $H$ . We will then be able to conclude that  $\mathcal{S}_g^{p-1} / \mathcal{K}_g^{p-1}$  acts trivially on  $A$ , so  $\mathcal{S}_g^p / \mathcal{K}_g^p$  is the abelian group

$$(\wedge^3 H) / H \oplus H^{p-1} \oplus A.$$

Using (1), a simple dimension count will then imply that  $A$  cannot be a proper quotient of  $H$ , and the lemma will follow.

The element of  $\mathcal{S}_g^p$  corresponding to  $\delta \in \pi_1(\Sigma_g^{p-1}, *)$  “drags”  $*$  around  $\delta$ . As shown in Figures 1.a–b, a simple closed curve  $\gamma \in \pi_1(\Sigma_g^{p-1}, *)$  corresponds to  $T_{\gamma_1} T_{\gamma_2}^{-1} \in \mathcal{S}_g^p$ , where  $\gamma_1$  and  $\gamma_2$  are the boundary components of a regular neighborhood of  $\gamma$ . In particular, if  $\gamma$  is a simple closed separating curve, then as shown in Figures 1.c–d, the corresponding element of  $\mathcal{S}_g^p$  is a product of

separating twists. Since  $[\pi_1(\Sigma_g^{p-1}, *), \pi_1(\Sigma_g^{p-1}, *)]$  is generated by simple closed separating curves (see, e.g., [12, Lemma A.1]), we deduce that  $[\pi_1(\Sigma_g^{p-1}, *), \pi_1(\Sigma_g^{p-1}, *)] \subset \pi_1(\Sigma_g^{p-1}, *) \cap \mathcal{K}_g^p$ . Thus  $A = \pi_1(\Sigma_g^{p-1}, *) / (\pi_1(\Sigma_g^{p-1}, *) \cap \mathcal{K}_g^p)$  is a quotient of  $H_1(\Sigma_g^{p-1}; \mathbb{Z})$ . Finally, as shown in Figures 1.e–f, all simple closed curves that are homotopic into punctures are also contained in  $\pi_1(\Sigma_g^{p-1}, *) \cap \mathcal{K}_g^p$ , so we conclude that  $A$  is a quotient of  $H = H_1(\Sigma_g; \mathbb{Z})$ , as desired.  $\square$

For the second lemma, define  $Q_{g,b}^p = \text{Mod}_{g,b}^p / \mathcal{K}_{g,b}^p$ .

**Lemma 4.4.** *For some  $g \geq 3$ , let  $Q' < Q_{g,b}^p$  satisfy  $[Q_{g,b}^p : Q'] < \infty$ . Then  $H_1(Q'; \mathbb{R}) = 0$ .*

*Proof.* Restricting the short exact sequence

$$1 \longrightarrow \mathcal{S}_{g,b}^p / \mathcal{K}_{g,b}^p \longrightarrow Q_{g,b}^p \longrightarrow \text{Sp}_{2g}(\mathbb{Z}) \longrightarrow 1$$

to  $Q'$ , we obtain a short exact sequence

$$1 \longrightarrow B \longrightarrow Q' \longrightarrow \overline{Q}' \longrightarrow 1,$$

where  $B$  and  $\overline{Q}'$  are finite index subgroups of  $\mathcal{S}_{g,b}^p / \mathcal{K}_{g,b}^p$  and  $\text{Sp}_{2g}(\mathbb{Z})$ , respectively. The last 3 terms of the associated 5-term exact sequence are

$$(H_1(B; \mathbb{R}))_{\overline{Q}'} \longrightarrow H_1(Q'; \mathbb{R}) \longrightarrow H_1(\overline{Q}'; \mathbb{R}) \longrightarrow 0.$$

By Theorem 4.1, we have  $H_1(\overline{Q}'; \mathbb{R}) = 0$ . Letting  $H = H_1(\Sigma_g; \mathbb{Z})$ , Lemma 4.3 says that

$$\mathcal{S}_{g,b}^p / \mathcal{K}_{g,b}^p \cong (\wedge^3 H) / H \oplus H^{b+p}.$$

Since  $B$  is a finite-index subgroup of  $\mathcal{S}_{g,b}^p / \mathcal{K}_{g,b}^p$ , we get that  $B$  is itself abelian and

$$H_1(B; \mathbb{R}) \cong B \otimes \mathbb{R} \cong (\mathcal{S}_{g,b}^p / \mathcal{K}_{g,b}^p) \otimes \mathbb{R} \cong (\wedge^3 H_{\mathbb{R}}) / H_{\mathbb{R}} \oplus H_{\mathbb{R}}^{b+p},$$

where  $H_{\mathbb{R}} = H_1(\Sigma_g; \mathbb{R})$ . Both  $(\wedge^3 H_{\mathbb{R}}) / H_{\mathbb{R}}$  and  $H_{\mathbb{R}}$  are nontrivial finite-dimensional irreducible representations of  $\text{Sp}_{2g}(\mathbb{R})$ , so Theorem 4.2 implies that  $(H_1(B; \mathbb{R}))_{\overline{Q}'} = 0$ , and we are done.  $\square$

### 4.3 The proof of Theorem B

The last 3 terms of the 5-term exact sequence associated to the short exact sequence

$$1 \longrightarrow \Gamma \cap \mathcal{K}_{g,b}^p \longrightarrow \Gamma \longrightarrow \Gamma / (\Gamma \cap \mathcal{K}_{g,b}^p) \longrightarrow 1$$

are

$$(H_1(\Gamma \cap \mathcal{K}_{g,b}^p; \mathbb{R}))_{\Gamma / (\Gamma \cap \mathcal{K}_{g,b}^p)} \xrightarrow{i} H_1(\Gamma; \mathbb{R}) \longrightarrow H_1(\Gamma / (\Gamma \cap \mathcal{K}_{g,b}^p); \mathbb{R}) \longrightarrow 0.$$

By assumption,  $[\Gamma \cap \mathcal{K}_{g,b}^p : K(\Gamma)] < \infty$ , so Lemma 2.1 implies that the map  $H_1(K(\Gamma); \mathbb{R}) \rightarrow H_1(\Gamma \cap \mathcal{K}_{g,b}^p; \mathbb{R})$  is surjective. Since  $K(\Gamma)$  is generated by powers of twists, Theorem A allows us to deduce that  $i = 0$ . Also,  $\Gamma / (\Gamma \cap \mathcal{K}_{g,b}^p)$  is a finite-index subgroup of  $Q_{g,b}^p$ , so Lemma 4.4 implies that  $H_1(\Gamma / (\Gamma \cap \mathcal{K}_{g,b}^p); \mathbb{R}) = 0$ , and we are done.

## References

- [1] J. S. Birman, Mapping class groups and their relationship to braid groups, *Comm. Pure Appl. Math.* **22** (1969), 213–238.
- [2] M. Boggi, Fundamental groups of moduli stacks of stable curves of compact type, *Geom. Topol.* **13** (2009), 247–276.
- [3] M. Bridson, Semisimple actions of mapping class groups on CAT(0) spaces, to appear.
- [4] K. S. Brown, *Cohomology of groups*, Corrected reprint of the 1982 original, Springer, New York, 1994.
- [5] R. M. Hain, Torelli groups and geometry of moduli spaces of curves, in *Current topics in complex algebraic geometry (Berkeley, CA, 1992/93)*, 97–143, Cambridge Univ. Press, Cambridge.
- [6] N. V. Ivanov, Fifteen problems about the mapping class groups, in *Problems on mapping class groups and related topics*, 71–80, Proc. Sympos. Pure Math., 74, Amer. Math. Soc., Providence, RI.
- [7] D. Johnson, An abelian quotient of the mapping class group  $\mathcal{S}_g$ , *Math. Ann.* **249** (1980), no. 3, 225–242.
- [8] D. Johnson, The structure of the Torelli group. II. A characterization of the group generated by twists on bounding curves, *Topology* **24** (1985), no. 2, 113–126.
- [9] D. Johnson, The structure of the Torelli group. III. The abelianization of  $\mathcal{T}$ , *Topology* **24** (1985), no. 2, 127–144.
- [10] M. Korkmaz, Low-dimensional homology groups of mapping class groups: a survey, *Turkish J. Math.* **26** (2002), no. 1, 101–114.
- [11] J. D. McCarthy, On the first cohomology group of cofinite subgroups in surface mapping class groups, *Topology* **40** (2001), no. 2, 401–418.
- [12] A. Putman, Cutting and pasting in the Torelli group, *Geom. Topol.* **11** (2007), 829–865.
- [13] R. J. Zimmer, *Ergodic theory and semisimple groups*, Birkhäuser, Basel, 1984.

Department of Mathematics; MIT, 2-306  
77 Massachusetts Avenue  
Cambridge, MA 02139-4307  
E-mail: andyp@math.mit.edu