

# MIYAOKA-YAU INEQUALITY FOR MINIMAL PROJECTIVE MANIFOLDS OF GENERAL TYPE

YUGUANG ZHANG

ABSTRACT. In this short note, we prove the Miyaoka-Yau inequality for minimal projective  $n$ -manifolds of general type by using Kähler-Ricci flow.

## 1. INTRODUCTION

If  $M$  is a projective  $n$ -manifold with ample canonical bundle  $\mathcal{K}_M$ , there exists a Kähler-Einstein metric  $\omega$  with negative scalar curvature by Yau's theorem on the Calabi conjecture ([14]), which was obtained by Aubin independently ([1]). As a consequence, there is an inequality for Chern numbers, the Miyaoka-Yau inequality,

$$(1.1) \quad \left(\frac{2(n+1)}{n}c_2(M) - c_1^2(M)\right) \cdot (-c_1(M))^{n-2} \geq 0,$$

where  $c_1(M)$  and  $c_2(M)$  are the first and the second Chern classes of  $M$  (c.f. [13]). Furthermore, if the equality in (1.1) holds, the Kähler-Einstein metric  $\omega$  is a complex hyperbolic metric, i.e. the holomorphic sectional curvature of  $\omega$  is a negative constant. If  $n = 2$ , (1.1) even holds for algebraic surfaces of general type (c.f. [5], [8], [9]), which may not admit any Kähler-Einstein metric. In [11], the inequality (1.1) is proved for any dimensional minimal projective manifold of general type by using singular Kähler-Einstein metrics. In this short note, we give a different proof of (1.1) for minimal projective  $n$ -manifolds of general type by using Kähler-Ricci flow, and study the extremal case of (1.1).

Let  $M$  be a minimal projective manifold of general type with  $\dim_{\mathbb{C}} M = n \geq 2$ . The canonical bundle  $\mathcal{K}_M$  of  $M$  is big, and semi-ample, i.e.  $\mathcal{K}_M^r > 0$ , and, for a positive integer  $m \gg 1$ , the linear system  $|m\mathcal{K}_M|$  is base point free (as quoted in [10]). For  $m \gg 1$ , the complete linear system  $|m\mathcal{K}_M|$  defines a holomorphic map  $\Phi : M \rightarrow \mathbb{C}P^N$ , which is birational onto its image  $M_{can}$ .  $M_{can}$  is called the canonical model of  $M$ , and  $\Phi$  is called the contraction map. Note that  $M$  may not admit any Kähler-Einstein metric. The Kähler-Ricci flow is an evolution equation of a family of Kähler metrics  $\omega_t$ ,  $t \in [0, T)$ , on  $M$ ,

$$(1.2) \quad \partial_t \omega_t = -Ric(\omega_t) - \omega_t,$$

where  $Ric(\omega_t)$  is the Ricci form of  $\omega_t$ . By [10] [12] [3] and [15], for any Kähler metric as initial metric, the solution  $\omega_t$  of the Kähler-Ricci flow equation exists for all time  $t \in [0, \infty)$ , and the scalar curvature of  $\omega_t$  is uniformly bounded. Thus we can prove (1.1) by using the technique developed in [6], where a Hitchin-Thorpe type inequality was proved for 4-manifolds which admit a long time solution to a normalized Ricci flow equation with bounded scalar curvature. Before proving the Miyaoka-Yau inequality, we show that the  $L^2$ -norm of the Einstein tensor tends to zero along a subsequence of a solution of the Kähler-Ricci flow equation (1.2).

**Theorem 1.1.** *Let  $M$  be a minimal projective manifold of general type with  $\dim_{\mathbb{C}} M = n \geq 2$ , and  $\omega_t$ ,  $t \in [0, \infty)$ , be a solution of the Kähler-Ricci flow equation (1.2). Then there exists a sequence of times  $t_k \rightarrow \infty$ , when  $k \rightarrow \infty$ , such that*

$$\lim_{k \rightarrow \infty} \int_M |\rho_{t_k}|^2 \omega_{t_k}^n = 0,$$

where  $\rho_{t_k} = \text{Ric}_{t_k} - \frac{R_{t_k}}{n} \omega_{t_k}$  denotes the Einstein tensor of  $\omega_{t_k}$ , and  $R_{t_k}$  denotes the scalar curvature of  $\omega_{t_k}$ .

As a corollary of this theorem, we obtain the Miyaoka-Yau inequality for minimal projective manifolds of general type.

**Corollary 1.2.** *If  $M$  is a minimal projective manifold of general type with  $\dim_{\mathbb{C}} M = n \geq 2$ , then*

$$\left(\frac{2(n+1)}{n} c_2(M) - c_1^2(M)\right) \cdot (-c_1(M))^{n-2} \geq 0.$$

Furthermore, if the equality holds, there is a complex hyperbolic metric on the smooth part  $M_0$  of the canonical model  $M_{can}$  of  $M$ .

**Acknowledgement:** The author thanks Zhou Zhang for explaining [15], and sending him the paper [12]. Thanks also goes to referees for their suggestions of improving the present paper. The author also thanks Valentino Tosatti for sending him [11].

## 2. PROOF OF THEOREM 1.1

Let  $M$  be a minimal projective manifold of general type with  $\dim_{\mathbb{C}} M = n \geq 2$ ,  $M_{can}$  be the canonical model of  $M$ , and  $\Phi : M \rightarrow M_{can}$  be the contraction map. Consider the Kähler-Ricci flow equation on  $M$ ,

$$(2.1) \quad \partial_t \omega_t = -\text{Ric}(\omega_t) - \omega_t,$$

with initial metric  $\omega_0$ . In [7], the short time existence of the solution of (2.1) is proved. Then, in [10] [12] and [3], it is proved that the solution  $\omega_t$  of (2.1) exists for all time, i.e.  $t \in [0, +\infty)$ , and, there exists a unique semi-positive current  $\omega_\infty$  on  $M$ , which satisfies that

- (1)  $\omega_\infty$  represents  $-2\pi c_1(M)$ .
- (2)  $\omega_\infty$  is a smooth Kähler-Einstein metric with negative scalar curvature on  $\Phi^{-1}(M_0)$ , where  $M_0$  is the smooth part of  $M_{can}$ .
- (3) On any compact subset  $K \subset \Phi^{-1}(M_0)$ ,  $\omega_t$   $C^\infty$ -converges to  $\omega_\infty$  when  $t \rightarrow \infty$ .

In [15], it is shown that there is a constant  $C > 0$  depending only on  $\omega_0$  such that

$$(2.2) \quad |R_t| < C,$$

where  $R_t$  is the scalar curvature of  $\omega_t$ .

First, we need evolution equations for volume forms and scalar curvatures as follows,

$$(2.3) \quad \partial_t \omega_t^n = -(R_t + n) \omega_t^n, \quad \text{and}$$

$$(2.4) \quad \partial_t R_t = \Delta_t R_t + |Ric_t|^2 + R_t = \Delta_t R_t + |Ric_t^\circ|^2 - (R_t + n),$$

where  $Ric_t^\circ = Ric_t + \omega_t$ , and  $|Ric_t^\circ|^2 = |Ric_t|^2 + 2R_t + n$  (c.f. Lemma 2.38 in [4]).

**Lemma 2.1.** *There are two constants  $t_0 > 0$  and  $c > 0$  independent of  $t$  such that, for  $t > t_0$ ,*

$$\check{R}_t = \inf_{x \in M} R_t(x) \leq -n + e^{-t}c < -\frac{n}{2} < 0.$$

*Proof.* If we define  $\alpha_t = [\omega_t] \in H^{1,1}(M, \mathbb{R})$ , from (2.1) we have

$$\partial_t \alpha_t = -2\pi c_1(M) - \alpha_t, \quad \text{and}$$

$$(2.5) \quad \alpha_t = -2\pi c_1(M) + e^{-t}(2\pi c_1(M) + \alpha_0).$$

Thus

$$(2.6) \quad [\omega_\infty] = \alpha_\infty = \lim_{t \rightarrow \infty} \alpha_t = -2\pi c_1(M).$$

Since

$$\check{R}_t \int_M \omega_t^n \leq \int_M R_t \omega_t^n = n \int_M Ric_t \wedge \omega_t^{n-1} = n 2\pi c_1(M) \cdot \alpha_t^{n-1},$$

we obtain

$$\check{R}_t \leq n \frac{2\pi c_1(M) \cdot \alpha_t^{n-1}}{\alpha_t^n} = n \frac{2\pi c_1(M) \cdot \alpha_t^{n-1}}{-2\pi c_1(M) \cdot \alpha_t^{n-1} + e^{-t}(2\pi c_1(M) + \alpha_0) \cdot \alpha_t^{n-1}} = \frac{-n}{1 + e^{-t}A_t},$$

where  $A_t = -\frac{(2\pi c_1(M) + \alpha_0) \cdot \alpha_t^{n-1}}{2\pi c_1(M) \cdot \alpha_t^{n-1}}$ . Note that  $(-c_1(M))^n > 0$ . Thus there is a  $t_1 > 0$  such that, if  $t > t_1$ ,  $A_t < \left| \frac{(\alpha_\infty + \alpha_0) \cdot \alpha_\infty^{n-1}}{\alpha_\infty^n} \right| + 1 = A$ , and we obtain that

$$\check{R}_t \leq \frac{-n}{1 + e^{-t}A} < -n + e^{-t}c,$$

where  $c = -n\left(\frac{A}{1 + e^{-t_1}A}\right)$ . By taking  $t_0 > t_1$  such that  $e^{-t_0}c < \frac{n}{2}$ , we obtain the conclusion.  $\square$

**Lemma 2.2.**

$$\int_0^\infty \int_M |R_t + n| \omega_t^n dt < \infty.$$

*Proof.* By (2.4) and the maximal principle,  $\partial_t \check{R}_t \geq -(\check{R}_t + n)$ , and so,

$$(2.7) \quad n + \check{R}_t \geq C e^{-t},$$

for a constant  $C$  independent of  $t$ . Note that, by Lemma 2.1, (2.7) and (2.5), when  $t > t_0$ ,

$$\begin{aligned} \int_M |R_t + n| \omega_t^n &\leq \int_M (R_t - \check{R}_t) \omega_t^n + \int_M |n + \check{R}_t| \omega_t^n \\ &\leq \int_M (R_t + n) \omega_t^n + 2 \int_M |n + \check{R}_t| \omega_t^n \\ &\leq \int_M (R_t + n) \omega_t^n + C_3 e^{-t} \\ &= n(2\pi c_1 \cdot \alpha_t^{n-1} + \alpha_t^n) + C_3 e^{-t} \\ &= n e^{-t}(2\pi c_1 + \alpha_0) \cdot \alpha_t^{n-1} + C_3 e^{-t} \\ &\leq C_4 e^{-t}, \end{aligned}$$

for two constants  $C_3$  and  $C_4$  independent of  $t$ . Thus

$$\int_0^\infty \int_M |R_t + n|\omega_t^n dt = \int_0^{t_0} \int_M |R_t + n|\omega_t^n dt + \int_{t_0}^\infty \int_M |R_t + n|\omega_t^n dt < \infty.$$

□

*Proof of Theorem 1.1.* From (2.4), (2.3), (2.6), (2.2), and Lemma 2.2, we obtain

$$\begin{aligned} \int_0^\infty \int_M |Ric^\circ_t|^2 \omega_t^n dt &= \int_0^\infty \int_M \left(\frac{\partial}{\partial t} R_t\right) \omega_t^n dt + \int_0^\infty \int_M (R_t + n) \omega_t^n dt \\ &= \int_0^\infty \frac{\partial}{\partial t} \left(\int_M R_t \omega_t^n\right) dt + \int_0^\infty \int_M (R_t + 1)(R_t + n) \omega_t^n dt \\ &\leq n\alpha_\infty^n - \int_M R_0 \omega_0^n + C \int_0^\infty \int_M |R_t + n|\omega_t^n dt \\ &< \infty. \end{aligned}$$

If  $\rho_t = Ric_t - \frac{R_t}{n}\omega_t$  is the Einstein tensor of  $\omega_t$ , then  $|\rho_t|^2 = |Ric^\circ_t|^2 - \frac{1}{n}(R_t + n)^2$ , and, from the above estimation,

$$\int_0^\infty \int_M |\rho_t|^2 \omega_t^n dt \leq \int_0^\infty \int_M |Ric^\circ_t|^2 \omega_t^n dt < \infty.$$

Thus there is a sequence  $t_k \rightarrow \infty$  such that

$$\lim_{k \rightarrow \infty} \int_M |\rho_{t_k}|^2 \omega_{t_k}^n = 0.$$

□

*Proof of Corollary 1.2.* Note that the Kähler curvature tensor has a decomposition

$$Rm_t = \frac{R_t}{2n^2} \omega_t \otimes \omega_t + \frac{1}{n} \omega_t \otimes \rho_t + \frac{1}{n} \rho_t \otimes \omega_t + B_t$$

(c.f. (2.63) and (2.38) in [2]). By Chern-Weil theory,

$$\left(\frac{2(n+1)}{n}c_2(M) - c_1^2(M)\right) \cdot [\omega_t]^{n-2} = \frac{(n-2)!}{4\pi^2 n!} \int_M \left(\frac{n+1}{n}|B_{0,t}|^2 - \frac{(n^2-2)}{n^2}|\rho_t|^2\right) \omega_t^n$$

(c.f. (2.82a) and (2.67) in [2]), where  $B_{0,t} = B_t - \frac{tr B_t}{n^2-1}Id$  is the tensor given by (2.64) in [2] corresponding to  $\omega_t$ . By Theorem 1.1, there is a sequence  $t_k \rightarrow \infty$  such that

$$\lim_{k \rightarrow \infty} \int_M |\rho_{t_k}|^2 \omega_{t_k}^n = 0.$$

Hence

$$\begin{aligned} \left(\frac{2(n+1)}{n}c_2(M) - c_1^2(M)\right) \cdot (-2\pi c_1(M))^{n-2} &= \left(\frac{2(n+1)}{n}c_2(M) - c_1^2(M)\right) \cdot [\omega_\infty]^{n-2} \\ &= \lim_{k \rightarrow \infty} \left(\frac{2(n+1)}{n}c_2(M) - c_1^2(M)\right) \cdot [\omega_{t_k}]^{n-2} \\ &= \lim_{k \rightarrow \infty} \frac{(n-2)!}{4\pi^2 n!} \int_M \left(\frac{n+1}{n}|B_{0,t_k}|^2\right) \omega_{t_k}^n \\ &\geq 0. \end{aligned}$$

If the equality holds, on any compact subset  $K \subset \Phi^{-1}(M_0)$ ,

$$\int_K |B_{0,\infty}|^2 \omega_\infty^n \leq \lim_{k \rightarrow \infty} \int_M |B_{0,t_k}|^2 \omega_{t_k}^n = 0,$$

by the smooth convergence of  $\omega_t$  to  $\omega_\infty$ . Thus  $B_{0,\infty} \equiv 0$ . Since  $\omega_\infty$  is a Kähler-Einstein metric with negative scalar curvature on  $\Phi^{-1}(M_0)$ , the holomorphic sectional curvature is a negative constant by Section 2.66 in [2], i.e.  $\omega_\infty$  is a complex hyperbolic metric.  $\square$

## REFERENCES

1. T. Aubin, *Equations du type Monge-Ampere sur les variete kaehleriennes compactes*. C. R. Acad. Paris. **283** (1976), 119-121.
2. A. L. Besse, *Einstein manifolds*. Ergebnisse der Math. Springer-Verlag, Berlin-New York 1987.
3. P.Cascini, P.La Nave, *Kähler-Ricci flow and the minimal model program for projective varieties*. arXiv:math/0603064.
4. B.Chow, S.C.Chu, D.Glickenstein, C.Guenther, J.Isenberg, T.Ivey, D.Knopf, P.Lu, F.Luo, L.Ni, *The Ricci Flow: Techniques and Applications, Part I: Geometric Aspect*. Mathematical Surveys and Monographs, Vol. 135, (2007).
5. S.Y.Cheng, S.T.Yau, *Inequality between Chern numbers of singular Kähler surfaces and characterization of orbit space of discrete group of  $SU(2, 1)$* . Contemporary Math., **49** (1986), 31-43.
6. F. Fang, Y. Zhang and Z. Zhang, *Non-singular solutions to the normalized Ricci flow equation*. Math. Ann. **340** (2008), 647-674.
7. R. Hamilton, *Three-manifolds with positive Ricci curvature*. J. Diff. Geom., **17** (1982), 255-306.
8. R.Kobayashi, *Einstein-Kähler V-metrics on open Satake V-surfaces with isolated quotient singularities*. Math. Ann. **272**, no.3 (1985), 385-398.
9. Y.Miyaoka, *On the Chern numbers of surfaces of general type*. Inv. math., **32** (1977), 225-237.
10. H.Tsuji, *Existence and degeneration of Kähler-Einstein metrics on minimal algebraic varieties of general type*. Math. Ann. **281** (1988), 123-133.
11. H.Tsuji, *Stability of tangent bundles of minimal algebraic varieties*. Topology, Vol.27 **4**, (1988), 429-442.
12. G.Tian, Z.Zhang, *On the Kähler-Ricci flow on projective manifolds of general type*. Chinese Annals of Mathematics B, **27** (2006), 179-192.
13. S.T.Yau, *Calabi's conjecture and some new results in algebraic geometry*. Proc. Natl. Acad. Sci. USA **74**, (1977), 1789-1799.
14. S.T.Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampere equations. I*. Comm. Pure Appl. Math. **31**, (1978), 339-411.
15. Z.Zhang, *Scalar Curvature Bound for Kähler-Ricci Flows over Minimal Manifolds of General Type*. arXiv:math/0801.32481.

DEPARTMENT OF MATHEMATICS, CAPITAL NORMAL UNIVERSITY, BEIJING, P.R.CHINA

*Current address:* Department of Mathematical Sciences, Korea Advanced Institute of Science and Technology, Daejeon, Republic of Korea

*E-mail address:* zhangyuguang76@yahoo.com