

EQUATIONS RESOLVING A CONJECTURE OF RADO ON PARTITION REGULARITY

BORIS ALEXEEV AND JACOB TSIMERMAN

ABSTRACT. The *degree of regularity* of a linear equation L is the least number of colors in a coloring of the positive integers with no monochromatic solution to L . Richard Rado conjectured in his 1933 thesis that for every positive integer k , there exists a linear equation that has degree of regularity k . We prove this conjecture by showing that the equation $\sum_{i=1}^{k-1} \frac{2^i}{2^i-1} x_i = \left(-1 + \sum_{i=1}^{k-1} \frac{2^i}{2^i-1}\right) x_0$ has this property.

This conjecture is part of problem E14 in Richard Guy's book "Unsolved problems in number theory".

In 1916, Schur [Sch16] proved that in any coloring of the positive integers with finitely many colors, there is a monochromatic solution to $x + y = z$. In 1927, van der Waerden [vdW27] proved his celebrated theorem that every finite coloring of the positive integers contains arbitrarily long monochromatic arithmetic progressions. In his famous 1933 thesis, Richard Rado [Rad33] generalized these results by classifying the systems of linear equations with monochromatic solutions in every finite coloring. His thesis also contained conjectures regarding equations that *do* have a finite coloring with no monochromatic solutions.

Definition. The *degree of regularity* of a linear equation L is the least number of colors in a coloring of the positive integers with no monochromatic solution to L .

Remark. Some authors require that the values of the variables be distinct in solutions to L . We follow Rado and Guy in not including this extra condition.

Conjecture (Rado [Rad33, Guy04]). For every positive integer k , there exists a linear equation that has degree of regularity k .

Fox and Radoičić [FR05] conjectured that the family of equations M_k given by $\sum_{i=0}^{k-2} 2^i x_i = 2^{k-1} x_{k-1}$ has this property. We prove Rado's conjecture by using the related family of equations,

$$\sum_{i=1}^{k-1} \frac{2^i}{2^i-1} x_i = \left(-1 + \sum_{i=1}^{k-1} \frac{2^i}{2^i-1}\right) x_0,$$

which we denote by L_k .

Theorem 1. *The degree of regularity of the equation L_k is k .*

Remark. This result and its proof carry over to colorings of the nonzero rationals.

Proof. We first use the power of 2-adic valuations to prove that there exists a k -coloring with no monochromatic solutions to L_k . This idea was proposed by Fox and Radoičić for the equation M_k ; later, Alexeev, Fox, and Graham [AFG07] proved that these colorings actually were minimal if $k \leq 7$, but only conjectured the result in general.

Date: November 18, 2018.

2000 Mathematics Subject Classification. 05D10.

Key words and phrases. Colorings, partition regularity, Ramsey theory.

If r is a nonzero rational number, let $\text{ord}_2 r$ denote the 2-adic valuation of r , that is, the unique m such that $r = 2^m \frac{a}{b}$ for odd a and b ; also, let $\text{ord}_2 0 = \infty$. Note that the 2-adic valuation of the coefficient of x_i in L_k is i . This is immediate if $i > 0$ while in the case of x_0 , this follows from the (non-Archimedean) 2-adic triangle inequality¹ because -1 has valuation 0 while the rest of the terms in the summation have positive valuation. We claim this implies that the k -coloring $C_k(r) = \text{ord}_2 r \bmod k$ has no monochromatic solutions to L_k .

Assume to the contrary that C_k admits a monochromatic solution to L_k . Then the terms of L_k would be a collection of numbers with distinct 2-adic valuations that sum to zero, which is impossible by the 2-adic triangle inequality.

We now show there is no coloring to L_k with fewer than k colors. Indeed, in *any* coloring with no monochromatic solutions to L_k , the color of x and $2^j x$ must be different if $0 < j < k$. (In the referenced literature, the number 2^j is thus said to be a *forbidden ratio*.) If there were an x so that x and $2^j x$ were the same color, then

$$x_i = \begin{cases} x & \text{if } i = j, \\ 2^j x & \text{if } i \neq j, \end{cases}$$

would be a monochromatic solution to L_k ; all but the j^{th} terms cancel, leaving $\frac{2^j}{2^j-1}x = \left(-1 + \frac{2^j}{2^j-1}\right)2^j x$. This implies that the k numbers $1, 2, 4, \dots, 2^{k-1}$ are colored with distinct colors. \square

ACKNOWLEDGMENTS

The authors wish to thank Owen Biesel for helpful comments on the exposition.

REFERENCES

- [AFG07] Boris Alexeev, Jacob Fox, and Ron Graham, *On minimal colorings without monochromatic solutions to a linear equation*, Combinatorial number theory, de Gruyter, Berlin, 2007, pp. 1–22.
- [FR05] Jacob Fox and Rados Radoičić, *The axiom of choice and the degree of regularity of equations over the reals*, preprint, Dec 2005.
- [Guy04] Richard K. Guy, *Unsolved problems in number theory*, third ed., Problem Books in Mathematics, ch. E14, p. 328, Springer-Verlag, New York, 2004.
- [Rad33] Richard Rado, *Studien zur Kombinatorik*, Math. Z. **36** (1933), no. 1, 424–470.
- [Sch16] Issai Schur, *Über die Kongruenz $x^m + y^m \equiv z^m \pmod{p}$* , Jahresber Deutsch. Math. Verein. **25** (1916), 114–117.
- [vdW27] Bartel Leendert van der Waerden, *Beweis einer Baudet'schen Vermutung*, Nieuw Arch. Wiskunde **15** (1927), 212–216.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, FINE HALL, WASHINGTON ROAD, PRINCETON, NJ 08544-1000

E-mail address: balixeev@math.princeton.edu

E-mail address: jtsimerm@math.princeton.edu

¹Namely, $\text{ord}_2(x + y) \geq \min(\text{ord}_2 x, \text{ord}_2 y)$. Moreover, it follows that equality holds if $\text{ord}_2(x) \neq \text{ord}_2(y)$.