

# Maximum Matchings in Random Bipartite Graphs and the Space Utilization of Cuckoo Hashtables

Alan Frieze\*, Páll Melsted  
 Department of Mathematical Sciences  
 Carnegie Mellon University  
 Pittsburgh PA 15213  
 U.S.A.

## Abstract

We study the the following question in Random Graphs. We are given two disjoint sets  $L, R$  with  $|L| = n = \alpha m$  and  $|R| = m$ . We construct a random graph  $G$  by allowing each  $x \in L$  to choose  $d$  random neighbours in  $R$ . The question discussed is as to the size  $\mu(G)$  of the largest matching in  $G$ . When considered in the context of Cuckoo Hashing, one key question is as to when is  $\mu(G) = n$  **whp**? We answer this question exactly when  $d$  is at least four. We also establish a precise threshold for when Phase 1 of the Karp-Sipser Greedy matching algorithm suffices to compute a maximum matching **whp**.

## 1 Introduction

For a graph  $G$  we let  $\mu(G)$  denote the size of the maximum matching in  $G$ . In essence this paper provides an analysis of  $\mu(G)$  in the following model of a random bipartite graph. We have two disjoint sets  $L, R$  where  $L = [n], R = [m]$  where  $n = \alpha m$ . Each  $v \in L$  independently chooses  $d$  random vertices of  $R$  as neighbours. Our assumptions are that  $\alpha > 0, d \geq 3$  are fixed and  $n \rightarrow \infty$ . One motivation for this study comes from Cuckoo Hashing.

Briefly each one of  $n$  items  $x \in L$  has  $d$  possible locations  $h_1(x), h_2(x), \dots, h_d(x) \in R$ , where  $d$  is typically a small constant and the  $h_i$  are hash functions, typically assumed to behave as independent fully random hash functions. (See [19] for some justification of this assumption.) We are thus led to consider the bipartite graph  $G$  which has vertex set  $L \cup R$  and edge set  $\{(x, h_j(x)) : x \in L, j = 1, 2, \dots, d\}$ . Under the assumption that the hash functions are completely random we see that  $G$  has the same distribution as the random graph defined in the previous paragraph.

We assume each location can hold only one item. When an item  $x$  is inserted into the table, it can be placed immediately if one of its  $d$  locations is currently empty. If not, one of the items in its  $d$  locations must be displaced and moved to another of its  $d$  choices to make room for  $x$ . This item in turn may need to displace another item out of one its  $d$  locations. Inserting an item may require a sequence of moves, each maintaining the invariant that each item remains in one of its  $d$  potential locations, until no further evictions are needed. Thus having inserted  $k$  items, we have constructed a matching  $M$  of size  $k$  in  $G$ . Adding a  $(k + 1)$ 'th item is tantamount to constructing an augmenting path with respect to  $M$ . All  $n$  items will be insertable in this way iff  $G$  contains a matching of size  $n$ .

---

\*Supported in part by NSF Grant DMS0753472.

The case of  $d = 2$  choices is notably different from that for other values of  $d$  and the theory for the case where there are  $d = 2$  bucket choices for each item is well understood at this point [8, 18, 20]. We will therefore assume that  $d \geq 3$ .

We will now revert to the abstract question posed in first paragraph of the paper.

## 2 Definitions and Results

This question was studied to some extent by Fotakis, Pagh, Sanders and Spirakis [13]. They show in the course of their analysis of Cuckoo hashing that the following holds:

**Lemma 1** *Suppose that  $0 < \varepsilon < 1$  and  $d \geq 2(1 + \varepsilon) \log(e/\varepsilon)$ . Suppose also that  $m = (1 + \varepsilon)n$ . Then **whp**  $G$  contains a matching of size  $n$  i.e. a matching of  $L$  into  $R$ .*

□

In particular, if  $d = 3$  and  $m \approx 1.57n$  then Lemma 1 shows that there is a matching of  $L$  into  $R$  **whp**.

This lemma is not tight and recently Mitzenmacher et al [11] observed a connection with a result of Dubois and Mandler on Random 3-XORSAT [9] that enables one to essentially answer the question as to when  $\mu(G) \geq n$  for the case  $d = 3$ . More recently, Fountoulakis and Panagiotou [10] have established thresholds for when there is a matching of  $L$  into  $R$  **whp**, for all  $d \geq 3$ .

We begin with a simple observation that is the basis of the Karp-Sipser Algorithm [14, 2]. If  $v$  is a vertex of degree one in  $G$  and  $e$  is its unique incident edge, then there exists a maximum matching of  $G$  that includes  $e$ . Karp and Sipser exploited this via a simple greedy algorithm:

---

**Algorithm 1** Karp-Sipser Algorithm

---

```

1: procedure KSGREEDY( $G$ )
2:    $M \leftarrow \emptyset$ 
3:   while  $\Gamma \neq \emptyset$  do
4:     if  $\Gamma$  has vertices of degree one then
5:       Select a vertex  $\xi$  uniformly at random from the set of vertices of degree one
6:       Let  $e = (\xi, \eta)$  be the edge incident to  $\xi$ 
7:     else
8:       Select an edge  $e = (v, u)$  uniformly at random
9:     end if
10:     $M \leftarrow M \cup \{e\}$ 
11:     $\Gamma \leftarrow \Gamma \setminus \{\xi, \eta\}$ 
12:  end while
13:  return  $M$ 
14: end procedure

```

---

Phase 1 of the Karp-Sipser Algorithm ends and Phase 2 begins when the graph remaining has minimum degree at least two. So if  $\Gamma_1$  denotes the graph  $\Gamma$  remaining at the end of Phase 1 and  $\tau_1$  is the number of iterations involved in Phase 1 then

$$\mu(G) = \tau_1 + \mu(\Gamma_1). \tag{1}$$

Our approach to estimating  $\mu(G)$  is to (i) obtain an asymptotic expression for  $\tau_1$  that holds **whp** and then (ii) show that **whp**  $\Gamma_1$  has a (near) perfect matching and then apply (1).

We summarise our results as follows: Let  $z_1$  satisfy

$$z_1 = \frac{e^{z_1} - 1}{d - 1} \quad (2)$$

and let

$$\alpha_1 = \frac{z_1}{d(1 - e^{-z_1})^{d-1}}. \quad (3)$$

**Theorem 2** *If  $\alpha \leq \alpha_1$  then whp  $\mu(G) = \tau_1 = n$ .*

Thus **whp** Phase 1 of the Karp-Sipser Algorithm finds a (near) maximum matching if  $\alpha \leq \alpha_1$ . In particular, if  $d = 3$  then  $z_1 \approx 1.251$  and  $\alpha_1 \approx .818$  and thus  $m \approx 1.222n$  is enough for a matching of  $L$  into  $R$ .

Now consider larger  $\alpha$ . Let  $z^*$  be the largest non-negative solution to

$$\left(\frac{z}{\alpha d}\right)^{\frac{1}{d-1}} + e^{-z} - 1 = 0.$$

**Theorem 3** *If  $\alpha > \alpha_1$  then whp*

(a)  $z^* > 0$ .

(b)  $\tau_1 \sim n \left(1 - \left(\frac{z^*}{\alpha d}\right)^{\frac{d}{d-1}}\right)$ .

(c) *If  $d \geq 4$  then*

$$\mu(\Gamma_1) = \min\{|L_1|, |R_1|\} = \min\left\{n - \tau_1, (1 - (1 + z^*)e^{-z^*})m + o(m)\right\}. \quad (4)$$

*Here  $L_1 \subseteq L, R_1 \subseteq R$  are the two sides of the bipartition of  $\Gamma_1$ , after deleting any isolated vertices from the  $R$ -side.*

What distinguishes this paper from [10] in the case  $d \geq 4$  is that we also deal with the case where  $|L_1| \geq |R_1|$ .

### 3 Structure of the paper

We first prove Theorem 2. This involves studying Phase 1 of the Karp-Sipser Algorithm. For this we first describe the distribution of the graph  $G$ . This is done in Section 4. The distribution of  $\Gamma$  is determined by a few parameters and these evolve as a Markov chain. To study this chain, we introduce and solve a set of differential equations. This is done in Section 5. We show that the chains trajectory and the solution to the equations are close. By analysing the equations we can tell when Phase 1 is sufficient to solve the problem. This is done in Section 6. If Phase 1 is not sufficient then the graph  $\Gamma_1$  that remains has degree  $d$  on the  $L$ -side and minimum degree at least two on the  $R$ -side. We conjecture that **whp**  $\Gamma_1$  has a matching of size equal to the minimum set size of the partition. We can only prove this for  $d \geq 4$ . The argument is given in Section 7. We are left with the case  $d = 3$  and this has been half answered by [11] and [8] and [10].

## 4 Probability Model for Phase 1

We will represent  $G$  and more generally  $\Gamma$  by a random sequence  $\mathbf{x} \in \Omega_{L,R} = (R^d \cup \{\star\})^L$ . A sequence  $\mathbf{x} \in \Omega_{L,R}$  is to be viewed as  $n$  subsequences  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$  where  $\mathbf{x}_j = (x_{j,1}, x_{j,2}, \dots, x_{j,d}) \in R^d$  or  $\mathbf{x}_j = \sigma = (\star, \star, \dots, \star)$ . The  $\star$ 's represent edges that have been deleted by the Karp-Sipser algorithm. For  $\mathbf{x} \in \Omega_{L,R}$  we define the bipartite (multi-)graph  $\Gamma_{\mathbf{x}}$  as follows: Its vertex set consists of a bipartition  $L_{\mathbf{x}} = \{j \in L : \mathbf{x}_j \neq \sigma\}$  (the *left side*) and  $R$  (the *right side*). The edges incident with  $j \in L_{\mathbf{x}}$  are  $(j, x_{j,i}), i \in [d]$  i.e. we read the sequence  $\mathbf{x}$  from left to right and add edges to  $\Gamma_{\mathbf{x}}$  in blocks of size  $d$ . Each block being assigned to a unique vertex of  $L_{\mathbf{x}}$ .

We should be clear now that our probability space is  $\Omega_{L,R}$  with uniform measure and not  $\mathcal{G}_{L,R} = \{\Gamma_{\mathbf{x}} : \mathbf{x} \in \Omega_{L,R}\}$ .

Given a graph  $\Gamma_{\mathbf{x}}$  we let  $v_j = v_j(\mathbf{x}) = |R_j(\mathbf{x})|$  where  $R_j(\mathbf{x})$  is the set of vertices in  $R$  that have degree  $j \geq 0$ . We let  $v = v(\mathbf{x}) = |R_{\mathbf{x}}|$  where  $R_{\mathbf{x}} = \bigcup_{j \geq 2} R_j(\mathbf{x})$ .

For the graph  $G$  we choose  $\mathbf{x}(0)$  uniformly at random from  $(R^d)^L$  and put  $G = \Gamma_{\mathbf{x}(0)}$ . Next let  $\Gamma(0) = G$  and let  $\Gamma(t) = \Gamma_{\mathbf{x}(t)}$  be the graph  $\Gamma$  that we have after  $t$  steps of Phase 1 of the Karp-Sipser algorithm. The sequence  $\mathbf{x}(t)$  is defined as follows: Observe first that the vertex  $\xi$  of degree one is always in  $R$ . Suppose that it is incident to the unique edge  $(\eta, \xi), \eta \in L$ . Then we simply replace  $\mathbf{x}_{\eta}$  in  $\mathbf{x}(t-1)$  by  $\sigma$  to obtain  $\mathbf{x}(t)$ . We should thus think of the Karp-Sipser Algorithm as acting on sequences  $\mathbf{x}$  and not on graphs. We write  $\mathbf{x} \rightarrow \mathbf{y}$  to mean that  $\mathbf{y}$  can be obtained from  $\mathbf{x}$  by a single Phase 1 step of the Karp-Sipser Algorithm.

Let  $\vec{v}(t) = (w(t), v_1(t), v(t))$  where  $w(t) = |L_{\mathbf{x}(t)}|$  is the number of vertices on the left side of the bipartition of  $\Gamma(t)$ . Assuming that we have only run the Karp-Sipser algorithm up to the end of Phase 1, we have  $w(t) = n - t$ . Also

$$\vec{v}(0) \sim (n, \alpha d e^{-\alpha d} m, (1 - e^{-\alpha d} - \alpha d e^{-\alpha d}) m) \quad \text{whp.} \quad (5)$$

We will omit the parameter  $t$  from  $\vec{v}(t)$  when it is clear from the context. Let  $\mathcal{X}_{\vec{v}}$  be the set of all  $\mathbf{x} \in \Omega_{L,R}$  with parameters  $\vec{v}$ .

**Lemma 4** *Suppose  $\mathbf{x}(0)$  is a random member of  $\mathcal{X}_{\vec{v}(0)}$ . Then given  $\vec{v}(0), \dots, \vec{v}(t)$ ,  $\mathbf{x}(t)$  is a random member of  $\mathcal{X}_{\vec{v}(t)}$  for all  $t \geq 0$ .*

**Proof:** We prove this by induction on  $t$ . It is true for  $t = 0$  by assumption and so assume it is true for some  $t \geq 0$ . Let  $\vec{v}(t) = (w, v_1, v)$  and now fix a triple  $\vec{v}' = (w' = w - 1, v'_1, v')$  as a possible value for  $\vec{v}(t+1)$ . Fix  $\mathbf{y} \in \mathcal{X}_{\vec{v}'(t)}$ . We first compute the number of  $\mathbf{x} \in \mathcal{X}_{\vec{v}(t)}$  such that  $\mathbf{x} \rightarrow \mathbf{y}$ . Let  $b = v - v'$  be the number of vertices in  $R_{\mathbf{x}} \setminus R_{\mathbf{y}}$ . Some of these will be in  $R_0(\mathbf{y})$  and some will be in  $R_1(\mathbf{y})$ . So we choose non-negative integers  $b_0, b_1$  such that  $b_0 + b_1 = b$ . Next let  $a = v_1 - v'_1 + b_1$  be the number of vertices in  $R_1(\mathbf{x}) \cap R_0(\mathbf{y})$ . We can choose a vertex  $\eta$  so that  $L_{\mathbf{x}} \setminus L_{\mathbf{y}} = \{\eta\}$  in  $t$  ways and now let us enumerate the ways of choosing  $\mathbf{x}_{\eta} = (\zeta_1, \zeta_2, \dots, \zeta_d)$ . Our choices for  $\zeta_i$  are (i) distinctly from  $R_0(\mathbf{y})$  (i.e.  $\zeta_i$  is distinct from rest of the  $\zeta_j$ ), (ii) non-distinctly from  $R_0(\mathbf{y})$  (i.e.  $\zeta = \zeta_i$  is chosen more than once in the construction), (iii) from  $R_1(\mathbf{y})$  and (iv) from  $R_{\mathbf{y}}$ . We must exercise choice (i) exactly  $a$  times, choice (ii) at least twice for each of  $b_0$  distinct values, choice (iii) at least once for each of  $b_1$  distinct values and choice (iv) the remaining times. \*\*\*\*

The number of choices for  $\mathbf{x}_{\eta}$  depends only on  $\vec{v}$  and  $\vec{v}'$ , i.e. for each  $\mathbf{y} \in \mathcal{X}_{\vec{v}'}$  we have that  $D(\vec{v}, \vec{v}') = |\{\mathbf{x} \in \mathcal{X}_{\vec{v}} : \mathbf{x} \rightarrow \mathbf{y}\}|$  is independent of  $\mathbf{y}$ , given  $\vec{v}$  and  $\vec{v}'$ .

Similarly given  $\mathbf{x}$  there is a unique  $i \in R_1(\mathbf{x})$ , which when removed determines  $\mathbf{y}$ . Thus

$N(\vec{v}) = |\{\mathbf{y} : \mathbf{x} \rightarrow \mathbf{y}\}|$  is fixed given  $\vec{v}$ . Thus if  $\mathbf{x}(t)$  is a random member of  $\mathcal{X}_{\vec{v}}$  then

$$\begin{aligned}
& \mathbb{P}(\mathbf{x}(t+1) = \mathbf{y} | \vec{v}(0), \dots, \vec{v}(t)) \\
&= \sum_{\mathbf{x} \in \mathcal{X}_{\vec{v}(t)}} \mathbb{P}(\mathbf{x}(t) = \mathbf{x} | \vec{v}(0), \dots, \vec{v}(t)) \cdot \mathbb{P}(\mathbf{x}(t+1) = \mathbf{y} | \vec{v}(0), \dots, \vec{v}(t-1), \mathbf{x}(t) = \mathbf{x}) \\
&= \sum_{\mathbf{x} \in \mathcal{X}_{\vec{v}(t)}} \mathbb{P}(\mathbf{x}(t+1) = \mathbf{y} | \mathbf{x}(t) = \mathbf{x}) \cdot |\mathcal{X}_{\vec{v}(t)}|^{-1} \\
&= \sum_{\mathbf{x} \in \mathcal{X}_{\vec{v}(t)}} \frac{D(\vec{v}(t), \vec{v}(t+1))}{N(\vec{v}(t))} \cdot |\mathcal{X}_{\vec{v}(t)}|^{-1}
\end{aligned}$$

which is independent of  $\mathbf{y}$  given  $\vec{v}(t)$  and so  $\mathbf{y}$  is a random member of  $\mathcal{X}_{\vec{v}(t+1)}$ .  $\square$

**Lemma 5** *The random sequence  $\vec{v}(t), t = 0, 1, 2, \dots$  is a Markov chain.*

**Proof:** As in [2],

$$\begin{aligned}
\mathbb{P}(\vec{v}(t+1) | \vec{v}(0), \vec{v}(1), \dots, \vec{v}(t)) &= \sum_{\mathbf{x}' \in \mathcal{X}_{\vec{v}(t+1)}} \mathbb{P}(\mathbf{x}' | \vec{v}(0), \vec{v}(1), \dots, \vec{v}(t)) \\
&= \sum_{\mathbf{x}' \in \mathcal{X}_{\vec{v}(t+1)}} \sum_{\mathbf{x} \in \mathcal{X}_{\vec{v}(t)}} \mathbb{P}(\mathbf{x}', \mathbf{x} | \vec{v}(0), \vec{v}(1), \dots, \vec{v}(t)) \\
&= \sum_{\mathbf{x}' \in \mathcal{X}_{\vec{v}(t+1)}} \sum_{\mathbf{x} \in \mathcal{X}_{\vec{v}(t)}} \mathbb{P}(\mathbf{x}' | \vec{v}(0), \vec{v}(1), \dots, \vec{v}(t-1), \mathbf{x}) \\
&\quad \times \mathbb{P}(\mathbf{x} | \vec{v}(0), \vec{v}(1), \dots, \vec{v}(t)) \\
&= \sum_{\mathbf{x}' \in \mathcal{X}_{\vec{v}(t+1)}} \sum_{\mathbf{x} \in \mathcal{X}_{\vec{v}(t)}} \mathbb{P}(\mathbf{x}' | \mathbf{x}) |\mathcal{X}_{\vec{v}(t)}|^{-1},
\end{aligned}$$

which depends only on  $\vec{v}(t), \vec{v}(t+1)$ .  $\square$

**Lemma 6** *Conditional on  $\vec{v}$  if  $\mathbf{x}$  is selected uniformly at random from  $\mathcal{X}_{\vec{v}}$  then each vertex  $i \in R_{\mathbf{x}}$  has degree  $Y_i$  where  $Y_i = \text{Poi}(z; \geq 2)$ , a Poisson random variable conditioned to take a value at least two, and  $z$  satisfies*

$$\frac{z(e^z - 1)}{f(z)} = \frac{dw - v_1}{v} \tag{6}$$

where  $f(z) = e^z - z - 1$ .

The  $Y_i$  are also conditioned to satisfy  $\sum_{i=1}^v Y_i = wd - v_1$ .

**Proof:** Suppose we first fix the edges incident with vertices of degree one in  $\mathbf{x}$ . Then we randomly fill in the remaining  $dw - v_1$  non- $\star$  positions in  $\mathbf{x}$  with values from some fixed  $v$ -subset  $R_{\mathbf{x}}$  of  $R$ , subject to each of these  $v$  vertices having degree at least two. The degrees  $Y_i$  of these vertices will have the description described in the lemma (for a proof see Lemma 4 of [2]).  $\square$

From [2] we can use the following lemma

**Lemma 7** [2]

(a) *Assume that  $\log n = O((vz)^{\frac{1}{2}})$ . For every  $j \in R_{\mathbf{x}}$  and  $2 \leq k \leq \log n$ ,*

$$\mathbb{P}(Y_j = k | \vec{v}) = \frac{z^k}{k! f(z)} \left( 1 + O\left(\frac{k^2 + 1}{vz}\right) \right) \tag{7}$$

(b) For all  $k \geq 2, j \in R_x$

$$\mathbb{P}(Y_j = k | \vec{v}) = O\left(\frac{(vz)^{\frac{1}{2}} z^k}{k! f(z)}\right)$$

□

## 5 Differential Equations

Let  $\vec{v}$  be the current parameter tuple and  $\vec{v}'$  be the tuple after one step of the Karp-Sipser algorithm. The following lemma gives  $E[\vec{v}' - \vec{v} | \vec{v}]$  for each step of Phase 1 of the Karp-Sipser algorithm.

**Lemma 8** *Assuming  $\log v = O((vz)^{\frac{1}{2}})$  and  $v_1 > 0$  we have*

$$\begin{aligned} \mathbb{E}[v'_1 - v_1 | \vec{v}] &= -1 - \frac{d-1}{dw} v_1 + \frac{d-1}{dw} \frac{vz^2}{f} + O\left(\frac{1}{vz}\right) \\ \mathbb{E}[v' - v | \vec{v}] &= -\frac{d-1}{dw} \frac{vz^2}{f} + O\left(\frac{1}{vz}\right) \end{aligned}$$

**Proof:** First note that  $v_1 > 0$ , one vertex  $\xi \in R$  with  $\deg(\xi) = 1$  will be picked and  $\xi$  and its neighbor  $\eta \in L$  will be removed from  $G_t$ . This implies that  $w$  decreases by 1 and the number of edges removed is  $d$ , i.e. all edges incident to  $\eta$ . Let  $\delta$  be the number of multiple edges incident to  $\eta$ . Then we have

$$\begin{aligned} \mathbb{E}[\delta | \vec{v}] &\leq d \cdot \mathbb{P}(\eta \text{ is incident to parallel edges}) \leq d \binom{d}{2} \frac{1}{(wd)_2} \sum_j \mathbb{E}[\deg(j)_2] \\ &\leq \frac{d^3}{2w^2} v \mathbb{E}[Y(Y-1) | \vec{v}] = O\left(\frac{1}{w}\right) \end{aligned} \quad (8)$$

where  $Y$  has distribution (7).

**Explanation:** The  $dw$  choices of neighbours for the remaining vertices in  $L$  form a list with  $v_1$  unique names and  $wd - v_1$  non-unique names and where the number of times a vertex appears among the  $wd - v_1$  has distribution (7). Also, if we construct this list vertex by vertex, it will appear in a random order. So the probability that  $j$  appears in two of the choices for  $\eta$  is bounded by  $\frac{\mathbb{E}[\deg(j)_2]}{(wd)_2}$  and this justifies (8).

The change in  $v_1$  comes from  $\xi$  being removed, minus the number of other degree one vertices adjacent to  $\eta$  and plus the number of vertices adjacent to  $\eta$  of degree exactly two. Any change from vertices of degree three or more is absorbed by the  $O\left(\frac{1}{w}\right)$  term for multiple edges.

The expected change is then

$$\begin{aligned} \mathbb{E}[v'_1 - v_1 | \vec{v}] &= -1 - \frac{d-1}{dw-1} (v_1 - 1) + \frac{2(d-1)}{dw-1} \mathbb{E}[v_2 | \vec{v}] + O\left(\frac{1}{w}\right) \\ &= -1 - \frac{d-1}{dw} v_1 + \frac{d-1}{dw} \frac{z^2}{f} v + O\left(\frac{1}{vz}\right) \end{aligned} \quad (9)$$

Similarly for  $v$ , the change is only due to vertices adjacent to  $y$  of degree exactly two, modulo multiple edges. Thus

$$\mathbb{E}[v' - v | \vec{v}] = -\frac{d-1}{dw} \frac{z^2}{f} v + O\left(\frac{1}{vz}\right)$$

□

Lemma 8 suggests that we consider the following pair of differential equations

$$\frac{dy_1}{dt} = -1 - \frac{d-1}{dw}y_1 + \frac{d-1}{dw}\frac{y\zeta^2}{f(\zeta)} \quad (10)$$

$$\frac{dy}{dt} = -\frac{d-1}{dw}\frac{y\zeta^2}{f(\zeta)} \quad (11)$$

where  $w = n - t$  and  $\zeta$  satisfies

$$\frac{\zeta(e^\zeta - 1)}{f(\zeta)} = \frac{dw - y_1}{y}. \quad (12)$$

The boundary conditions are (see (5))

$$\zeta(0) = \alpha d, \quad y_1(0) = m\alpha d e^{-\alpha d}, \quad y(0) = m(1 - (1 + \alpha d)e^{-\alpha d}). \quad (13)$$

The  $y_1, y, \zeta$  are of course the deterministic counterparts of  $v_1, v, z$  respectively.

**Lemma 9** *The solution to (10), (11) and (13) is*

$$w = \left(\frac{\zeta}{\alpha d}\right)^{\frac{d}{d-1}} n. \quad (14)$$

$$t = n \left(1 - \left(\frac{\zeta}{\alpha d}\right)^{\frac{d}{d-1}}\right) \quad (15)$$

$$y = e^{-\zeta} f(\zeta) m \quad (16)$$

$$y_1 = m\zeta \left( \left(\frac{\zeta}{\alpha d}\right)^{\frac{1}{d-1}} + e^{-\zeta} - 1 \right) \quad (17)$$

**Proof:** We take the derivative of (6) with respect to  $t$ . The RHS becomes, using (10) and (11)

$$\begin{aligned} \frac{d}{dt} \left( \frac{dw - y_1}{y} \right) &= \frac{1}{y} \left( -d - \left( -1 - \frac{d-1}{d} \frac{y_1}{w} + \frac{d-1}{d} \frac{y}{w} \frac{\zeta^2}{f(\zeta)} \right) - \frac{dw - y_1}{y} \frac{1}{y} \left( -\frac{d-1}{d} \frac{y}{w} \frac{\zeta^2}{f(\zeta)} \right) \right) \\ &= -\frac{d-1}{dw} \left( \frac{dw - y_1}{y} + \frac{\zeta^2}{f(\zeta)} - \frac{dw - y_1}{y} \frac{\zeta^2}{f(\zeta)} \right) \\ &= -\frac{d-1}{dw} \left( \frac{\zeta(e^\zeta - 1)f(\zeta) + \zeta^2 f(\zeta) - \zeta^3(e^\zeta - 1)}{f(\zeta)^2} \right) \\ &= -\frac{d-1}{dw} \left( \frac{\zeta(e^\zeta - 1)^2 - \zeta^3 e^\zeta}{f(\zeta)^2} \right) \end{aligned} \quad (18)$$

On the other hand, on differentiating the LHS of (6) (with  $z$  replaced by  $\zeta$ ) we get

$$\begin{aligned} \frac{d}{dt} \left( \frac{\zeta(e^\zeta - 1)}{f(\zeta)} \right) &= \frac{(\zeta e^\zeta + e^\zeta - 1)f(\zeta) - \zeta(e^\zeta - 1)^2}{f(\zeta)^2} \frac{d\zeta}{dt} \\ &= \frac{(e^\zeta - 1)^2 - \zeta^2 e^\zeta}{f(\zeta)^2} \frac{d\zeta}{dt}. \end{aligned} \quad (19)$$

Comparing (18) and (19) we see that

$$\frac{1}{\zeta} \frac{d\zeta}{dt} = -\frac{d-1}{dw} \quad (20)$$

Integrating yields

$$\frac{\zeta^d}{w^{d-1}} = \text{constant}$$

Plugging in  $\zeta(0) = \alpha d$  we see that

$$\frac{\zeta^d}{w^{d-1}} = \frac{(\alpha d)^d}{n^{d-1}}.$$

This verifies (14) and (15).

Going back to (11) and (20) we have

$$\frac{dy}{d\zeta} \frac{d\zeta}{dt} = \frac{dy}{dt} = -\frac{(d-1)y}{dw} \frac{\zeta^2}{f(\zeta)} = \frac{1}{\zeta} \frac{d\zeta}{dt} \frac{y\zeta^2}{f(\zeta)}.$$

So

$$\frac{1}{y} \frac{dy}{d\zeta} = \frac{\zeta}{f(\zeta)}$$

and integrating yields

$$\ln y = -\zeta + \ln f(\zeta) + C$$

Taking  $y(0) = m(1 - (1 + \alpha d)e^{-\alpha d})$  gives  $C = \ln m$  and we see that (16) holds.

We now solve for  $y_1$  in terms of  $\zeta$  as a function of  $d$ . It follows from (12) and (16) that

$$\begin{aligned} y_1 &= dw + \zeta e^{-\zeta} m - m\zeta \\ &= \left(\frac{\zeta}{\alpha d}\right)^{\frac{d}{d-1}} nd + \zeta e^{-\zeta} m - m\zeta \\ &= m\zeta \left( \left(\frac{\zeta}{\alpha d}\right)^{\frac{1}{d-1}} + e^{-\zeta} - 1 \right) \end{aligned}$$

□

At this point we wish to show that **whp** the sequence  $\vec{v}(t), t \geq 0$  closely follows the trajectory  $\vec{y}(t) = (w, y_1, y), t \geq 0$  described in Lemma 9. One possibility is to use Theorem 5.1 of Wormald [22], but there is a problem with an “unbounded” Lipschitz coefficient. One can allow for this in [22], but it is unsatisfactory to ask the reader to check this. We have decided to use an approach suggested in Bohman [4].

Next let  $K$  be a large positive constant and let  $\gamma \ll 1/K$ . Then let

$$g(x) = (1-x)^{-K} + K(1-x)^{-1}$$

and

$$Err(t) = n^{2/3} g(t/n).$$

Then define the event

$$\mathcal{E}(t) = \left\{ v(\tau)z(\tau) \geq n^{1/2} \text{ and } \zeta(\tau) \geq \zeta(t_1) + n^{-\gamma} \text{ and } |\vec{v}(\tau) - \vec{y}(\tau)|_\infty \leq 2Err(\tau) \text{ for } \tau \leq t \right\}$$

where

$$t_1 = \min \{t > 0 : y_1(t) = 0\}. \tag{21}$$

Now define four sequences of random variables:

$$\begin{aligned} X_1^\pm(t) &= \begin{cases} v_1(t) - y_1(t) \pm Err(t) & \mathcal{E}(t-1) \text{ holds} \\ X_1^\pm(t-1) & \text{otherwise} \end{cases} \\ X^\pm(t) &= \begin{cases} v(t) - y(t) \pm Err(t) & \mathcal{E}(t-1) \text{ holds} \\ X^\pm(t-1) & \text{otherwise} \end{cases} \end{aligned}$$

Because  $(1-x)^{-L}$  is convex we have

$$\frac{1}{(1-(x+h))^L} \geq \frac{1}{(1-x)^L} + \frac{hL}{(1-x)^{L+1}}$$

for  $L > 0$  and  $0 < x < x+h < 1$ . So,

$$g((t+1)/n) - g(t/n) \geq \frac{K}{n-t} g(t/n). \quad (22)$$

Suppose that  $\mathcal{E}(t)$  holds. We write

$$\begin{aligned} \left| \frac{dw - y_1}{y} - \frac{dw - v_1}{v} \right| &= \left| \frac{(dw - y_1)(v - y)}{yv} + \frac{v_1 - y_1}{v} \right| \\ &= O\left(\frac{Err(t)}{v}\right). \end{aligned} \quad (23)$$

(For this we need  $(dw - y_1)/y = O(1)$ . But this follows from (12) and the fact that  $\zeta$  is decreasing – see (20)).

Putting  $F(x) = \frac{x(e^x-1)}{f(x)}$  we have (see (19))  $F'(x) = \frac{(e^x-1)^2 - x^2 e^x}{f(x)^2}$  and since

$$(e^x - 1)^2 - x^2 e^x = \sum_{k=4}^{\infty} (2^k - 2 - k(k-1)) \frac{x^k}{k!} \quad (24)$$

we see that  $F'(x) = \Omega(1)$  in any bounded interval  $[0, L]$ .

Hence from (23) we have

$$O\left(\frac{Err(t)}{v}\right) = F(\zeta) - F(z) = \Omega(|\zeta - z|)$$

or

$$|\zeta - z| = O\left(\frac{Err(t)}{v}\right). \quad (25)$$

Using (25) we obtain

$$\begin{aligned} \left| \frac{vz^2}{f(z)} - \frac{y\zeta^2}{f(\zeta)} \right| &\leq \frac{|v-y|z^2}{f(z)} + y \left| \frac{z^2}{f(z)} - \frac{\zeta^2}{f(\zeta)} \right| \\ &\leq K_1 Err(t) \end{aligned} \quad (26)$$

for some  $K_1 = K_1(\alpha, d) > 0$ .

For the second term we use

$$\left(\frac{x^2}{f(x)}\right)' = \frac{2xf(x) - x^2(e^x - 1)}{f(x)^2} = -\frac{1}{f(x)^2} \sum_{k=4}^{\infty} \frac{k-3}{(k-1)!} x^k.$$

This implies that  $\left(\frac{x^2}{f(x)}\right)' = O(1)$  for  $x \geq 0$ .

Now with  $f = f(\zeta)$ ,

$$\begin{aligned} y_1''(t) &= \frac{d-1}{dw} \left( 1 - \frac{y_1}{w} + \frac{y\zeta^2}{wf} + \frac{d-1}{dw} \left( y_1 - \frac{y\zeta^2}{f} - \frac{y\zeta^4}{f^2} - y\zeta^2 \left( \frac{2}{f} - \frac{e^\zeta - 1}{f^2} \right) \right) \right) \\ &= O(\zeta^{-O(1)}n^{-1}). \\ y''(t) &= \frac{d-1}{dw} \left( \frac{y\zeta^2}{wf} + \frac{d-1}{dw} \left( \frac{y\zeta^4}{f^2} + y\zeta \left( \frac{2\zeta}{f} - \frac{\zeta(e^\zeta - 1)}{f^2} \right) \right) \right) \\ &= O(\zeta^{-O(1)}n^{-1}). \end{aligned}$$

If  $\mathcal{E}(t)$  holds then, where  $\rho_t = \frac{d-1}{dw} = \frac{d-1}{d(n-t)}$ ,

$$\begin{aligned} &\mathbb{E}(X_1^+(t+1) - X_1^+(t) \mid \vec{v}(t)) = \\ &\mathbb{E}(v_1(t+1) - v_1(t) \mid \vec{v}(t)) - (y_1(t+1) - y_1(t)) + n^{2/3}(g((t+1)/n) - g(t/n)) \geq \\ &- \rho_t(v_1(t) - y_1(t)) + \rho_t \left( \frac{v(t)z(t)^2}{f(z(t))} - \frac{y(t)\zeta(t)^2}{f(\zeta(t))} \right) + O\left(\frac{1}{v(t)z(t)}\right) - y_1''(t+\theta) + \frac{Kn^{2/3}}{n-t}g(t/n) \\ &\geq \frac{n^{2/3}g(t/n)}{n-t}(-K_1 - 2 + K) + O\left(\frac{1}{v(t)z(t)}\right) - y_1''(t+\theta) \\ &\geq 0. \end{aligned}$$

This shows that  $X_1^+(t), t \geq 0$  is a sub-martingale. Also,

$$\begin{aligned} |X_1^+(t+1) - X_1^+(t)| &\leq \\ &|v_1(t+1) - v_1(t)| + \sup_{0 \leq \theta \leq 1} |y_1'(t+\theta)| + n^{2/3}(g((t+1)/n) - g(t/n)) = O(1). \end{aligned}$$

It follows from the Azuma-Hoeffding inequality that we can write

$$\mathbb{P}(\exists 1 \leq t \leq t_1 : X_1^+(t) \leq X_1^+(0) - n^{3/5}) \leq e^{-\Omega(n^{1/5})}.$$

By almost identical arguments we have

$$\mathbb{P}(\exists 1 \leq t \leq t_1 : X_1^-(t) \geq X_1^-(0) + n^{3/5}) \leq e^{-\Omega(n^{1/5})}.$$

$$\mathbb{P}(\exists 1 \leq t \leq t_1 : X^+(t) \leq X^+(0) - n^{3/5}) \leq e^{-\Omega(n^{1/5})}.$$

$$\mathbb{P}(\exists 1 \leq t \leq t_1 : X^-(t) \geq X^-(0) + n^{3/5}) \leq e^{-\Omega(n^{1/5})}.$$

It follows that **whp**, when  $\mathcal{E}(t)$  holds, we have

$$|v_1(t) - y_1(t)| \leq Err(t) + n^{3/5} + |v_1(0) - y_1(0)| < 2Err(t). \quad (27)$$

$$|v(t) - y(t)| \leq Err(t) + n^{3/5} + |v(0) - y(0)| < 2Err(t). \quad (28)$$

Now by construction,  $\mathcal{E}(t)$  will fail at some time  $t_2 \leq t_1$ . It follows from (27), (28) that **whp** it will fail either because (i)  $v(t_2)z(t_2) < n^{1/2}$  or  $\zeta(t_2) < \zeta(t_1) + n^{-\gamma}$ . We claim the latter. Observe that if  $\zeta(\tau) \geq \zeta(t_1) + n^{-\gamma}$  then (14)–(17) imply  $w, y_1 = \Omega(n^{1-d\gamma/(d-1)})$  and  $y = \Omega(n^{1-2\gamma})$ . Together with (27), (28), this implies that  $v(t_2)z(t_2) = \Omega(n^{1-3\gamma}) \geq n^{1/2}$ .

In summary then, **whp** the process satisfies

$$|\vec{v}(t) - \vec{y}(t)|_\infty < 2Err(t_2) = O\left(\frac{n^{2/3}}{(1 - t_2/n)^K}\right) = O(n^{2/3 + Kd\gamma/(d-1)}) \quad \text{for } 1 \leq t \leq t_2 \quad (29)$$

and

$$v_1(t_2) = O(n^{1-\gamma}) \text{ where } t_2 = t_1 + O(n^{1-\gamma}). \quad (30)$$

We use (14) for (29) and (17), (25) for (30).

## 6 Analysis of Phase 1

We will first argue that **whp** Phase 1 is sufficient to find a matching from  $L$  to  $R$  when there is no solution  $0 < \zeta \leq \alpha d$  to

$$\left(\frac{\zeta}{\alpha d}\right)^{\frac{1}{d-1}} + e^{-\zeta} - 1 = 0. \quad (31)$$

It follows from (17), (21) and (30) that in this case Phase 1 ends with there being at most  $O(n^{1-\gamma})$  vertices of  $L$  left unmatched, **whp**. Furthermore at time  $t_2$  we will have

$$w \sim \left(\frac{\zeta}{\alpha d}\right)^{\frac{d}{d-1}} n, v_1 \sim dw \text{ and } v = O(\zeta^{\frac{d-2}{d-1}} v_1)$$

where  $\zeta = \zeta(t_2) = O(n^{-\gamma})$ .

**Lemma 10** *Suppose that  $t_1 = 0$ . Then **whp** at time  $t_2$ ,  $\Gamma$  is a forest.*

**Proof:** Let  $R_{\geq 2}$  denote the set of vertices of degree at least two in the  $R$ -side of  $\Gamma$ . Let  $P(d_1, \dots, d_k)$  denote  $\mathbb{P}(X_1 = d_1, \dots, X_k = d_k)$  where  $X_1, \dots, X_v$  are truncated Poisson conditioned only to sum to  $dw - v_1$ . For large  $k$  we use the bound

$$\mathbb{P}(X_1 = d_1, \dots, X_k = d_k) \leq O(n^{1/2}) \prod_{i=1}^k \frac{z^{d_i}}{d_i! f(z)}. \quad (32)$$

For  $k = O(1)$  and  $d_1, \dots, d_k = O(\log n)$  we write

$$\mathbb{P}(X_1 = d_1, \dots, X_k = d_k) = \prod_{i=1}^k \mathbb{P}(X_i = d_i \mid X_j = d_j, j < i) = (1 + o(1)) \prod_{i=1}^k \frac{z^{d_i}}{d_i! f(z)}. \quad (33)$$

It is equation (7) that allows us to write the final equality in (33). The extra conditioning  $X_j = d_j, j < i$  only changes the required sum.

Thus let  $B_k$  denote  $O(n^{1/2})$  for  $k \geq \frac{2(d-1)}{(d-2)^\gamma}$  and  $1 + o(1)$  otherwise. The expected number of cycles can be bounded by  $o(1) = (\mathbb{P}(\exists \text{ vertex of degree } \geq \log n))$  plus

$$\sum_{k \geq 2} \sum_{\substack{S \subseteq R_{\geq 2} \\ |S|=k}} \sum_{2 \leq d_1, \dots, d_k \leq \log n} P(d_1, \dots, d_k) \times \binom{w}{k} (d(d-1))^k (k!)^2 \prod_{i=1}^k \frac{d_i(d_i-1)}{(dw-2i+2)(dw-2i+1)} \leq \quad (34)$$

$$\begin{aligned} & \sum_{k \geq 2} B_k \binom{v}{k} \sum_{2 \leq d_1, \dots, d_k \leq \log n} \prod_{i=1}^k \frac{z^{d_i}}{d_i! f(z)} \times \binom{w}{k} \binom{d}{2}^k (k!)^2 \prod_{i=1}^k \frac{d_i(d_i-1)}{(dw-2i+2)(dw-2i+1)} \leq \\ & \sum_{k \geq 2} B_k \left( \frac{vwz^2 d^2}{f(z)(dw-2k)^2} \right)^k \sum_{2 \leq d_1, \dots, d_k} \prod_{i=1}^k \frac{z^{d_i-2}}{(d_i-2)!} = \\ & \sum_{k \geq 2} B_k \left( \frac{vwz^2 e^z d^2}{f(z)(dw-2k)^2} \right)^k = \\ & \sum_{k \geq 2} B_k O(\zeta^{\frac{d-2}{d-1}})^k = \\ & o(1). \end{aligned}$$

□

**Explanation of (34):** We condition on the degree sequence. Having fixed the degree sequence, we swap to the configuration model [5]. Having chosen  $S \subseteq R$  and  $k$  vertices  $W$  in  $L$  and their degrees, we can work within this model. We then choose a cycle through these vertices in  $(k!)^2$  ways. We then choose the configuration points associated with our  $k$ -cycle in  $(d(d-1))^k \prod_{i=1}^k d_i(d_i-1)$  ways. We then multiply by the probability  $\prod_{i=1}^k \frac{1}{(dw-2i+2)(dw-2i+1)}$  of choosing the pairings associated with the edges of the cycle.

**Corollary 11** *Suppose that  $t_1 = 0$ . Then **whp** at time  $t_2$ ,  $\Gamma$  contains a matching from  $L_\Gamma$  into  $R_\Gamma$ . Furthermore, such a matching will be constructed in Phase 1.*

**Proof:** We can assume from Lemma 10 that  $\Gamma$  is a forest. Each vertex of  $L_\Gamma$  has degree  $d$  and so Hall's theorem will show that the required matching exists. (Any Hall witness would induce a cycle). Finally note that Phase 1 of the Karp-Sipser algorithm is exact on a forest. □

## 6.1 Threshold for Phase 1 to be sufficient

Put  $A = (\alpha d)^{-1/(d-1)}$  and  $B = 1/(d-1)$  so that (31) can be written as

$$A\zeta^B + e^{-\zeta} - 1 = 0. \quad (35)$$

Assume  $B$  is fixed. We find a threshold for  $A$  in terms of  $B$  for there to be no positive solution to (35). We find the place  $\zeta^*$  where the curve  $y = 1 - e^{-\zeta}$  touches the curve  $y = A\zeta^B$  i.e. where

$$\begin{aligned} A\zeta^B &= 1 - e^{-\zeta} \\ AB\zeta^{B-1} &= e^{-\zeta} \end{aligned}$$

In which case

$$\frac{\zeta^*}{B} = e^{\zeta^*} - 1 \text{ and } A^* = \frac{1 - e^{-\zeta^*}}{(\zeta^*)^B} \quad (36)$$

or

$$\zeta^* = \frac{e^{\zeta^*} - 1}{d - 1} \text{ and } \alpha^* = \frac{\zeta^*}{d(1 - e^{-\zeta^*})^{d-1}}. \quad (37)$$

In general, keeping  $B < 1$  fixed let

$$f_A(\zeta) = A\zeta^B + e^{-\zeta} - 1 \text{ and } L_A = A^{-1/B}.$$

We must show that if  $A \geq A^*$  then the only solution to  $f_A(\zeta) = 0$ ,  $0 \leq L_A$  is  $\zeta = 0$ .

Observe that  $f_A(L_A) = e^{-L_A} > 0$  and  $f_A(0) = 0$ . Also, if  $A < 1$  then  $1 < L_A$  and if  $A < 1 - e^{-1}$  then  $f_A(1) < 0$  and there must be a positive solution to  $f_A(\zeta) = 0$ .

Observe that  $A' > A$  implies (i)  $L_{A'} < L_A$  and that (ii)  $f_{A'}(\zeta) > f_A(\zeta)$  for  $\zeta \neq 0$ . So if  $f_A(\zeta)$  has no positive solution then neither has  $f_{A'}$ . We argue that  $f'_A(\zeta) = 0$  has at most 2 solutions, which implies that  $f_A(\zeta) = 0$  has at most two positive solutions. As we increase  $A$  to  $A^*$  these solutions must converge, by (ii).

Now

$$f'_A(\zeta) = 0 \text{ iff } \frac{e^\zeta}{\zeta^{1-B}} = \frac{1}{AB}.$$

But the function  $g(\zeta) = e^\zeta/\zeta^\xi$  is convex for any  $\xi > 0$ . Indeed

$$g''(\zeta) = \frac{e^\zeta((\zeta - \xi)^2 + \xi)}{\zeta^{\xi+2}} > 0$$

and so  $g(\zeta) = a$  has at most two solutions for any  $a > 0$ .

## 6.2 Finishing the proof of Theorem 2

We now have to relate the above results to the actual process. We know from our analysis of the differential equations that for some  $A > 0$ ,

$$v_1(t_2 = t_1 + An^{-\gamma}) = O(n^{1-\gamma}).$$

When  $\zeta_1 = \zeta(t_1) = 0$ , Lemma 10 and Corollary 11 imply Theorem 2.

So assume that  $\zeta_1 > 0$ . Thus  $\alpha > \alpha^*$  and  $A < A^*$ . We argue that if  $z_A$  is the solution to (35) then  $z_A$  decreases monotonically with  $A$ . Indeed, if  $A' > A$  then  $f_{A'}(\zeta) > f_A(\zeta)$  for  $\zeta_A \leq \zeta \leq L_{A'} < L_A$ . Now  $\frac{(d-1)z}{e^z-1}$  is strictly monotone decreasing with  $z$  and so

$$\frac{(d-1)z_A}{e^{z_A}-1} < \frac{(d-1)z_{A^*}}{e^{z_{A^*}}-1} = 1. \quad (38)$$

The second equation in (38) is the first equation in (36).

At time  $t_2$  we will have  $v_1 = O(n^{1-\gamma})$  and  $v = \Omega(n)$ . For the next  $o(n)$  steps we have from (9) that

$$\mathbb{E}(v'_1 - v_1 \mid \vec{v}) = -1 + (1 + o(1)) \frac{d-1}{dw} \frac{z^2}{f} v = -1 + (1 + o(1)) \frac{(d-1)z_A}{e^{z_A}-1} \leq -\varepsilon \quad (39)$$

for some small positive  $\varepsilon$ . In which case, **whp**,  $v_1$  will become zero in  $O(n^{1-\gamma} \log n)$  steps. Indeed (39) implies that the sequence

$$X_k = \begin{cases} v_1(t_2 + k) + \varepsilon k & \text{if } v_1(t_2 + k) > 0 \\ X_{k-1} & \text{otherwise} \end{cases}$$

is a supermartingale that cannot change by more than  $d$  in any step. The Azuma-Hoeffding inequality implies that for  $T = 2X_0/\varepsilon$  we have

$$\mathbb{P}(v_1(t_2 + T) > 0) \leq \mathbb{P}(X_T - X_0 \geq \varepsilon T - X_0) \leq \exp \left\{ -\frac{(\varepsilon T - X_0)^2}{2Td^2} \right\} = o(1).$$

I.e. **whp**  $v_1(t_2 + T) = 0$  and  $v(t_2 + T) = v(t_2) - o(n) = \Omega(n)$ .

## 7 Proof of Theorem 3

Let us summarize what we have to prove. We have a random bipartite graph  $\Gamma_1$  with partition  $L_1, R_1$  and  $|L_1| = n_1 = \alpha_1 m_1, |R_1| = m_1$ . Each vertex in  $L_1$  has degree  $d$  and each vertex in  $R$  has degree at least 2. At this point it is convenient to drop the suffix 1. So from now on,  $m, n, \alpha, \Gamma$  etc. refer to the graph left at the end of Phase 1.

The degrees of  $\Gamma$  satisfy,  $d_L(a) = d$  for  $a \in L$ . The degrees of vertices in  $R$  are distributed as the box occupancies  $X_1, X_2, \dots, X_n$  in the following experiment. We throw  $dn$  balls randomly into  $n$  boxes and condition that each box gets at least two balls. In these circumstance the  $X_j$ 's are independent truncated Poisson, subject to the condition that  $X_1 + X_2 + \dots + X_n = dn$ , see Lemma 6 with  $v_1 = 0$ . Thus for any  $S = \{b_1, b_2, \dots, b_s\} \subseteq R$  and any set of positive integers  $k_i \geq 2, i \in S$  we have

$$\mathbb{P}(d_{R_1}(b_i) = k_i, i \in S) \leq O(n^{1/2}) \prod_{i \in S} \frac{z^{k_i}}{k_i! f(z)}$$

for  $k \geq 2$  where  $z$  satisfies

$$\frac{z(e^z - 1)}{f(z)} = \frac{nd}{m}.$$

The  $O(n^{1/2})$  term accounts for the conditioning  $\sum_{b \in R} d_R(b) = dn$  We will prove

**Theorem 12** *Let  $\Gamma$  be a bipartite graph chosen uniformly from the sets of graphs with bipartition  $L, R, |L| = n, |R| = m$  such that each vertex of  $L$  has degree  $d \geq 4$  and each vertex of  $R$  has degree at least two. Then **whp***

$$\mu(\Gamma) = \min \{m, n\}.$$

### 7.1 Useful Lemmas

Define the function  $\zeta(\gamma), \gamma > 0$  to be the unique solution to

$$\frac{u(e^u - 1)}{f(u)} = \gamma.$$

Let  $g$  be defined by

$$g(x) = (e^{\zeta(x)} - 1)^x f(\zeta(x))^{1-x}.$$

Observe that

$$\frac{f(\zeta(x))}{\zeta(x)^x} = \frac{g(x)}{x^x}. \tag{40}$$

**Lemma 13** *The function  $g(x)$  is log-concave as a function of  $x$*

**Proof:** We will write  $\zeta$  for  $\zeta(x)$  and  $f$  for  $f(\zeta)$  throughout this proof. Now  $\frac{\zeta(e^\zeta-1)}{f} = x$  from which we get

$$\frac{d\zeta}{dx} = \frac{f^2}{(e^\zeta - 1)^2 - \zeta^2 e^\zeta} \quad (41)$$

and note that  $\frac{d\zeta}{dx} > 0$  for  $\zeta > 0$ . Taking the derivative of  $\log(g(x))$  we get

$$\begin{aligned} \frac{d}{dx} \log(g(x)) &= \frac{d}{dx} \left( x \log(e^\zeta - 1) + (1 - x) \log(e^\zeta - \zeta - 1) \right) \\ &= \log\left(\frac{e^\zeta - 1}{f}\right) + \frac{d\zeta}{dx} \left( x \frac{e^\zeta}{e^\zeta - 1} + (1 - x) \frac{e^\zeta - 1}{f} \right) \end{aligned}$$

Now  $x = \frac{\zeta(e^\zeta-1)}{f}$  so

$$\begin{aligned} x \frac{e^\zeta}{e^\zeta - 1} + (1 - x) \frac{e^\zeta - 1}{f} &= \frac{\zeta e^\zeta}{f} + \frac{f - \zeta(e^\zeta - 1) e^\zeta - 1}{f} \\ &= \frac{\zeta e^\zeta (e^\zeta - \zeta - 1) + (e^\zeta - \zeta - 1 - \zeta e^\zeta + \zeta)(e^\zeta - 1)}{f^2} \\ &= \frac{(e^\zeta - 1)^2 - \zeta^2 e^\zeta}{f^2} \\ &= \frac{dx}{d\zeta} \end{aligned}$$

Thus we have

$$\frac{d}{dx} \log(g(x)) = \log\left(\frac{e^\zeta - 1}{f}\right) + 1 \quad (42)$$

Taking the second derivative we get

$$\begin{aligned} \frac{d^2}{dx^2} \log(g(x)) &= \frac{d}{dx} \left( \log\left(\frac{e^\zeta - 1}{f}\right) + 1 \right) \\ &= \frac{f}{e^\zeta - 1} \frac{e^\zeta (e^\zeta - \zeta - 1) - (e^\zeta - 1)^2 \frac{d\zeta}{dx}}{f^2} \\ &= \frac{1}{(e^\zeta - 1)f} \frac{d\zeta}{dx} \left( -(\zeta - 1)e^\zeta - 1 \right) \end{aligned}$$

and since  $-(\zeta - 1)e^\zeta - 1$  is strictly negative for  $\zeta > 0$  we get that  $g(x)$  is log-concave  $\square$

**Lemma 14**  $\zeta(x)$  is concave as a function of  $x$ .

**Proof:** We begin with (41). We note from (24) that the denominator

$$(e^\zeta - 1)^2 - \zeta^2 e^\zeta \geq 0.$$

Then we have

$$\frac{d^2 \zeta}{dx^2} = \frac{2(1 + \zeta) + e^\zeta(-6 - e^{2\zeta}(2 + \zeta(\zeta - 4)) + \zeta^2(5 + \zeta(\zeta + 2)) + e^\zeta(6 - 2\zeta(2\zeta + 3)))}{((e^\zeta - 1)^2 - \zeta^2 e^\zeta)^2} \frac{d\zeta}{dx}.$$

Now let

$$\phi(u) = \sum_{n=0}^{\infty} \phi_n u^n = 2(1 + u) + \psi(u)$$

where

$$\psi(u) = \sum_{n=0}^{\infty} \psi_n u^n = e^u(-6 - e^{2u}(2 + u(u-4)) + u^2(5 + u(u+2)) + e^u(6 - 2u(2u+3))).$$

We check that  $\psi_0 = -2$  and  $\psi_1 = 0$  which implies that  $\phi_0 = \phi_1 = 0$ . One can finish the argument by checking that

$$\psi_n = -\frac{3^{n-2}(n^2 - 13n + 18) + 2^n(n^2 + 2n - 6) - (n^4 - 4n^3 + 10n^2 - 7n - 6)}{n!} \leq 0$$

for  $n \geq 2$ . This is simply a matter of checking for small values until the  $3^n$  term dominates.  $\square$   
Next let

$$H(u) = \log f(u) - u - 2 \log u = \log \left( \frac{e^u - u - 1}{u^2 e^u} \right)$$

**Lemma 15**  $H(u)$  is convex as a function of  $u$ .

**Proof:**

$$\begin{aligned} \frac{d^2}{du^2} H(u) &= \frac{d}{du} \left( \frac{e^u - 1}{f(u)} - 1 - \frac{2}{u} \right) \\ &= \frac{e^u(e^u - 1 - u) - (e^u - 1)^2}{f^2(u)} + \frac{2}{u^2} \\ &= \frac{e^u - 1 - ue^u}{f^2(u)} + \frac{2}{u^2} \\ &= \frac{u^2(e^u - 1 - ue^u) + 2(e^u - 1 - u)^2}{u^2 f^2(u)} \\ &= \frac{2e^{2u} + u^2 e^u + u^2 + 4u + 2 - u^3 e^u - 4ue^u - 4e^u}{u^2 f^2(u)} \end{aligned}$$

Let

$$\phi(u) = 2e^{2u} + u^2 e^u + u^2 + 4u + 2 - u^3 e^u - 4ue^u - 4e^u = \sum_{n=0}^{\infty} \phi_n u^n.$$

Direct computation gives  $\phi_0 = \phi_1 = \phi_2 = 0$  and for  $n \geq 3$

$$\phi_n = \frac{1}{n!} (2^{n+1} + n(n-1) - n(n-1)(n-2) - 4n - 4).$$

One can then check that  $\phi_3 = \phi_4 = \phi_5 = 0 < \phi_n$  for  $n \geq 6$ . Thus  $\frac{d^2}{du^2} H(u) \geq 0$  implying that  $H(u)$  is convex.  $\square$

## 7.2 The case $m = n$

We will first prove Theorem 12 under the assumption that  $m = n$  and then in Sections 7.3 and 7.4 we will extend the result to arbitrary  $m$ . We will as usual prove that Hall's Condition holds **whp**. We will therefore estimate the probability of the existence of sets  $A, B$  where  $|A| = k$  and  $|B| = k - 1$  such that  $N_{\Gamma}(A) \subseteq B$ . Here  $N_{\Gamma}(S)$  is the set of neighbours of  $S$  in  $\Gamma$ . We call such a pair of sets, a *witness* to the non-existence of a perfect matching. There are two possibilities to

consider: (i)  $A \subseteq L$  and  $B \subseteq R$  or (ii)  $A \subseteq R$  and  $B \subseteq L$ . We deal with both cases in order to help extend the results to  $m \neq n$ . We observe that if there exist a pair  $A, B$  then there exist a minimal pair and in this case each  $b \in B$  has at least two neighbours in  $A$ . We deal first with the existence probability for a witness in Case (i) and leave Case (ii) until Section 7.2.2. We then combine these results to finish the case  $m = n$  in Section 7.2.3.

### 7.2.1 Case (i)

We consider a minimal witness  $A, B$  with  $A \subseteq L$ . Here we mean that  $A$  is minimal with respect to inclusion. We observe that each vertex in  $N(S)$  has degree at least 2 in  $S$ . For if  $v$  has a unique neighbour  $w$  in  $A$  then  $A \setminus \{w\}, B \setminus \{v\}$  is also a witness. We will need the estimates below for  $m \neq n$  later and so we begin with  $m$  as a parameter: We estimate

$$\pi_L(k, D) = \mathbb{P}(\exists A, B : |A| = k, |B| = k - 1, N_\Gamma(A) = B, d(B) = D, d_A(b) \geq 2, b \in B) \leq$$

$$O(n^{1/2}) \binom{n}{k} \binom{m}{k-1} \sum_{\substack{2 \leq x_b \leq d_b, b \in [k-1] \\ \sum_b x_b = kd \\ \sum_{b \in [k-1]} d_b = D \\ \sum_{b \notin [k-1]} d_b = dn - D}} \prod_{b=1}^m \frac{z^{d_b}}{d_b! f(z)} \binom{d_b}{x_b} (kd)! \prod_{i=0}^{dk-1} \frac{1}{dn-i} = \quad (43)$$

$$O(n^{1/2}) \binom{n}{k} \binom{m}{k-1} \frac{(d(n-k))! (kd)! z^{dn}}{(dn)! f(z)^m} \times$$

$$\left( \sum_{\substack{2 \leq x_b, b \in [k-1] \\ \sum_b x_b = kd}} \prod_{b=1}^{k-1} \frac{1}{x_b!} \right) \left( \sum_{\substack{2 \leq d_b, b \notin [k-1] \\ \sum_b d_b = dn - D}} \prod_{b=k}^m \frac{1}{d_b!} \right) \left( \sum_{\substack{0 \leq y_b, b \in [k-1] \\ \sum_b y_b = D - kd}} \prod_{b=1}^{k-1} \frac{1}{y_b!} \right) =$$

$$O(n^{1/2}) \binom{n}{k} \binom{m}{k-1} \frac{(d(n-k))! (kd)! z^{dn}}{(dn)! f(z)^m} \times$$

$$\left( [u^{kd}] (e^u - 1 - u)^{k-1} \right) \left( [u^{dn-D}] (e^u - 1 - u)^{m-k+1} \right) \left( [u^{D-kd}] e^{u(k-1)} \right) \leq \quad (44)$$

$$O(n^{1/2}) \binom{n}{k} \binom{m}{k-1} \frac{(d(n-k))! (kd)! z^{dn}}{(dn)! f(z)^m} \frac{f(z)^{k-1}}{z^{kd}} \frac{f(\zeta_1)^{m-k+1}}{\zeta_1^{dn-D}} \frac{(k-1)^{D-dk}}{(D-kd)!} \leq \quad (45)$$

$$O\left(\frac{k}{m^{1/2}}\right) \frac{\binom{n}{k} \binom{m}{k}}{\binom{dn}{dk}} \left( \frac{z^d}{f(z)^{\frac{m-k}{n-k}}} \frac{f(\zeta_1)^{\frac{m-k}{n-k}}}{\zeta_1^{\frac{dn-D}{n-k}}} \right)^{n-k} \left( \frac{ek}{D-dk} \right)^{D-dk}.$$

Putting  $k = an$  and  $m = \beta n$  and  $h(x) = x^x(1-x)^{1-x}$  we obtain

$$\pi_L(k, D) = O\left(\frac{k}{n^{1/2}}\right) \left(\frac{h(a)^{d-1}}{h(a/\beta)^\beta}\right)^n \left(\frac{z^d}{f(z)^{\frac{\beta-a}{1-a}}} \frac{f(\zeta_1)^{\frac{\beta-a}{1-a}}}{\zeta_1^{d-x}} \left(\frac{e \frac{a}{1-a}}{x}\right)^x\right)^{n-k}. \quad (46)$$

Here  $\zeta_1 = \zeta(y)$  where  $y = \frac{dn-D}{m-k+1} \geq 2$  due to our minimum degree assumption for  $R$ . Then let  $x = d - y$  where  $0 \leq x \leq d - 2$ .

**Explanation of (43):** Choose sets  $A, B$  in  $\binom{n}{k} \binom{m}{k-1}$  ways. Choose degrees  $d_b, b \in R$  with probability  $O(n^{1/2}) \prod_{b=1}^m \frac{z^{d_b}}{d_b! f(z)}$  such that  $\sum_{b \in B} d_b = D, \sum_{b \notin B} d_b = dn - D$  for some  $D \geq 2(k-1)$ . Choose the degrees  $x_a, a \in A$  in the sub-graph induced by  $A \cup B$ . Having fixed the degree sequence, we swap to the configuration model. Choose the configuration points associated the  $x_a, a \in A$  in

$\prod_{a \in A} \binom{d}{x_a}$  ways. Assign these  $D$  choices of points points associated with  $A$  in  $D!$  ways. Then multiply by the probability  $(kd)! \prod_{i=0}^{kd-1} \frac{1}{dn-i}$  of a given pairing of points in  $A$ .

**Explanation of (44) to (45):** If  $A(x) = \sum_{n=0}^{\infty} a_n x^n$  where  $a_n \geq 0$  for  $n \geq 0$  then  $a_n \leq A(\zeta)/\zeta^n$  for any positive  $\zeta$  and  $A(\zeta)/\zeta^n$  is minimised at  $\zeta$  satisfying  $\zeta A'(\zeta)/A(\zeta) = n$ .

We now assume that  $m = n$  for the remainder of the section. In which case (46) becomes

$$\pi_L(k, D) = O\left(\frac{k}{n^{1/2}}\right) \left(h(a)^{(d-2)n}\right) \left(\frac{z^d}{f(z)} \frac{f(\zeta_1)}{\zeta_1^{d-x}} \left(\frac{e^{\frac{a}{1-a}}}{x}\right)^x\right)^{n-k}. \quad (47)$$

Observe (see (40)) that

$$\frac{z^d}{f(z)} \frac{f(\zeta_1)}{\zeta_1^{d-x}} = \frac{d^d}{g(d)} \frac{g(d-x)}{(d-x)^{d-x}}$$

where  $g(x)$  is as defined in Lemma 13.

It follows from (42) that

$$-\log\left(\frac{g(d-x)}{g(d)}\right) = \int_{d-x}^d \frac{d}{dt} \log(g(t)) dt \geq \int_{d-x}^d (\log(1 + \zeta e^{-\zeta}) + 1) dt. \quad (48)$$

Now  $\zeta e^{-\zeta} \leq e^{-1}$  which implies that  $\log(1 + \zeta e^{-\zeta}) \geq \zeta e^{-\zeta}/10$ . Also,

$$\frac{\zeta(t)}{t} = 1 - \frac{\zeta}{e^\zeta - 1} \geq 1 - \frac{2}{\zeta + 2} = \frac{\zeta}{\zeta + 2}.$$

And so  $\zeta \leq t \leq \zeta + 2$ . Thus

$$\int_{d-x}^d (\log(1 + \zeta e^{-\zeta}) + 1) dt \geq \int_{d-x}^d \left(1 + \frac{t-2}{10e^d}\right) dt = x + \frac{x(2d-4-x)}{20e^d}.$$

This implies that  $\frac{g(d-x)}{g(d)} \leq e^{-x} \psi(x)$  where  $\psi(x) = e^{-\varepsilon_d(2d-4-x)x}$  and  $\varepsilon_d = \frac{1}{20e^d}$ . Note that  $d-x \geq 2$  and so  $\psi(x) \leq e^{-(d-2)\varepsilon_d x}$  in the range of interest. Plugging this into the last parenthesis of (47) gives

$$\pi_L(k, D) = O\left(\frac{k}{n^{1/2}}\right) \psi(x)^n \left(h(a)^{d-2} \left(d^d \frac{1}{(d-x)^{d-x}} \left(\frac{a}{x}\right)^x\right)^{1-a}\right)^n \quad (49)$$

This immediately yields

$$A_{50} = \sum_{k=\varepsilon_L n}^{n(1-2/d)} \sum_{D=dk}^{dk+n^{1/5}} \pi_L(k, D) \leq \sum_{k=\varepsilon_L n}^{n(1-2/d)} \sum_{D=dk}^{dk+n^{1/5}} O\left(\frac{k \log n}{n^{1/2}}\right) h(a)^{(d-2)n} e^{o(n)} = o(1). \quad (50)$$

We use the notation  $A_{50}$  so that the reader can easily refer back to the equation giving its definition.

We will work with  $D \leq k \log n$  because it is easy to show that **whp** the maximum degree in  $\Gamma$  is  $o(\log n)$ . The bound for  $A_{50}$  comes from (49), using the fact that  $h(a)$  is bounded away from 1 and  $x = o(1)$  in this summation.  $A_{50}$  is the first of several sums that together show the unlikelihood chance of a witness. We will display them as they become available and use them in Sections 7.2.3, 7.3 and 7.4.

The main term  $h(a)^{d-2} \left( d^d \frac{1}{(d-x)^{d-x}} \left( \frac{a}{x} \right)^x \right)^{1-a}$  in (49) is maximized when  $x = ad$ , provided  $ad \leq d-2$  or  $k \leq n(1 - \frac{2}{d})$ . This in turn gives

$$\begin{aligned} \pi_L(k, D) &= O\left(\frac{k}{n^{1/2}}\right) \psi(x)^n \left( h(a)^{d-2} \left( d^d \frac{1}{(d-ad)^{d-ad}} \frac{1}{((1-a)d)^{ad}} \right)^{1-a} \right)^n \\ &= O\left(\frac{k}{n^{1/2}}\right) \psi(x)^n \left( h(a)^{d-2} \left( \frac{d^d}{d^{d-ad} d^{ad}} (1-a)^{-d} \right)^{1-a} \right)^n \\ &\leq O\left(\frac{k}{n^{1/2}}\right) \psi(x)^n \left( a^{a(d-2)} (1-a)^{-2(1-a)} \right)^n \end{aligned} \quad (51)$$

The function  $\rho(a) = a^{a(d-2)}(1-a)^{-2(1-a)}$  is at most 1 and is log-convex in  $a$  on  $[0, 1 - \frac{2}{d}]$ . Indeed, if  $L_1(a) = \log \rho(a)$  then

$$\frac{dL_1}{da} = d + (d-2) \log a + 2 \log(1-a) \quad (52)$$

$$\frac{d^2 L_1}{da^2} = \frac{d-2-da}{a(1-a)} \quad (53)$$

We have  $L_1(0) = 0$  and  $L_1'(0) = -\infty$ . It follows that for every  $K > 0$  there exists a constant  $\varepsilon_L(K) > 0$  such that

$$\rho(a) \leq e^{-Ka} \quad \text{for } \rho \leq \varepsilon_L(K). \quad (54)$$

We let  $\varepsilon_L = \varepsilon_L(1)$ .

Furthermore  $(1 - \frac{2}{d})^{(d-2)^2/d} (\frac{2}{d})^{-4/d}$  decreases in  $d$  and is  $\leq 1$  for  $d \geq 4$ .

We can immediately write

$$A_{55} = \sum_{k=1}^{n^{1/5}} \sum_{D=dk}^{k \log n} \pi_L(k, D) = \sum_{k=1}^{n^{1/5}} \sum_{D=dk}^{k \log n} O\left(\frac{k \log n}{n^{1/2}}\right) = o(1). \quad (55)$$

The bound for  $A_{55}$  is derived from (51) using  $\psi(x), \rho(a) \leq 1$ .

Along the same lines we have

$$A_{56} = \sum_{k=n^{1/5}}^{\varepsilon_L n} \sum_{D=dk}^{k \log n} \pi_L(k, D) = \sum_{k=n^{1/5}}^{\varepsilon_L n} \sum_{D=dk}^{k \log n} O\left(\frac{k \log n}{n^{1/2}}\right) e^{-k} = o(1). \quad (56)$$

The bound for  $A_{56}$  comes from (51) and (54).

Similarly,

$$A_{57} = \sum_{k=\varepsilon_L n}^{n(1-2/d)} \sum_{D=dk+n^{1/5}}^{k \log n} \pi_L(k, D) = \sum_{k=\varepsilon_L n}^{n(1-2/d)} \sum_{D=dk+n^{1/5}}^{k \log n} O\left(\frac{k \log n}{n^{1/2}}\right) \psi(n^{-4/5})^n \rho(a)^n = o(1). \quad (57)$$

The bound for  $A_{57}$  comes from (51) using the fact that  $\rho(a) \leq 1$  and  $x \geq n^{-4/5}$  in this summation.

For  $a \geq 1 - \frac{2}{d}$  the maximising value for  $x$  in (49) is at  $x = d-2$  (recall that  $0 \leq x \leq d-2$ ), so plugging into (49) gives

$$\begin{aligned}
\pi_L(k, D) &= O\left(\frac{k}{n^{1/2}}\right) \left( h(a)^{d-2} \left( d^d \frac{1}{2^2} \left( \frac{a}{d-2} \right)^{d-2} \right)^{1-a} \right)^n \\
&= O\left(\frac{k}{n^{1/2}}\right) \left( a^{d-2} \left( \frac{d^d}{(d-2)^{d-2} 2^2} \right)^{1-a} \right)^n
\end{aligned} \tag{58}$$

Let  $L_2(a) = \log \left( a^{d-2} \left( \frac{d^d}{(d-2)^{d-2} 2^2} \right)^{1-a} \right)$ . Then

$$\begin{aligned}
\frac{d}{da} L_2(a) &= \frac{d}{da} \left( (d-2) \log a + (1-a) \log \left( \frac{d^d}{(d-2)^{d-2} 2^2} \right) \right) \\
&= \frac{d-2}{a} - d \log d + (d-2) \log(d-2) + 2 \log(2)
\end{aligned}$$

Assume for now that  $d \geq 6$ . Then the derivative with respect to  $d$ , for of the last expression is

$$\frac{1}{a} - \log \left( \frac{d}{d-2} \right) \geq 1 - \log \left( \frac{6}{4} \right) > 0$$

so it takes a minimum at  $d = 6$  with a value

$$\frac{4}{a} - 6 \log 6 + 4 \log 4 + 2 \log 2 \geq 0.18 > 0.$$

Now  $L_2(1) = 0$  and so for  $a \geq 1 - \frac{2}{d}$  and  $d \geq 6$  we have

$$A_{59} = \sum_{k=n(1-2/d)}^{n-n^{1/5}} \sum_{D=dk}^{k \log n} \pi_L(k, D) \leq \sum_{k=n(1-2/d)}^{n-n^{1/5}} \sum_{D=dk}^{k \log n} O\left(\frac{k \log n}{n^{1/2}}\right) e^{-.18(n-k)} = o(1). \tag{59}$$

For  $d = 4, 5$  we use the following

**Claim 16** For  $x \geq 2$  we have  $\frac{f(\zeta(x))}{\zeta(x)^x} \leq \frac{3}{4}$ .

Substituting this into (47) gives

$$\pi_L(k, D) = O\left(\frac{k}{n^{1/2}}\right) \left( h(a)^{d-2} \left( \frac{z^d}{f(z)} \frac{3}{4} \left( \frac{e \frac{a}{1-a}}{x} \right)^x \right)^{1-a} \right)^n \tag{60}$$

The maximum of  $L_3(x) = h(a)^{d-2} \left( \frac{z^d}{f(z)} \frac{3}{4} \left( \frac{e \frac{a}{1-a}}{x} \right)^x \right)^{1-a}$  is taken when either  $x = \frac{a}{1-a}$  for  $a \in [1 - \frac{2}{d}, 1 - \frac{1}{d-1}]$  or at  $x = d-2$  for  $a \in [1 - \frac{1}{d-1}, 1]$ .

So for  $a \in [1 - \frac{2}{d}, 1 - \frac{1}{d-1}]$  we get

$$\pi_L(k, D) \leq O\left(\frac{k}{n^{1/2}}\right) \left( h(a)^{d-2} e^a \left( \frac{3z^d}{4f(z)} \right)^{1-a} \right)^n.$$

The expression  $L_4(a) = h(a)^{d-2} e^a \left( \frac{3z^d}{4f(z)} \right)^{1-a}$  is log-convex on  $[1 - \frac{2}{d}, 1 - \frac{1}{d-1}]$  and  $\log L_4 \leq -1/5$  at both ends of the interval for both  $d = 4$  and  $d = 5$ . We can therefore write

$$A'_{61} = \sum_{k=n(1-2/d)}^{n(1-1/(d-1))} \sum_{D=dk}^{k \log n} \pi_L(k, D) \leq \sum_{k=n(1-2/d)}^{n(1-1/(d-1))} \sum_{D=dk}^{k \log n} O\left(\frac{k \log n}{n^{1/2}}\right) e^{-n/5} = o(1). \tag{61}$$

for  $a \in [1 - \frac{2}{d}, 1 - \frac{1}{d-1}]$ .

For  $a \in [1 - \frac{1}{d-1}, 1]$  from (60) we get

$$\begin{aligned}\pi_L(k, D) &= O\left(\frac{k}{n^{1/2}}\right) \left(h(a)^{d-2} e^a \left(\frac{3z^d}{4f(z)}\right)^{1-a} \left(\frac{e^{\frac{a}{1-a}}}{d-2}\right)^{(1-a)(d-2)}\right)^n \\ &= O\left(\frac{k}{n^{1/2}}\right) \left(a^{d-2} \left(\frac{3z^d e^{d-2}}{4f(z)(d-2)^{d-2}}\right)^{1-a}\right)^n\end{aligned}$$

The expression  $L_5(a) = a^{d-2} \left(\frac{3z^d e^{d-2}}{4f(z)(d-2)^{d-2}}\right)^{1-a}$  is log-convex on  $[1 - \frac{1}{d-1}, 1]$ . We have  $\log L_5(1 - 1/d) \leq -1/10$  for both  $d = 4$  and  $d = 5$ . The derivative of  $\log L_5$  at  $a = 1$  is at least 1.9 for both  $d = 4$  and  $d = 5$ . Consequently

$$A''_{62} = \sum_{k=n(1-1/(d-1))}^{n-n^{1/5}} \sum_{D=dk}^{k \log n} \pi_L(k, D) \leq \sum_{k=n(1-2/d)}^{n-n^{1/5}} \sum_{D=dk}^{k \log n} O\left(\frac{k \log n}{n^{1/2}}\right) e^{-n^{1/5}/10} = o(1). \quad (62)$$

for  $a \in [1 - \frac{1}{d-1}, 1]$ .

**Proof of Claim 16:** Recall from Lemma 14 that  $\zeta(x)$  is concave and thus  $\frac{\zeta(x)-\zeta(2)}{x-2}$  is decreasing. Since  $\zeta(2) = 0$  we have

$$\begin{aligned}\frac{\zeta(x)}{x-2} &= \frac{\zeta(x) - \zeta(2)}{x-2} \\ &\leq \lim_{x \rightarrow 2} \frac{d}{dx} \zeta(x) \\ &= \lim_{z \rightarrow 0} \frac{e^z - z - 1}{(e^z - 1)^2 - z^2 e^z} \\ &= 3\end{aligned}$$

Thus we have that  $\zeta(x) \leq 3(x-2)$  for all  $x$ . For  $x \geq 3$  we can upper bound

$$\frac{f(\zeta(x))}{\zeta(x)^x} \leq \frac{f(x)}{x^x} \leq \frac{e^x}{x^x} \leq \frac{e^3}{3^3} \leq \frac{3}{4}$$

For  $x \in [2, 3]$  we have that  $3(x-2) \leq x$  and so we can bound

$$\frac{f(\zeta(x))}{\zeta(x)^x} \leq \frac{f(3(x-2))}{(3(x-2))^x}$$

Taking the logarithm of this expression and substituting  $u = 3(x-2)$  we get

$$\log f(u) - \left(\frac{u}{3} + 2\right) \log u = u - \frac{u}{3} \log(u) + (\log f(u) - u - 2 \log(u)) = u - \frac{u}{3} \log(u) + H(u) \quad (63)$$

where  $H(u)$  is from Lemma 15.

Since  $H(u)$  is convex we have

$$H(u) \leq H(0) + u \frac{H(3) - H(0)}{3}$$

Plugging this into (63) we get

$$u - \frac{u}{3} \log u + H(0) + u \frac{H(3) - H(0)}{3} = H(0) + \frac{1}{3} ((H(3) - H(0)) + 3)u - u \log u$$

which is concave in  $u$  and takes a maximum value of  $H(0) + u/3$  when

$$u = \exp(H(3) - H(0) + 3 - 1) = \frac{e^3 - 4}{18e} \leq 0.33$$

Pluggin this back in we see that for  $x \in [2, 3]$ , which is  $u \in [0, 3]$  we have

$$\frac{f(\zeta(x))}{\zeta(x)^x} \leq \exp(H(0) + .11) = \exp(0.11)/2 < \frac{3}{4}$$

□

## 7.2.2 Case (ii)

Now let us estimate the probability of a violation of Hall's condition with  $A \subseteq R$ . We once again begin with arbitrary  $m$ . Let

$$\begin{aligned} \pi_R(k, D) &= \mathbb{P}(\exists A \subseteq R, B \subseteq L : |A| = k, |B| = k - 1, N_\Gamma(A) \subseteq B, d_B(b) \geq 2, b \in B, d_R(A) = D) \leq \\ O(n^{1/2}) &\binom{m}{k} \binom{n}{k-1} \sum_{\substack{2 \leq d_a, a \in [m] \\ 2 \leq x_b \leq d, b \in [k-1] \\ \sum_{a \in [k]} d_a = \sum_{b \in [k-1]} x_b = D \\ \sum_{a \notin [k]} d_a = dn - D}} \prod_{a=1}^m \frac{z^{d_a}}{d_a! f(z)} \prod_{b=1}^{k-1} \binom{d}{x_b} D! \prod_{i=0}^{D-1} \frac{1}{dn - i} = \quad (64) \\ O\left(\frac{k}{m^{1/2}}\right) &\binom{n}{k} \binom{m}{k} \frac{z^{dn} D! (dn - D)!}{f(z)^m (dn)!} \\ &\times \left( \sum_{\substack{2 \leq d_a, a \in [k] \\ \sum_a d_a = D}} \prod_{a=1}^k \frac{1}{d_a!} \right) \left( \sum_{\substack{2 \leq d_a, a \notin [k] \\ \sum_a d_a = dn - D}} \prod_{a=k+1}^n \frac{1}{d_a!} \right) \left( \sum_{\substack{2 \leq x_b \leq d, b \in [k-1] \\ \sum_b x_b = D}} \prod_{b=1}^{k-1} \binom{d}{x_b} \right) = \\ O\left(\frac{k}{m^{1/2}}\right) &\binom{n}{k} \binom{m}{k} \frac{z^{dn}}{f(z)^m} \frac{1}{\binom{dn}{D}} \\ &\times \left( [u^D] (e^u - 1 - u)^k \right) \left( [u^{dn-D}] (e^u - 1 - u)^{m-k} \right) \left( [u^D] ((1+u)^d - (1+du))^k \right) \leq \\ O\left(\frac{k}{m^{1/2}}\right) &\binom{n}{k} \binom{m}{k} \frac{z^{dn}}{f(z)^m} \frac{1}{\binom{dn}{D}} \frac{f(\zeta_1)^k}{\zeta_1^D} \frac{f(\zeta_2)^{m-k}}{\zeta_2^{dn-D}} \binom{dk}{D} = \\ &\text{where } \zeta_1 = \zeta(D/k) \text{ and } \zeta_2 = \zeta\left(\frac{dn-D}{m-k}\right) - \text{actually any value for } \zeta_1, \zeta_2 \text{ is valid -} \end{aligned}$$

$$O\left(\frac{k}{m^{1/2}}\right) \binom{n}{k} \binom{m}{k} \frac{\binom{dk}{D}}{\binom{dn}{D}} \left(\frac{f(\zeta_1)}{f(z)}\right)^k \left(\frac{f(\zeta_2)}{f(z)}\right)^{m-k} \left(\frac{z}{\zeta_1}\right)^D \left(\frac{z}{\zeta_2}\right)^{dn-D} \quad (65)$$

$$= O\left(\frac{k}{m^{1/2}}\right) \left( \frac{h(\theta a/d)^d}{h(a)h(a/\beta)^\beta h(\theta/d)^{ad}} \left(\frac{f(\zeta_1)}{f(z)}\right)^a \left(\frac{f(\zeta_2)}{f(z)}\right)^{\beta-a} \left(\frac{z}{\zeta_1}\right)^{\theta a} \left(\frac{z}{\zeta_2}\right)^{d-\theta a} \right)^n \quad (66)$$

$$= O\left(\frac{k}{m^{1/2}}\right) \left( \frac{h(\theta a/d)^d}{h(a)h(a/\beta)^\beta h(\theta/d)^{ad}} \frac{z^d}{f(z)^\beta} \frac{f(\zeta_1)^a}{\zeta_1^{\theta a}} \frac{f(\zeta_2)^{\beta-a}}{\zeta_2^{d-\theta a}} \right)^n \quad (67)$$

where  $a = k/n$ ,  $m = \beta n$  and  $D = \theta k$ .

**Explanation of (64):** Choose sets  $A, B$  in  $\binom{[m]}{k} \binom{[n]}{k-1}$  ways. Choose degrees  $d_a, a \in R$  with probability  $O(n^{1/2}) \prod_{a=1}^n \frac{z^{d_a}}{d_a! f(z)}$  such that  $\sum_{a \in A} d_a = D, \sum_{a \notin A} d_a = dn - D$  for some  $D \geq 2k$ . Choose the degrees  $x_b, b \in B$  in the sub-graph induced by  $A \cup B$ . Having fixed the degree sequence, swap to the configuration model [5]. Choose the configuration points associated the  $x_b, b \in B$  in  $\prod_{b \in B} \binom{d}{x_b}$  ways. Then multiply by the probability  $D! \prod_{i=0}^{D-1} \frac{1}{d^{n-i}}$  of a given pairing of points in  $A$ .

We now assume that  $m = n$  for the remainder of the section. In which case (67) becomes

$$\pi_R(k, D) \leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{h(\theta a/d)^d}{h(a)^2 h(\theta/d)^{ad}} \frac{z^d}{f(z)} \frac{f(\zeta_1)^a}{\zeta_1^{\theta a}} \frac{f(\zeta_2)^{1-a}}{\zeta_2^{d-\theta a}}\right)^n \quad (68)$$

It follows from Lemma 13 that we can upper bound

$$\begin{aligned} \frac{z^d}{f(z)} \left(\frac{f(\zeta_1)}{\zeta_1^\theta}\right)^a \left(\frac{f(\zeta_2)}{\zeta_2^{\frac{d-a\theta}{1-a}}}\right)^{1-a} &= \frac{d^d}{g(d)} \frac{g(\theta)^a g\left(\frac{d-a\theta}{1-a}\right)^{1-a}}{\theta^{a\theta} \left(\frac{d-a\theta}{1-a}\right)^{d-a\theta}} \\ &\leq \frac{g(a\theta + (1-a)\frac{d-a\theta}{1-a})}{g(d)} \frac{d^d}{\theta^{a\theta} \left(\frac{d-a\theta}{1-a}\right)^{d-a\theta}} \\ &= \frac{a^{a\theta} (1-a)^{d-a\theta}}{\left(\left(\frac{a\theta}{d}\right)^{\frac{a\theta}{d}} \left(1 - \frac{a\theta}{d}\right)^{1-\frac{a\theta}{d}}\right)^d} \\ &= \frac{a^{a\theta} (1-a)^{d-a\theta}}{h\left(\frac{a\theta}{d}\right)^d} \end{aligned}$$

Plugging this into (68) gives

$$\pi_R(k, D) \leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{a^{a\theta} (1-a)^{d-a\theta}}{h(a)^2 h\left(\frac{\theta}{d}\right)^{ad}}\right)^n \quad (69)$$

Now let  $R_1(\theta) = \log\left(\frac{a^{a\theta} (1-a)^{d-a\theta}}{h(a)^2 h\left(\frac{\theta}{d}\right)^{ad}}\right)$ . Then

$$R_1'(\theta) = a \log a - a \log(1-a) - a \log \theta + a \log(d-\theta).$$

$$R_1''(\theta) = -\frac{ad}{\theta(d-\theta)} < 0.$$

Thus  $R_1(\theta)$  is concave and is maximized when  $\theta = ad$ . Because  $\theta \geq 2$  we can only use this for  $a \geq 2/d$ . Thus if  $a \in [\frac{2}{d}, 1]$  then

$$\begin{aligned} \pi_R(k, D) &\leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{a^{a^2 d} (1-a)^{d-a^2 d}}{h(a)^{2+ad}}\right)^n \\ &= O\left(\frac{k}{n^{1/2}}\right) \left(a^{a^2 d - a(2+ad)} (1-a)^{d-a^2 d - (1-a)(2+ad)}\right)^n \\ &= O\left(\frac{k}{n^{1/2}}\right) \left(a^{-2a} (1-a)^{(d-2)(1-a)}\right)^n \\ &= O\left(\frac{k}{n^{1/2}}\right) \rho (1-a)^n \end{aligned} \quad (70)$$

where the function  $\rho$  is defined following (51).

We find that

$$\rho(1 - 2/d) = \left( \frac{d^4}{16} \left( 1 - \frac{2}{d} \right)^{(d-2)^2} \right)^{1/d} \leq .9 \text{ for } d \geq 5. \quad (71)$$

Now  $\rho(1 - 2/d) < 1$  for  $d \geq 5$  and  $\rho(1/2) = 1$ . So,

$$B_{72} = \sum_{k=2n/d}^{n-n^{3/4}} \sum_{D=2k}^{k \log n} \pi_R(k, D) \leq \sum_{k=2n/d}^{n-n^{3/4}} O\left(\frac{k}{n^{1/2}}\right) \sum_{D=2k}^{k \log n} (.9)^{n^{3/4}} \leq e^{-\Omega(n^{3/4})} \quad \text{for } d \geq 5. \quad (72)$$

$$B_{73} = \sum_{k=2.01n/d}^{n-n^{3/4}} \sum_{D=2k}^{k \log n} \pi_R(k, D) \leq \sum_{k=2.01n/d}^{n-n^{3/4}} O\left(\frac{k}{n^{1/2}}\right) \sum_{D=2k}^{k \log n} (.9)^{n^{3/4}} \leq e^{-\Omega(n^{3/4})} \quad \text{for } d = 4. \quad (73)$$

When  $a \leq \frac{2}{d}$  the expression in (69) is maximized at  $\theta = 2$ . Then

$$\begin{aligned} \pi_R(k, D) &\leq O\left(\frac{k}{n^{1/2}}\right) \left( \frac{a^{2a}(1-a)^{d-2a}}{h(a)^2 h\left(\frac{2}{d}\right)^{ad}} \right)^n \\ &= O\left(\frac{k}{n^{1/2}}\right) \left( \frac{(1-a)^{d-2a-2(1-a)}}{h\left(\frac{2}{d}\right)^{ad}} \right)^n \\ &= O\left(\frac{k}{n^{1/2}}\right) \left( \frac{(1-a)^{d-2}}{h\left(\frac{2}{d}\right)^{ad}} \right)^n \end{aligned}$$

Let  $R_2(a) = \log\left(\frac{(1-a)^{d-2}}{h\left(\frac{2}{d}\right)^{ad}}\right)$ . Then

$$\begin{aligned} R_2'(a) &= -\frac{d-2}{1-a} - d \log h(2/d) < 0 \quad \text{for } d \geq 6. \\ R_2''(a) &= -\frac{d-2}{(1-a)^2} < 0. \end{aligned}$$

Thus  $R_2(a)$  is strict concave and its maximum is taken at  $a = 0$  and  $R_2(a) \leq R_2(0)a$  for all  $a \in [0, \frac{2}{d}]$ . Furthermore,  $R_2(0) < -3/10$  for  $d \geq 6$ . It follows that if  $d \geq 6$  then

$$B_{74} = \sum_{k=0}^{n^{1/5}} \sum_{D=2k}^{k \log n} \pi_R(k, D) \leq \sum_{k=0}^{n^{1/5}} \sum_{D=2k}^{k \log n} O\left(\frac{k}{n^{1/2}}\right) = o(1). \quad (74)$$

$$B_{75} = \sum_{k=n^{3/4}}^{2n/d} \sum_{D=2k}^{k \log n} \pi_R(k, D) \leq \sum_{k=n^{3/4}}^{2n/d} \sum_{D=2k}^{k \log n} O\left(\frac{k}{n^{1/2}}\right) e^{-3n^{3/4}/10} = e^{-\Omega(n^{3/4})}. \quad (75)$$

For  $d = 4$  and  $d = 5$  we use a better bound on  $[u^D]((1+u)^d - 1 - du)^k$  in (44).

**Case 1:**  $d = 5$ .

$$\begin{aligned}
[u^D]((1+u)^5 - 1 - 5u)^k &= [u^D](10u^2 + 10u^2 + 5u^4 + u^5)^k \\
&= [u^{D-2k}](10 + 10u + 5u^2 + u^3)^k \\
&= 10^k [u^{D-2k}] \left(1 + u + \frac{u^2}{2} + \frac{u^3}{10}\right)^k \\
&\leq 10^k [u^{D-2k}] \left(1 + \frac{u}{2}\right)^{3k} \\
&= 10^k \frac{\binom{3k}{D-2k}}{2^{D-2k}}
\end{aligned}$$

Replacing the  $\frac{1}{h(\frac{\theta}{d})^{ad}}$  factor in (69) which comes from  $\binom{dk}{D}$  gives, for  $d=5$ ,

$$\begin{aligned}
\pi_R(k, D) &\leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{a^{a\theta}(1-a)^{5-a\theta}}{h(a)^2}\right)^n \left(\frac{10}{2^{\theta-2}h\left(\frac{\theta-2}{3}\right)^3}\right)^k \\
&= O\left(\frac{k}{n^{1/2}}\right) \left(\frac{10^a a^{a(\theta-2)}(1-a)^{5-2-a(\theta-2)}}{\left(2^{\theta-2}h\left(\frac{\theta-2}{3}\right)^3\right)^a}\right)^n \\
&= O\left(\frac{k}{n^{1/2}}\right) \left(10^a(1-a)^3 \left(\frac{\left(\frac{a}{1-a}\right)^{\frac{\theta-2}{3}}}{2^{\frac{\theta-2}{3}}h\left(\frac{\theta-2}{3}\right)}\right)^{3a}\right)^n \tag{76}
\end{aligned}$$

Let  $p(x) = \frac{q^x}{h(x)}$  for any  $x \in [0, 1]$ , note that if  $P(x) = \log p(x)$  then

$$\begin{aligned}
P'(x) &= \log q - \log x + \log(1-x) \\
P''(x) &= -\frac{1}{x} - \frac{1}{1-x} < 0
\end{aligned}$$

and so  $p(x)$  is maximized when  $\log q = \log\left(\frac{x}{1-x}\right)$  or  $x = \frac{q}{1+q}$  and the maximum value is  $1+q$

Thus from (76) we get

$$\pi_R(k, D) \leq O\left(\frac{k}{n^{1/2}}\right) \left(10^a(1-a)^3 \left(1 + \frac{a}{2(1-a)}\right)^{3a}\right)^n$$

Let  $R_3(a) = \log\left(10^a(1-a)^3 \left(1 + \frac{a}{2(1-a)}\right)^{3a}\right)$ . Then

$$\begin{aligned}
R_3'(a) &= \log 10 - \frac{6}{2-a} + 3 \log\left(\frac{2-a}{2-2a}\right) \\
R_3''(a) &= \frac{3a}{(2-a)^2(1-a)} > 0.
\end{aligned}$$

So  $R_3(a)$  is log-convex on  $[0, \frac{2}{5}]$ . We have  $R_3(0) = 1$  and  $H'(0) = \log 10 - 3 \leq -3/4$  and  $R_3(2/5) < -1/4$ . It follows that

$$B_{77} = \sum_{k=n^{3/4}}^{2n/d} \sum_{D=2k}^{k \log n} \pi_R(k, D) \leq \sum_{k=n^{3/4}}^{2n/d} \sum_{D=2k}^{k \log n} O\left(\frac{k}{n^{1/2}}\right) e^{-3n^{3/4}/4} = e^{-\Omega(n^{3/4})}. \tag{77}$$

**Case 2:**  $d = 4$ .

$$\begin{aligned}
[u^D]((1+u)^4 - 1 - 4u)^k &= [u^{D-2k}](6 + 4u + u^2)^k \\
&= 6^k [u^{D-2k}] \left(1 + \frac{4}{6}u + \frac{u^2}{6}\right)^k \\
&\leq 6^k [u^{D-2k}] \left(1 + \frac{u}{2}\right)^k \\
&= 6^k \frac{\binom{2k}{D-2k}}{2^{D-2k}}.
\end{aligned}$$

$$\begin{aligned}
\pi_R(k, D) &\leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{a^{a\theta}(1-a)^{4-a\theta}}{h(a)^2}\right)^n \left(\frac{6}{2^{\theta-2}h\left(\frac{\theta-2}{2}\right)^2}\right)^k \\
&= O\left(\frac{k}{n^{1/2}}\right) \left(6^a a^{a(\theta-2)}(1-a)^{2-a(\theta-2)} \left(\frac{1}{2^{\frac{\theta-2}{2}}h\left(\frac{\theta-2}{2}\right)}\right)^{2a}\right)^n \\
&= O\left(\frac{k}{n^{1/2}}\right) \left(6^a(1-a)^2 \left(\frac{\left(\frac{a}{1-a}\right)^{\frac{\theta-2}{2}}}{2^{\frac{\theta-2}{2}}h\left(\frac{\theta-2}{2}\right)}\right)^{2a}\right)^n \\
&\leq O\left(\frac{k}{n^{1/2}}\right) \left(6^a(1-a)^2 \left(1 + \frac{a}{2(1-a)}\right)^{2a}\right)^n.
\end{aligned}$$

Now if  $R_4(a) = \log\left(6^a(1-a)^2\left(1 + \frac{a}{2(1-a)}\right)^{2a}\right)$  then

$$\begin{aligned}
R_4'(a) &= \log 6 - \frac{4}{2-a} + 2 \log\left(\frac{2-a}{2-2a}\right) \\
R_4''(a) &= \frac{2a}{(2-a)^2(1-a)} > 0.
\end{aligned}$$

Thus  $R_4$  is log-convex on  $[0, \frac{2.01}{4}]$ . We have  $R_4(0) = 1$  and  $R_4'(0) = \log 6 - 2 \leq -1/5$  and  $R_4(2.01/4) < -1/20$ . It follows that

$$B_{78} = \sum_{k=n^{3/4}}^{2.01n/d} \sum_{D=2k}^{k \log n} \pi_R(k, D) \leq \sum_{k=n^{3/4}}^{2.01n/d} \sum_{D=2k}^{k \log n} O\left(\frac{k}{n^{1/2}}\right) e^{-n^{3/4}/5} = e^{-\Omega(n^{3/4})}. \quad (78)$$

### 7.2.3 Finishing the case $m = n$

We observe first that the maximum degree in  $\Gamma$  is  $o(\log n)$  **whp**. Therefore.

$$\mathbb{P}(\mu(\Gamma) < n) \leq o(1) + A_{50} + A_{55} + A_{56} + A_{57} + A_{59} + B_{74}$$

where  $A_{59} = A'_{61} + A''_{62}$  in the case  $d = 4, 5$ .

**Explanation:** We can use  $B_{74}$  to handle  $k \geq n - n^{1/5}$  because if  $A, B$  are a witness with  $A \subseteq L$  then  $R \setminus B, L \setminus A$  are a witness, looking at things from the  $R$  side.

We point out for use in the next section that our bounds on  $A_{50}, A_{56}, A_{57}, A_{59}$  allow us to claim that we have

$$\sum_{k=n^{3/4}}^{n-n^{3/4}} \sum_{D=dk}^{k \log n} \pi_L(k, D) = O(e^{-\Omega(n^{3/4})}). \quad (79)$$

In the case  $d = 4, 5$  this requires us to replace  $n^{1/5}$  by  $n^{3/4}$  everywhere in (62).

Our bounds on  $B_{72}, B_{73}, B_{75}, B_{77}, B_{78}$  allow us to claim that

$$\sum_{k=n^{3/4}}^{n-n^{3/4}} \sum_{D=2k}^{k \log n} \pi_R(k, D) = O(e^{-\Omega(n^{3/4})}). \quad (80)$$

### 7.3 The case $m > n$

Let  $\mathcal{G}(n, m)$  denote the set of bipartite graphs with  $|L| = n, |R| = m$  that are  $d$ -regular on  $L$  and degree at least 2 on  $R$ . Here  $n \leq m \leq dn/2$ . In fact suppose first that  $m \leq \xi dn$  where  $\xi < 1/2$  is a constant. Suppose that  $G(n, m)$  is chosen uniformly at random from  $\mathcal{G}(n, m)$ .

Let  $d_i(n, m)$  denote the number of  $R$ -vertices of degree  $i \geq 2$  in  $G(n, m)$  and let  $D_i(n, m) = E(d_i(n, m))$ .

We define three events:

$$\mathcal{A}_1(n, m) = \left\{ G(n, m) : \exists i : |d_i(n, m) - D_i(n, m)| > n^{3/5}/i^3, 2 \leq i \leq \log^2 n \right\} \quad (81)$$

$$\mathcal{A}_2(n, m) = \left\{ G(n, m) : \exists i : d_i(n, m) \neq 0, i > \log^2 n \right\} \quad (82)$$

$$\mathcal{B}(n, m+1) = \left\{ G(n, m+1) : |d_2(n, m+1) - D_2(n, m+1)| > 2n^{3/5} \right\} \quad (83)$$

We argue next that if  $\mathcal{A}(n, m) = \mathcal{A}_1(n, m) \cup \mathcal{A}_2(n, m)$  then

$$\mathbb{P}(\mathcal{A}(n, m) \cup \mathcal{B}(n, m+1)) = e^{-\Omega(\log^2 n)}. \quad (84)$$

Indeed, for any  $t > 0$  we have

$$\mathbb{P}(|d_i(n, m) - D_i(n, m)| > t) \leq O(n^{1/2})\mathbb{P}(\text{Bin}(n, q_i) > t)$$

where  $q_i = \frac{z^i}{i!f(z)}$ .

We will now use the following bounds (see for example [1])

$$\mathbb{P}(|\text{Bin}(n, p) - np| \geq t) \leq 2e^{-t^2/n}, \quad (85)$$

$$\mathbb{P}(\text{Bin}(n, p) \geq \alpha np) \leq (e/\alpha)^{\alpha np}. \quad (86)$$

If  $i \leq \log^2 n$  then we can use (85) with  $t = n^{3/5}/i^3$  to deal with  $\mathcal{A}_1(n, m)$  and also with  $\mathcal{B}(n, m+1)$ . If  $i \geq \log^2 n$  then  $nq_i \leq e^{-\Omega(\log^2 n)}$ . We can therefore use (86) with  $\alpha = 1/nq_i$  to deal with  $\mathcal{A}_2(n, m)$ . This concludes the proof of (84).

Now consider a set of pairs  $X \subseteq \mathcal{G}(n, m) \times \mathcal{G}(n, m+1)$ . We place  $(G_1, G_2)$  into  $X$  if  $G_2$  is obtained from  $G_1$  in the following manner: Choose a vertex  $x \in R$  of degree at least four in  $G_1$ . Suppose that its neighbours are  $y_i, i = 1, 2, \dots, k$  in any order. To create  $G_2$  we (i) replace  $x$  by two vertices  $x$  and  $m+1$  and then (ii) let the neighbours of  $x$  in  $G_2$  be  $y_1, y_2$  and let the neighbours of  $m+1$  be  $y_3, \dots, y_k$ .

Note that if  $(G_1, G_2) \in X$  and  $G_1$  has a matching of  $L$  into  $R$  then so does  $G_2$ .

For  $G \in \mathcal{G}(n, m)$  let

$$\pi_1(G) = |\{G_2 : (G, G_2) \in X\}|$$

and for  $G \in \mathcal{G}(n, m+1)$  let

$$\pi_2(G) = |\{G_1 : (G_1, G) \in X\}|.$$

We note that if

$$\Sigma_1 = \sum_{i \geq 4} \binom{i}{2} D_i(n, m)$$

then

- $G \notin \mathcal{A}(n, m)$  implies that  $|\pi_1(G) - \Sigma_1| \leq O(n^{3/5})$ .
- $\pi_1(G) \leq \binom{m}{2}$  for all  $G \in \mathcal{G}(n, m)$ .
- $G \notin \mathcal{B}(n, m+1)$  implies that  $|\pi_2(G) - D_2(n, m+1)| \leq n^{3/5}$ .
- $\pi_2(G) \leq m+1$  for all  $G \in \mathcal{G}(n, m+1)$ .

We then note that

$$\begin{aligned} \Sigma_1 - O(n^{3/5}) &\leq \frac{|X|}{|\mathcal{G}(n, m)|} \leq \Sigma_1 + O(n^{3/5}) + m^2 e^{-\Omega(\log^2 n)} = \Sigma_1 + O(n^{3/5}). \\ D_2(n, m+1) - n^{3/5} &\leq \frac{|X|}{|\mathcal{G}(n, m+1)|} \leq D_2(n, m+1) + n^{3/5} + m e^{-\Omega(\log^2 n)}. \end{aligned}$$

Now let  $\mathcal{P}$  be a property such that if  $(G_1, G_2) \in X$  and  $G_2 \in \mathcal{P}$  then  $G_1 \in \mathcal{P}$ . Let  $(G_1, G_2)$  be chosen uniformly from  $X$  and let  $\mathbb{P}_X$  denote probabilities computed w.r.t. this choice. Let  $\mathcal{P}_m = \mathcal{P} \cap \mathcal{G}(n, m)$  and  $\mathcal{P}_{m+1} = \mathcal{P} \cap \mathcal{G}(n, m+1)$ . Then

$$\mathbb{P}_X(G_2 \in \mathcal{P}) \leq \mathbb{P}_X(G_1 \in \mathcal{P}) \leq \frac{|\mathcal{P}_m|(\Sigma_1 + O(n^{3/5})) + m|\mathcal{A}(n, m)|}{|X|}$$

and

$$\mathbb{P}_X(G_2 \in \mathcal{P}) \geq \frac{(|\mathcal{P}_{m+1}| - |\mathcal{B}(n, m+1)|)(D_2(n, m+1) - n^{3/5})}{|X|}$$

So,

$$\frac{(|\mathcal{P}_{m+1}| - |\mathcal{B}(n, m+1)|)(D_2(n, m+1) - n^{3/5})}{|\mathcal{G}(n, m+1)|(D_2(n, m+1) + n^{3/5} + m e^{-\Omega(\log^2 n)})} \leq \frac{|\mathcal{P}_m|(\Sigma_1 + O(n^{3/5})) + m|\mathcal{A}(n, m)|}{|\mathcal{G}(n, m)|(\Sigma_1 - O(n^{3/5}))}.$$

So,

$$\frac{|\mathcal{P}_{m+1}|}{|\mathcal{G}(n, m+1)|} \leq (1 + O(n^{-2/5})) \frac{|\mathcal{P}_m|}{|\mathcal{G}(n, m)|}.$$

So

$$\frac{|\mathcal{P}_m|}{\mathcal{G}(n, m)} \leq (1 + O(n^{-2/5}))^{m-n} \frac{|\mathcal{P}_n|}{\mathcal{G}(n, n)}. \quad (87)$$

We use (87) in the following way: Let  $\mathcal{P}(k)$  be the property that  $G$  contains a witness  $A, B$  with  $A \subseteq L, |A| = k$ . If  $(G_1, G_2) \in X$  and  $G_2 \in \mathcal{P}(k)$  then  $G_1 \in \mathcal{P}(k)$ . Applying (87) and (79) we see

that **whp** there is no witness with  $n^{3/4} \leq |A| \leq n - n^{3/4}$ . So we only have to deal with small and very large  $k$ .

We go back to (46). We see that  $f(\zeta_1) < f(z)$  implies that the term  $\frac{z^d}{f(z)^{\frac{\beta-a}{1-a}}} \frac{f(\zeta_1)^{\frac{\beta-a}{1-a}}}{\zeta_1^{d-x}} \left(\frac{e^{-a}}{x}\right)^x$  is maximised over  $\beta \geq 1$  when  $\beta = 1$ . Next let  $H(\beta) = \beta \log h(a/\beta)$  then  $H'(\beta) = \log(1 - a/\beta)$  and  $H''(\beta) = \frac{a}{\beta(\beta-a)}$ . Thus  $h(a/\beta)^\beta$  is log-convex in  $\beta$  and so

$$h(a/\beta)^\beta \geq \exp \{H(1) + H'(1)(\beta - 1)\} = h(a)(1 - a)^{\beta-1}. \quad (88)$$

Going back to (51) we see that now we have

$$\pi_L(k, D) \leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{\rho(a)}{(1-a)^{\beta-1}}\right)^n. \quad (89)$$

By taking  $\varepsilon_L(\beta)$  in place of  $\varepsilon_L(1)$  we can replace (54) by  $\rho(a) \leq e^{-\beta a}$  and plugging this into (89) we see that

$$\sum_{k=1}^{n^{3/4}} \sum_{D=dk}^{k \log n} \pi_L(k, D) \leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{e^{-\beta a}}{(1-a)^{\beta-1}}\right)^n = o(1). \quad (90)$$

To deal with  $k \geq n - n^{3/4}$  we treat this as  $k \leq n^{3/4}$  in Section 7.2.2. We can do this because if  $A' = R \setminus B$  and  $B' = L \setminus A$  then  $N_\Gamma(A') \subseteq B'$  and  $|A'| = m - k + 1 > |B'| = n - k$ .

We now simplify our estimate of  $\pi_R(k, D)$  by removing some terms involving  $\beta$  from (67).

$$\begin{aligned} \pi_R(k, D) &\leq \mathbb{P}(\exists A \subseteq R, B \subseteq L : |A| = k, |B| = k - 1, N_\Gamma(A) \subseteq B, d_B(b) \geq 2, b \in B) \leq \\ &O(n^{1/2}) \binom{m}{k} \binom{n}{k-1} \sum_{\substack{2 \leq d_a, a \in [m] \\ 2 \leq x_b \leq d, b \in [k-1] \\ \sum_{a \in [k]} d_a = \sum_{b \in [k-1]} x_b = D}} \prod_{a=1}^k \frac{z^{d_a}}{d_a! f(z)} \prod_{b=1}^{k-1} \binom{d}{x_b} D! \prod_{i=0}^{D-1} \frac{1}{dn - i} = \\ &O\left(\frac{k}{m^{1/2}}\right) \binom{n}{k} \binom{m}{k} \frac{z^D D! (dn - D)!}{f(z)^k (dn)!} \left( \sum_{\substack{2 \leq d_a, a \in [k] \\ \sum_a d_a = D}} \prod_{a=1}^k \frac{1}{d_a!} \right) \left( \sum_{\substack{2 \leq x_b \leq d, b \in [k-1] \\ \sum_b x_b = D}} \prod_{b=1}^{k-1} \binom{d}{x_b} \right) = \\ &O\left(\frac{k}{m^{1/2}}\right) \binom{n}{k} \binom{m}{k} \frac{z^D}{f(z)^k} \frac{1}{\binom{dn}{D}} \left([u^D](e^u - 1 - u)^k\right) \left([u^D]((1+u)^d - (1+du))^k\right) \leq \\ &O\left(\frac{k}{m^{1/2}}\right) \binom{n}{k} \binom{m}{k} \frac{z^D}{f(z)^k} \frac{1}{\binom{dn}{D}} \frac{f(\zeta_1)^k}{\zeta_1^D} \binom{dk}{D} = \\ &O\left(\frac{k}{m^{1/2}}\right) \binom{n}{k} \binom{m}{k} \frac{\binom{dk}{D}}{\binom{dn}{D}} \left(\frac{f(\zeta_1)}{f(z)}\right)^k \left(\frac{z}{\zeta_1}\right)^D \\ &= O\left(\frac{k}{m^{1/2}}\right) \left(\frac{h(\theta a/d)^d}{h(a)h(a/\beta)^\beta h(\theta/d)^{ad}} \left(\frac{f(\zeta_1)}{\zeta_1^\theta} \frac{z^\theta}{f(z)}\right)^a\right)^n. \end{aligned} \quad (91)$$

Now

$$\begin{aligned} \frac{h(\theta a/d)^d}{h(a)h(a/b)^b h(\theta/d)^{ad}} &= \frac{\left(\frac{\theta a}{d}\right)^{\theta a} \left(1 - \frac{\theta a}{d}\right)^{d-\theta a}}{a^a (1-a)^{1-a} \left(\frac{a}{\beta}\right)^a \left(1 - \frac{a}{\beta}\right)^{\beta-a} \left(\frac{\theta}{d}\right)^{\theta a} \left(1 - \frac{\theta}{d}\right)^{da-\theta a}} = \\ &= a^{(\theta-2)a} \exp \left\{ -(d-\theta a) \sum_{k=1}^{\infty} \frac{\theta^k a^k}{k d^k} + a - \sum_{k=2}^{\infty} \frac{a^k}{k(k-1)} + \frac{a}{\beta} - \sum_{k=2}^{\infty} \frac{a^k}{\beta^{k-1} k(k-1)} + (da - \theta a) \sum_{k=1}^{\infty} \frac{\theta^k}{k d^k} \right\} \\ &= a^{(\theta-2)a} \exp \left\{ a \left( 1 + \frac{1}{\beta} - (d-\theta) \log(\theta/d) - \theta \right) + O(a^2) \right\} \end{aligned} \quad (92)$$

So from (91) we can write

$$\pi_R(k, D) \leq O \left( \frac{k}{m^{1/2}} \right) \left( \left( \frac{az}{\zeta_1} \right)^{\theta-2} \exp \left\{ 1 + \frac{1}{\beta} - (d-\theta) \log(1-\theta/d) - \theta + O(a) \right\} \frac{f(\zeta_1)}{\zeta_1^2} \frac{z^2}{f(z)} \right)^{an}. \quad (93)$$

Now we claim that

$$\zeta_1^{\theta-2} \geq \frac{1}{2} \text{ and that } f(x)x^{-2} \text{ is monotone increasing in } x. \quad (94)$$

First notice that  $f(x)x^{-2} = \sum_{i=2}^{\infty} \frac{x^{i-2}}{i!}$  which is clearly monotone increasing. Second note that  $\zeta_1 = \zeta(\theta)$  and since  $\frac{d\zeta(x)}{dx} > 0$  we have

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{d\zeta(x)}{dx} &= \lim_{x \rightarrow \infty} \frac{f(\zeta(x))^2}{(e^{\zeta(x)} - 1)^2 - \zeta(x)^2 e^{\zeta(x)}} \\ &= \lim_{\zeta \rightarrow \infty} \frac{f(\zeta)^2}{(e^{\zeta} - 1)^2 - \zeta^2 e^{\zeta}} = 1 \end{aligned}$$

and since  $\zeta(x)$  is concave we have  $\frac{d\zeta(x)}{dx} \geq 1$ . This, along with  $\lim_{x \rightarrow 2^-} \zeta(x) = 0$ , implies that  $\zeta(x) \geq x - 2$ . We can then lower bound

$$\zeta_1^{\theta-2} = \zeta(\theta)^{\theta-2} \geq (\theta - 2)^{\theta-2} \geq e^{-e^{-1}} \geq 0.69$$

Using this we see from (93) that if

$$\theta \geq \theta_0 = 2 + \frac{4}{\log(1/az)}$$

then

$$\pi_R(k, D) \leq O \left( \frac{k}{m^{1/2}} \right) e^{-k}.$$

In which case we have

$$\sum_{k=1}^{n^{3/4}} \sum_{\theta \geq \theta_0} \pi_R(k, D) \leq O \left( \frac{k}{m^{1/2}} \right) e^{-k} = o(1). \quad (95)$$

When  $\theta < \theta_0$  we have  $\theta = 2 + o(1)$ ,  $f(\zeta_1)/\zeta_1^2 = 1/2 + o(1)$ . Therefore

$$\pi_R(k, D) \leq O \left( \frac{k}{m^{1/2}} \right) \left( \frac{z^2 e^{2-(d-2) \log(1-2/d) - \theta + o(1)}}{2f(z)} \right)^k.$$

Now for  $d \geq 4$  we have

$$\frac{z^2 e^{2-(d-2)\log(1-2/d)-\theta+o(1)}}{2f(z)} \leq \frac{3}{5} \quad (96)$$

and so

$$\sum_{k=1}^{n^{3/4}} \sum_{\theta \leq \theta_0} \pi_R(k, D) \leq O\left(\frac{k}{m^{1/2}}\right) \left(\frac{3}{5}\right)^k = o(1). \quad (97)$$

It remains to consider  $m$  close to  $dn/2$  i.e. where  $\xi$  defined at the beginning of this section is close  $1/2$ . Observe first that the number of edges incident with vertices of degree greater than two is at most  $3dn(1-2\xi)$ . If there are  $d_i$  vertices of degree  $i = 2, \geq 3$  then  $d_2 + d_3 = m = \xi dn$  and  $2d_2 + 3d_3 \leq dn$  which implies that  $d_2 \geq dn(3\xi - 1)$ . So the number of edges incident with vertices of degree greater than two is at most  $dn - 2dn(3\xi - 1)$ .

Now consider a witness  $A, B$  where  $|A| = \gamma n$ . We must have  $\gamma dn \leq 2\gamma n + 3dn(1-2\xi)$  which implies that  $\gamma \leq \frac{3d(1-2\xi)}{d-2}$  which can be made arbitrarily small. Now the estimate in (89) will suffice up to  $k \leq \varepsilon_L n$  and so we only need to make  $\xi$  close enough to  $1/2$  so that  $\gamma < \varepsilon_L$  (which depends only on  $d$  and not  $\gamma$ ).

#### 7.4 The case $m < n$

Now consider a set of pairs  $Y \subseteq \mathcal{G}(n, m) \times \mathcal{G}(n+1, m)$ . We place  $(G_1, G_2)$  into  $Y$  if  $G_2$  is obtained from  $G_1$  in the following manner: Choose  $0 \leq k \leq n$ . Replace edges  $(\ell, y)$  by  $(\ell+1, y)$  for all  $\ell > k$  and all  $y$ . Add vertex  $k+1$  and  $d$  edges  $(k+1, y_j), j = 1, 2, \dots, d$ .

Note that if  $(G_1, G_2) \in Y$  and  $G_1$  has a matching of  $R$  into  $L$  then so does  $G_2$ .

For  $G \in \mathcal{G}(n, m)$  let now

$$\pi_1(G) = |\{G_2 : (G, G_2) \in Y\}|$$

and for  $G \in \mathcal{G}(n+1, m)$  let

$$\pi_2(G) = |\{G_1 : (G_1, G) \in Y\}|.$$

Let

$$\Sigma_2 = (n+1) \left(1 - \frac{z^2}{2f(z)}\right)^d$$

and for  $G \in \mathcal{G}(n+1, m)$  let

$$L_3(G) = |\{v \in L : \text{all neighbours of } v \text{ have degree at least } 3\}|.$$

Let

$$\mathcal{C}(n+1, m) = \{G \in \mathcal{G}(n+1, m) : |L_3(G) - \Sigma_2| \leq n^\gamma\}.$$

Let

We note that

- $G \in \mathcal{G}(n, m)$  implies that  $\pi_1(G) = (n+1) \binom{m}{d}$ .
- $G \notin \mathcal{C}(n, m+1)$  implies that  $|\pi_2(G) - \Sigma_2| \leq n^\gamma$ .
- $\pi_2(G) \leq n+1$  for all  $G \in \mathcal{G}(n+1, m)$ .

We then note that

$$\frac{|Y|}{|\mathcal{G}(n, m)|} = (n+1) \binom{m}{d}.$$

$$\Sigma_2 - n^\gamma \leq \frac{|Y|}{|\mathcal{G}(n+1, m)|} \leq \Sigma_2 + n^\gamma + (n+1)e^{-\Omega(\log^2 n)}.$$

Now let  $\mathcal{P}$  be a property such that if  $(G_1, G_2) \in Y$  and  $G_2 \in \mathcal{P}$  then  $G_1 \in \mathcal{P}$ . Let  $(G_1, G_2)$  be chosen uniformly from  $Y$  and let  $\mathbf{P}_Y$  denote probabilities computed w.r.t. this choice. Let  $\mathcal{P}_n = \mathcal{P} \cap \mathcal{G}(n, m)$  and  $\mathcal{P}_{n+1} = \mathcal{P} \cap \mathcal{G}(n+1, m)$ . Then

$$\mathbf{P}_Y(G_2 \in \mathcal{P}) \leq \mathbf{P}_Y(G_1 \in \mathcal{P}) = \frac{|\mathcal{P}_n|(n+1) \binom{m}{d}}{|Y|}$$

and

$$\mathbf{P}_Y(G_2 \in \mathcal{P}) \geq \frac{(|\mathcal{P}_{n+1}| - |\mathcal{C}(n+1, m)|)(\Sigma_2 - n^\gamma)}{|Y|}$$

Arguing as in Section 7.3 we see that

$$\frac{|\mathcal{P}_n|}{|\mathcal{G}(n, m)|} \leq (1 + O(n^{-2/5}))^{n-m} \frac{|\mathcal{P}_m|}{|\mathcal{G}(m, m)|}. \quad (98)$$

Let  $\mathcal{Q}(k)$  be the property that  $G$  contains a witness  $A, B$  with  $A \subseteq R, |A| = k$ . If  $(G_1, G_2) \in Y$  and  $G_2 \in \mathcal{Q}(k)$  then  $G_1 \in \mathcal{Q}(k)$ . Applying (98) and (80) we see that **whp** there is no witness with  $n^{3/4} \leq |A| \leq n - n^{3/4}$ . So we only have to deal with small and very large  $k$ .

We first go back to (91) and deal with  $\pi_R(k, D)$  for  $k \leq n^{3/4}$  as we did in Section 7.3. For  $k \geq n - n^{3/4}$  we deal with  $\pi_L(k, D)$  for  $k \leq n^{3/4}$ . We will go back to (46) and write

$$\pi_L(k, D) = O\left(\frac{k}{n^{1/2}}\right) \left(\frac{h(a)^{d-1}}{h(a/\beta)^\beta}\right)^n \left(\frac{z^d f(\zeta_1)}{f(z) \zeta_1^{d-x}} \left(\frac{e^{-\frac{a}{1-a}}}{x}\right)^x \left(\frac{f(z)}{f(\zeta_1)}\right)^{\frac{1-\beta}{1-a}}\right)^{n-k}$$

Now  $x = \frac{d(m-n)+(D-dk)+1}{m-k+1} \geq 0$  implies that

$$1 - \beta \leq \frac{D - dk + 1}{n} = O\left(\frac{k \log n}{n}\right) \text{ and that } x = O\left(\frac{k \log n}{n}\right).$$

Also,  $\zeta_1 = \zeta(d-x)$  implies that  $f(\zeta_1) = f(z)(1 - O(x))$ . Therefore,

$$\left(\frac{f(z)}{f(\zeta_1)}\right)^{\frac{1-\beta}{1-a}} = e^{O(a^2 \log^2 n)}.$$

Arguing as for (89) we get

$$\pi_L(k, D) \leq O\left(\frac{k}{n^{1/2}}\right) \left(\frac{\rho(a)e^{O(a^2 \log^2 n)}}{(1-a)^{\beta-1}}\right)^n.$$

Taking  $\rho(a) \leq e^{-a}$  as in (56) and noting that  $b \leq 1$  here we get Thus

$$\sum_{k=1}^{n^{3/4}} \sum_{D=dk}^{k \log n} \pi_L(k, D) \leq \sum_{k=1}^{n^{3/4}} \sum_{D=dk}^{k \log n} O\left(\frac{k}{n^{1/2}}\right) \left(e^{-a+O(a^2 \log^2 n)}\right)^n = o(1). \quad (99)$$

## References

- [1] N. Alon and J. Spencer, *The Probabilistic Method*, Wiley-Interscience, 2008.
- [2] J. Aronson, A. Frieze, B.G. Pittel. Maximum matchings in sparse random graphs: Karp-Sipser re-visited. *Random Structures and Algorithms*, 12(2):111-178, 1998.
- [3] Y. Azar, A. Broder, A. Karlin, and E. Upfal. Balanced Allocations. *SIAM Journal on Computing*, 29(1):180-200, 1999.
- [4] T. Bohman, The Triangle-Free Process , *Advances in Mathematics*, to appear.
- [5] B. Bollobás, A probabilistic proof of an asymptotic formula for the number of labelled regular graphs, *European Journal of Combinatorics* 1 (1980) 311-316.
- [6] A. Broder and A. Karlin. Multilevel Adaptive Hashing. In *Proceedings of the 1st ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pp. 43-53, 1990.
- [7] A. Broder and M. Mitzenmacher. Using Multiple Hash Functions to Improve IP Lookups. *Proceedings of the 20th IEEE International Conference on Computer Communications (INFOCOM)*, pp. 1454-1463, 2001.
- [8] L. Devroye and P. Morin. Cuckoo Hashing: Further Analysis. *Information Processing Letters*, 86(4):215-219, 2003.
- [9] O. Dubois and J. Mandler, The 3-XORSAT Threshold, *Proceedings of the 43rd IEEE Symposium on Foundations of Computing (2002)* 779-788.
- [10] N. Fountoulakis and K. Panagiotou, Sharp Thresholds for Cuckoo Hashing, arXiv:0910.5147v1 [cs.DS].
- [11] M. Mitzenmacher, Private Communication.
- [12] M. Dietzfelbinder and C. Weidling. Balanced Allocation and Dictionaries with Tightly Packed Constant Size Bins. *Theoretical Computer Science*, 380(1-2):47-68, 2007.
- [13] D. Fotakis, R. Pagh, P. Sanders, and P. Spirakis. Space Efficient Hash Tables With Worst Case Constant Access Time. *Theory of Computing Systems*, 38(2):229-248, 2005.
- [14] R.M. Karp and M. Sipser, Maximum Matchings in Sparse Random Graphs, *Proceedings of the 22nd Annual IEEE Symposium on Foundations of Computer Science (1981)* 364-375.
- [15] A. Kirsch and M. Mitzenmacher. Using a Queue to De-amortize Cuckoo Hashing in Hardware. In *Proceedings of the Forty-Fifth Annual Allerton Conference on Communication, Control, and Computing*, 2007.
- [16] A. Kirsch, M. Mitzenmacher, and U. Wieder. More Robust Hashing: Cuckoo Hashing with a Stash. In *Proceedings of the 16th Annual European Symposium on Algorithms*, pp. 611-622, 2008.
- [17] A. Kirsch and M. Mitzenmacher. The Power of One Move: Hashing Schemes for Hardware. In *Proceedings of the 27th IEEE International Conference on Computer Communications (INFOCOM)*, pp. 565-573, 2008.

- [18] R. Kutzelnigg. Bipartite Random Graphs and Cuckoo Hashing. In *Proceedings of the Fourth Colloquium on Mathematics and Computer Science*, 2006.
- [19] M. Mitzenmacher and S. Vadhan. Why Simple Hash Functions Work: Exploiting the Entropy in a Data Stream. In *Proceedings of the Nineteenth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pp. 746-755, 2008.
- [20] R. Pagh and F. Rodler. Cuckoo Hashing. *Journal of Algorithms*, 51(2):122-144, 2004.
- [21] B. Vöcking. How Asymmetry Helps Load Balancing. *Journal of the ACM*, 50(4):568-589, 2003.
- [22] N.C. Wormald, The differential equation method for random graph processes and greedy algorithms, in *Lectures on Approximation and Randomized Algorithms* (M. Karonski and H.J. Proemel, eds), PWN, Warsaw, (1999) 73-155.