Direction of negative curvature for regularized SQP

Phillip Gill, Vyacheslav Kungurtsev, Daniel Robinson

May 22, 2018

1. Introduction

This note discusses the computation and use of a direction of negative curvature in the regularized sequential quadratic programming primal-dual augmented Lagrangian method (pdSQP) of Gill and Robinson [7], [8] for the purpose of ensuring convergence towards second-order optimal points. Section 2 discusses how to compute a direction of negative curvature using appropriate matrix factorizations. Section 3 discusses the specific relevant changes to the algorithm. Section 4 discusses the changes in the convergence results established by Gill and Robinson [8], showing that the desired convergence results continue to hold. Section 5 discusses global convergence to points satisfying the second-order necessary optimality conditions.

2. Direction of negative curvature

2.1. The active-set estimate

An index set \mathcal{W}_k is maintained that consists of the variable indices that estimate which components of x on their bounds. This set determines the the space in which to calculate the directions of negative curvature. The tolerance for an index to be in \mathcal{W}_k must converge to zero. A test such as $i \in \mathcal{W}_k$ if $[x_k]_i \leq \min\{\mu_k, \epsilon_a\}$, would be appropriate for the purpose of forming a \mathcal{W}_k for convexification, initializing the QP, and obtaining a direction of negative curvature. Otherwise, it would be necessary to use three different factorizations.

2.2. Calculating the direction

Recall that in pdSQP, the QP must use a Lagrangian Hessian \tilde{H} such that $\tilde{H} + \frac{1}{\mu}J^T J$ is positive definite. The process for forming the requisite \tilde{H} , as well as calculating a direction of negative curvature begins with the inertia-controlling factorization of the KKT matrix (see Forsgren [4]). Consider the KKT matrix,

$$\begin{pmatrix} H_F & J_F^T \\ J_F & -\mu I_{|F|} \end{pmatrix}, \tag{2.1}$$

with F the set of estimated free variables (those not in \mathcal{W}_k), and $I_{|F|}$ the identity matrix with |F| rows and columns.

The algorithm begins an LBL^T factorization of the KKT matrix, where L is lower triangular and B is a symmetric diagonal with 1×1 and 2×2 diagonal blocks. Standard pivoting strategies are described in the literature (see Bunch and Parlett [2], Fletcher [3], and Bunch and Kaufman [1]). Let the lower-right block be defined as $D = -\mu I_{|F|}$.

At step k of the factorization, let the partially factorized matrix have the following structure:

$$\begin{pmatrix} L_1 & 0 \\ L_2 & I \end{pmatrix} \begin{pmatrix} B & 0 \\ 0 & A \end{pmatrix} \begin{pmatrix} L_1^T & L_2^T \\ 0 & I \end{pmatrix}$$

with L_1 being lower triangular, I the identity of appropriate size, and A the matrix remaining to be factorized. Let A be partitioned as $A = \begin{pmatrix} a & b^T \\ b & C \end{pmatrix}$. If the top left element is chosen as a 1×1 pivot, at the next step,

$$\begin{pmatrix} L_1 & 0 & 0 \\ L_3 & 1 & 0 \\ L_4 & a^{-1}b & I \end{pmatrix} \begin{pmatrix} B & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & C - ba^{-1}b^T \end{pmatrix} \begin{pmatrix} L_1^T & L_3^T & L_4^T \\ 0 & 1 & a^{-1}b^T \\ 0 & 0 & I \end{pmatrix}.$$

Let $S = C - ba^{-1}b$ be the *Schur complement* of the factorization. The matrix S is factorized at the next step.

For inertia control, this factorization has two stages. In the first stage, we restrict the factorization to allow only for pivots of type H^+ , D^- or HD. This means that an element (i, j) of H is selected such that $H_{ij} > 0$, a diagonal element of D is selected, or (i_1, i_2, j_1, j_2) is selected such that (i_1, j_1) is an element of H, (i_2, j_2) is an element of D and $S_k[(i_1, i_2), (j_1, j_2)]$ has mixed eigenvalues. This procedure is continued until there are no such remaining pivots.

The KKT matrix can be partitioned as

$$\begin{pmatrix} H_{11} & H_{12} & J_1^T \\ H_{21} & H_{22} & J_2^T \\ J_1 & J_2 & -\mu I \end{pmatrix},$$

where, all of the pivots have come from the rows and columns of H_{11} , J_1 , and $-\mu I$. At the end of the first stage, the factorization can be written as:

$$\begin{pmatrix} L_1 & 0 \\ L_2 & I \end{pmatrix} \begin{pmatrix} B & 0 \\ 0 & H_{22} - K_{21} K_{11}^{-1} K_{12} \end{pmatrix} \begin{pmatrix} L_1^T & L_2^2 \\ 0 & I \end{pmatrix}.$$
 (2.2)

Let $S = H_{22} - K_{21}K_{11}^{-1}K_{12}$. Proposition 3 of Forsgren [4] shows that if δI is added to H_{22} such that $\delta > ||S||$ then K_F has the correct inertia. In practice this δ is excessively large for the purpose of constructing the appropriate matrix with the required eigenvalues, but this result does indicate that such a constant exists.

Instead of proceeding to the second phase of this factorization, the procedure of Lemma 2.4 in Forsgren et al. [6] is applied to S to compute \hat{u} , a direction of negative curvature for S. The procedure to calculate this \hat{u} is as follows:

Let $\rho = \max_{i,j} |S_{ij}|$ with $|S_{qr}| = \rho$. Define \hat{u} as the solution to:

$$\begin{pmatrix} L_1 & 0\\ L_2 & I \end{pmatrix} \widehat{u} = \sqrt{\rho}h, \qquad (2.3)$$

where

$$h = \begin{cases} e_q & \text{if } q = r, \\ \frac{1}{\sqrt{2}}(e_q - \operatorname{sgn}(b_{qr})e_r) & \text{otherwise.} \end{cases}$$

This \hat{u} satisfies $\hat{u}^T S \hat{u} \leq \gamma \lambda_{\min}(S) ||\hat{u}||^2$, with γ independent of S.

The following bounds are important for the subsequent second-order convergence theory.

Lemma 2.1. Let \hat{u} be defined as in (2.3), S be the Schur complement of the partially factorized matrix (2.2), J_F and H_F defined as in (2.1), and Z a matrix consisting of columns for the basis of the null-space of J_F , then

$$\frac{\hat{u}^T S \hat{u}}{\gamma ||\hat{u}||^2} \le \lambda_{\min}(S) \le \lambda_{\min}(H_F + \frac{1}{\mu} J_F^T J_F) \le \lambda_{\min}(Z^T H_F Z).$$

Proof. Lemma 2.4 in Forsgren et al. [6] directly implies that $\hat{u}^T S \hat{u} / \gamma || \hat{u} ||^2 \leq \lambda_{\min}(S)$.

The proof that $\lambda_{\min}(S) \leq \lambda_{\min}(H_F + \frac{1}{\mu}J_F^T J_F)$ is given in the proof of Theorem 4.5 in Forsgren and Gill [5]. For the final inequality, let w = Zv, with $Z^T H_F Zv = \lambda_{\min}(Z^T H_F Z)v$ and ||v|| = 1. Then

$$\lambda_{\min}(H_F + \frac{1}{\mu}J_F^T J_F) \le \frac{w^T (H_F + \frac{1}{\mu}J_F^T J_F)w}{w^T w} = w^T H_F w = v^T Z^T H_F Z v = \lambda_{\min}(Z^T H_F Z).$$

3. Implementing Directions of Negative Curvature

3.1. Step of negative curvature

Several changes must be made to the algorithm of Gill and Robinson [8]. In order to minimize the number of factorizations, the computation of the direction of negative curvature should be followed by a test of second-order optimality. In addition, it is necessary that the direction of negative curvature is bounded, and a feasible direction with respect to both the linearized equalities and the bound constraints. Finally, the line-search must be extended to allow for this additional step of negative curvature.

In the description below, the subscript k denoting the step number in the sequence of iterations is suppressed.

The following procedure satisfies these requirements.

- 1. The first step computes the direction of negative curvature for the free KKTmatrix as described in Section 2, denoted as \hat{u}_F , then defines \hat{u} to be $[\hat{u}]_F = \hat{u}_F$ and $[\hat{u}]_A = 0$. If no such direction of negative curvature exists, then \hat{u} is set to zero.
- 2. The second step uses \hat{u} in a test of second-order optimality. This is described in Section 3.2.

3. The corresponding change in the multipliers corresponding to the definition for \hat{u} is defined as $\hat{w} = -\frac{1}{\mu}J\hat{u}$. This ensures that the linearized equality constraints are satisfied, i.e.,

$$0 = Jp + c + \mu q = J(p + \hat{u}) + c + \mu (q - \frac{1}{\mu}J\hat{u}).$$

The final resulting (u, w) is shown below in Section 3.3 to be a direction of negative curvature for $\nabla^2 \mathcal{M}$.

- 4. Since both (\hat{u}, \hat{w}) and $-(\hat{u}, \hat{w})$ are directions of negative curvature, the sign is chosen so that the step is a descent direction for $\nabla \mathcal{M}$, i.e. $\nabla \mathcal{M}^T \begin{pmatrix} \hat{u} \\ \hat{w} \end{pmatrix} \leq 0$.
- 5. Compute $\Delta v = (p, q)$, the solution of the convex QP.
- 6. The direction of negative curvature is scaled so that it is both bounded by $\max(u_{\max}, 2||p||)$ and also, in conjunction with the QP step, satisfies the bound constraints $x \ge 0$.

Specifically, u and w are set as $u = \beta \hat{u}$ and $w = \beta \hat{w}$, where

$$\beta = \left\{ \max \hat{\beta} \mid x + p + \hat{\beta} \widehat{u} \ge 0, \, ||\hat{\beta} \widehat{u}|| \le \max(u_{\max}, 2||p||) \right\}.$$

Note that this implies that if $[x + p]_i = 0$ and $[u]_i < 0$, then u is set to zero.

3.2. Optimality measures

Recall that in Gill and Robinson [8], with

$$\phi_S(v) = \eta(x) + 10^{-5}\omega(v)$$
 and $\phi_L(v) = 10^{-5}\eta(x) + \omega(v)$,

where

$$\eta(x) = \|c(x)\|$$
 and $\omega(x, y) = \|\min\left(x, g(x) - J(x)^T y\right)\|,$

an iterate is an S-iterate if $\phi_S(v) \leq \frac{1}{2}\phi_S^{max}$ and an L-iterate if $\phi_L(v) \leq \frac{1}{2}\phi_L^{max}$. Otherwise, an iterate is an M-iterate if

$$||\nabla_y \mathcal{M}(v_{k+1}; y_k^E, \mu_k^R)|| \le \tau_k \text{ and } ||\min(x_{k+1}, \nabla_x \mathcal{M}^{\nu}(v_{k+1}; y_k^E, \mu_k^R))|| \le \tau_k.$$

If none of these conditions hold, then an iterate v_k is an F-iterate.

In order to force convergence to a second-order optimal point, it is necessary to change the function $\omega(x, y)$ that appears in ϕ_S and ϕ_L , as well as the test for an iteration being an M-iterate.

Ideally, the minimum eigenvalue of H in the null-space for J_F should be found, as well as the minimum eigenvalue of $\nabla^2_{xx}\mathcal{M}$. However, this would require extensive computation. Instead, these quantities are estimated based on the value of the negative curvature. Recall that

$$\frac{\widehat{u}^T (H + \frac{1}{\mu} J^T J) \widehat{u}}{\gamma ||\widehat{u}||^2} \le \lambda_{\min} (H + \frac{1}{\mu} J^T J),$$

where we suppress the suffix F. Since γ is bounded from below and above, if $\hat{u}^T (H + \frac{1}{\mu} J^T J) \hat{u} / ||\hat{u}||^2 \to 0$, the estimate for \hat{u} implies $\lim \lambda_{\min} (H + \frac{1}{\mu} J^T J) \ge 0$. Hence, the test for M-iterate optimality is changed to:

$$\begin{aligned} ||\nabla_{y}\mathcal{M}(v_{k+1}; y_{k}^{E}, \mu_{k}^{R})|| &\leq \tau_{k} \\ \text{and } ||\min(x_{k+1}, \nabla_{x}\mathcal{M}^{\nu}(v_{k+1}; y_{k}^{E}, \mu_{k}^{R}))|| &\leq \tau_{k} \\ \text{and } \frac{\widehat{u}_{k+1}^{T}(H + \frac{1}{\mu}J^{T}J)\widehat{u}_{k+1}}{||\widehat{u}_{k+1}||^{2}} &\geq \tau_{k}. \end{aligned}$$

Similarly, for the filter functions,

$$\phi_S(v) = \eta(x) + 10^{-5}\omega(v)$$
 and $\phi_L(v) = 10^{-5}\eta(x) + \omega(v)$

the optimality tests become

$$\eta(x) = ||c(x)|| \text{ and } \omega(x,y) = \min(||\min(x,g(x) - J(x)^T y)|, -\frac{\widehat{u}_{k+1}^T (H + \frac{1}{\mu} J^T J) \widehat{u}_{k+1}}{||\widehat{u}_{k+1}||^2}).$$

3.3. Merit function

The line-search must also be changed to include the direction of negative curvature. First, it will be shown that the full primal-dual step is a step of negative curvature for the merit function Hessian.

Lemma 3.1. The vector (u, w) defined as in 3.1 is a direction of negative curvature for $\nabla^2 \mathcal{M}$.

Proof. Consider the calculation of $\begin{pmatrix} u \\ w \end{pmatrix}^T \nabla^2 \mathcal{M} \begin{pmatrix} u \\ w \end{pmatrix}$.

$$\begin{pmatrix} u \\ w \end{pmatrix}^T \begin{pmatrix} H + \frac{1}{\mu}(1+\nu)J^T J & \nu J^T \\ \nu J & \nu \mu I \end{pmatrix} \begin{pmatrix} u \\ w \end{pmatrix}$$

=
$$\begin{pmatrix} u \\ w \end{pmatrix}^T \begin{pmatrix} Hu + \frac{1}{\mu}(1+\nu)J^T Ju + \nu J^T w \\ \nu Ju + \nu \mu w \end{pmatrix}$$

=
$$u^T Hu + \frac{1}{\mu}(1+\nu)u^T J^T Ju + 2\nu u^T J^T w + \nu \mu ||w||^2.$$

From the definition above, $u = \beta \hat{u}$ and $\hat{u}^T (H + \frac{1}{\mu} J^T J) \hat{u} \leq \gamma \lambda_{\min} (H + \frac{1}{\mu} J^T J) ||\hat{u}||^2$, so multiplying both sides by β^2 , the expression becomes $u^T (H + \frac{1}{\mu} J^T J) u \leq \gamma \lambda_{\min} (H + \frac{1}{\mu} J^T J) ||u||^2$. Let $\bar{\gamma} = \gamma \lambda_{\min} (H + \frac{1}{\mu} J^T J)$. Using $w \triangleq -\frac{1}{\mu} J u$, $u^T H u + \frac{1}{\mu} (1 + \nu) u^T J^T J u + 2\nu u^T J^T w + \nu \mu ||w||^2 \leq -\bar{\gamma} ||u||^2 - 2\frac{\nu}{\mu} u^T J^T J u + \frac{\nu}{\mu} ||Ju||^2 = -\bar{\gamma} ||u||^2 - \frac{\nu}{\mu} ||Ju||^2 \leq -\bar{\gamma} ||u||^2 - \nu \mu ||w||^2$.

I		

For the line-search, let $R_k \triangleq u_k^T \nabla^2 \mathcal{M}^{\nu}(v_k; y_k^E, \mu_k^R) u_k \leq 0$. Define $\alpha_k = 2^{-j}$ such that

$$\mathcal{M}^{\nu}(v_k + \alpha_k u_k + \alpha_k^2 \Delta v_k; y_k^E, \mu_k^F) \le \mathcal{M}^{\nu} + \alpha_k^2 \eta_S N_k + \alpha_k \eta_S R_k.$$
(3.1)

Letting $\bar{\alpha} \triangleq \min(\alpha_{\min}, \alpha_k)$ and $\hat{\mu} \triangleq \max(\frac{1}{2}\mu_k, \mu_{k+1}^R)$, the update for the penalty parameter becomes:

$$\mu_{k+1} = \begin{cases} \mu_k, & \mathcal{M}^{\nu}(v_{k+1}; y_k^E, \mu_k) \le \mathcal{M}^{\nu}(v_k; y_k^E, \mu_k) + \bar{\alpha}\eta_S R_k + \bar{\alpha}^2 \eta_S N_k \\ \widehat{\mu}, & \text{otherwise}, \end{cases}$$
(3.2)

4. Consistency with established convergence theory

In their first-order analysis, Gill and Robinson [8] make the following assumptions:

Assumption 4.1. Each $\overline{H}(x_k, y_k)$ is chosen so that the sequence $\{\overline{H}(x_k, y_k)\}_{k\geq 0}$ is bounded, with $\{\overline{H}(x_k, y_k) + (1/\mu_k^R)J(x_k)^TJ(x_k)\}_{k\geq 0}$ uniformly positive definite.

Assumption 4.2. The functions f and c are twice continuously differentiable.

Assumption 4.3. The sequence $\{x_k\}_{k\geq 0}$ is contained in a compact set.

Since $\nabla \mathcal{M}^{\nu}$ does not involve any term involving the objective or constraint Hessians, much of the first-order convergence theory holds. Incorporating the direction of negative curvature, Theorem 4.1 changes to:

Theorem 4.1. If there exists an integer \hat{k} such that $\mu_k^R \equiv \mu^R > 0$ and k is an \mathcal{F} -iterate for all $k \geq \hat{k}$, then the following hold:

- 1. $\{||\Delta v_k|| + ||u_k||\}_{k \ge \hat{k}}$ is bounded away from zero
- 2. There exists an ϵ such that for all $k \geq \hat{k}$, it holds that

$$\nabla \mathcal{M}^{\nu}(v_k; y_k^E, \mu_k^R)^T \Delta v_k) \leq -\epsilon \text{ or } u_k^T \nabla^2 \mathcal{M}^{\nu}(v_k; y_k^E, \mu_k^R) u_k \leq -\epsilon.$$

Proof. If all iterates $k \ge \hat{k}$ are \mathcal{F} -iterates, then,

$$\tau_k \equiv \tau > 0, \ \mu_k^R = \mu^R, \text{ and } y_k^E = y^E \text{ for all } k \ge \hat{k}$$

Proof of the first result: Assume the contrary, i.e., there exists a subsequence $S_1 \subset \left\{k \mid k \geq \hat{k}\right\}$ such that $\lim_{k \in S_1} \Delta v_k = 0$ and $\lim_{k \in S_1} u_k = 0$. The solution Δv_k to the QP subproblem satisfies

$$\begin{pmatrix} z_k \\ 0 \end{pmatrix} = H_M^{\nu}(v_K; \mu^R) \Delta v_k + \nabla \mathcal{M}^{\nu}(v_k; y^E, \mu^R) \text{ and } 0 = \min(x_k + p_k, z_k).$$

As H_M^{ν} is uniformly bounded, eventually for some $k \in S_1$ sufficiently large, Δv_k satisfies the first-order conditions of an M-iterate, i.e.,

$$||\nabla_y \mathcal{M}(v_{k+1}; y_k^E, \mu_k^R)|| \le \tau_k \text{ and } ||\min(x_{k+1}, \nabla_x \mathcal{M}^{\nu}(v_{k+1}; y_k^E, \mu_k^R))|| \le \tau_k.$$

In the construction of u_k , recall that ||u|| is the largest possible value, subject to an upper bound, that is feasible. This implies that if $\lim u_k \to 0$, then eventually, uis constrained by feasibility, or set to zero.

In the first case, i.e. the limiting upper bound constraint on u_k must be $x_k + p_k + u_k \ge 0$, eventually since $u_k \to 0$ and $p_k \to 0$, if *i* is a blocking bound for u_k , $x_i \le \min(\mu, \epsilon_x)$ and $i \in \mathcal{W}_k$, which implies that $[u]_i \equiv 0$. Hence, by construction and the fact that the set of possible indices is finite, u_k is eventually identically zero. This implies that the second-order conditions of an M-iterate are also satisfied trivially, i.e.,

$$\frac{\widehat{u}_{k+1}^{T}(H + \frac{1}{\mu}J^{T}J)\widehat{u}_{k+1}}{||\widehat{u}_{k+1}||^{2}} \ge \tau_{k};$$

and μ_k^R is decreased. This contradicts the assumption that μ_k^R is held fixed at $\mu_k^R \equiv \mu^R$ for all $k \geq \hat{k}$.

Proof of part (2): Assume, to the contrary, that there exists a subsequence S_2 of $\{k : k \ge \hat{k}\}$ such that

$$\lim_{k \in \mathcal{S}_2} \nabla \mathcal{M}^{\nu}(v_k; y^E, \mu^R)^T \Delta v_k = 0$$
(4.1)

and

$$\lim_{k \in \mathcal{S}_2} u_k^T \nabla^2 \mathcal{M}^{\nu}(v_k; y_k^E, \mu_k^R) u_k = 0.$$

Consider the matrix

$$L_k = \begin{pmatrix} I & 0\\ \frac{1}{\mu^R} J_k & I \end{pmatrix}$$

Since the $\Delta v = 0$ is feasible and Δv_k a solution for the convex problem, it follows that

$$-\nabla \mathcal{M}^{\nu}(v_k; y^E, \mu^R)^T \Delta v_k \ge \frac{1}{2} \Delta v_k^T H_M^{\nu}(v_k; \mu^R) \Delta v_k$$

$$= \frac{1}{2} \Delta v_k^T L_k^{-T} L_k^T H_M^{\nu}(v_k; \mu^R) L_k L_k^{-1} \Delta v_k$$

$$= \begin{pmatrix} p_k \\ q_k + \frac{1}{\mu^R} J_k p_k \end{pmatrix}^T \begin{pmatrix} \bar{H}_k + \frac{1}{\mu^R} J_k^T J_k & 0 \\ 0 & \nu \mu^R \end{pmatrix} \begin{pmatrix} p_k \\ q_k + \frac{1}{\mu^R} J_k p_k \end{pmatrix}$$

Since H_M^{ν} is bounded,

$$\Delta v_k^T L_k^{-T} L_k^T H_M^{\nu}(v_k; \mu^R) L_k L_k^{-1} \Delta v_k \ge \bar{\lambda}_{\min} \|p_k\|^2 + \nu \mu^R \|q_k + (1/\mu^R) J_k p_k\|^2,$$

for some $\bar{\lambda}_{\min} > 0$. Combining this with (4.1) it follows that

$$\lim_{k \in \mathcal{S}_2} p_k = \lim_{k \in \mathcal{S}_2} \left(q_k + \frac{1}{\mu^R} J_k p_k \right) = 0,$$

in which case $\lim_{k \in S_2} q_k = 0$. Hence $\Delta v_{k \in S_2} \to 0$.

Since $\lim_{k \in S_2} u^T_k \nabla^2 \mathcal{M}^{\nu}(x_k, y_k; y^E, \mu) u_k = 0$, there exists a \hat{k}_2 , such that for all $k \geq \hat{k}_2$, $u_k^T \nabla^2 \mathcal{M}^{\nu}(x_k, y_k; y^E, \mu) u_k / \gamma ||u_k||^2 > -\tau$ or $u_k \to 0$. The former, by the same argument as for part (1), together with $\Delta v_k \to 0$, implies that eventually k is an M-iterate. The latter, together with $\lim \Delta_k = 0$, contradicts the statement of part (1) of the theorem, so part (3) must hold.

The proofs of the first result of Theorem 4.1 and Theorem 4.2 of Gill and Robinson [8] do not change.

5. Global convergence to second-order optimal points

5.1. Filter Convergence

Definition 5.1. The Weak Constant Rank (WCR) condition holds at x if there is a neighborhood M(x) for which the rank of $\begin{pmatrix} J(z) \\ E_A^T \end{pmatrix}$ is constant for all $z \in M(x)$, where E_A is the columns of the identity corresponding to the indices of x active at x (as in $i \in A$ if $x_i = 0$).

Theorem 5.1. Assume there is a subsequence v_k of S- and L-iterates converging to v^* , with $v^* = (x^*, y^*)$ satisfying the first-order KKT conditions. Furthermore, assume that MFCQ and WCR hold at v^* . Then v^* satisfies the necessary secondorder necessary optimality conditions.

Proof. Let $d \in T(x^*) \equiv \{d \mid J(x^*)d = 0 \text{ and } E_{w^*}^T d = 0\}$ with ||d|| = 1. By Lemma 3.1 of Andreani et al. ([9]) there exists $\{d_k\}$ such that $d_k \in T(x_k)$ and $d_k \to d$, where

$$T(x_k) = \{ d \mid J(x_k)d = 0 \text{ and } E_{w^*}^T d = 0 \}$$

Without loss of generality, we may let $||d_k|| = 1$. Since $x_k \to x^*$, eventually $\mathcal{W}_k = \mathcal{A}^*$, where \mathcal{A}^* is the active set at x^* . Then, by the definition of the S- and Literates, and Lemma 2.1, $d_k^T (\nabla^2 f(x_k) + \sum y_k \nabla^2 c(x_k)) d_k > \lambda_{\min}(Z_k H_k Z_k) > -\xi_k$, where $0 < \xi_k \to 0$. Taking limits, it follows that $d^T (\nabla^2 f(x_k) + \sum y^* \nabla^2 c(x^*)) d \ge 0$.

References

- J. R. Bunch and L. Kaufman. A computational method for the indefinite quadratic programming problem. *Linear Algebra Appl.*, 34:341–370, 1980.
- [2] J. R. Bunch and B. N. Parlett. Direct methods for solving symmetric indefinite systems of linear equations. SIAM J. Numer. Anal., 8:639–655, 1971.
- [3] Roger Fletcher. Factorizing symmetric indefinite matrices. Linear Algebra Appl., 14:257–272, 1976.
- [4] Anders Forsgren. Inertia-controlling factorizations for optimization algorithms. Appl. Num. Math., 43:91–107, 2002.
- [5] Anders Forsgren and Philip E. Gill. Primal-dual interior methods for nonconvex nonlinear programming. SIAM J. Optim., 8:1132–1152, 1998.

- [6] Anders Forsgren, Philip E. Gill, and Walter Murray. Computing modified Newton directions using a partial Cholesky factorization. Report TRITA-MAT-1993-9, Division of Optimization and Systems Theory, Department of Mathematics, Royal Institute of Technology, 1993.
- [7] Philip E. Gill and Daniel P. Robinson. A primal-dual augmented Lagrangian. Computational Optimization and Applications, pages 1–25, 2010.
- [8] Philip E. Gill and Daniel P. Robinson. Regularized sequential quadratic programming methods. Numerical Analysis Report 11-02, Department of Mathematics, University of California, San Diego, La Jolla, CA, 2011.
- [9] R. Andreani J.M. Martinez and M.L. Schuverdt. On second-order optimality conditions for nonlinear programming. 2010.