

FROM STATE INTEGRALS TO q -SERIES

STAVROS GAROUFALIDIS AND RINAT KASHAEV

ABSTRACT. It is well-known to the experts that multi-dimensional state integrals of products of Faddeev's quantum dilogarithm which arise in Quantum Topology can be written as finite sums of products of basic hypergeometric series in $q = e^{2\pi i\tau}$ and $\tilde{q} = e^{-2\pi i/\tau}$. We illustrate this fact by giving a detailed proof for a family of one-dimensional integrals which includes state-integral invariants of 4_1 and 5_2 knots.

CONTENTS

1. Introduction	1
1.1. State-integrals and their q -series	2
1.2. q -difference equations	4
1.3. The case of the 4_1 knot	4
1.4. The case of the 5_2 knot	5
2. Proofs	6
2.1. A residue computation	6
2.2. The Taylor series of $\phi_m(x)$ and $\tilde{\phi}_n(x)$	8
2.3. The connection with the differential operators δ_l and $\tilde{\delta}_l$	8
2.4. Proof of Theorem 1.1	10
2.5. Some auxiliary power series	10
2.6. Proof of Proposition 2.2	12
2.7. Proof of Lemma 1.5	13
3. An application: state-integrals of the 4_1 and 5_2 knots	14
3.1. Proof of Corollary 1.7	14
3.2. Proof of Corollary 1.8	14
Acknowledgments	15
References	15

1. INTRODUCTION

Date: April 3, 2013.

S.G. was supported in part by grant DMS-0805078 of the US National Science Foundation. R.K. was supported in part by the Swiss National Science Foundation.

2010 Mathematics Subject Classification: Primary 57N10. Secondary 57M25, 33F10, 39A13.

Key words and phrases: state-integrals, q -series, quantum dilogarithm, Euler triangular numbers, Nahm equation, gluing equations, 4_1 , 5_2 .

1.1. State-integrals and their q -series. Multi-dimensional state integrals of products of Faddeev's quantum dilogarithm appear in abundance in Quantum Topology, and were studied among others by Hikami [Hik01], Dimofte–Gukov–Lennels–Zagier [DGLZ09], Andersen–Kashaev [AK], and Kashaev–Luo–Vartanov [KLV12]. It is well-known to the experts that such state-integrals can be written as finite sums of products of pairs of q -series and \tilde{q} -series. The reason for this is a factorized structure of Faddeev's quantum dilogarithm, the structure of the set of its poles, and the specific form of exponential factors of the integrand of the state-integrals, while its derivation is based on an application of the residue theorem. Instead of formulating a general theorem for multi-dimensional integrals which obscures the principle, we will give a detailed proof for the case of a family of 1-dimensional integrals and illustrate it with some concrete examples taken from [AK, KLV12].

To state our results, recall that *Faddeev's quantum dilogarithm function* $\Phi_b(x)$ is given by [Fad95]

$$(1) \quad \Phi_b(x) = \frac{(e^{2\pi b(x+c_b)}; q)_\infty}{(e^{2\pi b^{-1}(x-c_b)}; \tilde{q})_\infty},$$

where

$$q = e^{2\pi i b^2}, \quad \tilde{q} = e^{-2\pi i b^{-2}}, \quad c_b = \frac{i}{2}(b + b^{-1}), \quad \Im(b^2) > 0.$$

Remarkably, this function admits an extension to all values of b with $b^2 \notin \mathbb{R}_{\leq 0}$. $\Phi_b(x)$ is a meromorphic function of x with

$$\text{poles: } c_b + i\mathbb{N}b + i\mathbb{N}b^{-1}, \quad \text{zeros: } -c_b - i\mathbb{N}b - i\mathbb{N}b^{-1}.$$

The functional equation

$$\Phi_b(x)\Phi_b(-x) = e^{\pi i x^2} \Phi_b(0)^2, \quad \Phi_b(0) = q^{\frac{1}{24}} \tilde{q}^{-\frac{1}{24}}$$

allows us to move $\Phi_b(x)$ from the denominator to the numerator of the integrand of a state-integral.

For natural numbers A, B with $B > A > 0$, we consider the absolutely convergent integral

$$\mathcal{I}_{A,B}(b) = \int_{\mathbb{R}+i\epsilon} \Phi_b(x)^B e^{-A\pi i x^2} dx$$

with small positive ϵ . The condition $B > A > 0$ ensures not only the convergence of $\mathcal{I}_{A,B}(b)$ for $\Im(b^2) > 0$, but also the convergence of the q -series and the \tilde{q} -series (for $|q|, |\tilde{q}| < 1$) that appear in Theorem 1.1 below.

To express the above state-integral in terms of series, consider the generating series

$$(2) \quad F_{A,B}(q, x) = \sum_{m=0}^{\infty} \frac{(-1)^{Am} q^{A \frac{m(m+1)}{2}}}{(q)_m^B} x^m, \quad \tilde{F}_{A,B}(q, x) = F_{B-A,B}(q, x).$$

Consider the operators δ and δ_k (for k a positive natural number) which act on the space of functions of x as follows

$$(3) \quad (\delta F)(x) = x \partial_x F(x), \quad (\delta_k F)(x) = \sum_{s=1}^{\infty} \frac{s^{k-1} q^s}{1 - q^s} F(q^s x).$$

Likewise, there are operators $\tilde{\delta}$ and $\tilde{\delta}_k$ which act on the space of functions of \tilde{x} and with q replaced by \tilde{q} . It is easy to see that any two of the operators $\delta, \delta_k, \tilde{\delta}, \tilde{\delta}_k$ commute and they freely generate over \mathbb{Q} a commutative ring $\mathcal{D} \otimes \tilde{\mathcal{D}}$, where

$$\mathcal{D} = \mathbb{Q}[\delta, \delta_1, \delta_2, \dots], \quad \tilde{\mathcal{D}} = \mathbb{Q}[\tilde{\delta}, \tilde{\delta}_1, \tilde{\delta}_2, \dots].$$

Let

$$\mathcal{D}_b = \mathcal{D}[(2\pi i)^{-1}, b^{\pm 1}, e_2, e_4, e_6, \dots], \quad \tilde{\mathcal{D}}_b = \tilde{\mathcal{D}}[(2\pi i)^{-1}, b^{\pm 1}, e_2, e_4, e_6, \dots],$$

where $e_l = e_l(\tilde{q}) = \tilde{\delta}_l(1) \in \mathbb{Z}[[\tilde{q}]]$. Consider the following *operator valued polynomial*:

$$(4) \quad P_{A,B} = \text{Res}_{w=0} \left(e^{\frac{1}{4\pi i} w^2 + Aw(b(\delta + \frac{1}{2}) + b^{-1}(\tilde{\delta} + \frac{1}{2}))} \right)^A \left(\frac{\phi(bw, \delta_\bullet) \tilde{\phi}(b^{-1}w, \tilde{\delta}_\bullet)}{b(1 - e^{b^{-1}w})} \right)^B \in \mathcal{D}_b \otimes \tilde{\mathcal{D}}_b,$$

where

$$(5a) \quad \phi(w, \delta_\bullet) = \exp \left(- \sum_{l=1}^{\infty} \frac{\delta_l}{l!} w^l \right)$$

$$(5b) \quad \tilde{\phi}(w, \tilde{\delta}_\bullet) = \exp(-\tilde{\delta}w) \exp \left(2 \sum_{l=\text{even}>0} e_l(\tilde{q}) \frac{w^l}{l!} \right) \exp \left(- \sum_{l=1}^{\infty} \frac{\tilde{\delta}_l}{l!} (-w)^l \right).$$

For a series $F(x, \tilde{x})$, we define:

$$(6) \quad \langle F(x, \tilde{x}) \rangle = F(1, 1).$$

Theorem 1.1. *We have:*

$$(7) \quad \mathcal{I}_{A,B}(b) = \left(\frac{\tilde{q}}{q} \right)^{\frac{B-3A}{24}} e^{\pi i \frac{B+2(A+1)}{4}} \left\langle P_{A,B} \left(F_{A,B}(q, x) \tilde{F}_{A,B}(\tilde{q}, \tilde{x}) \right) \right\rangle.$$

Corollary 1.2. Writing $P_{A,B} = \sum_k p_k P_k$ (a finite sum), for $p_k \in \mathcal{D}_b$ and $P_k \in \tilde{\mathcal{D}}_b$, it follows that

$$(8) \quad \mathcal{I}_{A,B}(b) = \left(\frac{\tilde{q}}{q} \right)^{\frac{B-3A}{24}} e^{\pi i \frac{B+2(A+1)}{4}} \sum_k g_k(q) G_k(\tilde{q})$$

where

$$(9) \quad g_k(q) = \langle p_k F_{A,B} \rangle, \quad G_k(\tilde{q}) = \langle P_k \tilde{F}_{A,B} \rangle.$$

Remark 1.3. The left hand side of Equation (8) has analytic continuation to the cut plane $\mathbb{C} \setminus \{b^2 \mid b^2 < 0\}$ whereas each of the series g_k and G_k is only well-defined in the upper-half plane $\{b^2 \mid \Im(b^2) > 0\}$.

Remark 1.4. $P_{A,B}$, as a polynomial in the variables e_2, e_4, \dots has degree $B - 1$, where the degree of e_l is l . $P_{A,B}$ as a Laurent polynomial in b has b -monomials of degrees in $\{-B + 1, -B + 3, \dots, B - 3, B - 1\}$.

1.2. q -difference equations. Next we describe a linear q -difference equation of $F_{A,B}(q, x)$. Consider the operators \hat{x} and \hat{E} which act on $f(x) \in \mathbb{Q}(q)[[x]]$ by:

$$(\hat{E}f)(x) = f(qx), \quad (\hat{x}f)(x) = xf(x).$$

Observe that $\hat{E}\hat{x} = q\hat{x}\hat{E}$.

Lemma 1.5. (a) We have:

$$(10) \quad F_{A,B}(q^{-1}, x) = \tilde{F}_{A,B}(q, x).$$

(b) $F_{A,B}$ satisfies the linear q -difference equation

$$(11) \quad \left((1 - \hat{E})^B - (-1)^A q^A x \hat{E}^A \right) F_{A,B}(q, x) = 0.$$

Corollary 1.6. (a) If we define $\omega(q, x) = F_{A,B}(q, qx)/F_{A,B}(q, x)$ and $\omega(q, x)_n = \prod_{j=1}^n \omega(q, q^j x)$, then ω satisfies the nonlinear equation

$$\sum_{j=0}^B (-1)^j \binom{B}{j} \omega(q, x)_j - (-1)^A q^A x \omega(q, x)_A = 0.$$

(b) F is an admissible power series in the sense of Kontsevich-Soibelman [KS11, Sec.6], the limit $\lim_{q \rightarrow 1} \omega(q, x) = \omega(x) \in \overline{\mathbb{Q}}[[x]]$ exists and satisfies the algebraic equation (also known as the Nahm equation or the gluing equation)

$$(12) \quad (1 - \omega(x))^B = (-1)^A x \omega(x)^A.$$

The Nahm equation has been studied by several authors including [Zag07, Sec.3], [Vla, VZ11], [RV, Sec.4].

1.3. The case of the 4_1 knot. We now specialize Corollary 1.2 to the invariant of the 4_1 and 5_2 knots is given by [KLV12, AK]

$$\mathcal{I}_{1,2} = \mathcal{I}_{4_1} \quad \mathcal{I}_{2,3} = \mathcal{I}_{5_2}.$$

In this section, let

$$(13) \quad F(q, x) = F_{1,2}(q, x) = \sum_{n=0}^{\infty} (-1)^n \frac{q^{\frac{1}{2}n(n+1)}}{(q)_n^2} x^n.$$

Corollary 1.7. (a) We have:

$$(14) \quad \mathcal{I}_{4_1}(b) = -\frac{i}{2} \left(\frac{q}{\tilde{q}} \right)^{\frac{1}{24}} (bG(q)g(\tilde{q}) - b^{-1}G(\tilde{q})g(q))$$

where

$$(15a) \quad g(q) = \sum_{n=0}^{\infty} (-1)^n \frac{q^{\frac{1}{2}n(n+1)}}{(q)_n^2}$$

$$(15b) \quad G(q) = \sum_{m=0}^{\infty} \left(1 + 2m - 4 \sum_{s=1}^{\infty} \frac{q^{s(m+1)}}{1 - q^s} \right) (-1)^m \frac{q^{\frac{1}{2}m(m+1)}}{(q)_m^2}$$

(b) The series $g(q)$ and $G(q)$ are given in terms of $F(q, x)$ by:

$$(16a) \quad g(q) = \langle F \rangle$$

$$(16b) \quad G(q) = \langle (2 + 2\delta - 4\delta_1)F \rangle$$

(c) F satisfies the linear q -difference equation

$$(17) \quad F(q, q^{-1}x) + F(q, qx) = (2 - x)F(q, x)$$

The series $g(q)$ that appears in Theorem 1.7 was known to the first author and Zagier to be closely related to the 4_1 knot. For a detailed discussion of experimental facts below, see [GZ]. Empirically, it appears that

- the pair $(g(q), G(q))$ is related to the 3D index of the 4_1 knot,
- the radial asymptotics of the pair $(g(q), G(q))$ are related to the asymptotics of the Kashaev invariant of the 4_1 knot,
- the above observations for 4_1 also hold for the case of 5_2 knot discussed below.

Recall that the index of an ideal triangulation was introduced in [DGGb, DGGa], necessary and sufficient conditions for its convergence was established in [Gar] and its topological invariance was proven in [GHRs]. For a detailed discussion of the above experimental facts, see [GZ].

1.4. **The case of the 5_2 knot.** In this section, let

$$F(q, x) = F_{2,3}(q, x) = \sum_{m=0}^{\infty} t_m(q)x^m, \quad \tilde{F}(q, \tilde{x}) = F_{1,3}(q, \tilde{x}) = \sum_{m=0}^{\infty} T_m(q)\tilde{x}^m$$

where

$$t_m(q) = \frac{q^{m(m+1)}}{(q)_m^3}, \quad T_n(q) = (-1)^n \frac{q^{\frac{1}{2}n(n+1)}}{(q)_n^3} = t_n(q^{-1}).$$

Let

$$\begin{aligned} R_{m,n}(q, \tilde{q}) &= -\frac{b^2}{2} \left(1 + 4m + 4m^2 - 6E_1^{(m)}(q) - 12mE_1^{(m)}(q) + 9E_1^{(m)2}(q) - 3E_2^{(m)}(q) \right) \\ &\quad - \frac{1}{2\pi i} + \frac{1}{2} \left(1 + 2m - 3E_1^{(m)}(q) \right) \left(1 + 2n - 6E_1^{(n)}(\tilde{q}) \right) \\ &\quad + \frac{b^{-2}}{2} \left(-n - n^2 - 6E_2^{(0)}(\tilde{q}) + 3E_1^{(n)}(\tilde{q}) + 6nE_1^{(n)}(\tilde{q}) - 9E_1^{(n)2}(\tilde{q}) + 3E_2^{(n)}(\tilde{q}) \right), \end{aligned}$$

where $E_i^{(m)}(q)$ are defined in Equation (29a). For $k = 1, \dots, 4$ let

$$(18) \quad g_k(q) = \sum_{m=0}^{\infty} p_k(m)t_m(q), \quad G_k(\tilde{q}) = \sum_{n=0}^{\infty} P_k(n)T_n(\tilde{q}),$$

where

$$(19a) \quad p_{1,m}(q) = 1 + 4m + 4m^2 - 6E_1^{(m)}(q) - 12mE_1^{(m)}(q) + 9E_1^{(m)2}(q) - 3E_2^{(m)}(q)$$

$$(19b) \quad p_{2,m}(q) = 1 + 2m - 3E_1^{(m)}(q)$$

$$(19c) \quad p_{3,m}(q) = 1$$

and

$$(20a) \quad P_{1,m}(q) = 1$$

$$(20b) \quad P_{2,m}(q) = 1 + 2n - 6E_1^{(n)}(\tilde{q})$$

$$(20c) \quad P_{3,m}(q) = -n - n^2 - 6E_2^{(0)}(\tilde{q}) + 3E_1^{(n)}(\tilde{q}) + 6nE_1^{(n)}(\tilde{q}) - 9E_1^{(n)2}(\tilde{q}) + 3E_2^{(n)}(\tilde{q}).$$

Corollary 1.8. (a) We have:

(21)

$$\begin{aligned} \mathcal{I}_{2,3}(q) &= -e^{\frac{3\pi i}{4}} \left(\frac{q}{\tilde{q}} \right)^{\frac{1}{8}} \sum_{m,n=0}^{\infty} R_{m,n}(q, \tilde{q}) t_m(q) T_n(\tilde{q}) \\ &= -e^{\frac{3\pi i}{4}} \left(\frac{q}{\tilde{q}} \right)^{\frac{1}{8}} \left(-\frac{b^2}{2} g_1(q) G_1(\tilde{q}) - \frac{1}{2\pi i} g_3(q) G_1(\tilde{q}) + \frac{1}{2} g_2(q) G_2(\tilde{q}) + \frac{b^{-2}}{2} g_3(q) G_3(\tilde{q}) \right) \end{aligned}$$


(b) F and \tilde{F} satisfy the linear q -difference equations

$$\begin{aligned} F(q, q^3x) - (3 - q^2x)F(q, q^2x) + 3F(q, qx) - F(q, x) &= 0 \\ \tilde{F}(q, q^3x) - 3\tilde{F}(q, q^2x) + (3 - q^2x)\tilde{F}(q, qx) - \tilde{F}(q, x) &= 0. \end{aligned}$$

Remark 1.9. A computation gives that $P(A, B) = P(B - A, B)$ for $(A, B) = (1, 2)$ and $(A, B) = (2, 3)$ corresponding to the invariants of the 4_1 and 5_2 knots. In all other cases that we tried, we found that $P(A, B)$ is not equal to $P(B - A, B)$.

2. PROOFS

2.1. A residue computation. To relate the state-integral $\mathcal{I}_{A,B}$ to a sum, we will apply the residue theorem on a semicircle γ_R with center 0 and radius R , oriented counterclockwise in the upper half-plane:



Then, we will take the limit $R \rightarrow \infty$. To compute the residue of the integrand, we need to expand $\Phi_b(x)$ near the pole

$$x_{m,n} = c_b + ibm + ib^{-1}n$$

for natural numbers m and n . Let

$$(23) \quad \phi_m(x) = \frac{(q^{m+1}e^x; q)_\infty}{(q^{m+1}; q)_\infty}$$

$$(24) \quad \tilde{\phi}_n(x) = \frac{(\tilde{q}; \tilde{q})_\infty (\tilde{q}^{-1}; \tilde{q}^{-1})_n}{(\tilde{q}e^x; \tilde{q})_\infty (\tilde{q}^{-1}e^x; \tilde{q}^{-1})_n}$$

Lemma 2.1. We have:

$$(25) \quad \Phi_b(x + x_{m,n}) = \frac{(q; q)_\infty}{(\tilde{q}; \tilde{q})_\infty} \frac{1}{(q; q)_m} \frac{1}{(\tilde{q}^{-1}; \tilde{q}^{-1})_n} \frac{\phi_m(2\pi bx) \tilde{\phi}_n(2\pi b^{-1}x)}{1 - e^{2\pi b^{-1}x}}.$$

Proof. Equation (1) implies the functional equations

$$\frac{\Phi_b(x + c_b + ib)}{\Phi_b(x + c_b)} = \frac{1}{1 - qe^{2\pi bx}}$$

$$\frac{\Phi_b(x + c_b + ib^{-1})}{\Phi_b(x + c_b)} = \frac{1}{1 - \tilde{q}^{-1}e^{2\pi b^{-1}x}}$$

which give

$$\Phi_b(x + x_{m,n}) = \Phi_b(x + c_b) \frac{1}{(qe^{2\pi bx}; q)_m} \frac{1}{(\tilde{q}^{-1}e^{2\pi b^{-1}x}; \tilde{q}^{-1})_n}$$

$$\Phi_b(x + c_b) = \frac{1}{1 - e^{2\pi b^{-1}x}} \frac{(qe^{2\pi bx}; q)_\infty}{(\tilde{q}e^{2\pi b^{-1}x}; \tilde{q})_\infty}.$$

Thus,

$$\begin{aligned} \Phi_b(x + x_{m,n}) &= \frac{(q; q)_\infty}{(\tilde{q}; \tilde{q})_\infty} \frac{1}{(q; q)_m} \frac{1}{(\tilde{q}^{-1}; \tilde{q}^{-1})_n} \\ &= \frac{1}{1 - e^{2\pi b^{-1}x}} \frac{(qe^{2\pi bx}; q)_\infty}{(q; q)_\infty} \frac{(\tilde{q}; \tilde{q})_\infty}{(\tilde{q}e^{2\pi b^{-1}x}; \tilde{q})_\infty} \frac{(q; q)_m}{(qe^{2\pi bx}; q)_m} \frac{(\tilde{q}^{-1}; \tilde{q}^{-1})_n}{(\tilde{q}^{-1}e^{2\pi b^{-1}x}; \tilde{q}^{-1})_n} \\ &= \frac{(q; q)_\infty}{(\tilde{q}; \tilde{q})_\infty} \frac{1}{(q; q)_m} \frac{1}{(\tilde{q}^{-1}; \tilde{q}^{-1})_n} \\ &= \frac{1}{1 - e^{2\pi b^{-1}x}} \frac{(q^{m+1}e^{2\pi bx}; q)_\infty}{(q^{m+1}; q)_\infty} \frac{(\tilde{q}; \tilde{q})_\infty}{(\tilde{q}e^{2\pi b^{-1}x}; \tilde{q})_\infty} \frac{(\tilde{q}^{-1}; \tilde{q}^{-1})_n}{(\tilde{q}^{-1}e^{2\pi b^{-1}x}; \tilde{q}^{-1})_n} \end{aligned}$$

The result follows. \square

The decoupling of (m, n) in the quadratic form comes as follows: since A, m, n are integers, $e^{A\pi imn} = 1$ and a computation gives

$$e^{-A\pi i(x+x_n, m)^2} = i^A \left(\frac{q}{\tilde{q}} \right)^{\frac{A}{8}} t_m^A(q) \tilde{t}_n^A(\tilde{q}) e^{-A\pi ix^2 + 2A\pi x(b(m+\frac{1}{2}) + b^{-1}(n+\frac{1}{2}))}$$

where

$$t_m^A(q) = (-1)^{Am} q^{A\frac{m(m+1)}{2}}, \quad \tilde{t}_n^A(\tilde{q}) = (-1)^{An} \tilde{q}^{-A\frac{n(n+1)}{2}}.$$

The Dedekind function $\eta(\tau) = q^{1/24}(q; q)_\infty$ (with $q = e^{2\pi i\tau}$) satisfies the modular equation $\eta(-\tau^{-1}) = \sqrt{-i\tau}\eta(\tau)$ [And76]. It follows that

$$(26) \quad \frac{(q; q)_\infty}{(\tilde{q}; \tilde{q})_\infty} = e^{\frac{\pi i}{4}} \left(\frac{\tilde{q}}{q} \right)^{\frac{1}{24}} b^{-1}.$$

After we set $w = x/(2\pi)$, the above discussion implies that

$$(27) \quad \mathcal{I}_{A,B}(b) = \left(\frac{\tilde{q}}{q} \right)^{\frac{B-3A}{24}} e^{\pi i \frac{B+2(A+1)}{4}} \sum_{m,n=0}^{\infty} (\text{Res}_{w=0} F_{A,B,m,n}(w)) \frac{t_m^A(q)}{(q; q)_m^B} \frac{\tilde{t}_n^A(\tilde{q})}{(\tilde{q}^{-1}; \tilde{q}^{-1})_n^B},$$

where

$$(28) \quad F_{A,B,m,n}(w) = e^{\frac{A}{4\pi i}w^2 + Aw(b(m+\frac{1}{2})+b^{-1}(n+\frac{1}{2}))} \left(\frac{\phi_m(bw) \tilde{\phi}_n(b^{-1}w)}{b(1-e^{b^{-1}w})} \right)^B.$$

2.2. The Taylor series of $\phi_m(x)$ and $\tilde{\phi}_n(x)$. In this section we express the Taylor series of $\phi_m(x)$ and $\tilde{\phi}_n(x)$ in terms of the q -series $E_l^{(m)}(q)$ and $\tilde{E}_l^{(m)}(\tilde{q})$ defined by:

$$(29a) \quad E_l^{(m)}(q) = \sum_{s=1}^{\infty} \frac{s^{l-1} q^{s(m+1)}}{1-q^s} = \langle \delta_l(x^m) \rangle$$

$$(29b) \quad \tilde{E}_l^{(n)}(\tilde{q}) = \begin{cases} -n + E_1^{(n)}(\tilde{q}) & \text{if } l = 1 \\ E_l^{(n)}(\tilde{q}) & \text{if } l > 1 \text{ is odd} \\ 2E_l^{(0)}(\tilde{q}) - E_l^{(n)}(\tilde{q}) & \text{if } l > 1 \text{ is even} \end{cases}$$

Proposition 2.2. We have:

$$(30a) \quad \phi_m(x) = \exp \left(- \sum_{l=1}^{\infty} \frac{1}{l!} E_l^{(m)}(q) x^l \right)$$

$$(30b) \quad \tilde{\phi}_n(x) = \exp \left(\sum_{l=1}^{\infty} \frac{1}{l!} \tilde{E}_l^{(m)}(\tilde{q}) x^l \right).$$

The proof of this proposition is given in Section 2.6. The first few terms in Equations (30a)-(30a) are given by:

$$(31a) \quad \phi_m(x) = \exp \left(-E_1^{(m)}x - \frac{1}{2}E_2^{(m)}x^2 - \frac{1}{6}E_3^{(m)}x^3 - \frac{1}{24}E_4^{(m)}x^4 - \dots \right)$$

$$(31b) \quad = 1 - E_1^{(m)}x + \frac{1}{2}(E_1^{(m)2} - E_2^{(m)})x^2 + \frac{1}{6}(-E_1^{(m)3} + 3E_1^{(m)}E_2^{(m)} - E_3^{(m)})x^3 + \frac{1}{24}(E_1^{(m)4} - 6E_1^{(m)2}E_2^{(m)} + 3E_2^{(m)2} + 4E_1^{(m)}E_3^{(m)} - E_4^{(m)})x^4 + \dots$$

$$(31c) \quad \tilde{\phi}_n(x) = \exp \left(\tilde{E}_1^{(n)}x + \frac{1}{2}\tilde{E}_2^{(n)}x^2 + \frac{1}{6}\tilde{E}_3^{(n)}x^3 + \frac{1}{24}\tilde{E}_4^{(n)}x^4 - \dots \right)$$

$$= 1 + \tilde{E}_1^{(n)}x + \frac{1}{2}(\tilde{E}_1^{(n)2} + \tilde{E}_2^{(n)})x^2 + \frac{1}{6}(\tilde{E}_1^{(n)3} + 3\tilde{E}_1^{(n)}\tilde{E}_2^{(n)} + \tilde{E}_3^{(n)})x^3 +$$

$$(31d) \quad \frac{1}{24}(\tilde{E}_1^{(n)4} + 6\tilde{E}_1^{(n)2}\tilde{E}_2^{(n)} + 3\tilde{E}_2^{(n)2} + 4\tilde{E}_1^{(n)}\tilde{E}_3^{(n)} + \tilde{E}_4^{(n)})x^4 + \dots$$

where $E_l^{(m)} = E_l^{(m)}(q)$ and $\tilde{E}_l^{(m)} = \tilde{E}_l^{(m)}(\tilde{q})$.

2.3. The connection with the differential operators δ_l and $\tilde{\delta}_l$. In this section we connect the series $E_l^{(m)}(q)$ and $\tilde{E}_l^{(m)}(\tilde{q})$ with the action of the differential operators δ_l and $\tilde{\delta}_l$ on a series $F(x)$ and $\tilde{F}(\tilde{x})$ respectively. Consider formal power series

$$F(x) = \sum_{m=0}^{\infty} t(m)x^m \quad \tilde{F}(\tilde{x}) = \sum_{m=0}^{\infty} \tilde{t}(m)\tilde{x}^m.$$

Lemma 2.3. We have:

$$(32) \quad \sum_{m=0}^{\infty} \left(\prod_{j=1}^r E_{l_j}^{(m)}(q) \right) t(m) = \left\langle \prod_{j=1}^r \delta_{l_j} F \right\rangle$$

$$(33) \quad \sum_{m=0}^{\infty} m^r t(m) = \langle \delta^r F \rangle$$

and

$$(34) \quad \sum_{n=0}^{\infty} \left(\prod_{j=1}^r \tilde{E}_{l_j}^{(n)}(\tilde{q}) \right) \tilde{t}(n) = \left\langle \prod_{j=1}^r \tilde{\delta}_{l_j} \tilde{F} \right\rangle$$

$$(35) \quad \sum_{n=0}^{\infty} n^r \tilde{t}(n) = \langle \tilde{d}^r \tilde{F} \rangle.$$

Proof. For a positive natural number l we have:

$$\sum_{m=0}^{\infty} E_l^{(m)}(q) t(m) = \sum_{m=0}^{\infty} \langle \delta_l(x^m) \rangle t(m) = \left\langle \delta_l \left(\sum_{m=0}^{\infty} t(m) x^m \right) \right\rangle = \langle \delta_l F \rangle.$$

Moreover, for positive natural numbers l, l' we have:

$$\begin{aligned} \sum_{m=0}^{\infty} E_l^{(m)}(q) E_{l'}^{(m)}(q) t(m) &= \sum_{m=0}^{\infty} \langle \delta_l(x^m) \rangle \langle \delta_{l'}(x^m) \rangle t(m) \\ &= \left\langle \delta_l \left(\sum_{m=0}^{\infty} \langle \delta_{l'}(x^m) \rangle t(m) x^m \right) \right\rangle. \end{aligned}$$

Now,

$$\langle \delta_{l'}(x^m) \rangle t(m) x^m = \sum_{s=1}^{\infty} \frac{s^{l'-1} q^s}{1 - q^s} q^{sm} t(m) x^m = \delta_{l'}(x^m) t(m)$$

and summing up over m , we obtain that

$$\sum_{m=0}^{\infty} \langle \delta_{l'}(x^m) \rangle t(m) x^m = \delta_{l'} F(q, x).$$

It follows that

$$\sum_{m=0}^{\infty} E_l^{(m)}(q) E_{l'}^{(m)}(q) t(m) = \langle \delta_l \delta_{l'} F \rangle.$$

The general case of Equation (32) follows by induction on r . Equation (33) is obvious. \square

2.4. **Proof of Theorem 1.1.** Fix natural numbers A and B with $B > A \geq 1$, and let

$$t(m) = \frac{(-1)^{Am} q^{A \frac{m(m+1)}{2}}}{(q)_m^B}, \quad F(q, x) = \sum_{m=0}^{\infty} t(m) x^m$$

and

$$\tilde{t}(n) = \frac{(-1)^{(B-A)n} \tilde{q}^{(B-A) \frac{n(n+1)}{2}}}{(\tilde{q})_n^B}, \quad \tilde{F}(\tilde{q}, \tilde{x}) = \sum_{n=0}^{\infty} \tilde{t}(n) \tilde{x}^n.$$

Use Equations (27) and (28) and Proposition 2.2 to expand $F_{A,B,m,n}(w)$ as a power series with coefficients polynomials in the variables $m, E_l^{(m)}(q)$ and $n, \tilde{E}_l^{(n)}(\tilde{q})$ and $b^{\pm 1}$ and $(2\pi i)^{-1}$. Now apply Lemma 2.3 to convert the variables $m, E_l^{(m)}(q), n, \tilde{E}_l^{(n)}(\tilde{q})$ in terms of the action of the operators $\delta, \delta_l, \tilde{\delta}, \tilde{\delta}_l$ respectively. This concludes the proof of Theorem 1.1. \square

2.5. **Some auxiliary power series.** Consider the auxiliary series

$$(36) \quad \frac{1}{ae^x - 1} = \sum_{n=0}^{\infty} p_n(a) x^n$$

where

$$p_n(a) = -\frac{a}{n!(1-a)^{n+1}} \sum_{m=0}^{n-1} A_{n,m} a^m \quad p_0(a) = -\frac{1}{1-a}$$

and $A_{n,m}$ are *Euler triangular numbers* (sequence A008292 in the online encyclopedia of integer sequences [Slo]) that satisfy the recursion

$$A_{n,m} = (n-m)A_{n-1,m-1} + (m+1)A_{n-1,m}$$

and also given by the sum

$$A_{n,m} = \sum_{k=0}^m (-1)^k \binom{n+1}{k} (m+1-k)^n.$$

For a detailed discussion on this subject, see [FS70]. A table of the first few numbers $A_{n,m}$ is given by

$n \setminus m$	0	1	2	3	4	5	6	7	8
1	1								
2	1	1							
3	1	4	1						
4	1	11	11	1					
5	1	26	66	26	1				
6	1	57	302	302	57	1			
7	1	120	1191	2416	1191	120	1		
8	1	247	4293	15619	15619	4293	247	1	
9	1	502	14608	88234	156190	88234	14608	502	1

Lemma 2.4. For $l \geq 1$, we have:

$$(37) \quad \frac{d^l}{dx^l} \log(1 - q^k e^{bx})|_{x=0} = b^l p_{l-1}(q^k) + b \delta_{l,1}$$

Proof. It follows from

$$\frac{d}{dx} \log(1 - q^k e^{bx}) = b \left(1 + \frac{1}{q^k e^{bx} - 1} \right)$$

and Equation (36). \square

For positive natural numbers l, r with $l \geq r$ and m consider the q -series $E_{l,r}^{(m)}(q)$ defined by

$$(38) \quad E_{l,r}^{(m)}(q) = \sum_{k=m+1}^{\infty} \frac{q^{kr}}{(1 - q^k)^l}$$

Lemma 2.5. (a) We have

$$(39) \quad E_{l,r}^{(m)}(q) = \sum_{s=r}^{\infty} a_{l,s} \frac{q^{s(m+1)}}{1 - q^s}$$

where

$$\frac{x^r}{(1 - x)^l} = \sum_{s=r}^{\infty} a_{l,s} x^s$$

(b) It follows that

$$(40) \quad \sum_{r=0}^{l-1} A_{l-1,r} E_{l,r+1}^{(m)}(q) = E_l^{(m)}(q)$$

Proof. For (a), interchange k and s summation:

$$E_{l,r}^{(m)}(q) = \sum_{k=m+1}^{\infty} \sum_{s=r}^{\infty} a_{l,s} q^{sk} = \sum_{s=r}^{\infty} \sum_{k=m+1}^{\infty} a_{l,s} q^{sk} = \sum_{s=r}^{\infty} q^{(m+1)s} \sum_{k=0}^{\infty} a_{l,s} q^{sk} = \sum_{s=r}^{\infty} a_{l,s} \frac{q^{(m+1)s}}{1 - q^s}$$

(b) follows from (a) and the fact that

$$\frac{\sum_{r=0}^{l-1} A_{l-1,r} x^r}{(1 - x)^l} = \sum_{s=1}^{\infty} s^{l-1} x^s.$$

\square

Lemma 2.6. We have:

$$(41) \quad \phi_m(x) = \exp \left(- \sum_{l=1}^{\infty} \frac{1}{l!} \sum_{r=0}^{l-1} A_{l-1,r} E_{l,r+1}^{(m)}(q) x^l \right)$$

Proof. It follows from Lemma 2.4 combined with

$$\log(\phi_m(x)) = \log \left(\frac{(q^{m+1} e^x; q)_{\infty}}{(q^{m+1}; q)_{\infty}} \right) = \sum_{l=m+1}^{\infty} (\log(1 - q^l e^x) - \log(1 - q^l))$$

\square

2.6. **Proof of Proposition 2.2.** Part (a) of Proposition 2.2 follows from Lemma 2.5 and Lemma 2.6. For part (b), we will use the series

$$E_l^{[m]}(q) = \sum_{s=1}^{\infty} \frac{s^{k-1} q^{s(m+1)}}{1 - q^s}$$

Using

$$\log(\tilde{\phi}_n(x)) = \log\left(\frac{(\tilde{q}; \tilde{q})_{\infty}}{(\tilde{q}e^x; \tilde{q})_{\infty}}\right) + \log\left(\frac{(\tilde{q}^{-1}; \tilde{q}^{-1})_n}{(\tilde{q}^{-1}e^x; \tilde{q}^{-1})_n}\right)$$

and the proof of part (a) of Proposition 2.2, it follows that

$$\begin{aligned} \log(\tilde{\phi}_n(x)) &= \log\left(\frac{(\tilde{q}; \tilde{q})_{\infty}}{(\tilde{q}e^x; \tilde{q})_{\infty}}\right) + \log\left(\frac{(\tilde{q}^{-1}; \tilde{q}^{-1})_n}{(\tilde{q}^{-1}e^x; \tilde{q}^{-1})_n}\right) \\ &= \sum_{l=1}^{\infty} \frac{1}{l!} \sum_{r=0}^{l-1} A_{l-1,r} E_{l,r+1}^{(0)}(\tilde{q}) x^l + \sum_{l=1}^{\infty} \frac{1}{l!} \sum_{r=0}^{l-1} A_{l-1,r} E_{l,r+1}^{[n]}(\tilde{q}^{-1}) x^l \\ &= \sum_{l=1}^{\infty} \frac{1}{l!} \sum_{r=0}^{l-1} A_{l-1,r} \left(E_{l,r+1}^{(0)}(\tilde{q}) + E_{l,r+1}^{[n]}(\tilde{q}^{-1}) \right) x^l \end{aligned}$$

where

$$(42) \quad E_{l,r}^{[n]}(q) = \sum_{k=1}^n \frac{q^{kr}}{(1 - q^k)^l}.$$

Let

$$(43) \quad \tilde{E}_{l,r}^{(n)}(\tilde{q}) = \begin{cases} -n + E_{1,1}^{(n)}(\tilde{q}) & \text{if } l = r = 1 \\ E_{l,r}^{(n)}(\tilde{q}) & \text{if } l > 1 \text{ is odd} \\ 2E_{l,r}^{(0)}(\tilde{q}) - E_{l,r}^{(n)}(\tilde{q}) & \text{if } l > 1 \text{ is even} \end{cases}$$

We claim that

$$(44) \quad E_{l,r}^{(0)}(\tilde{q}) + E_{l,l-r}^{[n]}(\tilde{q}^{-1}) = \tilde{E}_{l,r}^{(n)}(\tilde{q})$$

for $l > r \geq 1$ and

$$(45) \quad E_{1,1}^{(0)}(\tilde{q}) + E_{1,1}^{[n]}(\tilde{q}^{-1}) = \tilde{E}_{1,1}^{(n)}(\tilde{q})$$

Assuming Equations (44) and (45), it follows that

$$\begin{aligned} \log(\tilde{\phi}_n(x)) &= \sum_{l=1}^{\infty} \frac{1}{l!} \sum_{r=0}^{l-1} A_{l-1,r} \tilde{E}_{l,r+1}^{(n)}(\tilde{q}) x^l \\ &= \sum_{l=1}^{\infty} \frac{1}{l!} \tilde{E}_l^{(n)}(\tilde{q}) x^l \end{aligned}$$

where the last step follows from part (b) of Lemma 2.5.

It remains to prove Equations (44) and (45). Equation (44) follows from the definition of $\widetilde{E}_{1,1}^{(n)}(\tilde{q})$ and

$$\begin{aligned} E_{l,r}^{(0)}(\tilde{q}) + E_{l,l-r}^{[n]}(\tilde{q}^{-1}) &= \sum_{k=1}^{\infty} \frac{\tilde{q}^{kr}}{(1-\tilde{q}^k)^l} + \sum_{k=1}^n \frac{\tilde{q}^{-k(l-r)}}{(1-\tilde{q}^{-k})^l} \\ &= \sum_{k=1}^{\infty} \frac{\tilde{q}^{kr}}{(1-\tilde{q}^k)^l} + (-1)^l \sum_{k=1}^n \frac{\tilde{q}^{kr}}{(1-\tilde{q}^k)^l} \\ &= (1 + (-1)^l) \sum_{k=1}^n \frac{\tilde{q}^{kr}}{(1-\tilde{q}^k)^l} + \sum_{k=n+1}^{\infty} \frac{\tilde{q}^{kr}}{(1-\tilde{q}^k)^l} \end{aligned}$$

Equation (45) follows from

$$\begin{aligned} E_{1,1}^{(0)}(\tilde{q}) + E_{1,1}^{[n]}(\tilde{q}^{-1}) &= \sum_{k=1}^{\infty} \frac{\tilde{q}^k}{1-\tilde{q}^k} + \sum_{k=1}^n \frac{\tilde{q}^{-k}}{1-\tilde{q}^{-k}} \\ &= \sum_{k=1}^{\infty} \frac{1-1+\tilde{q}^k}{1-\tilde{q}^k} - \sum_{k=1}^n \frac{1}{1-\tilde{q}^k} = -n + \sum_{k=n+1}^{\infty} \frac{\tilde{q}^k}{1-\tilde{q}^k} \end{aligned}$$

This completes the proof of Proposition 2.2. \square

2.7. Proof of Lemma 1.5. Part (a) of Lemma 1.5 follows from the definition of $F_{A,B}$ and $\widetilde{F}_{A,B}$.

Part (b) follows from an application of Zeilberger's creative telescoping [Zei91]. To apply the method, define

$$t(m, x) = \frac{(-1)^{Am} q^{A \frac{m(m+1)}{2}}}{(q)_m^B} x^m$$

Then, observe that t satisfies the recursions with respect to m and x :

$$(1 - q^{m+1})^B t(m+1, x) = (-1)^A q^{A(m+1)} t(m, x) \quad t(m, qx) = q^m t(m, x).$$

Now, we eliminate q^m from the above equations as follows. The second equation implies that $t(m+1, q^j x) = q^{j(m+1)} t(m+1, x)$. Expanding the first equation, it follows that

$$\sum_{j=0}^B (-1)^j \binom{B}{j} t(m+1, q^j x) = (-1)^A q^A x t(m, q^A x)$$

Summing for $m \geq 0$ implies (b). \square

Proof. (of Corollary 1.6) The admissibility of F in the sense of Kontsevich-Soibelman, follows from [KS11, Sec.6.1] and [KS11, Thm.9]. Given this, the Nahm Equation (12) for ω follows easily from part (b) of Lemma 1.5. \square

3. AN APPLICATION: STATE-INTEGRALS OF THE 4_1 AND 5_2 KNOTS

3.1. **Proof of Corollary 1.7.** Assume now that $(A, B) = (1, 2)$. Then,

$$\begin{aligned} \frac{1}{(b(1 - e^{b^{-1}w}))^2} &= \frac{1}{w^2} - \frac{b^{-1}}{w} + O(1) \\ (\phi_m(bw))^2 &= 1 - 2E_1^{(m)}(q)bw + O(w^2) \\ (\tilde{\phi}_n(b^{-1}w))^2 &= 1 + 2\tilde{E}_1^{(n)}(\tilde{q})b^{-1}w + O(w^2) \\ e^{\frac{1}{4\pi i}w^2 + w(b(m+1/2) + b^{-1}(n+1/2))} &= 1 + \left(\frac{1}{2} + m\right)bw + \left(\frac{1}{2} + n\right)b^{-1}w + O(w^2) \end{aligned}$$

Combined with $\tilde{E}_1^{(n)}(\tilde{q}) = -n + E_1^{(n)}(\tilde{q})$, it follows that the residue $R = \text{Res}_{w=0}(F_{1,2,m,n}(w))$ is given by

$$R = \left(b \left(\frac{1}{2} + m - 2E_1^{(m)}(q) \right) - b^{-1} \left(\frac{1}{2} + n - 2E_1^{(n)}(\tilde{q}) \right) \right)$$

The above, together with the fact that $t_n(q) = (-1)^n \frac{q^{\frac{1}{2}n(n+1)}}{(q)_n^2}$ satisfies $t_n(q^{-1}) = t_n(q)$ implies Equation (14). Equation (17) follows from Equation (11) for $(A, B) = (1, 2)$.

3.2. **Proof of Corollary 1.8.** Assume now that $(A, B) = (2, 3)$. Then,

$$\begin{aligned} \frac{1}{(b(1 - e^{b^{-1}w}))^3} &= -\frac{1}{w^3} + \frac{3b^{-1}}{2w^2} - \frac{b^{-2}}{w} + O(1) \\ (\phi_m(bw))^3 &= 1 - 3E_1^{(m)}(q)bw + \frac{3}{2} \left(3E_1^{(m)2}(q) - E_2^{(m)}(q) \right) b^2w^2 + O(w^3) \\ (\tilde{\phi}_n(b^{-1}w))^3 &= 1 + 3\tilde{E}_1^{(n)}(\tilde{q})b^{-1}w + \frac{3}{2} \left(3\tilde{E}_1^{(n)2}(\tilde{q}) + \tilde{E}_2^{(n)}(\tilde{q}) \right) b^{-2}w^2 + O(w^3) \\ e^{\frac{2}{4\pi i}w^2 + 2w(b(m+1/2) + b^{-1}(n+1/2))} &= 1 + ((1 + 2m)b + (1 + 2n)b^{-1})w + \\ &\quad \left(1 + \frac{b^2 + b^{-2}}{2} + \frac{1}{2\pi i} + 2b^2m^2 + 2b^{-2}n^2 + 4mn \right. \\ &\quad \left. + 2(1 + b^2)m + 2(1 + b^{-2})n \right) w^2 + O(w^3) \end{aligned}$$

If $R = \text{Res}_{w=0}(F_{2,3,m,n}(w))$, then

$$\begin{aligned} R_{m,n} &= -\frac{b^2}{2} \left(1 + 4m + 4m^2 - 6E_1^{(m)}(q) - 12mE_1^{(m)}(q) + 9E_1^{(m)2}(q) - 3E_2^{(m)}(q) \right) \\ &\quad - \frac{1}{2\pi i} + \frac{1}{2} \left(1 + 2m - 3E_1^{(m)}(q) \right) \left(1 + 2n - 6E_1^{(n)}(\tilde{q}) \right) \\ &\quad + \frac{b^{-2}}{2} \left(-n - n^2 - 6E_2^{(0)}(\tilde{q}) + 3E_1^{(n)}(\tilde{q}) + 6nE_1^{(n)}(\tilde{q}) - 9E_1^{(n)2}(\tilde{q}) + 3E_2^{(n)}(\tilde{q}) \right), \end{aligned}$$

This proves part (a) of Corollary 1.8. Part (b) follows from Equation (11) for $(A, B) = (2, 3)$ and $(A, B) = (1, 3)$. Note that Theorem 1.1 states that

$$(46) \quad \mathcal{I}_{2,3}(q) = -e^{\frac{3\pi i}{4}} \langle P_{2,3}(F\tilde{F}) \rangle$$

where

$$\begin{aligned}
P_{2,3} = & -\frac{b^2}{2} \left(1 + 4\delta + 4\delta^2 - 6\delta_1 - 12\delta\delta_1 + 9\delta_1^2 - 3\tilde{\delta}_2 \right) \\
& + \frac{1}{2} \left(1 + 2\delta + \frac{i}{\pi} + 2\tilde{\delta} + 4\delta\tilde{\delta} - 3\delta_1 - 6\tilde{\delta}\delta_1 - 6e_2(\tilde{q}) - 6\tilde{\delta}_1 - 12\delta\tilde{\delta}_1 + 18\delta_1\tilde{\delta}_1 \right) \\
& + \frac{b^{-2}}{2} \left(-\tilde{\delta} - \tilde{\delta}^2 + 3\tilde{\delta}_1 + 6\tilde{\delta}\tilde{\delta}_1 - 9\tilde{\delta}_1^2 + 3\tilde{\delta}_2 \right).
\end{aligned}$$

Acknowledgments. The paper was conceived during a visit of the first author in Geneva in the spring of 2013. The first author wishes to thank the University of Geneva for its hospitality, and Don Zagier for encouragement and stimulating conversations.

REFERENCES

- [AK] Jørgen E. Andersen and Rinat M. Kashaev, *A TQFT from quantum Teichmüller theory*, arXiv:1109.6295, Preprint 2011.
- [And76] George E. Andrews, *The theory of partitions*, Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976, Encyclopedia of Mathematics and its Applications, Vol. 2.
- [DGGa] Tudor Dimofte, Davide Gaiotto, and Sergei Gukov, *3-manifolds and 3d indices*, arXiv:1112.5179, Preprint 2011.
- [DGGb] ———, *Gauge theories labelled by three-manifolds*, arXiv:1108.4389, Preprint 2011.
- [DGLZ09] Tudor Dimofte, Sergei Gukov, Jonatan Lenells, and Don Zagier, *Exact results for perturbative Chern-Simons theory with complex gauge group*, Commun. Number Theory Phys. **3** (2009), no. 2, 363–443.
- [Fad95] L. D. Faddeev, *Discrete Heisenberg-Weyl group and modular group*, Lett. Math. Phys. **34** (1995), no. 3, 249–254.
- [FS70] Dominique Foata and Marcel-P. Schützenberger, *Théorie géométrique des polynômes eulériens*, Lecture Notes in Mathematics, Vol. 138, Springer-Verlag, Berlin, 1970.
- [Gar] Stavros Garoufalidis, *The 3D index of an ideal triangulation and angle structures*, arXiv:1208.1663, Preprint 2012.
- [GHRS] Stavros Garoufalidis, Craig D. Hodgson, Hyam Rubinstein, and Henry Segerman, *1-efficient triangulations and the index of a cusped hyperbolic 3-manifold*, arXiv:1303.5278, Preprint 2013.
- [GZ] Stavros Garoufalidis and Don Zagier, *Empirical relations between q -series and kashaev’s invariant of knots*, Preprint 2013.
- [Hik01] Kazuhiro Hikami, *Hyperbolic structure arising from a knot invariant*, Internat. J. Modern Phys. A **16** (2001), no. 19, 3309–3333.
- [KLV12] Rinat M. Kashaev, Feng Luo, and Grigory Vartanov, *A TQFT of turaev-viro type on shaped triangulations*, 2012, arXiv:1210.8393, Preprint.
- [KS11] Maxim Kontsevich and Yan Soibelman, *Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson-Thomas invariants*, Commun. Number Theory Phys. **5** (2011), no. 2, 231–352.
- [RV] Fernando Rodriguez-Villegas, *A refinement of the A -polynomial of quivers*, Preprint 2011.
- [Slo] N.J.A. Sloane, *The online encyclopedia of integer sequences*, <http://oeis.org>.
- [Vla] Masha Vlasenko, *Computation of an asymptotics related to nahm’s conjecture*, Preprint 2010.
- [VZ11] Masha Vlasenko and Sander Zwegers, *Nahm’s conjecture: asymptotic computations and counterexamples*, Commun. Number Theory Phys. **5** (2011), no. 3, 617–642.
- [Zag07] Don Zagier, *The dilogarithm function*, Frontiers in number theory, physics, and geometry. II, Springer, Berlin, 2007, pp. 3–65.
- [Zei91] Doron Zeilberger, *The method of creative telescoping*, J. Symbolic Comput. **11** (1991), no. 3, 195–204.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160, USA
<http://www.math.gatech.edu/~stavros>

E-mail address: stavros@math.gatech.edu

SECTION DE MATHÉMATIQUES, UNIVERSITÉ DE GENÈVE, 2-4 RUE DU LIÈVRE, CASE POSTALE 64,
1211 GENÈVE 4, SWITZERLAND

<http://www.unige.ch/math/folks/kashaev>

E-mail address: Rinat.Kashaev@unige.ch