

April 21, 2013

## EVALUATION OF ONE EXOTIC FURDUI TYPE SERIES

**Khristo N. Boyadzhiev**

Department of Mathematics and Statistics, Ohio Northern University, Ada, OH 45810  
[k-boyadzhiev@onu.edu](mailto:k-boyadzhiev@onu.edu)

**Abstract.** In this note we present a closed form evaluation of one interesting alternating series whose value is a combination of three mathematical constants,  $\pi$ ,  $\ln 2$ , and  $\zeta(3)$ . The series is expressed as a symmetrical double integral and this integral is evaluated in several steps.

**Mathematics Subject Classification 2010:** 11B83, 40A25.

**Keywords:** Mathematical constants, dilogarithm, trilogarithm, harmonic numbers, double integrals.

### 1. Introduction

The following two series were evaluated recently

$$\sigma_1 = \sum_{n=0}^{\infty} (-1)^n \left( \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{n+k} \right)^2 = \frac{\pi^2}{24},$$

(problem 11682 in the American Mathematical Monthly [4]) and the more challenging one

$$\sigma_2 = \sum_{n=1}^{\infty} (-1)^n \left( \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{2n+k} \right)^2 = \frac{G}{2} + \frac{\pi^2}{48} - \frac{7}{8}(\ln 2)^2 - \frac{\pi}{8} \ln 2,$$

where  $G$  is the Catalan constant. The evaluation of  $\sigma_2$  was published in [2]. This series is remarkable with its connection to the three important constants  $\pi$ ,  $\ln 2$ , and  $G$ .

In this note we shall evaluate another interesting series

$$\sigma = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \left( \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{n+k} \right)^2 = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \left( \ln 2 - \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \right)^2.$$

Namely, we have the following.

**Proposition.** The series  $\sigma$  converges and its value is

$$\sigma = \frac{\pi^2}{12} \ln 2 + \frac{1}{3} (\ln 2)^3 - \frac{1}{2} \zeta(3) \approx 0.08007,$$

involving the three constants  $\pi$ ,  $\ln 2$ , and  $\zeta(3)$ .

The above series can be expressed as double integrals. Thus

$$\int_0^1 \int_0^1 \frac{dx dy}{(1+xy)(1+x)(1+y)} = \sigma_1 = \frac{\pi^2}{24},$$

$$\int_0^1 \int_0^1 \frac{x^2 y^2 dx dy}{(1+x^2 y^2)(1+x)(1+y)} = -\sigma_2 = \frac{7}{8} (\ln 2)^2 + \frac{\pi}{8} \ln 2 - \frac{G}{2} - \frac{\pi^2}{48},$$

$$\int_0^1 \int_0^1 \frac{\ln(1+xy)}{(1+x)(1+y)} dx dy = \sigma = \frac{\pi^2}{12} \ln 2 + \frac{1}{3} (\ln 2)^3 - \frac{1}{2} \zeta(3).$$

For related results see [3].

## 2. Evaluation of the series

Using the representation

$$\frac{1}{n+k} = \int_0^1 x^{n+k-1} dx,$$

we write

$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{n+k} = \int_0^1 x^n \left\{ \sum_{k=1}^{\infty} (-1)^{k-1} x^{k-1} \right\} dx = \int_0^1 \frac{x^n}{1+x} dx.$$

Then

$$\begin{aligned} \sigma &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \left( \int_0^1 \frac{x^n dx}{1+x} \right)^2 = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \left( \int_0^1 \frac{x^n dx}{1+x} \right) \left( \int_0^1 \frac{y^n dy}{1+y} \right) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \int_0^1 \int_0^1 \frac{x^n y^n dx dy}{(1+x)(1+y)} \\ &= \int_0^1 \int_0^1 \left\{ \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (xy)^n}{n} \right\} \frac{dx dy}{(1+x)(1+y)} = \int_0^1 \int_0^1 \frac{\ln(1+xy)}{(1+x)(1+y)} dx dy \\ &= \int_0^1 \frac{1}{1+x} \left\{ \int_0^1 \frac{\ln(1+xy)}{1+y} dy \right\} dx. \end{aligned}$$

Now set

$$H_n^- = 1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{(-1)^{n-1}}{n}.$$

We shall evaluate the inside integral by using series expansion. For  $|x| < 1$  we write

$$\begin{aligned} \int_0^1 \frac{\ln(1+xy)}{1+y} dy &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n} \left\{ \int_0^1 \frac{y^n}{1+y} dy \right\} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n} \left\{ \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{n+k} \right\} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n} (\ln 2 - H_n^-) (-1)^n \end{aligned}$$

and so,

$$\int_0^1 \frac{\ln(1+xy)}{1+y} dy = \ln 2 \ln(1-x) + \sum_{n=1}^{\infty} \frac{x^n}{n} H_n^- . \quad (1)$$

**Remark 1.** At this point it is good to notice that a slight modification of entry (5.5.27) in [5] yields

$$\ln 2 \ln(1-x) + \sum_{n=1}^{\infty} \frac{x^n}{n} H_n^- = \text{Li}_2\left(\frac{1-x}{2}\right) - \text{Li}_2(-x) + \frac{1}{2}(\ln 2)^2 - \frac{\pi^2}{12},$$

and as a result from (1) we have the closed form evaluation

$$\int_0^1 \frac{\ln(1+xy)}{1+y} dy = \text{Li}_2\left(\frac{1-x}{2}\right) - \text{Li}_2(-x) + \frac{1}{2}(\ln 2)^2 - \frac{\pi^2}{12} \quad (2)$$

true on the entire unit disk  $|x| \leq 1$ . Integrals of this type have been considered in [1] and in Levin's book [6, A.3.1 (2)], but the form given here is simpler and more appropriate for our work.

Now we can write

$$\begin{aligned} \sigma &= \int_0^1 \frac{1}{1+x} \left\{ \int_0^1 \frac{\ln(1+xy)}{1+y} dy \right\} dx \\ &= \int_0^1 \frac{1}{1+x} \text{Li}_2\left(\frac{1-x}{2}\right) dx - \int_0^1 \frac{1}{1+x} \text{Li}_2(-x) dx + \left\{ \frac{1}{2}(\ln 2)^2 - \frac{\pi^2}{12} \right\} \int_0^1 \frac{dx}{1+x} \end{aligned}$$

and we shall evaluate these integrals one by one. Making first the substitution  $t = \frac{1}{2}(1-x)$  and then integrating by parts,

$$J_1 = \int_0^1 \frac{1}{1+x} \operatorname{Li}_2\left(\frac{1-x}{2}\right) dx = \int_0^{1/2} \frac{\operatorname{Li}_2(t)}{1-t} dt = \ln 2 \operatorname{Li}_2\left(\frac{1}{2}\right) - \int_0^{1/2} \frac{\ln^2(1-t)}{t} dt.$$

We now use the antiderivative

$$\int_0^t \frac{\ln^2(1-t)}{t} dt = \ln(t) \ln^2(1-t) + 2 \ln(1-t) \operatorname{Li}_2(1-t) - 2 \operatorname{Li}_3(1-t) + 2\zeta(3) \quad (3)$$

(see [5, (6.27), p.159]) and setting  $t = 1/2$  we find

$$\int_0^{1/2} \frac{\ln^2(1-t)}{t} dt = -(\ln 2)^3 - 2 \ln 2 \operatorname{Li}_2\left(\frac{1}{2}\right) - 2 \operatorname{Li}_3\left(\frac{1}{2}\right) + 2\zeta(3).$$

It is well known that

$$\operatorname{Li}_2\left(\frac{1}{2}\right) = \frac{\pi^2}{12} - \frac{(\log 2)^2}{2},$$

$$\operatorname{Li}_3\left(\frac{1}{2}\right) = \frac{7}{8} \zeta(3) - \frac{\pi^2 \log 2}{12} + \frac{(\log 2)^3}{6},$$

and we compute

$$J_1 = \frac{\pi^2}{12} \ln 2 - \frac{1}{6} (\ln 2)^3 - \frac{1}{4} \zeta(3).$$

Next we continue with

$$J_2 = -\int_0^1 \frac{\operatorname{Li}_2(-x) dx}{1+x} = \int_0^{-1} \frac{\operatorname{Li}_2(t)}{1-t} dt = -\ln 2 \operatorname{Li}_2(-1) - \int_0^{-1} \frac{\ln^2(1-t)}{t} dt.$$

Here  $\operatorname{Li}_2(-1) = \frac{-\pi^2}{12}$  is easy, but for the integral on the RHS we cannot use the antiderivative in (3), because it brings to the expressions  $\ln(-1)$ ,  $\operatorname{Li}_2(2)$ ,  $\operatorname{Li}_3(2)$  which need to be defined additionally. Instead of discussing these values we shall follow a different approach. Let

$$H_n^{(2)} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2}.$$

Expanding  $(1-t)^{-1}$  in power series and using Cauchy's rule for the product of two power series we obtain

$$\frac{\text{Li}_2(t)}{1-t} = \sum_{n=1}^{\infty} H_n^{(2)} t^n,$$

and therefore,

$$J_2 = \int_0^{-1} \frac{\text{Li}_2(t)}{1-t} dt = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n+1} H_n^{(2)} = \frac{\pi^2}{12} \ln 2 - \frac{1}{4} \zeta(3).$$

The evaluation of this series is given in the next section.

Finally we have

$$\sigma = J_1 + J_2 + \left( \frac{1}{2} (\ln 2)^2 - \frac{\pi^2}{12} \right) \ln 2$$

and after simplifying we come to the desired result

$$\sigma = \frac{\pi^2}{12} \ln 2 + \frac{1}{3} (\ln 2)^3 - \frac{1}{2} \zeta(3).$$

**Remark 2.** As byproducts from (1) and (2) with  $x = -1$  we have

$$\int_0^1 \frac{\ln(1-x)}{1+x} dx = \frac{1}{2} (\ln 2)^2 - \frac{\pi^2}{12},$$

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} H_n^- = \frac{1}{2} (\ln 2)^2 + \frac{\pi^2}{12}.$$

### 3. Additional results

**Lemma.** For  $|t| \leq 1$ ,  $t \neq 1$ , we have

$$\sum_{n=1}^{\infty} H_n^{(2)} \frac{t^n}{n} + 2 \sum_{n=1}^{\infty} H_n \frac{t^n}{n^2} = 3\text{Li}_3(t) - \text{Li}_2(t) \ln(1-t). \quad (4)$$

*Proof.* We already know that

$$\sum_{n=1}^{\infty} H_n^{(2)} t^n = \frac{\text{Li}_2(t)}{1-t}.$$

From this, dividing by  $t$  and integrating,

$$\sum_{n=1}^{\infty} H_n^{(2)} \frac{t^n}{n} = \int_0^t \frac{\text{Li}_2(t)}{t(1-t)} dt = \int_0^t \frac{\text{Li}_2(t)}{t} dt + \int_0^t \frac{\text{Li}_2(t)}{1-t} dt .$$

Clearly,

$$\int_0^t \frac{\text{Li}_2(t)}{t} dt = \text{Li}_3(t) ,$$

and integrating by parts,

$$\int_0^t \frac{\text{Li}_2(t)}{1-t} dt = -\ln(1-t)\text{Li}_2(t) - \int_0^t \frac{\ln^2(1-t)}{t} dt ,$$

so that

$$\sum_{n=1}^{\infty} H_n^{(2)} \frac{t^n}{n} = \text{Li}_3(t) - \ln(1-t)\text{Li}_2(t) - \int_0^t \frac{\ln^2(1-t)}{t} dt \quad (5)$$

At the same time

$$\sum_{n=1}^{\infty} H_n t^n = \frac{-\ln(1-t)}{1-t} .$$

Dividing this by  $t$  and integrating we find

$$\sum_{n=1}^{\infty} H_n \frac{t^n}{n} = \int_0^t \frac{-\ln(1-t)}{t(1-t)} dt = \int_0^t \frac{-\ln(1-t)}{1-t} dt + \int_0^t \frac{-\ln(1-t)}{t} dt ,$$

that is,

$$\sum_{n=1}^{\infty} H_n \frac{t^n}{n} = \frac{1}{2} \ln^2(1-t) + \text{Li}_2(t) .$$

Multiplying this by 2 and then dividing by  $t$  we find

$$2 \sum_{n=1}^{\infty} H_n \frac{t^{n-1}}{n} = \frac{\ln^2(1-t)}{t} + 2 \sum_{n=1}^{\infty} \frac{t^{n-1}}{n^2} ,$$

which integrated becomes

$$2 \sum_{n=1}^{\infty} H_n \frac{t^n}{n^2} = \int_0^t \frac{\ln^2(1-t)}{t} dt + 2\text{Li}_3(t) . \quad (6)$$

Adding equations (5) and (6) gives the desired result.

We continue now with the evaluation of the series

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n+1} H_n^{(2)} = \frac{\pi^2}{12} \ln 2 - \frac{1}{4} \zeta(3).$$

Setting  $t = -1$  in (4) yields

$$\sum_{n=1}^{\infty} H_n^{(2)} \frac{(-1)^{n-1}}{n} + 2 \sum_{n=1}^{\infty} H_n \frac{(-1)^{n-1}}{n^2} = 3\text{Li}_3(-1) - \text{Li}_2(-1) \ln 2 = \frac{9}{4} \zeta(3) - \frac{\pi^2}{12} \ln 2.$$

The following evaluation is due to Sitaramachandra Rao [7]

$$\sum_{n=1}^{\infty} H_n \frac{(-1)^{n-1}}{n^2} = \frac{5}{8} \zeta(3).$$

Therefore, the above equation implies

$$\sum_{n=1}^{\infty} H_n^{(2)} \frac{(-1)^{n-1}}{n} = \zeta(3) - \frac{\pi^2}{12} \log 2.$$

From here we easily compute

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n+1} H_n^{(2)} = - \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} H_n^{(2)} - \text{Li}_3(-1) = \frac{\pi^2}{12} \ln 2 - \frac{1}{4} \zeta(3)$$

which is the value of  $J_2$  needed in section 2.

## References

- [1] **Tewodoros Amdeberhan, Victor H. Moll, and Armin Straub**, Closed-form evaluation of integrals appearing in positronium decay, *J. Math. Physics*, 50, 103528 (2009).
- [2] **Khristo N. Boyadzhiev**, On a series of Furdui and Qin and some related integrals, (Solution of Problem H-691 (The Fibonacci Quarterly, 50 (1) 2012) with corrections and details.) [arXiv:1203.4618v3](https://arxiv.org/abs/1203.4618v3) [math.NT]
- [3] **Khristo N. Boyadzhiev**, Power series with skew-harmonic numbers, dilogarithms, and double integrals, *Tatra Mountain Math. Publications*, 56, (2013), 93–108.
- [4] **Ovidiu Furdui**, Problem 11682, *Amer. Math. Monthly*, 119 (10) (2012). p.881.
- [5] **Eldon R. Hansen**, *A Table of Series and Products*, Prentice Hall, 1975.
- [6] **Leonard Lewin**, *Polylogarithms and associated functions*, North Holland, 1981.
- [7] **R. Sitaramachandra Rao**, A formula of S. Ramanujan, *J. Number Theory*, 25 (1987), 1-19.