# MINIMAL ZERO-SUM SEQUENCE OF LENGTH FIVE OVER FINITE CYCLIC GROUPS OF PRIME POWER ORDER

LI-MENG XIA<sup>1</sup>, YUANLIN LI\*,<sup>2</sup>, AND JIANGTAO PENG<sup>3</sup>

<sup>1</sup>Faculty of Science, Jiangsu University, Zhenjiang, 212013, Jiangsu Pro., P.R. China <sup>2</sup>Department of Mathematics, Brock University, St. Catharines, ON, Canada L2S 3A1 <sup>3</sup>College of Science, Civil Aviation University of China, Tianjin, 300300, P.R. China

ABSTRACT. Let G be a finite cyclic group. Every sequence S of length l over G can be written in the form  $S=(x_1g)\cdot\ldots\cdot(x_lg)$  where  $g\in G$  and  $x_1,\ldots,x_l\in[1,\operatorname{ord}(g)]$ , and the index  $\operatorname{ind}(S)$  of S is defined to be the minimum of  $(x_1+\cdots+x_l)/\operatorname{ord}(g)$  over all possible  $g\in G$  such that  $\langle g\rangle=G$ . Recently the second and the third authors determined the index of any minimal zero-sum sequence S of length S over a cyclic group of a prime order where  $S=g^2(x_2g)(x_3g)(x_4g)$ . In this paper, we determine the index of any minimal zero-sum sequence S of length S over a cyclic group of a prime power order. It is shown that if  $G=\langle g\rangle$  is a cyclic group of prime power order  $n=p^\mu$  with  $p\geq 7$  and  $p\geq 2$ , and  $p\geq 2$ , and  $p\geq 2$ , and  $p\geq 2$ , and  $p\geq 3$ , and  $p\geq 3$ , and  $p\geq 4$ , and  $p\geq 3$ , and  $p\geq 4$ , then  $p\geq 4$  if and only if  $p\geq 4$  if  $p\geq 4$ . Then  $p\geq 4$  if and only if  $p\geq 4$  if  $p\geq 4$  if  $p\geq 4$  if  $p\geq 4$  in  $p\geq 4$  if and only if  $p\geq 4$  if  $p\geq 4$  if  $p\geq 4$  if  $p\geq 4$  in p> 4 in p>

#### 1. Introduction

Throughout the paper G is assumed to be a finite cyclic group of order n written additively. Denote by  $\mathcal{F}(G)$ , the free abelian monoid with basis G and elements of  $\mathcal{F}(G)$  are called sequences over G. A sequence of length l of not necessarily distinct elements from G can be written in the form  $S = (x_1g) \cdot \ldots \cdot (x_lg)$  for some  $g \in G$ . S is called a zero-sum sequence if the sum of S is zero (i.e.  $\sum_{i=1}^{l} x_i g = 0$ ). If S is a zero-sum sequence, but no proper nontrivial subsequence of S has sum zero, then S is called a minimal zero-sum sequence. The index of a sequence S over G is defined as follows.

**Definition 1.1.** For a sequence over G

$$S = (x_1 g) \cdot \ldots \cdot (x_l g)$$
, where  $1 \leq x_1, \ldots, x_l \leq \operatorname{ord}(g)$ ,

E-mail: yli@brocku.ca (Y. Li)

.

<sup>2000</sup> Mathematics Subject Classification. Primary 11B30, 11B50, 20K01.

Key words and phrases: minimal zero-sum sequences, index of sequences.

<sup>\*</sup>Corresponding author: Yuanlin Li, Department of Mathematics, Brock University, St. Catharines, Ontario Canada L2S 3A1, Fax:(905) 378-5713;

the index of S is defined by  $\operatorname{ind}(S) = \min\{\|S\|_g \mid g \in G \text{ with } G = \langle g \rangle\}$  where

$$||S||_g = \frac{x_1 + \dots + x_l}{\operatorname{ord}(g)}.$$

Clearly, S has sum zero if and only if ind(S) is an integer. There are also slightly different definitions of the index in the literature, but they are all equivalent (see Lemma 5.1.2 in [7]).

The index of a sequence is a crucial invariant in the investigation of (minimal) zero-sum sequences (resp. of zero-sum free sequences) over cyclic groups. It was first addressed by Kleitman-Lemke (in the conjecture [9, page 344]), used as a key tool by Geroldinger ([6, page 736]), and then investigated by Gao [3] in a systematical way. Since then it has received a great deal of attention (see for example [1, 2, 4, 5, 7, 8, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21]).

A main focus of the investigation of index is to determine minimal zero-sum sequences of index 1. If S is a minimal zero-sum sequence of length |S| such that  $|S| \leq 3$  or  $|S| \geq \lfloor \frac{n}{2} \rfloor + 2$ , then  $\operatorname{ind}(S) = 1$  (see [1, 14, 20]). In contrast to that, it was shown that for each l with  $5 \leq l \leq \lfloor \frac{n}{2} \rfloor + 1$ , there is a minimal zero-sum sequence S of length |S| = l with  $\operatorname{ind}(S) \geq 2$  ([14, 20]) and that the same is true for l = 4 and  $\gcd(n, 6) \neq 1$  ([13]). In recent papers [10, 11, 18], the authors proved that  $\operatorname{ind}(S) = 1$  if |S| = 4 and  $\gcd(n, 6) = 1$  when n is a prime power or a product of two prime powers. When n is a product of at least 3 prime powers, some partial results were also obtained in [15, 16, 17]. However, the general case is still open.

It was mentioned in [12], in order to further investigate the index of a general minimal zero-sum sequence of length 4, it is helpful to determine the index of certain minimal zero-sum sequences of length 5. It is routine to check that if S is a minimal zero-sum sequence over G of length 5, then  $1 \leq \operatorname{ind}(S) \leq 2$ . Let h(S) be the maximal repetition of an element in S. In [12], the index of any minimal zero-sum sequence S of length 5 over acyclic group of a prime order with  $h(S) \geq 2$  was completely determined. In this paper, we continue the investigation on the index of minimal zero-sum sequences of length 5 over a cyclic group of prime power order. When G is a cyclic group of prime power order  $n = p^{\mu}$  with  $p \geq 7$  and  $p \geq 1$  and p

**Theorem 1.2.** Let G be a cyclic group of prime power order  $n = p^{\mu}$  with  $p \geq 7$  and  $\mu \geq 2$ , and  $S = (x_1g) \cdot (x_2g) \cdot (x_3g) \cdot (x_4g) \cdot (x_5g)$  be a minimal zero-sum sequence over G with  $gcd(n, x_1, x_2, x_3, x_4, x_5) = 1$  and  $x_1 = x_2$ , then ind(S) = 2 if and only if  $S = (mg)(mg)(m\frac{n-1}{2}g)(m\frac{n+3}{2}g)(m(n-3)g)$  where m is a positive integer such that gcd(m, n) = 1.

The paper is organized as follows. In the next section, we provide some preliminary results. In section 3, we state three main propositions and use them together with some preliminary results to give a proof for our main result. The proofs of the main propositions are given in the last section.

## 2. Preliminaries

We first prove some preliminary results which will be needed in the sequel. Let  $G = \langle g \rangle$  be a cyclic group of order n. Suppose that  $S = (x_1g) \cdot \ldots \cdot (x_lg)$ . Let  $||S||'_g = \operatorname{ord}(g)||S||_g = \sum_{i=1}^l x_i \in N_0$  and denote by  $|x|_n$  the least positive residue of x modulo n, where  $n \in \mathbb{N}$  and  $x \in \mathbb{Z}$ . Let mS denote the sequence  $(mx_1g) \cdot \ldots \cdot (mx_lg)$ . Since |g| = n, we have  $mS = (|mx_1|_ng) \cdot \ldots \cdot (|mx_l|_ng)$ . We note that if  $\gcd(n,m) = 1$ , then the multiplication by m is a group automorphism of G and hence  $\operatorname{ind}(S) = \operatorname{ind}(mS)$ . Two sequences S and S' are called equivalent, denoted by  $S \sim S'$ , if S = mS' for some m with  $\gcd(n,m) = 1$ . Clearly, if  $S \sim S'$ , then  $\operatorname{ind}(S) = \operatorname{ind}(S')$ . For all real numbers a < b, define  $[a,b] = \{k \in \mathbb{Z} | a \leq k \leq b\}$ .

From now on we always assume that  $G = \langle g \rangle$  is a cyclic group of order  $n = p^{\mu}$  with  $p \geq 7$  and  $\mu \geq 2$ , and  $S = (x_1g) \cdot \ldots \cdot (x_lg)$  is a minimal zero-sum sequence over G. We remark that if  $\gcd(n, x_1, x_2, x_3, x_4, x_5) = p^w > 1$ , let  $h = p^w g$  and  $x_i' = x_i/p^w$  for  $1 \leq i \leq 5$ . Then a minimal zero-sum sequence S over G can be rewritten as follows:

$$S = (x_1'h) \cdot \ldots \cdot (x_l'h),$$

which is a minimal zero-sum sequence over  $H = \langle h \rangle$ . Note that H is a cyclic group of prime power order with |H|||G| and  $\gcd(n, x_1', x_2', x_3', x_4', x_5') = 1$ . In what follows we may always assume that  $\gcd(n, x_1, x_2, x_3, x_4, x_5) = 1$ . Let U(n) denote the unit group of  $n = p^{\mu}$ , i.e.  $U(n) = \{m | 1 \le m \le n - 1, \gcd(m, n) = 1\}$ . It is well known that  $1 + t\alpha \in U(n)$  for any  $1 < \alpha = p^{\lambda} < n$ .

**Lemma 2.1.** Let  $n = p^{\mu}$  with  $\mu \ge 2$  and  $\alpha = \frac{n}{p}$ . If v is an integer such that  $1 \le v \le n-1$  and  $\gcd(v,p) = 1$ , then there exists  $y = 1 + t\alpha \in U(n)$  such that  $|vy|_n < \frac{n}{p}$ .

*Proof.* Note that  $y = 1 + t\alpha \in U(n)$  for any  $0 \le t \le p - 1$ . There exists a  $y_1 = 1 + t\alpha$  such that  $vy_1 < n$ , but vy > n where  $y = 1 + (t+1)\alpha$ . Thus  $|vy|_n < \alpha = \frac{n}{p}$ .

The following proposition is a generalization of [12, Proposition 2.2].

**Proposition 2.2.** Let  $G = \langle g \rangle$  be a cyclic group of order  $n = p^{\mu}$  with  $p \geq 5$ . If  $S = g^2 \cdot (\frac{p^{\mu}-1}{2}g) \cdot (\frac{p^{\mu}+3}{2}g) \cdot ((p^{\mu}-3)g) \in \mathcal{F}(G)$ , then  $\operatorname{ind}(S) = 2$ .

*Proof.* This lemma was proved in [12] for n = p and the same proof works for the prime power case.

**Proposition 2.3.** Let G be a cyclic group of prime power order  $n = p^{\mu}$  with  $p \geq 5$  and  $\mu \geq 2$ , and  $S \in \mathcal{F}(G)$  be a minimal zero-sum sequence of length 5. If  $h(S) \geq 3$ , then  $\operatorname{ind}(S) = 1$ .

*Proof.* If  $h(S) \ge 4$ , then we may assume that  $x_1 = x_2 = x_3 = x_4 = \gcd(n, x_1)$ . So we have that  $||S||_g' = \sum_{i=1}^5 x_i = 4x_1 + x_5 < 2n$ . Therefore  $\operatorname{ind}(S) = 1$ .

If h(S) = 3, then we may assume that  $x_1 = x_2 = x_3 = \gcd(n, x_1)$ . If  $x_1 = 1$ , then  $x_4, x_5 \le n-4$  and  $||S||'_g = \sum_{i=1}^5 x_i \le 3+2(n-4) < 2n$ , so we infer that  $||S||'_g = \sum_{i=1}^5 x_i = n$  and  $\inf(S) = 1$ .

If  $x_1 = p^s > 1$ , then  $\gcd(n, x_4) = 1$ . Let  $\alpha = p^{\mu - 1}$ . Then by Lemma 2.1 there exists  $y = 1 + t\alpha \in U(n)$  such that  $|yx_4|_n < \alpha$ . Since  $x_1 \le \alpha$  and y fixes  $x_1$  (i.e.  $|yx_1|_n = |x_1|_n$ ), we have  $||yS||_g' = |yx_1|_n + \dots + |yx_5|_n < 4\alpha + x_5 < 2n$ , so  $\operatorname{ind}(S) = 1$  and we are done.  $\square$ 

By using a computer search, we were able to verify the following proposition.

**Proposition 2.4.** Let  $G = \langle g \rangle$  be a cyclic group of prime power order  $n = p^{\mu}$  with  $p \geq 5$  and  $\mu \geq 2$ , and  $S = g^2 \cdot (x_3g) \cdot (x_4g) \cdot (x_5g)$  be a minimal zero-sum sequence of length 5. If n < 289, then  $\operatorname{ind}(S) = 2$  if and only if  $S = g^2 \cdot (\frac{n-1}{2}g) \cdot (\frac{n+3}{2}g) \cdot ((n-3)g)$ .

## 3. Main Propositions

In this section we present three main propositions and then use them together with some preliminary results to provide a proof for our main theorem. In terms of Proposition 2.3, from now on we may always assume that h(S) = 2 and  $x_1 = x_2 = \gcd(n, x_1)$ . The following proposition takes care of the case when  $x_1 > 1$ .

**Proposition 3.1.** Let  $G = \langle g \rangle$  be a cyclic group of prime power order  $n = p^{\mu}$  with  $p \geq 7$  and  $\mu \geq 2$ , and  $S = (x_1g) \cdot (x_2g) \cdot (x_3g) \cdot (x_4g) \cdot (x_5g)$  be a minimal zero-sum sequence of length 5. If  $x_1 (= x_2 = \gcd(n, x_1)) > 1$ , then  $\operatorname{ind}(S) = 1$ .

Next we assume that  $x_1 = x_2 = \gcd(n, x_1) = 1$  and  $S = g^2 \cdot (x_3 g) \cdot (x_4 g) \cdot (x_5 g)$  such that  $2 + x_3 + x_4 + x_5 = 2n$ . The following two propositions are crucial to prove the main theorem.

**Proposition 3.2.** Let G be a cyclic group of order n with  $n=p^{\mu} \geq 289, p \geq 7$  and  $\mu \geq 2$ . Let  $S=g^2 \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g)$  be a minimal zero-sum sequence with  $4 \leq a \leq b < c < \frac{n}{2}$  and c=b+a-2. Then  $\operatorname{ind}(S)=1$ .

**Proposition 3.3.** Let G be a cyclic group of order n with  $n=p^{\mu} \geq 289, p \geq 7$  and  $\mu \geq 2$ . Let  $S=g^2 \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g)$  be a minimal zero-sum sequence with  $3=a \leq b < b+1=c < \frac{n}{2}$ . Then  $\operatorname{ind}(S)=2$  if and only if  $c=\frac{n-1}{2}$ .

We are now ready to present a proof for our main result.

# Proof of Theorem 1.2.

In terms of Proposition 2.3, we may assume that h(S) = 2 and  $x_1 = x_2 = \gcd(n, x_1)$ . If  $x_1 > 1$ , then the result follows from Proposition 3.1. Next we assume  $x_1 = x_2 = \gcd(n, x_1) = 1$  and  $S = g^2 \cdot (x_3 g) \cdot (x_4 g) \cdot (x_5 g)$ . As discussed in [12], we may assume that  $S = g^2 \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g)$  with  $3 \le a \le b < c < \frac{n}{2}$  and c = a + b - 2, for otherwise, it is easy to show that  $\operatorname{ind}(S) = 1$ . If n < 289, the result follows from Proposition 2.4. So we may assume that  $n \ge 289$ . Now the result follows immediately from Propositions 3.2 and 3.3, and we are done.

## 4. Proofs For Main Propositions

We now give the proofs for the three crucial propositions stated in previous section. The proofs will be presented in the following three subsections.

## 4.1. Proof of Proposition 3.1.

Let  $x_1 = p^s > 1$  and  $\alpha' = \frac{n}{p^s} = \frac{n}{x_1}$ . We first assume that  $gcd(n, x_i) > 1$  for some i = 3, 4, 5, say  $gcd(n, x_3) > 1$ . We divide the proof into 3 cases.

Case 1. If  $x_1 > \gcd(n, x_3) > 1$ , let  $x_3' = \frac{x_3}{\gcd(n, x_3)}$ ,  $n' = \frac{n}{\gcd(n, x_3)}$  and  $\alpha' = \frac{n'}{p}$ . Then by Lemma 2.1, we may find a  $y = 1 + t\alpha' \in U(n')$  such that  $|yx_3'|_{n'} < \alpha'$ . Thus  $|yx_3|_n = \gcd(n, x_3)|yx_3'|_{n'} < \gcd(n, x_3)\alpha' = \frac{n}{p}$ . Since  $x_1 > \gcd(n, x_3)$ , y fixes  $x_1 = x_2$ , so by multiplying S with y, we may assume that  $x_1, x_2, x_3 \leq \frac{n}{p}$ . Since  $\gcd(n, x_1, x_2, x_3, x_4, x_5) = 1$ , we have  $\gcd(n, x_4) = 1$ . Thus by Lemma 2.1, there exists  $y' = 1 + t\frac{n}{p} \in U(n)$  such that  $|y'x_4|_n < \frac{n}{p}$ . Since y' fixes  $x_1, x_2$  and  $x_3$ , again we may assume that  $x_i \leq \frac{n}{p}$  for i = 1, 2, 3, 4. Thus  $||S||_g' = \sum_{i=1}^5 x_i \leq \frac{4n}{p} + x_5 < 2n$ , so  $\operatorname{ind}(S) = 1$  and we are done.

Case 2. If  $x_1 < \gcd(n, x_3)$ , then there exists  $u \in U(n)$  such that  $|ux_3|_n = \gcd(n, x_3)$ . So we may assume that  $x_3 = \gcd(n, x_3)$ . Note that  $x_1$  may not be  $\gcd(n, x_1)$  any longer. However, as in Case 1 we may find  $y = 1 + t\alpha'$ , where  $\alpha' = \frac{n}{\gcd(n, x_3)}$ , such that  $|yx_1|_n < \alpha' \le \frac{n}{p}$ . Since y fixes  $x_3$ , we may assume that  $x_1, x_2, x_3 \le \frac{n}{p}$ . Since  $\gcd(n, x_4) = 1$ , there exists  $y' = 1 + t\frac{n}{p} \in U(n)$  such that  $|y'x_4|_n < \frac{n}{p}$ . Since y' fixes y' fixes y' and y' so we may assume that y' and y' fixes y' fixes

Case 3. If  $x_1 = \gcd(n, x_3)$ , let  $x_3 = n - kx_1 = n - w$ . Since S is a minimal zero-sum sequence, we have  $k \geq 3$ . (Otherwise,  $x_1 + x_3$  or  $x_1 + x_2 + x_3$  has zero-sum.) If  $w < \frac{n}{2}$ , then 2w < n and  $|2x_3|_n = n - 2w$  and  $|2x_1|_n = 2x_1$ . We may replace S by the following equivalent sequence:

$$S' = (2x_1q) \cdot (2x_2q) \cdot ((n-w')q) \cdot (|2x_4|_nq) \cdot (|2x_5|_nq).$$

By repeating this process, we may assume that  $w > \frac{n}{2}$ . As before, since  $\gcd(n, x_4) = 1$ , we may assume that  $x_4 < \frac{n}{p}$ . Since  $k \ge 3, p > 6$  and kp - 2p - 2k = (k-2)(p-2) - 4 > 0,  $\frac{k-2}{k}w - x_4 > \frac{k-2}{k}\frac{n}{2} - \frac{n}{p} = \frac{n}{2kp}(kp - 2p - 2k) > 0$ , so we have

$$||S||'_g = \sum_{i=1}^5 x_i = n - (k-2)x_1 + x_4 + x_5$$
$$= n + x_5 - (\frac{k-2}{k}w - x_4) < 2n.$$

Therefore, ind(S) = 1 and we are done.

Next we assume that  $\gcd(n, x_i) = 1$  for all i = 3, 4, 5. Let  $x_1 = p^s > 1$  and  $\alpha' = \frac{n}{p^s}$ . As before, we may use a unit  $1 + t\alpha'(0 \le t \le x_1 - 1)$  to move  $x_3$  so that  $x_3 < \alpha'$ . Thus we may assume that  $x_3 < \alpha'$ . Let  $\beta = \frac{n}{p} \ge \max(x_1, \alpha')$ . Then  $y_1 = 1 + t_1 x_3^{-1} \beta \in U(n)$  and  $y_2 = 1 + t_2 x_3^{-1} \beta \in U(n)$  for  $t_1 \ne t_2 \in [0, p - 1]$  are distinct units. Note that every

such y fixes  $x_1$ . Note also that  $|(1 + tx_3^{-1}\beta)x_i|_n = |x_i' + (n - t'\beta)|_n \le x_i' + (n - t'\beta)$  where  $1 \le t' \le p$  and  $x_i' = x_i \mod \beta$ . If  $|(1 + t_1x_3^{-1}\beta)x_i|_n \ne |(1 + t_2x_3^{-1}\beta)x_i|_n$ , then  $|x_i' + (n - t_1'\beta)|_n \ne |x_i' + (n - t_2'\beta)|_n$ , implying  $t_1' \ne t_2'$ . Thus  $\sum_{t=0}^6 |(1 + tx_3^{-1}\beta)x_i|_n = \sum_{t=0}^6 |x_i' + (n - t'\beta)|_n \le 7x_i' + \sum_{t'=1}^7 (n - t'\beta) = 7x_i' + 7n - 28\beta$ . We now compute the following sum

$$\sum_{i=1}^{5} \sum_{t=0}^{6} |(1+tx_3^{-1}\beta)x_i|_n = 7x_1 + 7x_2 + \sum_{t=0}^{6} (x_3 + t\beta) + \sum_{i=4,5} \sum_{t=0}^{6} |(1+tx_3^{-1}\beta)x_i|_n$$

$$\leq 14x_1 + 7x_3 + 21\beta + 7x_4' + 7x_5' + 2\sum_{t=0}^{6} (n - (1+t)\beta)$$

$$= 7(2x_1 + x_3 + x_4' + x_5') + 14n - 35\beta.$$

Since  $2x_1 + x_3 + x'_4 + x'_5 < 5\beta$ , we have

$$\sum_{i=1}^{5} \sum_{t=0}^{6} |(1+tx_3^{-1}\beta)x_i|_n < 14n + 35\beta - 35\beta = 14n.$$

Then there must exist  $t \in [0,6]$  such that

$$\sum_{i=1}^{5} |(1+tx_3^{-1}\beta)x_i|_n < 2n.$$

So

$$\sum_{i=1}^{5} |(1+tx_3^{-1}\beta)x_i|_n = n,$$

and thus ind(S) = 1. This completes the proof.

# 4.2. Proof of Proposition 3.2.

In this subsection we will prove Proposition 3.2 through the following lemmas. Let  $s = \lfloor \frac{b}{a} \rfloor$ . Recall that from now on we always assume that  $n \geq 289$ .

**Lemma 4.1.** Let  $S = g^2 \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g)$  be the same sequence described in Proposition 3.2. If there exist integers M and k such that  $\gcd(n,M) = 1$ ,  $\frac{kn}{c} < M < \frac{kn}{b}$  and Ma < n, then  $\operatorname{ind}(S) = 1$ .

*Proof.* Note that

$$||MS||'_{g} = M + M + |Mc|_{n} + |M(n-b)|_{n} + |M(n-a)|_{n}$$

$$\leq M + M + Mc - kn + (kn - Mb) + (n - Ma)$$

$$= n + M(1 + 1 + c - b - a) = n.$$

We conclude that ind(S) = 1.

**Lemma 4.2.** If  $S = g^2 \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g)$  is a minimal zero-sum sequence such that  $3 \le a \le b < c < \frac{n}{2}$ , then  $\operatorname{ind}(S) = 1$  provided that one of the following two conditions holds:

(1) 
$$\frac{s(a-2)n}{bc} > 2$$
.

(2) 
$$\frac{(s-1)(a-2)n}{bc} > 1$$
 and  $\frac{n}{c} .$ 

*Proof.* (1). If  $\frac{s(a-2)n}{bc} > 2$ , then there exist two integers  $M_1, M_2 \in [\frac{sn}{c}, \frac{sn}{b}]$ , so one of them, say M, is co-prime to n. Since Ma < n, by Lemma 4.1 we have  $\operatorname{ind}(S) = 1$ .

(2). If  $\frac{(s-1)(a-2)n}{bc} > 1$ , then we can find two integers  $M_1$  and  $M_2$  such that  $M_1 \in \left[\frac{(s-1)n}{c}, \frac{(s-1)n}{b}\right]$  and  $M_2 \in \left[\frac{sn}{c}, \frac{sn}{b}\right]$ . We may assume that  $\frac{(s-1)n}{b} - M_1 \le 1$  and  $M_2 - \frac{sn}{c} \le 1$ ; for otherwise, by the proof in (1) we infer ind(S) = 1. Note that

$$M_2 - M_1 = (M_2 - \frac{sn}{c}) + (\frac{sn}{c} - \frac{(s-1)n}{b}) + (\frac{(s-1)n}{b} - M_1) \le 2 + \frac{n}{c} < p.$$

Thus one of  $M_1$  and  $M_2$  is co-prime to n, so by Lemma 4.1 we have  $\operatorname{ind}(S) = 1$ .  $\square$ 

Lemma 4.3. If the sequence

$$S = g^2 \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g),$$

is a minimal zero-sum sequence such that  $6 \le a \le b < c < \frac{n}{2}$  and  $s \ge 4$ , then  $\operatorname{ind}(S) = 1$ .

*Proof.* We divide the proof into the following 3 cases.

Case 1.  $\frac{n}{c} \ge 4$ . Since  $\frac{s(a-2)n}{bc} > \frac{s(a-2)n}{(s+1)ac} \ge (\frac{4}{5})(\frac{4}{6})4 > 2$ , by Lemma 4.2 we inferind(S) = 1.

Case 2.  $3 \le \frac{n}{c} < 4$ . Then  $\frac{(s-1)(a-2)n}{bc} > \frac{(s-1)(a-2)n}{(s+1)ac} \ge (\frac{3}{5})(\frac{4}{6})3 > 1$ . Since  $\frac{n}{c} < 4 < p-2$ , it follows from Lemma 4.2 that  $\operatorname{ind}(S) = 1$ .

Case 3.  $2 < \frac{n}{c} < 3$ . If  $\frac{n}{c} < 3 < \frac{n}{b}$ , then 3a < n and thus by Lemma 4.1  $\operatorname{ind}(S) = 1$ . So we may assume that  $2 < \frac{n}{c} < \frac{n}{b} < 3$ . If  $s \ge 7$ , then  $\frac{(s-1)(a-2)n}{bc} > \frac{(s-1)(a-2)n}{(s+1)ac} \ge (\frac{6}{8})(\frac{4}{6})2 = 1$ . By Lemma 4.2  $\operatorname{ind}(S) = 1$ . If  $4 \le s \le 6$ ,  $n < 3b < 3(s+1)a \le 21a$ , so  $a \ge 14$ . Again,  $\frac{(s-1)(a-2)n}{bc} > \frac{2(s-1)(a-2)}{(s+1)a} \ge (\frac{3}{5})(\frac{12}{14})2 > 1$ . By Lemma 4.2 we infer  $\operatorname{ind}(S) = 1$ .

**Lemma 4.4.** If  $S = (g) \cdot (g) \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g)$  is a minimal zero-sum sequence such that  $6 \le a \le b < c < \frac{n}{2}$  and  $s \le 3$ , then  $\operatorname{ind}(S) = 1$ .

*Proof.* We divide the proof into two cases.

Case 1.  $\frac{n}{c} \ge 5$ . If  $s \ge 2$ , then  $\frac{s(a-2)n}{bc} > \frac{s(a-2)n}{(s+1)ac} > (\frac{2}{3})(\frac{4}{6})5 > 2$ , so by Lemma 4.2, ind(S) = 1.

Next assume that s=1. If  $\frac{n}{c} \geq 6$ , then  $\frac{(a-2)n}{bc} > \frac{(a-2)n}{2ac} \geq 2$ , so again we infer  $\operatorname{ind}(S) = 1$ . If  $5 \leq \frac{n}{c} < 6$ , then  $3a-3 \geq c > \frac{n}{6}$ , so  $a \geq 18$ . Thus  $\frac{(a-2)n}{bc} > \frac{(a-2)n}{2ac} \geq (\frac{16}{36})5 > 2$ , so  $\operatorname{ind}(S) = 1$ .

Case 2.  $\frac{n}{c} < 5$ . We may assume that  $m < \frac{n}{c} < \frac{n}{b} < m + 1$  (\*) with  $m \in [2, 4]$  (for otherwise, it is easy to show that  $\operatorname{ind}(S) = 1$ ).

**Subcase 2.1.** s = 1. If  $m \ge 3$ , we have

$$\frac{n}{b} - \frac{n}{c} = \frac{(a-2)n}{bc} > \frac{3(a-2)}{2a} \ge 1,$$

giving a contradiction to (\*).

Next assume m=2. If  $b \leq 2a-4$ , then  $\frac{n}{b} - \frac{n}{c} = \frac{(a-2)n}{bc} \geq \frac{n}{2c} > 1$ , giving a contradiction. If  $b \geq 2a-3$ , then  $n \geq 2c+1 = 6a-9$ . Since  $\frac{2n}{b} - \frac{2n}{c} = \frac{2(a-2)n}{bc} \geq \frac{4(a-2)}{2a-1} > 1$ , we have  $\frac{2n}{c} < 5 < \frac{2n}{b}$ . Note that  $3a-3 \geq c > \frac{n}{3}$ , so  $a \geq 34$ . Thus 5a < 6a-9 < n, so by Lemma 4.1 ind(S) = 1.

**Subcase 2.2.** s=2. If m=2, then  $2<\frac{n}{c}<\frac{n}{b}<3$  and thus  $a>\frac{n}{9}>32$ . Since  $\frac{2n}{b}-\frac{2n}{c}=\frac{2(a-2)n}{bc}\geq\frac{4(a-2)}{3a-1}>1$ , we have  $\frac{2n}{c}<5<\frac{2n}{b}$ , implying  $\operatorname{ind}(S)=1$ .

Next assume m=3. Thus  $a>\frac{b}{3}>\frac{n}{12}>24$ . So  $\frac{2n}{b}-\frac{2n}{c}>\frac{2(a-2)n}{3ac}>\frac{(2\times 22)n}{(3\times 24)c}=\frac{11n}{18c}$ . If  $\frac{n}{c}>\frac{36}{11}$ , then  $\frac{2(a-2)n}{bc}>2$ , implying  $\operatorname{ind}(S)=1$ . Next assume that  $\frac{n}{c}\leq\frac{36}{11}$ . Then  $\frac{3n}{c}\leq\frac{108}{11}<10$ . Since  $\frac{3n}{b}-\frac{3n}{c}\geq\frac{3(a-2)n}{3ac}>1$ . We have  $\frac{3n}{c}<10<\frac{3n}{b}$ . We next show that 10a< n and thus  $\operatorname{ind}(S)=1$  (by Lemma 4.1). Note that  $3<\frac{n}{c}<\frac{n}{b}<4$  and a>24. Then  $a\leq\frac{24}{22}(a-2)\leq\frac{24}{22}(c-b)<\frac{24}{22}(\frac{n}{3}-\frac{n}{4})=\frac{n}{11}$ , so 10a< n as required.

If m = 4, then  $a \ge 20$  and thus  $\frac{2(a-2)n}{bc} > \frac{8(a-2)}{3a} > 2$ . By Lemma 4.2 ind(S) = 1.

**Subcase 2.3.** s=3. If  $m\geq 4$ , then  $\frac{3n}{b}-\frac{3n}{c}>\frac{3(a-2)n}{4ac}>2$ . It follows from Lemma 4.2 that  $\operatorname{ind}(S)=1$ . If m=3, then  $\frac{2n}{b}-\frac{2n}{c}>\frac{2(a-2)n}{4ac}>1$ , so by Lemma 4.2  $\operatorname{ind}(S)=1$ .

Next we assume that m=2 and  $2<\frac{n}{c}<\frac{n}{b}<3$ . If  $b\leq 4a-8$ , then  $\frac{2n}{b}-\frac{2n}{c}\geq \frac{2(a-2)n}{4(a-2)c}>1$ , so  $\operatorname{ind}(S)=1$ . Now assume that  $4a-7\leq b\leq 4a-1$ . Thus  $c\geq 5a-9$ , so  $n\geq 2c+1\geq 10a-17>9a$  (as  $a>\frac{n}{12}\geq 24$ ). Since  $\frac{3n}{b}-\frac{3n}{c}>\frac{3(a-2)}{2a}>1$ , we have  $6<\frac{3n}{c}< M<\frac{3n}{b}<9$ . If M=8, since 8a< n, we have  $\operatorname{ind}(S)=1$ . So we may assume  $6<\frac{3n}{c}<7<\frac{3n}{b}<8$ . Thus  $\frac{2n}{c}<\frac{14}{3}<5$ . If  $\frac{2n}{c}<5<\frac{2n}{b}$ , since 5a< n, we have  $\operatorname{ind}(S)=1$ . Thus  $4<\frac{2n}{c}<\frac{2n}{b}<5$ . Since  $\frac{4n}{b}-\frac{4n}{c}>1$ , we have  $8<\frac{4n}{c}<9<\frac{4n}{b}<10$ . Recall that 9a< n, so by Lemma 4.1  $\operatorname{ind}(S)=1$  and we are done.

#### Lemma 4.5. If

$$S = g^{2} \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g),$$

is a minimal zero-sum sequence such that  $a \in [4,5]$ ,  $a \le b < c < \frac{n}{2}$  and  $s \le 10$ , then  $\operatorname{ind}(S) = 1$ .

*Proof.* Note that  $b \le sa + 4$  and  $c \le (s+1)a + 2$ . If  $s \le 3$ , then  $\frac{n}{c} \ge \frac{289}{(s+1)a+2} \ge \frac{289}{22} > 13$ . Thus  $\frac{s(a-2)n}{bc} > \frac{13s(a-2)}{(s+1)a} \ge \frac{13\times 2}{8} > 2$ . If  $s \ge 4$ , then  $\frac{n}{c} \ge \frac{289}{(s+1)a+2} \ge \frac{289}{57} > 5$ . Thus  $\frac{s(a-2)n}{bc} > \frac{5s(a-2)}{(s+1)a} \ge \frac{5\times 4\times 2}{5\times 4} = 2$ . It follows from Lemma 4.2 that  $\operatorname{ind}(S) = 1$ .

**Lemma 4.6.** If  $S = g^2 \cdot (cg) \cdot ((n-b)g) \cdot ((n-a)g)$  is a minimal zero-sum sequence such that  $a \in [4,5]$ ,  $a \le b < c < \frac{n}{2}$  and  $s \ge 11$ , then  $\operatorname{ind}(S) = 1$ .

*Proof.* We divide the proof into two cases according to a = 5 and a = 4.

**Case 1.** a=5. In terms of Lemma 4.2, we may assume that  $\frac{s(a-2)n}{(s+1)ac} \leq 2$ . Thus  $\frac{3sn}{5(s+1)c} \leq 2$ , so  $\frac{n}{c} \leq \frac{10(s+1)}{3s} \leq \frac{120}{33} < 4 < p-2$ . Since  $\frac{(s-1)(a-2)n}{bc} > \frac{(s-1)(a-2)n}{(s+1)ac} > \frac{10\times 3\times 2}{12\times 5} = 1$ , by Lemma 4.2 we have  $\operatorname{ind}(S) = 1$ .

Case 2. a=4. In terms of Lemma 4.2, we may assume that  $\frac{s(a-2)n}{(s+1)ac} \leq 2$ . Thus  $\frac{n}{c} \leq \frac{8(s+1)}{2s} \leq \frac{96}{22} < 5 \leq p-2$ .

If  $\frac{n}{c} \ge 3$ , then  $\frac{(s-1)(a-2)n}{bc} > \frac{2(s-1)n}{(s+1)4c} > \frac{2\times 10\times 3}{12\times 4} > 1$ , by Lemma 4.2 we have  $\operatorname{ind}(S) = 1$ . Next as before we may assume  $2 < \frac{n}{c} < \frac{n}{b} < 3$ .

Since  $2 < \frac{n}{b} < 3$ , we have  $b > \frac{n}{3} > 96$  and  $b \ge 97$ . Assume that n = 2c + j = 2b + j + 4. If  $j \ge 17$ , we have  $2b^2 + (j - 10)b - 7(j + 4) - (2b^2 + 4b) = (j - 14)b - 7(j + 4) = j(b - 7) - 14(b + 2) = (j - 14)(b - 7) - 14 \times 9 \ge 3 \times 90 - 14 \times 9 = 144 > 0$  and

$$\frac{(s-1)n}{b} - \frac{(s-1)n}{c} = \frac{2(s-1)n}{bc} \ge \frac{(b-7)(2b+j+4)}{2b(b+2)} = \frac{2b^2 + (j-10)b - 7(j+4)}{2b^2 + 4b} > 1.$$

By Lemma 4.2 we have ind(S) = 1.

Next assume that  $j \leq 15$  (j is odd as n is odd). Let  $n = (j+3)t + r, r \in [0, j+2]$ .

First we claim that r < t. Since  $t = \lfloor \frac{n}{j+3} \rfloor \ge \lfloor \frac{n}{18} \rfloor \ge 16$  and  $r \le 17$ . If r = 17 = t+1, we have  $n = 18 \times 16 + 17 = 305$ , which is not a prime power. If r = 17 = t, then  $n = 18 \times 17 + 17 = 17 \times 19$ , which is not a prime power. If r = 16 = t, then  $n = 18 \times 16 + 16 = 16 \times 19$ , which is not a prime power. Hence our claim holds.

**Subcase 2.1.** If j = 1, let n = 5l + k with  $k \in [1, 4]$ , so  $l \ge 57$  and gcd(n, l) = 1. Then

$$S = (g) \cdot (g) \cdot (\frac{n-j}{2}g) \cdot (\frac{n+j+4}{2}g) \cdot ((n-4)g)$$

$$\sim (2g) \cdot (2g) \cdot ((n-1)g) \cdot (5g) \cdot ((n-8)g) \quad \text{(multiplied by 2)}$$

$$\sim (4lg) \cdot (4lg) \cdot ((3l+k)g) \cdot ((5l-k)g) \cdot ((4l+4k)g) \quad \text{(multiplied by 2l)}$$

$$\sim ((l+k)g) \cdot ((l+k)g) \cdot ((l-k)g) \cdot ((l+3k)g) \cdot ((l-3k)g) \quad \text{(multiplied by } n-1).$$

Since (l+k) + (l+k) + (l-k) + (l+3k) + (l-3k) = 5l + k = n, we have ind(S) = 1.

**Subcase 2.2.** If j = 3, let n = 7l + k with  $k \le 6$ , then  $l \ge 41 > k$  and gcd(n, l) = 1. If k > 0, then

$$S = (g) \cdot (g) \cdot (\frac{n-j}{2}g) \cdot (\frac{n+j+4}{2}g) \cdot ((n-4)g)$$

$$\sim (2g) \cdot (2g) \cdot ((n-3)g) \cdot (7g) \cdot ((n-8)g) \quad \text{(multiplied by 2)}$$

$$\sim (6lg) \cdot (6lg) \cdot ((5l+2k)g) \cdot ((7l-2k)g) \cdot ((4l+4k)g) \quad \text{(multiplied by 3l)}$$

$$\sim ((l+k)g) \cdot ((l+k)g) \cdot ((2l-k)g) \cdot (3kg) \cdot ((3l-3k)g) \quad \text{(multiplied by } n-1).$$

Since (l + k) + (l + k) + (2l - k) + 3k + (3l - 3k) = 7l + k = n, we have ind(S) = 1. If k = 0, then n is a power of 7.

$$S = (g) \cdot (g) \cdot (\frac{n-j}{2}g) \cdot (\frac{n+j+4}{2}g) \cdot ((n-4)g)$$

$$\sim (2g) \cdot (2g) \cdot ((n-3)g) \cdot (7g) \cdot ((n-8)g) \quad \text{(multiplied by 2)}$$

$$\sim (6(l-1)g) \cdot (6(l-1)g) \cdot ((5l+9)g) \cdot ((7l-21)g) \cdot ((4l+24)g)$$
(multiplied by  $3(l-1)$ )
$$\sim ((l+6)g) \cdot ((l+6)g) \cdot ((2l-9)g) \cdot (21g) \cdot ((3l-24)g)$$
(multiplied by  $n-1$ ),

where 3l - 24 = 3(l - 8) > 0. Since (l + 6) + (l + 6) + (2l - 9) + 21 + (3l - 24) = 7l = n, again we have ind(S) = 1.

**Subcase 2.3.** If gcd(n,t) = 1 and  $j \ge 5$ , we have

$$S = (g) \cdot (g) \cdot (\frac{n-j}{2}g) \cdot (\frac{n+j+4}{2}g) \cdot ((n-4)g)$$

$$\sim (2g) \cdot (2g) \cdot ((n-j)g) \cdot ((j+4)g) \cdot ((n-8)g) \quad \text{(multiplied by 2)}$$

$$\sim (2tg) \cdot (2tg) \cdot ((3t+r)g) \cdot ((t-r)g) \cdot (((j-5)t+r)g) \quad \text{(multiplied by } t).$$

Since 2t + 2t + (3t + r) + (t - r) + ((j - 5)t + r) = (j + 3)t + r = n, we have ind(S) = 1. If gcd(n,t) > 1 and j > 5, we have

$$S = (g) \cdot (g) \cdot (\frac{n-j}{2}g) \cdot (\frac{n+j+4}{2}g) \cdot ((n-4)g)$$

$$\sim (2g) \cdot (2g) \cdot ((n-j)g) \cdot ((j+4)g) \cdot ((n-8)g) \quad \text{(multiplied by 2)}$$

$$\sim (2(t+1)g) \cdot (2(t+1)g) \cdot ((3t+r-j)g) \cdot ((t-r+j+4)g) \cdot (((j-5)t+r-8)g)$$
(multiplied by  $(t+1)$ ),

where  $(j-5)t+r-9 \ge t+r-9 > 0$ . Since 2(t+1)+2(t+1)+(3t+r-j)+(t-r+j+4)+((j-5)t+r-8=(j+3)t+r=n, we infer ind(S)=1.

If gcd(n,t) > 1 and j = 5, we have

$$S = (g) \cdot (g) \cdot (\frac{n-j}{2}g) \cdot (\frac{n+j+4}{2}g) \cdot ((n-4)g)$$

$$\sim (2g) \cdot (2g) \cdot ((n-j)g) \cdot ((j+4)g) \cdot ((n-8)g) \quad \text{(multiplied by 2)}$$

$$\sim (2(t-1)g) \cdot (2(t-1)g) \cdot ((3t+r+j)g) \cdot ((t-r-j-4)g) \cdot (((j-5)t+r+8)g)$$
(multiplied by  $(t-1)$ ),

where  $t - r - j - 9 \ge t + r - 9 > 0$ . Since 2(t - 1) + 2(t - 1) + (3t + r + j) + (t - r = j - 4) + ((j - 5)t + r + 8 = (j + 3)t + r = n, again we infer ind(S) = 1.

Now Proposition 3.2 follows immediately from Lemmas 4.3, 4.4, 4.5 and 4.6.

# 4.3. Proof of Proposition 3.3.

In this subsection, we will provide a proof for Proposition 3.3. In terms of Proposition 2.2, from now on we may always assume that  $S = g^2 \cdot (cg) \cdot ((n-b)g)((n-3)g)$  where  $a = 3 \le b < c \le \frac{n-3}{2}$  and  $n = p^{\mu} \ge 289$  with  $p \ge 7$  and  $\mu \ge 2$  and show that  $\operatorname{ind}(S) = 1$ .

**Lemma 4.7.** If S is a minimal zero-sum sequence such that  $\frac{n}{c} > 4$ , then  $\operatorname{ind}(S) = 1$ .

*Proof.* If  $\frac{n}{c} > 8$ , we have  $\frac{sn}{bc} \ge \frac{sn}{3(s+1)c} > 2$  (for if  $s \le 2, \frac{n}{c} > 12$ ), so by Lemma 4.2 ind(S) = 1.

Next we assume that  $4 < \frac{n}{c} < 8$ . We divide our proof into two cases.

Case 1.  $\frac{n}{c} < m < \frac{n}{b}$  for some integer  $m \le 8$ . If  $\gcd(m,n) = 1$ , by Lemma 4.1,  $\operatorname{ind}(S) = 1$ . If  $\gcd(m,n) > 1$ , since  $m \le 8$ , we must have m = p = 7, so  $n \ge 7^3 = 343$ . If  $\frac{n}{c} < 6$ , we may take m = 6 and thus  $\operatorname{ind}(S) = 1$ . Thus we may assume  $\frac{n}{c} < 7 < \frac{n}{b}$ . Since  $n < 7c \le 21(s+1)$ , we have  $s \ge 16$ . Since  $\frac{sn}{bc} > \frac{7s}{3(s+1)} \ge \frac{112}{51} > 2$ , it follows from Lemma 4.2 that  $\operatorname{ind}(S) = 1$ .

Case 2.  $m < \frac{n}{c} < \frac{n}{b} < m+1$  with  $m \in [4,7]$ . If  $\frac{sn}{bc} > 2$ , By Lemma 4.2 we have  $\operatorname{ind}(S) = 1$ . So we may assume that  $\frac{sn}{bc} \le 2$ . Since  $s \ge 12$ , we have  $\frac{kn}{b} - \frac{kn}{c} = \frac{kn}{bc} > \frac{4k}{3(s+1)} \ge \frac{4(s-1)}{3(s+1)} > 1$  for k = s-1, s. Thus there exist two integers  $M_{s-1} \in [\frac{(s-1)n}{c}, \frac{(s-1)n}{b}]$  and  $M_s \in [\frac{sn}{c}, \frac{sn}{b}]$ . Note that  $|M_s - M_{s_1}| \le 2 + \frac{sn}{c} - \frac{(s-1)n}{b} = 2 + \frac{(c-s)n}{bc} \le 2 + \frac{(2s+3)n}{bc} \le 2 + 4 + \frac{3n}{bc} < 6 + \frac{n}{12c} < 7$ . We conclude that at least one of  $M_{s-1}, M_s$  is cop-prime to n. Since  $M_s a < n$ , it follows from Lemma 4.1 that  $\operatorname{ind}(S) = 1$ .

In terms of Lemma 4.7 and its proof, we may assume that  $m < \frac{n}{c} < \frac{n}{b} < m+1$  with  $m \in [2,3]$ .

**Lemma 4.8.** If S is a minimal zero-sum sequence such that  $3 < \frac{n}{c} < \frac{n}{b} < 4$ , then ind(S) = 1.

*Proof.* Suppose that n = 3c + j and n = 3b + j + 3, where  $1 \le j \le b - 4$  and gcd(j, 3) = 1. If  $j \ge 17$ , as in Case 2 of Lemma 4.6 we have

$$\frac{kn}{b} - \frac{kn}{c} = \frac{kn}{bc} \ge \frac{(b-5)(3b+j+3)}{3b(b+1)} > 1,$$

for k = s - 1, s. By Lemma 4.2 we have ind(S) = 1.

Next assume that  $j \leq 16$ . Note that

$$S = (g) \cdot (g) \cdot (\frac{n-j}{3}g) \cdot (\frac{n+j+3}{3}g)((n-3)g)$$
  
  $\sim (3g) \cdot (3g) \cdot ((n-j)g) \cdot ((j+3)g) \cdot ((n-9)g)$  (multiplied by 3).

Let S' = 3S. Next we show that ind(S) = ind(S') = 1.

**Case 1.**  $8 \le j \le 16$ . Let n = (j+1)t + r with  $r \in [0,j]$ . Since  $\frac{n}{j+1} \ge \frac{289}{17} = 17$ , we have  $t \ge 17 \ge j > r$ . If gcd(n,t) = 1, we have

$$S' \sim (3tg) \cdot (3tg) \cdot ((2t+r)g) \cdot ((t-r)g) \cdot (((j-8)t+r)g)$$
 (multiplied by t).

Since (3t) + (3t) + (2t + r) + (t - r) + ((j - 8)t + r) = (j + 1)t + r = n, we have ind(S) = 1. If gcd(n, t) > 1, we have

$$S' \sim (3(t-1)g) \cdot (3(t-1)g) \cdot ((t+r+j)g) \cdot ((2t-r-j-3)g) \cdot (((j-8)t+r+9)g)$$
 multiplied by  $t-1$ .

Note that  $t \ge 17 \ge j+1 \ge r+2$ . We always have 2t-r-j-3 > 0 for all  $j \in [8,16]$ . Again, we have 3(t-1)+3(t=1)+(t+r+j)+(2t-r-j-3)+((j-8)t+r+9)=(j+1)t+r=n, so  $\operatorname{ind}(S)=1$ .

Case 2. j = 7. Let n = 9t + r with  $r \in [1, 8]$ . Then  $t \ge 32 > r$ . If gcd(n, t) = 1, we have

$$S' \sim (3tg) \cdot (3tg) \cdot ((2t+r)g) \cdot ((t-r)g) \cdot (rg)$$
(multiplied by  $t$ ).

Since (3t) + (3t) + (2t + r) + (t - r) + r = 9t + r = n, we have  $\operatorname{ind}(S) = 1$ .

If gcd(n,t) > 1, then gcd(n,t-1) = 1. So we have

$$S' \sim (3(t-1)g) \cdot (3(t-1)g) \cdot ((2t+r+7)g) \cdot ((t-r-10)g) \cdot ((r+9)g)$$
 (multiplied by  $t-1$ ).

Since (3(t-1)) + (3(t-1)) + (2t+r+7) + (t-r-10) + (r+9) = 9t + r = n, we again have  $\operatorname{ind}(S) = 1$ .

Case 3. j = 5. Let n = 5t + r with  $r \in [1, 4]$ . Then gcd(n, t) = 1 and  $t \ge 57 > 3r$ . We have

$$S' \sim ((t-r)g) \cdot ((t-r)g) \cdot (2rg) \cdot ((t-3r)g) \cdot ((2t+4r)g)$$
(multiplied by  $2t$ ).

Since (t-r) + (t-r) + (2r) + (t-3r) + (2t+4r) = 5t + r = n, we have ind(S) = 1.

Case 4. 
$$j=4$$
. Let  $n=7t+r$  with  $r \in [0,6]$ . If  $r \neq 0$ , we have  $\gcd(n,t)=1$  
$$S' \sim (6tg) \cdot (6tg) \cdot ((6t+2r)g) \cdot ((7t-r)g) \cdot ((3t+3r)g)$$
 (multiplied by  $2t$ ) 
$$\sim ((t+r)g) \cdot ((t+r)g) \cdot ((t-r)g) \cdot (2rg) \cdot ((4t-2r)g)$$
 (multiplied by  $n-1=7t+r-1$ ).

Since 
$$(t+r) + (t+r) + (t-r) + (2r) + (4t-2r) = 7t + r = n$$
, ind $(S) = 1$ .  
If  $r = 0$ , we have  $p = 7$  and  $gcd(n, t-1) = 1$ 

(multiplied by n-1=7t+r-1).

$$S' \sim (6(t-1)g) \cdot (6(t-1)g) \cdot ((6t+2r+8)g) \cdot ((7t-r-14)g) \cdot ((3t+3r+18)g)$$
(multiplied by  $2(t-1)$ )
$$\sim ((t+r+6)g) \cdot ((t+r+6)g) \cdot ((t-r-8)g) \cdot ((2r+14)g) \cdot ((4t-2r-18)g)$$

Again, we have ind(S) = 1.

Case 5. j=2. Let n=5t+r with  $r\in[1,4]$ . Then  $\gcd(n,t)=1$ . We have

$$S' \sim (tg) \cdot (tg) \cdot ((t+r)g) \cdot ((5t-r)g) \cdot ((2t+2r)g)$$

$$(\text{multiplied by } 2t)$$

$$\sim (4tg) \cdot (4tg) \cdot ((4t+4r)g) \cdot ((5t-4r)g) \cdot ((3t+7r)g)$$

$$(\text{multiplied by } 4)$$

$$\sim ((t+r)g) \cdot ((t+r)g) \cdot ((t-3r)g) \cdot ((4r)g) \cdot ((2t-6r)g)$$

$$(\text{multiplied by } n-1=5t+r-1).$$

Since 
$$(t+r) + (t+r) + (t-3r) + (4r) + (2t-6r) = 5t + r = n$$
, ind $(S) = 1$ .

Case 6. j = 1. Let n = 4t + r with  $r \in [1, 3]$ . Then gcd(n, t) = 1. We have

$$S' \sim (3tg) \cdot (3tg) \cdot ((3t+r)g) \cdot ((4t)g) \cdot ((3t+3r)g)$$
(multiplied by t)
$$\sim ((t+r)g) \cdot ((t+r)g) \cdot (tg) \cdot (rg) \cdot ((t-2r)g)$$
(multiplied by  $n-1=4t+r-1$ ).

Since 
$$(t+r) + (t+r) + t + r + (t-2r) = 4t + r = n$$
, we have  $ind(S) = 1$ .

**Lemma 4.9.** If S is a minimal zero-sum sequence such that  $2 < \frac{n}{c} < \frac{n}{b} < 3$ , then ind(S) = 1.

*Proof.* We will show that there exist integers  $x, y \in [1, \lfloor \frac{b}{3} \rfloor]$  such that

$$\frac{n}{c} < 2 + \frac{x}{y} < \frac{n}{b}.$$

Then  $(2y+x)a < \frac{yn}{b}3 \le n$ . If  $\gcd(2y+x,n) = 1$ , then by Lemma 4.1  $\operatorname{ind}(S) = 1$  and we are done.

Suppose  $n=2b+b_0$ , where  $1 \le b_0 \le b-1$ . Since n is prime power and  $c=b+1 < \frac{n-1}{2} = b + \frac{b_0-1}{2}$ , we infer that

$$b_0 \equiv 1 \pmod{2}$$
 and  $b_0 > 3$ .

It suffices to show there exist  $x, y \in [1, \lfloor \frac{b}{3} \rfloor]$  such that

$$\frac{b_0-2}{b+1}<\frac{x}{y}<\frac{b_0}{b}$$
 and  $\gcd(2y+x,n)=1.$ 

If it is not easy to determine whether or not gcd(2y + x, n) = 1, we will show that there exist two more integers  $z, w \in [1, \lfloor \frac{b}{3} \rfloor]$  such that

$$\frac{b_0-2}{b+1} < \frac{z}{w} < \frac{b_0}{b}$$
, and  $1 \le |(2y+x) - (2w+z)| \le p-1$ .

Thus either gcd(2y + x, n) = 1 or gcd(2w + y, n) = 1, again we have ind(S) = 1. We divide the proof into three cases.

Case 1.  $b \equiv 0 \pmod{3}$ . Since n is prime power, we infer that  $b_0 \not\equiv 0 \pmod{3}$ . Suppose b = 3s.

If  $b_0 = 3t + 1$ , then let x = t and y = s. We infer that  $\frac{3t-1}{3s+1} < \frac{x}{y} < \frac{3t+1}{3s}$ . Since  $n = 2b + b_0 = 3(2s + t) + 1 = 3(2y + x) + 1$ , we have  $\gcd(2y + x, n) = 1$  and we are done.

If  $b_0 = 3t + 2$ , let x = t and y = s. Then  $\frac{3t}{3s+1} < \frac{x}{y} < \frac{3t+2}{3s}$ . Since  $n = 2b + b_0 = 3(2s+t) + 2 = 3(2y+x) + 2$ , we have gcd(2y+x,n) = 1 and we are done.

Case 2.  $b \equiv 1 \pmod{3}$ . Since  $n = p^{\mu}$  with a prime  $p \geq 7$ , we infer that  $b_0 \not\equiv 1 \pmod{3}$ . Suppose b = 3s + 1. Since  $289 \leq n < 9s + 3$ ,  $s \geq 32$ .

**Subcase 2.1.**  $b_0 = 3t$ . Since  $b_0 \equiv 1 \pmod{2}$  and  $b_0 > 3$ ,  $t \ge 3$ .

If s < 2t - 2, let x = t - 1, y = s. Then  $\frac{3t - 2}{3s + 2} < \frac{x}{y} < \frac{3t}{3s + 1}$ . Since  $n = 2b + b_0 = 3(2s + t) + 2 = 3(2y + x) + 5$ , we have  $\gcd(2y + x, n) = 1$  and we are done.

Next assume that  $s \ge 2t - 2$ . Choose  $y = s - \lceil \frac{s - 2t + 3}{3t - 2} \rceil$  and x = t - 1. Then  $\frac{3t - 2}{3s + 2} < \frac{x}{y} < \frac{3t}{3s + 1}$ . If  $s \ge 3t + 1$ , let w = y - 1, z = t - 1, then  $\frac{3t - 2}{3s + 2} < \frac{z}{w} < \frac{3t}{3s + 1}$  and |(2y + x) - (2w + z)| = 2, so we are done. Now assume that

$$2t - 2 < s < 3t$$
.

Since  $32 \le s \le 3t$ ,  $t \ge 11$ . If  $s \le \frac{11(t-1)}{4}$ , let  $y = s - \lceil \frac{s-2t+3}{3t-2} \rceil = s-1$  and x = t-1 as before. Let  $w = s - \lceil \frac{4s-2t+5}{3t-2} \rceil = s-3$  and z = t-2. Then  $\frac{3t-2}{3s+2} < \frac{x}{y}, \frac{z}{w} < \frac{3t}{3s+1}$  and |(2y+x)-(2w+z)| = 5, so we are done. If  $\frac{11(t-1)}{4} < s \le 3t$ , let y = s-2, x = t-1, w = s-4 and z = t-2. Then  $\frac{3t-2}{3s+2} < \frac{x}{y}, \frac{z}{w} < \frac{3t}{3s+1}$  and |(2y+x)-(2w+z)| = 5, so we are done.

**Subcase 2.2.**  $b_0 = 3t + 2$ . Let x = t and y = s. We infer that  $\frac{3t}{3s+2} < \frac{x}{y} < \frac{3t+2}{3s+1}$ . Since  $n = 2b + b_0 = 3(2s + t) + 4 = 3(2y + x) + 4$ , we have gcd(2y + x, n) = 1 and we are done.

Case 3.  $b \equiv 2 \pmod{3}$ . Since  $n = p^{\mu}$  with a prime  $p \geq 7$ , we infer that  $b_0 \not\equiv 2 \pmod{3}$ . Suppose b = 3s + 2. Since  $289 \leq 3b - 1 = 9s + 5$ ,  $s \geq 32$ .

**Subcase 3.1.**  $b_0 \equiv 0 \pmod{3}$ . Suppose  $b_0 = 3t$ . Recall that  $b_0 = 3t \equiv 1 \pmod{2}$  and  $b_0 > 3$ , we infer that  $t \geq 3$  and  $t \equiv 1 \pmod{2}$ .

If  $s < \frac{3t-4}{2}$ , then let  $x=t-1, \ y=s; \ z=t-2, \ w=s-1$ . We infer that  $\frac{3t-2}{3s+3} < \frac{z}{w} < \frac{x}{y} < \frac{3t}{3s+2}$  and |(2y+x)-(2w+z)|=3, so we are done. Next assume that  $s \geq \frac{3t-4}{2}$ . If s < 3t-3, then  $t \geq 13$ . Let  $x=t-1, \ y=s; \ z=t-2, \ w=s-2$ . Then  $\frac{3t-2}{3s+3} < \frac{x}{y}, \frac{z}{w} < \frac{3t}{3s+2}$  and |(2y+x)-(2w+z)|=5, so we are done.

If  $s \geq 3t-3$ , choose  $y=s-\lceil\frac{s-3t+4}{3t-2}\rceil$  and x=t-1. Then  $\frac{3t-2}{3s+3}<\frac{x}{y}<\frac{3t}{3s+2}$ . Let w=y-1, z=t-1. Note that if s<3t then t>11. Thus we have either  $s\geq 3t$  or  $t\geq 9$ . Therefore,  $\frac{3t-2}{3s+3}<\frac{z}{w}<\frac{3t}{3s+2}$ . Since |(2y+x)-(2w+z)|=2, we are done.

**Subcase 3.2.**  $b_0 \equiv 1 \pmod{3}$ . Suppose  $b_0 = 3t + 1$ . Recall that  $b_0 = 3t + 1 \equiv 1 \pmod{2}$  and  $b_0 > 3$ . Hence  $t \equiv 0 \pmod{2}$  and thus  $t \geq 2$ .

If s > 5t + 1, let x = t, y = s; z = t, w = s - 1. Then  $\frac{3t - 1}{3s + 3} < \frac{x}{y} < \frac{z}{w} < \frac{3t + 1}{3s + 2}$  and |(2y + x) - (2w + z)| = 2, so we are done. Hence we may assume that  $s \le 5t + 1$ .

If  $s \geq 3t-2$ , let  $x=z=t-1, y=s-\lceil \frac{2s-3t+4}{3t-1} \rceil, w=y-1$ . If  $t \geq 10$ , then  $\frac{3t-1}{3s+3} < \frac{z}{w}, \frac{x}{y} < \frac{3t+1}{3s+2}$  and |(2y+x)-(2w+z)|=2, so we are done. Hence we may assume that  $t \leq 9$  and  $3t-2 \leq s \leq 5t+1$ . Note that  $t \equiv 0 \pmod{2}$  and  $s \geq 33$ . Hence t=8 and then  $33 \leq s \leq 41$ . Let x=z=t-1, y=s-3, w=s-4. Then  $\frac{3t-1}{3s+3} < \frac{z}{w}, \frac{x}{y} < \frac{3t+1}{3s+2}$  and |(2y+x)-(2w+z)|=2, so we are done.

If 2t < s < 3t - 2, let x = t, y = s; z = t - 1, w = s - 1. Then  $\frac{3t - 1}{3s + 3} < \frac{x}{y}$ ,  $\frac{z}{w} < \frac{3t + 1}{3s + 2}$  and |(2y + x) - (2w + z)| = 3, so we are done.

Hence we may assume that  $s \leq 2t$ . Note that  $t \geq \frac{s}{2} \geq \frac{33}{2}$ , so  $t \geq 17$ .

If  $s < \frac{6t-7}{5}$ , let x = t-1, y = s; z = t-2, w = s-1. Then  $\frac{3t-1}{3s+3} < \frac{z}{w} < \frac{x}{y} < \frac{3t+1}{3s+2}$  and |(2y+x)-(2w+z)|=3, so we are done. Hence we may assume that  $s \ge \frac{6t-7}{5}$ .

If  $s < \frac{3t-3}{2}$ , let x = t-1, y = s; z = t-2,  $w = s - \lceil \frac{5s-3t+3}{3t-1} \rceil$ . Then  $\frac{3t-1}{3s+3} < \frac{z}{w}, \frac{x}{y} < \frac{3t+1}{3s+2}$  and  $1 \le |(2y+x)-(2w+z)| \le 5$ , so we are done. Hence we may assume that  $s \ge \frac{3t-3}{2}$ .

Next assume that  $\frac{3t-3}{2} \le s \le 2t$ .

Let x=t-1 and y=s-1. We infer that  $\frac{3t-1}{3s+3}<\frac{t-1}{s-1}<\frac{3t+1}{3s+2}$ . Let  $z=t-2,\,w=s-\left\lceil\frac{5s-3t+3}{3t-1}\right\rceil$ . Then  $\frac{3t-1}{3s+3}<\frac{z}{w},\frac{x}{y}<\frac{3t+1}{3s+2}$  and  $1\leq |(2y+x)-(2w+z)|\leq 5$ , so we are done.

Now Proposition 3.3 follows immediately from Lemmas 4.7, 4.8, 4.9 and Proposition 2.2.

We remark that the case when h(S) = 1 is much more complicated and  $\operatorname{ind}(S)$  is not yet determined (even when n = p is a prime number). We conclude the paper by listing this case as an open problem.

**Open Problem.** Let S be a minimal zero-sum sequence of length five over a cyclic group of order n. Determine  $\operatorname{ind}(S)$  when the height h(S) = 1.

#### ACKNOWLEDGEMENTS

The first author is supported by the NNSF of China (Grant No. 11001110 and 11271131) and *Jiangsu Government Scholarship for Overseas Studies*. The second author is supported by a Discovery Grant from the Natural Science and Engineering Research Council of Canada. The third author is supported by the NSFC (Grant No. 11271207 and 11301531)

#### REFERENCES

- [1] S.T. Chapman, M. Freeze, and W.W Smith, Minimal zero sequences and the strong Davenport constant, Discrete Math. 203(1999), 271-277.
- [2] S.T. Chapman, and W.W Smith, A characterization of minimal zero-sequences of index one in finite cyclic groups, Integers 5(1)(2005), Paper A27, 5p.
- [3] W. Gao, Zero sums in finite cyclic groups, Integers 0 (2000), Paper A14, 9p.
- [4] W. Gao and A. Geroldinger, On products of k atoms, Monatsh. Math. 156 (2009), 141-157.
- [5] W. Gao, Y. Li, J. Peng, P. Plyley and G. Wang On the index of sequences over cyclic groups (English), Acta Arith. 148, No. 2, (2011) 119-134.
- [6] A. Geroldinger, On non-unique factorizations into irreducible elements. II, Number Theory, Vol II Budapest 1987, Colloquia Mathematica Societatis Janos Bolyai, vol. 51, North Holland, 1990, 723-757.
- [7] A. Geroldinger, Additive group theory and non-unique factorizations, Combinatorial Number Theory and Additive Group Theory (A. Geroldinger and I. Ruzsa, eds.), Advanced Courses in Mathematics CRM Barcelona, Birkhäuser, 2009, pp. 1-86.
- [8] A. Geroldinger and F. Halter-Koch, Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory, Pure and Applied Mathematics, Vol. 278, Chapman & Hall/CRC, 2006.
- [9] D. Kleitman and P. Lemke, An addition theorem on the integers modulo n, J. Number Theory 31(1989), 335-345.
- [10] Y. Li and J. Peng, Minimal zero-sum sequences of length four over finite cyclic groups II, Int. J. Number Theory, 9(2013), 845-866
- [11] Y. Li, C. Plyley, P. Yuan and X. Zeng, Minimal zero sum sequences of length four over finite cyclic groups, J. Number Theory. 130 (2010), 2033-2048.
- [12] J. Peng, Y. Li, Minimal zero-sum sequences of length five over finite cyclic groups, Ars Combinatoria, 112(2013), 373-384.
- [13] V. Ponomarenko, Minimal zero sequences of finite cyclic groups, Integers 4(2004), Paper A24, 6p.
- [14] S. Savchen and F. Chen, Long zero-free sequences in finite cyclic groups, Discrete Math. 307(2007), 2671-2679.

- [15] Caixia Shen, Li-meng Xia, On the index-conjecture of length four minimal zero-sum sequences II, Int. J. Number Theory, to appear. DOI: 10.1142/S179304211350111X.
- [16] Caixia Shen, Li-meng Xia, Yuanlin Li, On the index of length four minimal zero-sum sequences, preprint.
- [17] Li-meng. Xia, On the index-conjecture of length four minimal zero-sum sequences, Int. J. Number Theory, 9(2013), 1505-1528.
- [18] Li-meng Xia, Caixia Shen, Minimal zero-sum sequences of length four over cyclic group with order  $n = p^{\alpha}q^{\beta}$ , J. Number Theory 133 (2013), 4047-4068.
- [19] X. Xia and P. Yuan, Indexes of insplitable minimal zero-sum sequences of length  $l(C_n) 1$ , Discrete Math., 310(2010)1127-1133.
- [20] P. Yuan, On the index of minimal zero-sum sequences over finite cyclic groups, J. Combin. Theory Ser. A114(2007), 1545-1551.
- [21] P. Yuan and X. Zeng, Indexes of long zero-sum free sequences over cyclic groups, Eur. J. Comb. 32(2011), 1213-1221.