EXCEPTIONAL SETS IN WARING'S PROBLEM: TWO SQUARES AND s BIQUADRATES

LILU ZHAO

ABSTRACT. Let $R_s(n)$ denote the number of representations of the positive number n as the sum of two squares and s biquadrates. When s=3 or 4, it is established that the anticipated asymptotic formula for $R_s(n)$ holds for all $n \leq X$ with at most $O(X^{(9-2s)/8+\varepsilon})$ exceptions.

1. Introduction

Waring's problem for sums of mixed powers involving one or two squares has been widely investigated. In 1987-1988, Brüdern [1, 2] considered the representation of n in the form

$$n = x_1^2 + x_2^2 + y_1^{k_1} + \dots + y_s^{k_s},$$

with $k_1^{-1} + \cdots + k_s^{-1} > 1$. Linnik [8] and Hooley [6] investigated sums of two squares and three cubes. In 2002, Wooley [11] investigated the exceptional set related to the asymptotic formula in Waring's problem involving one square and five cubes. Recently, Brüdern and Kawada [3] established the asymptotic formula for the number of representations of the positive number n as the sum of one square and seventeen fifth powers.

Let $R_s(n)$ denote the number of representations of the positive number n as the sum of two squares and s biquadrates. Very recently, subject to the truth of the Generalised Riemann Hypothesis and the Elliott-Halberstam Conjecture, Friedlander and Wooley [4] established that $R_3(n) > 0$ for all large n under certain congruence conditions. They also showed that if one is prepared to permit a small exceptional set of natural numbers n, then the anticipated asymptotic formula for $R_s(n)$ can be obtained. To state their results precisely, we introduce some notations. We define

(1.1)
$$\mathfrak{S}_s(n) = \sum_{q=1}^{\infty} \sum_{\substack{a=1\\(a,q)=1}}^{q} q^{-2-s} S_2(q,a)^2 S_4(q,a)^s e(-na/q),$$

²⁰¹⁰ Mathematics Subject Classification. Primary 11P05; Secondary 11P55, 11N37. Key words and phrases. circle method, Waring's problem, exceptional sets, asymptotic formula.

where the Gauss sum $S_k(q, a)$ is defined as

(1.2)
$$S_k(q, a) = \sum_{r=1}^{q} e(ar^k/q).$$

As in [4], we refer a function $\psi(t)$ as being a sedately increasing function when $\psi(t)$ is a function of a positive variable t, increasing monotonically to infinity, and satisfying the condition that when t is large, one has $\psi(t) = O(t^{\delta})$ for a positive number δ sufficiently small in the ambient context. Then we introduce $E_s(X, \psi)$ to denote the number of integers n with $1 \leq n \leq X$ such that

(1.3)
$$\left| R_s(n) - c_s \Gamma(\frac{5}{4})^4 \mathfrak{S}_s(n) n^{s/4} \right| > n^{s/4} \psi(n)^{-1},$$

where $c_3 = \frac{2}{3}\sqrt{2}$ and $c_4 = \frac{1}{4}\pi$. Friedlander and Wooley [4] established the upper bounds

$$(1.4) E_3(X,\psi) \ll X^{1/2+\varepsilon}\psi(X)^2$$

and

$$(1.5) E_4(X,\psi) \ll X^{1/4+\varepsilon} \psi(X)^4,$$

where $\varepsilon > 0$ is arbitrary small.

The main purpose of this note is to establish the following result.

Theorem 1.1. Suppose that $\psi(t)$ is a sedately increasing function. Let $E_s(X, \psi)$ be defined as above. Then for each $\varepsilon > 0$, one has

(1.6)
$$E_3(X,\psi) \ll X^{3/8+\varepsilon} \psi(X)^2$$

and

(1.7)
$$E_4(X,\psi) \ll X^{1/8+\varepsilon} \psi(X)^2,$$

where the implicit constants may depend on ε .

We establish Theorem 1.1 by means of the Hardy-Littlewood method. In order to estimate the corresponding exceptional sets effectively, we employ the method developed by Wooley [10, 11].

As usual, we write e(z) for $e^{2\pi iz}$. Whenever ε appears in a statement, either implicitly or explicitly, we assert that the statement holds for each $\varepsilon > 0$. Note that the "value" of ε may consequently change from statement to statement. We assume that X is a large positive number, and $\psi(t)$ is a sedately increasing function.

2. Preparations

Throughout this section, we assume that $X/2 < n \le X$. For $k \in \{2, 4\}$, we define the exponential sum

$$f_k(\alpha) = \sum_{1 \le x \le P_k} e(\alpha x^k),$$

where $P_k = X^{1/k}$. We take s to be either 3 or 4. By orthogonality, we have

(2.1)
$$R_s(n) = \int_0^1 f_2(\alpha)^2 f_4(\alpha)^s e(-n\alpha) d\alpha.$$

When Q is a positive number, we define $\mathfrak{M}(Q)$ to be the union of the intervals

$$\mathfrak{M}_Q(q, a) = \{ \alpha : |q\alpha - a| \leqslant QX^{-1} \},$$

with $1 \leqslant a \leqslant q \leqslant Q$ and (a,q) = 1. Whenever $Q \leqslant X^{1/2}/2$, the intervals $\mathfrak{M}_Q(q,a)$ are pairwise disjoint for $1 \leqslant a \leqslant q \leqslant Q$ and (a,q) = 1. Let ν be a sufficiently small positive number, and let $R = P_4^{\nu}$. We take $\mathfrak{M} = \mathfrak{M}(R)$ and $\mathfrak{m} = (R/N, 1 + R/N) \setminus \mathfrak{M}$. Write

$$v_k(\beta) = \int_0^{P_k} e(\gamma^k \beta) d\gamma.$$

One has the estimate

$$v_k(\beta) \ll P_k(1+X|\beta|)^{-1/k}.$$

For $\alpha \in \mathfrak{M}_{X^{1/2}/2}(q, a) \subseteq \mathfrak{M}(X^{1/2}/2)$, we define

(2.2)
$$f_k^*(\alpha) = q^{-1} S_k(q, a) v_k(\alpha - a/q).$$

It follows from Theorem 4.1 [9] that whenever $\alpha \in \mathfrak{M}_{X^{1/2}/2}(q,a)$, one has

$$(2.3) f_k(\alpha) - f_k^*(\alpha) \ll q^{1/2} (1 + X|\alpha - a/q|)^{1/2} X^{\varepsilon}.$$

We define the multiplicative function $w_k(q)$ by taking

$$w_k(p^{uk+v}) = \begin{cases} kp^{-u-1/2}, & \text{when } u \geqslant 0 \text{ and } v = 1, \\ p^{-u-1}, & \text{when } u \geqslant 0 \text{ and } 2 \leqslant v \leqslant k. \end{cases}$$

Note that $q^{-1/2} \leqslant w_k(q) \ll q^{-1/k}$. Whenever (a,q) = 1, we have

$$q^{-1}S_k(q,a) \ll w_k(q).$$

Therefore for $\alpha = a/q + \beta \in \mathfrak{M}_{X^{1/2}/2}(q, a) \subseteq \mathfrak{M}(X^{1/2}/2)$, one has

$$(2.4) f_k^*(\alpha) \ll w_k(q) P_k(1+X|\beta|)^{-1/k} \ll P_k q^{-1/k} (1+X|\beta|)^{-1/k}.$$

The following conclusion is (4.1) in [4].

Lemma 2.1. One has

$$\int_{\mathfrak{M}} f_2(\alpha)^2 f_4(\alpha)^s e(-n\alpha) d\alpha = c_s \Gamma(\frac{5}{4})^4 \mathfrak{S}_s(n) n^{s/4} + O(n^{s/4-\kappa+\varepsilon}),$$

for a suitably small positive number κ .

The next result provides the value of the Gauss sum $S_2(q, a)$.

Lemma 2.2. The Gauss sum $S_2(q, a)$ satisfies the following properties.

(i) If (2a, q) = 1, then

$$S_2(q, a) = \left(\frac{a}{q}\right) S_2(q, 1).$$

Here by $\left(\frac{a}{q}\right)$ we denote the Jacobi symbol.

(ii) If q is odd, then

$$S_2(q,1) = \begin{cases} q^{1/2}, & \text{if } q \equiv 1 \pmod{4}, \\ iq^{1/2}, & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

(iii) If (2, a) = 1, then

$$S_2(2^m, a) = \begin{cases} 0, & \text{if } m = 1, \\ 2^{m/2}(1 + i^a), & \text{if } m \text{ is even,} \\ 2^{(m+1)/2}e(a/8), & \text{if } m > 1 \text{ and } m \text{ is odd.} \end{cases}$$

(iv) If $(q_1, q_2) = 1$, then

$$S_2(q_1q_2, a_1q_2 + a_2q_1) = S_2(q_1, a_1)S_2(q_2, a_2).$$

Proof. These properties can be found in Lemma 2 [5].

3. The Proof of Theorem 1.1

Let τ be a fixed sufficiently small positive number. Set $Y = P_4^{3/2+\tau}\psi(X)^2$. We define $\mathfrak{m}_1 = \mathfrak{m} \setminus \mathfrak{M}(X^{1/2}/2)$, $\mathfrak{m}_2 = \mathfrak{M}(X^{1/2}/2) \setminus \mathfrak{M}(Y)$, $\mathfrak{m}_3 = \mathfrak{M}(Y) \setminus \mathfrak{M}(P_4)$ and $\mathfrak{m}_4 = \mathfrak{M}(P_4) \setminus \mathfrak{M}$. Let $\eta(n)$ be sequence of complex numbers satisfying $|\eta(n)| = 1$. Let \mathcal{Z} be a subset of $\{n \in \mathbb{N} : X/2 < n \leq X\}$. We abbreviate $\operatorname{card}(\mathcal{Z})$ to Z. Then we introduce the exponential sum $\mathcal{E}(\alpha)$ by

$$\mathcal{E}(\alpha) = \sum_{n \in \mathcal{Z}} \eta(n) e(-n\alpha).$$

For $1 \leq j \leq 4$, we define

(3.1)
$$\mathcal{I}_{j} = \int_{\mathfrak{m}_{j}} \left| f_{2}(\alpha)^{2} f_{4}(\alpha)^{s} \mathcal{E}(\alpha) \right| d\alpha.$$

Lemma 3.1. Let \mathcal{I}_1 be defined in (3.1). Then we have

$$\mathcal{I}_1 \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z^{1/2} + P_4^{s-\frac{1}{4}+\varepsilon} Z.$$

Proof. For any $\alpha \in \mathfrak{m}_1$, there exist a and q with $1 \leqslant a \leqslant q \leqslant 2X^{1/2}$ and (a,q)=1 such that $|q\alpha-a|\leqslant X^{-1/2}/2$. Since $\alpha \in \mathfrak{m}_1$, we conclude that $q>X^{1/2}/2$. It follows from Weyl's inequality (Lemma 2.4 [9]) that

$$f_2(\alpha) \ll P_2^{1/2+\varepsilon}$$
 for $\alpha \in \mathfrak{m}_1$.

Thus we have

$$\mathcal{I}_{1} \ll P_{2}^{1+\varepsilon} \int_{\mathfrak{m}_{1}} \left| f_{4}(\alpha)^{s} \mathcal{E}(\alpha) \right| d\alpha$$

$$\ll P_{2}^{1+\varepsilon} \left(\int_{0}^{1} \left| f_{4}(\alpha)^{6} \right| d\alpha \right)^{\frac{1}{2}} \left(\int_{0}^{1} \left| f_{4}(\alpha)^{2(s-3)} \mathcal{E}(\alpha)^{2} \right| d\alpha \right)^{\frac{1}{2}}.$$

By Hua's inequality (Lemma 2.5 [9]) and Schwartz's inequality,

$$\int_0^1 |f_4(\alpha)^6| d\alpha \ll \Big(\int_0^1 |f_4(\alpha)^4| d\alpha\Big)^{1/2} \Big(\int_0^1 |f_4(\alpha)^8| d\alpha\Big)^{1/2} \ll P_4^{7/2+\varepsilon}.$$

When s=4, one has the bound $\int_0^1 |f_4(\alpha)^{2(s-3)} \mathcal{E}(\alpha)^2| d\alpha \ll P_4 Z + P_4^{\varepsilon} Z^2$. Then we can conclude that

(3.2)
$$\mathcal{I}_1 \ll P_4^{4 - \frac{1}{4} + \frac{s - 3}{2} + \varepsilon} Z^{1/2} + P_4^{s - \frac{1}{4} + \varepsilon} Z.$$

Indeed when s=3, the estimate (3.2) holds with $P_4^{s-\frac{1}{4}+\varepsilon}Z$ omitted.

Lemma 3.2. Let \mathcal{I}_2 be given by (3.1). Then one has

$$\mathcal{J}_2 \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z^{1/2} + P_4^{s-\tau/2+\varepsilon} \psi(X)^{-1} Z.$$

Proof. We introduce

$$\mathcal{J}_1 = \int_{\mathfrak{m}_2} \left| \left(f_2(\alpha) - f_2^*(\alpha) \right)^2 f_4(\alpha)^s \mathcal{E}(\alpha) \right| d\alpha$$

and

$$\mathcal{J}_2 = \int_{\mathfrak{m}_2} \left| f_2^*(\alpha)^2 f_4(\alpha)^s \mathcal{E}(\alpha) \right| d\alpha.$$

Note that $|f_2(\alpha)|^2 \ll |f_2(\alpha) - f_2^*(\alpha)|^2 + |f_2^*(\alpha)|^2$, where $f_2^*(\alpha)$ is defined in (2.2). Then one has

$$(3.3) \mathcal{I}_2 \ll \mathcal{J}_1 + \mathcal{J}_2.$$

In view of (2.3), we know $f_2(\alpha) - f_2^*(\alpha) \ll P_2^{1/2+\varepsilon}$ for $\alpha \in \mathfrak{m}_2$. The argument leading to (3.2) also implies

(3.4)
$$\mathcal{J}_1 \ll P_4^{4 - \frac{1}{4} + \frac{s - 3}{2} + \varepsilon} Z^{1/2} + P_4^{s - \frac{1}{4} + \varepsilon} Z.$$

One has, by Schwartz's inequality, that

$$\mathcal{J}_2 \leqslant \left(\int_{\mathfrak{m}_2} \left| f_4(\alpha)^6 \right| d\alpha \right)^{1/2} \mathcal{J}^{1/2} \ll P_4^{7/4 + \varepsilon} \mathcal{J}^{1/2},$$

where \mathcal{J} is defined as

$$\mathcal{J} = \int_{\mathfrak{m}_2} \left| f_2^*(\alpha)^4 f_4(\alpha)^{2(s-3)} \mathcal{E}(\alpha)^2 \right| d\alpha.$$

In order to handle \mathcal{J} , we need the following estimate

(3.5)
$$\int_{\mathfrak{m}_2} |f_2^*(\alpha)^4| e(-h\alpha) d\alpha = \begin{cases} O(P_4^{4+\varepsilon} Y^{-1}), & \text{when } 0 < |h| \leqslant 2X, \\ O(P_4^{4+\varepsilon}), & \text{when } h = 0. \end{cases}$$

Recalling the definition of $f_2^*(\alpha)$, we conclude that

$$\int_{\mathfrak{m}_{2}} \left| f_{2}^{*}(\alpha)^{4} \right| e(-h\alpha) d\alpha$$

$$= \sum_{q \leqslant X^{1/2}/2}^{*} \int_{|\beta| \leqslant \frac{1}{2qX^{1/2}}}^{*} q^{-4} \left(\sum_{\substack{a=1\\ (a,q)=1}}^{q} |S_{2}(q,a)|^{4} e(-ha/q) \right) |v_{2}(\beta)|^{4} e(-h\beta) d\beta,$$

where the notations \sum^* and \int^* mean either q > Y or $Xq|\beta| > Y$. Whenever (a,q) = 1, one has by Lemma 2.2 that

$$|S_2(q,a)| = |S_2(q,1)| \le (2q)^{1/2}$$
.

We obtain

$$\left| \sum_{\substack{a=1\\(a,q)=1}}^{q} |S_2(q,a)|^4 e(-ha/q) \right| = |S_2(q,1)|^4 \left| \sum_{\substack{a=1\\(a,q)=1}}^{q} e(-ha/q) \right|$$

$$\leq 4q^2 \left| \sum_{\substack{a=1\\(a,q)=1}}^{q} e(-ha/q) \right| \leq 4q^2 (q,h),$$

whence

$$\int_{\mathfrak{m}_2} \left| f_2^*(\alpha)^4 \right| e(-h\alpha) d\alpha \ll P_2^4 \sum_{q \leqslant X^{1/2}/2}^* \int_{|\beta| \leqslant \frac{1}{2qX^{1/2}}}^* \frac{q^{-2}(q,h)}{(1+X|\beta|)^2} d\beta.$$

When h = 0, we have

$$\int_{\mathfrak{m}_{2}} \left| f_{2}^{*}(\alpha)^{4} \right| e(-h\alpha) d\alpha \ll P_{2}^{4} \sum_{q \leqslant X^{1/2}/2} \int_{|\beta| \leqslant \frac{1}{2qX^{1/2}}} q^{-1} (1+X|\beta|)^{-2} d\beta$$
$$\ll P_{2}^{4} X^{-1} \log X.$$

When $h \neq 0$, we get

$$\int_{\mathfrak{m}_{2}} \left| f_{2}^{*}(\alpha)^{4} \right| e(-h\alpha) d\alpha \ll P_{2}^{4} Y^{-1} \sum_{q \leqslant X^{1/2}/2} \int_{|\beta| \leqslant \frac{1}{2qX^{1/2}}} \frac{q^{-1}(q,h)}{1 + X|\beta|} d\beta$$
$$\ll P_{2}^{4} Y^{-1} X^{-1} (\log X) \sum_{q \leqslant X^{1/2}/2} q^{-1}(q,h)$$
$$\ll P_{2}^{4} Y^{-1} X^{-1+\varepsilon}.$$

The conclusion (3.5) is established. Now we are able to estimate \mathcal{J} . When s=4,

$$\mathcal{J} = \sum_{\substack{1 \leqslant x_1, x_2 \leqslant P_4 \\ n_1, n_2 \in \mathcal{Z}}} \eta(n_1) \overline{\eta(n_2)} \int_{\mathfrak{m}_2} |f_2^*(\alpha)^4| e(-(x_1^4 - x_2^4 + n_1 - n_2)\alpha) d\alpha.$$

On applying (3.5), we can deduce that

$$\mathcal{J} \ll \sum_{\substack{1 \leqslant x_1, x_2 \leqslant P_4, \ n_1, n_2 \in \mathcal{Z} \\ x_1^4 - x_2^4 + n_1 - n_2 \neq 0}} P_4^{4+\varepsilon} Y^{-1} + \sum_{\substack{1 \leqslant x_1, x_2 \leqslant P_4, \ n_1, n_2 \in \mathcal{Z} \\ x_1^4 - x_2^4 + n_1 - n_2 = 0}} P_4^{4+\varepsilon}$$

$$\ll P_4^{6+\varepsilon} Z^2 Y^{-1} + P_4^{4+\varepsilon} Z^2 + P_4^{5+\varepsilon} Z.$$

Substituting $Y = P_4^{3/2+\tau} \psi(X)^2$, we finally obtain

$$\mathcal{J} \ll P_4^{4+1/2-\tau+\varepsilon} \psi(X)^{-2} Z^2 + P_4^{5+\varepsilon} Z,$$

whence

$$\mathcal{J}_2 \ll P_4^{4-\tau/2+\varepsilon} \psi(X)^{-1} Z + P_4^{4+1/4+\varepsilon} Z^{1/2}$$

Similarly, when s = 3, one has

$$\mathcal{J} \ll P_4^{5/2-\tau+\varepsilon} \psi(X)^{-2} Z^2 + P_4^{4+\varepsilon} Z$$

whence

$$\mathcal{J}_2 \ll P_4^{3-\tau/2+\varepsilon} \psi(X)^{-1} Z + P_4^{4-1/4+\varepsilon} Z^{1/2}.$$

Therefore, we conclude that

(3.6)
$$\mathcal{J}_2 \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z^{1/2} + P_4^{s-\tau/2+\varepsilon} \psi(X)^{-1} Z.$$

Combining (3.3), (3.4) and (3.6), we conclude that

(3.7)
$$\mathcal{I}_2 \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z^{1/2} + P_4^{s-\tau/2+\varepsilon} \psi(X)^{-1} Z.$$

We complete the proof.

Lemma 3.3. Let \mathcal{I}_3 be defined in (3.1). Then we have

$$\mathcal{I}_3 \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z^{1/2} + P_4^{s-\tau+\varepsilon} \psi(X)^{-1} Z.$$

Proof. Similarly to (3.3) and (3.4), we can derive that

(3.8)
$$\mathcal{I}_3 \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z^{1/2} + P_4^{s-\frac{1}{4}+\varepsilon} Z + \mathcal{K},$$

where

$$\mathcal{K} = \int_{\mathfrak{m}_3} \left| f_2^*(\alpha)^2 f_4(\alpha)^s \mathcal{E}(\alpha) \right| d\alpha.$$

One has

$$\mathcal{K} \leqslant \sup_{\alpha \in \mathfrak{m}_3} |f_4(\alpha)| \Big(\int_{\mathfrak{m}_3} \left| f_2^*(\alpha)^2 f_4(\alpha)^4 \right| d\alpha \Big)^{1/2}$$

$$\times \Big(\int_{\mathfrak{m}_3} \left| f_2^*(\alpha)^2 f_4(\alpha)^{2(s-3)} \mathcal{E}(\alpha)^2 \right| d\alpha \Big)^{1/2}.$$

In view of (2.3) and (2.4), we have for $\alpha \in \mathfrak{m}_3$ that

$$f_4(\alpha) \ll P_4 q^{-1/4} (1 + X|\alpha - a/q|)^{-1/4} + Y^{1/2} X^{\varepsilon} \ll P_4^{3/4 + \tau/2 + \varepsilon} \psi(X).$$

Since $f_2^*(\alpha) - f_2(\alpha) \ll P_2^{1/2}$ for $\alpha \in \mathfrak{m}_3$, we easily deduce that

$$\int_{\mathfrak{m}_{3}} \left| f_{2}^{*}(\alpha)^{2} f_{4}(\alpha)^{4} \right| d\alpha \ll P_{2}^{1/2} \int_{0}^{1} \left| f_{2}(\alpha) f_{4}(\alpha)^{4} \right| d\alpha + \int_{0}^{1} \left| f_{2}(\alpha)^{2} f_{4}(\alpha)^{4} \right| d\alpha \ll P_{4}^{4+\varepsilon}.$$

Therefore we arrive at

$$\mathcal{K} \ll P_4^{11/4+\tau/2+\varepsilon} \psi(X) \Big(\int_{\mathfrak{m}_3} \left| f_2^*(\alpha)^2 f_4(\alpha)^{2(s-3)} \mathcal{E}(\alpha)^2 \right| d\alpha \Big)^{1/2}.$$

Similarly to (3.5), we have the following estimate

(3.9)
$$\int_{\mathfrak{M}(Y)} \left| f_2^*(\alpha)^2 \right| e(-h\alpha) d\alpha = \begin{cases} O(P_4^{\varepsilon}), & \text{when } 0 < |h| \leqslant 2X, \\ O(P_4^{\varepsilon}Y), & \text{when } h = 0. \end{cases}$$

Note that

$$\int_{\mathfrak{M}(Y)} |f_{2}^{*}(\alpha)|^{2} |e(-h\alpha)d\alpha$$

$$= \sum_{q \leqslant Y} \int_{|\beta| \leqslant \frac{Y}{qX}} q^{-2} \Big(\sum_{\substack{a=1\\ (a,q)=1}}^{q} |S_{2}(q,a)|^{2} e(-ha/q) \Big) |v_{2}(\beta)|^{2} e(-h\beta)d\beta$$

$$\ll P_{2}^{2} \sum_{q \leqslant Y} \int_{|\beta| \leqslant \frac{Y}{qX}} q^{-1}(q,h) (1+X|\beta|)^{-1} d\beta$$

$$\ll (\log X) \sum_{q \leqslant Y} q^{-1}(q,h).$$

The desired estimate (3.9) follows easily from above. For s = 4, we derive that

$$\int_{\mathfrak{m}_{3}} \left| f_{2}^{*}(\alpha)^{2} f_{4}(\alpha)^{2} \mathcal{E}(\alpha)^{2} \right| d\alpha$$

$$\leq \int_{\mathfrak{M}(Y)} \left| f_{2}^{*}(\alpha)^{2} f_{4}(\alpha)^{2} \mathcal{E}(\alpha)^{2} \right| d\alpha$$

$$= \sum_{\substack{n_{1}, n_{2} \in \mathcal{Z} \\ 1 \leqslant x_{1}, x_{2} \leqslant P_{4}}} \eta(n_{1}) \overline{\eta(n_{2})} \int_{\mathfrak{M}(Y)} \left| f_{2}^{*}(\alpha)^{2} \right| e(-(n_{1} - n_{2} + x_{1}^{4} - x_{2}^{4})\alpha) d\alpha$$

$$\ll P_{4}^{2+\varepsilon} Z^{2} + P_{4}^{\varepsilon} Y(P_{4}^{\varepsilon} Z^{2} + P_{4} Z)$$

$$\ll \left(P_{4}^{2+\varepsilon} + P_{4}^{3/2+\tau+\varepsilon} \psi(X)^{2} \right) Z^{2} + P_{4}^{5/2+\tau+\varepsilon} \psi(X)^{2} Z,$$

whence

$$\mathcal{K} \ll (P_4^{15/4+\tau/2+\varepsilon}\psi(X) + P_4^{7/2+\tau+\varepsilon}\psi(X)^2)Z + P_4^{4+\tau+\varepsilon}\psi(X)^2Z^{1/2}.$$

In particular, we have

$$\mathcal{K} \ll P_4^{4 + \frac{1}{4} + \varepsilon} Z^{1/2} + P_4^{4 - \tau + \varepsilon} \psi(X)^{-1} Z$$

provided that $\psi(X) \ll X^{1/64-\tau}$. For s=3, by (3.9) we have

$$\int_{\mathfrak{m}_3} \left| f_2^*(\alpha)^2 \mathcal{E}(\alpha)^2 \right| d\alpha \ll P_4^{\varepsilon} Z^2 + P_4^{3/2 + \tau + \varepsilon} \psi(X)^2 Z,$$

whence

$$\mathcal{K} \ll P_4^{11/4+\tau/2+\varepsilon} \psi(X) Z + P_4^{7/2+\tau+\varepsilon} \psi(X)^2 Z^{1/2}.$$

When $\psi(X) \ll X^{1/64-\tau}$, one has

$$\mathcal{K} \ll P_4^{4-\frac{1}{4}+\varepsilon} Z^{1/2} + P_4^{3-\tau+\varepsilon} \psi(X)^{-1} Z.$$

We conclude from above that

(3.10)
$$\mathcal{K} \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z^{1/2} + P_4^{s-\tau+\varepsilon} \psi(X)^{-1} Z.$$

By (3.8) and (3.10), we obtain

(3.11)
$$\mathcal{I}_3 \ll P_4^{4 - \frac{1}{4} + \frac{s - 3}{2} + \varepsilon} Z^{1/2} + P_4^{s - \tau + \varepsilon} \psi(X)^{-1} Z.$$

We complete the proof.

Lemma 3.4. Let \mathcal{I}_4 be defined in (3.1). Then we have

$$\mathcal{I}_4 \ll ZP_4^{s-(s-2)\nu/4+\varepsilon}$$
.

Proof. In view of (2.3) and (2.4), for $\alpha \in \mathfrak{M}_{P_4}(q, a)$, one has

$$f_4(\alpha) \ll P_4 w_4(q) (1 + X|\alpha - a/q|)^{-1/4} + P_4^{1/2+\varepsilon}$$

 $\ll P_4^{1+\varepsilon} w_4(q) (1 + X|\alpha - a/q|)^{-1/4},$

and

$$f_2(\alpha) \ll P_2 q^{-1/2} (1 + X|\alpha - a/q|)^{-1/2}$$

Therefore we obtain

$$\mathcal{I}_{4} \ll Z \sup_{\alpha \in \mathfrak{m}_{4}} |f_{4}(\alpha)|^{s-2} \int_{\mathfrak{M}(P_{4})} |f_{4}(\alpha)f_{2}(\alpha)|^{2} d\alpha$$

$$\ll Z P_{4}^{(s-2)(1-\nu/4)+\varepsilon} P_{4}^{2} P_{2}^{2} \sum_{q \leqslant P_{4}} w_{4}(q)^{2} \int_{|\beta| \leqslant \frac{P_{4}}{qX}} (1+X|\beta|)^{-3/2} d\beta$$

$$\ll Z P_{4}^{2+(s-2)(1-\nu/4)+\varepsilon} \sum_{q \leqslant P_{4}} w_{4}(q)^{2}.$$

In light of Lemma 2.4 by Kawada and Wooley [7], one can conclude that

(3.12)
$$\mathcal{I}_4 \ll Z P_4^{2+(s-2)(1-\nu/4)+\varepsilon} \ll Z P_4^{s-(s-2)\nu/4+\varepsilon}.$$

The desired estimate is established.

Proof of Theorem 1.1. We denote by $Z_s(X)$ the set of integers n with $X/2 < n \leq X$ for which the lower bound

$$\left| R_s(n) - c_s \Gamma(\frac{5}{4})^4 \mathfrak{S}_s(n) n^{s/4} \right| > n^{s/4} \psi(n)^{-1}$$

holds, and we abbreviate $\operatorname{card}(Z_s(X))$ to Z_s . It follows from (2.1) and Lemma 2.1 that for $n \in Z_s(X)$,

$$\left| \int_{\mathfrak{m}} f_2(\alpha)^2 f_4(\alpha)^s e(-n\alpha) d\alpha \right| \gg X^{s/4} \psi(X)^{-1},$$

whence

$$\sum_{n \in Z_s(X)} \left| \int_{\mathfrak{m}} f_2(\alpha)^2 f_4(\alpha)^s e(-n\alpha) d\alpha \right| \gg Z_s X^{s/4} \psi(X)^{-1}.$$

We choose complex numbers $\eta(n)$, with $|\eta(n)| = 1$, satisfying

$$\left| \int_{\mathfrak{m}} f_2(\alpha)^2 f_4(\alpha)^s e(-n\alpha) d\alpha \right| = \eta(n) \int_{\mathfrak{m}} f_2(\alpha)^2 f_4(\alpha)^s e(-n\alpha) d\alpha.$$

Then we define the exponential sum $\mathcal{E}_s(\alpha)$ by

$$\mathcal{E}_s(\alpha) = \sum_{n \in Z_s(X)} \eta(n) e(-n\alpha).$$

One finds that

$$(3.13) Z_s X^{s/4} \psi(X)^{-1} \ll \int_{\mathfrak{m}} \left| f_2(\alpha)^2 f_4(\alpha)^s \mathcal{E}_s(\alpha) \right| d\alpha.$$

Note that $\mathfrak{m} = \mathfrak{m}_1 \cup \mathfrak{m}_2 \cup \mathfrak{m}_3 \cup \mathfrak{m}_4$. Now we conclude from Lemmata 3.1-3.4 and (3.13) that

$$Z_s X^{s/4} \psi(X)^{-1} \ll P_4^{4-\frac{1}{4}+\frac{s-3}{2}+\varepsilon} Z_s^{1/2} + P_4^{s-\delta} \psi(X)^{-1} Z_s$$

for some sufficiently small positive number δ . Therefore we have

(3.14)
$$Z_s X^{s/4} \psi(X)^{-1} \ll X^{1-\frac{1}{16} + \frac{s-3}{8} + \varepsilon} Z_s^{1/2}.$$

The estimate (3.14) implies $Z_3 \ll X^{3/8+\varepsilon}\psi(X)^2$ and $Z_4 \ll X^{1/8+\varepsilon}\psi(X)^2$. The proof of Theorem 1.1 is completed by summing over dyadic intervals.

Acknowledgements. The author would like to thank the referees for many helpful comments and suggestions.

References

- [1] J. Brüdern, Sums of squares and higher powers, J. Lond. Math. Soc. (2) **35** (1987), 233–243.
- [2] J. Brüdern, A problem in additive number theory, Math. Proc. Cambridge Philos. Soc. **103** (1988), 27–33.
- [3] J. Brüdern and K. Kawada, The asymptotic formula in Waring's problem for one square and seventeen fifth powers, Monatshefte für Mathematik **162** (2011), 385–407.
- [4] J. B. Friedlander and T. D. Wooley, On Waring's problem: two squares and three biquadrates, Mathematika (in press), arXiv: 1211.1823.
- [5] D. R. Heath-Brown and D. I. Tolev, Lagrange's four squares theorem with one prime and three almost prime variables, J. Reine Angew. Math. 558 (2003), 159-224.
- [6] C. Hooley, On Waring's problem for two squares and three cubes, J. Reine Angew. Math. 328 (1981), 161-207.
- [7] K. Kawada and T. D. Wooley, On the Waring-Goldbach problem for fourth and fifth powers, Proc. Lond. Math. Soc. (3) 83 (2001), 1–50.
- [8] Yu. V. Linnik, Additive problems involving squares, cubes and almost primes, Acta Arith. 21 (1972), 413-422.
- [9] R. C. Vaughan, *The Hardy-Littlewood method*, 2nd ed. Cambridge University Press, Cambridge 1997.
- [10] T. D. Wooley, Slim exceptional sets for sums of four squares, Proc. Lond. Math. Soc. (3) 85 (2002), 1-21.
- [11] T. D. Wooley, Slim exceptional sets in Waring's problem: one square and five cubes, Quart. J. Math. 53 (2002), 111-118.

School of Mathematics, Hefei University of Technology, Heifei 230009, People's Republic of China

E-mail address: zhaolilu@gmail.com

٦