Effects of time-varying β in SNLS3 on constraining interacting dark energy models

Shuang Wang,^{1, *} Yong-Zhen Wang,^{1, [†](#page-0-1)} Jia-Jia Geng,^{1, [‡](#page-0-2)} and Xin Zhang^{§1, 2, [¶](#page-0-3)}

¹Department of Physics, College of Sciences, Northeastern University, Shenyang 110004, China

² Center for High Energy Physics, Peking University, Beijing 100080, China

It has been found that, for the Supernova Legacy Survey three-year (SNLS3) data, there is strong evidence for the redshift-evolution of color-luminosity parameter β . In this paper, adopting the w -cold-dark-matter ($wCDM$) model and considering its interacting extensions (with three kinds of interaction between dark sectors), we explore the evolution of β and its effects on parameter estimation. In addition to the SNLS3 data, we also use the latest Planck distance priors data, the galaxy clustering (GC) data extracted from sloan digital sky survey (SDSS) data release 7 (DR7) and baryon oscillation spectroscopic survey (BOSS), as well as the direct measurement of Hubble constant H_0 from the Hubble Space Telescope (HST) observation. We find that, for all the interacting dark energy (IDE) models, adding a parameter of β can reduce χ^2 by ~ 34 , indicating that a constant β is ruled out at 5.8 σ confidence level (CL). Furthermore, it is found that varying β can significantly change the fitting results of various cosmological parameters: for all the dark energy models considered in this paper, varying β yields a larger fractional CDM densities Ω_{c0} and a larger equation of state w; on the other side, varying β yields a smaller reduced Hubble constant h for the wCDM model, but has no impact on h for the three IDE models. This implies that there is a degeneracy between h and coupling parameter γ . Our work shows that the evolution of β is insensitive to the interaction between dark sectors, and then highlights the importance of considering β 's evolution in the cosmology fits.

PACS numbers: 98.80.-k, 98.80.Es, 95.36.+x Keywords: Cosmology, type Ia supernova, dark energy

I. INTRODUCTION

Cosmic acceleration is one of the biggest puzzles in modern cosmology [\[1–](#page-7-0)[7\]](#page-8-0). There are mainly two approaches to explain this extremely counterintuitive phenomenon: dark energy (DE) [\[8](#page-8-1)[–22\]](#page-9-0) and modified gravity (MG) [\[23–](#page-10-0)[30\]](#page-10-1). For recent reviews, see [\[31](#page-10-2)[–40\]](#page-10-3).

Cosmological observations are of essential importance to understanding cosmic acceleration, and one of the most important observations is Type Ia supernovae (SNe Ia) [\[41–](#page-10-4)[44\]](#page-10-5). In 2010, the Supernova Legacy Survey (SNLS) group released their three years data, i.e. SNLS3 dataset [\[45\]](#page-10-6). Soon after, using this dataset, Conley et al. [\[46\]](#page-10-7) and Sullivan et al. [\[47\]](#page-10-8) presented the SN-only cosmological results and the joint cosmological constraints, respectively. Unlike other supernova (SN) group, the SNLS team treated two important quantities, stretch-luminosity parameter α and color-luminosity parameter β of SNe Ia, as free model parameters.

There are many factors that can lead to systematic uncertainties of SNe Ia. One of the most important factors is the potential SN evolution, i.e. the possibility for the redshift evolution of α and β . The current studies show that α is consistent with a constant, but the hints for the evolution of β have been found in [\[48–](#page-10-9)[52\]](#page-10-10). For example, in [\[53\]](#page-10-11), using a linear $\beta(z) = \beta_0 + \beta_1 z$, Mohlabeng and Ralston studied the case of Union2.1 dataset and found that β deviates from a constant at 7σ confidence levels (CL). In [\[54\]](#page-10-12), Wang & Wang found that, for the SNLS3 data, β increases significantly with z at the 6σ CL; moreover, they proved that this conclusion is insensitive to the lightcurve fitter models, or the functional form of $\beta(z)$ assumed [\[54\]](#page-10-12). Therefore, the evolution of β is a common phenomenon for various SN datasets, and should be taken into account seriously.

It is very interesting to study the effects of a time-varying β on parameter estimation. In [\[55\]](#page-10-13), Wang, Li & Zhang explored this issue by considering the Λ-cold-dark-matter (ΛCDM) model, the wCDM model, and the Chevallier-Polarski-Linder (CPL) model. Soon after, Wang, Geng, Hu & Zhang [\[56\]](#page-10-14) studied the case of holographic dark energy (HDE) model, which is a physically plausible DE candidate based on the holographic principle. It is found that, for

[§] Corresponding author

[∗]Electronic address: wjysysgj@163.com

[†]Electronic address: [w˙avin@163.com](mailto:w_avin@163.com)

[‡]Electronic address: gengjiajia163@163.com

[¶]Electronic address: zhangxin@mail.neu.edu.cn

corresponding discussions to the case of interacting dark energy (IDE) models. In this paper, we explore the effects of a time-varying β on the cosmological constraints of the IDE model. Three kinds of interaction terms are taken into account. In addition to the SNLS3 data, we also use the Planck distance prior data [\[57\]](#page-10-15), the galaxy clustering (GC) data from SDSS DR7 [\[58\]](#page-10-16) and BOSS [\[59\]](#page-10-17), as well as the direct measurement of Hubble constant $H_0 = 73.8 \pm 2.4 \text{km/s/Mpc}$ from the Hubble Space Telescope (HST) observation [\[7\]](#page-8-0).

We describe our method in Sec. II, present our results in Sec. III, and conclude in Sec. IV. In this paper, we assume today's scale factor $a_0 = 1$, thus the redshift $z = a^{-1} - 1$. The subscript "0" always indicates the present value of the corresponding quantity, and the natural units are used.

II. METHODOLOGY

A. Theoretical models

In this paper, we consider a non-flat universe. The Friedmann equation can be written as

$$
3M_{pl}^2H^2 = \rho_c + \rho_{de} + \rho_r + \rho_b + \rho_k,\tag{1}
$$

where $M_{pl} \equiv 1/$ √ $8\pi G$ is the reduced Planck mass, ρ_c , ρ_{de} , ρ_r , ρ_b and ρ_k are the energy densities of CDM, DE, radiation, baryon and curvature, respectively. The reduced Hubble parameter $E(z) \equiv H(z)/H_0$ satisfies

$$
E^{2} = \Omega_{c0} \frac{\rho_{c}}{\rho_{c0}} + \Omega_{de0} \frac{\rho_{de}}{\rho_{de0}} + \Omega_{r0} \frac{\rho_{r}}{\rho_{r0}} + \Omega_{b0} \frac{\rho_{b}}{\rho_{b0}} + \Omega_{k0} \frac{\rho_{k}}{\rho_{k0}},
$$
\n(2)

where Ω_{c0} , Ω_{de0} , Ω_{ro} , Ω_{bo} and Ω_{ko} are the present fractional densities of CDM, DE, radiation, baryon and curvature, respectively. Since $\Omega_{c0} + \Omega_{de0} + \Omega_{b0} + \Omega_{b0} + \Omega_{k0} = 1$, we do not treat Ω_{de0} as an independent parameter in this paper. In addition, $\rho_r = \rho_{r0}(1+z)^4$, $\rho_b = \rho_{b0}(1+z)^3$, $\rho_k = \rho_{k0}(1+z)^2$, $\Omega_{r0} = \Omega_{m0}/(1+z_{eq})$, where $\Omega_{m0} = \Omega_{c0} + \Omega_{b0}$ and $z_{\text{eq}} = 2.5 \times 10^4 \Omega_{m0} h^2 (T_{\text{cmb}}/2.7 \text{ K})^{-4}$ (here we take $T_{\text{cmb}} = 2.7255 \text{ K}$).

Considering the interaction between dark sectors, the dynamical evolutions of CDM and DE become

$$
\dot{\rho}_c + 3H\rho_c = Q,\tag{3}
$$

$$
\dot{\rho}_{de} + 3H(\rho_{de} + p_{de}) = -Q,\tag{4}
$$

where the over dot denotes the derivative with respect to the cosmic time t, $p_{de} = w \rho_{de}$ is the pressure of DE, w is the equation of state of DE, and Q is the interaction term, which describes the energy transfer rate between CDM and DE. Notice that $a = \frac{1}{1+z}$ and $H = \frac{\dot{a}}{a}$, we have $\frac{d}{dt} = -H(1+z)\frac{d}{dz}$. Then Eqs. [\(3\)](#page-1-0) and [\(4\)](#page-1-0) can be rewritten as

$$
(1+z)\frac{d\rho_c}{dz} - 3\rho_c = -Q/H,\tag{5}
$$

$$
(1+z)\frac{d\rho_{de}}{dz} - 3(1+w)\rho_{de} = Q/H.
$$
\n(6)

The solutions of these two equations depend on the specific forms of Q.

So far, the microscopic origin of interaction between dark sectors is still a big puzzle to us. To study the issue of interaction, one has to write down the possible forms of Q by hand. In this paper we consider the following four cases:

$$
Q_0 = 0,\t\t(7)
$$

$$
Q_1 = 3\gamma H \rho_c,\tag{8}
$$

$$
Q_2 = 3\gamma H \rho_{de},\tag{9}
$$

$$
Q_3 = 3\gamma H \frac{\rho_c \rho_{de}}{\rho_c + \rho_{de}},\tag{10}
$$

where γ is a dimensionless coupling parameter describing the strength of interaction. Notice that the model with Q_0 denotes the case without dark sector interaction; the models with Q_1 and Q_2 are very popular, and both of them have been widely studied in the literature (see, e.g., $[60-63]$ $[60-63]$); the model with Q_3 is proposed in Ref. [\[64\]](#page-10-20), and it can solve

.

the early-time superhorizon instability and future unphysical CDM density problems at the same time. For simplicity, hereafter we call them wCDM model, IwCDM1 model, IwCDM2 model, and IwCDM3 model, respectively.

For the wCDM model, the reduced Hubble parameter $E(z) \equiv H(z)/H_0$ can be written as

$$
E(z) = (\Omega_{r0}(1+z)^4 + (\Omega_{c0} + \Omega_{b0})(1+z)^3 + \Omega_{k0}(1+z)^2 + \Omega_{de0}(1+z)^{3(1+w)})^{1/2}.
$$
 (11)

For the IwCDM1 model, Eq. (5) has a general solution

$$
\rho_c = \rho_{c0} (1+z)^{3(1-\gamma)}.
$$
\n(12)

Substituting Eq. [\(12\)](#page-2-0) into Eq. [\(6\)](#page-1-1) and using the initial condition $\rho_{de}(z=0) = \rho_{de0}$, we get

$$
\rho_{de} = \frac{\gamma \rho_{c0}}{\gamma + w} \left((1+z)^{3(1+w)} - (1+z)^{3(1-\gamma)} \right) + \rho_{de0} (1+z)^{3(1+w)}.
$$
\n(13)

Then, substituting Eqs. (12) and (13) into Eq. (2) , we obtain

$$
E(z) = \left(\Omega_{r0}(1+z)^4 + \Omega_{b0}(1+z)^3 + \Omega_{k0}(1+z)^2 + \Omega_{de0}(1+z)^{3(1+w)} + \Omega_{c0}\left(\frac{\gamma}{w+\gamma}(1+z)^{3(1+w)} + \frac{w}{w+\gamma}(1+z)^{3(1-\gamma)}\right)\right)^{1/2}
$$
\n(14)

For the IwCDM2 model, Eq. (6) has a general solution

$$
\rho_{de} = \rho_{de0} (1+z)^{3(1+w+\gamma)}.
$$
\n(15)

Substituting Eq. [\(15\)](#page-2-2) into Eq. [\(5\)](#page-1-1) and using the initial condition $\rho_c(z=0) = \rho_{c0}$, we get

$$
\rho_c = \rho_{c0}(1+z)^3 + \frac{\gamma \rho_{de0}}{w+\gamma}(1+z)^3 - \frac{\gamma \rho_{de0}}{w+\gamma}(1+z)^{3(1+w+\gamma)}.
$$
\n(16)

Then, substituting Eqs. (15) and (16) into Eq. (2) , we have

$$
E(z) = \left(\Omega_{r0}(1+z)^4 + (\Omega_{c0} + \Omega_{b0})(1+z)^3 + \Omega_{k0}(1+z)^2 + \Omega_{de0}\left(\frac{\gamma}{w+\gamma}(1+z)^3 + \frac{w}{w+\gamma}(1+z)^{3(1+w+\gamma)}\right)\right)^{1/2}.
$$
 (17)

For the IwCDM3 model, the energy densities of CDM and DE satisfy

$$
\rho_c = \rho_{c0} (1+z)^3 \left(\frac{\rho_{c0}}{\rho_{c0} + \rho_{de0}} + \frac{\rho_{de0}}{\rho_{c0} + \rho_{de0}} (1+z)^{3(w+\gamma)} \right)^{-\frac{\gamma}{w+\gamma}},\tag{18}
$$

$$
\rho_{de} = \rho_{de0}(1+z)^{3(1+w+\gamma)} \left(\frac{\rho_{c0}}{\rho_{c0} + \rho_{de0}} + \frac{\rho_{de0}}{\rho_{c0} + \rho_{de0}}(1+z)^{3(w+\gamma)}\right)^{-\frac{\gamma}{w+\gamma}}.
$$
\n(19)

Substituting Eqs. (18) and (19) into Eq. (2) , we get

$$
E(z) = (\Omega_{c0}C(z)(1+z)^3 + \Omega_{de0}C(z)(1+z)^{3(1+w+\gamma)} + \Omega_{r0}(1+z)^4 + \Omega_{b0}(1+z)^3 + \Omega_{k0}(1+z)^2)^{1/2},
$$
 (20)

where

$$
C(z) = \left(\frac{\Omega_{c0}}{\Omega_{c0} + \Omega_{de0}} + \frac{\Omega_{de0}}{\Omega_{c0} + \Omega_{de0}} (1+z)^{3(w+\gamma)}\right)^{-\frac{\gamma}{w+\gamma}}.
$$
\n(21)

Note that in Eqs. [\(11\)](#page-2-6), [\(14\)](#page-2-7), [\(17\)](#page-2-8), [\(20\)](#page-2-9), and [\(21\)](#page-2-10), Ω_{de0} is not an independent parameter, which is given by $\Omega_{de0} = 1 - \Omega_{c0} - \Omega_{b0} - \Omega_{r0} - \Omega_{k0}.$

B. Observational data

In this subsection, we introduce how to include the SNLS3 data into the χ^2 analysis.

For the SNLS3 sample, the observable is m_B , which is the rest-frame peak B-band magnitude of the SN. By considering three functional forms (linear case, quadratic case, and step function case), Wang & Wang [\[54\]](#page-10-12) showed that the evolutions of α and β are insensitive to functional form of α and β assumed. So in this paper, we just adopt a constant α and a linear $\beta(z) = \beta_0 + \beta_1 z$. Then, the predicted magnitude of an SN becomes

$$
m_{\text{mod}} = 5\log_{10}\mathcal{D}_L(z) - \alpha(s-1) + \beta(z)\mathcal{C} + \mathcal{M},\tag{22}
$$

where s and C are the stretch measure and the color measure for the SN light curve. Here $\mathcal M$ is a parameter representing some combination of SN absolute magnitude M and Hubble constant H_0 . It must be emphasized that, to include host-galaxy information in the cosmological fits, Conley et al. [\[46\]](#page-10-7) split the SNLS3 sample based on hostgalaxy stellar mass at $10^{10}M_{\odot}$, and made M to be different for the two samples. Therefore, unlike other SN samples, there are two values of M , M_1 and M_2 , for the SNLS3 data Moreover, Conley et al. removed M_1 and M_2 from cosmology-fits by analytically marginalizing over them (for more details, see the appendix C of [\[46\]](#page-10-7), as well as the the public code which is available at https://tspace.library.utoronto.ca/handle/1807/24512). In this paper, we just follow the recipe of Ref. [\[46\]](#page-10-7). The luminosity distance $\mathcal{D}_L(z)$ is defined as

$$
\mathcal{D}_L(z) \equiv H_0(1 + z_{\text{hel}})r(z),\tag{23}
$$

where z and z_{hel} are the CMB restframe and heliocentric redshifts of SN. In addition, the comoving distance $r(z)$ is given by

$$
r(z) = H_0^{-1} |\Omega_{k0}|^{-1/2} \sin \left(|\Omega_{k0}|^{1/2} \Gamma(z) \right), \tag{24}
$$

where $\Gamma(z) = \int_0^z \frac{dz'}{E(z')}$, and $\sinh(x) = \sin(x)$, x, $\sinh(x)$ for $\Omega_{k0} < 0$, $\Omega_{k0} = 0$, and $\Omega_{k0} > 0$ respectively.

For a set of N SNe with correlated errors, the χ^2 function is

$$
\chi_{SN}^2 = \Delta \mathbf{m}^T \cdot \mathbf{C}^{-1} \cdot \Delta \mathbf{m},\tag{25}
$$

where $\Delta m \equiv m_B - m_{\text{mod}}$ is a vector with N components, and C is the $N \times N$ covariance matrix of the SN, given by

$$
\mathbf{C} = \mathbf{D}_{\text{stat}} + \mathbf{C}_{\text{stat}} + \mathbf{C}_{\text{sys}}.\tag{26}
$$

 D_{stat} is the diagonal part of the statistical uncertainty, given by [\[46\]](#page-10-7)

$$
\mathbf{D}_{\text{stat},ii} = \sigma_{m_{B},i}^{2} + \sigma_{\text{int}}^{2} + \sigma_{\text{lensing}}^{2} + \sigma_{\text{host correction}}^{2} + \left[\frac{5(1+z_{i})}{z_{i}(1+z_{i}/2)\ln 10} \right]^{2} \sigma_{z,i}^{2} + \alpha^{2} \sigma_{s,i}^{2} + \beta(z_{i})^{2} \sigma_{c,i}^{2} + 2\alpha C_{m_{B}s,i} - 2\beta(z_{i}) C_{m_{B}c,i} - 2\alpha\beta(z_{i}) C_{s,c,i},
$$
\n(27)

where $C_{m_B,s,i}$, $C_{m_B,c,i}$, and $C_{s,c,i}$ are the covariances between m_B , s, and C for the *i*-th SN, $\beta_i = \beta(z_i)$ are the values of $\bar{\beta}$ for the *i*-th SN. Notice that $\sigma_{z,i}^2$ includes a peculiar velocity residual of 0.0005 (i.e., 150 km/s) added in quadrature. Following the Ref. [\[46\]](#page-10-7), we fix the intrinsic scatter σ_{int} to ensure that $\chi^2/dof = 1$. Varying σ_{int} could have a significant impact on parameter estimation, see [\[65\]](#page-10-21) for details.

We define $V \equiv C_{stat} + C_{sys}$, where C_{stat} and C_{sys} are the statistical and systematic covariance matrices, respectively. After treating β as a function of z, **V** is given in the form,

$$
\mathbf{V}_{ij} = V_{0,ij} + \alpha^2 V_{a,ij} + \beta_i \beta_j V_{b,ij} + \alpha V_{0a,ij} + \alpha V_{0a,ji} - \beta_j V_{0b,ij} - \beta_i V_{0b,ji} - \alpha \beta_j V_{ab,ij} - \alpha \beta_i V_{ab,ji}.
$$
 (28)

It must be stressed that, while V_0 , V_a , V_b , and V_{0a} are the same as the "normal" covariance matrices given by the SNLS data archive, V_{0b} , and V_{ab} are not the same as the ones given there. This is because the original matrices of SNLS3 are produced by assuming β is constant. We have used the V_{0b} and V_{ab} matrices for the "Combined" set that are applicable when varying $\beta(z)$ (A. Conley, private communication, 2013).

To improve the cosmological constraints, we also use some other cosmological observations, including the Planck distance prior data [\[57\]](#page-10-15), the galaxy clustering (GC) data extracted from SDSS DR7 [\[58\]](#page-10-16) and BOSS [\[59\]](#page-10-17), as well as the direct measurement of Hubble constant $H_0 = 73.8 \pm 2.4 \text{km/s/Mpc}$ from the HST observations [\[7\]](#page-8-0). For the details of including Planck and GC data into the χ^2 analysis, see Ref. [\[55\]](#page-10-13). Now the total χ^2 function is

$$
\chi^2 = \chi_{SN}^2 + \chi_{CMB}^2 + \chi_{GC}^2 + \chi_{H_0}^2.
$$
\n(29)

TABLE I: Fitting results for various constant β and linear $\beta(z)$ cases, where the SNe+CMB+GC+H₀ data are used.

	wCDM		IwCDM1		IwCDM2		IwCDM3	
Parameters	Const β	Linear $\beta(z)$	Const β	Linear $\beta(z)$	Const β	Linear $\beta(z)$	Const β	Linear $\beta(z)$
α	$1.444_{ -0.115}^{ +0.079}$	$1.423 + 0.087$ -0.095	$1.424 + 0.104$ -0.094	$1.398^{+0.110}_{-0.066}$	$1.427^{+0.096}_{-0.097}$	$1.421 + 0.084$ -0.101	$1.445 + 0.082$ -0.115	$1.393^{+0.121}_{-0.068}$
β_0	$3.251^{+0.113}_{-0.098}$	$1.518^{+0.326}_{-0.378}$	$3.272^{+0.087}_{-0.116}$	$1.438^{+0.367}_{-0.372}$	$3.275 + 0.084$ -0.112	$1.474^{+0.308}_{-0.369}$	$3.248 + 0.110$ -0.084	$1.505 + 0.292$ -0.402
β_1		$4.926^{+1.011}_{-0.869}$		$5.102^{+0.988}_{-0.924}$		$4.970 + 1.015$ -0.819		$4.886^{+1.191}_{-0.747}$
Ω_{c0}	$0.224_{-0.010}^{+0.010}$	$0.231 + 0.011$ -0.009	$0.232^{+0.012}_{-0.015}$	$0.244_{ -0.013}^{ +0.016}$	$0.226 + 0.012$ -0.013	$0.238^{+0.020}_{-0.012}$	$0.225 + 0.014$ -0.011	$0.244_{ -0.016}^{ +0.013}$
Ω_{b0}	$0.042^{+0.002}_{-0.002}$	$0.044_{0.002}^{+0.002}$ -0.002	$0.041 + 0.003$ -0.002	$0.040 + 0.003$ -0.002	$0.041_{-0.002}^{+0.003}$	$0.041_{ -0.002}^{ +0.003}$	$0.042^{+0.002}_{-0.002}$	$0.042_{-0.003}^{+0.002}$
Ω_{k0}	$0.00046^{+0.004}_{-0.003}$	$0.0032^{+0.0038}_{-0.0041}$	$0.0039^{+0.0044}_{-0.0061}$	$0.0095 + 0.0050$ -0.0059	$0.0061_{-0.0162}^{+0.0142}$	$0.0192^{+0.0180}_{-0.0165}$	$0.0046 + 0.0160$ -0.0131	$0.0194 + 0.0194$ -0.0159
γ			$-0.0028 + 0.0043$	$-0.0053^{+0.0035}$	$-0.0105 + 0.0310$ -0.0295	$-0.0322^{+0.0300}_{-0.0396}$	$^{-0.0198}^{+0.0613}_{-0.0752}$	$-0.0732^{+0.0684}_{-0.0984}$
\boldsymbol{w}	$-1.118 + 0.065$ -0.071	$-1.042 + 0.068$ -0.072	$-1.105 + 0.075$ $' = 0.069$	$-1.016 + 0.075$ -0.063	$-1.124^{+0.070}_{-0.062}$	$-1.052 + 0.070$ -0.068	$-1.116 + 0.059$ -0.072	$-1.038 + 0.068$ $' = 0.080$
\boldsymbol{h}	$0.725 + 0.014$ -0.014	$0.716 + 0.014$ -0.015	$0.739 + 0.019$ -0.023	$0.743 + 0.016$ -0.024	$0.734 + 0.018$ -0.025	$0.735_{-0.024}^{+0.017}$	$0.732^{+0.022}_{-0.021}$	$0.729 + 0.027$ -0.018
x_{min}^2	422.696	388.508	422.376	387.128	422.674	387.814	422.642	387.716

In addition, assuming the measurement errors are Gaussian, the likelihood function satisfies

$$
\mathcal{L} \propto e^{-\chi^2/2}, \quad \text{Likelihood} \equiv \mathcal{L}/\mathcal{L}_{max} = \mathcal{L}/\mathcal{L}(\chi^2_{min}). \tag{30}
$$

It should be mentioned that, in this paper we just use the purely geometric measurements, and do not consider the cosmological perturbations in the DE models. As analyzed in detail in Ref. [\[66\]](#page-10-22), adopting a new framework for calculating the perturbations, the cosmological perturbations will always be stable in all IDE models (For a related discussion concerning the stability, see Ref. [\[64\]](#page-10-20)). Therefore, the use of the Planck distance prior is sufficient for our purpose.

Finally, we perform an MCMC likelihood analysis [\[67\]](#page-10-23) to obtain $\mathcal{O}(10^6)$ samples for each model considered in this paper.

III. RESULTS

A. Evolution of β

In this subsection, we explore the evolution of β in the frame of IDE.

In Table [I,](#page-4-0) we list the fitting results for various constant β and linear $\beta(z)$ cases, where the SNe+CMB+GC+H₀ data are used. An obvious feature of this table is that varying β can significantly improve the fitting results: for all the models, adding a parameter of β can reduce the best-fit values of χ^2 by ~ 34. Based on the Wilk's theorem, 34 units of χ^2 is equivalent to a Gaussian fluctuation of 5.8 σ . This means that the result of $\beta_1=0$ is ruled out at 5.8 σ confidence level (CL) . As shown in Refs. [\[55\]](#page-10-13) and [\[56\]](#page-10-14), for the cases of various DE models (such as ΛCDM, wCDM, CPL, and HDE model) without interaction, β deviates from a constant at 6σ CL. Therefore, we further confirm the redshift-evolution of β for the SNLS3 data.

In Fig. [1,](#page-5-0) using the SNe+CMB+GC+H₀ data, we plot the 1σ confidence constraints of $\beta(z)$, for the wCDM model, the IwCDM1 model, the IwCDM2 model, and the IwCDM3 model. For comparison, we also plot the best-fit result of constant β case for the wCDM model. From this figure one can see that, the 1σ regions of $\beta(z)$ of all these models are almost overlapping. This shows that the evolution of β is independent of the interacting dark energy models. In addition, for all the models, $\beta(z)$ rapidly increases with z. This result is consistent with the results of Refs. [\[55\]](#page-10-13) and [\[56\]](#page-10-14), showing that the evolution of β is insensitive to dark energy models including those with interaction between dark sectors.

It should be pointed out that the evolutionary behaviors of $\beta(z)$ depends on the SN samples used. In [\[53\]](#page-10-11), Mohlabeng and Ralston found that, for the Union2.1 SN data, $\beta(z)$ decreases with z. This is similar to the case of Pan-STARRS1 SN data [\[52\]](#page-10-10).

It is interesting to study how different segments of the SNLS3 dataset give rise to different behavior of β . To do this, we perform the following test: (1) Per [\[46\]](#page-10-7), we evenly divide the redshift range $[0,1]$ into 9 bins and assume that both α and β are constant within each bin. (2) For each redshift bin, we make a small covariance matrix corresponding to only SNe in that bin. (3) Since we have already proved that the evolution of β is insensitive to dark energy models, per [\[46\]](#page-10-7), we just adopt a fixed cosmological background (a flat ΛCDM model with $\Omega_{m0} = 0.26$) to do this test. (4) We fit α and β separately for the 9 redshift bins. Based on the best-fit analysis, it is found that β is relatively flat till the 7th bin, and then it rapidly increases along with redshift z. In other words, the rapid increase of $\beta(z)$ is mainly due to the contributions from high-redshift $(z > 0.7)$ SN samples of SNLS3 dataset. It should be mentioned that, to keep this paper focus on its main purpose, here we just briefly present the conclusion, instead of describing all the detailed

FIG. 1: 1σ confidence constraints for the evolution of $\beta(z)$, given by the SNe+CMB+GC+H₀ data, for the wCDM model, the IwCDM1 model, the IwCDM2 model, and the IwCDM3 model. The solid black lines denote the wCDM model, the dashed red lines denote the IwCDM1 model, the dotted blue lines denote the IwCDM2 model, and the dashed-dotted pink lines denote the IwCDM3 model. To make a comparison, for the wCDM model, the best-fit result of constant β case is also plotted, shown as the horizontal dashed black line.

results of the test. To understand why high-redshift SNLS3 samples will yield this kind of evolutionary behavior of β , some numerical simulation studies may be needed. We will study this issue in future works.

B. Effects of time-varying β

In this subsection, we discuss the effects of varying β on parameter estimation of IDE models.

In Fig. [2,](#page-6-0) using $SNe+CMB+GC+H_0$ data, we plot the 1D marginalized probability distributions of Ω_{c0} , for all the models considered in this paper. We find that varying β yields a larger Ω_{c0} : for the constant β case, the best-fit results are $\Omega_{c0} = 0.224, 0.232, 0.226,$ and 0.225, for the wCDM, the IwCDM1, the IwCDM2, and the IwCDM3 model, respectively; while for the linear $\beta(z)$ case, the best-fit results are $\Omega_{c0} = 0.231, 0.244, 0.238,$ and 0.244, for the wCDM, the IwCDM1, the IwCDM2, and the IwCDM3 model, respectively. In addition, as shown in Refs. [\[55\]](#page-10-13) and [\[56\]](#page-10-14), for various DE models without interaction term, a time-varying β also yields a larger fractional matter density $\Omega_{m0} \equiv \Omega_{c0} + \Omega_{b0}$. Therefore, we can conclude that the effects of varying β on the present fractional matter density are insensitive to the interaction between dark sectors.

For all the models considered in this paper, the 1D marginalized probability distributions of w are plotted in Fig. [3.](#page-7-1) It is found that varying β yields a larger w: for the constant β case, $w = -1.118^{+0.065}_{-0.071}$, $-1.105^{+0.075}_{-0.069}$, $-1.124^{+0.070}_{-0.062}$, and $-1.116^{+0.059}_{-0.072}$, for the wCDM model, the IwCDM1 model, the IwCDM2 model, and the IwCDM3 model, respectively; while for the linear $\beta(z)$ case, $w = -1.042_{-0.072}^{+0.068}$, $-1.016_{-0.063}^{+0.075}$, $-1.052_{-0.068}^{+0.070}$, and $-1.038_{-0.080}^{+0.068}$, for the wCDM model, the IwCDM1 model, the IwCDM2 model, and the IwCDM3 model, respectively. In other words, $w < -1$ is preferred at more than 1σ CL for the constant β case, while w is consistent with -1 at 1σ CL for the linear $\beta(z)$ case. This means that, compared to the constant β case, the results from varying β case are in better agreement with a cosmological constant. This conclusion is consistent with the noninteracting cases [\[55,](#page-10-13) [56\]](#page-10-14), showing that the effects of varying β on w are insensitive to the interaction between dark sectors.

In Fig. [4,](#page-8-2) we plot the 1D marginalized probability distributions of h , for all the models considered in this paper. It can be seen that, for the wCDM model, varying β yields a smaller h; this result is consistent with the noninteracting cases [\[55,](#page-10-13) [56\]](#page-10-14). However, for all the IDE models, the 1D distribution results of h of the linear β case are almost same with those of the constant β case. In other words, once considering the interaction between dark sectors, varying β will not change the fitting results of h. This result is quite different from the results of Fig. [2](#page-6-0) and Fig. [3,](#page-7-1) showing that there is a degeneracy between h and γ .

Next, we turn to the constraints on interaction parameter γ . In Fig. [5,](#page-9-1) we plot 1σ and 2σ confidence contours for $\{\gamma, h\}$, for all the IDE models. Again, one can see that varying β has no impact on h. In contrary, varying β yields a smaller γ : for the constant β case, the best-fit results are $\gamma = -0.0028, -0.0105,$ and -0.0198 , for the IwCDM1 model, the IwCDM2 model, and the IwCDM3 model, respectively; while for the linear $\beta(z)$ case, the best-fit results are $\gamma = -0.0053, -0.0322,$ and -0.0732 , for the IwCDM1 model, the IwCDM2 model, and the IwCDM3 model, respectively. In other words, $\gamma < 0$ is slightly more favored in the linear $\beta(z)$ case. This means that energy will

FIG. 2: The 1D marginalized probability distributions of Ω_{c0} , given by the SNe+CMB+GC+H₀ data, for the wCDM model and the three IDE models. Both the results of constant β (dashed black lines) and linear $\beta(z)$ (solid red lines) cases are presented.

transfer from dark matter to dark energy. In addition, we find that γ and h are anti-correlated, showing that there is a degeneracy between h and γ .

In Fig. [6,](#page-9-2) to make a visual comparison among three interaction forms, we plot the 2σ confidence contours for $\{\Omega_{c0}, \gamma\}$, based on the linear $\beta(z)$ case, for all the IDE models. From this figure one can see that, γ is tightly constrained in the IwCDM1 model; in contrary, γ cannot be well constrained in the IwCDM2 and IwCDM3 models. This result is consistent with the result of [\[68\]](#page-10-24), in which only the constant β case was considered.

IV. DISCUSSION AND SUMMARY

In recent years, more and more SNe Ia have been discovered, and the systematic errors of SNe Ia have drawn more and more attentions. One of the most important systematic uncertainties for SNe Ia is the potential SN evolution. The hints for the evolution of β have been found [\[48–](#page-10-9)[52\]](#page-10-10). For examples, Mohlabeng and Ralston [\[53\]](#page-10-11) studied the case of Union2.1 and found that β deviates from a constant at 7 σ CL. In [\[54\]](#page-10-12), Wang & Wang found that, for the SNLS3 data, β increases significantly with z at the 6 σ CL; moreover, they proved that this conclusion is insensitive to the lightcurve fitter models, or the functional form of $\beta(z)$ assumed [\[54\]](#page-10-12).

It is clear that a time-varying β will have significant impact on parameter estimation. Adopting a constant α and a linear $\beta(z) = \beta_0 + \beta_1 z$, Wang, Li & Zhang [\[55\]](#page-10-13) explored this issue by considering the ΛCDM model, the wCDM model, and the CPL model. Soon after, Wang, Geng, Hu & Zhang [\[56\]](#page-10-14) studied this issue in the frame of HDE model, which is a physically plausible DE candidate based on the holographic principle. It is found that, for all these DE models, β deviates from a constant at 6σ CL; in addition, considering the evolution of β is helpful in reducing the tension between SN and other cosmological observations. It should be pointed out that, in principle there is always an important possibility that DE directly interacts with CDM. This factor was not considered in Refs. [\[55\]](#page-10-13) and [\[56\]](#page-10-14).

In this paper, we extend the corresponding discussions to the case of IDE model. To perform the cosmology-fits, the wCDM model is adopted. Moreover, three kinds of interaction forms are considered: $Q_1 = 3\gamma H \rho_c$, $Q_2 = 3\gamma H \rho_{de}$, and $Q_3 = 3\gamma H \frac{\rho_c \rho_{de}}{\rho_c + \rho_{de}}$. In addition to the SNLS3 SN data, we also use the Planck distance priors data, the GC data extracted from SDSS DR7 and BOSS, as well as the direct measurement of Hubble constant from the HST observation.

We further confirm the redshift-evolution of β for the SNLS3 data: for all the IDE models, adding a parameter of β can reduce χ^2 by ~ 34 , indicating that $\beta_1 = 0$ is ruled out at 5.8 σ CL. In addition, we find that the 1 σ regions of $\beta(z)$

FIG. 3: The 1D marginalized probability distributions of w, given by the $SNe+CMB+GC+H_0$ data, for the wCDM model and the three IDE models. Both the results of constant β (dashed black lines) and linear $\beta(z)$ (solid red lines) cases are presented.

of all these models are almost overlapping, showing that the evolution of β is insensitive to the interaction between dark sectors. These results further verify the importance of considering the evolution of β in the cosmology-fits.

Furthermore, we find that a time-varying β has significant effects on the results of parameter estimation: for all the models considered in this paper, varying β yields a larger Ω_{c0} and a larger w; on the other side, varying β yields a smaller h for the wCDM model, while varying β has no influence on h for the three IDE models. Moreover, we find that γ and h are anti-correlated, showing that there is a degeneracy between h and γ . In addition, we find that γ is tightly constrained in the IwCDM1 model, but cannot be well constrained in the IwCDM2 and IwCDM3 models.

In all, these results show that the evolution of β is insensitive to the interaction between dark sectors, and highlight the importance of considering β 's evolution in the cosmology fits.

So far, only the effects of varying β on DE models are considered. It is of great interest to study the effects of varying β on parameter estimation in MG models. In addition, some other factors, such as the evolution of σ_{int} [\[65\]](#page-10-21), may also cause the systematic uncertainties of SNe Ia. These issues will be studied in future works.

Acknowledgments

We thank the referee for valuable suggestions, which help us to improve this work. We also thank Dr. Daniel Scolnic for helpful discussions. We are grateful to Dr. Alex Conley for providing us with the SNLS3 covariance matrices that allow redshift-dependent β . We acknowledge the use of CosmoMC. SW is supported by the Fundamental Research Funds for the Central Universities under Grant No. N130305007. XZ is supported by the National Natural Science Foundation of China under Grant No. 11175042 and the Fundamental Research Funds for the Central Universities under Grant No. N120505003.

- [1] A. G. Riess et al., AJ. 116, 1009 (1998); S. Perlmutter et al., ApJ. 517, 565 (1999).
- [2] D. N. Spergel et al., ApJS 148, 175 (2003); C. L. Bennet et al., ApJS. 148, 1 (2003); D. N. Spergel et al., ApJS 170, 377 (2007); L. Page et al., ApJS 170, 335 (2007); G. Hinshaw et al., ApJS 170, 263 (2007).
- [3] M. Tegmark et al., Phys. Rev. D 69, 103501 (2004); ApJ 606, 702 (2004); Phys. Rev. D 74, 123507 (2006).
- [4] E. Komatsu et al., ApJS. 180, 330 (2009); E. Komatsu et al., ApJS. 192, 18 (2011).

FIG. 4: The 1D marginalized probability distributions of h, given by the $SNe+CMB+GCH_0$ data, for the wCDM model and the three IDE models. Both the results of constant β (dashed black lines) and linear $\beta(z)$ (solid red lines) cases are presented.

- [5] W. J. Percival et al., MNRAS 401, 2148 (2010); A. G. Sanchez, et al., [arXiv:1203.6616,](http://arxiv.org/abs/1203.6616) MNRAS accepted.
- [6] M. Drinkwater *et al.*, MNRAS 401 , 1429 (2010); C. Blake *et al.*, [arXiv:1108.2635,](http://arxiv.org/abs/1108.2635) MNRAS accepted.
- [7] A. G. Riess et al., ApJ. **730**, 119 (2011).
- [8] P. J. E. Peebles and B. Ratra, ApJ 325, L17 (1988); C. Wetterich, Nucl. Phys. B 302, 668 (1988); R. R. Caldwell, R. Dave and P. J. Steinhardt, Phys. Rev. Lett. 80, 1582 (1998); I. Zlatev, L. Wang and P. J. Steinhardt, Phys. Rev. Lett. 82, 896 (1999).
- [9] R. R. Caldwell, Phys. Lett. B 545, 23 (2002); S. M. Carroll, M. Hoffman and M. Trodden, Phys. Rev. D 68, 023509 (2003); R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, Phys. Rev. Lett. 91, 071301 (2003); X. Zhang, Eur. Phys. J. C 59, 755 (2009); X. Zhang, Eur. Phys. J. C 60, 661 (2009); X. D. Li et al., Sci. China Phys. Mech. Astron. 55, 1330 (2012).
- [10] C. Armendariz-Picon, T. Damour and V. Mukhanov, Phys. Lett. B 458, 209 (1999); C. Armendariz-Picon, V. Mukhanov and P. J. Steinhardt, Phys. Rev. D 63, 103510 (2001); T. Chiba, T. Okabe and M. Yamaguchi, Phys. Rev. D 62, 023511 $(2000).$
- [11] A. Y. Kamenshchik, U. Moschella and V. Pasquier, Phys. Lett. B 511, 265 (2001); M. C. Bento, O. Bertolami and A. A. Sen, Phys. Rev. D 66, 043507 (2002).
- [12] X. Zhang, F. Q. Wu and J. Zhang, JCAP 01, 003 (2006); K. Liao, Y. Pan and Z. H. Zhu, Res. Astron. Astrophys. 13, 159 (2013).
- [13] T. Padmanabhan, Phys. Rev. D 66, 021301 (2002); J. S. Bagla, H. K. Jassal, and T. Padmanabhan, Phys. Rev. D 67, 063504 (2003).
- [14] M. Li, Phys. Lett. B 603, 1 (2004); Q. G. Huang and M. Li, JCAP 08, 013 (2004). X. Zhang and F. Q. Wu, Phys. Rev. D 72, 043524 (2005); Z. Chang, F. Q. Wu and X. Zhang, Phys. Lett. B 633, 14 (2006); X. Zhang and F. Q. Wu, Phys. Rev. D 76, 023502 (2007); J. -F. Zhang, X. Zhang and H. -Y. Liu, Eur. Phys. J. C 52, 693 (2007); M. Li, C. S. Lin and Y. Wang, JCAP 05, 023 (2008); M. Li, X. D. Li, S. Wang and X. Zhang, JCAP 06, 036 (2009); X. Zhang, Phys. Lett. B 683, 81 (2010); Y. H. Li, S. Wang, X. D. Li and X. Zhang, JCAP 02, 033 (2013).
- [15] H. Wei, R. G. Cai, and D. F. Zeng, Class. Quant. Grav. 22, 3189 (2005); H. Wei, and R. G. Cai, Phys. Rev. D 72, 123507 (2005); H. Wei, N. Tang, and S. N. Zhang, Phys. Rev. D75, 043009 (2007).
- [16] W. Zhao and Y. Zhang, Class. Quant. Grav. 23, 3405 (2006); T. Y. Xia and Y. Zhang, Phys. Lett. B 656, 19 (2007); S. Wang, Y. Zhang and T. Y. Xia, JCAP 10, 037 (2008); S. Wang and Y. Zhang, Phys. Lett. B 669, 201 (2008).
- [17] X. Zhang, Phys. Lett. B 648, 1 (2007); X. Zhang, Phys. Rev. D 74, 103505 (2006); J. Zhang, X. Zhang and H. Liu, Phys. Lett. B 651, 84 (2007); J. Zhang, X. Zhang and H. Liu, Eur. Phys. J. C 54, 303 (2008); X. Zhang, Phys. Rev. D 79, 103509 (2009).
- [18] D. Comelli, M. Pietroni and A. Riotto, Phys. Lett. B 571, 115 (2003); X. Zhang, Mod. Phys. Lett. A 20, 2575 (2005); X. Zhang, Phys. Lett. B 611, 1 (2005).
- [19] J. A. Frieman, C. T. Hill, A. Stebbins, and I. Waga, Phys. Rev. Lett. 75, 2077 (1995); M. Chevallier and D. Polarski, Int.

FIG. 5: The 1 σ and 2σ confidence contours for $\{\gamma, h\}$, for the three IDE models. Both the results of constant β (red regions) and linear $\beta(z)$ (dark cyan regions) cases are presented.

FIG. 6: The 2σ confidence contours for $\{\Omega_{c0}, \gamma\}$, based on the linear $\beta(z)$ case, for the IwCDM1 model (solid black line), the IwCDM2 model (dashed red line), and the IwCDM3 model (dotted blue line).

J. Mod. Phys. D 10, 213 (2001); E. V. Linder, Phys. Rev. Lett. 90 091301 (2003); D. Huterer and G. Starkman, Phys. Rev. Lett. 90, 031301 (2003); D. Huterer and A. Cooray, Phys. Rev. D 71, 023506 (2005).

- [20] Y. Wang and M. Tegmark, Phys. Rev. Lett. 92, 241302 (2004); Y. Wang and M. Tegmark, Phys. Rev. D 71, 103513 (2005); Y. Wang, and K. Freese, Phys. Lett. B 632, 449 (2006); Y. Wang and P. Mukherjee, ApJ. 650, 1 (2006); Y. Wang and P. Mukherjee, Phys. Rev. D 76, 103533 (2007); Y. Wang,Phys. Rev. D 78, 123532 (2008).
- [21] U. Alam, V. Sahni, T. D. Saini and A. A. Starobinsky, Mon. Not. Roy. Astron. Soc. 344, 1057 (2003); U. Alam, V. Sahni, T. D. Saini and A. A. Starobinsky, Mon. Not. Roy. Astron. Soc. 354, 275 (2004); A. Shafieloo, U. Alam, V. Sahni and A. A. Starobinsky, Mon. Not. Roy. Astron. Soc. 366, 1081 (2006); U. Alam, V. Sahni and A. A. Starobinsky, JCAP 02, 011 (2007); V. Sahni, A. Shafieloo and A. A. Starobinsky, Phys. Rev. D 78, 103502, (2008); A. Shafieloo, V. Sahni and A. A. Starobinsky, Phys. Rev. D 80, 101301(R) (2009); A. Shafieloo, V. Sahni and A. A. Starobinsky, Phys. Rev. D 86, 103527 (2012).
- [22] J. F. Zhang, X. Zhang and H. Y. Liu, Mod. Phys. Lett. A 23, 139 (2008); Q. G. Huang, M. Li, X. D. Li and S. Wang,

Phys. Rev. D 80, 083515 (2009); S. Wang, X. D. Li and M. Li, Phys. Rev. D 82, 103006 (2010); M. Li, X. D. Li and X. Zhang, Sci. China Phys. Mech. Astron. 53, 1631 (2010); S. Wang, X. D. Li and M. Li, Phys. Rev. D 83, 023010 (2011); Y. H. Li and X. Zhang, Eur. Phys. J. C 71, 1700 (2011); X. D. Li et al., JCAP 07, 011 (2011); J. Z. Ma and X. Zhang, Phys. Lett. B 699, 233 (2011); H. Li and X. Zhang, Phys. Lett. B 713, 160 (2012).

- [23] V. Sahni and S. Habib, Phys. Rev. Lett. 81, 1766 (1998).
- [24] L. Parker and A. Raval, Phys. Rev. D 60, 063512 (1999).
- [25] G. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B 485, 208 (2000).
- [26] S. Nojiri, S. D. Odintsov, and M. Sasaki, Phys. Rev. D 71, 123509 (2005).
- [27] A. Nicolis, R. Rattazzi, and E. Trincherini, Phys. Rev. D 79 , 064036 (2009).
- [28] W. Hu and I. Sawicki, Phys. Rev. D 76 , 064004 (2007); A. A. Starobinsky, J. Exp. Theor. Phys. Lett. 86, (2007) 157.
- [29] G. R. Bengochea and R. Ferraro, Phys. Rev. D 79, 124019 (2009); E. V. Linder, Phys. Rev. D 81, (2010) 127301.
- [30] T. Harko, F. S. N. Lobo, S. Nojiri and S. D. Odintsov, Phys. Rev. D 84, 024020 (2011).
- [31] E. J. Copeland, M. Sami and S. Tsujikawa, Int. J. Mod. Phys. D 15, 1753 (2006).
- [32] J. Frieman, M. Turner and D. Huterer, Ann. Rev. Astron. Astrophys 46, 385 (2008).
- [33] E. V. Linder, Rept. Prog. Phys. 71, 056901 (2008).
- [34] R. R. Caldwell and M. Kamionkowski, Ann. Rev. Nucl. Part. Sci. 59, 397 (2009).
- [35] J.-P. Uzan, [arXiv:0908.2243.](http://arxiv.org/abs/0908.2243)
- [36] S. Tsujikawa, [arXiv:1004.1493.](http://arxiv.org/abs/1004.1493)
- [37] S. Nojiri and S. D. Odintsov, Phys. Rept. 505, 59 (2011).
- [38] M. Li, X. D. Li, S. Wang and Y. Wang, Commun. Theor. Phys. 56, 525 (2011).
- [39] T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, Phys. Rept. 513, 1 (2012).
- [40] Y. Wang, *Dark Energy*, Wiley-VCH (2010).
- [41] M. Kowalski, *et al.*, ApJ. **686**, 749 (2008).
- [42] M. Hicken, et al., ApJ. 700, 1097 (2009); M. Hicken, et al., ApJ. 700, 331 (2009).
- [43] R. Amanullah, et al., ApJ. **716**, 712 (2010).
- [44] N. Suzuki, et al., ApJ **746**, 85 (2012).
- [45] J. Guy, et al., A&A, **523**, 7 (2010).
- [46] A. Conley, et al., ApJS. 192 1 (2011).
- [47] M. Sullivan, et al., [arXiv:1104.1444.](http://arxiv.org/abs/1104.1444)
- [48] Astier, *et al.*, Astron. Astrophys. **447**, 31 (2006).
- [49] R. Kessler, et al., ApJS. 185, 32 (2009).
- [50] Marriner, *et al.*, [arXiv:1107.4631.](http://arxiv.org/abs/1107.4631)
- [51] D. Scolnic, *et al.*, [arXiv:1306.4050,](http://arxiv.org/abs/1306.4050) ApJ in press.
- [52] D. Scolnic, et al., [arXiv:1310.3824.](http://arxiv.org/abs/1310.3824)
- [53] G.Mohlabeng and J. Ralston, [arXiv:1303.0580.](http://arxiv.org/abs/1303.0580)
- [54] S. Wang and Y. Wang, Phys. Rev. D 88, 043511 (2013).
- [55] S. Wang, Y. H. Li and X. Zhang, Phys. Rev. D 89, 063524 (2014).
- [56] S. Wang, J. J. Geng, Y. L. Hu and X. Zhang, [arXiv:1312.0184.](http://arxiv.org/abs/1312.0184)
- [57] Y. Wang and S. Wang, Phys. Rev. D 88, 043522 (2013).
- [58] C. H. Chuang and Y. Wang, MNRAS, 426, 226 (2012).
- [59] C. H. Chuang, et al., [arXiv:1303.4486.](http://arxiv.org/abs/1303.4486)
- [60] G. R. Farrar and P. J. E. Peebles, Astrophys. J. 604, 1 (2004); G. Olivares, F. Atrio-Barandela, and D. Pavon, Phys. Rev. D 71, 063523 (2005); T. Koivisto, Phys. Rev. D 72, 043516 (2005); H. M. Sadjadi and M. Alimohammadi, Phys. Rev. D 74, 103007 (2006); Z. K. Guo, N. Ohta, and S. Tsujikawa, Phys. Rev. D 76, 023508 (2007).
- [61] C. G. Boehmer, G. Caldera-Cabral, R. Lazkoz, and R. Maartens, Phys. Rev. D 78, 023505 (2008); M. Quartin, M. O. Calvao, S. E. Joras, R. R. R. Reis, and I. Waga, JCAP 05, 007 (2008); J. Valiviita, E. Majerotto, and R. Maartens, J. Cosmol. Astropart. Phys. 07 020 (2008); R. Bean, E. E. Flanagan, I. Laszlo, and M. Trodden, Phys. Rev. D 78, 123514 (2008); S. Chongchitnan, Phys. Rev. D 79, 043522 (2009); B. M. Jackson, A. Taylor, and A. Berera, Phys. Rev. D 79, 043526 (2009).
- [62] J. Zhang, H. Liu, and X. Zhang, Phys. Lett. B 659, 26 (2008); M. Li, X. -D. Li, S. Wang, Y. Wang and X. Zhang, JCAP 0912, 014 (2009); L. Zhang, J. Cui, J. Zhang and X. Zhang, Int. J. Mod. Phys. D 19, 21 (2010); J. Cui and X. Zhang, Phys. Lett. B 690, 233 (2010); Y. Li, J. Ma, J. Cui, Z. Wang and X. Zhang, Sci. China Phys. Mech. Astron. 54, 1367 (2011); T. -F. Fu, J. -F. Zhang, J. -Q. Chen and X. Zhang, Eur. Phys. J. C 72, 1932 (2012); Z. Zhang, S. Li, X. -D. Li, X. Zhang and M. Li, JCAP 1206, 009 (2012); T. Clemson, K. Koyama, G. B. Zhao, R. Maartens, and J. Valiviita, Phys. Rev. D 85, 043007 (2012); J. Zhang, L. Zhao and X. Zhang, Sci. China Phys. Mech. Astron. 57, 387 (2014).
- [63] J. H. He and B. Wang, JCAP. 06, 010 (2008); J. H. He, B. Wang, and P. Zhang, Phys. Rev. D 80, 063530 (2009); J. H. He, B. Wang, E. Abdalla, and D. Pavon, JCAP. 12, 022 (2010); X. D. Xu, B. Wang, and E. Abdalla, Phys. Rev. D 85, 083513 (2012); X. D. Xu, B. Wang, P. Zhang, and F. Atrio-Barandela, JCAP. 12, 001 (2013).
- [64] Y. H. Li and X. Zhang, Phys. Rev. D 89, 083009 (2014).
- [65] A. Kim, [arXiv:1101.3513;](http://arxiv.org/abs/1101.3513) J. Marriner, et al., [arXiv:1107.4631.](http://arxiv.org/abs/1107.4631)
- [66] Y. H. Li, J. F. Zhang and X. Zhang, [arXiv:1404.5220.](http://arxiv.org/abs/1404.5220)
- [67] A. Lewis and S. Bridle, Phys. Rev. D 66, 103511 (2002).
- [68] J. J. Geng, J. F. Zhang and X. Zhang, JCAP 07, 006 (2014).