

# Asymptotic Theory of Bayes Factor in Stochastic Differential Equations: Part II

Trisha Maitra and Sourabh Bhattacharya\*

## Abstract

The problem of model selection in the context of a system of stochastic differential equations (*SDE*'s) has not been touched upon in the literature. Indeed, properties of Bayes factors have not been studied even in single *SDE* based model comparison problems.

In this article, we first develop an asymptotic theory of Bayes factors when two *SDE*'s are compared, assuming the time domain expands. Using this we then develop an asymptotic theory of Bayes factors when systems of *SDE*'s are compared, assuming that the number of equations in each system, as well as the time domain, increase indefinitely. Our asymptotic theory covers situations when the observed processes associated with the *SDE*'s are independently and identically distributed (*iid*), as well as when they are independently but not identically distributed (*non-iid*). Quite importantly, we allow inclusion of available time-dependent covariate information into each *SDE* through a multiplicative factor of the drift function in a random effects set-up; different initial values for the *SDE*'s are also permitted.

Thus, our general model-selection framework includes simultaneously the variable selection problem associated with time-varying covariates, as well as choice of the part of the drift function free of covariates. It is to be noted that given that the underlying process is wholly observed, the diffusion coefficient becomes known, and hence is not involved in the model selection problem.

For both *iid* and *non-iid* set-ups we establish almost sure exponential convergence of the Bayes factor. As we show, the Bayes factor is inconsistent for comparing individual *SDE*'s, in the sense that the log-Bayes factor converges only in expectation, while the relevant variance does not converge to zero. Nevertheless, it has been possible to exploit this result to establish almost sure exponential convergence of the Bayes factor when, in addition, the number of individuals are also allowed to increase indefinitely.

We carry out simulated and real data analyses to demonstrate that Bayes factor is a suitable candidate for covariate selection in our *SDE* models even in non-asymptotic situations.

**Keywords:** *Bayes factor consistency; Kullback-Leibler divergence; Martingale; Stochastic differential equations; Time-dependent covariates and random effects; Variable selection.*

## 1 Introduction

Stochastic differential equations (*SDE*'s) have important standing in statistical applications where “within” subject variability is caused by some random component varying continuously in time. It also seems worthwhile to incorporate available time-dependent covariate information into the subject-wise *SDE*'s. Apart from the covariates there may also be random effects associated with the individuals, which may be useful in modeling variabilities between the individuals.

*SDE*-based models with time-dependent covariates are considered in Oravec *et al.* (2011), Overgaard *et al.* (2005), Leander *et al.* (2015); moreover, Oravec *et al.* (2011) analyse their covariate-based *SDE* model in the hierarchical Bayesian paradigm. In the literature, random effects *SDE* models without covariates seem to be more popular than those based on covariates. A brief overview of random effects *SDE* models is provided in Delattre *et al.* (2013) who undertake theoretical and classical asymptotic investigation of a class of random effects models based on *SDE*'s. Specifically, they model the *i*-th individual by

$$dX_i(t) = b(X_i(t), \phi_i)dt + \sigma(X_i(t))dW_i(t), \quad (1.1)$$

where, for  $i = 1, \dots, n$ ,  $X_i(0) = x^i$  is the initial value of the stochastic process  $X_i(t)$ , which is assumed to be continuously observed on the time interval  $[0, T_i]$ ;  $T_i > 0$  assumed to be known. The function  $b(x, \varphi)$ , which is the drift function, is a known, real-valued function on  $\mathbb{R} \times \mathbb{R}^d$  ( $\mathbb{R}$  is the real

---

\*Trisha Maitra is a PhD student and Sourabh Bhattacharya is an Associate Professor in Interdisciplinary Statistical Research Unit, Indian Statistical Institute, 203, B. T. Road, Kolkata 700108. Corresponding e-mail: sourabh@isical.ac.in.

line and  $d$  is the dimension), and the function  $\sigma : \mathbb{R} \mapsto \mathbb{R}$  is the known diffusion coefficient. The  $SDE$ 's given by (1.1) are driven by independent standard Wiener processes  $\{W_i(\cdot); i = 1, \dots, n\}$ , and  $\{\phi_i; i = 1, \dots, n\}$ , which are to be interpreted as the random effect parameters associated with the  $n$  individuals, which are assumed by Delattre *et al.* (2013) to be independent of the Brownian motions and independently and identically distributed (*iid*) random variables with some common distribution. For the sake of convenience Delattre *et al.* (2013) (see also Maitra and Bhattacharya (2016) and Maitra and Bhattacharya (2015)) assume  $b(x, \phi_i) = \phi_i b(x)$ . Thus, the random effect is a multiplicative factor of the drift function. In this work, we generalize this to a random effects  $SDE$  set-up consisting of time-dependent covariates.

Note that model selection constitutes an important part of research in both Bayesian and classical paradigms; see, for example, Dey *et al.* (2000), Jiang (2007), Claeskens and Hjort (2008), Müller *et al.* (2013). In the case of  $SDE$ -based mixed effects models as well, model selection constitutes an important issue involving the choice of the drift function and selection of the appropriate subset of (time-dependent) covariates. Here Bayes factors are expected to play the central role as their effectiveness in model selection in complex problems is well-established (see, for example, Kass and Raftery (1995) for a good account of Bayes factors). Unavailability of closed form expressions in the traditional  $SDE$  set-ups usually prompt usage of numerical approximations based on Markov chain Monte Carlo or related criteria such as the Akaike Information Criterion (Akaike (1973)) and Bayes Information Criterion (Schwarz (1978)). For details, see, for example, Fuchs (2013), Iacus (2008). But we are not aware of any research existing in the literature that attempts to address covariate selection in  $SDE$ 's.

We are also not aware of any existing literature on asymptotic investigation of Bayes factors in the  $SDE$  context although Sivaganesan and Lingham (2002) present some asymptotic investigation of intrinsic and fractional Bayes factors in the context of three specific diffusion models. The only investigation available in this context seems to be that of Maitra and Bhattacharya (2018), who model a multiplicative part of the drift function using time-varying covariates, and address Bayes factor asymptotics in a general set-up consisting of the covariate selection problem as well as selection of the part of the drift function independent of the covariates. Different initial values and domains of observations pertaining to different individuals, are also considered in their set-up. Assuming that only the number of individuals increase without bound, Maitra and Bhattacharya (2018) establish almost sure exponential convergence of Bayes factor in both *iid* and non-*iid* situations. Here we recall that the *iid* set-up is the case when there is no covariate associated with the model and when the initial values and the domains of observations are the same for every individual. The non-*iid* set-up, on the other hand, consists of time-varying covariates, different initial values and domains of observations; in this work we also consider random effects. Thus, unlike the *iid* case, here the model selection problem also deals with covariate selection apart from selection of the part of the drift functions free of the covariates.

In this article, we prove almost sure exponential convergence of the relevant Bayes factors in both *iid* and non-*iid* cases, assuming that the number of individuals, as well as the domains of observations, increase without bound. Hence, for our current purpose, the asymptotic theory developed by Maitra and Bhattacharya (2018) when only the number of individuals tends to infinity, is clearly inapplicable. Indeed, incorporation of random effects is asymptotically feasible only in our current asymptotic framework; Maitra and Bhattacharya (2018) elucidate that inclusion of random effects does not make sense asymptotically unless the domains of observations are also increased indefinitely. Also, only our current asymptotic framework allows different sets of time-dependent covariates for different individuals.

It is important to remark that the diffusion coefficient becomes known once the continuous process is completely observed; see Roberts and Stramer (2001). Hence, following Maitra and Bhattacharya (2018) we assume that the diffusion coefficient is known, and is not involved in the model selection problem.

We begin by establishing an asymptotic theory of Bayes factor for two competing individual  $SDE$ 's, and then extend the theory to systems of  $SDE$ 's. In this context it is important to draw attention to the fact that even this relatively simple problem of comparing any two individual  $SDE$ 's using Bayes factors has not yet been considered in the literature. Our investigation in this simpler case, however,

faced with an apparently negative result; the associated Bayes factor failed to be consistent in the sense that the relevant variance failed to converge to zero, even though convergence of the log-Bayes factor in expectation is ensured. Despite this, we have been able to utilise this result to establish almost sure exponential convergence of the Bayes factor when the number of individuals are also allowed to increase indefinitely.

The rest of our article is structured as follows. We begin with formalization of our set-up in Section 2, while we provide the necessary assumptions and results in Section 3. In Section 4 we investigate the asymptotics of Bayes factor for comparing two individual *SDE*'s. We illustrate our results with a special case in Section 5. In Section 6 we exploit the asymptotic theory of Bayes factors developed for comparing individual *SDE*'s to construct a convergence theory of Bayes factors comparing systems of *SDE*'s in both *iid* and non-*iid* cases. In Section 7 we carry out two simulation studies to demonstrate that Bayes factor yields the correct set of covariates in our *SDE* models even in non-asymptotic cases, and in Section 8, we model a real, company-wise national stock exchange data set, using a system of *SDE*'s, each consisting of a plausible set of covariates, and obtain the best possible sets of covariate combinations for the companies, using Bayes factor. We summarize our contributions and provide concluding remarks in Section 9.

## 2 Formalization of the model selection problem in the *SDE* set-up when $n \rightarrow \infty$ and $T_i \rightarrow \infty$ for every $i$

That the systems considered by us are well-defined and the exact likelihoods are computable, are guaranteed by assumption (H2'') in Section 3. For our purpose we consider the filtration  $(\mathcal{F}_t^W, t \geq 0)$ , where  $\mathcal{F}_t^W = \sigma(W_i(s), s \leq t)$ . Each process  $W_i$  is a  $(\mathcal{F}_t^W, t \geq 0)$ -adapted Brownian motion.

Here we consider the set-up where, for  $i = 1, 2, \dots, n$ ,

$$dX_i(t) = \phi_{i, \xi_0^{(i)}}(t) b_{\beta_0^{(i)}}(t, X_i(t)) dt + \sigma(t, X_i(t)) dW_i(t) \quad (2.1)$$

and

$$dX_i(t) = \phi_{i, \xi_1^{(i)}}(t) b_{\beta_1^{(i)}}(t, X_i(t)) dt + \sigma(t, X_i(t)) dW_i(t), \quad (2.2)$$

where,  $X_i(0) = x^i$  is the initial value of the stochastic process  $X_i(t)$ , which is assumed to be continuously observed on the time interval  $[0, T_i]$ ;  $T_i > 0$ . We consider (2.1) as representing the true model and (2.2) is any other model.

It is useful to remark that we must analyze the same data set with respect to two different models for the purpose of model selection. Hence, even though the distribution of the underlying stochastic process under the two models are different, for notational convenience we denote the process by  $X_i(t)$  under both the models, relying on the context and the model-specific parameters to naturally clarify the distinction.

### 2.1 Inclusion of time-dependent covariates

We model  $\phi_{i, \xi_j^{(i)}}(t)$  for  $j = 0, 1$ , and  $i = 1, \dots, n$ , as

$$\phi_{i, \xi_j^{(i)}}(t) = \phi_{i, \xi_j^{(i)}}(\mathbf{z}_i(t)) = \xi_{0j}^{(i)} + \xi_{1j}^{(i)} g_1(z_{i1}(t)) + \xi_{2j}^{(i)} g_2(z_{i2}(t)) + \dots + \xi_{pj}^{(i)} g_p(z_{ip}(t)), \quad (2.3)$$

where  $\mathbf{z}_i(t) = (z_{i1}(t), z_{i2}(t), \dots, z_{ip}(t))$  is the set of available covariate information corresponding to the  $i$ -th individual, depending upon time  $t$ . Following Maitra and Bhattacharya (2018) we assume  $\mathbf{z}_i(t)$  is continuous in  $t$ ,  $z_{il}(t) \in \mathcal{Z}_l$  where  $\mathcal{Z}_l$  is compact and  $g_l : \mathcal{Z}_l \rightarrow \mathbb{R}$  is continuous, for  $l = 1, \dots, p$ . We let  $\mathcal{Z} = \mathcal{Z}_1 \times \dots \times \mathcal{Z}_p$ , and  $\mathfrak{Z} = \{\mathbf{z}(t) \in \mathcal{Z} : t \in [0, \infty) \text{ such that } \mathbf{z}(t) \text{ is continuous in } t\}$ . Hence,  $\mathbf{z}_i \in \mathfrak{Z}$  for all  $i$ .

## 2.2 The random effects set-up

In (2.1),  $\theta_0^{(i)} = (\beta_0^{(i)}, \xi_{00}^{(i)}, \xi_{10}^{(i)}, \dots, \xi_{p0}^{(i)}) = (\beta_0^{(i)}, \xi_0^{(i)})$  stands for the true parameters, and  $\theta_1^{(i)} = (\beta_1^{(i)}, \xi_{01}^{(i)}, \xi_{11}^{(i)}, \dots, \xi_{p1}^{(i)}) = (\beta_1^{(i)}, \xi_1^{(i)})$  are the parameters associated with (2.2). Let  $\theta_j^{(i)} \in \Theta = \mathfrak{B} \times \Gamma$  for all  $i$ , where both  $\mathfrak{B}$  and  $\Gamma$  are compact spaces. We also assume that for  $i = 1, 2, \dots, n$ ,

$$\theta_1^{(i)} \stackrel{iid}{\sim} \pi,$$

where  $\pi$  is some specified distribution on  $\Theta$ .

Hence, the above describes a random effects set-up. Observe that if  $\xi_{lj}^{(i)} = 0$  for  $l = 1, \dots, p$ , and for  $i = 1, \dots, n$ , then it reduces to the random effects model of Delattre *et al.* (2013), showing that the latter is a special case of our model.

As is well-known, even though the term ‘‘prior’’ is not appropriate for the random effects coefficients, operationally there is no difference between a prior and a distribution for random effects in the Bayesian paradigm. Somewhat abusing the terminology, we continue to refer to the distribution of the *iid* random effects coefficients,  $\pi$ , as the relevant prior.

## 2.3 Covariate and drift function selection

The key difference between our current model selection idea and that of Maitra and Bhattacharya (2018) is that here, for every individual, there is an independent model selection problem. In other words, for each  $i$ , one needs to choose between  $\theta_0^{(i)}$  and  $\theta_1^{(i)}$ . This involves selection of perhaps different sets of covariates for different  $i$  with respect to the coefficients  $\xi_j^{(i)}$ , and different drift functions  $b_{\beta_j^{(i)}}$ .

Obviously, the dimensions of  $\xi_0^{(i)}$  and  $\xi_1^{(i)}$  are allowed to differ for each  $i$ ; likewise, for every  $i$ , the dimensions of  $\beta_0^{(i)}$  and  $\beta_1^{(i)}$  may be different as well. Thus, from this perspective, our current model selection framework appears to be more general compared to that of Maitra and Bhattacharya (2018), who consider the same set of parameters  $\xi_j$  and  $\beta_j$  for all the individuals, allowing only a fixed set of covariates for every subject.

## 2.4 Form of the Bayes factor in our set-up

For  $j = 0, 1$ , we first define the following quantities:

$$U_{i, \theta_j^{(i)}} = \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) b_{\beta_j^{(i)}}(s, X_i(s))}{\sigma^2(s, X_i(s))} dX_i(s), \quad V_{i, \theta_j^{(i)}} = \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}^2(s) b_{\beta_j^{(i)}}^2(s, X_i(s))}{\sigma^2(s, X_i(s))} ds \quad (2.4)$$

for  $j = 0, 1$  and  $i = 1, \dots, n$ .

Let  $\mathcal{C}_{T_i}$  denote the space of real continuous functions  $(x(t), t \in [0, T_i])$  defined on  $[0, T_i]$ , endowed with the  $\sigma$ -field  $\mathcal{C}_{T_i}$  associated with the topology of uniform convergence on  $[0, T_i]$ . We consider the distribution  $P_j^{x_i, T_i, z_i}$  on  $(\mathcal{C}_{T_i}, \mathcal{C}_{T_i})$  of  $(X_i(t), t \in [0, T_i])$  given by (2.1) and (2.2) for  $j = 0, 1$ . We choose the dominating measure  $P_i$  as the distribution of (2.1) and (2.2) with null drift. So, for  $j = 0, 1$ ,

$$\frac{dP_j^{x_i, T_i, z_i}}{dP_i} = f_{i, \theta_j^{(i)}}(X_i) = \exp \left( U_{i, \theta_j^{(i)}} - \frac{V_{i, \theta_j^{(i)}}}{2} \right), \quad (2.5)$$

where  $f_{i, \theta_0^{(i)}}(X_i)$  denotes the true density and  $f_{i, \theta_1^{(i)}}(X_i)$  stands for the other density associated with the modeled SDE.

For each  $i = 1, \dots, n$ , letting  $X_{i, a, b}$  denote the  $i$ -th process observed on  $[a, b]$  for any  $0 \leq a < b <$

$\infty$ ,

$$I_{x^i, T_i, z_i} = \int_{\Theta} \frac{f_{i, \theta_1^{(i)}}(X_{i,0, T_i})}{f_{i, \theta_0^{(i)}}(X_{i,0, T_i})} \pi(\theta_1^{(i)}) d\theta_1^{(i)} \quad (2.6)$$

denotes the Bayes factor associated with the  $i$ -th equation of the above two systems of equations. Assuming that the  $SDE$ 's (2.1) and (2.2) are independent for  $i = 1, \dots, n$ ,

$$I_{n, T_1, \dots, T_n} = \prod_{i=1}^n I_{x^i, T_i, z_i}$$

is the Bayes factor comparing the entire systems of  $SDE$ 's (2.1) and (2.2).

Comparisons between a collection of different models using Bayes factor, none of which may be the true model, is expected to favour that model which minimizes the Kullback-Leibler divergence from the true model.

## 2.5 The *iid* and the non-*iid* cases

We are interested in studying the properties of  $I_{n, T_1, \dots, T_n}$  in both *iid* and non-*iid* cases when  $n \rightarrow \infty$  and  $T_i \rightarrow \infty$ . In the *iid* set-up, we assume that  $x^i = x$ ,  $T_i = T$  and  $\theta_j^{(i)} = (\beta_j^{(i)}, \xi_{0j}^{(i)})$ , for  $i = 1, \dots, n$  and  $j = 0, 1$ . In the non-*iid* case we relax these assumptions. However, for simplicity, we assume  $T_i = T$  for each  $i$ , even in the non-*iid* set-up, so that in our asymptotic framework we study convergence of

$$\tilde{I}_{n, T} = \prod_{i=1}^n I_{i, T}, \quad (2.7)$$

as  $n \rightarrow \infty$  and  $T \rightarrow \infty$ , where  $I_{i, T} = I_{x^i, T, z_i}$ .

## 2.6 A key relation between $U_{i, \theta_j^{(i)}}$ and $V_{i, \theta_j^{(i)}}$ in the context of model selection using Bayes factors

An useful relation between  $U_{i, \theta_j^{(i)}}$  and  $V_{i, \theta_j^{(i)}}$  which we will often make use of in this paper is as follows.

$$\begin{aligned} U_{i, \theta_j^{(i)}} &= \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) b_{\beta_j^{(i)}}(X_i(s))}{\sigma^2(X_i(s))} dX_i(s) \\ &= \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) b_{\beta_j^{(i)}}(X_i(s))}{\sigma^2(X_i(s))} \left[ \phi_{i, \xi_0^{(i)}}(s) b_{\beta_0^{(i)}}(X_i(s)) ds + \sigma(X_i(s)) dW_i(s) \right] \\ &= \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) \phi_{i, \xi_0^{(i)}}(s) b_{\beta_j^{(i)}}(X_i(s)) b_{\beta_0^{(i)}}(X_i(s))}{\sigma^2(X_i(s))} ds + \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) b_{\beta_j^{(i)}}(X_i(s))}{\sigma(X_i(s))} dW_i(s) \\ &= V_{i, \theta_0^{(i)}, \theta_j^{(i)}} + \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) b_{\beta_j^{(i)}}(X_i(s))}{\sigma(X_i(s))} dW_i(s), \end{aligned} \quad (2.8)$$

with

$$V_{i, \theta_0^{(i)}, \theta_j^{(i)}} = \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) \phi_{i, \xi_0^{(i)}}(s) b_{\beta_j^{(i)}}(X_i(s)) b_{\beta_0^{(i)}}(X_i(s))}{\sigma^2(X_i(s))} ds. \quad (2.9)$$

Note that  $V_{i, \theta_0^{(i)}} = V_{i, \theta_0^{(i)}, \theta_0^{(i)}}$  and  $V_{i, \theta_1^{(i)}} = V_{i, \theta_1^{(i)}, \theta_1^{(i)}}$ . Also note that, for  $j = 0, 1$ , for each  $i$ ,

$$E_{\theta_0^{(i)}} \left[ \int_0^{T_i} \frac{\phi_{i, \xi_j^{(i)}}(s) b_{\beta_j^{(i)}}(X_i(s))}{\sigma(X_i(s))} dW_i(s) \right] = 0, \quad (2.10)$$

so that  $E_{\theta_0^{(i)}} \left( U_{i, \theta_j^{(i)}} \right) = E_{\theta_0^{(i)}} \left( V_{i, \theta_0^{(i)}, \theta_j^{(i)}} \right)$ .

### 3 Requisite assumptions and results for the asymptotic theory of Bayes factor when $n \rightarrow \infty$ and $T \rightarrow \infty$

All our following assumptions and results are true for each  $i$ , in particular true for each  $\beta_j^{(i)}, \xi_j^{(i)}$  and consequently for  $U_{i, \theta_j^{(i)}}, V_{i, \theta_j^{(i)}}, V_{i, \theta_0^{(i)}, \theta_j^{(i)}}$ . For the sake of notational simplicity we provide all the assumptions and results without mentioning  $i$  at every stage. We make the following assumptions:

(H1'') The parameter space  $\Theta = \mathfrak{B} \times \Gamma$  such that  $\Gamma$  and  $\mathfrak{B}$  are compact.

(H2'') For  $j = 0, 1$ , given any  $s, \beta_j, b_{\beta_j}(s, \cdot), \sigma(s, \cdot)$  are  $C^1$  on  $\mathbb{R}$ ; we also assume that  $b_{\beta_j}^2(s, x) \leq K_1(1 + x^2 + \|\beta_j\|^2)$  and  $\sigma^2(x) \leq K_2(1 + x^2)$  for all  $s \in [0, T], x \in \mathbb{R}$ , for some  $K_1, K_2 > 0$ . By (H1'') it follows as before that for  $s \in [0, T], b_{\beta_j}^2(s, x) \leq K(1 + x^2)$  and  $\sigma^2(s, x) \leq K(1 + x^2)$  for all  $x \in \mathbb{R}$ , for some  $K > 0$ .

Because of (H2'') it follows from Theorem 4.4 of Mao (2011), page 61, that for all  $T > 0$ , and any  $k \geq 2$ ,

$$E \left( \sup_{s \in [0, T]} |X_i(s)|^k \right) \leq \left( 1 + 3^{k-1} E |X_i(0)|^k \right) \exp(\tilde{\vartheta} T), \quad (3.1)$$

where

$$\tilde{\vartheta} = \frac{1}{6} (18K)^{\frac{k}{2}} T^{\frac{k-2}{2}} \left[ T^{\frac{k}{2}} + \left( \frac{k^3}{2(k-1)} \right)^{\frac{k}{2}} \right].$$

Specifically, for any  $k \geq 2$ , we can write, as  $T \rightarrow \infty$ ,

$$E \left( \sup_{s \in [0, T]} |X(s)|^k \right) = o \left( \exp \{ T^{k+1} \} \right). \quad (3.2)$$

We further assume the following conditions.

(H3'')  $b_{\beta_j}(s, x)$  is continuous in  $(x, \beta_j)$ .

(H4'') For  $s \in [0, T]$  and  $j = 0, 1$ ,  $\frac{b_{\beta_j}^2(s, x)}{\sigma^2(s, x)}$  and  $\frac{b_{\beta_1}(s, x)b_{\beta_0}(s, x)}{\sigma^2(s, x)}$  satisfy the following:

$$\begin{aligned} \kappa_j(\beta_j) + \frac{K_{\beta_j} (1 + x^2 + \|\beta_j\|^2)}{c_j + \exp(T^5)} &< \frac{b_{\beta_j}^2(s, x)}{\sigma^2(s, x)} \\ &< \kappa_j(\beta_j) + \frac{M_{\beta_j} (1 + x^2 + \|\beta_j\|^2)}{d_j + \exp(T^5)}, \end{aligned} \quad (3.3)$$

and

$$\begin{aligned} \bar{\kappa}(\beta_0, \beta_1) + \frac{K_{\beta_0, \beta_1} (1 + x^2 + \|\beta_0\|^2 + \|\beta_1\|^2)}{\bar{c} + \exp(T^5)} &< \frac{b_{\beta_1}(s, x)b_{\beta_0}(s, x)}{\sigma^2(s, x)} \\ &< \bar{\kappa}(\beta_0, \beta_1) + \frac{M_{\beta_0, \beta_1} (1 + x^2 + \|\beta_0\|^2 + \|\beta_1\|^2)}{\bar{d} + \exp(T^5)}, \end{aligned} \quad (3.4)$$

where  $0 < c_j, d_j, \bar{c}, \bar{d} < \infty$ , are some constants;  $\kappa_j(\beta_j)$  are positive, continuous functions of  $\beta_j$ ;  $\bar{\kappa}(\beta_0, \beta_1)$  is a continuous function of  $(\beta_0, \beta_1)$ ;  $K_{\beta_j}, M_{\beta_j}$  are continuous in  $\beta_j$ , for  $j = 0, 1$ , and  $K_{\beta_0, \beta_1}, M_{\beta_0, \beta_1}$  are continuous in  $(\beta_0, \beta_1)$ .

(H5'') (i) We assume that  $\mathcal{Z} = \mathbf{Z}_1 \times \mathbf{Z}_2 \times \cdots \times \mathbf{Z}_p$  is the space of the covariates where  $\mathbf{Z}_l$  is compact for  $l = 1, \dots, p$ , and for every  $t \geq 0$ ,  $\mathbf{z}_i(t) = (z_{i1}(t), z_{i2}(t), \dots, z_{ip}(t)) \in \mathcal{Z}$  for  $i = 1, \dots, n$ . Also, we assume that  $\mathbf{z}_i(t)$  are continuous in  $t$  for every  $i$ , so that  $\mathbf{z}_i \in \mathfrak{Z}$ , for every  $i$ .

(ii) For  $j = 0, 1$ , and for  $t \geq 0$ , we assume that the vector of covariates  $\mathbf{z}_i(t)$  is related to the  $i$ -th SDE of the  $j$ -th model via

$$\phi_{i, \xi_j^{(i)}}(t) = \phi_{i, \xi_j^{(i)}}(\mathbf{z}_i(t)) = \xi_{0j}^{(i)} + \sum_{l=1}^p \xi_{lj}^{(i)} g_l(\mathbf{z}_i(t)),$$

where, for  $l = 1, \dots, p$ ,  $g_l : \mathbf{Z}_l \rightarrow \mathbb{R}$  is continuous. Notationally, when reference to the  $i$ -th individual is self-explanatory, we shall denote the function  $\xi_{0j} + \sum_{l=1}^p \xi_{lj} g_l$  by  $\phi_{\xi_j}$ .

(iii) For  $l = 1, 2, \dots, p$ , for  $i = 1, \dots, n$ , and for  $t \geq 0$ ,

$$\frac{1}{n} \sum_{i=1}^n g_l(z_{il}(t)) \rightarrow c_l(t), \quad (3.5)$$

and

$$\frac{1}{n} \sum_{i=1}^n g_l(z_{il}(t)) g_m(z_{im}(t)) \rightarrow c_l(t) c_m(t), \quad (3.6)$$

as  $n \rightarrow \infty$ , where  $\{c_l(t) : t \geq 0\}$  are real constants for  $l = 1, \dots, p$ .

(iv) For  $l = 1, 2, \dots, p$ , and for  $i = 1, \dots, n$ ,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T g_l(z_{il}(s)) ds = \bar{c}_{il}^{(1)} \quad (3.7)$$

and

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T g_l(z_{il}(s)) g_m(z_{im}(s)) ds = \bar{c}_{ilm}^{(2)}, \quad (3.8)$$

where  $\bar{c}_{il}^{(1)}$  and  $\bar{c}_{ilm}^{(2)}$  are real constants.

**Remark 1** Observe that although (H4'') is seemingly restrictive in the sense that the ratios  $\frac{b_{\beta_j}^2(s, x)}{\sigma^2(s, x)}$  and  $\frac{b_{\beta_1}(s, x) b_{\beta_0}(s, x)}{\sigma^2(s, x)}$  are approximately independent of the underlying stochastic process, assumption (H5'') attempts to compensate for the restrictions by providing a rich structure to  $\phi_{\xi_j}$  consisting of covariate information varying continuously with time. Hence, assumption (H4'') need not be viewed as restrictive.

Maitra and Bhattacharya (2018) argue that (3.5) and (3.6) hold if one assumes that for  $i = 1, \dots, n$ , and  $l = 1, \dots, p$ , the covariates  $z_{il}$  are observed realizations of stochastic processes that are *iid* for  $i = 1, \dots, n$ , for all  $l = 1, \dots, p$ , and that for  $l \neq m$ , the processes generating  $z_{il}$  and  $z_{im}$  are independent. In other words, although we assume the covariates to be non-random, in essence, it may be assumed  $g_l(z_{il}(t))$  and  $g_m(z_{im}(t))$  are uncorrelated for  $l \neq m$ .

In order that (H5'') (iv) holds, one needs to further assume that the relevant stochastic processes converge to appropriate stationary distributions. For example,  $z_{il}(t)$  may be realizations of Markov processes which are irreducible (with respect to some appropriate measure), aperiodic, positive recurrent and possess invariant distributions; see, for example, Kontoyiannis and Meyn (2003).

It follows from (H5'') (iv), that,

$$\begin{aligned}
& \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi_{i, \xi_j^{(i)}}(z_i(s)) ds \\
&= \xi_{0j}^{(i)} + \sum_{l=1}^p \xi_{lj}^{(i)} \bar{c}_{il}^{(1)} \\
&= \bar{\phi}_{i, \xi_j^{(i)}}^{(1)} \text{ (say),}
\end{aligned} \tag{3.9}$$

$$\begin{aligned}
& \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi_{i, \xi_j^{(i)}}^2(z_i(s)) ds \\
&= \left\{ \xi_{0j}^{(i)} \right\}^2 + 2\xi_{0j}^{(i)} \sum_{l=1}^p \xi_{lj}^{(i)} \bar{c}_{il}^{(1)} + \sum_{l=1}^p \sum_{m=1}^p \xi_{lj}^{(i)} \xi_{mj}^{(i)} \bar{c}_{ilm}^{(2)} \\
&= \bar{\phi}_{i, \xi_j^{(i)}}^{(2)} \text{ (say),}
\end{aligned} \tag{3.10}$$

and

$$\begin{aligned}
& \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \phi_{i, \xi_0^{(i)}}(z_i(s)) \phi_{i, \xi_1^{(i)}}(z_i(s)) ds \\
&= \xi_{00}^{(i)} \xi_{01}^{(i)} + \xi_{00}^{(i)} \sum_{l=1}^p \xi_{l1}^{(i)} \bar{c}_{il}^{(1)} + \xi_{01}^{(i)} \sum_{l=1}^p \xi_{l0}^{(i)} \bar{c}_{il}^{(1)} + \sum_{l=1}^p \sum_{m=1}^p \xi_{l0}^{(i)} \xi_{m1}^{(i)} \bar{c}_{ilm}^{(2)} \\
&= \bar{\phi}_{i, \xi_0^{(i)}, \xi_1^{(i)}}^{(2)} \text{ (say).}
\end{aligned} \tag{3.11}$$

When  $i$  is clear from the context, we shall often use the notations  $\bar{\phi}_{\xi_j}^{(1)}$ ,  $\bar{\phi}_{\xi_j}^{(2)}$  and  $\bar{\phi}_{\xi_0, \xi_1}^{(2)}$ .

Note that, (3.9), (3.10) and (3.11) are limits of expectations with respect to the uniform distribution on  $[0, T]$ . Hence, by the Cauchy-Schwartz inequality it follows that

$$\bar{\phi}_{\xi_0, \xi_1}^{(2)} \leq \sqrt{\bar{\phi}_{\xi_0}^{(2)} \bar{\phi}_{\xi_1}^{(2)}}. \tag{3.12}$$

The following lemmas will be useful in our proceedings. The proofs of these lemmas are provided in sections S-1, S-2 and S-3 of the supplement.

**Lemma 2** *The limits  $\bar{\phi}_{\xi_1}^{(1)}$ ,  $\bar{\phi}_{\xi_1}^{(2)}$  and  $\bar{\phi}_{\xi_0, \xi_1}^{(2)}$  are continuous in  $\xi_1$ .*



**Lemma 3** Assume  $(H1'')$  –  $(H5'')$ . Then, the following hold:

$$(i) \quad E_{\theta_0} \left( \frac{V_{\theta_j, T}}{T} \right) \rightarrow \bar{\phi}_{\xi_j}^{(2)} \kappa_j(\beta_j); j = 0, 1. \quad (3.13)$$

$$(ii) \quad E_{\theta_0} \left( \frac{V_{\theta_0, \theta_1, T}}{T} \right) \rightarrow \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1). \quad (3.14)$$

$$(iii) \quad E_{\theta_0} \left( \frac{U_{\theta_0, T}}{T} \right) \rightarrow \bar{\phi}_{\xi_0}^{(2)} \kappa_0(\beta_0). \quad (3.15)$$

$$(iv) \quad E_{\theta_0} \left( \frac{U_{\theta_1, T}}{T} \right) \rightarrow \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1). \quad (3.16)$$

$$(v) \quad \frac{1}{T} \int_0^T \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} dW(s) \xrightarrow{a.s.} 0; j = 0, 1, \quad (3.17)$$

$$(vi) \quad \frac{V_{\theta_j, T}}{T} \xrightarrow{a.s.} \bar{\phi}_{\xi_j}^{(2)} \kappa_j(\beta_j); j = 0, 1, \quad (3.18)$$

$$(vii) \quad \frac{V_{\theta_0, \theta_1, T}}{T} \xrightarrow{a.s.} \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1), \quad (3.19)$$

$$(viii) \quad \frac{U_{\theta_0, T}}{T} \xrightarrow{a.s.} \bar{\phi}_{\xi_0}^{(2)} \kappa_0(\beta_0), \quad (3.20)$$

$$(ix) \quad \frac{U_{\theta_1, T}}{T} \xrightarrow{a.s.} \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1), \quad (3.21)$$

In the above, “ $\xrightarrow{a.s.}$ ” denotes convergence “almost surely” as  $T \rightarrow \infty$  with respect to  $X$  (under  $\theta_0$ ), and the expectations are also with respect to  $X$  (under  $\theta_0$ ).

**Lemma 4** Assume  $(H1'')$  –  $(H5'')$ . Then, the following holds:

$$\bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1) \leq \sqrt{\bar{\phi}_{\xi_0}^{(2)} \bar{\phi}_{\xi_1}^{(2)}} \times \sqrt{\kappa_0(\beta_0) \kappa_1(\beta_1)}. \quad (3.22)$$

## 4 Convergence of Bayes factor with respect to time when two individual $SDE$ 's are compared

From the system of  $SDE$ 's defined by (2.1) and (2.2) we now consider the  $i$ -th individual only. To avoid notational complexity we denote  $X_i$  simply by  $X$ . Consequently,  $\phi_{i, \xi_j}(t)$  and  $T_i$  will be denoted by  $\phi_{\xi_j}(t)$  and  $T$ , respectively. In connection with the  $i$ -th individual we consider the following two  $SDE$ 's:

$$dX(t) = \phi_{\xi_0}(t) b_{\beta_0}(t, X(t)) dt + \sigma(t, X(t)) dW(t) \quad (4.1)$$

and

$$dX(t) = \phi_{\xi_1}(t) b_{\beta_1}(t, X(t)) dt + \sigma(t, X(t)) dW(t). \quad (4.2)$$

For any  $t \in [0, T]$ , for  $j = 0, 1$ , let

$$\begin{aligned} U_{\theta_j, t} &= \int_0^t \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma^2(s, X(s))} dX(s), & V_{\theta_j, t} &= \int_0^t \frac{\phi_{\xi_j}^2(s) b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} ds, \\ V_{\theta_0, \theta_1, t} &= \int_0^t \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s)) \phi_{\xi_0}(s) b_{\beta_0}(s, X(s))}{\sigma^2(s, X(s))} ds. \end{aligned} \quad (4.3)$$

Note that  $V_{\theta_0, t} = V_{\theta_0, \theta_0, t}$  and  $V_{\theta_1, t} = V_{\theta_1, \theta_1, t}$ . We also let

$$f_{\theta_j, t}(X_{0, t}) = \exp \left( U_{\theta_j, t} - \frac{V_{\theta_j, t}}{2} \right). \quad (4.4)$$

Here we are interested in asymptotic properties of the Bayes factor, given by

$$I_T = \int \frac{f_{\theta_1, T}(X_{0, T})}{f_{\theta_0, T}(X_{0, T})} \pi(d\theta_1), \quad (4.5)$$

as  $T \rightarrow \infty$ .

For our purpose, let us define, for any  $h > 0$ ,

$$\begin{aligned} U_{\theta_j, t, t+h} &= \int_t^{t+h} \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma^2(s, X(s))} dX(s), & V_{\theta_j, t, t+h} &= \int_t^{t+h} \frac{\phi_{\xi_j}^2(s) b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} ds, \\ V_{\theta_0, \theta_j, t, t+h} &= \int_t^{t+h} \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s)) \phi_{\xi_0}(s) b_{\beta_0}(s, X(s))}{\sigma^2(s, X(s))} ds. \end{aligned} \quad (4.6)$$

Observe, as before, that  $V_{\theta_0, t, t+h} = V_{\theta_0, \theta_0, t, t+h}$  and  $V_{\theta_1, t, t+h} = V_{\theta_1, \theta_1, t, t+h}$ . We let

$$f_{\theta_j, t, t+h}(X_{t, t+h}) = \exp\left(U_{\theta_j, t, t+h} - \frac{V_{\theta_j, t, t+h}}{2}\right), \quad (4.7)$$

where, for any  $0 \leq a < b < \infty$ ,  $X_{a, b}$  denotes a path of the process  $X$  from  $a$  to  $b$ . For any  $t > 0$  and  $h > 0$ , we define

$$\tilde{\mathcal{K}}(f_{\theta_0, t, t+h}, f_{\theta_1, t, t+h}) = E_{\theta_0} \left[ \log \frac{f_{\theta_0, t, t+h}}{f_{\theta_1, t, t+h}} \right], \quad (4.8)$$

where  $E_{\theta_0} \equiv E_{f_{\theta_0, t, t+h}}$ . Note that although the expectation is with respect to  $f_{\theta_0, t, t+h}$ , which is not the same as  $f_{\theta_0, t, t+h}$ , (4.8) is still the Kullback-Leibler divergence between  $f_{\theta_0, t, t+h}$  and  $f_{\theta_1, t, t+h}$ . Also since in our case, for  $j = 0, 1$ ,

$$\begin{aligned} f_{\theta_j, t, t+h}(X_{t, t+h}) &= \exp\left(U_{\theta_j, t, t+h} - \frac{V_{\theta_j, t, t+h}}{2}\right) \\ &= \exp\left((U_{\theta_j, t, t+h} - U_{\theta_j, t}) - \frac{(V_{\theta_j, t, t+h} - V_{\theta_j, t})}{2}\right), \end{aligned} \quad (4.9)$$

it follows that

$$\tilde{\mathcal{K}}(f_{\theta_0, t, t+h}, f_{\theta_1, t, t+h}) = \mathcal{K}(f_{\theta_0, t, t+h}, f_{\theta_1, t, t+h}) - \mathcal{K}(f_{\theta_0, t}, f_{\theta_1, t}), \quad (4.10)$$

where  $\mathcal{K}(f_{\theta_0, t, t+h}, f_{\theta_1, t, t+h})$  and  $\mathcal{K}(f_{\theta_0, t}, f_{\theta_1, t})$  are proper Kullback-Leibler divergences between  $f_{\theta_0, t, t+h}$ ,  $f_{\theta_1, t, t+h}$ , and  $f_{\theta_0, t}$ ,  $f_{\theta_1, t}$ , respectively. We now define

$$\begin{aligned} \tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) &= \lim_{h \rightarrow 0} \frac{\tilde{\mathcal{K}}(f_{\theta_0, t, t+h}, f_{\theta_1, t, t+h})}{h} \\ &= \frac{1}{2} \frac{d}{dt} E_{\theta_0}(V_{\theta_0, t}) - \frac{d}{dt} E_{\theta_0}(V_{\theta_0, \theta_1, t}) + \frac{1}{2} \frac{d}{dt} E_{\theta_0}(V_{\theta_1, t}). \end{aligned} \quad (4.11)$$

The expression (4.11) easily follows using (4.9), the relation (2.8) and (2.10).

#### 4.1 Pseudo Kullback-Leibler ( $\delta$ ) property

We make the following assumption:

(H6'') For a fixed  $\delta \geq 0$ , the prior  $\pi$  satisfies

$$\pi\left(\theta_1 \in \Theta : \inf_t \tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) \geq \delta\right) = 1. \quad (4.12)$$

Let us define

$$\bar{\mathcal{K}}^\infty(f_{\theta_0}, f_{\theta_1}) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) dt. \quad (4.13)$$

We assume the following:

(H7'') Given  $\delta$  associated with (H6''), for any  $c \geq 0$ , the prior  $\pi$  satisfies

$$\pi(\theta_1 \in \Theta : \delta \leq \bar{\mathcal{K}}^\infty(f_{\theta_0}, f_{\theta_1}) \leq \delta + c) > 0. \quad (4.14)$$

We refer to property (H7'') as the pseudo Kullback-Leibler ( $\delta$ ) property of the prior  $\pi$ . Note that, (4.11), (3.13) and (3.14) imply

$$\begin{aligned} \bar{\mathcal{K}}^\infty(f_{\theta_0}, f_{\theta_1}) &= \frac{\bar{\phi}_{\xi_0}^{(2)}}{2} \kappa_0(\beta_0) - \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1) + \frac{\bar{\phi}_{\xi_1}^{(2)}}{2} \kappa_1(\beta_1) \\ &\geq \frac{1}{2} \left( \sqrt{\bar{\phi}_{\xi_0}^{(2)} \kappa_0(\beta_0)} - \sqrt{\bar{\phi}_{\xi_1}^{(2)} \kappa_1(\beta_1)} \right)^2 \\ &\geq 0, \end{aligned} \quad (4.15)$$

by Lemma 4. Provided that (4.12) holds and the prior  $\pi$  is dominated by the Lebesgue measure, the pseudo Kullback-Leibler ( $\delta$ ) property holds because of continuity of (4.15) in  $\theta_1 = (\beta_1, \xi_1)$  ensured by Lemma 2.

## 4.2 $Q^*$ property

For  $t \geq 0$ , let  $\mathcal{F}_t$  be the  $\sigma$ -algebra generated by  $X(0)$  and the history of the process upto (and including) time  $t$ , and let  $\pi_t(\theta_1) = \pi(\theta_1 | \mathcal{F}_t)$  be the posterior of  $\theta_1$  given  $\mathcal{F}_t$ . Also, let

$$\begin{aligned} &\hat{f}_{t-h,t}(X_{t-h,t}) \\ &= \int_{\theta_1 \in \Theta} \exp \left( \int_{t-h}^t \frac{\phi_{\xi_1}(s) b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_1}^2(s) b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} ds \right) \pi_{t-h}(d\theta_1) \\ &= E \left[ \exp \left( \int_{t-h}^t \frac{\phi_{\xi_1}(s) b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_1}^2(s) b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} ds \right) \middle| \mathcal{F}_{t-h} \right] \end{aligned} \quad (4.16)$$

be the posterior predictive density. Further, for any Borel set  $A$  such that  $\pi(A) > 0$ , let

$$\begin{aligned} &\hat{f}_{t-h,t,A}(X_{t-h,t}) \\ &= \int_{\theta_1 \in A} \exp \left( \int_{t-h}^t \frac{\phi_{\xi_1}(s) b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_1}^2(s) b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} ds \right) \pi_{t-h,A}(d\theta_1) \\ &= E \left[ \exp \left( \int_{t-h}^t \frac{\phi_{\xi_1}(s) b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_1}^2(s) b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} ds \right) \middle| \mathcal{F}_{t-h}, A \right], \end{aligned} \quad (4.17)$$

where

$$\pi_{t,A}(d\theta_1) = \frac{\mathbf{I}_A(\theta_1) \pi_t(d\theta_1)}{\int_A \pi_t(d\theta_1)}$$

is the posterior restricted to the set  $A$ . We assume the following:

(H8'')

$$\liminf_t E_{\theta_0} \left[ \tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}_{A_t(\delta)}) \right] \geq \delta, \quad (4.18)$$

whenever

$$A_t(\delta) = \left\{ \theta_1 \in \Theta : \tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) \geq \delta \right\}. \quad (4.19)$$

We refer to (H8'') as the  $Q^*$  property.

### 4.3 Main result on convergence of Bayes factor when two individual SDE's are compared

Let  $I_0 \equiv 1$  and for  $t > 0$ , let us define, analogous to (4.5),

$$I_t = \int \frac{f_{\theta_1, t}(X_{0, t})}{f_{\theta_0, t}(X_{0, t})} \pi(d\theta_1). \quad (4.20)$$

The following lemma, proved in section S-4 of the supplement, will prove useful in proving our main theorem on convergence of Bayes factor.

#### Lemma 5

$$E_{\theta_0} \left[ \log \frac{I_t}{I_{t-h}} \middle| \mathcal{F}_{t-h} \right] = E_{\theta_0} \left[ \log \frac{\hat{f}_{t-h, t}(X_{t-h, t})}{f_{\theta_0, t-h, t}(X_{t-h, t})} \middle| \mathcal{F}_{t-h} \right] = -\tilde{\mathcal{K}}(f_{\theta_0, t-h, t}, \hat{f}_{t-h, t}). \quad (4.21)$$

We make the following further assumption:

(H9'') For any  $t \geq 0$ ,  $\tilde{\mathcal{K}}(f_{\theta_0, t, t+h_n}, \hat{f}_{t, t+h_n})$  converges in expectation for all sequences  $\{h_n\}$  converging to zero as  $n \rightarrow \infty$ , with limit independent of  $\{h_n\}$ . We refer to the limiting process as  $\tilde{\mathcal{K}}'_t$ . In other words,

$$\lim_{n \rightarrow \infty} E \left( \frac{\tilde{\mathcal{K}}(f_{\theta_0, t, t+h_n}, \hat{f}_{t, t+h_n})}{h_n} \right) = E \left( \tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}) \right), \quad (4.22)$$

for any sequence  $\{h_n\}$  such that  $h_n \rightarrow 0$  as  $n \rightarrow \infty$ .

Because of Lemma 5 it follows from (H9''), using uniform integrability (which is easily seen to hold because of (H1'') – (H4'') and (3.2)), that  $J_{h_n}(t) = \frac{\log I_{t+h_n} - \log I_t}{h_n}$  converges in expectation for all sequences  $\{h_n\}$  converging to zero as  $n \rightarrow \infty$ , with limit independent of  $\{h_n\}$ . We refer to the limiting process as  $J'_t$ . That is, for any  $t \geq 0$ ,

$$\lim_{n \rightarrow \infty} E(J_{h_n}(t)) = E(J'_t) = \frac{d}{dt} E(\log I_t). \quad (4.23)$$

Now,

$$\hat{f}_{t, t+h} = E \left[ \exp \left( \int_t^{t+h} \frac{\phi_{\xi_1}(s) b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \frac{1}{2} \int_t^{t+h} \frac{\phi_{\xi_1}^2(s) b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} ds \right) \middle| \mathcal{F}_t \right], \quad (4.24)$$

so that Lemma 5 implies

$$E \left[ \log \frac{I_{t+h}}{I_t} \middle| \mathcal{F}_t \right] = E \left[ \log \frac{\hat{f}_{t, t+h}(X_{t, t+h})}{f_{\theta_0, t, t+h}(X_{t, t+h})} \middle| \mathcal{F}_t \right] = -\tilde{\mathcal{K}}(f_{\theta_0, t, t+h}, \hat{f}_{t, t+h}). \quad (4.25)$$

It follows, using (H9''), that

$$E(J'_t | \mathcal{F}_t) = -\tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}). \quad (4.26)$$

Note that for all sequences  $\{h_n\}$  such that  $h_n \rightarrow 0$  as  $n \rightarrow \infty$ ,  $J_{h_n}(t) = \frac{\log I_{t+h_n} - \log I_t}{h_n}$  is measurable with respect to  $\mathcal{F}_{t^*} = \sigma(X(s) : 0 \leq s \leq t^*)$ , for all  $t^* > t \geq 0$ . Hence,  $E(J'_t | \mathcal{F}_t) \neq J'_t$ .

Regarding convergence of  $I_T$ , we are now ready to present our main theorem whose proof is provided in section S-5 of the supplement.

**Theorem 6** *Assume the SDE set-up and conditions (H1'') – (H9''). Then*

$$\frac{1}{T} E_{\theta_0} (\log I_T) \rightarrow -\delta, \quad (4.27)$$

but

$$\frac{1}{T^2} \text{Var}_{\theta_0} (\log I_T) = O(1), \quad (4.28)$$

as  $T \rightarrow \infty$ .

**Corollary 7** *For  $j = 1, 2$ , let  $R_{jT}(\theta_j) = \frac{f_{\theta_j, T}(X_T)}{f_{\theta_0, T}(X_T)}$ , where  $\theta_1$  and  $\theta_2$  are two different finite sets of parameters, perhaps with different dimensionalities, associated with the two models to be compared. For  $j = 1, 2$ , let*

$$I_{jT} = \int R_{jT}(\theta_j) \pi_j(d\theta_j),$$

where  $\pi_j$  is the prior on  $\theta_j$ . Let  $B_T = I_{1T}/I_{2T}$  denote the Bayes factor for comparing the two models associated with  $\pi_1$  and  $\pi_2$ . Assume that both the models satisfy (H1'') – (H9''), and have the pseudo Kullback-Leibler property with  $\delta = \delta_1$  and  $\delta = \delta_2$  respectively. Then

$$E_{\theta_0} \left( \frac{1}{T} \log B_T \right) \rightarrow \delta_2 - \delta_1, \quad (4.29)$$

as  $T \rightarrow \infty$ .

## 5 Illustration of our asymptotic result for comparing two individual SDE's with a special case

Let the parameter space  $\Theta$  be compact, so that (H1'') holds. Let  $b_{\beta_j}$  and  $\sigma$  satisfy (H2'') such that

$$\frac{b_{\beta_j}(s, x)}{\sigma(s, x)} \equiv \eta_j(\beta_j); \quad j = 0, 1, \quad (5.1)$$

so that

$$\frac{b_{\beta_1}(s, x) b_{\beta_0}(s, x)}{\sigma^2(s, x)} \equiv \eta_0(\beta_0) \eta_1(\beta_1). \quad (5.2)$$

In the above,  $\eta_1(\beta_1)$  is continuous in  $\beta_1$ . Hence, (H3'') and (H4'') are satisfied. We assume that the relevant covariates and the functions  $g_l$  are such that (H5'') holds.

Letting  $\kappa_j(\beta_j) = \{\eta_j(\beta_j)\}^2$  and  $\bar{\kappa}(\beta_0, \beta_1) = \eta(\beta_0)\eta(\beta_1)$ , equations (5.1) – (5.2) entail

$$V_{\theta_j, t} = \kappa_j(\beta_j) \int_0^t \phi_{\xi_j}^2(s) ds; \quad (5.3)$$

$$V_{\theta_0, \theta_1, t} = \bar{\kappa}(\beta_0, \beta_1) \int_0^t \phi_{\xi_0}(s) \phi_{\xi_1}(s) ds; \quad (5.4)$$

$$U_{\theta_0, t} = \kappa_0(\beta_0) \int_0^t \phi_{\xi_0}^2(s) ds + \eta_0(\beta_0) \int_0^t \phi_{\xi_0}(s) dW(s); \quad (5.5)$$

$$U_{\theta_0, \theta_1, t} = \bar{\kappa}(\beta_0, \beta_1) \int_0^t \phi_{\xi_0}(s) \phi_{\xi_1}(s) ds + \eta_1(\beta_1) \int_0^t \phi_{\xi_1}(s) dW(s); \quad (5.6)$$

$$V_{\theta_j, t, t+h} = \kappa_j(\beta_j) \int_t^{t+h} \phi_{\xi_j}^2(s) ds; \quad (5.7)$$

$$V_{\theta_0, \theta_1, t, t+h} = \bar{\kappa}(\beta_0, \beta_1) \int_t^{t+h} \phi_{\xi_0}(s) \phi_{\xi_1}(s) ds; \quad (5.8)$$

$$U_{\theta_0, t, t+h} = \kappa_0(\beta_0) \int_t^{t+h} \phi_{\xi_0}^2(s) ds + \eta_0(\beta_0) \int_t^{t+h} \phi_{\xi_0}(s) dW(s); \quad (5.9)$$

$$U_{\theta_0, \theta_1, t, t+h} = \bar{\kappa}(\beta_0, \beta_1) \int_t^{t+h} \phi_{\xi_0}(s) \phi_{\xi_1}(s) ds + \eta_1(\beta_1) \int_t^{t+h} \phi_{\xi_1}(s) dW(s). \quad (5.10)$$

Due to (5.3) and (5.4), we obtain

$$\begin{aligned} \tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) &= \lim_{h \rightarrow 0} \frac{\tilde{\mathcal{K}}(f_{\theta_0, t, t+h}, f_{\theta_1, t, t+h})}{h} \\ &= \frac{1}{2} \frac{d}{dt} E_{\theta_0}(V_{\theta_0, t}) - \frac{d}{dt} E_{\theta_0}(V_{\theta_0, \theta_1, t}) + \frac{1}{2} \frac{d}{dt} E_{\theta_0}(V_{\theta_1, t}) \\ &= \frac{\phi_{\xi_0}^2(t)}{2} \kappa_0(\beta_0) - \phi_{\xi_1}(t) \phi_{\xi_0}(t) \bar{\kappa}(\beta_0, \beta_1) + \frac{\phi_{\xi_1}^2(t)}{2} \kappa_1(\beta_1) \\ &= \frac{1}{2} (\phi_{\xi_0}(t) \eta_0(\beta_0) - \phi_{\xi_1}(t) \eta_1(\beta_1))^2. \end{aligned} \quad (5.11)$$

Note that

$$\begin{aligned} \inf_t \tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) &\geq \frac{1}{2} \inf_{\beta_1 \in \mathfrak{B}, \xi_1 \in \Gamma, z \in \mathfrak{Z}} (\phi_{\xi_0}(z) \eta_0(\beta_0) - \phi_{\xi_1}(z) \eta_1(\beta_1))^2 \\ &= \delta. \end{aligned} \quad (5.12)$$

It follows from (5.12) that (H6'') holds for any prior on  $\theta_1$ .

Also, it follows directly from (5.11), that

$$\bar{\mathcal{K}}^\infty(f_{\theta_0}, f_{\theta_1}) = \frac{\bar{\phi}_{\xi_0}^{(2)}}{2} \kappa_0(\beta_0) - \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1) + \frac{\bar{\phi}_{\xi_1}^{(2)}}{2} \kappa_1(\beta_1), \quad (5.13)$$

which is continuous in  $(\beta_1, \xi_1)$ , due to the continuity assumption of  $\eta_1(\beta_1)$  in  $\beta_1$  and Lemma 2, which guarantees continuity of  $\bar{\phi}_{\xi_0, \xi_1}^{(2)}$  and  $\bar{\phi}_{\xi_1}^{(2)}$  in  $\xi_1$ . Since the right-most side of (5.13) is a continuous function of  $\theta_1$ , it follows that (H7'') is clearly satisfied if the prior  $\pi$  is dominated by the Lebesgue measure.

We now verify the  $Q^*$  property (H8''). Recall that  $\hat{f}_{A_t(\delta)} = \hat{f}_{t, t+h}$ , since  $\pi(A_t(\delta)) = 1$ . Since

$$\hat{f}_{t, t+h}(X_{t, t+h}) \leq \sup_{\theta_1 \in A_t(\delta)} f_{\theta_1, t, t+h}(X_{t, t+h}) = \hat{f}_{\hat{\theta}_1(X_{t, t+h}), t, t+h}(X_{t, t+h}), \quad (5.14)$$

where  $\hat{\theta}_1(X_{t,t+h}) \in A_t(\delta)$ , is the maximizer of  $f_{\theta_1,t,t+h}$  in the compact set  $A_t(\delta)$ . Hence,

$$\begin{aligned}
& \tilde{\mathcal{K}}\left(f_{\theta_0,t,t+h}, \hat{f}_{A_t(\delta)}\right) \\
&= E_{\theta_0}(\log f_{\theta_0,t,t+h}(X_{t,t+h})) - E_{\theta_0}\left(\log \hat{f}_{A_t(\delta)}(X_{t,t+h})|\mathcal{F}_t\right) \\
&\geq E_{\theta_0}(\log f_{\theta_0,t,t+h}(X_{t,t+h})) - E_{\theta_0}\left(\log \hat{f}_{\hat{\theta}_1(X_{t,t+h}),t,t+h}(X_{t,t+h})|\mathcal{F}_t\right) \\
&= E_{\theta_0}\left(\log \frac{f_{\theta_0,t,t+h}(X_{t,t+h})}{f_{\hat{\theta}_1(X_{t,t+h}),t,t+h}(X_{t,t+h})}\middle|\mathcal{F}_t\right) \\
&= E_{\hat{\theta}_1(X_{t,t+h})|\theta_0} E_{X_{t,t+h}|\hat{\theta}_1(X_{t,t+h}),\theta_0}\left(\log \frac{f_{\theta_0,t,t+h}(X_{t,t+h})}{f_{\hat{\theta}_1(X_{t,t+h}),t,t+h}(X_{t,t+h})}\middle|\mathcal{F}_t\right). \tag{5.15}
\end{aligned}$$

Now, let  $f_{\theta_0,tt+h}(X_{t,t+h}|Y) = \frac{f_{\theta_0,tt+h}(X_{t,t+h})}{g_{\theta_0,tt+h}(Y)} \mathbf{I}_{\{\hat{\theta}_1(X_{t,t+h})=Y\}}(Y)$  be the conditional density of  $X_{t,t+h}$  given  $Y = \hat{\theta}_1(X_{t,t+h})$ , the latter having density  $g_{\theta_0,tt+h}(Y)$ . The dominating probability measure associated with this conditional density is  $P_{0,t,t+h}$ , which is the same dominating probability measure associated with  $f_{\theta_0,t,t+h}$ . Then as in Maitra and Bhattacharya (2018) we have

$$\begin{aligned}
& E_{X_{t,t+h}|Y,\theta_0}\left(\log \frac{f_{\theta_0,t,t+h}(X_{t,t+h})}{f_{\hat{\theta}_1(X_{t,t+h}),t,t+h}(X_{t,t+h})}\middle|\mathcal{F}_t\right) \\
&= \int \log\left(\frac{f_{\theta_0,t,t+h}(X_{t,t+h}|Y)}{f_{Y,t,t+h}(X_{t,t+h})}\right) f_{\theta_0,t,t+h}(X_{t,t+h}|Y) dP_{0,t,t+h} + \log g_{\theta_0,tt+h}(Y). \tag{5.16}
\end{aligned}$$

Since the first term of (4.11) is the Kullback-Leibler divergence between  $f_{\theta_0,t,t+h}(X_{t,t+h}|Y)$  and  $f_{Y,t,t+h}(X_{t,t+h})$ , it is positive for almost all  $Y$ . Hence,

$$E_{Y|\theta_0} E_{X_{t,t+h}|Y,\theta_0}\left(\log \frac{f_{\theta_0,t,t+h}(X_{t,t+h}|Y)}{f_{Y,t,t+h}(X_{t,t+h})}\right) > 0. \tag{5.17}$$

Also, by Jensen's inequality,  $E_{Y|\theta_0}[\log g_{\theta_0,tt+h}(Y)] \geq -\log E_{Y|\theta_0}\left(\frac{1}{g_{\theta_0,tt+h}(Y)}\right)$ . Assuming the distribution of  $Y$  is dominated by the Lebesgue measure, we have  $E_{Y|\theta_0}\left(\frac{1}{g_{\theta_0,tt+h}(Y)}\right) = |A_t(\delta)| = \int_{A_t(\delta)} dy$ . As in Maitra and Bhattacharya (2018), once again we argue that the compact space  $\Theta$  can be rescaled appropriately with respect to suitable reparameterization such that for all  $h > 0$ ,  $\sup_t |A_t(\delta)| < \exp(-\delta h)$ . Hence,  $E_{Y|\theta_0}[\log g_{\theta_0,tt+h}(Y)] \geq \delta h$ , which finally implies in accordance with (5.17), that  $\tilde{\mathcal{K}}\left(f_{\theta_0,t,t+h}, \hat{f}_{A_t(\delta)}\right) \geq \delta h$ . Hence,  $\tilde{\mathcal{K}}'\left(f_{\theta_0}, \hat{f}_{A_t(\delta)}\right) \geq \delta$ , showing that the  $Q^*$  property is satisfied.

To see that (H9'') holds, first observe that it follows from the proof of Lemma 5 that  $\frac{I_{t+h}}{I_t} = \frac{\hat{f}_{t,t+h}}{f_{\theta_0,t,t+h}}$ , which implies

$$\frac{\tilde{\mathcal{K}}\left(f_{\theta_0,t,t+h}, \hat{f}_{t,t+h}\right)}{h} = \frac{E_{\theta_0}\left(\log f_{\theta_0,t,t+h}(X_{t,t+h}) - \log \hat{f}_{t,t+h}(X_{t,t+h})|\mathcal{F}_t\right)}{h} \tag{5.18}$$

Now,

$$E_{\theta_0}(\log f_{\theta_0,t,t+h}(X_{t,t+h})|\mathcal{F}_t) = \frac{\kappa_0(\beta_0)}{2} \int_t^{t+h} \phi_{\xi_0}^2(s) ds = \frac{\kappa_0(\beta_0)}{2} h \phi_{\xi_0}^2(s^*(h)), \tag{5.19}$$

by the mean value theorem for integrals, where  $s^*(h) \rightarrow t$ , as  $h \rightarrow 0$ . Hence, using continuity of  $\phi_{\xi_0}(t)$

in  $t$ , we obtain

$$\lim_{h \rightarrow 0} \frac{1}{h} E_{\theta_0} (\log f_{\theta_0, t, t+h}(X_{t, t+h}) | \mathcal{F}_t) = \frac{\kappa_0(\beta_0)}{2} \lim_{h \rightarrow 0} \phi_{\xi_0}^2(s^*(h)) = \frac{\kappa_0(\beta_0)}{2} \phi_{\xi_0}^2(t). \quad (5.20)$$

To deal with  $E_{\theta_0} (\log \hat{f}_{t, t+h}(X_{t, t+h}) | \mathcal{F}_t)$ , note that for any  $X_{t, t+h}$ , by the mean value theorem for integrals,

$$\hat{f}_{t, t+h}(X_{t, t+h}) = f_{\check{\theta}(X_{0, t}, X_{t, t+h}), t, t+h}(X_{t, t+h}),$$

where  $\check{\theta}(X_{0, t}, X_{t, t+h}) \in \Theta$ . It is clear that  $\check{\theta}(X_{0, t}, X_{t, t+h}) \rightarrow \check{\theta}(X_{0, t}, X_t) = \check{\theta}(X_{0, t})$  almost surely, as  $h \rightarrow 0$ . Hence,

$$\begin{aligned} & E_{\theta_0} (\log \hat{f}_{t, t+h}(X_{t, t+h}) | \mathcal{F}_t) \\ &= E_{\check{\theta}(X_{0, t}, X_{t, t+h}) | \theta_0} E_{X_{t, t+h} | \check{\theta}(X_{0, t}, X_{t, t+h}) = \alpha, \theta_0} (\log f_{\{\check{\theta}(X_{0, t}, X_{t, t+h}) = \alpha\}, t, t+h}(X_{t, t+h}) | \mathcal{F}_t), \end{aligned}$$

where

$$\begin{aligned} & E_{\theta_0} (\log \hat{f}_{t, t+h}(X_{t, t+h}) | \mathcal{F}_t) \\ &= E_{\theta_0} (\log f_{\{\check{\theta}(X_{0, t}, X_{t, t+h})\}, t, t+h}(X_{t, t+h}) | \mathcal{F}_t) \\ &= E_{\theta_0} \left( \bar{\kappa}(\beta_0, \check{\beta}_1(X_{0, t}, X_{t, t+h})) \int_t^{t+h} \phi_{\xi_1(X_{0, t}, X_{t, t+h})}(s) \phi_{\xi_0}(s) ds \right. \\ &\quad \left. - \frac{\kappa_1(\check{\beta}_1(X_{0, t}, X_{t, t+h}))}{2} \int_t^{t+h} \phi_{\xi_1(X_{0, t}, X_{t, t+h})}^2(s) ds \right) \\ &= E_{\theta_0} \left( \bar{\kappa}(\beta_0, \check{\beta}_1(X_{0, t}, X_{t, t+h})) h \phi_{\xi_1(X_{0, t}, X_{t, t+h})}^{\check{z}}(s_1(h)) \phi_{\xi_0}(s_1(h)) \right. \\ &\quad \left. - \frac{\kappa_1(\check{\beta}_1(X_{0, t}, X_{t, t+h}))}{2} h \phi_{\xi_1(X_{0, t}, X_{t, t+h})}^2(s_2(h)) \right), \quad (5.21) \end{aligned}$$

where  $s_1(h), s_2(h) \in [t, t+h]$ , associated with the mean value theorem for integrals. Hence,  $s_1(h) \rightarrow t$  and  $s_2(h) \rightarrow t$ , almost surely as  $h \rightarrow 0$ .

Continuity of  $\bar{\kappa}(\cdot), \kappa_1(\cdot)$  and the results  $\check{\theta}(X_{0, t}, X_{t, t+h}) \rightarrow \check{\theta}(X_{0, t}), s_1(h) \rightarrow t, s_2(h) \rightarrow t$ , almost surely, as  $h \rightarrow 0$ , in conjunction with the dominated convergence theorem exploiting boundedness of the functions  $\phi_{\xi_0}, \phi_{\xi_1}^{\check{z}}, \bar{\kappa}$  and  $\kappa_j$ , imply, using continuity of  $\phi_{\xi_1}^{\check{z}}(t)$  in  $t$ , that

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{1}{h} E_{\theta_0} (\log \hat{f}_{t, t+h}(X_{t, t+h}) | \mathcal{F}_t) \\ &= \phi_{\xi_0}(t) \phi_{\xi_1(X_{0, t})}^{\check{z}}(t) \bar{\kappa}(\beta_0, \check{\beta}(X_{0, t})) - \frac{\phi_{\xi_1(X_{0, t})}^2(t)}{2} \kappa_1(\check{\beta}(X_{0, t})). \end{aligned}$$

In other words, the limit of (5.18) exists and is unique as  $h \rightarrow 0$ . Now, equations (5.19) and (5.21) along with dominated convergence theorem imply that (H9'') holds.

Thus, all the assumptions required for Theorem 6 and Corollary 7 are satisfied. Hence, both (4.27) and (4.29) hold.



## 6 Asymptotic convergence of Bayes factor in the *SDE* set-up with respect to number of individuals and time

### 6.1 Convergence of Bayes factor in the *iid* set-up

Although Theorem 6 fails to ensure consistency of the Bayes factor as  $T \rightarrow \infty$  in the sense that the relevant variance is asymptotically positive, the theorem is useful to prove almost sure consistency when  $T \rightarrow \infty$  as well as  $n \rightarrow \infty$ , for both *iid* and non-*iid* situations. Theorem 8 formalizes this for the *iid* set-up, while Theorem 12 establishes almost sure consistency of the Bayes factor in the non-*iid* situation. Proofs of these theorems are contained in section S-6 and S-9 respectively in the supplement.

**Theorem 8** *Assume the iid set-up; also assume that conditions (H1'') – (H9'') hold for each SDE in the systems (2.1) and (2.2). Then*

$$\frac{1}{nT} \log \tilde{I}_{n,T} \rightarrow -\delta, \quad (6.1)$$

almost surely, as  $n \rightarrow \infty$  and  $T \rightarrow \infty$ .

The following corollary is obvious.

**Corollary 9** *For  $j = 1, 2$ , and  $i = 1, \dots, n$ , let  $R_{j,i,T}(\boldsymbol{\theta}_j^{(i)}) = \frac{f_{\boldsymbol{\theta}_j^{(i)},i,T}(X_{i,0,T})}{f_{\boldsymbol{\theta}_0^{(i)},i,T}(X_{i,0,T})}$ , where, for each  $i$ ,  $\boldsymbol{\theta}_1^{(i)}$  and  $\boldsymbol{\theta}_2^{(i)}$  are two different finite sets of parameters, perhaps with different dimensionalities, associated with the two systems (2.1) and (2.2) to be compared. For  $j = 1, 2$ , let*

$$\tilde{I}_{j,n,T} = \prod_{i=1}^n \int R_{j,i,T}(\boldsymbol{\theta}_j^{(i)}) \pi_j(d\boldsymbol{\theta}_j^{(i)}),$$

where  $\pi_j$  is the prior on  $\boldsymbol{\theta}_j^{(i)}$ , for  $i = 1, 2, \dots$ . Let  $B_{n,T} = \tilde{I}_{1,n,T} / \tilde{I}_{2,n,T}$  denote the Bayes factor for comparing the two models associated with  $\pi_1$  and  $\pi_2$ . Assume the *iid* case and suppose that both the systems satisfy (H1'') – (H9''), and have the pseudo Kullback-Leibler property with  $\delta = \delta_1$  and  $\delta = \delta_2$  respectively. Then

$$\frac{1}{nT} \log B_{n,T} \rightarrow \delta_2 - \delta_1,$$

almost surely, as  $n \rightarrow \infty$  and  $T \rightarrow \infty$ .

### 6.2 Convergence of Bayes factor in the non-*iid* set-up

We now relax the assumptions  $x^i = x$  and  $\xi_{1j}^{(i)} = \xi_{2j}^{(i)} = \xi_{3j}^{(i)} = \dots = \xi_{pj}^{(i)} = 0$  for  $j = 0, 1$ . Thus, we are now in a non-*iid* situation where the processes  $X_i(\cdot)$ ;  $i = 1, \dots, n$ , are independently, but not identically distributed. As mentioned in Section 2.1 we assume that  $\boldsymbol{\theta}_1^{(i)} \stackrel{iid}{\sim} \pi$ . In this set-up, for each  $\mathbf{z} \in \mathfrak{Z} = \{\mathbf{z}(t) \in \mathfrak{Z} : t \in [0, \infty)\}$ , it holds, due to Theorem 6, that

$$\frac{1}{T} E_{\boldsymbol{\theta}_0} (\log I_{x,T,\mathbf{z}}) \rightarrow -\delta(x, \mathbf{z}), \quad (6.2)$$

as  $T \rightarrow \infty$ , where  $\delta(x, \mathbf{z})$  depends upon the initial value  $x \in \mathfrak{X}$  and the set of time-dependent covariates  $\mathbf{z} \in \mathfrak{Z}$ . The following lemma shows that  $\delta(x, \mathbf{z})$  is continuous in  $(x, \mathbf{z}) \in \mathfrak{X} \times \mathfrak{Z}$ .

**Lemma 10** *Assume the conditions of Theorem 6. Then,  $\delta(x, \mathbf{z})$  is continuous in  $(x, \mathbf{z}) \in \mathfrak{X} \times \mathfrak{Z}$ .*

Now consider the following limit:

$$\delta^\infty = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta(x^i, \mathbf{z}_i). \quad (6.3)$$

The following lemma shows that the above limit exists for all sequences  $\{(x^i, z_i)\}_{i=1}^\infty \in \mathfrak{X} \times \mathfrak{Z}$ .

**Lemma 11** *The limit (6.3) exists for all sequences  $\{(x^i, z_i)\}_{i=1}^\infty \in \mathfrak{X} \times \mathfrak{Z}$ .*

Proof of these two lemmas are provided in section S-7 and S-8 respectively in the supplement. Now, we have the following theorem.

**Theorem 12** *Assume the non-iid set-up, and conditions (H1'') – (H9''), for each SDE in the systems (2.1) and (2.2). Then*

$$\frac{1}{nT} \log \tilde{I}_{n,T} \rightarrow -\delta^\infty, \quad (6.4)$$

almost surely, as  $T \rightarrow \infty$  and  $n \rightarrow \infty$ .

We then have the following corollary for the non-iid case.

**Corollary 13** *For  $j = 1, 2$ , and  $i = 1, \dots, n$ , let  $R_{j,i,T}(\theta_j^{(i)}) = \frac{f_{\theta_j^{(i)},i,T}(X_{i,0,T})}{f_{\theta_0^{(i)},i,T}(X_{i,0,T})}$ , where, for each  $i$ ,  $\theta_1^{(i)}$  and  $\theta_2^{(i)}$  are two different finite sets of parameters, perhaps with different dimensionalities, associated with the two systems (2.1) and (2.2) to be compared. For  $j = 1, 2$ , let*

$$\tilde{I}_{j,n,T} = \prod_{i=1}^n \int R_{j,i,T}(\theta_j^{(i)}) \pi_j(d\theta_j^{(i)}),$$

where  $\pi_j$  is the prior on  $\theta_j^{(i)}$ ;  $i = 1, 2, \dots$ . Let  $B_{n,T} = \tilde{I}_{1,n,T} / \tilde{I}_{2,n,T}$  denote the Bayes factor for comparing the two models associated with  $\pi_1$  and  $\pi_2$ . Assume the non-iid case and suppose that both the systems satisfy (H1'') – (H9''), and have the pseudo Kullback-Leibler property with  $\delta_i = \delta_{1i}$  and  $\delta_i = \delta_{2i}$  respectively. Let, for  $j = 1, 2$ ,

$$\delta_j^\infty = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta_{ji}.$$

Then

$$\frac{1}{nT} \log B_{n,T} \rightarrow \delta_2^\infty - \delta_1^\infty,$$

almost surely, as  $n \rightarrow \infty$  and  $T \rightarrow \infty$ .

## 7 Simulation studies

### 7.1 Covariate selection when $n = 1, T = 5$

We first demonstrate with simulation study the finite sample analogue of Bayes factor analysis associated with a single individual, when  $T \rightarrow \infty$ . In this regard, we consider modeling a single individual by

$$dX(t) = (\xi_1 + \xi_2 z_1(t) + \xi_3 z_2(t) + \xi_4 z_3(t))(\xi_5 + \xi_6 X(t))dt + \sigma dW(t), \quad (7.1)$$

where we fix our diffusion coefficient as  $\sigma = 20$ . We consider the initial value  $X(0) = 0$  and the time interval  $[0, T]$  with  $T = 5$ .

To achieve numerical stability of the marginal likelihood corresponding to data we choose the true values of  $\xi_i$ ;  $i = 1, \dots, 6$  as follows:  $\xi_i \stackrel{iid}{\sim} N(\mu_i, 0.001^2)$ , where  $\mu_i \stackrel{iid}{\sim} N(0, 1)$ . This is not to be interpreted as the prior; this is just a means to set the true values of the parameters of the data-generating model.

We assume that the time dependent covariates  $z_i(t)$  satisfy the following *SDEs*

$$\begin{aligned} dz_1(t) &= (\tilde{\theta}_1 + \tilde{\theta}_2 z_1(t))dt + dW_1(t) \\ dz_2(t) &= \tilde{\theta}_3 dt + dW_2(t) \\ dz_3(t) &= \tilde{\theta}_4 z_3(t)dt + dW_3(t), \end{aligned} \tag{7.2}$$

where  $W_i(\cdot)$ ;  $i = 1, 2, 3$ , are independent Wiener processes, and  $\tilde{\theta}_i \stackrel{iid}{\sim} N(0, 0.01^2)$  for  $i = 1, \dots, 4$ .

We obtain the covariates by first simulating  $\tilde{\theta}_i \stackrel{iid}{\sim} N(0, 0.01^2)$  for  $i = 1, \dots, 4$ , fixing the values, and then by simulating the covariates using the *SDEs* (7.2) by discretizing the time interval  $[0, 5]$  into 500 equispaced time points. In all our applications we have standardized the covariates over time so that they have zero means and unit variances.

Once the covariates are thus obtained, we assume that the data are generated from the (true) model where all the covariates are present. For the true values of the parameters, we simulated  $(\xi_1, \dots, \xi_6)$  from the prior and treated the obtained values as the true set of parameters  $\theta_0$ . We then generated the data using (7.1) by discretizing the time interval  $[0, 5]$  into 500 equispaced time points.

As we have three covariates so we will have  $2^3 = 8$  different models. Denoting a model by the presence and absence of the respective covariates, it then is the case that  $(1, 1, 1)$  is the true, data-generating model, while  $(0, 0, 0)$ ,  $(0, 0, 1)$ ,  $(0, 1, 0)$ ,  $(0, 1, 1)$ ,  $(1, 0, 0)$ ,  $(1, 0, 1)$ , and  $(1, 1, 0)$  are the other 7 possible models.

As per our theory, for a single individual, the Bayes factor is not consistent for increasing time domain. However, we have shown that

$$\frac{1}{T} E_{\theta_0}(\log I_T) \rightarrow -\delta$$

as  $T \rightarrow \infty$ . Thus, the Bayes factor is consistent with respect to the expectation. Our simulation results show that this holds even for the time domain  $[0, 5]$ , where we approximate the expectation with the average of 1000 realizations of  $I_T$  associated with as many simulated data sets.

### 7.1.1 Case 1: the true parameter set $\theta_0$ is fixed

#### Prior on $\theta$

We first obtain the maximum likelihood estimator (*MLE*) of  $\theta$  using simulated annealing and then consider a normal prior with the *MLE* as the mean and variance  $0.8^2 \mathbb{I}_6$ , where  $\mathbb{I}_6$  is the identity matrix of order 6.

#### Form of the Bayes factor

In this case the related Bayes factor has the form

$$I_T = \int \frac{f_{\theta_1, T}(X_{0, T})}{f_{\theta_0, T}(X_{0, T})} \pi(d\theta_1), \tag{7.3}$$

where  $\theta_0 = (\xi_{0,1}, \xi_{0,2}, \xi_{0,3}, \xi_{0,4}, \xi_{0,5}, \xi_{0,6})$  is the true parameter set and  $\theta_1 = (\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6)$  is the unknown set of parameters corresponding to any other model. Table 7.1 describes the results of our Bayes factor analyses. It is clear from the 7 values of the table that the correct model  $(1, 1, 1)$  is always preferred.

Table 7.1: Bayes factor results

Model	Averaged $\frac{1}{5} \log I_5$
(0, 0, 0)	-2.5756029
(0, 0, 1)	-0.913546
(0, 1, 1)	-0.5454860
(0, 1, 0)	-0.763952
(1, 0, 0)	-2.5774163
(1, 0, 1)	-0.9312218
(1, 1, 0)	-0.7628154

### 7.1.2 Case 2: the parameter set $\theta_0$ is random and has the prior distribution $\pi$

As before, we consider the same form of the prior as in Section 7.1.1, but with variance  $0.1^2 \mathbb{I}_6$ . In this case we calculate marginal likelihood of the 8 possible models, and approximate

$$\frac{1}{5} E_{\theta_0} \left( \log \int f_{i, \theta_{1,5}}(X_{0,5}) \pi(d\theta_1) \right)$$

for  $i = 1, \dots, 8$  by averaging over 1000 replications of the data obtained from the true model. Denoting its values by  $\ell_i$ , Table 7.2 shows that  $\ell_8$  is the highest, implying consistency of the averaged Bayes factor.

Table 7.2: Averages of  $\frac{1}{5} \times$  marginal log-likelihood

Model	$\ell_i$
(0, 0, 0)	-1.21923
(0, 0, 1)	-0.21428
(0, 1, 0)	1.47992
(0, 1, 1)	2.102966
(1, 0, 0)	-1.222362
(1, 0, 1)	-0.21898
(1, 1, 0)	1.459921
(1, 1, 1)	2.121237 (true model)

## 7.2 Bayes factor analysis for $n = 15$ and $T = 5$

In this case we allow our parameter and the covariate sets to vary from individual to individual. We consider 15 individuals modeled by

$$dX_i(t) = (\xi_1^i + \xi_2^i z_1(t) + \xi_3^i z_2(t) + \xi_4^i z_3(t)) (\xi_5^i + \xi_6^i X_i(t)) dt + \sigma_i dW_i(t) \quad (7.4)$$

for  $i = 1, \dots, 15$ . We fix our diffusion coefficients as  $\sigma_{i+1} = \sigma_i + 5$  for  $i = 1 \dots, 14$  where  $\sigma_1 = 10$ . We consider the initial value  $X(0) = 0$  and the interval  $[0, T]$ , with  $T = 5$ . As before, we generated the observed data after discretizing the time interval into 500 equispaced time points. Here our covariates and the parameter set  $\theta_0^i = (\xi_{0,1}^i, \xi_{0,2}^i, \xi_{0,3}^i, \xi_{0,4}^i, \xi_{0,5}^i, \xi_{0,6}^i)$ ;  $i = 1, \dots, 15$ , are simulated in a similar way as mentioned in Section 7.1.

For each of the 15 individuals, the true set of covariate combination is randomly selected. Thus, for a given model, there are 15 sets of covariate combinations to be compared with other models consisting of 15 different sets of covariate combinations. To decrease computational burden we compare the true model with 100 other models consisting of different sets of covariate combinations.




Figure 8.1: Some company-wise time-series data.

The Bayes factor corresponding to the  $j$ -th covariate combination is given by

$$I_{nT}^j = \prod_{i=1}^n \int \frac{f_{i,\theta_1^{(i)}}^j(X_{i,0,T})}{f_{i,\theta_0^{(i)}}(X_{i,0,T})} \pi(\theta_1^{(i)}) d\theta_1^{(i)} \quad (7.5)$$

for  $j = 1, \dots, 100$ , where  $n = 15$ ,  $T = 5$  and  $\theta_0^{(i)}$  is the true parameter set corresponding to the  $i$ -th individual.

We obtain the *MLE* of the 15 parameter sets by simulated annealing. Then we calculate the Bayes factor with the prior such that the parameter components are independent normal with means as the respective *MLEs* and variances 1. In all the cases corresponding to 100 covariate combinations we obtain  $\frac{1}{nT} \log I_{nT}^j < 0$  for  $j = 1, \dots, 100$ . Thus, Bayes factor indicated the correct covariate combination in all the cases considered. We also considered the case when a normal prior is considered for the parameters of the true model. In this case with respect to the component-wise independent normal prior with individual mean as obtained from simulated annealing and component-wise variance  $0.1^2$ , we obtain

$$\frac{1}{15 \times 5} \left[ \log \left( \prod_{i=1}^{15} \int f_{i,\theta_1^{(i)}}^j(X_{i,0,T}) \pi(\theta_1^{(i)}) d\theta_1^{(i)} \right) - \log \left( \prod_{i=1}^{15} \int f_{i,\theta_0^{(i)}}(X_{i,0,T}) \pi(\theta_0^{(i)}) d\theta_0^{(i)} \right) \right] < 0, \quad (7.6)$$

for  $j = 1, \dots, 100$ . Indeed, it turned out that  $\frac{1}{15 \times 5} \log \left( \prod_{i=1}^{15} \int f_{i,\theta_0^{(i)}}(X_{i,0,T}) \pi(\theta_0^{(i)}) d\theta_0^{(i)} \right) = 0.4865$  and the maximum of  $\frac{1}{15 \times 5} \log \left( \prod_{i=1}^{15} \int f_{i,\theta_1^{(i)}}^j(X_{i,0,T}) \pi(\theta_1^{(i)}) d\theta_1^{(i)} \right)$  is 0.4127. In other words, the Bayes factor consistently selects the correct model even in this situation.

## 8 Company-wise data from national stock exchange

To deal with real data we collect the stock market data (467 observations during the time range August 5, 2013, to June 30, 2015) for 15 companies which is available on [www.nseindia.com](http://www.nseindia.com). The nature of some company-wise data are shown in Figure 8.1.

Each company-wise data is modeled by various available standard financial *SDE* models with the available “fitsde” package in *R*. After obtaining the BIC (Bayesian Information Criterion) for each company corresponding to each available financial model, we find that the minimum value of BIC corresponds to the *CKLS* model, given, for process  $X(t)$ , by

$$dX(t) = (\theta_1 + \theta_2 X(t))dt + \theta_3 X(t)^{\theta_4} dW(t).$$




Figure 8.2: Covariates.

As per our theory we treat the diffusion coefficient as a fixed quantity. So, after obtaining the estimated value of the coefficients by the “fitsde” function, we fix the values of  $\theta_3$  and  $\theta_4$ , so that the diffusion coefficient becomes fixed. We let  $\theta_3 = A$ ,  $\theta_4 = B$ .

In this *CKLS* model, we now wish to include time varying covariates. In our work we consider the “close price” of each company. The stock market data is assumed to be dependent on IIP general index, bank interest rate, US dollar exchange rate and on various other quantities. But we assume only these three quantities as possibly the most important time dependent covariates.

Briefly, IIP, that is, index of industrial production, is a measurement which represents the status of production in the industrial sector for a given period of time compared to a reference period of time. It is one of the best statistical data, which helps us measure the level of industrial activity in Indian economy. Its importance lies in the fact that low industrial production will result in lower corporate sales and profits, which will directly affect stock prices. So a direct impact of weak IIP data is a sudden fall in stock prices.

As the IIP data is purely industrial data, banking sector is not included in it. So, we also consider the bank interest rate as another covariate. Note that, higher the bank interest rate, fixed deposits become more attractive and one will preferably deposit money in bank rather than invest in stock market. Besides, companies with a high amount of loans in their balance sheets would be affected very seriously. Interest cost on existing debt would go up affecting their EPS (Earning per Share) and ultimately the stock prices. But during low interest rate these companies would stand to gain. Banking sector is likely to benefit most due to high interest rates. The Net Interest Margins (it is the difference between the interest they earn on the money they lend and the interest they pay to the depositors) for banks is likely to increase leading to growth in profits and the stock prices. Hence, it is clear that, the interest rates and stock markets are inversely related. As the interest rates go up, stock market activities tend to come down.

Finally, exchange rates directly affect the realized return on an investment portfolio with overseas holdings. If one own stock in a foreign company and the local currency goes up, the value of the investment also goes up. Foreign investment is also related very much to US dollar exchange rate.

Hence, we collect the values of the aforementioned time varying covariates during the time range August 5, 2013, to June 30, 2015. The pattern of the covariates are displayed in Figure 8.2.

We denote these three covariates by  $c_1, c_2, c_3$  respectively. Now, our considered *SDE* models for national stock exchange data associated with the 15 companies are the following:

$$dX_i(t) = (\theta_1^i + \theta_2^i c_1(t) + \theta_3^i c_2(t) + \theta_4^i c_3(t))(\theta_5^i + \theta_6^i X_i(t))dt + A^i X_i(t)^{B^i} dW_i(t), \quad (8.1)$$

for  $i = 1, \dots, 15$ .

## 8.1 Selection of covariates by Bayes factor

Among the considered three time varying covariates we now select the best set of covariate combinations for the 15 companies among 100 such sets through Bayes factor, computing the log-marginal-likelihoods with respect to the normal prior on the parameter set, assuming *a priori* independence of the parameter components with individual means being the corresponding *MLE* (based on simulated annealing) and  $0.01^2$  variance (relatively small variance ensured numerical stability of the marginal likelihood). Table 8.1 provides the sets of covariates for the 15 companies obtained by our Bayes factor analysis. Also observe that each of the three covariates occurs about 50% times among the companies, demonstrating that overall impact of these on national stock exchange is undeniable.

Table 8.1: Company-wise covariates obtained by Bayes factor analysis

Company	Covariates
1	Bank rate
2	US dollar exchange rate
3	None
4	None
5	Bank rate and US dollar exchange rate
6	Bank rate and US dollar exchange rate
7	IIP general index and US dollar exchange rate
8	Bank rate
9	IIP general index and Bank rate
10	IIP general index
11	IIP general index, Bank rate and US dollar exchange rate
12	IIP general index and Bank rate
13	US dollar exchange rate
14	IIP general index, Bank rate and US dollar exchange rate
15	IIP general index

## 9 Summary and discussion

This article establishes the asymptotic theory of Bayes factors when the models to be compared are systems of *SDE*'s consisting of time-dependent covariates and random effects, assuming that the number of individuals as well as the domains of observations of the individuals increase indefinitely. Different initial values for different *SDE*'s are also allowed. The only instance of related effort in this direction is that of Maitra and Bhattacharya (2018). The main difference of our undertaking with that of Maitra and Bhattacharya (2018) is that they assumed the domains of observations to be fixed for the individuals, a consequence being that incorporation of random effects in their model was not possible from the asymptotic perspective. Moreover, in their case, a single set of covariates was associated with all the individuals, but here our random effects set-up allows different sets of time-dependent covariates for different individuals.

To proceed, we first needed to build an asymptotic theory of Bayes factors for comparing two individual *SDE*'s, rather than two systems of *SDE*'s, as the domain of observation expands. Our results in this regard, which help formulate our asymptotic theory for comparing two systems of *SDE*'s using Bayes factors, are perhaps also of independent interest, being possibly the first ever results in this direction of research. Although the relevant variance did not converge to zero when two individual *SDE*'s are compared, we are able to establish almost sure exponential convergence of the Bayes factor when the number of subjects are allowed to increase indefinitely. Importantly, our theory covers both *iid* and non-*iid* cases.

Our simulation studies associated with covariate selection demonstrate that Bayes factor yields consistent results even in non-asymptotic situations. Bayes factor analysis of a real data on company-wise national stock exchange also yielded plausible sets of covariates for the companies.

Note that our current asymptotic Bayes factor theory remains valid for comparison between *iid* and non-*iid* models. For instance, if the true model constitutes an *iid* system, then  $f_{0i} \equiv f_0 \equiv f_{\theta_0}$ ; the rest remains the same as the theory for our non-*iid* setting. The situation is analogous when the other model forms an *iid* system.

## **Acknowledgments**

We are thankful to Dr. Diganta Mukherjee for drawing our attention towards the website [www.nseindia.com](http://www.nseindia.com) at which the real data was available. The first author gratefully acknowledges her CSIR Fellowship, Govt. of India.



## Supplementary Material

Throughout, we refer to our main manuscript as MB.

### S-1 Proof of Lemma 2 of MB

Due to compactness of  $\Gamma$  it follows, using the form of  $\phi_{\xi_j}$  provided in (H5''), that the convergences (3.9), (3.10) and (3.11) of MB are uniform over  $\Gamma$ . The same form shows that the above integrals are continuous in  $\xi_1$ , for every  $T > 0$ . Hence, due to uniform convergence, the limits  $\bar{\phi}_{\xi_1}^{(1)}$ ,  $\bar{\phi}_{\xi_1}^{(2)}$  and  $\bar{\phi}_{\xi_0, \xi_1}^{(2)}$  are continuous in  $\xi_1$ .

### S-2 Proof of Lemma 3 of MB

The proofs of (i) – (iv) follow from (H1''), (H4''), the results (3.10) and (3.11) of MB following from (H5''), (3.1) and its asymptotic form (3.2) (with  $k = 2$ ), using the relation (2.8) of MB. To prove (v), note that, since for any  $k \geq 1$ , it holds, due to (H4''), (3.1) of MB and boundedness of  $\phi_{\xi_j}$  on  $[0, T]$ , that for  $j = 0, 1$ ,

$$E \int_0^T \left| \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} \right|^{2k} ds < \infty,$$

it follows from Theorem 7.1 of Mao (2011), page 39, that

$$E \left| \int_0^T \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} dW(s) \right|^{2k} \leq (k(2k-1))^k T^{k-1} E \int_0^T \left| \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} \right|^{2k} ds. \quad (\text{S-2.1})$$

Hence, using Chebychev's inequality, it follows that for any  $\epsilon > 0$ ,

$$\begin{aligned} P \left( \left| \frac{1}{T} \int_0^T \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} dW(s) \right| > \epsilon \right) \\ < \epsilon^{-2k} (k(2k-1))^k T^{-(k+1)} E \int_0^T \left| \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} \right|^{2k} ds. \end{aligned} \quad (\text{S-2.2})$$

In particular, if  $k = 2$  is chosen, then it follows from the above inequality, (H4''), (3.2) of MB, and boundedness of  $\phi_{\xi_j}$  on  $[0, T]$ , that

$$\sum_{T=1}^{\infty} P \left( \left| \frac{1}{T} \int_0^T \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} dW(s) \right| > \epsilon \right) < \infty,$$

proving that

$$\frac{1}{T} \int_0^T \frac{\phi_{\xi_j}(s) b_{\beta_j}(s, X(s))}{\sigma(s, X(s))} dW(s) \xrightarrow{a.s.} 0.$$

To prove (vi), first note that

$$\begin{aligned}
& E \left| \frac{1}{T} \int_0^T \left[ \frac{\phi_{\xi_j}^2(s) b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} - \phi_{\xi_j}^2(s) \kappa_j(\beta_j) \right] ds \right|^{2k} \\
& \leq T^{-1} E \int_0^T \phi_{\xi_j}^{4k}(s) \left| \frac{b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} - \kappa_j(\beta_j) \right|^{2k} ds \\
& \leq K_4 T^{-1} E \int_0^T \left| \frac{b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} - \kappa_j(\beta_j) \right|^{2k} ds, \tag{S-2.3}
\end{aligned}$$

for some finite constant  $K_4 > 0$ . The second last inequality is by Hölder's inequality, and the last inequality holds because  $\phi_{\xi_j}(t)$  is uniformly bounded on  $[0, \infty]$  thanks to compactness of  $\mathcal{Z}$  and continuity of the functions  $g_l; l = 1, \dots, p$ . Hence, for any  $\epsilon > 0$ ,

$$\begin{aligned}
& P \left( \left| \frac{1}{T} \int_0^T \frac{\phi_{\xi_j}^2(s) b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} ds - \kappa_j(\beta_j) \frac{1}{T} \int_0^T \phi_{\xi_j}^2(s) ds \right| > \epsilon \right) \\
& < K_4 \epsilon^{-2k} T^{-1} E \int_0^T \left| \frac{b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} - \kappa_j(\beta_j) \right|^{2k} ds.
\end{aligned}$$

In the same way as the proof of (v), it follows, using the above inequality, (3.3) and (3.2) of MB, that

$$\sum_{T=1}^{\infty} P \left( \left| \frac{1}{T} \int_0^T \frac{\phi_{\xi_j}^2(s) b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} ds - \kappa_j(\beta_j) \frac{1}{T} \int_0^T \phi_{\xi_j}^2(s) ds \right| > \epsilon \right) < \infty.$$

That is,

$$\frac{1}{T} \int_0^T \frac{\phi_{\xi_j}^2(s) b_{\beta_j}^2(s, X(s))}{\sigma^2(s, X(s))} ds - \kappa_j(\beta_j) \frac{1}{T} \int_0^T \phi_{\xi_j}^2(s) ds \rightarrow 0,$$

almost surely, as  $T \rightarrow \infty$ . Since, as  $T \rightarrow \infty$ ,  $\frac{1}{T} \int_0^T \phi_{\xi_j}^2(s) ds \rightarrow \bar{\phi}_{\xi_j}^{(2)}$  by (3.10) of MB, the result follows. Using (3.4) instead of (3.3), (vii) can be proved in the same way as (vi). The proofs of (viii) and (ix) follow from (v), (vi) and (vii), using the relation (2.8).

### S-3 Proof of Lemma 4 of MB

Using the Cauchy-Schwartz inequality twice we obtain

$$\begin{aligned}
& \frac{1}{T} \int_0^T E \left( \frac{\phi_{\xi_0}(s) \phi_{\xi_1}(s) b_{\beta_0}(s, X(s)) b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} \right) ds \\
& \leq \frac{1}{T} \int_0^T \sqrt{E \left( \frac{\phi_{\xi_0}^2(s) b_{\beta_0}^2(s, X(s))}{\sigma^2(s, X(s))} \right)} \times \sqrt{E \left( \frac{\phi_{\xi_1}^2(s) b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} \right)} ds \\
& = \sqrt{\kappa_0(\beta_0) \kappa_1(\beta_1) + O(\exp(T^3 - T^5))} \times \frac{1}{T} \int_0^T |\phi_{\xi_0}(s)| \times |\phi_{\xi_1}(s)| ds \\
& \leq \sqrt{\kappa_0(\beta_0) \kappa_1(\beta_1) + O(\exp(T^3 - T^5))} \times \left( \sqrt{\frac{1}{T} \int_0^T \phi_{\xi_0}^2(s) ds} \right) \times \left( \sqrt{\frac{1}{T} \int_0^T \phi_{\xi_1}^2(s) ds} \right). \tag{S-3.1}
\end{aligned}$$

Taking the limit of both sides of (S-3.1) as  $T \rightarrow \infty$ , using (ii) of Lemma 3 and the limits (3.10), the result follows.

## S-4 Proof of Lemma 5 of MB

For any  $h \in (0, t)$ ,

$$\begin{aligned}
\frac{I_t}{I_{t-h}} &= \frac{\int_{\Theta} \frac{f_{\theta_1,t}(X_{0,t})}{f_{\theta_0,t}(X_{0,t})} \pi(d\theta_1)}{\int_{\Theta} \frac{f_{\theta_1,t-h}(X_{0,t-h})}{f_{\theta_0,t-h}(X_{0,t-h})} \pi(d\theta_1)} \\
&= \frac{\int_{\Theta} \exp\left((U_{\theta_1,t} - U_{\theta_0,t}) - \frac{(V_{\theta_1,t} - V_{\theta_0,t})}{2}\right) \pi(d\theta_1)}{\int_{\Theta} \exp\left((U_{\theta_1,t-h} - U_{\theta_0,t-h}) - \frac{(V_{\theta_1,t-h} - V_{\theta_0,t-h})}{2}\right) \pi(d\theta_1)} \\
&= \int_{\Theta} \exp\left((U_{\theta_1,t-h} - U_{\theta_0,t-h}) - \frac{(V_{\theta_1,t-h} - V_{\theta_0,t-h})}{2}\right) \\
&\quad \exp\left(U_{\theta_1,t-h,t} - U_{\theta_0,t-h,t} - \frac{(V_{\theta_1,t-h,t} - V_{\theta_0,t-h,t})}{2}\right) \pi(d\theta_1) \\
&\quad \times \frac{\int_{\Theta} \exp\left((U_{\theta_1,t-h} - U_{\theta_0,t-h}) - \frac{(V_{\theta_1,t-h} - V_{\theta_0,t-h})}{2}\right) \pi(d\theta_1)}{\int_{\Theta} \exp\left((U_{\theta_1,t-h} - U_{\theta_0,t-h}) - \frac{(V_{\theta_1,t-h} - V_{\theta_0,t-h})}{2}\right) \pi(d\theta_1)} \\
&= E \left[ \exp\left(\int_{t-h}^t \frac{\phi_{\xi_1}(s)b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \int_{t-h}^t \frac{\phi_{\xi_0}(s)b_{\beta_0}(s, X(s))}{\sigma^2(s, X(s))} dX(s) \right. \right. \\
&\quad \left. \left. - \left(\frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_1}^2(s)b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} ds - \frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_0}^2(s)b_{\beta_0}^2(s, X(s))}{\sigma^2(s, X(s))} ds\right) \right) \middle| \mathcal{F}_{t-h} \right] \\
&= \frac{E \left[ \exp\left(\int_{t-h}^t \frac{\phi_{\xi_1}(s)b_{\beta_1}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_1}^2(s)b_{\beta_1}^2(s, X(s))}{\sigma^2(s, X(s))} ds\right) \middle| \mathcal{F}_{t-h} \right]}{\exp\left(\int_{t-h}^t \frac{\phi_{\xi_0}(s)b_{\beta_0}(s, X(s))}{\sigma^2(s, X(s))} dX(s) - \frac{1}{2} \int_{t-h}^t \frac{\phi_{\xi_0}^2(s)b_{\beta_0}^2(s, X(s))}{\sigma^2(s, X(s))} ds\right)} \\
&= \frac{\hat{f}_{t-h,t}(X_{t-h,t})}{f_{\theta_0,t-h,t}(X_{t-h,t})}. \tag{S-4.1}
\end{aligned}$$

Hence, the result holds.

## S-5 Proof of Theorem 6 of MB

Let us consider

$$S_{Tq_n} = Tq_n \sum_{r=0}^{n(T)-1} \left( J'_{rTq_n} + \tilde{\mathcal{K}}'_{rTq_n} \right), \tag{S-5.1}$$

where  $q_n = \frac{1}{n(T)}$ , where, given  $T > 0$ ,  $n(T)$  is the number of intervals partitioning  $[0, T]$  each of length  $\frac{T}{n(T)}$ . We assume that as  $T \rightarrow \infty$ ,  $\frac{T}{n(T)} \rightarrow 0$ .

It follows, using (H9''), that for any  $T > 0$ ,

$$\begin{aligned}
E \left( \frac{S_{Tq_n}}{T} \right) &\rightarrow \frac{1}{T} \int_0^T \frac{d}{dt} E_{\theta_0} (\log I_t) dt + \frac{1}{T} \int_0^T E_{\theta_0} \left[ \tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}) \right] dt \\
&= E_{\theta_0} \left( \frac{1}{T} \log I_T \right) + \frac{1}{T} \int_0^T E_{\theta_0} \left[ \tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}) \right] dt, \tag{S-5.2}
\end{aligned}$$

as  $n(T) \rightarrow \infty$ , for any given  $T > 0$ . Also, since due to (4.26) of MB,  $E \left( J'_{rTq_n} + \tilde{\mathcal{K}}'_{rTq_n} | \mathcal{F}_{rTq_n} \right) = 0$ , we must have  $E \left( J'_{rTq_n} + \tilde{\mathcal{K}}'_{rTq_n} \right) = E \left[ E \left( J'_{rTq_n} + \tilde{\mathcal{K}}'_{rTq_n} | \mathcal{F}_{rTq_n} \right) \right] = 0$ , for any  $r, T, n(T)$ . Hence,  $E \left( \frac{S_{Tq_n}}{T} \right) = 0$  for any  $T, n(T)$ . Thus, it follows from (S-5.2), that

$$E_{\theta_0} \left( \frac{1}{T} \log I_T \right) + \frac{1}{T} \int_0^T E_{\theta_0} \left[ \tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}) \right] dt \rightarrow 0, \quad \text{as } T \rightarrow \infty. \quad (\text{S-5.3})$$

We now deal with the second term of the left hand side of (S-5.3). Since, by (H6''),

$$\pi \left( \theta_1 : \inf_t \tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) \geq \delta \right) = 1,$$

it holds that  $\tilde{\mathcal{K}}'_t(f_{\theta_0}, f_{\theta_1}) \geq \delta$  for all  $t$  with probability 1, so that

$$\tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}) = \tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}_{A_t(\delta)}),$$

where  $A_t(\delta)$  is given by (4.19) of MB. The  $Q^*$  property implies that

$$\liminf_T \frac{1}{T} \int_0^T E_{\theta_0} \left[ \tilde{\mathcal{K}}'_t(f_{\theta_0}, \hat{f}) \right] dt \geq \delta. \quad (\text{S-5.4})$$

The results (S-5.3) and (S-5.4) imply that

$$\limsup_T E_{\theta_0} \left( \frac{1}{T} \log I_T \right) \leq -\delta. \quad (\text{S-5.5})$$

Now observe that

$$\begin{aligned} I_T &= \int_{\Theta} \exp(U_{\theta_1, T} - U_{\theta_0, T}) \times \exp \left\{ -\frac{1}{2} (V_{\theta_1, T} - V_{\theta_0, T}) \right\} \pi(d\theta_1) \\ &\geq \int_{\mathcal{N}_0(c)} \exp \left\{ T \left( \frac{U_{\theta_1, T}}{T} - \frac{U_{\theta_0, T}}{T} \right) \right\} \\ &\quad \times \exp \left\{ -\frac{T}{2} \left( \frac{V_{\theta_1, T}}{T} - \frac{V_{\theta_0, T}}{T} \right) \right\} \pi(d\theta_1), \end{aligned} \quad (\text{S-5.6})$$

where  $c > 0$ , and

$$\begin{aligned} \mathcal{N}_0(c) &= \{ \theta_1 \in \Theta : \delta \leq \bar{K}^\infty(f_{\theta_0}, f_{\theta_1}) \leq \delta + c \} \\ &= \left\{ \theta_1 \in \Theta : \delta \leq \frac{\bar{\phi}_{\xi_0}^{(2)}}{2} \kappa_0(\beta_0) - \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1) + \frac{\bar{\phi}_{\xi_1}^{(2)}}{2} \kappa_1(\beta_1) \leq \delta + c \right\}, \end{aligned}$$

the second line following from (4.15) of MB. Using Jensen's inequality, we obtain

$$\begin{aligned} \frac{1}{T} \log(I_T) &\geq \int_{\mathcal{N}_0(c)} \left[ \left( \frac{U_{\theta_1, T}}{T} - \frac{U_{\theta_0, T}}{T} \right) \right. \\ &\quad \left. - \frac{1}{2} \left( \frac{V_{\theta_1, T}}{T} - \frac{V_{\theta_0, T}}{T} \right) \right] \pi(d\theta_1). \end{aligned} \quad (\text{S-5.7})$$

By (vi) – (ix) of Lemma 3 of MB, the integrand of the right hand side of the above inequality, which we denote by  $g_{X_T}(\theta_1)$ , converges to  $g(\theta_1) = - \left[ \frac{\bar{\phi}_{\xi_1}^{(2)}}{2} \kappa_1(\beta_1) - \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \beta_1) + \frac{\bar{\phi}_{\xi_0}^{(2)}}{2} \kappa_0(\beta_0) \right]$ , pointwise for every  $\theta_1$ , given any path of the process  $X$  in the complement of the null set. Due to (H1''), (H4'')

and (3.2) of MB,  $\sup_T E_{\theta_1} [g_{X_T}(\theta_1)]^2 < \infty$ , so that  $\{g_{X_T}(\theta_1)\}_{T=1}^\infty$  is uniformly integrable. Hence,

$$\int_{\mathcal{N}_0(c)} g_{X_T}(\theta_1) \pi(d\theta_1) \rightarrow \int_{\mathcal{N}_0(c)} g(\theta_1) \pi(d\theta_1),$$

given any path of the process  $X$  in the complement of the null set. Let us denote the left hand side of the above by  $H_{X_T}$  let  $H$  denote the right hand side. We just proved that  $H_{X_T}$  converges to  $H$  almost surely. Now observe that

$$\begin{aligned} \sup_T E_{\theta_0} [H_{X_T}]^2 &= \sup_T E_{\theta_0} \left[ \int_{\mathcal{N}_0(c)} g_{X_T}(\theta_1) \pi(d\theta_1) \right]^2 \\ &\leq \int_{\mathcal{N}_0(c)} \sup_T E_{\theta_0} [g_{X_T}(\theta_1)]^2 \pi(d\theta_1). \end{aligned}$$

Again, due to (H1''), (H4'') and (3.2) of MB, the last expression is finite, proving uniform integrability of  $\{H_{X_T}\}_{T=1}^\infty$ . Hence,

$$\lim_{T \rightarrow \infty} E_{\theta_0} (H_{X_T}) = E_{\theta_0} (H) = H.$$

It follows that

$$\begin{aligned} \liminf_{T \rightarrow \infty} E_{\theta_0} \left[ \frac{1}{T} \log(I_T) \right] &\geq \liminf_{T \rightarrow \infty} E_{\theta_0} (H_{X_T}) = E_{\theta_0} (H) \\ &= \int_{\mathcal{N}_0(c)} g(\theta_1) \pi(d\theta_1) \\ &\geq -(\delta + c) \pi(\mathcal{N}_0(c)) \\ &\geq -(\delta + c). \end{aligned} \tag{S-5.8}$$

Since the above holds for arbitrary  $c > 0$ , it holds that

$$\liminf_{T \rightarrow \infty} E_{\theta_0} \left( \frac{1}{T} \log I_T \right) \geq -\delta. \tag{S-5.9}$$

Thus (S-5.5) and (S-5.9) together help us conclude that

$$E_{\theta_0} \left( \frac{1}{T} \log I_T \right) \rightarrow -\delta,$$

as  $T \rightarrow \infty$ .

We now show that the variance of  $\frac{1}{T} \log I_T$  is  $O(1)$ , as  $T \rightarrow \infty$ . First note, due to compactness of  $\Theta$ , the mean value theorem for integrals ensure existence of  $\hat{\theta}_1 = (\hat{\beta}_1, \hat{\xi}_1) = \hat{\theta}_1(W) \in \Theta$ , depending on the Wiener process  $W$  such that

$$\log I_T = \left( U_{\hat{\theta}_1, T} - U_{\theta_0, T} \right) - \frac{1}{2} \left( V_{\hat{\theta}_1, T} - V_{\theta_0, T} \right). \tag{S-5.10}$$

Now note that the results presented in Lemma 3 of MB continue to hold even when  $\theta_1$  is replaced with  $\hat{\theta}_1$ . Specifically, the following hold in addition to the results of Lemma 3:

$$\begin{aligned} \frac{V_{\hat{\theta}_1, T}}{T} - \bar{\phi}_{\hat{\xi}_1}^{(2)} \kappa_1(\hat{\beta}_1) &\xrightarrow{a.s.} 0; \\ \frac{U_{\hat{\theta}_1, T}}{T} - \bar{\phi}_{\xi_0, \hat{\xi}_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) &\xrightarrow{a.s.} 0. \end{aligned}$$

It follows that, as  $T \rightarrow \infty$ ,

$$\ell_T = \frac{1}{T} \log I_T - \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) + \frac{1}{2} \bar{\phi}_{\xi_1}^{(2)} \kappa_1(\hat{\beta}_1) + \frac{1}{2} \bar{\phi}_{\xi_0}^{(2)} \kappa_0(\beta_0) \xrightarrow{a.s.} 0.$$

By uniform integrability arguments, which follow in similar lines as the proofs of Lemmas 1 and 10 of Maitra and Bhattacharya (2018) using (H4''), compactness, and Cauchy-Schwartz, it holds that

$$Var_{\theta_0}(\ell_T) \rightarrow 0, \text{ as } T \rightarrow \infty. \quad (\text{S-5.11})$$

Hence,

$$\begin{aligned} Var_{\theta_0} \left( \frac{1}{T} \log I_T \right) &= Var_{\theta_0} \left( \ell_T + \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_1}^{(2)} \kappa_1(\hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_0}^{(2)} \kappa_0(\beta_0) \right) \\ &= Var_{\theta_0}(\ell_T) + Var_{\theta_0} \left( \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_1}^{(2)} \kappa_1(\hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_0}^{(2)} \kappa_0(\beta_0) \right) \\ &\quad + 2Cov_{\theta_0} \left( \ell_T, \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_1}^{(2)} \kappa_1(\hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_0}^{(2)} \kappa_0(\beta_0) \right). \end{aligned} \quad (\text{S-5.12})$$

By (S-5.11), the first term of (S-5.12) goes to zero as  $T \rightarrow \infty$ , and the third, covariance term tends to zero by Cauchy-Schwartz and (S-5.11). In other words, as  $T \rightarrow \infty$ ,

$$\left| Var_{\theta_0} \left( \frac{1}{T} \log I_T \right) - Var_{\theta_0} \left( \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_1}^{(2)} \kappa_1(\hat{\beta}_1) \right) \right| \rightarrow 0. \quad (\text{S-5.13})$$

However,

$$Var_{\theta_0} \left( \bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_1}^{(2)} \kappa_1(\hat{\beta}_1) \right) \rightarrow 0,$$

unless  $\bar{\phi}_{\xi_0, \xi_1}^{(2)} \bar{\kappa}(\beta_0, \hat{\beta}_1) - \frac{1}{2} \bar{\phi}_{\xi_1}^{(2)} \kappa_1(\hat{\beta}_1)$  is constant almost surely. It then follows from (S-5.13) that

$$Var_{\theta_0} \left( \frac{1}{T} \log I_T \right) = O(1), \text{ as } T \rightarrow \infty. \quad (\text{S-5.14})$$

## S-6 Proof of Theorem 8 of MB

In our set-up it follows from (2.7) of MB that

$$\frac{1}{nT} \log \tilde{I}_{n,T} = \frac{1}{n} \sum_{i=1}^n \frac{1}{T} \log I_{i,T}. \quad (\text{S-6.1})$$

In the *iid* case, given  $T > 0$ , using the above form, it follows by the strong law of large numbers, that

$$\lim_{n \rightarrow \infty} \frac{1}{nT} \log \tilde{I}_{n,T} = E_{\theta_0} \left( \frac{1}{T} \log I_{i,T} \right), \quad (\text{S-6.2})$$

almost surely. Now, in the *iid* situation, for each  $i$ ,  $E_{\theta_0} \left( \frac{1}{T} \log I_{i,T} \right) \rightarrow -\delta$ , as  $T \rightarrow \infty$ . Hence, taking limit as  $T \rightarrow \infty$  on both sides of (S-6.2) yields

$$\lim_{T \rightarrow \infty} \lim_{n \rightarrow \infty} \frac{1}{nT} \log \tilde{I}_{n,T} = \lim_{T \rightarrow \infty} E_{\theta_0} \left( \frac{1}{T} \log I_{i,T} \right) = -\delta, \quad (\text{S-6.3})$$

almost surely, proving the theorem.

## S-7 Proof of Lemma 10 of MB

Note that, due to compactness of  $\mathfrak{X}$  and  $\mathfrak{Z}$  and continuity of the covariates in time  $t$ , there exists  $x^* \in \mathfrak{X}$  and  $z^* \in \mathfrak{Z}$ , such that

$$\sup_{x \in \mathfrak{X}, z \in \mathfrak{Z}} \left| \frac{1}{T} E_{\theta_0} (\log I_{x,T,z}) + \delta(x, z) \right| = \left| \frac{1}{T} E_{\theta_0} (\log I_{x^*,T,z^*}) + \delta(x^*, z^*) \right| \rightarrow 0, \quad (\text{S-7.1})$$

as  $T \rightarrow \infty$ , where the convergence is due to (6.2) of MB. Also,  $\frac{1}{T} E_{\theta_0} (\log I_{x,T,z})$  is clearly continuous in  $(x, z)$  for every  $T > 0$  (the proof of this follows in the same way as that of Theorem 5 of Maitra and Bhattacharya (2016)). Combining this with the uniform convergence (S-7.1) it follows that  $\delta(x, z)$  is also continuous in  $(x, z)$ .

## S-8 Proof of Lemma 11 of MB

Note that the limit (6.3) of MB can be represented as

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=0}^{n-1} \varrho_{\delta} \left( \frac{r}{n} \right), \quad (\text{S-8.1})$$

where  $\varrho_{\delta} : [0, 1] \mapsto \mathbb{R}^+$  is some continuous function satisfying  $\varrho_{\delta} \left( \frac{r}{n} \right) = \delta(x^{r+1}, z_{r+1})$  for  $r = 0, \dots, n-1$ . For the remaining points  $y \in [0, 1]$ , we set  $\varrho_{\delta}(y) = \delta(x, z)$ , where  $(x, z) \in \mathfrak{X} \times \mathfrak{Z}$  is such that  $\varrho_{\delta}(y)$  is continuous in  $y \in [0, 1]$ . Since  $\delta(x, z)$  is continuous in  $(x, z)$ ,  $\varrho_{\delta}(y)$  can be thus constructed. Note that, it is possible to relate  $y \in [0, 1]$  to  $(x, z) \in \mathfrak{X} \times \mathfrak{Z}$  by some continuous mapping  $G : \mathfrak{X} \times \mathfrak{Z} \mapsto [0, 1]$ , taking  $(x, z)$  to  $y$ . Thus,  $\delta^{\infty}$  in (6.3) of MB is the limit of the Riemann sum (S-8.1) associated with the continuous function  $\varrho_{\delta}$ ; the limit is given by the integral  $\int_0^1 \varrho_{\delta}(y) dy$ . Since the domain of integration is  $[0, 1]$ , it follows, using continuity of  $\varrho_{\delta}$ , that the integral is finite. Observe that for any given sequence  $\{(x^i, z_i)\}_{i=1}^{\infty}$ , one can construct a continuous function  $\varrho_{\delta}$  such that  $\delta^{\infty} = \int_0^1 \varrho_{\delta}(y) dy$ . In other words,  $\delta^{\infty}$  exists for all sequences  $\{(x^i, z_i)\}_{i=1}^{\infty}$ .

## S-9 Proof of Theorem 12 of MB

For given  $T > 0$ , it follows from (S-5.14), compactness of  $\Theta$ ,  $\mathfrak{X}$ ,  $\mathfrak{Z}$ , and continuity of the relevant functions  $\phi_{\xi_j}, b_{\beta_j}, g_1, \dots, g_p, \kappa_0, \kappa_1$  and  $\bar{\kappa}$ , that

$$\sup_{x \in \mathfrak{X}, z \in \mathfrak{Z}} \text{Var}_{\theta_0} \left( \frac{1}{T} \log I_{x,T,z} \right) < \infty. \quad (\text{S-9.1})$$

Hence, given  $T > 0$ ,

$$\sum_{i=1}^{\infty} \frac{\text{Var}_{\theta_0} \left( \frac{1}{T} \log I_{i,T} \right)}{i^2} < \infty.$$

It then follows due to Kolmogorov's strong law of large numbers for independent random variables that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{T} \log I_{i,T} \right) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n E_{\theta_0} \left( \frac{1}{T} \log I_{i,T} \right), \quad (\text{S-9.2})$$

almost surely.

Now observe that the right hand side of (S-9.2) admits the following representation

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=0}^{n-1} \check{\varrho} \left( \frac{r}{n}, T \right), \quad (\text{S-9.3})$$

where  $\check{\varrho}(\cdot, T) : [0, 1] \mapsto \mathbb{R}$  is some continuous function depending upon  $T$  with  $\check{\varrho}(\frac{r}{n}, T) = E_{\theta_0} \left( \frac{1}{T} \log I_{r+1, T} \right)$ .

Since  $E_{\theta_0} \left( \frac{1}{T} \log I_{x, T, z} \right)$  is continuous in  $(x, z)$ ,  $\check{\varrho}(y, T)$  can be constructed as in Lemma 11 of MB. Then, for almost all  $y \in [0, 1]$ ,  $\check{\varrho}(y, T) \rightarrow -\delta(x, z)$  as  $T \rightarrow \infty$ , for appropriate  $(x, z)$  associated with  $y$  via  $y = G(x, z)$  as in Lemma 11. Also, it follows from (S-5.10) that  $\check{\varrho}(\cdot, T)$  so constructed is uniformly bounded in  $T > 0$ . Thus, the conditions of the dominated convergence theorem are satisfied.

Since (S-9.3) is nothing but the Riemann sum associated with  $\check{\varrho}(\cdot)$ , it follows that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n E_{\theta_0} \left( \frac{1}{T} \log I_{i, T} \right) = \int_0^1 \check{\varrho}(y, T) dy. \quad (\text{S-9.4})$$

By construction of  $\check{\varrho}(y, T)$ , the dominated convergence theorem holds for the right hand side of (S-9.4). Hence,

$$\begin{aligned} \lim_{T \rightarrow \infty} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n E_{\theta_0} \left( \frac{1}{T} \log I_{i, T} \right) &= \lim_{T \rightarrow \infty} \int_0^1 \check{\varrho}(y, T) dy = \int_0^1 \lim_{T \rightarrow \infty} \check{\varrho}(y, T) dy \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \lim_{T \rightarrow \infty} E_{\theta_0} \left( \frac{1}{T} \log I_{i, T} \right) = - \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta(x^i, z_i) = -\delta^\infty. \end{aligned} \quad (\text{S-9.5})$$

Combining (S-9.5) with (S-9.2) it follows that

$$\lim_{T \rightarrow \infty} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{T} \log I_{i, T} \right) = \lim_{T \rightarrow \infty} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n E_{\theta_0} \left( \frac{1}{T} \log I_{i, T} \right) = -\delta^\infty, \quad (\text{S-9.6})$$

almost surely.

## References

- Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In B. N. Petrov and F. Csaki, editors, *Second International Symposium on Information Theory*, pages 267–281, Budapest. Akademiai Kiado. Reprinted in S. Kotz and N. L. Johnson (Eds) (1992). *Breakthroughs in Statistics Volume I: Foundations and Basic Theory*, pp. 610–624. Springer-Verlag.
- Claeskens, G. and Hjort, N. L. (2008). *Model Selection and Model Averaging*. Cambridge University Press, Cambridge, UK.
- Delattre, M., Genon-Catalot, V., and Samson, A. (2013). Maximum Likelihood Estimation for Stochastic Differential Equations with Random Effects. *Scandinavian Journal of Statistics*, **40**, 322–343.
- Dey, D. K., Ghosh, S. K., and Mallick, B. K. (2000). *Generalized Linear Models: A Bayesian Perspective*.
- Fuchs, C. (2013). *Inference for Diffusion Processes: With Applications in Life Sciences*. Springer, New York.
- Iacus, S. M. (2008). *Simulation and Inference for Stochastic Differential Equations: With R Examples*. Springer, New York.



- Jiang, J. (2007). *Linear and Generalized Linear Mixed Models and Their Applications*. Springer, New York, USA.
- Kass, R. E. and Raftery, R. E. (1995). Bayes factors. *Journal of the American Statistical Association*, **90**(430), 773–795.
- Kontoyiannis, I. and Meyn, S. P. (2003). Spectral Theory and Limit Theorems for Geometrically Ergodic Markov Processes. *The Annals of Applied Probability*, **13**, 304–362.
- Leander, J., Almquist, J., Ahlström, C., Gabrielsson, J., and Jirstrand, M. (2015). Mixed Effects Modeling Using Stochastic Differential Equations: Illustrated by Pharmacokinetic Data of Nicotinic Acid in Obese Zucker Rats. *The AAPS Journal*, **17**, 586–596.
- Maitra, T. and Bhattacharya, S. (2015). On Bayesian Asymptotics in Stochastic Differential Equations with Random Effects. *Statistics and Probability Letters*, **103**, 148–159. Also available at “<http://arxiv.org/abs/1407.3971>”.
- Maitra, T. and Bhattacharya, S. (2016). On Asymptotics Related to Classical Inference in Stochastic Differential Equations with Random Effects. *Statistics and Probability Letters*, **110**, 278–288. Also available at “<http://arxiv.org/abs/1407.3968>”.
- Maitra, T. and Bhattacharya, S. (2018). Asymptotic Theory of Bayes Factor in Stochastic Differential Equations: Part I. ArXiv Preprint.
- Mao, X. (2011). *Stochastic Differential Equations and Applications*. Woodhead Publishing India Private Limited, New Delhi, India.
- Müller, A., Scealy, J. L., and Welsh, A. H. (2013). Model Selection in Linear Mixed Models. *Statistical Science*, **28**, 135–167.
- Oravecz, Z., Tuerlinckx, F., and Vandekerckhove, J. (2011). A hierarchical latent stochastic differential equation model for affective dynamics. *Psychological Methods*, **16**, 468–490.
- Overgaard, R. V., Jonsson, N., Tornøe, C. W., and Madsen, H. (2005). Non-Linear Mixed-Effects Models with Stochastic Differential Equations: Implementation of an Estimation Algorithm. *Journal of Pharmacokinetics and Pharmacodynamics*, **32**, 85–107.
- Roberts, G. and Stramer, O. (2001). On Inference for Partially Observed Nonlinear Diffusion Models Using the Metropolis-hastings Algorithm. *Biometrika*, **88**, 603–621.
- Schwarz, G. (1978). Estimating the Dimension of a Model. *The Annals of Statistics*, **6**, 461–464.
- Sivaganesan, S. and Lingham, R. T. (2002). On the Asymptotic of the Intrinsic and Fractional Bayes Factors for Testing Some Diffusion Models. *Annals of the Institute of Statistical Mathematics*, **54**, 500–516.