Directional derivatives of the singular values of matrices depending on several real parameters

Juan-Miguel Gracia[∗]

May 14, 2020

Abstract

In this document I recapitulate some results by Hiriart-Urruty and Ye[\[2\]](#page-10-0) concerning the properties of differentiability and the existence of directional derivatives of the multiple eigenvalues of a complex Hermitian matrix function of several real variables, where the eigenvalues are supposed in a decreasing order. Another version of these results was obtained by Ji-guang $Sun[6-8]$ $Sun[6-8]$.

Contents

1 Differentiability of the eigenvalues of a complex Hermitian matrix

We will denote by $\Lambda(C)$ the spectrum or set of eigenvalues of any complex square matrix C. Let Ω be an open subset of \mathbb{R}^p and let $A: \Omega \to \mathbb{C}^{n \times n}$ be a matrix function of class C^1 such that for every $x \in \Omega$ the matrix $A(x)$ is Hermitian, i.e. $A(x)^* = A(x)$ where ^{*} denotes the conjugate transpose. As it is well known the eigenvalues of $A(x)$ are real numbers; thus, there exist n real functions defined on Ω , $\lambda_1, \ldots, \lambda_n$, such that for all $x \in \Omega$,

 $\lambda_1(x) \geq \lambda_2(x) \geq \cdots \geq \lambda_n(x)$

[∗]Department of Applied Mathematics and Statistics and O.R., University of the Basque Country UPV/EHU, Faculty of Pharmacy, Paseo de la Universidad, 7, 01006 Vitoria-Gasteiz, Spain, juanmiguel.gracia@ehu.es

are the eigenvalues of $A(x)$. Let $m \in \{1, \ldots, n\}$; it is easy to prove that the function $\lambda_m : \Omega \to \mathbb{R}$ is continuous. When the eigenvalue $\lambda_m(x_0)$ of $A(x_0)$ is simple, the function λ_m is differentiable at $x_0 \in \Omega$. But in case of $\lambda_m(x_0)$ is a multiple eigenvalue of $A(x_0)$, λ_m can be nondifferentiable at x_0 . For example [\[4\]](#page-10-3), let

$$
A(x_1, x_2) := \begin{pmatrix} x_1 & \text{i}x_2 \\ -\text{i}x_2 & -x_1 \end{pmatrix}
$$

be for $(x_1, x_2) \in \mathbb{R}^2$. It is obvious that for each $(x_1, x_2) \in \mathbb{R}^2$ the matrix $A(x_1, x_2)$ is Hermitian. Then

$$
\begin{vmatrix} \lambda - x_1 & -ix_2 \\ ix_2 & \lambda + x_1 \end{vmatrix} = \lambda^2 - x_1^2 - x_2^2;
$$

hence the eigenvalues of $A(x_1, x_2)$ are $\pm \sqrt{x_1^2 + x_2^2}$. Observe that the matrix

$$
A(0,0) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}
$$

has a double eigenvalue; but neither the function $\lambda_1(x_1, x_2) = \sqrt{x_1^2 + x_2^2}$, nor the function $\lambda_2(x_1, x_2) = -\sqrt{x_1^2 + x_2^2}$ are differentiable at $(0, 0)$.

Let $d \in \mathbb{R}^p$ be a unitary vector, i.e. $||d||_2 = 1$, where $||\cdot||_2$ denotes the Euclidean norm. The directional derivative of the function λ_m at the point x_0 with respect to d is defined as the limit

$$
\lambda'_m(x_0, d) := \lim_{t \to 0^+} \frac{\lambda_m(x_0 + td) - \lambda_m(x_0)}{t}
$$

whenever this limit exists.

Based on techniques and results from convex and nonsmooth analysis (in Clarke's sense), Hiriart-Urruty and Ye proved Theorems [1](#page-1-0) and [2.](#page-2-0) See [\[2,](#page-10-0) Theorem 4.5].

Theorem 1 For all $x_0 \in \Omega$, for all unitary vector $d \in \mathbb{R}^p$, and for all $m \in \mathbb{R}^p$ $\{1, \ldots, n\}$, there exists always

$$
\lambda_m'(x_0, d).
$$

Moreover, it can be proved that $\lambda'_m(x_0, d)$ is equal to a determined eigenvalue of a matrix constructed from $A(x_0)$ and d in the following way: For each $x_0 \in \Omega$, there is a unitary matrix $U = [u_1, \ldots, u_n]$ such that

$$
U^*A(x_0)U=\mathrm{diag}\left(\lambda_1(x_0),\ldots,\lambda_n(x_0)\right).
$$

Suppose that $\lambda_m(x_0)$ is a multiple eigenvalue of $A(x_0)$, of multiplicity r_m . Introduce two integers $i_m \geq 1, j_m \geq 0$ to precise the position that $\lambda_m(x_0)$ occupies among the r_m repeated eigenvalues that are equal to it. Consider the detailed arrangement of the eigenvalues of $A(x_0)$:

$$
\lambda_1(x_0) \geq \cdots \geq \lambda_{m-i_m}(x_0) > \lambda_{m-i_m+1}(x_0) = \cdots = \lambda_m(x_0)
$$

= $\lambda_{m+1}(x_0) = \cdots = \lambda_{m+j_m}(x_0) > \lambda_{m+j_m+1}(x_0) \geq \cdots \geq \lambda_n(x_0)$

That is to say, j_m is the number of eigenvalues placed after the subscript m that are equal to $\lambda_m(x_0)$; whereas i_m is the number of eigenvalues placed before m that are equal to $\lambda_m(x_0)$, plus one (we put $\lambda_m(x_0)$ in this list). Hence, j_m may be zero, $i_m \geq 1$, and $i_m + j_m = r_m$. When $m = 1$, i.e. if we are considering $\lambda_1(x_0)$, we have $i_1 = 1, j_1 = r_1 - 1$. When $m = n$, i.e. for $\lambda_n(x_0)$, we have $i_n = r_n$, $j_n = 0$. In case $\lambda_m(x_0)$ is a simple eigenvalue, $i_m = 1$, $j_m = 0$. Although the notation does not indicate it, the numbers i_m, j_m and r_m depend on x_0 .

Let U_2 be the $n \times r_m$ matrix formed by the $(m - i_m + 1)$ th, ..., $(m + j_m)$ th columns of the matrix U :

$$
U_2 := [u_{m-i_m+1}, \ldots, u_{m+j_m}];
$$

i.e. U_2 is formed by r_m orthonormal eigenvectors associated with the eigenvalue $\lambda_m(x_0)$ of $A(x_0)$. For each $j \in \{1, \ldots, p\}$ define

$$
\frac{\partial A}{\partial x_j}(x_0) = \left(\frac{\partial a_{ik}}{\partial x_j}(x_0)\right)
$$

 $a_{ik}(x)$ being the entries of $A(x)$. We will call $F'(d)$ to the $r_m \times r_m$ matrix

$$
F'(d) := U_2^* \left(\sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0) \right) U_2
$$

for every unitary vector $d = (d_1, \ldots, d_p) \in \mathbb{R}^p$. Given that

$$
\overline{\left(\frac{\partial a_{ik}}{\partial x_j}\right)} = \frac{\partial \bar{a}_{ik}}{\partial x_j} = \frac{\partial a_{ki}}{\partial x_j},
$$

we have that the matrix $\frac{\partial A}{\partial x_j}$ is Hermitian, and so is $F'(d)$; indeed,

$$
F'(d)^* = U_2^* \left(\sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0)\right)^* U_2
$$

=
$$
U_2^* \left(\sum_{j=1}^p d_j \left[\frac{\partial A}{\partial x_j}(x_0)\right]^*\right) U_2
$$

=
$$
U_2^* \left(\sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0)\right) U_2 = F'(d).
$$

Therefore, the eigenvalues of $F'(d)$ are real numbers.

Theorem 2 The directional derivative $\lambda'_m(x_0, d)$ is given by

$$
\lambda'_m(x_0, d) = \mu_{i_m}\big(F'(d)\big)
$$

where $\mu_{i_m}(F'(d))$ is the i_mth eigenvalue of $F'(d)$ when the eigenvalues are arranged in a decreasing order:

$$
\mu_1(F'(d)) \geqslant \cdots \geqslant \mu_{r_m}(F'(d)).
$$

A theorem related to Theorem [2](#page-2-0) was proved by Ji-guang Sun [\[8,](#page-10-2) Theorem 3.1] applying the implicit function theorem and the Rellich theorem.

Theorem 3 The function

$$
t_m(x) := \lambda_{m-i_m+1}(x) + \dots + \lambda_m(x) + \dots + \lambda_{m+j_m}(x), \quad x \in \Omega
$$

is differentiable at x_0 .

See [\[2,](#page-10-0) Corollary 4.3] for a proof of this theorem.

Corollary 4 There exists a neighborhood V of $x_0, V \subset \Omega$, in which the function

$$
t_m(x) := \lambda_{m-i_m+1}(x) + \dots + \lambda_m(x) + \dots + \lambda_{m+j_m}(x)
$$

is differentiable.

PROOF. Let $V \subset \Omega$ be a neighborhood of x_0 , sufficiently small so that the inequalities

$$
\lambda_{m-i_m}(x) > \lambda_{m-i_m+1}(x), \quad \lambda_{m+j_m}(x) > \lambda_{m+j_m+1}(x)
$$

hold when $x \in V$. Let x_1 be any point of V. Then the arrangement of the eigenvalues of $A(x_1)$

$$
\lambda_{m-i_m+1}(x_1) \geqslant \cdots \geqslant \lambda_{m+j_m}(x_1)
$$

may have groups of equalities. In view of Theorem [3,](#page-3-1) the sum of the functions λ_i corresponding to each one of these groups, is differentiable at x_1 ; therefore, as t_m is the sum of these sums, we deduce that t_m is differentiable at x_1 .

2 Differentiability of the singular values of a complex matrix

Notation. Let m, n be positive integers and let $q := \min(m, n)$. Given $B \in$ $\mathbb{C}^{m \times n}$, let

$$
\sigma_1(B) \geqslant \sigma_2(B) \geqslant \cdots \geqslant \sigma_q(B)
$$

be the singular values of B. For each $k \in \{1, \ldots, q\}$, it is said that a pair of vectors of unit length $y_k \in \mathbb{C}^{m \times 1}$, $z_k \in \mathbb{C}^{n \times 1}$ are left and right singular vectors of B associated with the singular value $\sigma_k(B)$ if $Bz_k = \sigma_k(B)y_k$ and $B^*y_k = \sigma_k(B)z_k.$

Let $A: \Omega \to \mathbb{C}^{m \times n}$ be a matrix function of class C^1 . For each $x \in \Omega \subset \mathbb{R}^p$, let

$$
s_1(x) \geqslant \cdots \geqslant s_q(x)
$$
, with $q := \min(m, n)$,

be the singular values of the matrix $A(x)$ arranged in a decreasing order. Thus, we can define q functions $s_i \colon \Omega \to \mathbb{R}, i \in \{1, \ldots, q\}$. We are going to establish the properties of differentiability of these functions. By Wielandt's lemma, the $m + n$ eigenvalues of the Hermitian matrix

$$
M(x) := \begin{pmatrix} 0 & A(x) \\ A(x)^* & 0 \end{pmatrix} \in \mathbb{C}^{(m+n)\times (m+n)}
$$

$$
s_1(x) \geqslant \cdots \geqslant s_q(x) \geqslant 0 = \cdots = 0 \geqslant -s_q(x) \geqslant \cdots \geqslant -s_1(x)
$$

(it may have repeated intermediate zeros), for all $x \in \Omega$. Hence, the analogous results to Theorems [1,](#page-1-0) [2](#page-2-0) and [3](#page-3-1) for Hermitian matrices are true.

Theorem 5 Let $k \in \{1, ..., q\}$, $x_0 \in \Omega$, and $d \in \mathbb{R}^p$ be a unitary vector. Then there exists the directional derivative

$$
s'_k(x_0, d).
$$

Let $u \in \mathbb{C}^{m \times 1}$, $v \in \mathbb{C}^{n \times 1}$, where $u \neq 0$ or $v \neq 0$. Then

$$
\begin{pmatrix} u \\ v \end{pmatrix}
$$

is an eigenvector of

$$
H:=\begin{pmatrix}0&B\\B^*&0\end{pmatrix}
$$

associated with its eigenvalue $\sigma_k(B)$ if and only if

$$
Bv = \sigma_k(B)u,\tag{1}
$$

$$
B^*u = \sigma_k(B)v.
$$
 (2)

So, if $(y_k, z_k) \in \mathbb{C}^{m \times 1} \times \mathbb{C}^{n \times 1}$ is a pair of (left, right)-singular vectors of B associated with the singular value $\sigma_k(B)$, then

$$
\begin{pmatrix} y_k \\ z_k \end{pmatrix}
$$

is an eigenvector of H corresponding to its eigenvalue $\sigma_k(B)$.

Let $x_0 \in \Omega$ be a fixed point, and let $W \in \mathbb{C}^{(m+n)\times (m+n)}$ a unitary matrix that diagonalizes $M(x_0)$:

$$
W^*M(x_0)W = \begin{pmatrix} s_1(x_0) & & & & & 0 \\ & \ddots & & & & & \\ & & s_k(x_0) & & & \\ & & & \ddots & & \\ & & & & -s_k(x_0) & \\ & & & & & \ddots \\ 0 & & & & & & -s_1(x_0) \end{pmatrix}
$$
 (3)

Suppose that

$$
s_1(x_0) \ge \dots \ge s_{k-i_k}(x_0) > s_{k-i_k+1}(x_0) = \dots = s_k(x_0)
$$

= $s_{k+1}(x_0) = \dots = s_{k+j_k}(x_0) > s_{k+j_k+1}(x_0) \ge \dots \ge s_q(x_0) \ge \dots \ge -s_1(x_0)$

are the eigenvalues of $M(x_0)$, where $s_k(x_0)$ is a multiple eigenvalue of multiplicity $r_k = i_k + j_k$, i_k being the number of eigenvalues equal to $s_k(x_0)$ placed

are

before the rank $k+1$, and j_k is the number of eigenvalues equal to $s_k(x_0)$ situate after the rank k .

Call W_2 to the $(m+n) \times r_k$ matrix formed by the $(k-i_k+1)$ th,..., $(k+j_k)$ th columns of the matrix W. For each unitary vector $d = (d_1, \ldots, d_p) \in \mathbb{R}^p$, define

$$
F'(d) := W_2^* \left(\sum_{j=1}^p d_j \begin{bmatrix} O & \frac{\partial A}{\partial x_j}(x_0) \\ \left(\frac{\partial A}{\partial x_j}(x_0)\right)^* & O \end{bmatrix} \right) W_2,
$$

which is an $r_k \times r_k$ Hermitian matrix. Then, by Theorem [2,](#page-2-0) we have the next result.

Theorem 6 For each unitary vector $d = (d_1, \ldots, d_p) \in \mathbb{R}^p$

$$
s'_{k}(x_0,d) = \mu_{i_k}(F'(d)),
$$

 $\mu_{i_k}\big(F'(d)\big)$ being the i_k th eigenvalue of the matrix $F'(d)$ when we arrange the eigenvalues of this matrix in a decreasing order.

To facilitate the writing let W_2 be partitioned thus:

$$
W_2 = \begin{bmatrix} U_2 \\ V_2 \end{bmatrix}
$$

where

$$
U_2 := [u_{k-i_k+1}, \dots, u_{k+j_k}], \quad V_2 := [v_{k-i_k+1}, \dots, v_{k+j_k}],
$$

$$
U_2 \in \mathbb{C}^{m \times r_k}, V_2 \in \mathbb{C}^{n \times r_k}.
$$

Corollary 7 For each unitary vector $d = (d_1, \ldots, d_p) \in \mathbb{R}^p$ we have

$$
s'_k(x_0,d)=\mu_{i_k},
$$

where μ_{i_k} is the i_k th eigenvalue of

$$
U_2^* \left(\sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0) \right) V_2 + \left(U_2^* \left(\sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0) \right) V_2 \right)^*,
$$

when the eigenvalues are ranked in a decreasing order.

PROOF. Given that

$$
W_2 := \begin{bmatrix} U_2 \\ V_2 \end{bmatrix},
$$

the matrix $F'(d)$ is given by

$$
[U_2^*, V_2^*] \left(\sum_{j=1}^p d_j \begin{bmatrix} O & \frac{\partial A}{\partial x_j}(x_0) \\ \frac{\partial A}{\partial x_j}(x_0) \end{bmatrix}^* O \right) \begin{bmatrix} U_2 \\ V_2 \end{bmatrix}
$$

\n
$$
= \sum_{j=1}^p d_j \left[V_2^* \left[\frac{\partial A}{\partial x_j}(x_0) \right]^*, U_2^* \frac{\partial A}{\partial x_j}(x_0) \right] \left[\frac{U_2}{V_2} \right]
$$

\n
$$
= V_2^* \sum_{j=1}^p d_j \left[\frac{\partial A}{\partial x_j}(x_0) \right]^* U_2 + U_2^* \sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0) V_2
$$

\n
$$
= U_2^* \sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0) V_2 + \left(U_2^* \sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0) V_2 \right)^*.
$$
 (4)

The sum of all singular values that coalesce with $s_k(x_0)$ at x_0 is differentiable at x_0 . Even more it is true as we can see in the next theorem.

Theorem 8 The function

$$
t_k(x) := s_{k-i_k+1}(x) + \cdots + s_k(x) + \cdots + s_{k+j_k}(x)
$$

is differentiable in a neighborhood $V \subset \Omega$ of x_0 .

The neighborhood V is determined by the $x \in \Omega$ sufficient close to x_0 in order that the inequalities

$$
s_{k-i_k}(x) > s_{k-i_k+1}(x)
$$
 and $s_{k+j_k}(x) > s_{k+j_k+1}(x)$

hold.

From Corollary [7](#page-5-0) we can give another description of $s'_k(x_0, d)$ in terms of singular vectors of $A(x_0)$ associated with $s_k(x_0)$.

Theorem 9 With the previous notation, let

$$
Y = [y_{k-i_k+1}, \dots, y_{k+j_k}] \in \mathbb{C}^{m \times r_k}, \quad Z = [z_{k-i_k+1}, \dots, z_{k+j_k}] \in \mathbb{C}^{n \times r_k}
$$

be matrices of orthonormal columns and such that $(y_{\ell}, z_{\ell}) \in \mathbb{C}^{m \times 1} \times \mathbb{C}^{n \times 1}$ is a pair of (left, right)-singular vectors of $A(x_0)$ associated with the singular value $s_k(x_0)$ for $\ell \in \{k - i_k + 1, \ldots, k + j_k\}$. Then $s'_k(x_0, d)$ is equal to the i_k th eigenvalue of the $r_k \times r_k$ Hermitian matrix

$$
G := \frac{1}{2} \left[Y^* \left(\sum_{j=1}^p d_j \frac{\partial A}{\partial x_j}(x_0) \right) Z + Z^* \left(\sum_{j=1}^p d_j \left[\frac{\partial A}{\partial x_j}(x_0) \right]^* \right) Y \right],
$$

when the eigenvalues are ranked in a decreasing order.

3 Function of Ikramov-Nazari

With the notations of [\[3\]](#page-10-4), let $(\xi_1, \xi_2, \xi_3, \xi_4) \in \mathbb{R}^4$, $A \in \mathbb{C}^{n \times n}$. Define

$$
Q(\xi_1, \xi_2, \xi_3, \xi_4) := \begin{pmatrix} A & \xi_1 I & (\xi_3 + i \xi_4)I \\ 0 & A & \xi_2 I \\ 0 & 0 & A \end{pmatrix}, \quad n \geq 3.
$$

Set

$$
f(\xi) := s_{3n-2}(Q(\xi)).
$$

Suppose that the function f attains a local maximum at a given $\xi^0 \in \mathbb{R}^4$, say $s_0 := s_{3n-2}(Q(\xi^0))$. Let us also assume that $s_0 > 0$ and it is a multiple singular of $Q(\xi^0)$. With the above notations, there are i_{3n-2} singular values before the place $3n - 2 + 1$ and j_{3n-2} singular values after the place $3n - 2$ equal to

 $s_{3n-2}(Q(\xi^0))$. To shorten notation, we let p and q stand for i_{3n-2} and j_{3n-2} , respectively. Thus, the multiplicity of s_0 is $m = p + q$. Hence,

$$
s_1(Q(\xi^0)) \ge \cdots \ge s_{3n-2-p}(Q(\xi^0))
$$

>
$$
s_{3n-2-p+1}(Q(\xi^0)) = \cdots = s_{3n-2}(Q(\xi^0))
$$

=
$$
s_{3n-2+1}(Q(\xi^0)) = \cdots = s_{3n-2+q}(Q(\xi^0))
$$

>
$$
s_{3n-2+q+1}(Q(\xi^0)) \ge \cdots \ge s_{3n}(Q(\xi^0)).
$$

Here $p \geqslant 1$ and $q \geqslant 0$. The function

$$
t(\xi) := s_{3n-2-p+1} (Q(\xi)) + \cdots + s_{3n-2+q} (Q(\xi))
$$

is differentiable in a neighborhood of ξ^0 . Also for each $k \in \{1, ..., 3n\}$ and each unitary vector $d \in \mathbb{R}^4$, the function

$$
g_k(\xi) := s_k(Q(\xi))
$$

admits the directional derivative

 $g'_k(\xi^0, d)$.

Observe that the used notation implies

$$
f(\xi) = g_{3n-2}(\xi), \quad \xi \in \mathbb{R}^4.
$$

Next, we determine the relationship between the directional derivatives $f'(\xi^0, d)$ and $f'(\xi^0, -d)$. Given that f has a local maximum at ξ^0 , it follows that for all $e \in \mathbb{R}^4$,

$$
f'(\xi^0, e) := \lim_{h \to 0^+} \frac{f(\xi^0 + he) - f(\xi^0)}{h} \le 0.
$$

Thus, $f'(\xi^0, d) \leq 0$ and $f'(\xi^0, -d) \leq 0$. What conditions must be satisfied in order for $f'(\xi^0, d) = 0$ to hold for all unit vector $d \in \mathbb{R}^4$? By Theorem [6,](#page-5-1) $f'(\xi^0, d)$ is equal to $\mu_p(d)$, pth eigenvalue of the $m \times m$ matrix

$$
F'(d) = [U_2^*, V_2^*] \left(\sum_{j=1}^4 d_j \begin{bmatrix} 0 & \frac{\partial Q}{\partial \xi_j} (\xi^0) \\ \frac{\partial Q^*}{\partial \xi_j} (\xi^0) & 0 \end{bmatrix} \right) \begin{bmatrix} U_2 \\ V_2 \end{bmatrix}
$$

with

$$
U_2 = [u_{3n-2-p+1}, \dots, u_{3n-2+q}]
$$

$$
V_2 = [v_{3n-2-p+1}, \dots, v_{3n-2+q}]
$$

where u_i and v_j are the left and right singular vectors

$$
\begin{aligned}\nQ(\xi^0)v_j &= s_0 u_j \\
Q(\xi^0)^* u_j &= s_0 v_j\n\end{aligned}\n\bigg\}\n\qquad\nj = 3n - 2 - p + 1, \dots, 3n - 2 + q,
$$

and the eigenvalues of $F'(d)$ are arranged in this way

$$
\mu_1(d) \geqslant \cdots \geqslant \mu_p(d) \geqslant \cdots \geqslant \mu_m(d) \tag{5}
$$

Therefore,

$$
f'(\xi^0, d) = \mu_p(d).
$$

Again by Theorem [6,](#page-5-1) we deduce that $f'(\xi^0, -d)$ is equal to the pth eigenvalue of the Hermitian matrix $F'(-d)$. But, it is worth noting that $f'(\xi^0, -d)$ is not necessarily equal to $\mu_p(-d)$. In fact, if

$$
\alpha_1 \geqslant \cdots \geqslant \alpha_m
$$

are the eigenvalues of $F'(-d)$, then

$$
f'(\xi^0, -d) = \alpha_p.
$$

As $F'(-d) = -F'(d)$, it follows

$$
-\mu_m(d) \geqslant \cdots \geqslant -\mu_p(d) \geqslant \cdots \geqslant -\mu_1(d) \tag{6}
$$

are the eigenvalues of $F'(-d)$; whence,

$$
f'(\xi^0, -d) = \alpha_p = -\mu_{m-(p-1)}(d). \tag{7}
$$

Now it is necessary to analyze the relative positions of the indices p and $m (p-1)$.

If $p \leq m - (p - 1)$, then $\mu_p(d) \leq 0$, and it follows that

$$
0 \geqslant \mu_p(d) \geqslant \cdots \geqslant \mu_{m-(p-1)}(d) \geqslant \cdots \geqslant \mu_m(d).
$$

Hence, $0 \geq \mu_{m-(p-1)}(d)$ and therefore $\alpha_p = -\mu_{m-(p-1)}(d) \geq 0$, but $\alpha_p =$ $f'(\xi^0, -d) \leq 0$. Thus, $\alpha_p = 0$; i.e. $f'(\xi^0, -d) = 0$. Given that f has a local maximum at ξ^0 , for all unit vector $e \in \mathbb{R}^4$, we have

$$
f'(\xi^0, e) = 0.
$$

Doutbful case: If $p > m - (p - 1)$, then $\mu_{m-(p-1)}(d) \geq \mu_p(d)$. But, although $\mu_p(d) \leq 0$, it is not guaranteed that the inequality $\mu_{m-(p-1)}(d) \leq 0$ holds.

4 Average of singular values

We know that the average of singular values of $Q(\xi)$ that coalesce with the mmultiple singular value $s_{3n-2}(Q(\xi^0))$ at $\xi = \xi^0$, is a differentiable function in a neighborhood of ξ^0 . Thus we consider the differentiable function

$$
H(\xi) := t(\xi) - ms_0;
$$

obviously, $H(\xi^0) = 0$. Hence, the point ξ^0 belongs to the level hypersurface of level 0 of the function $H(\xi)$. Let

$$
\nabla H(\xi^0) = \left(\frac{\partial H}{\partial \xi_1}(\xi^0), \frac{\partial H}{\partial \xi_2}(\xi^0), \frac{\partial H}{\partial \xi_3}(\xi^0), \frac{\partial H}{\partial \xi_4}(\xi^0) \right)
$$

be the gradient of $H(\xi)$ at ξ^0 . Let $d \in \mathbb{R}^4$ such that

$$
\nabla H(\xi^0) \cdot d = 0,
$$

where \cdot denotes the ordinary scalar product in \mathbb{R}^4 . Then, by the chain rule,

$$
H'(\xi^0, d) = \nabla H(\xi^0) \cdot d = 0.
$$

This implies

$$
0 = g'_{3n-2-p+1}(\xi^0, d) + \cdots + g'_{3n-2}(\xi^0, d) + \cdots + g'_{3n-2+q}(\xi^0, d);
$$

if we consider the $m \times m$ Hermitian matrix $F'(d)$, it means that the sum of its eigenvalues is zero:

$$
0 = \mu_1(d) + \dots + \mu_p(d) + \dots + \mu_m(d).
$$

When $p = 1$, this is equivalent to say that $s_{3n-2}(Q(\xi^0))$ is the first value of the chain of singular values equal to s_0 , then all the functions

$$
g_{3n-2}(\xi), g_{3n-2+1}(\xi), \ldots, g_{3n-2+q}(\xi)
$$

take the same value at ξ^0 , and it is equal to s_0 . Moreover, all these functions have at ξ^0 a local maximum, because of

$$
f(\xi) := g_{3n-2}(\xi) \geq g_{3n-2+1}(\xi) \geq \cdots \geq g_{3n-2+q}(\xi).
$$

This implies that for all unitary $d \in \mathbb{R}^4$,

$$
\forall k = 3n - 2, ..., 3n - 2 + q, \quad g'_{k}(\xi^{0}, d) \leq 0;
$$

therefore, $\mu_1(d) \leq 0, \ldots, \mu_m(d) \leq 0$, and, given that $t(\xi)$ has a local maximum at ξ^0 and is differentiable at ξ^0 , we have

$$
\nabla t(\xi^0) = \mathbf{0};
$$

whence $\nabla H(\xi^0) = \mathbf{0}$ and for all $k = 3n - 2, ..., 3n - 2 + q$, $g'_k(\xi^0, d) = 0$; in particular, $f'(\xi_0, d) = 0$. This is proved because $0 = \mu_1(d) + \cdots + \mu_m(d)$; since $\forall k, \mu_k(d) \leq 0$, we obtain $\forall k, \mu_k(d) = 0$; consequently, $\forall k, g'_k(\xi^0, d) = 0$.

From now on let p be any integer from the range we are considering. Furthermore, suppose that for all $k = 3n-2-p+1, \ldots, 3n-2+q$, all the functions $g_k(\xi)$ have a local maximum at ξ^0 . Then for all unitary $d \in \mathbb{R}^4, g'_k(\xi^0, d) \leq 0$. As $t(\xi)$ has a local maximum at ξ^0 , $t'(\xi^0, d) = 0$; but

$$
t'(\xi^0, d) = g'_{3n-2-p+1}(\xi^0, d) + \cdots + g'_{3n-2+q}(\xi^0, d);
$$

consequently, $f'(\xi^0, d) = 0$.

When some of the functions $g_k(\xi)$ have a local maximum at ξ^0 and any others have a local minimum at ξ^0 , the analysis becomes more complicated and I do not obtain any conclusion.

5 Remark

In January 31, 2005, I wrote an e-mail to J.B. Hiriart-Urruty asking him whether his results in [\[2\]](#page-10-0) for real symmetric matrices could be generalized to complex Hermitian matrices. He forwarded my message to M. Torki [\[10\]](#page-10-5), who answered affirmatively. Moreover, Torki told me that his results in [\[9\]](#page-10-6) for second order directional derivatives and real symmetric matrices were also true for the Hermitian case.

A particular case of Theorem [9](#page-6-1) about the right derivative of the function $t \mapsto \sigma_k(A+tB)$ at $t = 0$, where t is real and A, B are $n \times n$ complex matrices, was obtained by Lippert [\[5,](#page-10-7) Lemma A.5] by a different method. See also Corollary 11 in [\[1\]](#page-10-8).

Acknowledgments

I thank Gorka Armentia and Francisco E. Velasco for the talks we had about this topic.

References

- [1] G. Armentia, J. M. Gracia, F. E. Velasco. Perforated strict pseudospectra of Demmel's matrices. Linear Algebra Appl. 548:77–94, 2018. [11](#page-10-9)
- [2] J. Hiriart-Urruty and D. Ye. Sensitivity analysis of all eigenvalues of a symmetric matrix. Numer. Math., 70:45–72, 1995. [1,](#page-0-1) [2,](#page-1-1) [4,](#page-3-2) [10](#page-9-1)
- [3] K. D. Ikramov and A. M. Nazari. On the distance to the closest matrix with triple zero eigenvalue. Mathematical Notes, 73(4):511–520, 2003. [7](#page-6-2)
- [4] T. Kato. A Short Introduction to Perturbation Theory for Linear Operators. Springer, 1982. [2](#page-1-1)
- [5] R. A. Lippert. Fixing two eigwnvalues by a minimal perturbation. Linear Algebra Appl., 406:177–200, 2005. [11](#page-10-9)
- [6] J.-G. Sun. A note on simple non-zero singular values. Journal of Computational Mathematics, 6(3):258–266, 1988. [1](#page-0-1)
- [7] J.-G. Sun. Sensitivity analysis of zero singular values and multiple singular values. J. Comput. Math., 6(4):325–335, Oct. 1988.
- [8] J.-G. Sun. Multiple Eigenvalue Sensitivity Analysis. Linear Algebra Appl., 137/138:183–211, 1990. [1,](#page-0-1) [4](#page-3-2)
- [9] M. Torki. Second-order directional derivatives of all eigenvalues of a symmetric matrix. Nonlinear Analysis, 46:1133–1150, 2001. [10](#page-9-1)
- [[10](#page-9-1)] M. Torki. Private communication, 2005. 10