

# The Number of 1...d-Avoiding Permutations of Length $d+r$ for SYMBOLIC $d$ but Numeric $r$

By Shalosh B. EKHAD, Nathaniel SHAR, and Doron ZEILBERGER

*Dedicated to Ira Martin GESSEL (b. April 9, 1951), on his millionth<sub>2</sub> birthday*

**Preface: How many permutations are there of length googol+30 avoiding an increasing subsequence of length googol?**

This number is way too big for our physical universe, but the number of permutations of length googol+30 that *contain* at least one increasing subsequence of length googol is a certain integer that may be viewed in <http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimPDF/gessel64.pdf>. Hence the number of permutations of length googol+30 *avoiding* an increasing subsequence of length googol is  $(\text{googol} + 30)!$  *minus* the above small number.

## Counting the “Bad Guys”

Recall that thanks to Robinson-Schensted ([Rob][Sc]), the number of permutations of length  $n$  that **do not** contain an increasing subsequence of length  $d$  is given by

$$G_d(n) := \sum_{\substack{\lambda \vdash n \\ \#rows(\lambda) < d}} f_\lambda^2 \quad ,$$

where  $\lambda$  denotes a typical *Young diagram*, and  $f_\lambda$  is the number of *Standard Young tableaux* whose *shape* is  $\lambda$ .

Hence the number of permutations of length  $n$  that **do** contain an increasing subsequence of length  $d$  is

$$B_d(n) := \sum_{\substack{\lambda \vdash n \\ \#rows(\lambda) \geq d}} f_\lambda^2 \quad .$$

Since the total number of permutations of length  $n$  is  $n!$  ([B]), if we know how to find  $B_d(n)$ , we would know immediately  $G_d(n) = n! - B_d(n)$ , at least if we leave  $n!$  alone as a factorial, rather than spell it out.

Recall that the *Hook Length formula* (see [Wiki]) tells you that if  $\lambda$  is a Young diagram then

$$f_\lambda = \frac{n!}{\prod_{c \in \lambda} h(c)} \quad ,$$

where the product is over all the  $n$  cells of the Young diagram, and the *hook-length*,  $h(c)$ , of a cell  $c = (i, j)$ , is  $(\lambda_i - i) + (\lambda'_j - j) + 1$ , where  $\lambda'$  is the *conjugate* diagram, where the rows become columns and vice-versa.

Let  $r$  be a fixed integer, then for *symbolic*  $d$ , valid for  $d \geq r - 1$ , any Young diagram with at least  $d$  rows, and with  $d + r$  cells, can be written, for some Young diagram  $\mu = (\mu_1, \dots, \mu_r)$ , with  $\leq r$  cells, (where we add zeros to the end if the number of parts of  $\mu$  is less than  $r$ ) as

$$\lambda = (1 + \mu_1, \dots, 1 + \mu_r, 1^{d-r+r'}) \quad ,$$

where  $r' = r - |\mu|$ . For such a shape  $\lambda$ , with *at least*  $d$  rows,

$$\prod_{c \in \lambda} h(c) = \left( \prod_{c \in \mu} h(c) \right) \cdot ((d + r' + \mu_1)(d + r' - 1 + \mu_2) \cdots (d + r' - r + 1 + \mu_r)) \cdot (d - r + r')! \quad .$$

Hence  $f_\lambda$ , that is  $(d+r)!$  divided by the above, is a certain specific number times a certain polynomial in  $d$ . Since, for a specific, *numeric*,  $r$ , there are only *finitely* many Young diagrams with at most  $r$  cells, the computer can find all of them, compute the polynomial corresponding to each of them, square it, and add-up all these terms, getting an *explicit polynomial* expression, in the variable  $d$ , for  $B_d(d+r)$ , the number of permutations of length  $d+r$  that *contain* an increasing subsequence of length  $d$ . As we said above, from this we can find  $G_d(d+r) = (d+r)! - B_d(d+r)$ , valid for *symbolic*  $d \geq r - 1$ .

### **$B_d(d+r)$ for $r$ from 0 to 30**

$$B_d(d) = 1 \quad ,$$

$$B_d(d+1) = d^2 + 1 \quad ,$$

$$B_d(d+2) = \frac{1}{2} d^4 + d^3 + \frac{1}{2} d^2 + d + 3 \quad ,$$

$$B_d(d+3) = \frac{1}{6} d^6 + d^5 + \frac{5}{3} d^4 + \frac{2}{3} d^3 + \frac{19}{6} d^2 + \frac{31}{3} d + 11 \quad ,$$

$$B_d(d+4) = \frac{1}{24} d^8 + \frac{1}{2} d^7 + \frac{25}{12} d^6 + \frac{19}{6} d^5 + \frac{29}{24} d^4 + 9 d^3 + \frac{247}{6} d^2 + \frac{395}{6} d + 47 \quad ,$$

$$B_d(d+5) = \frac{1}{120} d^{10} + \frac{1}{6} d^9 + \frac{31}{24} d^8 + \frac{14}{3} d^7 + \frac{823}{120} d^6 + \frac{67}{30} d^5 + \frac{653}{24} d^4 + \frac{959}{6} d^3 + \frac{10459}{30} d^2 + \frac{3981}{10} d + 239 \quad .$$

For  $B_d(d+r)$  for  $r$  from 6 up to 30, see

<http://www.math.rutgers.edu/~zeilberg/tokhniot/oGessel64a> .

### **Sequences**

The sequence  $G_3(n)$  is the greatest *celeb* in the kingdom of combinatorial sequences [the subject of an entire book ([St]) by Ira Gessel's illustrious *academic father*, Richard Stanley], the super-famous **A000108** in Neil Sloane's legendary database ([Sl]).  $G_4(n)$ , while not in the same league as the Catalan sequence, is still moderately famous, **A005802**.  $G_5(n)$  is **A047889**,  $G_6(n)$  is **A047890**,  $G_7(n)$  is **A052399**,  $G_8(n)$  is **A072131**,  $G_9(n)$  is **A072132**,  $G_{10}(n)$  is **A072133**,  $G_{11}(n)$  is **A072167**, but  $G_d(n)$  for  $d \geq 12$  are absent (for a good reason, one must stop somewhere!). Also the *flattened version* of the *double-sequence*,  $\{G_d(n)\}$ , for  $1 \leq d \leq n \leq 45$  is **A047887**. Using the

polynomials  $B_d(d+r)$ , we computed the first  $2d+1$  terms of  $G_d(n)$  for  $d \leq 30$ . See <http://www.math.rutgers.edu/~zeilberg/tokhniot/oGessel64b>.

But this method can only go up to  $2d+1$  terms of the sequence  $G_d(n)$ , and of course, the first  $d-1$  terms are trivial, namely  $n!$ . Can we find the first 100 terms (or whatever) for the sequences  $G_d(n)$  for  $d$  up to 20, and beyond, **efficiently**?

### Encore: Efficient Computer-Algebra Implementation of Ira Gessel's AMAZING Determinant Formula

Recall Ira Gessel's [G] famous expression for the generating function of  $G_d(n)/n!^2$ , *canonized* in the *bible* ([W], p. 996, Eq. (5)). Here it is:

$$\sum_{n \geq 0} \frac{G_d(n)}{n!^2} x^{2n} = \det(I_{|i-j|}(2x))_{i,j=1,\dots,d} \quad ,$$

in which  $I_\nu(t)$  is (the modified Bessel function)

$$I_\nu(t) = \sum_{j=0}^{\infty} \frac{(\frac{1}{2}t)^{2j+\nu}}{j!(j+\nu)!} \quad .$$

Can we use this to compute the first 100 terms of, say,  $G_{20}(n)$ ?

While computing *numerical* determinants is very fast, computing *symbolic* ones is a different story. First, do not get scared by the “infinite” power series. If we are only interested in the first  $N$  terms of  $G_d(n)$ , then it is safe to truncate the series up to  $t^{2N}$ , and take the determinant of a  $d \times d$  matrix with *polynomial entries*. If you use the *vanilla* determinant in a computer-algebra system such as Maple, it would be very inefficient, since the degree of the determinant is much larger than  $2N$ . But a little cleverness can make things more efficient. The Maple package **Gessel64**, available free of charge from

<http://www.math.rutgers.edu/~zeilberg/tokhniot/Gessel64> ,

accompanying this article, has a procedure `SeqIra(k,N)` that computes the first  $N$  terms of  $G_k(n)$ , using a division-free algorithm (see [Rot]) over an appropriate ring to compute the determinant in Gessel's famous formula.

```
SeqIra:=proc(k,N) local ira,t,i,j, R:
```

```
  R := table():
```

```
  R['0'] := 0:
```

```
  R['1'] := 1:
```

```
  R['+'] := '+':
```

```

R['-'] := '-':
R['*'] := proc(p, q): return add(coeff(p*q, t, i)*t**i, i=0..2*N): end:
R['='] := proc(p, q): return evalb(p = q): end:
ira:=expand(LinearAlgebra[Generic][Determinant][R](Matrix([seq([seq(Iv(abs(i-j),t,2*N),
                                                                    j=1..k-1)],
                                                                    i=1..k-1)]))):
[seq(coeff(ira,t,2*i)*i!**2,i=1..N)]:
end:

```

In the above code, procedure  $Iv(v,t,N)$  computes the truncated modified Bessel function that shows up in Gessel's determinant, and it is short enough to reproduce here:

```
Iv:=proc(v,t,N) local j: add(t**(2*j+v)/j!/(j+v)!,j=0..trunc((N-v)/2)+1): end: .
```

Using this procedure, the first-named author computed (in 4507 seconds) the first 100 terms of each of the sequences  $G_d(n)$  for  $3 \leq d \leq 20$ , and could have gone much further.

See <http://www.math.rutgers.edu/~zeilberg/tokhniot/oGessel64c> .

## HAPPY 64th BIRTHDAY, IRA!

### References

- [B] Rabbi Levi Ben Gerson, *Sefer Maaseh Hoshev*, Avignon, 1321.
- [G] I. Gessel, *Symmetric functions and P-recursiveness*, Journal of Combinatorial Theory Series A **53** (1990), 257-285; <http://people.brandeis.edu/~gessel/homepage/papers/dfin.pdf> .
- [Rob] G. de B. Robinson, *On the representations of  $S_n$* , Amer. J. Math. **60** (1938), 745-760.
- [Rot] G. Rote, *Division-Free Algorithms for the Determinant and Pfaffian: Algebraic and Combinatorial Approaches*, *Computational Discrete Mathematics: Advanced Lectures* (2001), 119-135.
- [Sc] C. E. Schensted, *Largest increasing and decreasing subsequences*, Canad. J. Math **13** (1961), 179-191.
- [SI] N.J.A. Sloane, *The On-Line Encyclopedia of Integer Sequences*; <http://oeis.org> .
- [St] R. P. Stanley, *"Catalan Numbers"*, Cambridge University Press, 2015.

[Wiki] The Wikipedia Foundation, *Hook Length Formula*;  
[http://en.wikipedia.org/wiki/Hook\\_length\\_formula](http://en.wikipedia.org/wiki/Hook_length_formula) .

[W] H. S. Wilf, *Mathematics, an experimental science*, in: “Princeton Companion to Mathematics”,  
(W. Timothy Gowers, ed.), Princeton University Press, 2008, 991-1000;  
<http://www.math.rutgers.edu/~zeilberg/akherim/HerbMasterpieceEM.pdf> .

---

Shalosh B. Ekhad, c/o D. Zeilberger, Department of Mathematics, Rutgers University (New Brunswick),  
Hill Center-Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.

---

Nathaniel Shar, Department of Mathematics, Rutgers University (New Brunswick), Hill Center-  
Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.  
nshar at math dot rutgers dot edu ; <http://www.math.rutgers.edu/~nbs48/> .

---

Doron Zeilberger, Department of Mathematics, Rutgers University (New Brunswick), Hill Center-  
Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.  
zeilberg at math dot rutgers dot edu ; <http://www.math.rutgers.edu/~zeilberg/> .

---

Published in The Personal Journal of Shalosh B. Ekhad and Doron Zeilberger  
( <http://www.math.rutgers.edu/~zeilberg/pj.html> ) and [arxiv.org](http://arxiv.org).

---

**April 9, 2015**