# MINIMUM RIESZ ENERGY PROBLEMS FOR A CONDENSER WITH "TOUCHING PLATES"

P. D. DRAGNEV<sup>†</sup>, D. P. HARDIN<sup>\*</sup>, E. B. SAFF<sup>\*</sup>, AND N. ZORII<sup>\*\*</sup>

ABSTRACT. We consider minimum energy problems in the presence of an external field for a condenser with "touching plates"  $A_1$  and  $A_2$  in  $\mathbb{R}^n$ ,  $n \ge 3$ , relative to the  $\alpha$ -Riesz kernel  $|x - y|^{\alpha - n}$ ,  $0 < \alpha \le 2$ . An intimate relationship between such problems and minimal  $\alpha$ -Green energy problems for positive measures on  $A_1$  is shown. We obtain sufficient and/or necessary conditions for the solvability of these problems in both the unconstrained and the constrained settings, investigate the properties of minimizers, and prove their uniqueness. Furthermore, characterization theorems in terms of variational inequalities for the weighted potentials are established. The approach applied is mainly based on the establishment of a perfectness-type property for the  $\alpha$ -Green kernel with  $0 < \alpha < 2$  which enables us, in particular, to analyze the existence of the  $\alpha$ -Green equilibrium measure of a set. The results obtained are illustrated by several examples.

## 1. INTRODUCTION

The purpose of the paper is to study minimum energy problems in the presence of an external field for a condenser **A** with touching oppositely-charged plates  $A_1$  and  $A_2$  in  $\mathbb{R}^n$ ,  $n \ge 3$ , relative to the  $\alpha$ -Riesz kernel  $|x - y|^{\alpha - n}$ ,  $0 < \alpha \le 2$ . The difficulties appearing in the course of our investigation are caused by the fact that a short-circuit between  $A_1$  and  $A_2$  might occur, for the Euclidean distance between these conductors is zero.

Therefore, it is meaningful to ask what kind of conditions on the objects in question could prevent such a phenomenon so that a minimizer for the corresponding  $\alpha$ -Riesz energy problem might exist. One of the ideas, to be discussed for this purpose, is to impose upper constraints on the charges of the touching conductors.

We establish sufficient and/or necessary conditions for the existence of minimizing measures for both the unconstrained and the constrained problems, and prove their uniqueness. The conditions obtained are expressed in geometric-potential terms for  $A_1$  and  $A_2$ , or in measure

*Date*: October 24, 2018.

<sup>&</sup>lt;sup>†</sup> The research of this author was supported, in part, by a Simons Foundation grant no. 282207.

 $<sup>^{\</sup>ast}$  The research of these authors was supported, in part, by the U. S. National Science Foundation under grants DMS-1109266 and DMS-1412428.

 $<sup>^{\</sup>ast}$  The research of this author was supported, in part, by the Scholar-in-Residence program at IPFW .

The authors express their gratitude to Erwin Schrödinger International Institute for providing conducive research atmosphere during their stay when part of this manuscript was written.

 $\mathbf{2}$ 

theory terms for the constraints under consideration, or in terms of variational inequalities for the weighted potentials. We also provide a detailed analysis of the supports of the minimizers.

The approach developed in the paper is based on a newly discovered important relationship between, on the one hand, minimum  $\alpha$ -Riesz energy problems over signed measures associated with a condenser **A** and, on the other hand, minimum energy problems for nonnegative measures on  $A_1$  relative to the  $\alpha$ -Green function  $g_D^{\alpha}$  of the domain  $D := \mathbb{R}^n \setminus A_2$ .

Regarding the latter problems, crucial to the arguments applied in their investigation is the pre-Hilbert structure on the linear space  $\mathcal{E}_{g_D^{\alpha}}(D)$  of all (signed) Borel measures on D with finite  $g_D^{\alpha}$ -Green energy and, most importantly, a completeness theorem for certain metric subspaces of  $\mathcal{E}_{g_D^{\alpha}}(D)$  with nonnegative elements. This completeness theorem enables us, in particular, to analyze the existence of the  $\alpha$ -Green equilibrium measure of a set.

To formulate precisely the problems in question, we first need to introduce several notions, to discuss relations between them, and to recall some well-known results; this is the purpose of the next section. The scheme of the rest of the paper is described at the end of Section 3, after the formulations of the problems (see Problems 3.1 and 3.2).

# 2. Basic notions; relations between them. Known results

2.1. Measures, energies, potentials. Let X be a locally compact Hausdorff space, to be specified below, and  $\mathfrak{M} = \mathfrak{M}(X)$  the linear space of all Radon measures  $\mu$  on X, equipped with the *vague* (=*weak*<sup>\*</sup>) topology, i.e. the topology of pointwise convergence on the class of all continuous functions on X with compact support. We denote by  $\mu^+$  and  $\mu^-$  the positive and the negative parts in the Hahn–Jordan decomposition of a measure  $\mu \in \mathfrak{M}(X)$ , respectively, and by  $S_X^{\mu}$  its support. Given  $\mu$  and a  $\mu$ -measurable function  $\psi$ , for the sake of brevity we shall write  $\langle \psi, \mu \rangle := \int \psi \, d\mu$ .<sup>1</sup>

A kernel  $\kappa(x, y)$  on X is a symmetric, lower semicontinuous function  $\kappa : X \times X \to [0, \infty]$ . Given  $\mu, \mu_1 \in \mathfrak{M}$ , let  $E_{\kappa}(\mu, \mu_1)$  and  $U^{\mu}_{\kappa}(\cdot)$  denote the *mutual energy* and the *potential* relative to the kernel  $\kappa$ , respectively, i.e.

$$E_{\kappa}(\mu,\mu_{1}) := \int \kappa(x,y) \, d(\mu \otimes \mu_{1})(x,y),$$
$$U_{\kappa}^{\mu}(x) := \int \kappa(x,y) \, d\mu(y).$$

For  $\mu = \mu_1$ , the mutual energy  $E_{\kappa}(\mu, \mu_1)$  defines the energy  $E_{\kappa}(\mu, \mu) =: E_{\kappa}(\mu)$ .

<sup>&</sup>lt;sup>1</sup>When introducing notation, we assume the corresponding object on the right to be well-defined.

2.2. Strictly positive definite kernels. Capacities. Throughout this section, a kernel  $\kappa$  is assumed to be *strictly positive definite*, which means that  $E_{\kappa}(\mu), \mu \in \mathfrak{M}(X)$ , is nonnegative whenever defined and  $E_{\kappa}(\mu) = 0$  implies  $\mu = 0$ . Then the collection  $\mathcal{E}_{\kappa} = \mathcal{E}_{\kappa}(X)$ of all  $\mu \in \mathfrak{M}$  with  $E_{\kappa}(\mu) < \infty$  forms a pre-Hilbert space with the scalar product  $E_{\kappa}(\mu, \mu_1)$ and the norm  $\|\mu\|_{\kappa} := \sqrt{E_{\kappa}(\mu)}$  (see [10]). The topology on  $\mathcal{E}_{\kappa}$  defined by  $\|\cdot\|_{\kappa}$  is called *strong*. The following lemma from the geometry of the pre-Hilbert space  $\mathcal{E}_{\kappa}$  is often useful (see [10, Lemma 4.1.1]).

**Lemma 2.1.** Let  $\Gamma$  be a convex subset of  $\mathcal{E}_{\kappa}$ . If there exists  $\mu_0 \in \Gamma$  with minimal norm

$$\|\mu_0\|_{\kappa} = \inf_{\mu \in \Gamma} \|\mu\|_{\kappa},$$

then such a minimal element is unique. Moreover,

$$\|\mu - \mu_0\|_{\kappa}^2 \leq \|\mu\|_{\kappa}^2 - \|\mu_0\|_{\kappa}^2 \text{ for all } \mu \in \Gamma.$$

Given a set  $B \subset X$ , let  $\mathfrak{M}^+(B)$  be the convex cone of all nonnegative measures concentrated in B, and let  $\mathfrak{M}^+(B,b)$ , b > 0, consist of all  $\mu \in \mathfrak{M}^+(B)$  with  $\mu(B) = b$ . We also write  $\mathfrak{M}^+ := \mathfrak{M}^+(X), \mathcal{E}^+_{\kappa}(B) := \mathcal{E}_{\kappa} \cap \mathfrak{M}^+(B), \mathcal{E}^+_{\kappa} := \mathcal{E}^+_{\kappa}(X)$ , and  $\mathcal{E}^+_{\kappa}(B,b) := \mathcal{E}_{\kappa} \cap \mathfrak{M}^+(B,b)$ .

Let  $C_{\kappa}(B)$  denote the *interior capacity* of B relative to a kernel  $\kappa$ , defined by<sup>2</sup>

(2.1) 
$$1/C_{\kappa}(B) := w_{\kappa}(B) := \inf_{\mu \in \mathcal{E}_{\kappa}^{+}(B,1)} E_{\kappa}(\mu)$$

Note that, in consequence of Lemma 2.1, a measure  $\lambda_B = \lambda_B^{\kappa} \in \mathcal{E}_{\kappa}^+(B,1)$  with minimal energy  $\|\lambda_B\|_{\kappa}^2 = w_{\kappa}(B)$  is unique (provided it exists).

Following Fuglede [10], we call a kernel  $\kappa$  perfect if any strong Cauchy sequence in  $\mathcal{E}_{\kappa}^{+}$  converges strongly and, in addition, the strong topology on  $\mathcal{E}_{\kappa}^{+}$  is finer than the induced vague topology on  $\mathcal{E}_{\kappa}^{+}$ . Note that then the metric space  $\mathcal{E}_{\kappa}^{+}$  is complete in the induced strong topology. What is also important is that the solution  $\lambda_{B}^{\kappa}$  to the minimum energy problem appeared in (2.1) exists, provided that  $\kappa$  is perfect, B is closed, and  $0 < C_{\kappa}(B) < \infty$  (see [10, Theorem 4.1]).

If  $f : X \to [-\infty, \infty]$  is an external field, then the *f*-weighted potential  $W^{\mu}_{\kappa, f}$  and the *f*-weighted energy  $G_{\kappa, f}(\mu)$  of  $\mu \in \mathcal{E}_{\kappa}(X)$  are respectively given by

$$W^{\mu}_{\kappa,f}(x) := U^{\mu}_{\kappa}(x) + f(x),$$
  
$$G_{\kappa,f}(\mu) := E_{\kappa}(\mu) + 2\langle f, \mu \rangle = \langle W^{\mu}_{\kappa,f} + f, \mu \rangle.$$

We also define

$$\mathcal{E}_{\kappa,f}(\mathbf{X}) := \{ \mu \in \mathcal{E}_{\kappa}(\mathbf{X}) : G_{\kappa,f}(\mu) < \infty \}.$$

<sup>&</sup>lt;sup>2</sup>As usual, the infimum over the empty set is taken to be  $+\infty$ . We put  $1/(+\infty) = 0$  and  $1/0 = +\infty$ .

2.3.  $\alpha$ -Riesz and  $\alpha$ -Green kernels. Balayage. Fix  $n \ge 3$ , a domain  $D \subset \mathbb{R}^n$ , and  $\alpha \in (0, 2]$ . In the rest of the paper, unless stated otherwise, one of the following two cases is assumed to hold: either  $X = \mathbb{R}^n$  and  $\kappa$  is the  $\alpha$ -Riesz kernel  $\kappa_{\alpha}(x, y) := |x - y|^{\alpha - n}$  (where |x - y| is the Euclidean distance in  $\mathbb{R}^n$  between x and y), or X = D and  $\kappa$  is the generalized  $\alpha$ -Green function  $g_D^{\alpha}$  of D, defined by<sup>3</sup>

(2.2) 
$$g_D^{\alpha}(x,y) = U_{\kappa_{\alpha}}^{\varepsilon_y}(x) - U_{\kappa_{\alpha}}^{\beta_D^{\alpha}\varepsilon_y}(x) \quad \text{for all } x, y \in D,$$

where  $\varepsilon_y$  denotes the unit Dirac measure at a point y and  $\beta_{D^c}^{\alpha}$  the  $\alpha$ -Riesz balayage onto  $D^c := \mathbb{R}^n \setminus D$  (cf. [14, Chapter IV, Section 5], or see just below).

To avoid triviality, assume each component of  $D^c$  to have nonzero  $\alpha$ -Riesz capacity. Note that, if  $\alpha = 2$  and D is regular in the sense of the solvability of the (classical) Dirichlet problem, then  $g_D^2$  is, in fact, the classical Green function of D.

Let Q be a given closed subset of  $\mathbb{R}^n$ . By definition, the  $\alpha$ -Riesz balayage measure  $\beta_Q^{\alpha}\mu$  of  $\mu \in \mathfrak{M}(\mathbb{R}^n)$  onto Q is supported by Q and satisfies the relation

$$U^{\beta^{\alpha}_{Q}\mu}_{\kappa_{\alpha}}(x) = U^{\mu}_{\kappa_{\alpha}}(x)$$
 n.e. in  $Q$ 

where "n.e." (nearly everywhere) means that the equality holds everywhere in Q except for a subset with  $\alpha$ -Riesz capacity zero. Such a  $\beta_Q^{\alpha}\mu$  exists and it is unique among the *C*-absolutely continuous measures  $\nu \in \mathfrak{M}(\mathbb{R}^n)$ , namely those that  $\nu(K) = 0$  for every compact  $K \subset \mathbb{R}^n$  with  $C_{\kappa_\alpha}(K) = 0$ . Throughout the paper, when speaking of the  $\alpha$ -Riesz balayage measure, we always mean exactly this one. Then, by [14, Chapter IV, Section 5],

(2.3) 
$$\beta_Q^{\alpha} \mu = \int \beta_Q^{\alpha} \varepsilon_y \, d\mu(y).$$

If  $\mu \ge 0$ , then also  $U_{\kappa_{\alpha}}^{\beta_{Q}^{\alpha}\mu}(x) \le U_{\kappa_{\alpha}}^{\mu}(x)$  for all  $x \in \mathbb{R}^{n}$ . If, moreover,  $\mu \in \mathcal{E}_{\kappa_{\alpha}}^{+}(\mathbb{R}^{n})$ , then  $E_{\kappa_{\alpha}}(\beta_{Q}^{\alpha}\mu) \le E_{\kappa_{\alpha}}(\mu)$ 

and also

(2.4) 
$$\left\|\mu - \beta_Q^{\alpha} \mu\right\|_{\kappa_{\alpha}} < \left\|\mu - \nu\right\|_{\kappa_{\alpha}} \quad \text{for all} \quad \nu \in \mathcal{E}^+_{\kappa_{\alpha}}(Q), \ \nu \neq \beta_Q^{\alpha} \mu,$$

so that  $\beta_Q^{\alpha}\mu$  is, in fact, the orthogonal projection of  $\mu$  in the pre-Hilbert space  $\mathcal{E}_{\kappa_{\alpha}}(\mathbb{R}^n)$ onto the convex cone  $\mathcal{E}^+_{\kappa_{\alpha}}(Q)$ .

It is well known (see [14]) that, if  $\mu \in \mathfrak{M}^+(\mathbb{R}^n)$  is bounded (i.e.,  $\mu(\mathbb{R}^n) < \infty$ ), then  $\beta_O^{\alpha}\mu(\mathbb{R}^n) \leq \mu(\mathbb{R}^n)$ .

This general fact is specified by the following assertion (see [17, Theorem 4]; for  $\alpha = 2$ , see also [16, Theorem B]).

<sup>&</sup>lt;sup>3</sup>The function  $g^{\alpha}_{D}(x, y), x, y \in D$ , is nonnegative and symmetric (see [12] and [14, Chapter IV, Section 5]). It is also lower semicontinuous, which follows from the lower semicontinuity of the  $\alpha$ -Riesz kernel and the continuity of  $U^{\beta^{\alpha}_{D}c^{\varepsilon}y}_{\kappa_{\alpha}}(x), x, y \in D$ , the latter in turn being a consequence of  $U^{\beta^{\alpha}_{D}c^{\varepsilon}y}_{\kappa_{\alpha}}(x) = U^{\beta^{\alpha}_{D}c^{\varepsilon}x}_{\kappa_{\alpha}}(y), x, y \in D$ . Hence,  $g^{\alpha}_{D}$  can be treated as a kernel on the locally compact space D.

**Theorem 2.2.** Q is not  $\alpha$ -thin at the Alexandroff point  $\omega$  of  $\mathbb{R}^n$  if and only if, for every bounded measure  $\mu \in \mathfrak{M}^+(\mathbb{R}^n)$ ,

(2.5) 
$$\beta_O^{\alpha}\mu(\mathbb{R}^n) = \mu(\mathbb{R}^n).$$

By definition, Q is not  $\alpha$ -thin at  $\omega$  if  $Q^*$ , the inverse of Q relative to the unit sphere  $S(0,1) := \{x \in \mathbb{R}^n : |x| = 1\}$ , is not  $\alpha$ -thin at x = 0, or equivalently (see [14, Theorem 5.10]), if x = 0 is an  $\alpha$ -regular point for  $Q^*$ .<sup>4</sup>

If  $\mu \in \mathfrak{M}(D)$  and  $U^{\mu}_{g^{\alpha}_{D}}(x)$  is well defined at  $x \in D$ , then, due to (2.2) and (2.3),

(2.6) 
$$U^{\mu}_{g^{\alpha}_{D}}(x) = U^{\mu}_{\kappa_{\alpha}}(x) - U^{\beta^{\alpha}_{Dc}\mu}_{\kappa_{\alpha}}(x).$$

Likewise,

(2.7) 
$$E_{g_D^{\alpha}}(\mu) = E_{\kappa_{\alpha}}\left(\mu - \beta_{D^c}^{\alpha}\mu\right) = E_{\kappa_{\alpha}}(\mu) - E_{\kappa_{\alpha}}\left(\beta_{D^c}^{\alpha}\mu\right).$$

whenever  $E_{g_D^{\alpha}}(\mu)$  is well defined. See [3, Chapter XI, Section 10] and [14, Chapter IV, Section 1, n° 2], where (2.7) has been shown for  $\alpha = 2$ . For  $\alpha < 2$ , the proof is based on (2.2) and (2.3) and runs in a way similar to that in [14].

It is well known that the  $\alpha$ -Riesz kernel is strictly positive definite and, moreover, perfect (see, e.g., [5, 6, 7, 10, 14]); hence, the metric space  $\mathcal{E}^+_{\kappa_{\alpha}}(\mathbb{R}^n)$  is complete in the induced strong topology. However, by Cartan [5], the whole pre-Hilbert space  $\mathcal{E}_{\kappa_{\alpha}}(\mathbb{R}^n)$  is, in general, strongly incomplete, and this is the case even for the Coulomb kernel  $\kappa_2(x, y) = |x - y|^{-1}$  on  $\mathbb{R}^3$ ; compare with Theorem 2.4 below.

Thus, in consequence of (2.7), the  $\alpha$ -Green kernel is strictly positive definite as well. In the case  $\alpha = 2$ , the 2-Green kernel  $g_D^2$  is actually known to be perfect (see [9, 10, 14]), so that, by [10, Theorem 4.1], the measure  $\lambda_B \in \mathcal{E}_{g_D^2}^+(B, 1)$  with minimal  $g_D^2$ -Green energy

$$\|\lambda_B\|_{g_D^2}^2 = w_{g_D^2}(B) = \left[C_{g_D^2}(B)\right]^{-\frac{1}{2}}$$

exists provided that B is closed and  $0 < C_{g_D^2}(B) < \infty$ . For similar results related to the  $\alpha$ -Green kernel with  $\alpha < 2$ , see Theorems 10.1 and 11.1 below.

From now on we shall often write simply  $\alpha$  instead of  $\kappa_{\alpha}$  if it serves as an index. E.g.,  $C_{\alpha}(\cdot) = C_{\kappa_{\alpha}}(\cdot)$  and  $C_{g_{D}^{\alpha}}(\cdot)$  denote the  $\alpha$ -Riesz and the  $\alpha$ -Green interior capacities of a set, respectively.

**Lemma 2.3.** For any  $B \subset D$ ,  $C_{\alpha}(B) = 0$  if and only if  $C_{g_{D}^{\alpha}}(B) = 0$ .

*Proof.* We need the following general facts related to an arbitrary strictly positive definite kernel  $\kappa$  on a locally compact space X. First of all, for any  $B \subset X$ ,

(2.8) 
$$C_{\kappa}(B) = \sup_{K \in \{K\}_B} C_{\kappa}(K),$$

<sup>&</sup>lt;sup>4</sup>For  $\alpha = 2$ , this definition is due to Brelot (see [4]; cf. also [11, 13]). For  $\alpha \in (0, 2)$ , such a notion has been introduced in [17].

where  $\{K\}_B$  consists of all compact subsets of B (see [10]). Further, for a compact set  $K \subset X$  one has  $C_{\kappa}(K) < \infty$  and hence, by [10, Theorem 2.5],

(2.9) 
$$C_{\kappa}(K) = \sup \mu(K),$$

where  $\mu$  ranges over all nonnegative measures supported by K with the additional property

$$U^{\mu}_{\kappa}(x) \leq 1$$
 for all  $x \in S^{\mu}_{\mathcal{X}}$ .

Now we apply representation (2.9) to a compact set  $K \subset D$  and each of the  $\alpha$ -Riesz and the  $\alpha$ -Green kernels, which is possible in view of their strict positive definiteness. Since for every  $\mu \in \mathfrak{M}^+(K)$ ,  $U^{\mu}_{\kappa_{\alpha}}(x) - U^{\mu}_{g^{\alpha}_{D}}(x)$  is bounded on K in consequence of (2.6), the lemma for B = K follows. To prove the lemma for any  $B \subset D$ , it is thus left to apply (2.8).  $\Box$ 

Let  $B \subset D$ . By Lemma 2.3, if some expression E(x) is valid n.e. in B, then  $C_{g_D^{\alpha}}(N) = 0$ , N being the set of all  $x \in B$  with E(x) not to hold; and also the other way around.

2.4. Condensers. Existence of minimizers. By a *condenser* in  $\mathbb{R}^n$  we mean an ordered pair  $\mathbf{B} = (B_1, B_2)$  of nonintersecting sets  $B_1, B_2 \subset \mathbb{R}^n$  (so far of arbitrary topological structure), treated as the *positive* and the *negative* plates of  $\mathbf{B}$ , respectively. Define

$$\mathfrak{M}(\mathbf{B}) := \big\{ \nu \in \mathfrak{M}(\mathbb{R}^n) : \ \nu^+ \in \mathfrak{M}^+(B_1), \ \nu^- \in \mathfrak{M}^+(B_2) \big\}, \\ \mathcal{E}_{\alpha}(\mathbf{B}) := \mathfrak{M}(\mathbf{B}) \cap \mathcal{E}_{\alpha}(\mathbb{R}^n).$$

Then the following theorem on the strong completeness is true (see [18, Theorem 1]; compare with [5] or [14, Theorem 1.19]).<sup>5</sup>

**Theorem 2.4.** Assume  $B_1$  and  $B_2$  to be closed in  $\mathbb{R}^n$ . Then the metric space  $\mathcal{E}_{\alpha}(\mathbf{B})$  is complete in the induced strong topology, and the strong convergence in this space implies the vague convergence to the same limit.

From now on we fix a (particular) condenser  $\mathbf{A} = (A_1, A_2)$  in  $\mathbb{R}^n$  with the plates  $A_2 := D^c$ and  $A_1 := F$ , where  $F \subseteq D$  is closed in the relative topology of D and  $C_{\alpha}(F) > 0$ . Thus (see also Section 2.3),

(2.10) 
$$C_{\alpha}(A_i) > 0 \text{ for all } i = 1, 2.$$

Also fix a unit two-dimensional numerical vector  $\mathbf{1} = (1, 1)$ , and define

$$\mathcal{E}_{\alpha}(\mathbf{A}, \mathbf{1}) := \{ \mu \in \mathcal{E}_{\alpha}(\mathbf{A}) : \mu^{+}(A_{1}) = \mu^{-}(A_{2}) = 1 \}.$$

In view of (2.10), the class  $\mathcal{E}_{\alpha}(\mathbf{A}, \mathbf{1})$  is nonempty and, hence, it makes sense to consider the variational problem on the existence of  $\lambda_{\mathbf{A}} \in \mathcal{E}_{\alpha}(\mathbf{A}, \mathbf{1})$  with

(2.11) 
$$E_{\alpha}(\lambda_{\mathbf{A}}) = \inf_{\mu \in \mathcal{E}_{\alpha}(\mathbf{A}, \mathbf{1})} E_{\alpha}(\mu) \quad (=: E_{\alpha}(\mathbf{A}, \mathbf{1})).$$

<sup>&</sup>lt;sup>5</sup>In fact, Theorem 2.4 holds true also for  $\alpha \in (2, n)$ ; see [18, Theorem 1]. Its proof is based on Deny's theorem [6] stating that, for the Riesz kernels,  $\mathcal{E}_{\alpha}$  can be completed by making use of distributions with finite energy. Regarding the history of the question, see [16, Theorem A] and [17, Theorem 1].

7

In particular, the following theorem on the solvability holds (see [17, Theorems 5, 7]).<sup>6</sup>

# Theorem 2.5. Let

$$\inf_{x \in F, \ y \in D^c} |x - y| > 0.$$

If, moreover,  $C_{\alpha}(F) < \infty$  then, for a solution to the minimum energy problem (2.11) to exist, it is necessary and sufficient that either  $C_{\alpha}(D^c) < \infty$  or  $D^c$  be not  $\alpha$ -thin at  $\omega$ .<sup>7</sup>

In the paper, we are mainly interested in the case

$$F \cap \partial_{\overline{\mathbb{R}^n}} D \neq \emptyset$$

where  $\overline{\mathbb{R}^n} := \mathbb{R}^n \cup \{\omega\}$  is the one-point compactification of  $\mathbb{R}^n$  and  $\overline{F} := C\ell_{\overline{\mathbb{R}^n}}F$ . Then, in general, the infimum value in (2.11) can not be achieved among  $\mu \in \mathcal{E}_{\alpha}(\mathbf{A}, \mathbf{1})$ . Using the physical interpretation, which is possible for the Coulomb kernel, a short-circuit between the plates of the condenser might occur.

Therefore, it is meaningful to investigate what kind of conditions on the objects under consideration would prevent such a phenomenon, and a minimizer in the corresponding minimum  $\alpha$ -Riesz energy problem for the condenser **A** would, nevertheless, exist. One of the ideas, to be discussed, is to find out such an upper constraint on the measures (charges) from  $\mathcal{E}^+_{\alpha}(F, 1)$  which would not allow the "blow-up" effect between F and  $D^c$ . Note that we do not intend to impose any constraint on the measures on  $D^c$ .

Assume also the measures from  $\mathcal{E}^+_{\alpha}(F)$  to be influenced by some external field f, while  $\langle f, \mu \rangle = 0$  for all  $\mu \in \mathcal{E}^+_{\alpha}(D^c)$ . Then, what kind of external fields, acting on the charges on F only, would still guarantee the existence of minimizers?

Recently a similar problem for the logarithmic kernel in  $\mathbb{R}^2$  has been investigated by Beckermann and Gryson [1, Theorem 2.2]. Our study is related to the Riesz kernels, and the results obtained and the approaches applied are rather different from those in [1].

## 3. Constrained and unconstrained minimum energy problems

When speaking of the external field f, we shall tacitly assume that at least one of the following Cases I or II takes place:

I.  $f|_F$  is lower semicontinuous, and it is  $\ge 0$  unless F is compact, while f(x) = 0 n.e. in  $D^c$ ;

II. 
$$f(x) = U_{\alpha}^{\zeta - \beta_{D^c}^{\alpha} \zeta}(x)$$
 for all  $x \in \mathbb{R}^n$ , where a (signed) measure  $\zeta \in \mathcal{E}_{g_D^{\alpha}}(D)$  is given.

<sup>&</sup>lt;sup>6</sup>The first result of this type was obtained in [16, Theorem 1], where  $\alpha = 2$  and  $A_1$  was assumed to be compact in D. See also [18, Theorem 12] where Theorem 2.5 has been generalized to any  $\alpha \in (0, n)$ and any  $\mathbf{a} = (a_1, a_2)$  with  $a_i > 0$ , i = 1, 2. Instead of the balayage technique, which implicitly appears in Theorem 2.5, for  $\alpha \in (2, n)$  one should use the operator of orthogonal projection in the pre-Hilbert space  $\mathcal{E}_{\alpha}$ onto the convex cone  $\mathcal{E}^+_{\alpha}(D^c)$ .

<sup>&</sup>lt;sup>7</sup>We refer to [16, 22] for an example of a set with infinite Newtonian capacity, though 2-thin at  $\omega$ .

Note that Case II can certainly be reduced to Case I provided that  $\zeta^- = 0$ . The values  $G_{\alpha,f}(\mu), \mu \in \mathcal{E}_{\alpha}(\mathbf{A})$ , and  $G_{g_D^{\alpha},f}(\nu), \nu \in \mathcal{E}_{g_D^{\alpha}}^+(F)$ , are well defined and

(3.1) 
$$G_{\alpha,f}(\mu) = \|\mu\|_{\alpha}^2 + 2\langle f, \mu^+ \rangle > -\infty,$$

(3.2) 
$$G_{g_D^{\alpha}, f}(\nu) = \|\nu\|_{g_D^{\alpha}}^2 + 2\langle f, \nu \rangle > -\infty.$$

If Case II holds, then these values can alternatively be expressed in the form

(3.3) 
$$G_{\alpha,f}(\mu) = \|\mu\|_{\alpha}^{2} + 2E_{\alpha}(\zeta - \beta_{D^{c}}^{\alpha}\zeta, \mu) = \|\mu + \zeta - \beta_{D^{c}}^{\alpha}\zeta\|_{\alpha}^{2} - \|\zeta - \beta_{D^{c}}^{\alpha}\zeta\|_{\alpha}^{2},$$

(3.4) 
$$G_{g_D^{\alpha},f}(\nu) = \|\nu\|_{g_D^{\alpha}}^2 + 2E_{g_D^{\alpha}}(\zeta,\nu) = \|\nu+\zeta\|_{g_D^{\alpha}}^2 - \|\zeta\|_{g_D^{\alpha}}^2.$$

Write

$$\mathcal{E}_{\alpha,f}(\mathbf{A},\mathbf{1}) := \mathcal{E}_{\alpha}(\mathbf{A},\mathbf{1}) \cap \mathcal{E}_{\alpha,f}(\mathbb{R}^n) \quad \text{and} \quad \mathcal{E}^+_{g^{\alpha}_{D},f}(F,1) := \mathcal{E}^+_{g^{\alpha}_{D}}(F,1) \cap \mathcal{E}_{g^{\alpha}_{D},f}(D);$$

these classes of measures are convex. It is seen from (3.3) and (3.4) that, in Case II,

(3.5) 
$$\mathcal{E}_{\alpha,f}(\mathbf{A},\mathbf{1}) = \mathcal{E}_{\alpha}(\mathbf{A},\mathbf{1}) \text{ and } \mathcal{E}_{g_D^{\alpha},f}^+(F,1) = \mathcal{E}_{g_D^{\alpha}}^+(F,1)$$

We denote by  $\mathfrak{C}(F)$  the collection of all  $\xi \in \mathfrak{M}^+(D)$  with the properties

$$S_D^{\xi} = F \quad \text{and} \quad \xi(F) > 1;$$

such  $\xi$  will be treated as *constraints* for measures from the class  $\mathfrak{M}^+(F,1)$ . Let  $\mathfrak{C}_0(F)$  consist of all bounded  $\xi \in \mathfrak{C}(F)$ , i.e., with  $\xi(F) < \infty$ . Given  $\xi \in \mathfrak{C}(F)$ , write

$$\mathcal{E}_{\alpha,f}^{\xi}(\mathbf{A},\mathbf{1}) := \left\{ \mu \in \mathcal{E}_{\alpha,f}(\mathbf{A},\mathbf{1}) : \ \mu^{+} \leqslant \xi \right\},\\ \mathcal{E}_{g_{D}^{\alpha},f}^{\xi}(F,1) := \left\{ \nu \in \mathcal{E}_{g_{D}^{\alpha},f}^{+}(F,1) : \ \nu \leqslant \xi \right\},$$

where  $\nu_1 \leq \nu_2$  means that  $\nu_2 - \nu_1$  is a nonnegative Borel measure.

To combine (where this is possible) assertions related to extremal problems in both the constrained and unconstrained settings, we accept the notations

$$\mathcal{E}^{\infty}_{\alpha,f}(\mathbf{A},\mathbf{1}) := \mathcal{E}_{\alpha,f}(\mathbf{A},\mathbf{1}) \quad \text{and} \quad \mathcal{E}^{\infty}_{g^{\alpha}_{D},f}(F,1) := \mathcal{E}^{+}_{g^{\alpha}_{D},f}(F,1).$$

In all that follows, we consider a fixed  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$ , where the formal formula  $\sigma = \infty$  means that *no* upper constraint is allowed,<sup>8</sup> and we define

$$G_{\alpha,f}^{\sigma}(\mathbf{A},\mathbf{1}) := \inf_{\substack{\mu \in \mathcal{E}_{\alpha,f}^{\sigma}(\mathbf{A},\mathbf{1})}} G_{\alpha,f}(\mu),$$
  
$$G_{g_{D}^{\alpha},f}^{\sigma}(F,1) := \inf_{\substack{\nu \in \mathcal{E}_{g_{D}^{\alpha},f}^{\sigma}(F,1)}} G_{g_{D}^{\alpha},f}(\nu).$$

In the case  $\sigma = \infty$ , the upper index  $\infty$  will often be omitted, i.e., we shall write

$$G_{\alpha,f}(\mathbf{A},\mathbf{1}) := G_{\alpha,f}^{\infty}(\mathbf{A},\mathbf{1}) \quad \text{and} \quad G_{g_D^{\alpha},f}(F,1) := G_{g_D^{\alpha},f}^{\infty}(F,1).$$

<sup>&</sup>lt;sup>8</sup>It is natural to set  $\nu \leq \infty$  for all  $\nu \in \mathfrak{M}^+(F)$ .

Note that each of  $G^{\sigma}_{\alpha,f}(\mathbf{A}, \mathbf{1})$  and  $G^{\sigma}_{g^{\alpha}_{D},f}(F, 1)$  is  $> -\infty$ . One can also see from (2.10) and Lemma 2.3 that any of the classes  $\mathcal{E}^{\sigma}_{\alpha,f}(\mathbf{A}, \mathbf{1})$  and  $\mathcal{E}^{\sigma}_{g^{\alpha}_{D},f}(F, 1)$  is nonempty if and only if so is the other, and therefore the following two assumptions are equivalent:<sup>9</sup>

$$(3.6) G^{\sigma}_{\alpha,f}(\mathbf{A},\mathbf{1}) < \infty$$

$$(3.7) G^{\sigma}_{g^{\alpha}_{D},f}(F,1) < \infty$$

**Problem 3.1.** Under condition (3.6), does there exist  $\lambda_{\mathbf{A}}^{\sigma} \in \mathcal{E}_{\alpha,f}^{\sigma}(\mathbf{A}, \mathbf{1})$  whose *f*-weighted  $\alpha$ -Riesz energy is minimal in this class, i.e.

(3.8) 
$$G_{\alpha,f}(\lambda_{\mathbf{A}}^{\sigma}) = G_{\alpha,f}^{\sigma}(\mathbf{A}, \mathbf{1})?$$

Problem 3.1 turns out to be intricately related to the following one.

**Problem 3.2.** Under condition (3.7), does there exist  $\lambda_F^{\sigma} \in \mathcal{E}_{g_D^{\alpha}, f}^{\sigma}(F, 1)$  whose *f*-weighted  $\alpha$ -Green energy is minimal in this class, i.e.

(3.9) 
$$G_{g^{\alpha}_{D},f}(\lambda^{\sigma}_{F}) = G^{\sigma}_{q^{\alpha}_{D},f}(F,1)?$$

Note that, if  $\sigma = \infty$  and f = 0, then Problem 3.1 is in fact reduced to the minimum energy problem (2.11), while Problem 3.2 to that appeared in (2.1) for B = F and  $\kappa = g_D^{\alpha}$ .

The rest of the paper is organized as follows. Sufficient and/or necessary conditions for Problems 3.1 and 3.2 to be solvable are established in Sections 5, 6 and 7. In Section 5 they are formulated either in measure theory terms for the constraints under consideration, or in geometric-potential terms for F and  $D^c$ , while in Sections 6 and 7 they are given in terms of variational inequalities for the f-weighted potentials. Sections 6 and 7 provide also a detailed analysis of the supports of the minimizers. The results obtained are proved in Sections 8, 9, 12 and 13, and they are illustrated by the examples in Section 14.

E.g., by Theorem 5.2, in both Cases I and II, the condition  $C_{g_D^{\alpha}}(F) < \infty$  is close to be sufficient for Problem 3.2 to be solvable for every  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$ , while according to Theorem 5.3, it is also necessary for this to happen provided Case II with  $\zeta \ge 0$  holds. However, if we restricted our analysis to the constraints from the class  $\mathfrak{C}_0(F)$  then, in Case I, Problem 3.2 would already be always solvable (see Theorem 5.1). Further, if we assume  $D^c$  to be not  $\alpha$ -thin at  $\omega$ , then all this remains true for Problem 3.1 as well. See Section 5 for the strict formulations of the results just described.

A crucial key in the proofs is Theorem 10.1, providing us with a perfectness-type result for the  $\alpha$ -Green kernel, where  $\alpha < 2$  (cf. [9] for  $\alpha = 2$ ). It makes it possible, in particular, to establish Theorem 11.1 on the existence of the  $\alpha$ -Green equilibrium measure of a set.

The uniqueness of solutions to Problems 3.1 and 3.2 is shown by Lemma 4.1 in the next section. Another assertion of this section, Lemma 4.2, discovers an intimate relationship

 $<sup>^{9}</sup>$ See Lemmas 4.5, 4.7 and Remark 4.8 below, providing necessary and/or sufficient conditions for (3.6) and (3.7) to hold.

between their solvability (or unsolvability), as well as their minimizers (provided they exist), which turns out to be a powerful tool in the proofs of the above-mentioned results.

## 4. AUXILIARY ASSERTIONS

**Lemma 4.1.** A solution to Problem 3.1, as well as that to Problem 3.2, is unique (provided *it exists*).

*Proof.* We shall verify the latter part of the lemma. Assume there exist two solutions to Problem 3.2,  $\lambda_F^{\sigma}$  and  $\hat{\lambda}_F^{\sigma}$ . Since the class  $\mathcal{E}_{g_{D}^{\alpha},f}^{\sigma}(F,1)$  is convex, from (3.2) we get

$$4G_{g_D^{\alpha},f}^{\sigma}(F,1) \leqslant 4G_{g_D^{\alpha},f}\Big(\frac{\lambda_F^{\sigma} + \hat{\lambda}_F^{\sigma}}{2}\Big) = \|\lambda_F^{\sigma} + \hat{\lambda}_F^{\sigma}\|_{g_D^{\alpha}}^2 + 4\langle f, \lambda_F^{\sigma} + \hat{\lambda}_F^{\sigma} \rangle.$$

Applying the parallelogram identity in  $\mathcal{E}_{g_D^{\alpha}}(D)$ , we obtain

$$0 \leqslant \|\lambda_F^{\sigma} - \hat{\lambda}_F^{\sigma}\|_{g_D^{\alpha}}^2 \leqslant -4G_{g_D^{\alpha},f}^{\sigma}(F,1) + 2G_{g_D^{\alpha},f}(\lambda_F^{\sigma}) + 2G_{g_D^{\alpha},f}(\hat{\lambda}_F^{\sigma}),$$

so that  $\|\lambda_F^{\sigma} - \hat{\lambda}_F^{\sigma}\|_{g_D^{\alpha}} = 0$  by (3.9). As  $g_D^{\alpha}$  is strictly positive definite,  $\lambda_F^{\sigma} = \hat{\lambda}_F^{\sigma}$ .

Likewise, the former part of the lemma can be proved with the help of the convexity of the class  $\mathcal{E}^{\sigma}_{\alpha,f}(\mathbf{A},\mathbf{1})$  and the pre-Hilbert structure in the space  $\mathcal{E}_{\alpha}(\mathbb{R}^n)$ .

**Lemma 4.2.** Assume  $D^c$  to be not  $\alpha$ -thin at  $\omega$ . Then for every  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$ ,

(4.1) 
$$G_{g_D,f}^{\sigma}(F,1) = G_{\alpha,f}^{\sigma}(\mathbf{A},\mathbf{1}).$$

In addition, the solution to Problem 3.1 exists if and only if so does that to Problem 3.2, and then they are related to each other by the formula

(4.2) 
$$\lambda_{\mathbf{A}}^{\sigma} = \lambda_{F}^{\sigma} - \beta_{D^{c}}^{\alpha} \lambda_{F}^{\sigma}.$$

*Proof.* We begin by establishing the inequality

(4.3) 
$$G^{\sigma}_{g^{\alpha}_{D},f}(F,1) \ge G^{\sigma}_{\alpha,f}(\mathbf{A},\mathbf{1})$$

Having assumed  $G_{g_D^{\alpha},f}^{\sigma}(F,1) < \infty$ , we fix  $\nu \in \mathcal{E}_{g_D^{\alpha},f}^{\sigma}(F,1)$ ; then, by (2.5) for  $Q = D^c$ ,

$$\nu - \beta_{D^c}^{\alpha} \nu \in \mathcal{E}^{\sigma}_{\alpha,f}(\mathbf{A}, \mathbf{1}).$$

On account of (2.7), (3.1) and (3.2), we therefore get

$$G_{g_D^{\alpha},f}(\nu) = \|\nu\|_{g_D^{\alpha}}^2 + 2\langle f,\nu\rangle = \|\nu - \beta_{D^c}^{\alpha}\nu\|_{\alpha}^2 + 2\langle f,\nu\rangle = G_{\alpha,f}(\nu - \beta_{D^c}^{\alpha}\nu) \ge G_{\alpha,f}^{\sigma}(\mathbf{A},\mathbf{1}).$$

Since  $\nu \in \mathcal{E}_{g_D^{\alpha}, f}^{\sigma}(F, 1)$  has been chosen arbitrarily, this yields (4.3).

On the other hand, for any  $\mu \in \mathcal{E}^{\sigma}_{\alpha,f}(\mathbf{A},\mathbf{1})$  we have  $\mu^+ \in \mathcal{E}^{\sigma}_{g^{\alpha}_{\alpha},f}(F,1)$ . Thus, due to (2.4),

(4.4) 
$$G_{\alpha,f}(\mu) = \|\mu\|_{\alpha}^{2} + 2\langle f, \mu^{+} \rangle \ge \|\mu^{+} - \beta_{D^{c}}^{\alpha}\mu^{+}\|_{\alpha}^{2} + 2\langle f, \mu^{+} \rangle$$
$$= \|\mu^{+}\|_{g_{D^{c}}}^{2} + 2\langle f, \mu^{+} \rangle = G_{g_{D}^{\alpha},f}(\mu^{+}) \ge G_{g_{D}^{\alpha},f}^{\sigma}(F,1).$$

In view of the arbitrary choice of  $\mu \in \mathcal{E}^{\sigma}_{\alpha,f}(\mathbf{A}, \mathbf{1})$ , this proves (4.1) when combined with (4.3). Let now  $\lambda^{\sigma}_{F} \in \mathcal{E}^{\sigma}_{g^{\alpha}_{D},f}(F, 1)$  satisfy (3.9). Then, in consequence of (2.5),

$$\widehat{\mu} := \lambda_F^{\sigma} - \beta_{D^c}^{\alpha} \lambda_F^{\sigma} \in \mathcal{E}_{\alpha,f}^{\sigma}(\mathbf{A}, \mathbf{1}).$$

Substituting  $\hat{\mu}$  instead of  $\mu$  in relation (4.4), we see that all the inequalities therein are, in fact, equalities. Therefore, by (4.1),

$$G_{\alpha,f}(\widehat{\mu}) = G_{g_{D^c},f}^{\sigma}(F,1) = G_{\alpha,f}^{\sigma}(\mathbf{A},\mathbf{1}),$$

so that the measure  $\lambda_{\mathbf{A}}^{\sigma}$ , defined by (4.2), solves Problem 3.1.

To complete the proof, assume further that  $\lambda_{\mathbf{A}}^{\sigma} = \lambda^{+} - \lambda^{-} \in \mathcal{E}_{\alpha,f}^{\sigma}(\mathbf{A}, \mathbf{1})$  satisfies (3.8). Then, by (4.1) and (4.4), the latter with  $\lambda_{\mathbf{A}}^{\sigma}$  instead of  $\mu$ ,

$$G_{g_D^{\alpha},f}^{\sigma}(F,1) = G_{\alpha,f}(\lambda_{\mathbf{A}}^{\sigma}) \ge \|\lambda^+ - \beta_{D^c}^{\alpha}\lambda^+\|_{\alpha}^2 + 2\langle f, \lambda^+ \rangle$$
$$= \|\lambda^+\|_{g_{D^c}^{\alpha}}^2 + 2\langle f, \lambda^+ \rangle = G_{g_D^{\alpha},f}(\lambda^+) \ge G_{g_D^{\alpha},f}^{\sigma}(F,1).$$

Hence, all the inequalities here are, in fact, equalities. This shows that  $\lambda_F^{\sigma} := \lambda^+$  solves Problem 3.2 and, on account of (2.4), also that  $\lambda^- = \beta_{D^c}^{\alpha} \lambda^+ = \beta_{D^c}^{\alpha} \lambda_F^{\sigma}$ .

**Lemma 4.3.** Assume (3.7) holds. For a measure  $\lambda = \lambda_F^{\sigma} \in \mathcal{E}_{g_D^{\sigma},f}^{\sigma}(F,1)$  to solve Problem 3.2, it is necessary and sufficient that

(4.5) 
$$\langle W_{g_D^{\alpha},f}^{\lambda}, \nu - \lambda \rangle \ge 0 \quad \text{for all } \nu \in \mathcal{E}_{g_D^{\alpha},f}^{\sigma}(F,1).$$

*Proof.* By direct calculation, for any  $\nu, \mu \in \mathcal{E}_{g_D^{\alpha}, f}^{\sigma}(F, 1)$  and any  $h \in (0, 1]$  we obtain

(4.6) 
$$G_{g_D^{\alpha},f}(h\nu + (1-h)\mu) - G_{g_D^{\alpha},f}(\mu) = 2h \langle W_{g_D^{\alpha},f}^{\mu}, \nu - \mu \rangle + h^2 \|\nu - \mu\|_{g_D^{\alpha}}^2.$$

If  $\mu = \lambda_F^{\sigma}$  solves Problem 3.2, then the left (hence, the right) side of (4.6) is  $\geq 0$ , for the class  $\mathcal{E}_{g_{\Omega,f}^{\sigma}}^{\sigma}(F,1)$  is convex, which leads to (4.5) by letting  $h \to 0$ .

Conversely, if (4.5) holds, then (4.6) with  $\mu = \lambda$  and h = 1 gives  $G_{g_D^{\alpha}, f}(\nu) \ge G_{g_D^{\alpha}, f}(\lambda)$  for all  $\nu \in \mathcal{E}_{g_D^{\alpha}, f}^{\sigma}(F, 1)$ , which means that  $\lambda = \lambda_F^{\sigma}$  solves Problem 3.2.

**Lemma 4.4.**  $G_{g_D^{\alpha},f}(\cdot)$  is vaguely lower semicontinuous on  $\mathcal{E}_{g_D^{\alpha}}^+(D)$  if Case I takes place, and otherwise, it is strongly continuous.

*Proof.* If Case I holds, then the lemma follows from [10] (see Section 1.1 and Lemma 2.2.1 therein), while otherwise it is a direct consequence of relation (3.4).

Lemmas 4.5 and 4.7 below provide sufficient and/or necessary conditions that guarantee (3.6) and (3.7) (compare with Lemmas 4 and 5 from [20]). From now on we write

(4.7) 
$$F_0 := \{ x \in F : f(x) < \infty \}.$$

**Lemma 4.5.** Let  $\sigma = \infty$ . Then (3.7) (hence, also (3.6)) holds if and only if  $C_{\alpha}(F_0) > 0$ .

*Proof.* Suppose first that  $C_{\alpha}(F_0) > 0$ . On account of [10, Lemma 2.3.3], then one can choose a compact set  $K \subset F$  with  $C_{\alpha}(K) > 0$  so that  $f(x) \leq M < \infty$  for all  $x \in K$ . In turn, this yields that there exists  $\nu \in \mathcal{E}_{g_{D}^{\alpha}}^{+}(K, 1)$  with  $G_{g_{D}^{\alpha}, f}(\nu) < \infty$ , and (3.7) follows.

To prove the necessary part of the lemma, assume, on the contrary, that  $C_{\alpha}(F_0) = 0$ . Then for every  $\mu \in \mathcal{E}_{g_{D}^{\alpha}}^{+}(F, 1)$ , we necessarily have  $G_{g_{D}^{\alpha}, f}(\nu) = \infty$ , which contradicts (3.7).  $\Box$ 

**Definition 4.6.**  $\xi \in \mathfrak{C}(F)$  is called *admissible* if its restriction to any compact subset of F has finite  $\alpha$ -Riesz (hence,  $\alpha$ -Green) energy. Let  $\mathcal{A}(F)$  consist of all admissible constraints.

When considering admissibility of a measure, the parameter  $\alpha$  and the set F should be clear in each context. Observe that, for a constraint  $\xi \in \mathfrak{C}(F)$  to be admissible, it is sufficient that its  $\alpha$ -Riesz potential be continuous. Also note that any  $\xi \in \mathcal{A}(F)$  is C-absolutely continuous.

**Lemma 4.7.** If  $\xi \in \mathcal{A}(F)$ , then (3.7) (hence, also (3.6)) holds provided that  $\xi(F_0) > 1$ .

*Proof.* Choose a compact set  $K \subset F$  so that  $\xi(K) > 1$  and  $f(x) \leq M < \infty$  for all  $x \in K$ . Then  $\xi|_K / \xi(K) \in \mathcal{E}_{g_D^{\alpha}, f}^{\xi}(F, 1)$ , which yields (3.7).

**Remark 4.8.** If Case II takes place, then f(x) is finite n.e. in F and, hence, Lemma 4.5 (similarly, Lemma 4.7) remains true with  $C_{\alpha}(F_0) > 0$  (respectively,  $\xi(F_0) > 1$ ) dropped.

# 5. Criteria of the solvability, given either in measure theory terms for $\sigma$ , or in geometric-potential terms for F and $D^c$

Throughout this section and Sections 6 and 7, assume (3.6) or, equivalently, (3.7) to be satisfied. See Lemmas 4.5, 4.7 and Remark 4.8 above, providing necessary and/or sufficient conditions for these to hold.

**Theorem 5.1.** If Case I takes place, then Problem 3.2 is (uniquely) solvable for every constraint  $\xi \in \mathfrak{C}_0(F)$ .

In the next theorem, the following condition on the geometry of the condenser **A** is required to hold:

(\*) If  $\alpha < 2$ , then  $\overline{F} \cap \partial_{\overline{\mathbb{R}^n}} D$  consists of at most one point.

Note that we do not impose any restriction on  $\overline{F} \cap \partial_{\mathbb{R}^n} D$  provided that  $\alpha = 2$ .

Theorem 5.2. If, moreover,

(5.1)  $C_{g_D^{\alpha}}(F) < \infty$ 

then, in both Cases I and II, Problem 3.2 is (uniquely) solvable for every  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$ .<sup>10</sup>

 $<sup>^{10}</sup>$ Compare with [21, Theorem 2.2] and [22, Theorem 8.1].

**Theorem 5.3.** Suppose that Case II with  $\zeta \ge 0$  takes place. If, moreover,  $C_{g_D^{\alpha}}(F) = \infty$ , then Problem 3.2 is unsolvable for every  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$  such that  $\sigma \ge \xi_0$ , where  $\xi_0 \in \mathfrak{C}(F) \setminus \mathfrak{C}_0(F)$  is properly chosen.

Combining Theorems 5.2 and 5.3 shows that, if assumption (\*) and Case II with  $\zeta \ge 0$  both hold, then (5.1) is necessary and sufficient for Problem 3.2 to be solvable for every  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$ .

**Theorem 5.4.** Assume  $D^c$  to be not  $\alpha$ -thin at  $\omega$ . Under the hypotheses of Theorem 5.1 (similarly, Theorems 5.2 or 5.3), its conclusion remains true for Problem 3.1 as well.

Indeed, Theorem 5.4 is obtained from Theorems 5.1–5.3 with the help of Lemma 4.2.

In the next two sections we shall examine properties of the *f*-weighted potentials and the supports of the minimizers  $\lambda_F^{\sigma}$  and  $\lambda_{\mathbf{A}}^{\sigma}$ , whose existence has been ensured, e.g., by Theorems 5.1, 5.2, and 5.4.

6. Variational inequalities for the f-weighted  $\alpha$ -Green potentials

This section provides necessary and/or sufficient conditions for the solvability of Problem 3.2 in terms of variational inequalities for the f-weighted  $\alpha$ -Green potentials. It also presents a detailed analysis of properties of the supports of the minimizers.

Following [14, p. 164], we denote by  $\breve{F}$  the reduced kernel of F, i.e.

(6.1)  $\breve{F} := \{ x \in F : C_{\alpha}(B(x,\varepsilon) \cap F) > 0 \text{ for every } \varepsilon > 0 \}.$ 

Here  $B(x,\varepsilon) := \{y \in \mathbb{R}^n : |y-x| < \varepsilon\}$ . Observe that, if the constraint under consideration is admissible, then necessarily  $F = \breve{F}$ .

To simplify the formulations of the results obtained, throughout this section and Section 7 we assume  $\partial D$  to be simultaneously the boundary of the (open) set Int  $D^c$ . Here, the boundary and the interior are considered relative to  $\mathbb{R}^n$ . Notice that then  $m_n(D^c) > 0$ , where  $m_n$  is the *n*-dimensional Lebesgue measure.

6.1. Variational inequalities in the constrained  $\alpha$ -Green minimum energy problems. We start by studying Problem 3.2 in the constrained case (i.e., for  $\sigma \neq \infty$ ). In this section, we consider  $\xi \in \mathcal{A}(F)$  and assume that  $\xi(F_0) > 1$ . Note that, for any  $\nu \in \mathcal{E}_{g_D^{\alpha}}^+(F)$ ,  $W_{g_D^{\alpha},f}^{\nu}(x)$  is well defined and  $\neq -\infty$  n.e. in F, while it is finite n.e. in  $F_0$  (see (4.7)).

**Theorem 6.1.** Let Case I take place. Then a measure  $\lambda \in \mathcal{E}_{g_{D}^{\alpha},f}^{\xi}(F,1)$  solves Problem 3.2 if and only if there exists  $w_{\lambda} \in \mathbb{R}$  possessing the following two properties:

- (6.2)  $W_{q_{\alpha,f}^{\alpha}}^{\lambda}(x) \ge w_{\lambda} \quad (\xi \lambda) \text{-}a.e. \text{ in } F,$
- (6.3)  $W_{a_{\alpha,f}^{\lambda}}^{\lambda}(x) \leq w_{\lambda} \quad for \ all \ x \in S_D^{\lambda}.$

**Corollary 6.2.** Let  $f|_D = U_{g_D^{\alpha}}^{\chi}$  for some  $\chi \in \mathfrak{M}^+(D)$ . If  $\lambda$  solves Problem 3.2, then  $C_{\alpha}(\partial D \cap S_{\mathbb{R}^n}^{\xi-\lambda}) = 0.$ 

When speaking of the *non-weighted case* f = 0, we simply write

$$\mathcal{E}_{g_D^{\alpha}}^{\xi}(F,1) := \big\{ \nu \in \mathcal{E}_{g_D^{\alpha}}^+(F,1) : \ \nu \leqslant \xi \big\}.$$

Then Problem 3.2 is in fact reduced to that on the existence of  $\lambda_0 \in \mathcal{E}_{g^{\alpha}_D}^{\xi}(F, 1)$  with

(6.4) 
$$E_{g_D^{\alpha}}(\lambda_0) = \inf_{\nu \in \mathcal{E}_{g_D^{\alpha}}^{\xi}(F,1)} E_{g_D^{\alpha}}(\nu).$$

**Corollary 6.3.** Let f = 0. A measure  $\lambda_0 \in \mathcal{E}_{g_D^{\alpha}}^{\xi}(F,1)$  solves Problem 3.2 if and only if there exists  $w'_{\lambda_0} \in (0,\infty)$  such that

(6.5) 
$$U_{g_D^{\alpha}}^{\lambda_0}(x) = w_{\lambda_0}' \quad (\xi - \lambda_0) \text{-}a.e. \text{ in } F$$

(6.6) 
$$U_{q_{\alpha}}^{\lambda_0}(x) \leq w'_{\lambda_0} \quad for \ all \ x \in D.$$

If, moreover,  $\alpha < 2$ , then also

On account of the uniqueness of a solution to Problem 3.2, such a  $w'_{\lambda_0}$  is unique (provided it exists). Integrating (6.5) with respect to  $\xi - \lambda_0$ , we get

(6.8) 
$$w'_{\lambda_0} = \frac{E_{g^{\alpha}_D}(\lambda_0, \xi - \lambda_0)}{(\xi - \lambda_0)(D)}.$$

6.2. Variational inequalities in the unconstrained  $\alpha$ -Green minimum energy problems. Throughout this section, it is assumed that  $\sigma = \infty$ . We proceed with criteria of the solvability of Problem 3.2, given in terms of variational inequalities for the *f*-weighted  $\alpha$ -Green potentials. In the unconstrained case, the results obtained take a simpler form if compare with those in the constrained case, while provide us with much more detailed information about the potentials and the supports of the minimizers.

**Theorem 6.4.** Suppose that Case I takes place. For  $\lambda \in \mathcal{E}^+_{g^{\alpha}_{D},f}(F,1)$  to solve Problem 3.2, it is necessary and sufficient that there exist  $w_f \in \mathbb{R}$  possessing the properties

(6.9) 
$$W_{q_{D}^{\alpha},f}^{\lambda}(x) \ge w_{f} \quad n.e. \text{ in } F,$$

(6.10) 
$$W_{a_{\alpha,f}^{\alpha}}^{\lambda}(x) \leqslant w_f \quad \text{for all } x \in S_D^{\lambda}$$

Such a number  $w_f$  is unique (provided it exists) and can be given by the formula

$$w_f = \left\langle W_{g_D^{\alpha}, f}^{\lambda}, \lambda \right\rangle.$$

**Corollary 6.5.** Let Problem 3.2 be solvable. Then the following two assertions hold:

- (a) If Case II with  $\zeta \ge 0$  takes place, then  $C_{g^{\alpha}_{D}}(F) < \infty$ ;
- (b) If  $f|_{D} = U_{q_{D}^{\alpha}}^{\chi}$  for  $\chi \in \mathfrak{M}^{+}(D)$ , then  $C_{\alpha}(\partial D \cap C\ell_{\mathbb{R}^{n}}\check{F}) = 0$ .

**Corollary 6.6.** Let f = 0. Then  $\lambda_F \in \mathcal{E}^+_{g^+_D}(F, 1)$  solves Problem 3.2 if and only if there exists a number  $w \in (0, \infty)$  admitting the properties

(6.11) 
$$U_{g_D^{\lambda_F}}^{\lambda_F}(x) = w \quad n.e. \ in \ F,$$

(6.12) 
$$U_{g_D^{\Lambda}}^{\lambda_F}(x) \leq w \quad \text{for all } x \in D.$$

Such a number w is unique (provided it exists) and can be written in the form

$$w = E_{g_D^{\alpha}}(\lambda_F) = w_{g_D^{\alpha}}(F) = \left[C_{g_D^{\alpha}}(F)\right]^{-1}.$$

Furthermore, if the minimizer  $\lambda_F$  exists, then it is the unique measure in the class  $\mathcal{E}_{g_D^{\alpha}}^+(F, 1)$ whose  $\alpha$ -Green potential is constant n.e. in F. Namely, if  $\nu \in \mathcal{E}_{g_D^{\alpha}}^+(F, 1)$  and  $U_{g_D^{\alpha}}^{\nu}(x) = c$ n.e. in F, where  $c \in \mathbb{R}$ , then  $\nu = \lambda_F$ .

Recall that the reduced kernel  $\check{F}$  of F has been defined by (6.1). For the sake of simplicity, in the following assertion we assume that, if  $\alpha = 2$ , then  $D \setminus F$  is connected.

**Corollary 6.7.** Let f = 0. If  $\lambda_F$  solves Problem 3.2, then, in addition to (6.11) and (6.12), we have

(6.13) 
$$U_{g_D^{\alpha}}^{\lambda_F}(x) < w \quad for \ all \ x \in D \setminus \check{F}.$$

Furthermore,

(6.14) 
$$S_D^{\lambda_F} = \begin{cases} \breve{F} & \text{if } \alpha < 2, \\ \partial_D \breve{F} & \text{if } \alpha = 2. \end{cases}$$

6.3. Duality relation between non-weighted constrained and weighted unconstrained  $\alpha$ -Green minimum energy problems. Throughout this section, F is compact. Consider the non-weighted Problem 3.2 with a constraint  $\xi \in \mathfrak{C}(F)$  whose potential  $U_{g_{D}^{\alpha}}^{\xi}(x)$  is continuous.<sup>11</sup> Note that then  $E_{g_{D}^{\alpha}}(\xi) < \infty$ . Also assume that  $\lambda_{0}$  is its solution, i.e. both  $\lambda_{0} \in \mathcal{E}_{g_{D}^{\alpha}}^{\xi}(F, 1)$  and (6.4) hold. Write

$$\theta := q(\xi - \lambda_0), \quad \text{where} \quad q := \frac{1}{\xi(F) - 1}$$

Combining Corollary 6.3 and Theorem 6.4 allows us to formulate the following result.

<sup>&</sup>lt;sup>11</sup>When speaking of a continuous function, we understand that the values are *finite* numbers.

**Theorem 6.8.** The measure  $\theta$  solves Problem 3.2 with the external field  $f(x) := -qU_{g_D^{\alpha}}^{\xi}(x)$  in both the unconstrained and the  $q\xi$ -constrained settings, i.e.

$$\theta \in \mathcal{E}_{g_{D}^{\alpha},f}^{q\xi}(F,1) \subset \mathcal{E}_{g_{D}^{\alpha},f}^{+}(F,1) \quad and \quad G_{g_{D}^{\alpha},f}(\theta) = G_{g_{D}^{\alpha},f}^{q\xi}(F,1) = G_{g_{D}^{\alpha},f}(F,1).$$

Moreover,

(6.15) 
$$W^{\theta}_{g^{\alpha}_{D},f}(x) = -qw'_{\lambda_{0}} \quad on \ S^{\theta}_{D},$$

(6.16) 
$$W^{\theta}_{g^{\alpha}_{D},f}(x) \ge -qw'_{\lambda_{0}} \quad on \ D,$$

where  $w'_{\lambda_0}$  is the number determined by (6.8).

# 7. Variational inequalities for the f-weighted $\alpha$ -Riesz potentials

This section is devoted to necessary and/or sufficient conditions for the solvability of Problem 3.1 with  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$ , given in terms of variational inequalities for the *f*-weighted  $\alpha$ -Riesz potentials. Throughout this section, we assume  $D^c$  to be not  $\alpha$ -thin at  $\omega$ .

Then, by Lemma 4.2, for  $\lambda_{\mathbf{A}}^{\sigma} = \lambda^{+} - \lambda^{-}$  to solve Problem 3.1, it is necessary and sufficient that  $\lambda^{+}$  solve Problem 3.2 with the same  $\sigma$ . Furthermore, by (4.2),

(7.1) 
$$\lambda^{-} = \beta^{\alpha}_{D^{c}} \lambda^{+},$$

which yields

$$W_{\alpha,f}^{\lambda_{\mathbf{A}}^{\sigma}}(x) = U_{\alpha}^{\lambda^{+} - \beta_{D^{c}}^{\alpha} \lambda^{+}}(x) + f(x) = W_{g_{D}^{\alpha},f}^{\lambda^{+}}(x) \quad \text{for all} \quad x \in D.$$

For the sake of simplicity, in the next assertion we assume that in the case  $\alpha = 2$ , D is simply connected.

**Lemma 7.1.** If  $\lambda_{\mathbf{A}}^{\sigma} = \lambda^{+} - \lambda^{-}$  solves Problem 3.1, then

(7.2) 
$$S_{\mathbb{R}^n}^{\lambda^-} = \begin{cases} D^c & \text{if } \alpha < 2, \\ \partial D & \text{if } \alpha = 2. \end{cases}$$

Indeed, Lemma 7.1 follows from (7.1) and the description of the supports of the  $\alpha$ -Riesz balayage measures.

7.1. Variational inequalities in the constrained  $\alpha$ -Riesz minimum energy problems. In this section, consider  $\xi \in \mathcal{A}(F)$  and assume  $\xi(F_0) > 1$ , where  $F_0$  was given by (4.7). Combining what has been noticed just above with the assertions of Section 6.1 (for  $\lambda^+$  instead of  $\lambda$  or  $\lambda_0$ ) results in the following Theorem 7.2 and Corollaries 7.3 and 7.4. **Theorem 7.2.** Let Case I take place. Then  $\lambda_{\mathbf{A}}^{\xi} = \lambda^{+} - \lambda^{-} \in \mathcal{E}_{\alpha,f}^{\xi}(\mathbf{A}, \mathbf{1})$  is the (unique) solution to Problem 3.1 if and only if (7.1) holds and, in addition, there exists  $w_{\lambda_{\mathbf{A}}^{\xi}} \in \mathbb{R}$  possessing the following two properties:

$$\begin{split} W^{\lambda_{\mathbf{A}}^{\xi}}_{\alpha,f}(x) &\geqslant w_{\lambda_{\mathbf{A}}^{\xi}} \quad (\xi - \lambda^{+}) \text{-}a.e. \ in \ F, \\ W^{\lambda_{\mathbf{A}}^{\xi}}_{\alpha,f}(x) &\leqslant w_{\lambda_{\mathbf{A}}^{\xi}} \quad for \ all \ x \in S^{\lambda^{+}}_{D}. \end{split}$$

**Corollary 7.3.** Assume that  $f|_D = U_{g_D^{\alpha}}^{\chi}$  for some  $\chi \in \mathfrak{M}^+(D)$ . If  $\lambda_{\mathbf{A}}^{\xi} = \lambda^+ - \lambda^-$  solves Problem 3.1, then  $C_{\alpha}(\partial D \cap S_{\mathbb{R}^n}^{\xi-\lambda^+}) = 0$ .

**Corollary 7.4.** Let f = 0. A measure  $\lambda_{\mathbf{A}}^{\xi} = \lambda^{+} - \lambda^{-} \in \mathcal{E}_{\alpha,f}^{\xi}(\mathbf{A}, \mathbf{1})$  solves Problem 3.1 if and only if (7.1) holds and, in addition, there exists a (unique) number  $w'_{\lambda_{\mathbf{A}}^{\xi}} \in (0, \infty)$  such that

$$U_{\alpha}^{\lambda_{\mathbf{A}}^{\xi}}(x) = w'_{\lambda_{\mathbf{A}}^{\xi}} \quad (\xi - \lambda^{+}) \text{-a.e. in } F,$$
$$U_{\alpha}^{\lambda_{\mathbf{A}}^{\xi}}(x) \leq w'_{\lambda_{\mathbf{A}}^{\xi}} \quad \text{for all } x \in D.$$

Furthermore, if  $\alpha < 2$ , then also  $S_D^{\lambda^+} = F$  and  $S_{\mathbb{R}^n}^{\lambda^-} = D^c$ .

In the notations of Corollary 6.3 with  $\lambda^+$  in place of  $\lambda_0$ ,

$$w_{\lambda_{\mathbf{A}}^{\xi}}' = \frac{E_{\alpha}(\lambda_{\mathbf{A}}^{\xi}, \xi - \lambda^{+})}{(\xi - \lambda^{+})(D)} = \frac{E_{g_{D}^{\alpha}}(\lambda^{+}, \xi - \lambda^{+})}{(\xi - \lambda^{+})(D)} = w_{\lambda^{+}}'.$$

7.2. Variational inequalities in the unconstrained  $\alpha$ -Riesz minimum energy problems. In this section,  $\sigma = \infty$ . Similarly as it has been done just above, we derive the following corollaries from the assertions of Section 6.2.

**Corollary 7.5.** Assume Case I takes place. A measure  $\lambda_{\mathbf{A}} = \lambda^+ - \lambda^- \in \mathcal{E}_{\alpha,f}(\mathbf{A}, \mathbf{1})$  solves Problem 3.1 if and only if (7.1) holds and, in addition, there exists a (unique) number  $w'_f \in \mathbb{R}$  possessing the properties

$$\begin{split} W_{\alpha,f}^{\lambda_{\mathbf{A}}}(x) \geqslant w'_{f} \quad n.e. \ in \ F, \\ W_{\alpha,f}^{\lambda_{\mathbf{A}}}(x) \leqslant w'_{f} \quad for \ all \ x \in S_{D}^{\lambda^{+}}. \end{split}$$

Furthermore, then  $w'_f = w_f$ , where  $w_f$  is the number from Theorem 6.4, and assertions (a) and (b) of Corollary 6.5 both hold.

For the sake of simplicity, in the following assertion we assume that in the case  $\alpha = 2$ ,  $D \setminus F$  is simply connected.

**Corollary 7.6.** Let f = 0. A measure  $\lambda_{\mathbf{A}} = \lambda^+ - \lambda^- \in \mathcal{E}_{\alpha}(\mathbf{A}, \mathbf{1})$  solves Problem 3.1 if and only if (7.1) holds and there exists a (unique) number  $w' \in (0, \infty)$  such that

$$\begin{aligned} U_{\alpha}^{\lambda_{\mathbf{A}}}(x) &= w' \quad n.e. \ in \quad F, \\ U_{\alpha}^{\lambda_{\mathbf{A}}}(x) &\leq w' \quad for \ all \quad x \in D, \\ U_{\alpha}^{\lambda_{\mathbf{A}}}(x) &< w' \quad for \ all \quad x \in D \setminus \breve{F}. \end{aligned}$$

Furthermore, then w' = w, where w is the number from Corollary 6.6, i.e.

$$w' = E_{\alpha}(\lambda_{\mathbf{A}}, \lambda^{+}) = E_{\alpha}(\lambda_{\mathbf{A}}) = E_{g_{D}^{\alpha}}(\lambda^{+}) = w_{g_{D}^{\alpha}}(F) = \left[Cg_{D}^{\alpha}(F)\right]^{-1} = E_{\alpha}(\mathbf{A}, \mathbf{1}).$$

The descriptions of  $S_D^{\lambda^+}$  and  $S_{\mathbb{R}^n}^{\lambda^-}$  are given by (6.14) for  $\lambda^+$  in place of  $\lambda_F$  and (7.2), respectively.

# 8. Proof of Theorem 5.1

Consider an exhaustion of F by an increasing sequence of compact sets  $K_k, k \in \mathbb{N}$ . Since the constraint  $\xi$  is bounded, it holds

(8.1) 
$$\lim_{k \to \infty} \xi(F \setminus K_k) = 0.$$

Because of assumption (3.7), there exists  $\{\mu_\ell\}_{\ell\in\mathbb{N}}\subset \mathcal{E}^{\xi}_{g^{\alpha}_{\mathcal{D},f}}(F,1)$  such that

(8.2) 
$$\lim_{\ell \to \infty} G_{g_D^{\alpha}, f}(\mu_\ell) = G_{g_D^{\alpha}, f}^{\xi}(F, 1).$$

This sequence  $\{\mu_\ell\}_{\ell\in\mathbb{N}}$  is vaguely bounded; hence, by [2, Chapter III, Section 2, Prop. 9], it has a vague cluster point  $\mu_0$ . We assert that, in Case I,  $\mu_0$  is a solution to Problem 3.2.

Since  $\mathfrak{M}^+(F)$  is a vaguely closed subset of  $\mathfrak{M}^+(D)$ , we get  $\mu_0 \in \mathfrak{M}^+(F)$  and  $\mu_0 \leq \xi$ . Let, further,  $\{\mu_{\ell_m}\}_{m \in \mathbb{N}}$  be a subsequence of  $\{\mu_\ell\}_{\ell \in \mathbb{N}}$  converging vaguely to  $\mu_0$ . Then

$$1 = \lim_{m \to \infty} \mu_{\ell_m}(F) \ge \mu_0(F) = \lim_{k \to \infty} \mu_0(K_k)$$
$$\ge \lim_{k \to \infty} \limsup_{m \to \infty} \mu_{\ell_m}(K_k) = 1 - \lim_{k \to \infty} \liminf_{m \to \infty} \mu_{\ell_m}(F \setminus K_k).$$

On account of the fact that  $\mu_{\ell_m}(F \setminus K_k) \leq \xi(F \setminus K_k)$  for all  $m, k \in \mathbb{N}$ , combining the preceding chain of inequalities with (8.1) yields  $\mu_0(F) = 1$ .

To complete the proof, it thus remains to observe that  $G_{g_D^{\alpha},f}(\mu_0) \leq G_{g_D^{\alpha},f}^{\xi}(F,1)$ , which is seen from (8.2) in view of the lower semicontinuity of  $G_{g_D^{\alpha},f}$  on  $\mathcal{E}_{g_D^{\alpha}}^+(D)$  (see Lemma 4.4).

# 9. Proof of Theorem 5.3

Under the assumptions of the theorem, Case II with  $\zeta \ge 0$  takes place, and therefore

(9.1) 
$$G_{g_D^{\alpha},f}(\nu) = \|\nu\|_{g_D^{\alpha}}^2 + 2E_{g_D^{\alpha}}(\zeta,\nu) \ge E_{g_D^{\alpha}}(\nu) \ge 0 \quad \text{for all} \quad \nu \in \mathcal{E}_{g_D^{\alpha}}^+(D).$$

Consider an exhaustion of F by an increasing sequence of compact sets  $K_k$ ,  $k \in \mathbb{N}$ . Since  $C_{g_D^{\alpha}}(F) = \infty$ , it holds  $C_{g_D^{\alpha}}(F \setminus K_k) = \infty$  for all  $k \in \mathbb{N}$ . Hence, for every k one can choose a measure  $\nu_k \in \mathcal{E}_{q_D^{\alpha}}^+(F \setminus K_k, 1)$  with compact support so that

(9.2) 
$$\lim_{k \to \infty} \|\nu_k\|_{g_D^{\alpha}}^2 = 0$$

Certainly, there is no loss of generality in assuming  $K_k \cup S_D^{\nu_k} \subset K_{k+1}$ .

Fix  $\xi \in \mathfrak{C}(F)$  and write  $\xi_0 := \xi + \sum_{k \in \mathbb{N}} \nu_k$ ; then  $\xi_0 \in \mathfrak{C}(F) \setminus \mathfrak{C}_0(F)$ . Due to (3.5), for each  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$  such that  $\sigma \ge \xi_0$ , it holds

$$\nu_k \in \mathcal{E}^{\sigma}_{q^{\alpha}_{\infty},f}(F,1)$$
 for all  $k \in \mathbb{N}$ .

From (9.2) and the Cauchy–Schwarz inequality we get

$$\lim_{k \to \infty} G_{g_D^{\alpha}, f}(\nu_k) = \lim_{k \to \infty} \left[ \|\nu_k\|_{g_D^{\alpha}}^2 + 2E_{g_D^{\alpha}}(\zeta, \nu_k) \right] \leqslant 2 \|\zeta\|_{g_D^{\alpha}} \lim_{k \to \infty} \|\nu_k\|_{g_D^{\alpha}} = 0$$

Combined with (9.1), this yields  $G_{g_D^{\alpha},f}^{\sigma}(F,1) = 0$ . Repeated application of (9.1) shows also that such an infimum value can be attained only at zero measure. As  $0 \notin \mathcal{E}_{g_D^{\alpha},f}^{\sigma}(F,1)$ , Problem 3.2 with  $\sigma$  specified above is unsolvable.

#### 10. $\alpha$ -Green strong completeness theorem

A crucial point in our proof of Theorem 5.2, given in Section 12, is the following perfectnesstype result for the  $\alpha$ -Green kernel.

**Theorem 10.1.** Let  $E \subset \mathbb{R}^n$  be closed in D. If  $\alpha < 2$ , require additionally  $\overline{E} \cap \partial_{\overline{\mathbb{R}^n}} D$  to consist of at most one point. Then any strong Cauchy sequence  $\{\nu_k\}_{k\in\mathbb{N}} \subset \mathcal{E}_{g_D^{\alpha}}^+(E)$  with

(10.1) 
$$\sup_{k\in\mathbb{N}}\nu_k(E)<\infty$$

converges both strongly and vaguely to the unique  $\nu_0 \in \mathcal{E}_{q_{D}^{+}}^{+}(E)$ .

*Proof.* We can certainly assume that  $\alpha < 2$ , since otherwise the theorem holds true due to the perfectness of the  $g_D^2$ -kernel, established in [9].

The (strongly fundamental) sequence  $\{\nu_k\}_{k\in\mathbb{N}}\subset \mathcal{E}_{g^{\alpha}_{D}}^{+}(E)$  is strongly bounded, i.e.

$$\sup_{k\in\mathbb{N}} \|\nu_k\|_{g_D^\alpha} < \infty.$$

Besides, by (10.1),  $\{\nu_k\}_{k\in\mathbb{N}}$  is vaguely bounded, and hence it has a vague cluster point  $\nu_0$ . Since  $\mathcal{E}^+_{q^{\alpha}_{D}}(E)$  is a vaguely closed subset of  $\mathfrak{M}^+(D)$ , we have  $\nu_0 \in \mathcal{E}^+_{q^{\alpha}_{D}}(E)$ .

Let  $\{\nu_{k_{\ell}}\}_{\ell \in \mathbb{N}}$  be a subsequence of  $\{\nu_k\}_{k \in \mathbb{N}}$  such that

(10.2) 
$$\nu_{k_{\ell}} \to \nu_0$$
 vaguely as  $\ell \to \infty$ 

We next proceed to show that  $\nu_{k_{\ell}} \to \nu_0$  also strongly in  $\mathcal{E}_{g_D^{\alpha}}^+(E)$ , i.e.

(10.3) 
$$\lim_{\ell \to \infty} \|\nu_{k_{\ell}} - \nu_0\|_{g_D^{\alpha}} = 0.$$

As  $\{\nu_{k_\ell}\}_{\ell \in \mathbb{N}} \subset \mathcal{E}_{g_D^{\alpha}}^+(E)$ , being a subsequence of the strong Cauchy sequence  $\{\nu_k\}_{k \in \mathbb{N}}$ , is strongly fundamental as well, we see from (2.7) that

(10.4) 
$$\widetilde{\nu_{k_{\ell}}} := \nu_{k_{\ell}} - \beta_{D^c}^{\alpha} \nu_{k_{\ell}}, \quad \ell \in \mathbb{N}.$$

is strongly fundamental in  $\mathcal{E}_{\alpha}(\mathbb{R}^n)$ . The proof of (10.3) is given in two steps.

Step 1. Throughout this step, let  $\overline{E} \cap \partial_{\mathbb{R}^n} D$  either be empty or consist of only  $\omega$ . Then  $\nu_{k_\ell}$  and  $\beta_{D^c}^{\alpha} \nu_{k_\ell}$  are supported by the sets E and  $D^c$ , which due to the assumptions made are closed in  $\mathbb{R}^n$  and nonintersecting. Consider the condenser  $\mathbf{B} := (E, D^c)$ . The strong completeness theorem from [18] (or see Theorem 2.4 above) yields that there exists the unique measure  $\tilde{\nu} = \tilde{\nu}^+ - \tilde{\nu}^- \in \mathcal{E}_{\alpha}(\mathbf{B})$  such that

(10.5) 
$$\lim_{\ell \to \infty} \|\widetilde{\nu_{k_{\ell}}} - \widetilde{\nu}\|_{\alpha} = 0.$$

Furthermore, by this theorem,  $\tilde{\nu}^+$  and  $\tilde{\nu}^-$  are the vague limits of the positive and the negative parts of  $\tilde{\nu}_{k_{\ell}}$ ,  $\ell \in \mathbb{N}$ , respectively. In view of (10.2), we thus have

(10.6) 
$$\tilde{\nu}^+ = \nu_0,$$

for the vague topology is Hausdorff.

By the remark in [10, p. 166], it follows from (10.5) that there exists a subsequence of the sequence  $\{\widetilde{\nu_{k_\ell}}\}_{\ell \in \mathbb{N}}$  (denote it again by the same symbol) such that

$$U^{\widetilde{\nu}}_{\alpha}(x) = \lim_{\ell \to \infty} \, U^{\widetilde{\nu_{k_{\ell}}}}_{\alpha}(x) \quad \text{n.e. in} \ \ \mathbb{R}^n$$

On account of (10.4) and the countable subadditivity of  $C_{\alpha}(\cdot)$  over Borel sets, we see from the preceding relation that  $U^{\tilde{\nu}}_{\alpha}(x) = 0$  n.e. in  $D^c$  and, therefore,  $\tilde{\nu}^- = \beta^{\alpha}_{D^c} \tilde{\nu}^+$ . Combining this with (10.4), (10.5) and (10.6) implies

$$\lim_{\ell \to \infty} \left\| \left( \nu_{k_{\ell}} - \nu_0 \right) - \beta_{D^c}^{\alpha} \left( \nu_{k_{\ell}} - \nu_0 \right) \right\|_{\alpha} = 0,$$

which in view of (2.7) establishes (10.3).

**Step 2.** We next prove relation (10.3) in the case  $\overline{E} \cap \partial_{\mathbb{R}^n} D = \{x_0\}$ , where  $x_0 \neq \omega$ . Throughout this step, all the measures can be assumed to have zero mass at  $x_0$ , for we can restrict our consideration to those with finite energy. Define the inversion with respect to  $S(x_0, 1)$ , namely, each point  $x \neq x_0$  is mapped to the point  $x^*$  on the ray through x which issues from  $x_0$ , determined uniquely by

$$|x - x_0| \cdot |x^* - x_0| = 1.$$

This is a one-to-one, bicontinuous mapping of  $\mathbb{R}^n \setminus \{x_0\}$  onto itself; furthermore,

(10.7) 
$$|x^* - y^*| = \frac{|x - y|}{|x_0 - x||x_0 - y|}.$$

It can be extended to a one-to-one, bicontinuous map of  $\overline{\mathbb{R}^n}$  onto itself by setting  $x_0 \mapsto \omega$ .

To each  $\nu \in \mathfrak{M}(\mathbb{R}^n)$  (with  $\nu(\{x_0\}) = 0$ ) we correspond the Kelvin transform  $\nu^* \in \mathfrak{M}(\mathbb{R}^n)$  by means of the formula

$$d\nu^*(x^*) = |x - x_0|^{\alpha - n} d\nu(x), \quad x^* \in \mathbb{R}^n$$

Then, in view of (10.7),

(10.8) 
$$U_{\alpha}^{\nu^{*}}(x^{*}) = |x - x_{0}|^{n - \alpha} U_{\alpha}^{\nu}(x), \quad x^{*} \in \mathbb{R}^{n},$$

and therefore

(10.9)  $E_{\alpha}(\nu^*) = E_{\alpha}(\nu)$ 

(see [14, Chapter IV, Section 5, n° 19] and [14, Chapter V, Section 2, n° 8], respectively).

It is obvious that the Kelvin transformation is additive, i.e.

(10.10) 
$$(\nu_1 + \nu_2)^* = \nu_1^* + \nu_2^*.$$

We also observe that

(10.11) 
$$(\beta_{D^c}^{\alpha}\nu)^* = \beta_{(\overline{D^c})^*}^{\alpha}\nu^*,$$

where  $(\overline{D^c})^*$  is the image of  $\overline{D^c}$  under the inversion  $x \mapsto x^*$ .<sup>12</sup> Indeed, in view of (10.8) and the definition of the  $\alpha$ -Riesz balayage, we get

$$U_{\alpha}^{(\beta_{D^c}^{\alpha}\nu)^*}(x^*) = |x - x_0|^{n-\alpha} U_{\alpha}^{\beta_{D^c}^{\alpha}\nu}(x) = |x - x_0|^{n-\alpha} U_{\alpha}^{\nu}(x) = U_{\alpha}^{\nu^*}(x^*),$$

the relation being valid for nearly all  $x \in D^c$ . Consequently, it also holds for nearly all  $x^* \in (\overline{D^c})^*$ , because the inversion of a set with  $C_{\alpha}(\cdot) = 0$  has the interior  $\alpha$ -Riesz capacity zero as well (see [14, Chapter IV, Section 5, n° 19]). Since  $(\beta_{D^c}^{\alpha}\nu)^*$  is supported by  $(\overline{D^c})^*$ , identity (10.11) follows.

Applying [14, Lemma 4.3] to  $\nu_{k_{\ell}}$ ,  $\ell \in \mathbb{N}$ , and  $\nu_0$  (where  $\nu_{k_{\ell}}$ ,  $\ell \in \mathbb{N}$ , and  $\nu_0$  are as above), on account of (10.1) and (10.2) we have

(10.12) 
$$\nu_{k_{\ell}}^* \to \nu_0^*$$
 vaguely as  $\ell \to \infty$ .

<sup>12</sup>In fact,  $(\overline{D^c})^* = (D^c)^* \cup \{x_0\}.$ 

Also observe that, according to (10.9) and the fact that  $\{\widetilde{\nu_{k_\ell}}\}_{\ell \in \mathbb{N}}$  is strongly fundamental, so is the sequence  $(\widetilde{\nu_{k_\ell}})^* \in \mathcal{E}_{\alpha}(\mathbb{R}^n), \ell \in \mathbb{N}$ , which in consequence of (10.4), (10.10) and (10.11) can be rewritten in the form

(10.13) 
$$\left(\widetilde{\nu_{k_{\ell}}}\right)^* = \nu_{k_{\ell}}^* - \left(\beta_{D^c}^{\alpha}\nu_{k_{\ell}}\right)^* = \nu_{k_{\ell}}^* - \beta_{(\overline{D^c})^*}^{\alpha}\nu_{k_{\ell}}^*, \quad \ell \in \mathbb{N}.$$

The positive and the negative parts of  $(\widetilde{\nu_{k_\ell}})^*$  are supported by the sets  $E^*$  and  $(\overline{D^c})^*$ , respectively, which are closed in  $\mathbb{R}^n$  and nonintersecting; hence, the strong completeness theorem from [18] (see Theorem 2.4 above) can be applied. Therefore, there exists the unique measure  $\hat{\nu} = \hat{\nu}^+ - \hat{\nu}^- \in \mathcal{E}_{\alpha}(\mathbb{R}^n)$ , where  $\hat{\nu}^+$  and  $\hat{\nu}^-$  are supported by  $E^*$  and  $(\overline{D^c})^*$ , respectively, such that

(10.14) 
$$\lim_{\ell \to \infty} \left\| \left( \widetilde{\nu_{k_\ell}} \right)^* - \hat{\nu} \right\|_{\alpha} = 0.$$

Furthermore,  $\hat{\nu}^+$  and  $\hat{\nu}^-$  are the vague limits of the positive and the negative parts of  $(\widetilde{\nu_{k_\ell}})^*$ ,  $\ell \in \mathbb{N}$ , respectively. When combined with (10.12), (10.13) and the fact that the vague topology is Hausdorff, this implies

(10.15) 
$$\hat{\nu}^+ = \nu_0^*.$$

In view of (10.14) and the remark in [10, p. 166], one can choose a subsequence of the sequence  $\{(\widetilde{\nu_{k_{\ell}}})^*\}_{\ell \in \mathbb{N}}$  (denote it again by the same symbol) so that

$$U^{\hat{\nu}}_{\alpha}(x) = \lim_{\ell \to \infty} U^{(\widetilde{\nu_{k_{\ell}}})^*}_{\alpha}(x)$$
 n.e. in  $\mathbb{R}^n$ .

On account of (10.13), we thus have  $U^{\hat{\nu}}_{\alpha}(x) = 0$  n.e. in  $(\overline{D^c})^*$ , and therefore, by (10.15),

(10.16) 
$$\hat{\nu}^{-} = \beta^{\alpha}_{(\overline{D^{c}})^{*}} \hat{\nu}^{+} = \beta^{\alpha}_{(\overline{D^{c}})^{*}} \nu^{*}_{0}.$$

Using the fact that the Kelvin transformation is an involution and applying (10.9), (10.10) and (10.11) again, we conclude from (10.14), (10.15) and (10.16) that

$$\widetilde{\nu_{k_{\ell}}} \to \nu_0 - \beta_{D^c}^{\alpha} \nu_0 \quad (\text{as } \ell \to \infty) \quad \text{in } \mathcal{E}_{\alpha}(\mathbb{R}^n),$$

or equivalently, by the definition of  $\widetilde{\nu_{k_{\ell}}}$ ,

$$\lim_{\ell \to \infty} \left\| \left( \nu_{k_{\ell}} - \nu_0 \right) - \beta_{D^c}^{\alpha} \left( \nu_{k_{\ell}} - \nu_0 \right) \right\|_{\alpha} = 0.$$

Repeated application of (2.7) then proves relation (10.3) also in the case  $\overline{E} \cap \partial_{\mathbb{R}^n} D = \{x_0\}$ , where  $x_0 \neq \omega$ . This completes Step 2.

Since the sequence  $\{\nu_k\}_{k\in\mathbb{N}}$  is strongly fundamental,  $\nu_k \to \nu_0$  strongly by (10.3). It has thus been proved that  $\{\nu_k\}_{k\in\mathbb{N}}$  converges strongly to any of its vague cluster points. As the  $\alpha$ -Green kernel is strictly positive definite,  $\nu_0$  is the only vague cluster point of  $\{\nu_k\}_{k\in\mathbb{N}}$ , and so  $\nu_k \to \nu_0$  also vaguely.

#### 11. $\alpha$ -Green Equilibrium measure

**Theorem 11.1.** Let  $E \subset \mathbb{R}^n$  be closed in D. If  $\alpha < 2$ , require additionally  $\overline{E} \cap \partial_{\overline{\mathbb{R}^n}} D$  to consist of at most one point. If, moreover,  $C_{g_D^{\alpha}}(E) < \infty$ , then there exists an  $\alpha$ -Green interior equilibrium measure  $\gamma = \gamma_E$  on E, that is, a one possessing the properties  $\gamma \in \mathcal{E}_{g_D^{\alpha}}^+(E)$  and

(11.1) 
$$E_{g_D^{\alpha}}(\gamma) = \gamma(E) = C_{g_D^{\alpha}}(E),$$

(11.2) 
$$U_{g_D^{\alpha}}^{\gamma}(x) \ge 1$$
 n.e. in E,

(11.3) 
$$U_{a_{D}}^{\gamma}(x) \leq 1 \quad \text{for all} \ x \in S_{D}^{\gamma}.$$

This  $\gamma$  solves the problem of minimizing the energy  $E_{g_D^{\alpha}}(\nu)$  over the convex class  $\Gamma_E$  of all  $\nu \in \mathcal{E}_{g_D^{\alpha}}(D)$  such that  $U_{g_D^{\alpha}}^{\nu}(x) \ge 1$  n.e. in E, and hence it is unique.

*Proof.* This theorem needs to be proved only in the case  $\alpha < 2$ , since otherwise it is a special case of [10, Theorem 4.1] in view of the perfectness of the  $g_D^2$ -Green kernel. Also note that we can assume E to be noncompact in D, for if not, then the theorem follows from [10] (see Theorem 2.5 and Lemma 3.2.2 with t = 1 therein).

Consider an exhaustion of E by an increasing sequence of sets  $K_k \subset E$ ,  $k \in \mathbb{N}$ , compact in D, and let  $\gamma_k = \gamma_{K_k}$  be the  $\alpha$ -Green equilibrium measure on  $K_k$ . Then, by (11.1) with  $E = K_k$  and (2.8),

(11.4) 
$$\lim_{k \to \infty} \|\gamma_k\|_{g_D^{\alpha}}^2 = \lim_{k \to \infty} C_{g_D^{\alpha}}(K_k) = C_{g_D^{\alpha}}(E) < \infty.$$

Since  $\gamma_k \in \Gamma_{K_p}$  for all  $k \ge p$ , which is seen from the monotonicity of  $\{K_k\}_{k \in \mathbb{N}}$  and inequality (11.2) with  $E = K_k$ , Lemma 2.1 yields

$$\|\gamma_k - \gamma_p\|_{g_D^{\alpha}}^2 \leqslant \|\gamma_k\|_{g_D^{\alpha}}^2 - \|\gamma_p\|_{g_D^{\alpha}}^2 \quad \text{for all} \ k \ge p.$$

In consequence of the last two relations,  $\{\gamma_k\}_{k\in\mathbb{N}} \subset \mathcal{E}^+_{g^{\alpha}_D}(E)$  is strongly fundamental. In addition, by (11.1) with  $E = K_k$ ,

(11.5) 
$$\gamma_k(E) = C_{g_D^{\alpha}}(K_k) < C_{g_D^{\alpha}}(E) < \infty \quad \text{for all} \ k \in \mathbb{N},$$

so that all the assumptions of Theorem 10.1 for  $\{\gamma_k\}_{k\in\mathbb{N}}$  are satisfied. Hence, there exists the unique  $\gamma \in \mathcal{E}^+_{g^{\alpha}_{D}}(E)$  such that  $\gamma_k \to \gamma$  both strongly and vaguely.

On account of (11.4), we thus get

(11.6) 
$$\|\gamma\|_{g_D^{\alpha}}^2 = \lim_{k \to \infty} \|\gamma_k\|_{g_D^{\alpha}}^2 = C_{g_D^{\alpha}}(E)$$

According to [10] (see the remark on p. 166 therein), the strong convergence of  $\gamma_k$  to  $\gamma$  also yields that there exists a subsequence  $\gamma_{k_\ell} = \gamma_{K_{k_\ell}}$ ,  $\ell \in \mathbb{N}$ , of  $\{\gamma_k\}_{k \in \mathbb{N}}$  such that

$$\lim_{\ell \to \infty} \, U_{g_D^{\alpha}}^{\gamma_{k_\ell}}(x) = U_{g_D^{\alpha}}^{\gamma}(x) \quad \text{n.e. in} \ D$$

while by (11.2) for  $E = K_{k_{\ell}}$ ,

$$U_{g_D^{\alpha}}^{\gamma_{k_\ell}}(x) \ge 1$$
 n.e. in  $K_{k_\ell}$ .

Since the sets  $K_{k_{\ell}}$ ,  $\ell \in \mathbb{N}$ , increase and  $E = \bigcup_{\ell \in \mathbb{N}} K_{k_{\ell}}$ , the last two relations imply (11.2). Here we have used the fact that the  $\alpha$ -Green capacity of a countable union of Borel sets with zero  $\alpha$ -Green capacity is still zero; see [10].

Fix  $x \in S_D^{\gamma}$ . As  $\gamma_k \to \gamma$  vaguely, one can choose  $x_k \in S_D^{\gamma_k}$  so that  $x_k \to x$  as  $k \to \infty$ . Because of the fact (cf. [10, Lemma 2.2.1]) that  $U_{g_D^{\alpha}}^{\mu}(x)$  is lower semicontinuous on the product space  $D \times \mathfrak{M}^+(D)$ , where  $\mathfrak{M}^+(D)$  is equipped with the vague topology, we get

$$U_{g_D^{\alpha}}^{\gamma}(x) \leq \liminf_{k \to \infty} U_{g_D^{\alpha}}^{\gamma_k}(x_k).$$

Since, by (11.3) for  $E = K_k$ ,  $U_{g_D^{\gamma_k}}^{\gamma_k}(x_k) \leq 1$  for all  $k \in \mathbb{N}$ , inequality (11.3) follows.

In view of the vague convergence of  $\gamma_k$  to  $\gamma$ , we also have

$$\gamma(E) \leqslant \liminf_{k \to \infty} \gamma_k(E),$$

so that  $\gamma(E) \leq C_{g_D^{\alpha}}(E)$  by (11.5). When combined with (11.6), this shows that, in order to complete the proof of (11.1), it is left to establish the inequality  $\gamma(E) \geq C_{g_D^{\alpha}}(E)$ , but it follows at once by integrating (11.3) with respect to  $\gamma$ .

Finally, [10, Lemma 3.2.2] with t = 1 yields the very last assertion of the theorem.

**Remark 11.2.**  $\gamma_E$  coincides up to a constant factor with the solution  $\lambda_E$  of Problem 3.2 (for *E* in place of *F*) with  $\sigma = \infty$  and f = 0. See Corollaries 6.6 and 6.7 for a more detailed information about the properties of the  $\alpha$ -Green potential and the support of  $\lambda_E$ .

## 12. Proof of Theorem 5.2

In this section we follow methods developed in [21] (see Theorems 2.2 and 3.1 therein). Under the assumptions of Theorem 5.2, the following auxiliary result holds.

**Lemma 12.1.** For any  $\sigma \in \mathfrak{C}(F) \cup \{\infty\}$ , the metric space

$$\mathcal{E}_{g^{\alpha}_{D}}^{\sigma}(F,1) := \left\{ \mu \in \mathcal{E}_{g^{\alpha}_{D}}^{+}(F,1) : \ \mu \leqslant \sigma \right\}$$

is strongly complete. In more detail, any strong Cauchy sequence  $\{\mu_k\}_{k\in\mathbb{N}} \subset \mathcal{E}_{g_D^{\alpha}}^{\sigma}(F,1)$ converges both strongly and vaguely to the unique  $\mu_0 \in \mathcal{E}_{g_D^{\alpha}}^{\sigma}(F,1)$ .

*Proof.* Fix a strong Cauchy sequence  $\{\mu_k\}_{k\in\mathbb{N}} \subset \mathcal{E}_{g_D^{\alpha}}^{\sigma}(F,1)$ . According to Theorem 10.1, there exists the unique  $\mu_0 \in \mathcal{E}_{g_D^{\alpha}}^+(F)$  such that

 $\mu_k \rightarrow \mu_0$  strongly and vaguely.

Actually,  $\mu_0 \in \mathcal{E}_{g^{\alpha}_D}^{\sigma}(F)$ , since  $\mathcal{E}_{g^{\alpha}_D}^{\sigma}(F)$  is vaguely closed. Hence, it is left to show that (12.1)  $\mu_0(F) = 1.$  Assume F to be noncompact, for if not, then (12.1) is evident. Consider an exhaustion of F by an increasing sequence of sets  $K_m \subset F$ ,  $m \in \mathbb{N}$ , compact in D; then

$$1 = \lim_{k \to \infty} \mu_k(F) \ge \mu_0(F) = \lim_{m \to \infty} \mu_0(K_m) \ge \lim_{m \to \infty} \limsup_{k \to \infty} \mu_k(K_m)$$
$$= 1 - \lim_{m \to \infty} \liminf_{k \to \infty} \mu_k(F \setminus K_m).$$

Therefore, identity (12.1) will be established once we prove

(12.2) 
$$\lim_{m \to \infty} \liminf_{k \to \infty} \mu_k(F \setminus K_m) = 0.$$

Write  $K_m^* := C\ell_D(F \setminus K_m)$ . It is seen from Theorem 11.1 that, under the assumptions made, there exists the  $\alpha$ -Green equilibrium measure  $\gamma_m$  on  $K_m^*$ , and it solves the problem of minimizing  $E_{g_D^{\alpha}}(\nu)$  over the convex cone  $\Gamma_{K_m^*}$ . Since, by the monotonicity of  $K_m^*$ ,  $m \in \mathbb{N}$ , and relation (11.2) for  $E = K_m^*$ ,  $\gamma_m$  belongs to  $\Gamma_p$  for all  $p \ge m$ , Lemma 2.1 yields

$$\|\gamma_m - \gamma_p\|_{g_D^{\alpha}}^2 \leqslant \|\gamma_m\|_{g_D^{\alpha}}^2 - \|\gamma_p\|_{g_D^{\alpha}}^2 \quad \text{for all} \ p \ge m.$$

Furthermore, it is clear from (11.1) for  $E = K_m^*$  that the sequence  $\|\gamma_m\|_{g_D^{\alpha}}^2$ ,  $m \in \mathbb{N}$ , is bounded and nonincreasing, and hence it is fundamental in  $\mathbb{R}$ . The preceding inequality thus implies that  $\gamma_m$ ,  $m \in \mathbb{N}$ , is strongly fundamental in  $\mathcal{E}_{g_D^{\alpha}}^+(D)$ . Since it obviously converges vaguely to zero, zero is also its strong limit due to Theorem 10.1. Hence,

$$\lim_{m \to \infty} \|\gamma_m\|_{g_D^\alpha} = 0$$

Besides, by (11.2) for  $E = K_m^*$ ,

$$\mu_k(F \setminus K_m) \leqslant \mu_k(K_m^*) \leqslant \left\langle U_{g_D^{\alpha}}^{\gamma_m}, \mu_k \right\rangle \leqslant \|\gamma_m\|_{g_D^{\alpha}} \cdot \|\mu_k\|_{g_D^{\alpha}} \quad \text{for all} \ k, m \in \mathbb{N}.$$

As  $\|\mu_k\|_{g^{\alpha}_D}$ ,  $k \in \mathbb{N}$ , is bounded, combining the last two relations yields (12.2).

Now we are able to complete the proof of Theorem 5.2. In view of (3.7), one can choose  $\nu_k \in \mathcal{E}_{g_{D,f}^{\sigma}}^{\sigma}(F,1), k \in \mathbb{N}$ , so that

(12.3) 
$$\lim_{k \to \infty} G_{g_D^{\alpha}, f}(\nu_k) = G_{g_D^{\alpha}, f}^{\sigma}(F, 1) < \infty.$$

Based on the convexity of the class  $\mathcal{E}_{g_D^{\alpha},f}^{\sigma}(F,1)$  and the pre-Hilbert structure on  $\mathcal{E}_{g_D^{\alpha}}(D)$ , with the help of arguments similar to those in the proof of Lemma 4.1 we obtain

$$0 \leq \|\nu_k - \nu_p\|_{g_D^{\alpha}}^2 \leq -4G_{g_D^{\alpha},f}^{\sigma}(F,1) + 2G_{g_D^{\alpha},f}(\nu_k) + 2G_{g_D^{\alpha},f}(\nu_p) \quad \text{for all} \ k,p \in \mathbb{N}.$$

Substituting (12.3) into this relation implies that  $\{\nu_k\}_{k\in\mathbb{N}}$  is strongly fundamental in the metric space  $\mathcal{E}_{g_D}^{\sigma}(F,1)$ . By Lemma 12.1,  $\{\nu_k\}_{k\in\mathbb{N}}$  therefore converges both strongly and vaguely to the unique  $\nu_0 \in \mathcal{E}_{g_D}^{\sigma}(F,1)$ . On account of Lemma 4.4, we thus get

(12.4) 
$$G_{g_D^{\alpha},f}(\nu_0) \leqslant \lim_{k \to \infty} G_{g_D^{\alpha},f}(\nu_k) = G_{g_D^{\alpha},f}^{\sigma}(F,1) < \infty.$$

Hence,  $\nu_0 \in \mathcal{E}_{g_D^{\alpha}, f}^{\sigma}(F, 1)$  and, consequently,  $G_{g_D^{\alpha}, f}(\nu_0) \ge G_{g_D^{\alpha}, f}^{\sigma}(F, 1)$ . Combined with (12.4), this shows that  $\nu_0 =: \lambda_F^{\sigma}$  is the solution to Problem 3.2.

## 13. Proof of the assertions formulated in Section 6

13.1. **Proof of Theorem 6.1.** Fix  $\lambda \in \mathcal{E}_{g_D^{\alpha}, f}^{\xi}(F, 1)$ , and first assume that it solves Problem 3.2. Then inequality (6.2) holds for  $w_{\lambda} = L$ , where

$$L := \sup \left\{ q \in \mathbb{R} : \ W_{g^{\alpha}_{D},f}^{\lambda}(x) \geqslant q \quad (\xi - \lambda) \text{-a.e. in } F \right\}.$$

In turn, (6.2) with  $w_{\lambda} = L$  implies  $L < \infty$ , since  $W_{g_D^{\lambda}, f}^{\lambda}(x) < \infty$  holds n.e. in  $F_0$ , hence  $(\xi - \lambda)$ -a.e. in  $F_0$ , while  $(\xi - \lambda)(F_0) > 0$ . Also,  $L > -\infty$ , for f is bounded from below.

We proceed by establishing (6.3) for  $w_{\lambda} = L$ . Having denoted (cf. [8, 15])

$$F^{+}(w) := \left\{ x \in F : W^{\lambda}_{g^{\alpha}_{D}, f}(x) > w \right\} \text{ and } F^{-}(w) := \left\{ x \in F : W^{\lambda}_{g^{\alpha}_{D}, f}(x) < w \right\},$$

where  $w \in \mathbb{R}$  is arbitrary, we assume on the contrary that (6.3) for  $w_{\lambda} = L$  does not hold. In view of the lower semicontinuity of  $W_{g_{D}^{\alpha},f}^{\lambda}$  on F, then one can choose  $w_{1} \in (L,\infty)$  so that  $\lambda(F^{+}(w_{1})) > 0$ . At the same time, as  $w_{1} > L$ , relation (6.2) with  $w_{\lambda} = L$  yields  $(\xi - \lambda)(F^{-}(w_{1})) > 0$ . Therefore, there exist compact sets  $K_{1} \subset F^{+}(w_{1})$  and  $K_{2} \subset F^{-}(w_{1})$  such that

$$0 < \lambda(K_1) < (\xi - \lambda)(K_2).$$

Write  $\tau := (\xi - \lambda)|_{K_2}$ ; then  $\tau \in \mathcal{E}^+_{g^{\alpha}_D}(K_2)$ . Since  $\langle W^{\lambda}_{g^{\alpha}_D,f}, \tau \rangle \leq w_1 \tau(K_2) < \infty$ , we get  $\langle f, \tau \rangle < \infty$ . Define

$$\theta := \lambda - \lambda \big|_{K_1} + c\tau$$
, where  $c := \lambda(K_1) \big/ \tau(K_2) \in (0, 1)$ .

A straightforward verification shows that  $\theta(F) = 1$  and  $\theta \leq \xi$ , and so  $\theta \in \mathcal{E}_{g_D^{\alpha}, f}^{\xi}(F, 1)$ . On the other hand,

$$\begin{split} \left\langle W_{g_{D}^{\lambda},f}^{\lambda},\theta-\lambda\right\rangle &= \left\langle W_{g_{D}^{\lambda},f}^{\lambda}-w_{1},\theta-\lambda\right\rangle \\ &= -\left\langle W_{g_{D}^{\lambda},f}^{\lambda}-w_{1},\lambda\right|_{K_{1}}\right\rangle + c\left\langle W_{g_{D}^{\lambda},f}^{\lambda}-w_{1},\tau\right\rangle < 0, \end{split}$$

which is impossible in view of Lemma 4.3. This proves the necessary part of the theorem.

Next, let  $\lambda$  satisfy both (6.2) and (6.3) for some  $w_{\lambda} \in \mathbb{R}$ . Then  $\lambda(F^+(w_{\lambda})) = 0$  and  $(\xi - \lambda)(F^-(w_{\lambda})) = 0$ . For any  $\nu \in \mathcal{E}_{g_{\Omega,f}^{\xi}}^{\xi}(F, 1)$ , we therefore obtain

$$\langle W_{g_{D}^{\alpha},f}^{\lambda},\nu-\lambda\rangle = \langle W_{g_{D}^{\alpha},f}^{\lambda}-w_{\lambda},\nu-\lambda\rangle = \langle W_{g_{D}^{\alpha},f}^{\lambda}-w_{\lambda},\nu\big|_{F^{+}(w_{\lambda})}\rangle + \langle W_{g_{D}^{\alpha},f}^{\lambda}-w_{\lambda},(\nu-\xi)\big|_{F^{-}(w_{\lambda})}\rangle \ge 0.$$

Application of Lemma 4.3 shows that, indeed,  $\lambda$  is the solution to Problem 3.2.

13.2. **Proof of Corollary 6.2.** Under the conditions of the corollary,  $W_{g_D^{\alpha},f}^{\lambda}(x) > 0$  in D; hence, the number  $w_{\lambda}$  from Theorem 6.1 satisfies relation

(13.1) 
$$w_{\lambda} \in (0, \infty).$$

Furthermore, in the notation  $\Psi(x) := U_{\alpha}^{\lambda+\chi}(x) - U_{\alpha}^{\beta_{D^c}^{\alpha}(\lambda+\chi)}(x), x \in \mathbb{R}^n$ , inequality (6.2) can be rewritten in the form

$$\Psi(x) \ge w_{\lambda} > 0$$
 ( $\xi - \lambda$ )-a.e. in F.

We can certainly assume that there is  $y_0 \in \partial D \cap S_{\mathbb{R}^n}^{\xi-\lambda}$ , for if not, then the corollary is obvious. Since for every  $\varepsilon > 0$  it holds  $(\xi - \lambda) (B(y_0, \varepsilon)) > 0$ , one can choose  $x_{\varepsilon} \in F \cap B(y_0, \varepsilon)$  so that  $\Psi(x_{\varepsilon}) \ge w_{\lambda} > 0$ . Therefore,

$$\limsup_{x \to y_0, \ x \in D} \Psi(x) \ge w_{\lambda} > 0$$

On the other hand,  $\Psi(x) \ge 0$  for all  $x \in \mathbb{R}^n$  and  $\Psi(x) = 0$  n.e. in  $D^c$ ; hence,

$$\lim_{x \to y_0, \ x \in D^c} \Psi(x) = 0$$

Consequently,  $\Psi$  is discontinuous on  $\partial D \cap S_{\mathbb{R}^n}^{\xi-\lambda}$ , and Lusin's type theorem for the  $\alpha$ -Riesz potentials (see [14, Theorem 3.6]) establishes the corollary.

13.3. **Proof of Corollary 6.3.** Fix  $\lambda_0 \in \mathcal{E}_{g_D^{\alpha}, f}^{\xi}(F)$ . We first assume that it solves Problem 3.2, and let  $w'_{\lambda_0} \in (0, \infty)$  be the number from (6.2) and (6.3) for f = 0 (see also (13.1)). Then (6.3) can be rewritten in the form

$$U_{\alpha}^{\lambda_0}(x) \leqslant w'_{\lambda_0} + U_{\alpha}^{\beta_{D^c}^{\alpha}\lambda_0}(x) \quad \text{for all } x \in S_D^{\lambda_0}.$$

Note that the right-hand side of this relation is  $\alpha$ -superharmonic in  $\mathbb{R}^n$ . Applying [14, Theorems 1.27, 1.29], we see that, actually,

(13.2) 
$$U_{\alpha}^{\lambda_0}(x) \leq w_{\lambda_0}' + U_{\alpha}^{\beta_{\alpha}^{D_c}\lambda_0}(x) \quad \text{for all } x \in \mathbb{R}^n.$$

which gives (6.6). Combining (6.6) with (6.2) for  $\lambda = \lambda_0$  and  $w_{\lambda} = w'_{\lambda_0}$  results in (6.5).

Assuming now that both (6.5) and (6.6) hold for some  $w'_{\lambda_0} \in (0, \infty)$ , we conclude from Theorem 6.1 that  $\lambda_0$  solves Problem 3.2, as was to be proved.

Finally, let  $\alpha < 2$  and let  $\lambda_0$  solve Problem 3.2. To establish (6.7), assume on the contrary that there exists  $x_0 \in F$  such that  $x_0 \notin S_D^{\lambda_0}$ . Then one can choose r > 0 so that

$$\overline{B}(x_0,r) := \{ x \in \mathbb{R}^n : |x - x_0| \leqslant r \} \subset D \quad \text{and} \quad \overline{B}(x_0,r) \cap S_D^{\lambda_0} = \emptyset.$$

It follows that  $(\xi - \lambda_0) (B(x_0, r) \cap F) > 0$ . Therefore, by (6.5),

(13.3)  $U_{\alpha}^{\lambda_0}(x_1) = w_{\lambda_0}' + U_{\alpha}^{\beta_{D^c}^{\alpha}\lambda_0}(x_1) \quad \text{for some} \ x_1 \in B(x_0, r) \cap F.$ 

As  $U^{\lambda_0}_{\alpha}(\cdot)$  is  $\alpha$ -harmonic in  $B(x_0, r)$  and continuous on  $\overline{B}(x_0, r)$ , while  $w'_{\lambda_0} + U^{\beta^{\alpha}_{Dc}\lambda_0}_{\alpha}(\cdot)$  is  $\alpha$ -superharmonic in  $\mathbb{R}^n$ , we conclude from (13.2) and (13.3) with the help of [14, Theorem 1.28] that

$$U_{\alpha}^{\lambda_0}(x) = w'_{\lambda_0} + U_{\alpha}^{\beta_{D^c}^{\alpha}\lambda_0}(x) \quad m_n$$
-a.e. in  $\mathbb{R}^n$ .

This implies  $w'_{\lambda_0} = 0$ , for  $U^{\beta_{D^c}\lambda_0}_{\alpha}(x) = U^{\lambda_0}_{\alpha}(x)$  holds n.e. in  $D^c$ , hence, also  $m_n$ -a.e. in  $D^c$ . A contradiction.

13.4. **Proof of Theorem 6.4.** This theorem is a very particular case of [22, Theorems 7.1, 7.2, 7.3] (see also [19] and Theorems 1, 2 and Proposition 1 therein).  $\Box$ 

13.5. **Proof of Corollary 6.5.** Since (a) follows directly from Theorem 5.3, assume the conditions of assertion (b) to hold. In the same manner as in Section 13.2, then one can see that the number  $w_f$  from Theorem 6.4 is strictly positive. Hence, by (6.9),

$$\Psi(x) \ge w_f > 0$$
 n.e. in  $F$ ,

where  $\Psi$  has been defined in Section 13.2. We can certainly assume that there is  $y_0 \in \partial D \cap C\ell_{\mathbb{R}^n}\check{F}$ , for if not, then (b) is obvious. For every  $\varepsilon > 0$ , it holds  $C_{\alpha}(B(y_0,\varepsilon)\cap\check{F}) > 0$ , and therefore one can choose  $x_{\varepsilon} \in B(y_0,\varepsilon)\cap\check{F}$  so that  $\Psi(x_{\varepsilon}) \ge w_f > 0$ . This yields

$$\lim_{x \to y_0, \ x \in D} \Psi(x) \ge w_f > 0.$$

Likewise as in Section 13.2, we can thus see that  $\Psi$  is discontinuous on  $\partial D \cap C\ell_{\mathbb{R}^n} \check{F}$ , and Lusin's type theorem for the  $\alpha$ -Riesz potentials establishes the corollary.  $\Box$ 

13.6. **Proof of Corollary 6.6.** Let  $w := w_f$  be the number from Theorem 6.4 for f = 0; then w > 0. Proof of the statement that  $\lambda_F \in \mathcal{E}_{g_D^{+}}^+(F, 1)$  solves Problem 3.2 if and only if both (6.11) and (6.12) hold is based on Theorem 6.4 and runs in a way similar to that in the proof of Corollary 6.3. In particular, as an application of [14, Theorems 1.27, 1.29], we conclude from (6.10) with f = 0 that, if  $\lambda_F$  solves Problem 3.2, then

(13.4) 
$$U_{\alpha}^{\lambda_F}(x) \leqslant w + U_{\alpha}^{\beta_{Dc}^{\alpha}\lambda_F}(x) \quad \text{for all } x \in \mathbb{R}^n.$$

The very last statement of the corollary is obtained from (6.11) with the help of standard arguments (see, e.g., [14, pp. 137–138]), based on the strict positive definiteness of the  $\alpha$ -Green kernel.

13.7. Proof of Corollary 6.7. Having first assumed  $\alpha < 2$ , we start by showing that

(13.5) 
$$U_{g_D^{\alpha}}^{\lambda_F}(x) < w \quad \text{for all} \ x \in D \setminus S_D^{\lambda_F}.$$

Suppose to the contrary that (13.5) is not satisfied for some  $x_0 \in D \setminus S_D^{\lambda_F}$ . Then  $U_{g_D^{\alpha}}^{\lambda_F}(x_0) = w$  in accordance with (6.12), or equivalently

(13.6) 
$$U_{\alpha}^{\lambda_F}(x_0) = w + U_{\alpha}^{\beta_{Dc}^{\alpha}\lambda_F}(x_0).$$

Choose  $\varepsilon > 0$  so that  $\overline{B}(x_0, \varepsilon) \subset D \setminus S_D^{\lambda_F}$ . Since then  $U_{\alpha}^{\lambda_F}(\cdot)$  is  $\alpha$ -harmonic in  $B(x_0, \varepsilon)$ and continuous on  $\overline{B}(x_0, \varepsilon)$ , while  $w + U_{\alpha}^{\beta_{D^c}^{\alpha}\lambda_F}(\cdot)$  is  $\alpha$ -superharmonic in  $\mathbb{R}^n$ , we conclude from (13.4) and (13.6) with the help of [14, Theorem 1.28] that

$$U_{\alpha}^{\lambda_F}(x) = w + U_{\alpha}^{\beta_{D^c}^{\alpha}\lambda_F}(x)$$
 m<sub>n</sub>-a.e. in  $\mathbb{R}^n$ .

As  $U_{\alpha}^{\beta_{D^c}^{\alpha}\lambda_F}(x) = U_{\alpha}^{\lambda_F}(x)$  n.e. in  $D^c$ , we thus get w = 0. A contradiction.

We next proceed by proving the former identity in (6.14). Let, on the contrary, there exist  $x_1 \in \breve{F}$  such that  $x_1 \notin S_D^{\lambda_F}$ , and let  $V \subset D \setminus S_D^{\lambda_F}$  be an open neighborhood of  $x_1$ . Then, by (13.5),  $U_{g_D^{\alpha}}^{\lambda_F}(x) < w$  for all  $x \in V$ . On the other hand, since  $V \cap F$  has nonzero capacity,  $U_{g_D^{\alpha}}^{\lambda_F}(x_2) = w$  for some  $x_2 \in V$  by (6.11). The contradiction obtained shows that, indeed,  $S_D^{\lambda_F} = \breve{F}$ . Substituting this identity into (13.5) establishes (6.13) for  $\alpha < 2$ .

In the rest of the proof,  $\alpha = 2$ . To verify (6.13), assume, on the contrary, that it does not hold for some  $x_3$  in the domain  $D_0 := D \setminus \breve{F}$ . According to (6.12), then  $U_{g_D^2}^{\lambda_F}(x_3) = w$ , which in view of the harmonicity of  $U_{g_D^2}^{\lambda_F}$  in  $D_0$  implies, by the maximum principle, that

$$U_{g_D^2}^{\lambda_F}(x) = w \quad \text{for all} \ x \in D_0.$$

Thus,

$$\lim_{x \to z, \ x \in D_0} U_{g_D^2}^{\lambda_F}(x) = w > 0 \quad \text{for all} \ z \in \partial D_0.$$

Since  $C_{\alpha}(\partial D \cap \partial D_0) > 0$  in consequence of Corollary 6.5, (b), Lusin's type theorem for the Newtonian potentials shows that the preceding relation is impossible.

In view of (6.11), [14, Theorem 1.13] yields  $\lambda_F |_{\operatorname{Int} F} = 0$ , and so  $S_D^{\lambda_F} \subset \partial_D \check{F}$ . Thus, if we prove the converse inclusion, the latter identity in (6.14) follows. Assume, on the contrary, it not to hold; then one can choose a point  $y \in \partial_D \check{F}$  and a neighborhood  $V_1 \subset D$  of y so that  $V_1 \cap S_D^{\lambda_F} = \emptyset$ . As  $V_1 \cap F$  has nonzero capacity, we see from (6.11) that there exists  $y_1 \in V_1$  such that  $U_{g_D^2}^{\lambda_F}(y_1) = w$ . Taking (6.12) into account and applying the maximum principle to the harmonic in  $V_1$  function  $U_{g_D^2}^{\lambda_F}$ , we thus have  $U_{g_D^2}^{\lambda_F}(x) = w$  for all  $x \in V_1$ . This contradicts (6.13), because  $V_1 \cap D_0 \neq \emptyset$ .

13.8. Proof of Theorem 6.8. Since  $U_{g^{\alpha}_{D}}^{\xi}(x)$  is continuous, so is  $U_{g^{\alpha}_{D}}^{\lambda_{0}}(x)$ . Indeed,

$$U_{g_{D}^{\alpha}}^{\lambda_{0}}(x) = U_{g_{D}^{\alpha}}^{\xi}(x) - U_{g_{D}^{\alpha}}^{\xi-\lambda_{0}}(x),$$

which implies that  $U_{g_D^{\lambda_0}}^{\lambda_0}(x)$  is both lower semicontinuous and upper semicontinuous. Next, since  $\lambda_0$  solves the non-weighted Problem 3.2 with the constraint  $\xi$ , both (6.5) and (6.6)

are fulfilled. As  $U_{g_D^{\alpha}}^{\lambda_0}(\cdot)$  is continuous, equality in (6.5) holds in fact everywhere on  $S_D^{\xi-\lambda_0}$ . This allows us to rewrite (6.5) and (6.6) respectively as

$$\begin{split} U_{g_D^{\beta}}^{\xi-\lambda_0}(x) - U_{g_D^{\beta}}^{\xi}(x) &= -w'_{\lambda_0} \quad \text{on} \quad S_D^{\xi-\lambda_0} \\ U_{g_D^{\beta}}^{\xi-\lambda_0}(x) - U_{g_D^{\beta}}^{\xi}(x) &\ge -w'_{\lambda_0} \quad \text{on} \ D, \end{split}$$

or equivalently, in the notations accepted in Section 6.3,

$$\begin{split} W^{\theta}_{g^{\alpha}_{D},f}(x) &= -qw'_{\lambda_{0}} \quad \text{on } S^{\theta}_{D}, \\ W^{\theta}_{g^{\alpha}_{D},f}(x) &\geq -qw'_{\lambda_{0}} \quad \text{on } D. \end{split}$$

This establishes relations (6.15) and (6.16). Since  $\theta \in \mathcal{E}_{g^{\alpha}_{D},f}^{q\xi}(F,1) \subset \mathcal{E}_{g^{\alpha}_{D},f}^{+}(F,1)$ , application of Theorems 6.1 and 6.4 therefore completes the proof.

## 14. EXAMPLES

In this section, n = 3 and  $x = (x_1, x_2, x_3)$  is a point in  $\mathbb{R}^3$ . In the following Examples 14.1–14.3, consider  $0 < \alpha \leq 2$ , D := B(0, 1) and  $A_2 := D^c$ ; then  $A_2$  is not  $\alpha$ -thin at  $\omega$ .

**Example 14.1.** Write  $E := \{x \in B(0,1) : 0 \leq x_1 < 1, x_2 = x_3 = 0\}$ . Since  $C_{\alpha}(E) = 0$ , Lemma 2.3 yields  $C_{g_D^{\alpha}}(E) = 0$ . Consequently, there exists a neighborhood F of E, closed in D, with  $0 < C_{g_D^{\alpha}}(F) < \infty$ . We can certainly assume that  $\partial D \cap C\ell_{\mathbb{R}^3}F = \{(1,0,0)\}$ . Consider an external field f such that  $f(x) < \infty$  n.e. in F unless Case II holds. Application of Lemma 4.7, Theorems 5.2 and 5.4 then shows that, in both Cases I and II, Problem 3.1 is (uniquely) solvable for every  $\sigma \in \mathcal{A}(F) \cup \{\infty\}$ .

**Example 14.2.** Let F = D. Define  $\xi := m_3|_F$ ; then  $\xi \in \mathfrak{C}_0(F)$  and has finite  $\alpha$ -Riesz energy and thus it is admissible (see Definition 4.6). Consider an external field f such that Case I holds and  $f(x) < \infty m_3|_F$ -a.e. Hence, by Lemma 4.7, assumptions (3.6) and (3.7) hold and so we can apply Theorems 5.1 and 5.4 to conclude that Problem 3.1 is solvable; that is, no short-circuit between the conductors F and  $D^c$  occurs, though they touch each other over the whole sphere S(0, 1).

**Example 14.3.** Let  $F = S(x_0, 1/2) \cap D$ , where  $x_0 = (1/2, 0, 0)$ . Consider an external field f such that Case I holds and  $f(x) < \infty m_2|_F$ -a.e. We further assume  $1 < \alpha \leq 2$ . Define  $\xi := m_2|_F$ ; then (since  $\alpha > 1$ )  $\xi$  has finite  $\alpha$ -Riesz energy and so, as in the previous example, we can apply Theorems 5.1 and 5.4 to obtain the solvability of Problem 3.1.

**Example 14.4.** Let  $\alpha = 2$ ,  $D = \{x \in \mathbb{R}^3 : x_1 > 0\}$  and  $F = \{x \in D : x_1 = 1\}$ ; then  $A_2 = D^c$  is not 2-thin at  $\omega$ , while  $C_{g_D^2}(F) = \infty$ , for  $C_2(F) = \infty$  by [14, Chapter II, Section 3, n° 14]. Let, in addition, Case II with  $\zeta \ge 0$  hold. Then, by Theorems 5.3 and 5.4, Problem 3.1 is nonsolvable for every  $\sigma \in \mathcal{A}(F) \cup \{\infty\}$  such that  $\sigma \ge \xi_0$ , where  $\xi_0 \in \mathfrak{C}(F) \setminus \mathfrak{C}_0(F)$  is properly chosen. Thus, for these  $\sigma$ , a short-circuit between F and  $D^c$ occurs at  $\omega$ . To construct a constraint which would not allow such a short-circuit, consider  $K_k, k \in \mathbb{N}$ , where  $K_k := \{x \in F : (k-1)^2 \leq x_2^2 + x_3^2 \leq k^2\}$  for all  $k \ge 2$  and  $K_1 := \{x \in F : x_2^2 + x_3^2 \leq 1\}$ , and write

$$\xi := \sum_{k \in \mathbb{N}} \frac{m_2|_{K_k}}{k^3}.$$

Then  $\xi$  is bounded and admissible, and so we can again use Theorems 5.1 and 5.4 to see that Problem 3.1 for this constraint  $\xi$  is solvable.




FIGURE 1. The condenser for Example 14.5.

**Example 14.5.** Let  $\alpha = 2$ , Case II with  $\zeta \ge 0$  hold, and let F and D be defined by

$$F := \left\{ x \in \mathbb{R}^3 : \quad 2 \leqslant x_1 < \infty, \quad x_2^2 + x_3^2 = \rho_1^2(x_1), \quad \text{where } \rho_1(x_1) = \exp(-x_1) \right\},$$
  
$$D := \left\{ x \in \mathbb{R}^3 : \quad 1 < x_1 < \infty, \quad x_2^2 + x_3^2 < \rho_2^2(x_1), \quad \text{where } \rho_2(x_1) = x_1^{-1} \right\}.$$

Then  $A_2 = D^c$  is not 2-thin at  $\omega$ , while  $C_{g_D^2}(F) = \infty$ , for  $C_2(F) = \infty$  by [16]. Hence, by Theorems 5.3 and 5.4, Problem 3.1 is nonsolvable for every  $\sigma \in \mathcal{A}(F) \cup \{\infty\}$  such that  $\sigma \geq \xi_0$ , where  $\xi_0 \in \mathfrak{C}(F) \setminus \mathfrak{C}_0(F)$  is properly chosen. However, Problem 3.1 with  $\xi := m_2|_F$ is already solvable, which is seen from Theorems 5.1 and 5.4.

## References

- Beckermann, B., Gryson, A.: Extremal rational functions on symmetric discrete sets and superlinear convergence of the ADI method. Constr. Approx. 32, 393–428 (2010)
- [2] Bourbaki, N.: Elements of Mathematics, Integration, Chapters 1–6. Springer, Berlin (2004)
- [3] Brelot, M.: Eléments de la théorie classique du potential, Les cours Sorbonne, Paris (1961)
- Brelot, M.: On Topologies and Boundaries in Potential Theory. Lectures Notes in Math., vol. 175. Springer, Berlin (1971)
- [5] Cartan, H.: Théorie du potentiel Newtonien: énergie, capacité, suites de potentiels. Bull. Soc. Math. Fr. 73, 74–106 (1945)
- [6] Deny, J.: Les potentiels d'énergie finite. Acta Math. 82, 107–183 (1950)
- [7] Deny, J.: Sur la définition de l'énergie en théorie du potentiel. Ann. Inst. Fourier Grenoble 2, 83–99 (1950)
- [8] Dragnev, P.D., Saff, E.B.: Constrained energy problems with applications to orthogonal polynomials of a discrete variable. J. Anal. Math. 72, 223-259 (1997)

- [9] Edwards, R.: Cartan's balayage theory for hyperbolic Riemann surfaces. Ann. Inst. Fourier 8, 263–272 (1958)
- [10] Fuglede, B.: On the theory of potentials in locally compact spaces. Acta Math. 103, 139–215 (1960)
- [11] Fuglede, B.: Asymptotic paths for subharmonic functions and polygonal connectedness of fine domains. Lectures Notes in Math., vol. 814, 97–115. Springer, Berlin (1980)
- [12] Frostman, O.: Sur les fonctions surharmoniques d'ordre fractionnaire, Ark. Mat., Astr. Fys. 26A, N16 (1939)
- [13] Hayman, W.K.: Subharmonic Functions. Vol. 2. Academic Press, London (1989)
- [14] Landkof, N.S.: Foundations of Modern Potential Theory. Springer, Berlin (1972)
- [15] Rakhmanov, E.A.: Equilibrium measure and the distribution of zeros of extremal polynomials of a discrete variable. Sb. Math. 187, 1213–1228 (1996)
- [16] Zorii, N.: An extremal problem of the minimum of energy for space condensers. Ukr. Math. J. 38, 365–369 (1986)
- [17] Zorii, N.: A problem of minimum energy for space condensers and Riesz kernels. Ukr. Math. J. 41, 29–36 (1989)
- [18] Zorii, N.: A noncompact variational problem in Riesz potential theory. I; II. Ukr. Math. J. 47, 1541– 1553 (1995); 48, 671–682 (1996)
- [19] Zorii, N.: Equilibrium potentials with external fields. Ukr. Math. J. 55, 1423–1444 (2003)
- [20] Zorii, N.: Equilibrium problems for potentials with external fields. Ukr. Math. J. 55, 1588–1618 (2003)
- [21] Zorii, N.: Constrained energy problems with external fields. Complex Anal. Oper. Theory 5, 775–785 (2011), DOI:10.1007/s11785-010-0070-9
- [22] Zorii, N.: Equilibrium problems for infinite dimensional vector potentials with external fields. Potential Anal. 38, 397–432 (2013), DOI:10.1007/s11118-012-9279-8

DEPARTMENT OF MATHEMATICAL SCIENCES INDIANA-PURDUE UNIVERSITY FORT WAYNE, IN 46805, USA

#### *E-mail address*: dragnevp@ipfw.edu

CENTER FOR CONSTRUCTIVE APPROXIMATION, DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240, USA

#### *E-mail address*: doug.hardin@vanderbilt.edu

#### *E-mail address*: edward.b.saff@vanderbilt.edu

DEPARTMENT OF MATHEMATICAL PHYSICS, INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, TERESHCHENKIVSKA 3, 01601, KYIV-4, UKRAINE

E-mail address: natalia.zorii@gmail.com