

A NOTE ON STANDARD MODULES AND VOGAN L -PACKETS

VOLKER HEIERMANN

ABSTRACT. Let F be a non-Archimedean local field of characteristic 0, let G be the group of F -rational points of a connected reductive group defined over F and let G' be the group of F -rational points of its quasi-split inner form. Given standard modules $I(\tau, \nu)$ and $I(\tau', \nu')$ for G and G' respectively with τ' a generic tempered representation, such that the Harish-Chandra μ -function of a representation in the supercuspidal support of τ agrees with the one of a generic essentially square-integral representation in some Jacquet module of τ' (after a suitable identification of the underlying spaces under which $\nu = \nu'$), we show that $I(\tau, \nu)$ is irreducible whenever $I(\tau', \nu')$ is.

The conditions are satisfied if the Langlands quotients $J(\tau, \nu)$ and $J(\tau', \nu')$ of respectively $I(\tau, \nu)$ and $I(\tau', \nu')$ lie in the same Vogan L -packet (whenever this Vogan L -packet is defined), proving that, for any Vogan L -packet, all the standard modules with Langlands quotient in a given Vogan L -packet are irreducible, if and only if this Vogan L -packet contains a generic representation.

This result for generic Vogan L -packets was proven for quasi-split orthogonal and symplectic groups by Mœglin-Waldspurger and used in their proof of the general case of the local Gan-Gross-Prasad conjectures for these groups.

Let F be a non-Archimedean local field of characteristic 0. Denote by G the group of F -rational points of a connected reductive group defined over F .

Fix a minimal F -parabolic subgroup $P_0 = M_0U_0$ of G . A F -parabolic subgroup $P = MU$ will be called standard, if it contains P_0 and M contains M_0 and semi-standard, if only M contains M_0 . One says then respectively that M is a standard or semi-standard F -Levi subgroup. If M is any semi-standard F -Levi subgroup of G , a standard F -parabolic subgroup of M will be a F -parabolic subgroup of M which contains $P_0 \cap M$ and analog for the F -Levi subgroups of M_0 .

Let $P = MU$ be a standard F -parabolic subgroup of G and T_M the maximal split torus in the center of M . We will write a_M^* for the dual of the real Lie-algebra a_M of T_M , $a_{M, \mathbb{C}}^*$ for its complexification and a_M^{*+} for the positive Weyl chamber in a_M^* defined with respect to P . The set of reduced roots for the action of T_M in the Lie algebra of G will be denoted $\Sigma(T_M)$, the subset of positive roots $\Sigma(P)$, and

The author has benefitted from a grant of Agence Nationale de la Recherche with reference ANR-13-BS01-0012 FERPLAY.

the corresponding sets of coroots respectively $\Sigma^\vee(T_M)$ and $\Sigma^\vee(P)$. They will be identified with subsets of a_M^* and of a_M respectively.

Following [W], one defines a map $H_M : M \rightarrow a_M$, such that, for every F -rational character $\chi \in a_M^*$ of M , $|\chi(m)|_F = q^{-\langle \chi, H_M(m) \rangle}$. If π is a smooth representation of M and $\nu \in a_{M, \mathbb{C}}^*$, we denote by π_ν the smooth representation of M defined by $\pi_\nu(m) = q^{-\langle \nu, H_M(m) \rangle} \pi(m)$. (Remark that, although the sign in the definition of H_M has been changed compared to the one due to Harish-Chandra, the meaning of π_ν is unchanged.) The symbol i_P^G will denote the functor of parabolic induction normalized such that it sends unitary representations to unitary representations, G acting on its space by right translations.

Let τ be an irreducible tempered representation of M and $\nu \in a_M^{*+}$. Then the induced representation $i_P^G \tau_\nu$ is called a standard module and will be denoted $I(P, \tau, \nu)$ in the sequel. It has a unique irreducible quotient $J(P, \tau, \nu)$, the so-called Langlands quotient. By the Langlands classification theorem, every irreducible smooth representation π of G appears as a Langlands quotient of a unique standard module $I(P, \tau, \nu)$. One says then that (P, τ, ν) are Langlands data for π and writes $(P_\pi, \tau_\pi, \nu_\pi)$ if one wants to underline the link to π .

Let G' be the group of F -rational points of a quasi-split connected reductive group defined over F . Fix a F -Borel subgroup $B' = T'V'$, which defines the notions of standard F -parabolic, standard F -Levi, and semi-standard F -Levi subgroup of G' and of its semi-standard F -Levi subgroups. Suppose that there is a standard F -parabolic subgroup $P'_0 = M'_0 U'_0$ of G' so that $(a_{M'_0}^*, \Sigma(T_{M'_0}), a_{M'_0}, \Sigma^\vee(T_{M'_0}))$ is a root datum and that the based root data $(a_{M'_0}^*, \Sigma(T_{M'_0}), a_{M'_0}, \Sigma^\vee(T_{M'_0}), \Delta(P'_0))$ and $(a_{M_0}^*, \Sigma(T_{M_0}), a_{M_0}, \Sigma^\vee(T_{M_0}), \Delta(P_0))$ are isomorphic. The isomorphism is a pair (φ, φ^\vee) , where φ is an isomorphism between $a_{M_0}^*$ and $a_{M'_0}^*$ sending $\Sigma(T_{M_0})$ onto $\Sigma(T_{M'_0})$ and $\Delta(P_0)$ onto $\Delta(P'_0)$, and φ^\vee is the inverse dual map that sends $\Sigma^\vee(T_{M_0})$ onto $\Sigma^\vee(T_{M'_0})$. (One may think of G' as an inner form of G (and it is then unique), but this is not required below.)

In particular, the isomorphism φ of based root data defines a bijection between standard F -parabolic subgroups $P' = M'U'$ of G' , $P' \supseteq P'_0$, and standard F -parabolic subgroups $P = MU$ of G with an isomorphism $a_M^* \rightarrow a_{M'}^*$, also denoted φ in the sequel, that sends the set of positive roots $\Sigma(P)$ onto $\Sigma(P')$. We will say that the standard parabolic subgroups $P = MU$ and $P' = M'U'$ correspond to each other by φ . Remark that the inverse dual map $\varphi^\vee : a_M \rightarrow a_{M'}$ sends then $\Sigma^\vee(P)$ onto $\Sigma^\vee(P')$.

Denote by W' the Weyl group of G' defined with respect to the maximal F -split torus T' in G' and by $w_0^{G'}$ the longest element in W' . After changing the splitting in V' , for any generic representation π' of G' , one can always find a non degenerate character ψ' of V' , which is compatible with $w_0^{G'}$, such that π' is ψ' -generic [Sh, section 3]. For any semi-standard Levi-subgroup M' of G' , we will still denote by ψ' the restriction of ψ' to $M' \cap V'$. It is compatible with $w_0^{M'}$. If we write in the

sequel that a representation of a F -semi-standard Levi subgroup of G' is ψ' -generic, then we always mean that ψ' is a non degenerate character of V' with the above properties.

Fix for the rest of this introduction a standard F -parabolic subgroup $P = MU$ of G and a standard F -parabolic subgroup $P' = M'U'$ of G' that correspond to each other by the above isomorphism of based root data φ .

Let τ be an irreducible tempered representation of M , σ a unitary irreducible supercuspidal representation of a standard F -Levi subgroup M_1 of M and $\lambda_\tau \in a_{M_1}^{M^*}$, so that σ_{λ_τ} lies in the supercuspidal support of τ . Denote by $P'_1 = M'_1U'_1$ the standard F -parabolic subgroup of G' that corresponds to P_1 by the isomorphism of root data φ . Assume that there is a ψ' -generic discrete series representation σ' of M'_1 , such that Harish-Chandra's μ -functions μ^M and $\mu^{M'}$ become equal under φ on the inertial orbits of σ and σ' , i.e. $\mu^M(\sigma_\lambda) = \mu^{M'}(\sigma'_{\varphi(\lambda)})$ for all $\lambda \in a_{M_1}^*$.

Our first result is then (cf. lemma **2.5**) that $i_{P'_1 \cap M'}^{M'} \sigma'_{\varphi(\lambda_\tau)}$ has a ψ' -generic tempered sub-quotient τ' .

The aim of this note is further to show that under these conditions the standard module $I(P, \tau, \nu)$ is irreducible, whenever the standard module $I(P', \tau', \varphi(\nu))$ is irreducible (cf. theorem **2.7**). We show in the annex **A.** that the above conditions are satisfied if G' is a quasi-split classical group and π and π' are two members of a generic Vogan L -packet, π' being the generic member, and (P, τ, ν) and (P', τ', ν') are Langlands data for them. (For a general quasi-split group G' , one may add this property to the conjectural properties of Vogan L -packets.) It follows, using the standard modules conjecture for G' proven in its whole generality in [HM] (subsequent to [HO]) that $I(P, \tau, \nu)$ is always irreducible when the Langlands quotient $J(P, \tau, \nu)$ lies in a ψ' -generic Vogan L -packet (cf. theorem **3.2**).

The above result for generic Vogan L -packets was proven for quasi-split orthogonal and symplectic groups by Mœglin-Waldspurger in [MW, 2.12] and used in their proof of the general case of the local Gan-Gross-Prasad conjectures for these groups. (The restrictions that we make in **3.1** on the literature hold also for their work.)

As the Vogan L -packets for unitary groups (with their conjectured properties) are known following the work of Mok [Mk] and Kaletha-Minguez-Shin-White [KMSW], our results apply in particular to standard modules corresponding to generic Vogan L -packets of unitary groups. This should be useful if one wants to prove the general case of the local Gan-Gross-Prasad conjectures for unitary groups, the conjectures for tempered representations having been established by Beuzart-Plessis [BP].

The method of proof of the present note is a generalization of the proof of the standard modules conjecture in [HM], based on the basic principle in [Mu], following the observation that everything is determined by the geometric combinatorial configuration of the parameters with respect to Harish-Chandra's μ -function. Remark that the result in [HM] depended on the tempered L -function conjecture which has been proven in its whole generality in [HO]. (The fact that the theory

of L -functions is only established in characteristic 0 is the reason why we had to make this assumption on the characteristic of F . Thanks to the work of L. Lomeli and others, this obstacle may be lifted soon.)

1. Recall that G' denotes the set of F -rational points of a quasi-split connected reductive group defined over F and that one has fixed a Borel subgroup $B' = T'V'$ of G' and a non-degenerate character ψ' of V' .

1.1 Theorem: (Rodier) *Let $P' = M'U'$ be a standard F -parabolic subgroup of G' and τ' an irreducible smooth representation of M' .*

The representation $i_{P'}^{G'}\tau'$ has an irreducible ψ' -generic subquotient, if and only if τ' is ψ' -generic. This irreducible ψ' -generic subquotient is then unique.

1.2 Let M' be a semi-standard Levi subgroup of G' and τ' an irreducible generic tempered representation of M' . In [Sh3], F. Shahidi attached to each irreducible component r'_i of the adjoint action of the L -group ${}^L M'$ of M' on $\text{Lie}({}^L U')$ an L -function $L(s, \tau', r'_i)$, which is a meromorphic function in the complex variable s .

Theorem: [HO, 5.1] *The meromorphic function $L(s, \tau', r'_i)$ is regular for $\Re(s) > 0$.*

1.3 The following statement was proven in [HM, 2.2] modulo theorem **1.2** (which was proven in its whole generality later).

Proposition: *Let σ' be a ψ' -generic supercuspidal representation of the standard Levi factor of a standard parabolic subgroup P' of G' .*

Suppose that $i_{P'}^{G'}\sigma'$ has an irreducible subquotient that is a tempered representation. Then, the ψ' -generic sub-quotient of $i_{P'}^{G'}\sigma'$ is a tempered representation.

1.4 Corollary: *Let σ' be a ψ' -generic representation of the standard Levi factor of a standard parabolic subgroup P' of G' .*

Suppose that $i_{P'}^{G'}\sigma'$ has an irreducible sub-quotient that is a tempered representation. Then, the generic subquotient of $i_{P'}^{G'}\sigma'$ is a tempered representation.

Proof: Fix a standard parabolic subgroup $P'_1 = M'_1U'_1$ of G' with $M'_1 \subseteq M'$ and an irreducible supercuspidal representation σ'_1 of M'_1 , such that σ' is a subrepresentation of $i_{P'_1 \cap M'}^{M'}\sigma'_1$. The representation σ'_1 is ψ' -generic by **1.1**. An irreducible sub-quotient of $i_{P'}^{G'}\sigma'$ is necessarily a sub-quotient of $i_{P'_1}^{G'}\sigma'_1$. In particular, $i_{P'_1}^{G'}\sigma'_1$ has a tempered sub-quotient. By **1.3**, the ψ' -generic sub-quotient π is tempered, too. By uniqueness of the ψ' -generic sub-quotient, π is also the ψ' -generic subquotient of $i_{P'}^{G'}\sigma'$. This proves the corollary. \square

2. Recall that G denotes the F -rational points of a connected reductive group defined over F and that one has fixed a minimal F -parabolic subgroup $P_0 = M_0U_0$ of G and a F -parabolic subgroup $P'_0 = M'_0U'_0$ of the quasi-split group G' so that the corresponding based root spaces are isomorphic by an isomorphism φ (see introduction).

2.1 The Harish-Chandra μ -function is the main ingredient of the Plancherel density for a p -adic reductive group H [W]. It assigns to every square-integrable representation of a Levi subgroup a complex number and can be analytically extended to a meromorphic function on the space of essentially square-integrable representations of Levi subgroups. What is important here, is the following: if $Q = NV$ is a parabolic subgroup of a connected reductive group H over F and σ an irreducible unitary supercuspidal representation of N , then the Harish-Chandra's μ -function μ^H corresponding to H defines a meromorphic function $a_{N,\mathbb{C}}^* \rightarrow \mathbb{C}$, $\lambda \mapsto \mu^H(\sigma_\lambda)$, on the \mathbb{C} -vector space $a_{N,\mathbb{C}}^*$ which can be written in the form

$$(2.1) \quad \mu^H(\sigma_\lambda) = f(\lambda) \prod_{\alpha \in \Sigma(Q)} \frac{(1 - q^{\langle \alpha^\vee, \lambda \rangle})(1 - q^{-\langle \alpha^\vee, \lambda \rangle})}{(1 - q^{\epsilon_{\alpha^\vee} + \langle \alpha^\vee, \lambda \rangle})(1 - q^{\epsilon_{\alpha^\vee} - \langle \alpha^\vee, \lambda \rangle})} \quad (\text{cf. [H1, Si]}),$$

where f is a meromorphic function without poles and zeroes on a_N^* and the ϵ_{α^\vee} are non negative rational numbers with $\epsilon_{\alpha^\vee} = \epsilon_{\alpha'^\vee}$ if α and α' are conjugate.

2.2 Definition: A point σ_{λ_0} , $\lambda_0 \in a_N^*$, is called a residue point for μ^H , if

$$|\{\alpha \in \Sigma(Q) | \langle \alpha^\vee, \lambda_0 \rangle = \pm \epsilon_{\alpha^\vee}\}| - 2|\{\alpha \in \Sigma(Q) | \langle \alpha^\vee, \lambda_0 \rangle = 0\}| = \dim(a_N^*/a_G^*).$$

2.3 Theorem: [H1] *The induced representation $i_Q^H \sigma_{\lambda_0}$ has a sub-quotient which is square integrable, if and only if σ_{λ_0} is a residue point of μ^H and the projection of λ_0 on a_H^* is 0.*

2.4 Lemma: *Let σ be a unitary irreducible supercuspidal representation of the standard F -Levi subgroup M of a standard parabolic subgroup P of G . Denote by $P' = M'U'$ the F -standard parabolic subgroup of G' that corresponds to P by the isomorphism of basic root data φ .*

Assume that there is a ψ' -generic discrete series representation σ' of M' such that the Harish-Chandra μ -functions of σ and σ' are directly proportional under φ , i.e. that there is some complex number $c \in \mathbb{C}$ such that $\mu^G(\sigma_\lambda) = c\mu^{G'}(\sigma'_{\varphi(\lambda)})$ for all $\lambda \in a_{M,\mathbb{C}}^$.*

Let $\lambda_0 \in a_M^$. If σ_{λ_0} is a residue point for μ^G , then $\sigma'_{\varphi(\lambda_0)}$ is a residue point for $\mu^{G'}$ and vice versa. In addition, the projection of λ_0 onto a_G^* is zero if and only if the projection of $\varphi(\lambda_0)$ is zero on $a_{G'}^*$.*

Remark: Although the notion of a residue point is only defined for μ -functions with respect to supercuspidal representations, the statement makes sense because the μ -function with respect to $\sigma'_{\lambda'}$, $\lambda' \in a_{M'}^*$, behaves by our assumptions like the one of a supercuspidal representation.

Proof: By the formula (2.1), there are a meromorphic function f without poles and zeroes on a_N^* and non negative rational numbers ϵ_{α^\vee} , $\epsilon_{\alpha^\vee} = \epsilon_{\alpha'^\vee}$ if α and α' are conjugate, such that, for $\lambda' \in a_{M'}^*$,

$$\mu^G(\sigma_{\varphi^{-1}(\lambda')}) = f(\varphi^{-1}(\lambda')) \prod_{\alpha \in \Sigma(P)} \frac{(1 - q^{\langle \alpha^\vee, \varphi^{-1}(\lambda') \rangle})(1 - q^{-\langle \alpha^\vee, \varphi^{-1}(\lambda') \rangle})}{(1 - q^{\epsilon_{\alpha^\vee} + \langle \alpha^\vee, \varphi^{-1}(\lambda') \rangle})(1 - q^{\epsilon_{\alpha^\vee} - \langle \alpha^\vee, \varphi^{-1}(\lambda') \rangle})}.$$

Applying the dual of φ^{-1} , this becomes

$$f(\varphi^{-1}(\lambda')) \prod_{\alpha \in \Sigma(P)} \frac{(1 - q^{\langle \varphi^\vee(\alpha^\vee), \lambda' \rangle})(1 - q^{-\langle \varphi^\vee(\alpha^\vee), \lambda' \rangle})}{(1 - q^{\epsilon_{\alpha^\vee} + \langle \varphi^\vee(\alpha^\vee), \lambda' \rangle})(1 - q^{\epsilon_{\alpha^\vee} - \langle \varphi^\vee(\alpha^\vee), \lambda' \rangle})}.$$

By our assumptions, this equals $\mu^{G'}(\sigma'_{\lambda'})$ up to a constant factor. Comparing this expression with the expression for $\mu^{G'}(\sigma'_{\lambda'})$ given by (2.1), one deduces that $\epsilon_{\alpha^\vee} = \epsilon_{\varphi^\vee(\alpha^\vee)}$. Looking at the zero and polar hyperplanes in both cases, it becomes clear that $\sigma'_{\lambda'}$ is a residue point of $\mu^{G'}$ if and only if $\sigma_{\varphi^{-1}(\lambda')}$ is one for μ^G .

The projection of an element $\lambda \in a_M^*$ onto a_G^* is 0, if and only if λ is a linear combination of elements in $\Sigma(T_M)$. As φ induces a bijection between $\Sigma(T_M)$ and $\Sigma(T_{M'})$, this is equivalent to say that $\varphi(\lambda)$ is a linear combination of elements in $\Sigma(T_{M'})$. Consequently, the projection of λ onto a_G^* is 0, if and only if the projection of $\varphi(\lambda)$ onto $a_{G'}^*$ is 0. \square

2.5 Lemma: (with the notations and assumptions of **2.4**) *Let $\lambda \in a_M^*$ and assume that $i_P^G \sigma_\lambda$ has a tempered sub-quotient. Then, $i_{P'}^{G'} \sigma'_{\varphi(\lambda)}$ has a ψ' -generic tempered sub-quotient.*

Proof: Let τ be a tempered sub-quotient of $i_P^G \sigma_\lambda$. Then, it is a sub-representation of a representation induced by a discrete series representation τ_2 of some standard F -Levi subgroup M_2 of G and σ_λ is conjugated to an element in the supercuspidal support of τ_2 . After conjugating σ_λ and M by a Weyl group element (which does not effect the Jordan-Hölder sequence for $i_P^G \sigma_\lambda$ neither for $i_{P'}^{G'} \sigma'_{\varphi(\lambda)}$ - if σ_λ is conjugated by w , $\sigma'_{\varphi(\lambda)}$ has to be conjugated by $w' := \varphi \circ w \circ \varphi^{-1}$), we can assume that $M \subseteq M_2$ and consequently σ_λ lies in the supercuspidal support of τ_2 . By **2.3**, this implies that σ_λ is a residue point of the Harish-Chandra μ -function μ^{M_2} and that the projection of $\Re(\lambda)$ onto $a_{M_2}^*$ is trivial.

Denote by M'_2 the standard F -Levi subgroup of G' that corresponds to M_2 by the isomorphism of root data φ . By **2.4**, $\sigma'_{\varphi(\lambda)}$ is a residue point of $\mu^{M'_2}$ and the projection of $\mathfrak{R}(\lambda)$ on $a_{M'_2}^*$ is trivial.

It remains to show that $i_{P' \cap M'_2}^{M'_2} \sigma'_{\varphi(\lambda)}$ has a discrete series sub-quotient. If σ' is not a supercuspidal representation, one cannot immediately apply theorem **2.3**. Let $P'_1 = M'_1 U'_1$ be a standard parabolic subgroup of G' contained in P' , σ'_1 a ψ' -generic unitary supercuspidal representation of M'_1 and $\lambda'_2 \in a_{M'_1}^{M'_1}$ such that σ'_{1, λ'_2} lies in the supercuspidal support of σ' . As σ' is a square-integrable representation, σ_{1, λ'_2} is a residue point of $\mu^{M'}$. One remarks that there is a complex number c , such that $c\mu^{M'_2}(\sigma'_{\lambda'}) = (\mu^{M'_2}/\mu^{M'}) (\sigma'_{1, \lambda'_2 + \lambda'})$ as functions in $\lambda' \in a_{M'}^*$ (cf. lemma **A.3** in the attachment). As $\sigma'_{\varphi(\lambda)}$ is a residue point for $\mu^{M'_2}$, one sees that $\sigma'_{1, \lambda'_2 + \varphi(\lambda)}$ is also a residue point for $\mu^{M'_2}$ and the projection on $a_{M'_2}^*$ is zero. From **2.3**, it follows that $i_{P'_1 \cap M'_2}^{M'_2} \sigma'_{1, \lambda'_2 + \varphi(\lambda)}$ has a square-integrable sub-quotient and consequently $i_{P'_1}^{G'} \sigma'_{1, \lambda'_2 + \varphi(\lambda)}$ has a tempered sub-quotient. As $\sigma'_{1, \lambda'_2 + \varphi(\lambda)}$ is ψ' -generic, $i_{P'_1}^{G'} \sigma'_{1, \lambda'_2 + \varphi(\lambda)}$ has by **1.4** a ψ' -generic tempered sub-quotient, which is by its uniqueness the ψ' -generic sub-quotient of $i_{P'}^{G'} \sigma'_{\varphi(\lambda)}$. This proves the lemma. \square

2.6 Définition: Let $I(P, \tau, \nu)$ be a standard module for G and $I(P', \tau', \nu')$ a ψ' -generic standard module for G' . We will say that $I(P, \tau, \nu)$ is related to $I(P', \tau', \nu')$ (by φ), if $\nu' = \varphi(\nu)$, the two F -standard parabolic subgroups $P = MU$ and $P' = M'U'$ correspond to each other by φ , and there are in addition standard parabolic subgroups $P_1 = M_1 U_1$ and $P'_1 = M'_1 U'_1$ of G and G' respectively, a unitary supercuspidal representation σ of M_1 , a ψ' -generic discrete series representation σ' of M'_1 such that P_1 and P'_1 correspond to each other by φ , $\mu^M(\sigma_\lambda)$ and $\mu^{M'}(\sigma'_{\varphi(\lambda)})$ are directly proportional as functions of $\lambda \in a_{M_1}^*$ (i.e. there is some complex number c such that $\mu^M(\sigma_\lambda) = c\mu^{M'}(\sigma'_{\varphi(\lambda)})$), and there is $\lambda_1 \in a_{M_1}^*$ such that τ is a subquotient of $i_{P_1 \cap M}^M \sigma_{\lambda_1}$ and τ' a subquotient of $i_{P'_1 \cap M'}^{M'} \sigma'_{\varphi(\lambda_1)}$.

Remark: 1) In other words, the supercuspidal representation σ_{λ_1} of M_1 is an element of the supercuspidal support of τ and σ' is a ψ' -generic discrete series representation of the standard Levi subgroup M'_1 of G' that corresponds to M'_1 and with μ -function directly proportional to the one of σ via the identification φ .

2) There is a distinguished class of inner forms of G' , called pure inner forms, defined by Vogan [V]. It is expected that $I(P, \tau, \nu)$ is related to $I(P', \tau', \nu')$, if G' is a pure inner form of G and $J(P, \tau, \nu)$ and $J(P', \tau', \nu')$ lie in the same Vogan L -packet (defined in [V]). If $G = G'$, this means that these representations lie in the same (Langlands) L -packet. In addition, if G' is a pure inner form of G , then every standard module $I(P, \tau, \nu)$ for G should be related to a ψ' -generic standard

module for G' . All this is known now for quasi-split classical groups (orthogonal, symplectic or unitary) by the stabilization of the Arthur-Selberg trace formula and work of different authors on it, see **3.1** for more details.

By a generalization of the Jacquet-Langlands correspondence, similar properties should hold for a general inner form G of G' .

2.7 Theorem: *Let $I(P', \tau', \nu')$ be a ψ' -generic standard module for G' . If $I(P', \tau', \nu')$ is irreducible, then every standard module $I(P, \tau, \nu)$ of G that is related to $I(P', \tau', \nu')$ is also irreducible. In particular, if $J(P', \tau', \nu')$ is ψ' -generic, each standard module $I(P, \tau, \nu)$ of G that is related to $I(P', \tau', \nu')$ is irreducible.*

Proof: Fix data $P_1 = M_1U_1, \sigma_1, \lambda_1$ and $P'_1 = M'_1U'_1, \sigma'_1$ associated to $I(P, \tau, \nu)$ and $I(P', \tau', \nu')$ as in definition **2.6**. In particular, σ_{λ_1} lies in the supercuspidal support of τ and $\sigma'_{\varphi(\lambda'_1)}$ in the supercuspidal support of τ' .

If $I(P, \tau, \nu)$ is reducible, then there is an irreducible subquotient π_2 of $I(P, \tau, \nu)$ that is not isomorphic to $J(P, \tau, \nu)$. By [BW, chapter XI, 2.13], there is a Langlands data (P_2, τ_2, ν_2) with $\nu_2 < \nu$ in a_T^* (the precise definition of this order does not matter here, only that it is stable by the isomorphism of based root data φ), so that $\pi_2 = J(P_2, \tau_2, \nu_2)$. Remark that $\sigma_{\lambda_1 + \nu}$ lies in the supercuspidal support of π , π_2 and τ_ν , and that it is at least conjugate to an element in the supercuspidal support of τ_{2, ν_2} . This element in the supercuspidal support of τ_{2, ν_2} can be written in the form $(w\sigma)_{w\lambda_1 + w\nu}$ where w is some Weyl group element which acts by conjugation so that $wM_1 := wM_1w^{-1}$ is a standard F -Levi subgroup. Put $w' = \varphi \circ w \circ \varphi^{-1}$. Then, the standard F -Levi subgroup of G' that corresponds to wM_1 by the isomorphism of root data G is $w'M'_1$.

As σ and σ' have directly proportional μ -functions by φ , the representation $i_{M' \cap P'_1}^{M'} \sigma'_{\varphi(\lambda_1)}$ has by lemma **2.5** a ψ' -generic tempered sub-quotient τ' . As the standard F -Levi subgroups wM_1 and $w'M'_1$ correspond to each other by the isomorphism of root data φ and Plancherel measures are invariant by conjugation, $w\sigma$ and $w'\sigma'$ have also directly proportional μ -functions by φ . Denote by $P'_2 = M'_2U'_2$ the standard F -parabolic subgroup of G' that corresponds to $P_2 = M_2U_2$ and by P'_1 the standard F -parabolic subgroup of G' with Levi subgroup M'_1 . It follows from lemma **2.5** that the induced representation $i_{wP'_1 \cap M'_2}^{M'_2} \sigma'_{\varphi(w\lambda_1 + w\nu - \nu_2)}$ has a ψ' -generic tempered sub-quotient τ'_2 . Consequently, the ψ' -generic sub-quotient π^{gen} of $I(P', \tau', \varphi(\nu))$ is also a ψ' -generic sub-quotient of $I(P'_2, \tau'_2, \varphi(\nu_2))$. As $\nu_2 < \nu$ implies $\varphi(\nu_2) < \varphi(\nu)$, this proves that $\nu_{\pi^{gen}} < \varphi(\nu)$, showing that π^{gen} cannot be the Langlands quotient of $I(P', \tau', \varphi(\nu))$. This implies that $I(P', \tau', \varphi(\nu))$ is reducible.

The last sentence follows from the standard modules conjecture for G' . \square

Remark: The opposite direction of the statement in the theorem is wrong. There are many examples of standard modules coming from the same L -packet, some of them being irreducible and others being reducible (see for example [T, 5.1 (vi)]).

3. We assume now that G' is the set of F -rational points of the quasi-split inner form of G and that G is a pure inner form of G' in the sense of Vogan [V]. There exists then a natural isomorphism of based root data as mentioned in the introduction that we will still denote φ .

3.1 The notion of Vogan L -packets is defined in [V]. Remark first that G' and G have as inner forms of each other the same L -group, denoted here \mathcal{G} . Two representations π and π' of G and G' respectively lie in the same Vogan L -packet if they correspond to the same, in general hypothetical, Langlands parameter $\eta : W_F \times SL_2(\mathbb{C}) \rightarrow \mathcal{G}$. (For quasi-split classical groups, Vogan L -packets are known thanks to the stabilization of the trace formula in [A, Mk, KMSW] and its application [M1, M2], see also [H4] for some summary, only the analogue of [A] for pure inner forms of orthogonal groups and of [M2] for pure inner forms of quasi-split unitary groups remains to be published.)

By the expected properties of the local Langlands correspondence (see for example [M1, M2] for the known case of quasi-split classical groups and [H2, H3] for the general philosophy), this implies in particular that, if (P, τ, ν) and (P', τ', ν') are Langlands data for π and π' , then the standard parabolic subgroups P and P' correspond to each other, the tempered representations τ and τ' lie in the same Vogan L -packet, τ' will be ψ' -generic if τ is ψ -generic, and equality $\nu' = \varphi(\nu)$ holds.

If G' is a quasi-split classical group, the following result is proven in the annex (cf. theorem **A.1**):

Theorem: *Suppose that G' is a quasi-split classical group. If σ and σ' are discrete series representations of standard Levi subgroups M and M' of G and G' respectively that correspond to each other, then their Harish-Chandra μ -functions are directly proportional, i.e. there exists a complex number c such that*

$$\mu(\sigma_\lambda) = c\mu(\sigma'_{\varphi(\lambda)})$$

as functions in $\lambda \in a_{M, \mathbb{C}}^*$.

For a general quasi-split group G' , one may put this theorem as a conjecture along the existence of Vogan L -packets. Remark that, for Langlands L -packets, this conjecture appeared in [Sh, 9.3]. In general, some evidence can be seen from the formal degree conjecture for Vogan L -packets by Hiraga-Ichino-Ikeda [HII]. More philosophically, one can say that μ -functions are essentially defined by certain L -functions attached to them and which should not differ in an L -packet (see [GT] for the case of $GS\mathfrak{p}_4$, where a more precise result is given).

3.2 Theorem: *Assume either that G' is a quasi-split classical group or that the conclusions of theorem 3.1 and the expected properties of Vogan L -packets hold for G' .*

Let Π be the Vogan L -packet of an irreducible smooth representation of G' . Fix an F -parabolic subgroup $P' = M'U'$ of G' , an irreducible ψ' -generic tempered representation τ' of M' and an element ν' in $a_{M'}^{+}$ so that the Langlands quotient $J(P', \tau', \nu')$ lies in Π .*

The following are equivalent:

- (i) the standard module $I(P', \tau', \nu')$ is irreducible;*
- (ii) for every π in Π and Langlands data (P, τ, ν) for π , the standard module $I(P, \tau, \nu)$ is irreducible.*
- (iii) the Langlands quotient $J(P', \tau', \nu')$ is ψ' -generic.*
- (iv) the Vogan L -packet Π contains a generic representation.*

Proof: The existence of the ψ' -generic tempered representation τ' of a standard F -Levi subgroup M' comes from the conjecture that tempered L -packets are generic (one can then always find a ψ' -generic representation [Sh, 3.]). This conjecture is known for quasi-split groups thanks to the above mentioned works on the trace formula. For general G' , they take part of our assumptions on the Vogan L -packets.

As the conclusions of theorem 3.1 hold by assumption, it follows from the discussion in 3.1 and the properties of the local Langlands correspondence (known for quasi-split groups) that the standard module $I(P, \tau, \nu)$ is related to $I(P', \tau', \nu')$ in the sense of definition 2.6 so that one can conclude from theorem 2.7. \square

3.3 One can reformulate things in the following way:

Theorem: *(with the assumptions of 3.2) Denote by \mathcal{T} the Vogan L -packet of a ψ' -generic tempered representation τ' of the standard F -Levi subgroup of a standard F -parabolic subgroup P' of G' . The following are equivalent:*

- (i) the standard module $I(P', \tau', \nu')$ is irreducible;*
- (ii) for every inner form G of G' and Langlands data (P, τ, ν) with τ in \mathcal{T} and ν corresponding to ν' by an appropriate isomorphism of based root data for G and G' , the standard module $I(P, \tau, \nu)$ is irreducible.*
- (iii) the Langlands quotient $J(P', \tau', \nu')$ is ψ' -generic.*

Remark: Recall that there exist many examples of standard modules which are irreducible, while the corresponding ψ' -generic standard module is reducible (cf. for example [T, 5.1 (vi)]).

A. Annex

The aim of this annex is to give a proof of theorem 3.1:

A.1 Theorem: *Suppose that G' is a quasi-split classical group. If τ' is an irreducible discrete series representation of a Levi subgroup of M' and τ is an element of the Vogan L -packet of τ' , then the μ -functions associated to τ and τ' become directly proportional after identification of the real Lie algebras of the split centers of the corresponding Levi subgroups.*

The proof of this theorem will occupy the rest of this section. It will be done in several steps.

A.2 That the μ -function is constant on L -packets of discrete series representations of a standard Levi subgroup of a quasi-split classical group is in [Co, 7.8]. (The proof holds for every group where L -packets of discrete series representations are known and a linear combination of their characters gives a stable distribution - this is the case for quasi-split orthogonal and symplectic groups thanks to the work of J. Arthur [A] and for quasi-split unitary groups thanks to the work of C.-P. Mok [Mk].)

A.3 Suppose now that τ is a discrete series representation of a standard Levi subgroup M of G . Denote by σ a unitary supercuspidal representation of a standard Levi subgroup M_1 of M such that, for some $\lambda_1 \in a_{M_1}^{M*}$, σ_{λ_1} is contained in the supercuspidal support of τ (or, in other words, τ is a subquotient of $i_{P_1 \cap M}^M \sigma_{\lambda_1}$, where P_1 is the standard parabolic subgroup with Levi factor M_1).

The following lemma follows immediately from the fact that Harish-Chandra's μ -function is defined, up to a constant, as the reciprocal of the scalar operator associated to the composition of the standard intertwining operators with respect to opposite parabolic subgroups [W, V.2] and the product formula [W, V.2.1]:

Lemma: *As functions in $\lambda \in a_{M, \mathbb{C}}^*$, $\mu(\tau_\lambda)$ and $(\mu/\mu^M)(\sigma_{\lambda_1+\lambda})$ are directly proportional, i.e. $\mu(\tau) \sim (\mu/\mu^M)(\sigma_{\lambda_1+\cdot})$.*

A.4 Suppose first that there is a supercuspidal representation σ' of the corresponding Levi subgroup M'_1 of G' in the Vogan L -packet of σ . Then, as the μ -function of a supercuspidal representation is determined, up to a constant, by the reducibility points of the representation, and, by the results of Mœglin [M1, M2] (and of Bernstein-Zelevinsky for GL_n) these reducibility points can be read off from the Langlands parameter, one sees that the μ -functions of σ and of σ' are directly proportional. This implies that, with P'_1 equal to the standard parabolic subgroup of G' with Levi factor M'_1 , $i_{M' \cap P'_1}^{M'} \sigma'_{\varphi(\lambda_1)}$ has a discrete series subquotient τ' , as σ_{λ_1} is a residue point for μ^G (cf. **2.3**). By the Mœglin-Tadic classification for classical groups and its links to the Langlands correspondence [M1, M2], one sees that τ and τ' have same Langlands parameters, i.e. that they lie in the same

Vogan L -packet. One concludes by the equality

$$\mu(\tau_\lambda) \sim (\mu/\mu^M)(\sigma_{\lambda+\lambda_1}) \sim (\mu/\mu^{M'}) (\sigma'_{\varphi(\lambda)+\varphi(\lambda_1)}) \sim \mu(\tau'_{\varphi(\lambda)}),$$

the first and the last equality following from lemma **A.3**.

A.5 In general, however there may not exist a supercuspidal representation of M'_1 in the Vogan L -packet of σ . This happens precisely (see the end of this subsection), if the generic member in the Vogan L -packet is induced from a parabolic subgroup included in the Siegel parabolic subgroup (the one whose standard Levi-factor is isomorphic to a general linear group). To deal with this exceptional case, we will slightly modify the Langlands parameter (and the group G') to have the existence of a supercuspidal representation with good properties and relate this to M' .

Before doing that, let us go a little bit in the details of the Langlands correspondence for quasi-split classical groups, which is known thanks to the work of C. Mœglin [M1, M2] on the stablized trace formula for these groups (summarized in [H4, 1.1 and C.3] with the restriction on the literature mentioned in **3.1**). Denote by \mathcal{M}_1 the common L -group of M_1 and M'_1 and by $\eta_1 : W_F \times SL_2(\mathbb{C}) \rightarrow \mathcal{M}_1$ the Langlands parameter corresponding to the Vogan L -packet defined by σ . Remark that \mathcal{M}_1 is a complex reductive group that is a product of general linear groups with a group \mathcal{H} of the same type as the L -group of G . This means that η_1 is a direct sum of irreducible representations of W_F and a Langlands parameter $\eta_{1,0} : W_F \times SL_2(\mathbb{C}) \rightarrow \mathcal{H}$. There is a canonical inclusion ι of \mathcal{H} into a general linear group. The representation $\iota \circ \eta_{1,0}$ decomposes into irreducible representations of $W_F \times SL_2(\mathbb{C})$.

By the restriction of a Langlands parameter to ΔW_F , in notations $\eta_{1|\Delta W_F}$ or $(\iota \circ \eta_1)|_{\Delta W_F}$, we will mean in the sequel the composition with the map $\Delta : W_F \rightarrow W_F \times SL_2(\mathbb{C})$, $\gamma \mapsto (\gamma, \text{diag}(|\gamma|_F^{1/2}, |\gamma|_F^{-1/2}))$, while $\eta_{1|W_F}$ or $(\iota \circ \eta_1)|_{W_F}$ will denote the restriction to the first component.

By the results of Mœglin (see [H4, 1.1, C.3]), it may happen that there is no supercuspidal representation of M'_1 in the Vogan L -packet of σ (by assumption σ is a supercuspidal representation of M_1 , but there may be none on the quasi-split inner form of M_1 inside this L -packet). In this case, all irreducible components of $(\iota \circ \eta_{1,0})|_{W_F}$ have even multiplicities. It can be easily deduced from the above mentioned result of Mœglin that this is equivalent to the property that the supercuspidal support of the ψ' -generic member of the Vogan L -packet comes from a standard Levi subgroup included in the Siegel parabolic subgroup of G' . In fact, the Langlands parameter of the supercuspidal support of the ψ' -generic member of the Vogan L -packet of σ takes the same values as $\eta_{1|\Delta W_F}$ and its target space is the smallest Levi subgroup \mathcal{M}_{gen} of \mathcal{M} , containing the image of $\eta_{1|\Delta W_F}$.

A.6 To circumvent the problem of not having a supercuspidal representation of M'_1 in the Vogan L -packet of σ , we will choose an irreducible representation ρ of W_F , which factors through a group of the same type as the L -group of G' and which is not a factor of $(\iota \circ \eta_1)|_{W_F}$. (As there are infinitely many, this is always possible.) Replacing \mathcal{H} by a suitable group \mathcal{H}'' of bigger rank, but of the same type as \mathcal{H} , one sees that there is a Langlands parameter $\eta''_{1,0} : W_F \times SL_2(\mathbb{C}) \rightarrow \mathcal{H}''$ such that $\iota \circ \eta''_{1,0} = (\iota \circ \eta_{1,0}) \oplus \rho$. Denote by M''_1 the complex group obtained from M_1 , by replacing the factor \mathcal{H} by \mathcal{H}'' , and by G'' a quasi-split classical group of the same type as G' such that M''_1 is the L -group of a Levi subgroup of G'' . Denote by η''_1 the Langlands parameter $W_F \times SL_2(\mathbb{C}) \rightarrow M''_1$ obtained from η_1 by replacing $\eta_{1,0}$ by $\eta''_{1,0}$.

Then η''_1 defines an L -packet of discrete series representations of M'' that contains a supercuspidal representation σ'' . As the non semi-simple factors of M_1 and M''_1 are the same, there is a canonical isomorphism $\varphi'' : a^*_{M_1, \mathbb{C}} \rightarrow a^*_{M''_1, \mathbb{C}}$, as for corresponding Levi subgroups of pure inner forms. Denote by M'' the standard Levi subgroup of G'' that corresponds to M by this identification. As ρ is not a direct summand of $(\iota \circ \eta_1)|_{W_F}$, the reducibility points of σ and σ'' via the isomorphism φ'' are the same by the results of Mœglin [M1, M2]. One concludes as in the above case $M'' = M'$ (cf. **A.4**) that, with P''_1 equal to the standard parabolic subgroup with Levi factor M''_1 , the representation $i_{P''_1 \cap M''}^{M''}(\sigma''_{\varphi''(\lambda_1)})$ has a discrete series subquotient τ'' which has, via φ'' , a μ -function directly proportional to the one of τ . As the μ -function is invariant on (Langlands) L -packets of quasi-split classical groups (cf. **A.2**), it follows that the μ -function of τ'' equals the one of the ψ' -generic member τ''_{gen} in its L -packet.

A.7 Denote by η the Langlands parameter of τ . Using the relation between the Mœglin-Tadic classification of discrete series representations and the local Langlands correspondence [M1, M2], one sees that the common Langlands parameter η'' of τ'' and τ''_{gen} satisfies $\iota \circ \eta'' = (\iota \circ \eta) \oplus \rho$ and that ρ is not a summand of $(\iota \circ \eta)|_{W_F}$. Denote by τ_{gen} the generic member of the Vogan L -packet of τ , which is necessarily a representation of M' . The values of $\eta|_{\Delta W_F}$ (resp. $\eta''|_{\Delta W_F}$) are equal to the values of the Langlands parameter of a representation in the supercuspidal support of τ_{gen} (resp. τ''_{gen}), and both differ by the summand ρ .

Again (as in **A.6**, **A.4**): as the μ -functions for supercuspidal representations are determined, up to a constant, by the reducibility points of the representations and these points can be read off by the results of Mœglin and Bernstein-Zelevinsky from the Langlands parameter, while ρ is not a direct summand of $(\iota \circ \eta)|_{W_F}$, the two generic supercuspidal representations in the supercuspidal support of respectively τ_{gen} and τ''_{gen} have directly proportional μ -functions via the identification $\varphi'' \circ \varphi^{-1}$ of $a^*_{M'}$ and $a^*_{M''}$. One concludes from this as in **A.4** by the lemma **A.3** that the μ -functions of τ_{gen} and τ''_{gen} are directly proportional after identification of the underlying spaces.

A.8 We have shown, via the identifications, in **A.6** that the μ -function of τ is directly proportional to the one of τ''_{gen} and in **A.7** that the μ -function of τ''_{gen} is directly proportional to the one of τ_{gen} . This implies that the μ -functions of τ and τ_{gen} are directly proportional.

In conclusion, one sees that, for every discrete series representation τ of M in the Vogan L -packet, there exists a discrete series representation τ' of M' in the same Vogan L -packet with directly proportional μ -function. As, as we recalled above in **A.2**, it is known that the members of a discrete series L -packet of a Levi subgroup of a quasi-split classical group have the same μ -function, this finishes the proof of the theorem **A.1**. \square

REFERENCES

- [A] J. Arthur, *The endoscopic classification of representations - orthogonal and symplectic groups*, Colloquium Publications, vol. 61, AMS, 2013.
- [BP] R. Beuzart-Plessis, *Endoscopie et conjecture locale raffinée de Gan-Gross-Prasad pour les groupes unitaires*, *Compositio Mathematica*, **151** (2015), 1309–1371.
- [BW] A. Borel, N. Wallach, *Continuous Cohomology, Discrete Subgroups and Representations of Reductive Groups* (. Princeton University Press, Princedeton, ed.), 1980.
- [Co] J. Cohen, *A spectral expression for a certain orbital integral*, arXiv:1407.4316v2, 34 p. (2014).
- [GT] W.-T. Gan, W. Tanton, *The local Langlands conjecture for $GSp(4)$: The case of inner forms*, *Amer. J. Math.* **136** (2014), 761–805.
- [H1] V. Heiermann, *Décomposition spectrale d'un groupe réductif p -adique*, *J. Inst. Math. Jussieu* **3** (2004), 327–395.
- [H2] V. Heiermann, *Orbites unipotentes et pôles d'ordre maximal de la fonction μ de Harish-Chandra*, *Cand. J. Math.* **58** (2006), 1203–1228.
- [H3] V. Heiermann, *Unipotent Orbits and Local L -functions*, *J. reine angew. Math.* **596**, (2006), 103–114.
- [H4] V. Heiermann, *Langlands correspondence for classical groups and affine Hecke algebras*, arXiv:1502.04357v2, 29 p. (2015).
- [HM] V. Heiermann, G. Muić, *Standard Modules Conjecture*, *Math. Z.* **255** (2007), 847–853.
- [HO] V. Heiermann, E. Opdam, *On the tempered L -function conjecture*, *Am. J. Math.* **135** (2013), 777–800.
- [HI] K. Hira, A. Ichino, T. Ikeda, *Formal degrees and adjoint γ -factors*, *J. Amer. Math. Soc.*, **21** (2008), 283–304.
- [KMSW] ■ Kaletha, A. Minguez, S.-W. Shin, P.-J. White, *Endoscopic Classification of Representations: Inner Forms of Unitary Groups*, arXiv:1409:3731v3 (2014).
- [M1] C. Moeglin, *Multiplicité 1 dans les paquets d'Arthur*, dans "On Certain L-Functions: Proceedings of a conference in honor of Freydoon Shahidi's 60th Birthday" (J. Arthur, J. Cogdell, S. Gelbart, D. Goldberg, S. Kudla, D. Ramakrishnan and J.-K- Yu, eds.), AMS/Clay Math Institute - Proceedings, 2011.
- [M2] C. Moeglin, *Classification et Changement de base pour les séries discrètes des groupes unitaires p -adiques*, *Pacific J. Math.* **233** (2007), 159–204.
- [MW] C. Moeglin, J.-L. Waldspurger, *La conjecture locale de Gross-Prasad pour les groupes spéciaux orthogonaux: le cas général*, *Astérisque* **347** (2012), 167–216.
- [Mk] C.-P. Mok, *Endoscopic Classification of representations of Quasi-Split Unitary Groups*, *Mem. AMS* **235**, no. **1108** (2015).

- [Mu] G. Muić, *A proof of Casselman-Shahidi's Conjecture for quasi-split classical groups*, *Canad. Math. Bull.* **43** (2000), 90–99.
- [Sh] F. Shahidi, *A Proof of Langlands' Conjecture on Plancherel Measures; Complementary Series of p -adic groups*, *Ann. Math.* **132** (1990), 273–330.
- [Si] A. Silberger, *Special representations of reductive p -adic groups are not integrable*, *Ann. of Math.* **111** (1980), 571–587.
- [T] M. Tadić, *On reducibility and unitarizability for classical p -adic groups, some general results*, *Canad. J. Math.* **61** (2009), 427–450.
- [V] D. Vogan, *The local Langlands conjecture*, *Contemporary Maths*, AMS **145** (1993), 305–379.
- [W] J.-L. Waldspurger, *La formule de Plancherel pour les groupes p -adiques (d'après Harish-Chandra)*, *J. Inst. Math. Jussieu* **2** (2003), 235–333.

AIX-MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 13453
MARSEILLE, FRANCE;

E-mail address: volker.heiermann@univ-amu.fr