

Diophantine Equations and Congruent Number Equation Solutions

Mamuka Meskhishvili

Abstract

By using pairs of nontrivial rational solutions of congruent number equation

$$C_N : y^2 = x^3 - N^2x,$$

constructed are pairs of rational right (Pythagorean) triangles with one common side and the other sides equal to the sum and difference of the squares of the same rational numbers.

The parametrizations are found for following Diophantine systems:

$$\begin{aligned}(p^2 \pm q^2)^2 - a^2 &= \square_{1,2}, \\ c^2 - (p^2 \pm q^2)^2 &= \square_{1,2}, \\ a^2 + (p^2 \pm q^2)^2 &= \square_{1,2}, \\ (p^2 \pm q^2)^2 - a^2 &= (r^2 \pm s^2)^2.\end{aligned}$$

Keywords. Diophantine equations, Pythagorean triangles, parametrization, congruent number equation, congruent curve.

2010 AMS Classification. 11D25, 11D41, 11D72, 14G05, 14H52.

1 Introduction

By using rational solution (x, y) of congruent number equation

$$C_N : y^2 = x^3 - N^2x,$$

it is possible to construct a rational right triangle (a, b, c) with area N [1]

$$\begin{aligned} a &= \sqrt{x + N} - \sqrt{x - N}, \\ b &= \sqrt{x + N} + \sqrt{x - N}, \\ c &= 2\sqrt{x}. \end{aligned} \tag{1}$$

This construction is not possible from every solution. From (1) follows: a rational right triangle exists, if and only if solution x satisfies:

$$x = \square, \quad x + N = \square, \quad x - N = \square. \tag{2}$$

These conditions we call Fibonacci conditions, because Fibonacci was the first to find such a number:

$$x = \left(\frac{41}{12}\right)^2, \quad N = 5.$$

Congruent equation solutions which satisfy (2) we call Fibonacci solutions, respectively. Fibonacci solution has three properties:

$$x = \left(\frac{L}{K}\right)^2, \quad K = 2M, \quad (L, N) = 1,$$

where L , K and M positive integers (as well N).

If there exists even one nontrivial ($y \neq 0$) rational solution of C_N , that means there exists infinitely many rational solutions. Among them are

infinitely many Fibonacci and non-Fibonacci solutions. The tangent and the secant methods can be used to construct infinitely many solutions from the original. Draw the tangent line to C_N curve at original point (x, y) , this line will meet the curve in a second point [1]:

$$\left(\left(\frac{x^2 + N^2}{2y} \right)^2; \frac{(x^2 + N^2)(x^4 + N^4 - 6x^2N^2)}{8y^3} \right).$$

Now using a secant is possible to produce a new rational point and so forth.

In this paper is shown the usage of congruent equation solutions to solve Diophantine systems, including Fibonacci and non-Fibonacci solutions as well. Through them are constructed pairs of rational right (Pythagorean) triangles with one common side and the other sides equal to the sum and difference of the squares of the same rational numbers $p^2 \pm q^2$.

2 Common Legs, Hypotenuses $p^2 \pm q^2$

The shared (common) legs denote $a = a_1 = a_2$ and hypotheses $c_{1,2} = p^2 \pm q^2$. This case is equivalent to solving Diophantine system:

$$\begin{cases} (p^2 + q^2)^2 - a^2 = \square, \\ (p^2 - q^2)^2 - a^2 = \square. \end{cases} \quad (3)$$

We try to find nontrivial integer solutions of system (3). Nontrivial solutions mean both Pythagorean triangles exist and they are distinct:

$$a_1 b_1 c_1 a_2 b_2 c_2 \neq 0,$$

or in system notations:

$$apq \neq 0, \quad p \neq q.$$

By using rational parametrization formulas for unit hyperbola:

$$\frac{p^2 + q^2}{a} = \frac{1 + \xi^2}{2\xi}, \quad \frac{p^2 - q^2}{a} = \frac{1 + \zeta^2}{2\zeta}, \quad (4)$$

where ξ and ζ are arbitrary nontrivial rationals. After simplification (4)

$$\begin{aligned} p^2 &= \frac{a(\xi + \zeta)(1 + \xi\zeta)}{4\xi\zeta}, \\ q^2 &= \frac{a(\zeta - \xi)(1 - \xi\zeta)}{4\xi\zeta}. \end{aligned} \quad (5)$$

From (5):

$$\begin{aligned} (\zeta^2 - \xi^2)(1 - (\xi\zeta)^2) &= \square, \\ \left(\frac{\xi}{\zeta}\right)(\xi\zeta)\left(1 - \left(\frac{\xi}{\zeta}\right)^2\right)(1 - (\xi\zeta)^2) &= \square. \end{aligned}$$

By using Property 2 in [2] it is possible to express ξ and ζ by arbitrary nontrivial different rational solutions x and z ($xz = \square$) of arbitrary C_N congruent number equation:

$$\frac{\xi}{\zeta} = \frac{x}{N}, \quad \xi\zeta = \frac{z}{N}.$$

Accordingly,

$$\xi = \frac{\sqrt{xz}}{N}, \quad \zeta = \sqrt{\frac{z}{x}}. \quad (6)$$

Substitute (6) in (5)

$$p^2 = \frac{a}{4N\sqrt{xz}} (x + N)(z + N),$$

$$q^2 = \frac{a}{4N\sqrt{xz}} (x - N)(z - N).$$

Theorem 1. *Nontrivial integer solutions of system (3) are given by formulas:*

$$a = k \cdot 4N\sqrt{xz},$$

$$p^2 = k \cdot (x + N)(z + N),$$

$$q^2 = k \cdot (x - N)(z - N),$$

where k is integer, x and z are nontrivial different rational solutions of arbitrary congruent number equation.

Integer k and solutions x , z must be chosen so that a , p and q are integers.

Direct calculations give:

$$a_1 = k \cdot 4N\sqrt{xz},$$

$$b_1 = k \cdot 2(xz - N^2),$$

$$c_1 = p^2 + q^2 = k \cdot 2(xz + N^2);$$

(7)

$$a_2 = k \cdot 4N\sqrt{xz},$$

$$b_2 = k \cdot 2N(x - z),$$

$$c_2 = p^2 - q^2 = k \cdot 2N(x + z).$$

Numerical Examples

To construct examples of Diophantine system (3), we use C_N solution from [1].

$$1) N = 6; \quad x = 18, z = \frac{19602}{47^2}.$$

$$(74^2 + 23^2)^2 - 4653^2 = 3796^2,$$

$$(74^2 - 23^2)^2 - 4653^2 = 1680^2.$$

$$2) N = 34; \quad x = 162, z = \frac{2178}{7^2}.$$

$$(217^2 + 64^2)^2 - 35343^2 = 37024^2,$$

$$(217^2 - 64^2)^2 - 35343^2 = 24480^2.$$

3 Common Hypotheses (Siblings), Legs $p^2 \pm q^2$

The hypotenuses denote $c = c_1 = c_2$ and legs $a_{1,2} = p^2 \pm q^2$. Diophantine system is:

$$\begin{cases} c^2 - (p^2 + q^2)^2 = \square, \\ c^2 - (p^2 - q^2)^2 = \square. \end{cases} \quad (8)$$

Again we seek nontrivial integer solutions of system (8)

$$a_1 b_1 c_1 a_2 b_2 c_2 \neq 0;$$

$$cpq \neq 0, \quad p \neq q.$$

By using parametrization formulas for unit circle:

$$\frac{p^2 + q^2}{c} = \frac{2\xi}{1 + \xi^2}, \quad \frac{p^2 - q^2}{a} = \frac{2\zeta}{1 + \zeta^2}. \quad (9)$$

After simplification (9):

$$\begin{aligned} p^2 &= \frac{c(\xi + \zeta)(1 + \xi\zeta)}{(1 + \xi^2)(1 + \zeta^2)}, \\ q^2 &= \frac{c(\xi - \zeta)(1 - \xi\zeta)}{(1 + \xi^2)(1 + \zeta^2)}. \end{aligned} \quad (10)$$

From (10)

$$(\xi^2 - \zeta^2)(1 - (\xi\zeta)^2) = \square.$$

Again using Property 2 in [2]:

$$\xi = \frac{\sqrt{xz}}{N}, \quad \zeta = \sqrt{\frac{z}{x}}, \quad (11)$$

where x and z are nontrivial different rational solutions of arbitrary congruent number equation.

Substitute (11) in (10):

$$\begin{aligned} p^2 &= \frac{c\sqrt{xz}(x + N)(z + N)}{(x + z)(xz + N^2)}, \\ q^2 &= \frac{c\sqrt{xz}(x - N)(z - N)}{(x + z)(xz + N^2)}. \end{aligned}$$

So,

Theorem 2. *Nontrivial integer solutions of system (8) are given by formulas:*

$$\begin{aligned}c &= k \cdot \sqrt{xz} (x + z)(xz + N^2), \\p^2 &= k \cdot xz(x + N)(z + N), \\q^2 &= k \cdot xz(x - N)(z - N),\end{aligned}$$

where k is integer, x and z are nontrivial different rational solutions of arbitrary congruent number equation.

Integer k and solutions x , z must be chosen so that c , p and q are integers.

Direct calculations give:

$$\begin{aligned}a_1 &= p^2 + q^2 = k \cdot 2xz(xz + N^2), \\b_1 &= k \cdot \sqrt{xz} (x - z)(xz + N^2), \\c_1 &= k \cdot \sqrt{xz} (x + z)(xz + N^2); \\a_2 &= p^2 - q^2 = k \cdot 2Nxz(x + z), \\b_2 &= k \cdot \sqrt{xz} (x + z)(xz - N^2), \\c_2 &= k \cdot \sqrt{xz} (x + z)(xz + N^2).\end{aligned}\tag{12}$$

Numerical Examples

To construct examples of Diophantine system (8), we use the same solutions.

$$1) \ N = 6; \quad x = 18, \ z = \frac{19602}{47^2}.$$

$$15358381995^2 - (114774^2 + 35673^2)^2 = 5215702800^2,$$

$$15358381995^2 - (114774^2 - 35673^2)^2 = 9708645804^2.$$

$$2) \ N = 34; \quad x = 162, \ z = \frac{2178}{7^2}.$$

$$3322469535^2 - (50127^2 + 14784^2)^2 = 1891797600^2,$$

$$3322469535^2 - (50127^2 - 14784^2)^2 = 2403264864^2.$$

4 Common Legs, Other Legs $p^2 \pm q^2$

Denote common legs $a = a_1 = a_2$ and other legs $b_{1,2} = p^2 \pm q^2$. Diophantine system is

$$\begin{cases} a^2 + (p^2 + q^2)^2 = \square, \\ a^2 + (p^2 - q^2)^2 = \square. \end{cases} \quad (13)$$

We seek nontrivial integer solutions of system (13):

$$a_1 b_1 c_1 a_2 b_2 c_2 \neq 0;$$

$$apq \neq 0, \quad p \neq q.$$

Rational parametrization for unit hyperbola gives:

$$\frac{p^2 + q^2}{a} = \frac{1 - \xi^2}{2\xi}, \quad \frac{p^2 - q^2}{a} = \frac{1 - \zeta^2}{2\zeta}. \quad (14)$$

Accordingly from (14):

$$\begin{aligned} p^2 &= \frac{a(\xi + \zeta)(1 - \xi\zeta)}{4\xi\zeta}, \\ q^2 &= \frac{a(\zeta - \xi)(1 + \xi\zeta)}{4\xi\zeta}. \end{aligned} \quad (15)$$

From (15)

$$(\zeta^2 - \xi^2)(1 - (\xi\zeta)^2) = \square.$$

Again using Property 2 in [2]:

$$\xi = \frac{\sqrt{xz}}{N}, \quad \zeta = \sqrt{\frac{z}{x}},$$

and after substitution in (15)

$$\begin{aligned} p^2 &= \frac{a(x + N)(z - N)}{4N\sqrt{xz}}, \\ q^2 &= \frac{a(x - N)(z + N)}{4N\sqrt{xz}}. \end{aligned}$$

Theorem 3. *Nontrivial integer solutions of system (13) are given by formulas:*

$$\begin{aligned} a &= k \cdot 4N\sqrt{xz}, \\ p^2 &= k \cdot (x + N)(z - N), \\ q^2 &= k \cdot (x - N)(z + N), \end{aligned}$$

where k is integer, x and z are nontrivial different rational solutions of arbitrary congruent number equation.

Integer k and solutions x , z must be chosen so that a , p and q are integers.

Direct calculations give:

$$\begin{aligned} a_1 &= k \cdot 4N\sqrt{xz}, \\ b_1 &= p^2 + q^2 = k \cdot 2(xz - N^2), \\ c_1 &= k \cdot 2(xz + N^2); \\ a_2 &= k \cdot 4N\sqrt{xz}, \\ b_2 &= p^2 - q^2 = k \cdot 2N(x - z), \\ c_2 &= k \cdot 2N(x + z). \end{aligned} \tag{16}$$

Solutions of system (13) are impossible to find using the preceding congruent number equation solutions, so we have to take other C_N solutions.

Numerical Examples

$$1) N = 5; \quad x = \frac{12005}{31^2}, \quad z = 45.$$

$$4557^2 + (82^2 + 60^2)^2 = 11285^2,$$

$$4557^2 + (82^2 - 60^2)^2 = 5525^2.$$

$$2) N = 34; \quad x = \frac{833}{4^2}, \quad z = \frac{153}{2^2}.$$

$$1344^2 + (17^2 + 9^2)^2 = 1394^2,$$

$$1344^2 + (17^2 - 9^2)^2 = 1360^2.$$

5 Common Legs, Hypotenuses $p^2 \pm q^2$, Other Legs $r^2 \pm s^2$

Preceding numerical examples are constructed by using non-Fibonacci solutions. Of course if we use Fibonacci solution (x, z) it is possible to construct solutions for each Diophantine systems (3), (8) and (13). It indicates that intersection case may exist.

Consider Pythagorean triangles with sides:

$$(a, r^2 + s^2, p^2 + q^2), \quad (a, r^2 - s^2, p^2 - q^2),$$

accordingly

$$a = a_1 = a_2, \quad b_{1,2} = r^2 \pm s^2, \quad c_{1,2} = p^2 \pm q^2.$$

Corresponding Diophantine system is:

$$\begin{cases} (p^2 + q^2)^2 - a^2 = (r^2 + s^2)^2, \\ (p^2 - q^2)^2 - a^2 = (r^2 - s^2)^2; \end{cases} \quad (17)$$

again we seek nontrivial integer solutions:

$$apqrs \neq 0, \quad p \neq q, \quad r \neq s.$$

Divide system (17) by two subsystems:

$$\begin{cases} (p^2 + q^2)^2 - a^2 = \square, \\ (p^2 - q^2)^2 - a^2 = \square \end{cases} \quad \text{and} \quad \begin{cases} a^2 + (r^2 + s^2)^2 = \square, \\ a^2 + (r^2 - s^2)^2 = \square. \end{cases}$$

For these subsystems use parametric formulas from Theorem 1 and Theorem 3, respectively:

$$\begin{aligned} a &= k_1 \cdot 4N_1 \sqrt{x_1 z_1}, \\ p^2 &= k_1 \cdot (x_1 + N_1)(z_1 + N_1), \\ q^2 &= k_1 \cdot (x_1 - N_1)(z_1 - N_1); \\ a &= k_2 \cdot 4N_2 \sqrt{x_2 z_2}, \\ r^2 &= k_2 \cdot (x_2 + N_2)(z_2 - N_2), \\ s^2 &= k_2 \cdot (x_2 - N_2)(z_2 + N_2). \end{aligned} \quad (18)$$

Expressions (7) and (16) gives:

$$k_1 \cdot 4N_1\sqrt{x_1z_1} = k_2 \cdot 4N_2\sqrt{x_2z_2}, \quad (19_1)$$

$$k_1 \cdot 2(x_1z_1 - N_1^2) = k_2 \cdot 2(x_2z_2 - N_2^2), \quad (19_2)$$

$$k_1 \cdot 2(x_1z_1 + N_1^2) = k_2 \cdot 2(x_2z_2 + N_2^2), \quad (19_3)$$

$$k_1 \cdot 2N_1(x_1 - z_1) = k_2 \cdot 2N_2(x_2 - z_2), \quad (19_4)$$

$$k_1 \cdot 2N_1(x_1 + z_1) = k_2 \cdot 2N_2(x_2 + z_2). \quad (19_5)$$

The sum and difference (19₂) and (19₃), (19₄) and (19₅) gives:

$$k_1N_1^2 = k_2N_2^2, \quad (20_1)$$

$$k_1x_1z_1 = k_2x_2z_2, \quad (20_2)$$

$$k_1N_1x_1 = k_2N_2x_2, \quad (20_3)$$

$$k_1N_1z_1 = k_2N_2z_2. \quad (20_4)$$

$$(20_3), (20_1) \implies \frac{x_1}{x_2} = \frac{k_2N_2}{k_1N_1} = \frac{N_1}{N_2}, \quad (21_1)$$

$$(20_4), (20_1) \implies \frac{z_1}{z_2} = \frac{k_2N_2}{k_1N_1} = \frac{N_1}{N_2}. \quad (21_2)$$

From (21₁) follows, exists rational t

$$N_1 = x_1t \quad \text{and} \quad N_2 = x_2t.$$

Because x_1 and x_2 are solutions C_{N_1} and C_{N_2} , respectively:

$$\begin{aligned} x_1(x_1^2 - N_1^2) &= x_1^3(1 - t^2) = \square, \\ x_2(x_2^2 - N_2^2) &= x_2^3(1 - t^2) = \square. \end{aligned} \quad (22)$$

From (22) follows

$$\frac{x_1}{x_2} = \square. \quad (23)$$

Expressions (23) and (21₁) gives

$$\frac{N_1}{N_2} = \square.$$

Because congruent numbers are squafree so,

$$N_1 = N_2. \quad (24)$$

From (21₁), (21₂) and (20₁) follows

$$x_1 = x_2, \quad z_1 = z_2, \quad k_1 = k_2. \quad (25)$$

Conditions (24) and (25) give parametric solution of this intersection case:

Theorem 4. *Nontrivial integer solutions of system (17) are given by formulas:*

$$\begin{aligned} p^2 &= k \cdot (x + N)(z + N), \\ q^2 &= k \cdot (x - N)(z - N), \\ a &= k \cdot 4N\sqrt{xz}, \\ r^2 &= k \cdot (x + N)(z - N), \\ s^2 &= k \cdot (x - N)(z + N), \end{aligned} \quad (26)$$

where k is integer, x and z are nontrivial different rational solutions of arbitrary congruent number equation.

Integer k and solutions x, z must be chosen so that p, q, a, r, s are integers. Because Fibonacci solutions satisfy (2), from them it is always possible to construct integer solution of (17).

Direct calculations give:

$$a_1 = k \cdot 4N\sqrt{xz},$$

$$b_1 = k \cdot 2(xz - N^2),$$

$$c_1 = k \cdot 2(xz + N^2);$$

$$a_2 = k \cdot 4N\sqrt{xz},$$

$$b_2 = k \cdot 2N(x - z),$$

$$c_2 = k \cdot 2N(x + z).$$

Numerical Examples

1) Non-Fibonacci pair:

$$N = 5; \quad x = \left(\frac{5}{2}\right)^2, \quad z = \left(\frac{41}{12}\right)^2.$$

$$(735^2 + 155^2)^2 - 492000^2 = (465^2 + 245^2)^2,$$

$$(735^2 - 155^2)^2 - 492000^2 = (465^2 - 245^2)^2.$$

2) Fibonacci pair:

$$N = 6; \quad x = \left(\frac{5}{2}\right)^2, \quad z = \left(\frac{1201}{140}\right)^2.$$

$$(8743^2 + 1151^2)^2 - 40353600^2 = (8057^2 + 1249^2)^2,$$

$$(8743^2 - 1151^2)^2 - 40353600^2 = (8057^2 - 1249^2)^2.$$

Diophantine system (17) is equivalent to:

$$\begin{cases} p^4 + q^4 - a^2 = r^4 + s^4, \\ pq = rs. \end{cases} \quad (27)$$

By Theorem 4 the second equation is satisfied automatically. This means: the parametric formulas in (26) are solution of the first equation in system (27).

Theorem 5. *Nontrivial integer solutions of equation*

$$p^4 + q^4 - a^2 = r^4 + s^2, \quad (28)$$

which satisfy $pq = rs$ condition, are given by parametric formulas (26).

Obviously there are solutions (28) which do not satisfy $pq = rs$ condition. For example, it is possible to construct such solutions by using the equal sum of biquadratics [3]:

$$\begin{aligned} 7^4 + 28^4 &= 3^4 + 20^4 + 26^4, \\ 51^4 + 76^4 &= 5^4 + 42^4 + 78^4, \\ 37^4 + 38^4 &= 25^4 + 26^4 + 42^4. \end{aligned}$$

Numerical Examples

$$\begin{aligned} 735^4 + 155^4 - 492000^2 &= 465^4 + 245^4, \\ 8743^4 + 1151^4 - 40353600^2 &= 8057^4 + 1249^4. \end{aligned}$$

References

1. Meskhishvili M., Three-Century Problem. *Tbilisi*, 2013.
2. Meskhishvili M., Perfect Cuboid and Congruent Number Equation Solutions. 2013, *Unsolved Problems in Number Theory, Logic, and Cryptography*, www.unsignedproblems.org; arxiv.org/pdf/1211.6548v2.pdf.
3. Dickson L. E., History of the Theory of Numbers, Volume II: Diophantine Analysis. *Dover, New York*, 2005.

Author's address:

Georgian-American High School, 18 Chkondideli Str., Tbilisi 0180, Georgia.
E-mail: director@gahs.edu.ge