

Random Walk in Changing Environment

Gideon Amir ^{*} Itai Benjamini [†] Ori Gurel-Gurevich [‡] Gady Kozma [§]

September 26, 2018

Abstract

In this paper we introduce the notion of *Random Walk in Changing Environment* - a random walk in which each step is performed in a different graph on the same set of vertices, or more generally, a weighted random walk on the same vertex and edge sets but with different (possibly 0) weights in each step. This is a very wide class of RW, which includes some well known types of RW as special cases (e.g. reinforced RW, true SAW). We define and explore various possible properties of such walks, and provide criteria for recurrence and transience when the underlying graph is \mathbb{N} or a tree. We provide an example of such a process on \mathbb{Z}^2 where conductances can only change from 1 to 2 (once for each edge) but nevertheless the walk is transient, and conjecture that such behaviour cannot happen when the weights are chosen in advance, that is, do not depend on the location of the RW.

1 Introduction

Theseus is thrown into Daedalus' labyrinth, this time without a ball of thread. Noticing that the labyrinth is a subgraph of \mathbb{Z}^2 , Theseus decides to simply random walk his way out - he knows that he will almost surely reach the exit eventually. What Theseus doesn't know is that Daedalus, aware of the recurrency of his labyrinth, is working relentlessly to amend this vulnerability. He is continually digging new passages throughout the labyrinth, following a carefully laid plan. He cannot, however, block existing passages, only create new ones and only between adjacent rooms, such that the labyrinth is a subgraph of \mathbb{Z}^2 at any point. Will Theseus find his way to the exit or is it possible that Daedalus' cunning plan will deceive him forever (with positive probability)?

It turns out that if Daedalus is aware of Theseus whereabouts he can devise a plan to lure poor Theseus further and further into the labyrinth with positive probability (see Thm 6.1). We conjecture that this is not the case if Daedalus is not aware of Theseus whereabouts.

^{*}Bar-Ilan University, Ramat-Gan, 5290002, Israel. Email: gidi.amir@gmail.com

[†]Weizmann Institute, Rehovot, 76100, Israel. Email: itai.benjamini@weizmann.ac.il

[‡]The Hebrew University of Jerusalem, 91904, Israel. Email: origurel@math.huji.ac.il

[§]Weizmann Institute, Rehovot, 76100, Israel. Email: gady.kozma@weizmann.ac.il

Conjecture 1.1. *If Daedalus is oblivious of Theseus location, then Theseus will almost surely reach the exit (infinitely many times, if he chooses to stay in the labyrinth). In other words, Theseus' Random Walk is recurrent.*

See Section 7 for formal statement and further open problems.

In this paper we introduce the notion of *Random Walk in Changing Environment* or *RWCE*. Generally speaking, a RWCE is a random walk in which each step is performed in a different graph on the same set of vertices. By different we may mean that the set of edges is different, but it is easier and more general to assume that the underlying graph is the same throughout the walk and what changes are the conductances of the edges. This is a very wide class of RW, which includes some well known types of RW as special cases, most notably the reinforced RW. We define and explore various possible properties of RWCE with the conclusion that the interesting case is when the walk is monotone (Daedalus can only create passages) and bounded (all the passages are edges of \mathbb{Z}^2). Under these assumptions we give criteria for recurrence and transience when the underlying graph is \mathbb{N} or a tree, even when the sequence of graphs may depend on the history of the walk (the latter we call "adaptive", see section 2 for exact definitions).

We show that the above criteria cannot hold for general graphs: we provide an example where the underlying graph is \mathbb{Z}^2 , but the RWCE is transient. This example is an RWCE on \mathbb{Z}^2 where each edge is started with weight 1 and at each stage we change the weight of the edge to the right of the walker to 2 if this was not already done. The idea behind this construction is to try and mimic the behaviour of excited random walk on two dimensions (see [2]) in which the walker gets a bias to the right whenever it visits a point for the first time, and was shown in [2] to be transient. However, it turns out that the proof carried out in [2] depends quite delicately on the model, and one must take care when working out the details. In particular, a similar attempt to mimic multi-excited RW in dimension 1 (see [21]) cannot succeed, as follows from our results on RWCEs on trees. Note that the above example was an adaptive RWCE, and we conjecture that such behaviour cannot occur in nonadaptive RWCE.

Related works: In recent years there have been a number of papers that studied related models. These works have some overlap with our model and some of the examples, but not with the results of this paper, and generally speaking the emphasis of these works are in different directions. Avin, Kouck and Lotker [1] studied RWCE on a sequence of finite unweighted graphs (which they called "evolving graphs") They were interested mainly in the problem of the cover time of the walk, showing, in particular, that contrary to a regular random walk on the graph, the RWCE may have exponential cover time. Dembo, Huang and Sidoravicius studied models on random walks on "monotone domains" - that is they assumed that the sequence of graphs in the RWCE is an increasing sequence of subgraphs of a pre-given graph, with a focus on \mathbb{Z}^d . They proved criteria for recurrence and transience of such walks, with one paper [7] focusing on the nonadaptive case (where they also consider a continuous analog for brownian motion), and the other [6] focusing on the adaptive case - that is when there is an interaction between the walk and the graph sequence.

As will be seen in section 3, the generality of these models implies that some further assumptions must be taken in order to get meaningful criteria, and Dembo, Huang and Sidoravicius focus on several interaction mechanisms (such as, e.g. the walker uncovering new edges when it approaches them) and give criteria for transience and recurrence as well as some conjectures, some of which carry a similar flavour to the ones in this paper.

Structure of the paper: In Section 2 we give the basic definitions and examples of known random walk models which falls into our framework. In Section 3 we give simple examples of Random Walks in Changing Environment which illustrate that if the environment is unbounded or nonmonotone then the random walk can have (almost) any behavior. Sections 4 and 5 give the main results about bounded monotone RWCE on \mathbb{N} and on trees, respectively. Section 6 gives an example of a bounded monotone (adaptive) RWCE on \mathbb{Z}^2 which is transient, thus showing that the results on recurrent trees cannot be extended to general recurrent graphs. We conclude with a conjecture and some open problems.

2 Definitions

We begin by giving a rigorous definition of what a random walk in changing environment is, in the broadest sense. In this paper, we use discrete time, see section 7 for a brief discussion of the continuous time version.

Definition 1. A Random Walk in Changing Environment (RWCE), on a graph $G = (V, E)$ is a stochastic process $\{\langle X_t, G_t \rangle\}_{t=0}^\infty$, where $G_t = (V, E, C_t)$ are graphs with a conductances function $C_t : E \rightarrow [0, \infty)$ over a fixed vertex set V and edge set E , and for all t , $X_t \in V$ and

$$\mathbb{P}(X_{t+1} = v \mid \langle X_0, G_0 \rangle, \dots, \langle X_t, G_t \rangle) = \frac{C_t(X_t, v)}{\sum_{\{e \in E \mid X_t \in e\}} C_t(e)} .$$

We call the sequence $\{X_t\}_{t=0}^\infty$ the Random Walk and the sequence $\{G_t\}_{t=0}^\infty$ the Environment.

In other words, the law of the process governs the changes in the environment, while the distribution of X_t , given the history, only depends only G_t and is the same as a random walk step on the graph G with weights C_t . Note that the conductances C_t may depend on the history of the process so far and on extra randomness.

In our Labyrinth example, Daedalus was creating new edges, not changing conductances. It is easy to see, however, that the definition using conductances is a generalization of this scenario.

Definition 2. A RWCE is called proper if $0 < C_t(e) < \infty$ for all $t \in \mathbb{N}$ and $e \in E$. It is called improper otherwise.

Definition 3. A RWCE is said to be bounded from above (below) by $G = (V, E, C)$ if $C_t(e) \leq C(e)$ ($C_t(e) \geq C$) for all $e \in E$ and all $t \in \mathbb{N}$, almost surely.

Note that a RWCE bounded from above and below is necessarily proper. All the RWCE in this paper are proper unless otherwise noted. Also note the requirement $C_t(e) < \infty$ in the definition is formally redundant as the conductances were defined to be real numbers. However, in the more naive approach of a changing graph, a conductance of infinity would correspond to merging (shorting) two vertices together.

Definition 4. *A RWCE is called nonadaptive if the distribution of G_{t+1} given G_0, \dots, G_t is independent of X_0, \dots, X_t . It is called adaptive otherwise.*

The Labyrinth example is nonadaptive if Daedalus is oblivious of Theseus whereabouts or adaptive if Daedalus responds to it.

Definition 5. *A RWCE is called monotone increasing (decreasing) if $C_{t+1} \geq C_t$ ($C_{t+1} \leq C_t$) almost surely.*

The Labyrinth example is monotone increasing, since Daedalus only adds new edges, i.e. raises the conductance by 1.

Note that the definition of a general RWCE is very broad. Actually, it is too broad, as an adaptive, improper, nonmonotone RWCE on the full graph can implement any behavior at any stage. But even with some restrictions, many interesting walks can be implemented as RWCE in a natural way. We next give several examples of well-known random walks and how they fit into our definition:

Example 2.1. *The once-reinforced random walk (see [8]) on \mathbb{Z}^2 , is a proper, adaptive monotone increasing RWCE. At the beginning the conductance of each edge is 1, and at each stage, if the RW traversed an edge with conductance 1, replace it with an edge of conductance c (for a fixed constant c). This RWCE is bounded between 1 and c . Other reinforced random walks also fit similarly into the RWCE framework. See [16] for a survey of such models.*

Example 2.2. *The Bridge Burning Random Walk (where the conductance of each edge the walk traverses is reduced to 0) on \mathbb{Z}^2 , is an improper, adaptive monotone decreasing RWCE. It is as the once-reinforced RW with $c = 0$.*

Example 2.3. *The Laplacian random walk from between v_0 and u (which is equivalent to the loop erased random walk from v_0 to u , see [13]), which starts at $X_0 = v_0$ and chooses which neighbour to move to at each step with probabilities proportional to the value of the harmonic function which is 0 on the path of the RW up to this time and 1 on u , can be described as a monotone bounded improper adaptive RWCE.*

Example 2.4. *The "true" self-avoiding walk with bond repulsion (see e.g. [18]) is a nearest neighbor random walk, for which the probability of jumping along an edge e is proportional to $e^{-ck(e)}$, where $k(e)$ is the number of times e has been traversed. This is an adaptive, monotone, proper RWCE.*

The main question we will be interested in, is whether a given RWCE is recurrent. Note that for RWCE, the dichotomy between recurrence and transience is not always as clear cut as for simple RW. There might be a difference between a.s. returning to the origin, a.s. visiting every vertex, a.s. returning to the origin infinitely many times and a.s. visiting every vertex infinitely many times. Also, since no 0-1 law holds in general for RWCE, we can have a RWCE which return to the origin infinitely many times with probability which is positive but less than 1.

In most natural cases, however, the various possible definitions of recurrence and transience for RWCE coincide. We will therefore use the strictest definitions.

Definition 6. *A RWCE on $G = (V, E)$ is called recurrent if it visits every vertex in V infinitely many times almost surely. A RWCE is called transient if it visits every vertex a finite number of times almost surely. The RWCE is said to be of mixed type otherwise.*

3 Simple examples

The aim of this section is to demonstrate the myriad possible behaviors of unrestricted RWCE. We begin with a simple example on general graphs. Let G be any graph and X_0 be a vertex in G .

Example 3.1. *For any distribution on paths in G (starting with X_0), there is an improper, adaptive, nonmonotone RWCE inducing this distribution on X .*

Since we have complete control over the conductances of the edges emerging from X_t , we can arbitrarily determine the distribution of the next step, and therefore the distribution of the sequence.

A distribution on paths in G is called *elliptic* if for every finite path in G , v_0, \dots, v_n , with $v_0 = X_0$, we have $\mathbb{P}(X_0 = v_0, \dots, X_n = v_n) > 0$.

Example 3.2. *For any elliptic distribution on paths in G (starting with X_0), there is a proper, adaptive, nonmonotone RWCE inducing this distribution on X .*

This example is the same as the previous one except you can't have probability 0 for any transition. Next, note that since multiplying the conductances by some constant does not change the next step distribution, the previous example can be made monotone, either increasing or decreasing. Also, the starting set of conductances C_0 can be arbitrary (except for conductances of edges emerging from X_0) and by monotonicity the RWCE is bounded (from above or below) by C_0 . Put together we have:

Example 3.3. *For any elliptic distribution on paths in $G = (V, E)$ (starting with X_0), and any (proper) choice of conductances C there is a proper, adaptive, monotone (increasing or decreasing) RWCE, bounded (from below or above, resp.) by (V, E, C) , inducing this distribution on X .*

If we drop monotonicity, but require boundedness instead then we can still produce any distribution that has *bounded conditional probabilities*, i.e. the probability for traversing a given edge is uniformly bounded away from zero. In particular, we have the following example on \mathbb{N} :

Example 3.4. *The RWCE with conductances $C_t(X_t, X_t + 1) = 2$ and $C_t(j, j + 1) = 1$ for $j \neq X_t$ is bounded from above and below by a recurrent graph, adaptive, nonmonotone and transient.*

Indeed, X_t is simply a biased RW and is therefore transient.

We have thus seen that neither boundedness nor monotonicity are enough to draw any significant conclusions about the RWCE, at least in the adaptive setting.

The next example shows that even in the nonadaptive setting, boundedness does not imply recurrence or transience.

Example 3.5. *The RWCE with conductances $C_t(j, j + 1) = 100$ for $t \equiv j \pmod{100}$ and $C_t(j, j + 1) = 1$ otherwise is bounded from above and below by a recurrent graph, nonadaptive, nonmonotone and transient.*

Sketch of proof. When $X_t \equiv t \pmod{100}$ the conductance to the right of X_t is 100 while to the left it is only 1. Therefore, with probability $100/101$, $X_{t+1} = X_t + 1$, in which case $X_{t+1} = t + 1 \pmod{100}$. This happens for an expected number of 101 times, after which the walk is simple until the next 100 conductance "catches up". This takes about 100 steps in which the expected displacement is 0. Bipartiteness of the graph ensures the walk never gets a bias to the left. All in all, the RW gets a strong bias to the right about half the time and so it is transient. ■

Note that the same conductances would work even if the RW had some probability of staying at the same vertex, thus nullifying the bipartiteness of the graph, though the calculation would be slightly more involved. The reason being that while the walker would sometime get a bias to the left, the wave would "pass" the walker once a step to the left was made.

Similarly, we can make the RW recurrent, even if it is bounded by a transient graph.

Example 3.6. *The RWCE with conductances $C_t(j, j + 1) = 1000 \cdot 2^j$ for $t \equiv -j \pmod{100}$ and $C_t(j, j + 1) = 2^j$ otherwise is bounded from above and below by a transient graph, nonadaptive, nonmonotone and recurrent.*

Sketch of proof. The argument is the same as in example 3.5, except that when $X_t \not\equiv -t \pmod{100}$ the RW gets a $(1/3, 2/3)$ bias to the right instead of being balanced. However, simple calculation shows that this bias is not enough to counter the bias to the left when $X_t \equiv -t \pmod{100}$, which once caught would persist as long as the walker keeps going left. ■

4 RWCE on \mathbb{N}

In this section we study RWCE whose underlying graph is \mathbb{N} (with edges between consecutive integers). All the theorems here apply equally to RWCE on \mathbb{Z} , but the proofs are slightly simpler for \mathbb{N} since there's only one way to infinity. For such graphs we can prove quite general conditions which ensure the RWCE is recurrent (or transient).

The main idea of the proofs in this section and the next is as follows. We will define a *potential sequence* - an adaptive sequence of functions $F_t : V \rightarrow \mathbb{R}^+$ satisfying:

1. *Harmonicity*: F_t is harmonic on (V, E, C_t) except at 0.
2. *Monotonicity*: $F_t(v)$ is either monotone increasing for all $v \in V$ or monotone decreasing for all $v \in V$.

Note that F_t may depend on H_t , the history of the RWCE up to time t , even if the RWCE itself is nonadaptive. The two properties above imply that $F_t(X_t)$ is either a supermartingale or a submartingale as long as $X_t \neq 0$. This is because $\mathbb{E}(F_t(X_{t+1})|H_t) = F_t(X_t)$ by harmonicity of F_t and because $F_{t+1}(X_{t+1}) \geq F_t(X_{t+1})$ (or \leq) by monotonicity. We will then use the optional stopping theorem to deduce bounds on the probability of return to 0. Note that related ideas were used by Vervoort [19] and even earlier by Davis [4] in the context of reinforced random walks.

The following theorems all require the RWCE be bounded from below and above by some graph. When this condition holds, the walk is *elliptic* (uniformly in time), that is, the probability of traversing each edge when the walk is at one of its endpoints is bounded away from 0. On \mathbb{N} this implies that such a walk cannot stay on a finite segment indefinitely - it will a.s. visit every vertex to the right of its current location. Therefore, when trying to determine whether the process is recurrent or transient, we can assume that the walk starts at any vertex of \mathbb{N} , as long as the conditions of the theorem still hold for the RWCE at that time. Ellipticity also means that the walk is recurrent (by our definition) exactly when it visits some vertex infinitely many times almost surely and transient exactly when it visits some vertex only finitely many times almost surely. Throughout this section we write $C(j)$ instead of $C(j, j + 1)$ to abbreviate notation.

Theorem 4.1. *If $\{(X_t, G_t)\}$ is a monotone increasing adaptive RWCE on \mathbb{N} , bounded above by some recurrent connected graph $G_\infty = (\mathbb{N}, C_\infty)$ then the walk is recurrent.*

Proof. Notice that since the RWCE may be adaptive, G_∞ is just a bound on G_i and not necessarily its limit.

Assume that the walk starts at some $X_0 > 0$. We will show that the walk almost surely hits 0. Since the conditions of the Theorem continue to hold at this hitting time, this implies the walk will

a.s. hit 0 infinitely often and is therefore recurrent. The potential sequence we use in this case is

$$F_t(v) = \sum_{j=0}^{v-1} \frac{1}{C_t(j)}$$

i.e. the resistance between 0 and v on the graph G_t . That F_t is harmonic on G_t is well known (and easily verified). Monotonicity follows from the monotonicity of the RWCE. Therefore, $F_t(X_t)$ is a super-martingale until the first time $X_t = 0$.

Since G_∞ is recurrent, we know that $\sum_{j=0}^{\infty} 1/C_\infty(j) = \infty$. Therefore, given any $A > 0$ there is a $v \in \mathbb{N}$ such that $F_\infty(v) \geq A$. Let τ to be the first time the walk hits either v or 0. By ellipticity, τ is finite almost surely. By the optional stopping theorem $F_0(X_0) \geq \mathbb{E}(F_\tau(X_\tau))$. Denote by p_v the probability that $X_\tau = v$, i.e. that the RW hits v before 0. Noting that $F_t(0) = 0$ for all t and that $F_t(v) \geq F_\infty(v) \geq A$ we have

$$F_0(X_0) \geq \mathbb{E}(F_\tau(X_\tau)) \geq Ap_v$$

and therefore

$$p_v \leq \frac{F_0(X_0)}{A}.$$

Since A was arbitrary, the proof is complete. ■

Theorem 4.2. *If $\{(X_t, G_t)\}$ is a monotone increasing adaptive RWCE on \mathbb{N} , with G_0 transient, and bounded above by some transient graph $G_\infty = (\mathbb{N}, C_\infty)$ then the walk is transient.*

Proof. Note that G_0 bounds the sequence G_t from below. The potential sequence is

$$F_t(v) = \sum_{j=v}^{\infty} \frac{1}{C_t(j)}$$

i.e. the resistance between v and infinity. Harmonicity and monotonicity hold as above and $F_t(X_t)$ is therefore a super-martingale. G_0 is transient, thus, given $\varepsilon > 0$ there is $v \in \mathbb{N}$ such that $F_0(v) < \varepsilon$. By ellipticity, we may assume that $X_0 = v$ and since $F_t(v)$ is decreasing we have $F_0(X_0) < \varepsilon$.

Let τ be the first time $X_t = 0$, or infinity if the walk never reaches 0. Let p_v be the probability that $\tau < \infty$. Since $F_t(X_t)$ is positive and using the optional stopping theorem we have

$$p_v F_\infty(0) \leq \mathbb{E}(F_\tau(X_\tau)) \leq F_0(X_0)$$

which implies

$$p_v \leq \frac{\varepsilon}{F_\infty(0)}.$$

Since ε was arbitrary, there exists a vertex $v \in \mathbb{N}$ such that $p_v < \frac{1}{2}$. Ellipticity implies that whenever the walk is at 0 it will almost surely visit v at some later time and thereafter it would never visit 0 again with probability $\frac{1}{2}$. Therefore, 0 would be visited only a finite number of times, almost surely. ■

Theorem 4.3. *If $\{(X_n, G_n)\}$ is a monotone decreasing adaptive RWCE on \mathbb{N} , bounded below by some transient graph $G_\infty = (\mathbb{N}, C_\infty)$, then the walk is transient.*

Proof. The potential sequence is

$$F_t(v) = \sum_{j=0}^{v-1} \frac{1}{C_t(j)}$$

i.e. the resistance between 0 and v . Harmonicity and monotonicity hold as above and $F_t(X_t)$ is therefore a sub-martingale. Obviously, this sub-martingale is bounded by 0 and $F_\infty(\infty) = \sum_{j=0}^{\infty} 1/C_\infty(j)$ which is finite.

Assume that the walk starts at $X_0 > 0$ and fix some $v > X_0$. Let τ be the first time the walk hits 0 or v , which, by ellipticity, happens almost surely. Let p_v be the probability that the walk hits v first. By the optional stopping theorem we have

$$F_0(X_0) \leq E(F_\tau(X_\tau)) \leq (1 - p_v)F_\infty(0) + p_v F_\infty(v) \leq p_v F_\infty(\infty)$$

and therefore

$$p_v \geq \frac{F_0(X_0)}{F_\infty(\infty)}.$$

This holds for all $v > X_0$ and thus the probability that the walk never visits 0 is at least $F_0(X_0)/F_\infty(\infty)$. Since F_i is increasing, this bound holds every time the walk returns to X_0 and therefore the walk will visit 0 only finitely many times, almost surely. ■

Theorem 4.4. *If $\{(X_t, G_t)\}$ is a monotone decreasing adaptive RWCE on \mathbb{N} , with G_0 recurrent and bounded below by $G_\infty = (\mathbb{N}, C_\infty)$ with $C_\infty = c C_0$ for some $0 < c$, then the walk is recurrent.*

Proof. Let X_0 be arbitrary. Given X_0 , let n be such that

$$\frac{1}{2} \sum_{j=0}^{n-1} \frac{1}{C_0(j)} \leq \sum_{j=X_0}^{n-1} \frac{1}{C_0(j)}. \quad (1)$$

This is possible since G_0 is recurrent. The potential sequence will be

$$F_t(v) = \sum_{j=v}^{n-1} \frac{1}{C_t(j)}$$

i.e. the resistance between v and n . Then $F_t(X_t)$ is a sub-martingale until the first time that the RW reaches either 0 or n . Let τ be that time and let p_0 be the probability that $X_\tau = 0$. By the optional stopping theorem we have

$$F_0(X_0) \leq \mathbb{E}(F_\tau(X_\tau)) = p_0 \mathbb{E}(F_\tau(0) | X_\tau = 0) + (1 - p_0)0 \leq p_0 F_\infty(0) = \frac{p_0 F_0(0)}{c}$$

Combining the above with (1) we conclude that $p \geq c/2$. This bound holds for any X_0 , i.e. regardless of the current state of the RWCE, the probability of reaching 0 in the future is at least $c/2$. A standard argument now shows that this probability must actually be 1. ■

Unlike the other theorems in this section, the last theorem requires the RWCE to have bounded ratio between G_0 and G_∞ . As the example below shows, this requirement is essential.

Example 4.5. *The RWCE with conductances $C_t(j) = 2^{-j}$ for $j < t$ and $C_t(j) = 1$ otherwise is monotone decreasing, nonadaptive, bounded from above and below by a recurrent graph and is of mixed type.*

Proof. Indeed, with probability $\prod_{t=0}^{\infty} 1/(1 + 2^{-t}) > 0$ the RW will always go to the right and otherwise it will eventually perform a simple random walk on the graph with conductances 2^{-t} , which is recurrent. ■

It is not too difficult to make this example transient. Let $D_t^n(j) = 2^{-j}$ for $n \leq j < t$ and 1 otherwise. So the conductances of the last example are $D_t^0(j)$.

Example 4.6. *There exists an increasing sequence t_n such that the RWCE with conductances $C_t(j) = \prod_{\{n|t_n < t\}} D_{t-t_n}^n(j)$ is monotone decreasing, nonadaptive, bounded from above and below by a recurrent graph and transient.*

Sketch of proof. In example 4.5 we had, in essence, a "wave" of conductances 2^{-j} threatening to carry the RW away. In this example there's a multitude of such waves, each starting one edge further, so that the final conductance of each edge is finite, and each have some fixed positive probability of carrying the RW away. The sequence t_n is chosen to be increasing fast enough, so that the RW would have a fixed positive probability of being to the right of the n -th "wave" when it starts. ■

5 RWCE on trees

Theorems 4.1 and 4.3 can be extended to the case where the underlying graph is a tree. In order to do that first notice that both proofs use the same potential sequence. Second, notice that these functions can be described as follows: Consider the trivial unit flow (on \mathbb{N}) from 0 to infinity and fix the potential at 0 to be 0. Then $F_t(v)$ is the potential of v in G_t . If the underlying graph is a tree, there are many possible choices of flows, each determining a potential. Harmonicity and monotonicity are true for any of these potential sequences, but some care in choosing the right flow is still needed. For general graphs, however, this method fails, since not every flow determines a potential. More precisely, if the graph contains cycles, then there are 2 distinct flows from the source to some vertex and the potential is well defined only when Kirchoff's cycle law is satisfied, which is not necessarily the case.

Theorem 5.1. *If $\{(X_t, G_t)\}$ is a monotone increasing adaptive RWCE on a tree T , bounded above by some recurrent tree $G_\infty = (T, C_\infty)$ then the walk is recurrent.*

Proof. Fix $A > 0$. Since G_∞ is recurrent, there is some $n \in \mathbb{N}$ such that the effective resistance (in G_∞) between the root of the tree (denoted 0) and the outside of the ball of radius n around 0 is at least A , i.e.

$$R = R_{\text{eff}}(0 \leftrightarrow \partial B_n(0); G_\infty) \geq A .$$

Fix such an n and let i be the unit current flow induced by putting a voltage difference of R between 0 and $\partial B_n(0)$ in G_∞ . Let

$$F_t(v) = \sum_e \frac{i(e)}{C_t(e)}$$

where the sum is over all edges e on the (unique) path connecting 0 and v . In words, $F_t(v)$ is the voltage which is induced by the flow i on G_t . Harmonicity follows, as usual, from Kirchhoff's law and monotonicity is trivial since $F_t(v)$ is a fixed positive linear combination of $C_t(e)$'s. Therefore, $F_t(X_t)$ is a super-martingale until the first time $X_t = 0$ or $X_t \in \partial B_n(0)$. From the definition of the flow $F_\infty(v) = R$ for any $v \in \partial B_n(0)$. Since $C_t \leq C_\infty$ we have $F_t(v) \geq R \geq A$ for all $v \in \partial B_n(0)$.

The rest of the proof is the same as in theorem 4.1. Let τ be the first time the walk hits either 0 or $\partial B_n(0)$. Denote by p the probability that the RW hits $\partial B_n(0)$ first. Since $F_t(0) = 0$ for all t , by the optional stopping theorem we have

$$F_0(X_0) \geq \mathbb{E}(F_\tau(X_\tau)) \geq Ap$$

and therefore

$$p \leq \frac{F_0(X_0)}{A} .$$

Since A was arbitrary, the proof is complete. ■

Theorem 5.2. *If $\{(X_t, G_t)\}$ is a monotone decreasing adaptive RWCE on a tree T , bounded below by some transient tree $G_\infty = (T, C_\infty)$, then the walk is transient.*

Proof. To prove transience, it is enough to show that under these conditions there is a vertex u such that such the RWCE, starting from $X_0 = u$, has at least some fixed probability of never returning to 0. Indeed, by ellipticity, every time the walk returns to 0 it visits u with some fixed probability and will therefore return to 0 only finitely many time, almost surely.

Since G_∞ transient, the effective resistance, R between between 0 and infinity is finite, that is, if i is the unit current flow from 0 to infinity then the corresponding potential is bounded by R . Let

$$F_t(v) = \sum_e \frac{i(e)}{C_t(e)}$$

where the sum is over all edges e on the (unique) path connecting 0 and v . This is the same as the previous proof except now $F_t(X_t)$ is a sub-martingale since C_t is decreasing.

Let u be a neighbor of 0 such that the flow from 0 to u is positive and assume that $X_0 = u$. By definition, $F_0(u) = i(0, u)/C_0(0, u)$ is positive too.

The rest of the proof is the same as in theorem 4.3. Let τ be the first time the walk hits either 0 or $\partial B_n(0)$. Denote by p_n the probability that the RW hits $\partial B_n(0)$ first. By the optional stopping theorem we have

$$F_0(X_0) \leq \mathbb{E}(F_\tau(X_\tau)) = p_n \mathbb{E}(F_\tau(X_\tau) | X_\tau \neq 0) \leq p_n R$$

and therefore

$$p_n \geq \frac{F_0(X_0)}{R}.$$

This holds for all n and thus the probability that the walk never visits 0 is at least $F_0(X_0)/R$. Since F_t is increasing, this bound holds every time the walk returns to u and therefore the walk will visit 0 only finitely many times, almost surely. ■

6 RWCE on \mathbb{Z}^2

One could hope that the conclusions of Theorem 5.1 would hold for any monotone RWCE, but unfortunately, this is not true, as the following example of an adaptive RWCE in 2 dimensions shows.

The example we build is a monotone increasing adaptive RWCE on \mathbb{Z}^2 , with $C_0 = 1$ for horizontal edges and $C_0 = 2$ for vertical edges and $C_n \leq 2$ for all n and all edges.

We shall try to mimic the behavior of excited random walk in our model as follows. When the walk reaches a vertex, we will try to give it a push to the right by increasing the conductance of the right edge to 2. If its left neighbor was never visited, this will make the transition probabilities for the next step equal to $\frac{1}{7}$ for the left and $\frac{2}{7}$ for all the other directions which will give a drift to the right. If the left neighbor has already been visited, all transition probabilities will be $\frac{1}{4}$ so the drift would be zero. Call this walk MAW for “Monotone Adaptive Walk” (the MAW is a specific example of a RWCE).

This is obviously quite similar to excited random walk so one is tempted to assume we will get transience, as in [2]. One should be careful, though, because in one dimension a similar attempt to mimic the results of [21] would fail, as Theorem 4.1 shows. So this kind of result is quite sensitive to specific details of the model.

Theorem 6.1. *MAW is transient.*

We present two proofs for Theorem 6.1. The first is based on a theorem of Meshnikov and Popov ([14]) on generalized excited random walks, while the second proof is based on the methods of Benjamini and Wilson ([2]) and on harmonic analysis of random walks in 2 dimensions. The second proof appears in the appendix to this paper. Part of the reason for keeping the second proof is that the methods in it were used and cited in other works (e.g. [6]).

Proof. To show that MAW is transient, we use a result of Menshikov and Popov ([14], Theorem 1.4) which gives bounds on the size of the range of so called “strongly directed” submartingales.

We will show that MAW is such a strongly directed submartingale. Indeed, following definition 1.1 of [14], we choose \mathcal{L} to be \mathbb{R}^2 so that the projection operator $P_{\mathcal{L}}$ is the identity. ℓ is the unit vector in the positive x -direction and $u = 1$. Then, $H_{\ell, \mathcal{L}}^u$ (in the paragraph above definition 1.1 of [14]) is simply the positive x -axis and we see that a submartingale is (u, ℓ, \mathcal{L}) -strongly directed if the drift at any step (conditioned on the history) is in the positive x -direction (including 0). Hence, MAW is (u, ℓ, \mathcal{L}) -strongly directed. It is also uniformly elliptic and has uniformly bounded jumps, thus satisfying the conditions of Theorem 1.4 of [14].

The conclusion of Theorem 1.4 of [14] is that there are constants $\gamma < \frac{1}{2}$ and $C_1, C_2, \delta > 0$ such that

$$\mathbb{P}(R_n < n^{1-\gamma}) \leq C_1 n e^{-C_2 n^\delta}, \quad (2)$$

where R_n denotes the number of distinct vertices visited up to time n . Define $S_{i,j}$ to be the 3×3 square centered at (i, j) , that is, $S_{i,j} = \{i-1, i, i+1\} \times \{j-1, j, j+1\}$. Consider all such squares centered at multiples of 3 - they are all disjoint. Denote by R'_n the number of such squares that are visited by the random walk until time n . Since each such square has only 9 distinct vertices, it follows that $R'_n \geq R_n/9$.

Let $\tau_{i,j}$ be the first hitting time of $S_{i,j}$. Call a square *good* if, during the 4 steps immediately after $\tau_{i,j}$, the walk stays in $S_{i,j}$ and visits $(i+1, j)$, but not (i, j) . It is straightforward to check that no matter where the walk enters $S_{i,j}$, it can hit $(i+1, j)$ in 4 steps, without leaving the square or hitting (i, j) and that each of those 4 steps has transition probability at least $\frac{1}{4}$. Thus, conditioned on the history up to time $\tau_{i,j}$, the probability that the square $S_{i,j}$ is good is at least $\frac{1}{256}$.

Let Q_k be the number of good squares among the first k squares visited (that are centered at multiples of 3). Q_k stochastically dominates a $\text{Binom}(k, 1/256)$ r.v. hence the probability that $Q_k < k/1000$ decays exponentially in k . Combining with (2) we deduce that there exist some $C_3, C_4, \delta_1 > 0$ such that

$$\mathbb{P}(R''_n < n^{1-\gamma}/1000) \leq C_3 e^{-C_4 n^{\delta_1}}, \quad (3)$$

where R''_n is the number of good squares visited by time n .

Let R'''_n be the number of times up to time n that the MAW visits a vertex before visiting its left neighbour. These are exactly the times the MAW takes a step to the right with probability $2/7$ and to the left with probability $1/7$, conditioned on the history until that time. At all other times, the MAW's step is balanced. Note that every time the MAW visits a good square $S_{i,j}$ and gets to $(i+1, j)$ for the first time, its left neighbour has not been visited, thus $R'''_n \geq R''_{n-4}$.

Let X_n be the x -coordinate of the MAW. Applying Azuma's inequality to the martingale $X_n - R'''_n/7$ we get that there exist some $C_5, C_6, \delta_2 > 0$ such that

$$\mathbb{P}(|X_n - R'''_n/7| < n^{1-\gamma}/14000) \leq C_5 e^{-C_6 n^{\delta_2}}. \quad (4)$$

Combining (4) with (3) we deduce that there exist some $C_7, C_8, \delta_3 > 0$ such that

$$\mathbb{P}(X_n < n^{1-\gamma}/14000) \leq C_7 e^{-C_8 n^{\delta_3}}.$$

Summing for all values of n and using the Borel-Cantelli lemma we conclude that $\lim_{n \rightarrow \infty} X_n = \infty$ almost surely and in particular the MAW is transient. ■

7 A conjecture and some open problems

The following conjecture seems the most interesting to us:

Conjecture 7.1. *Theorems 4.1, 4.2, 4.3 and 4.4 are true for nonadaptive RWCE on any graph.*

Remark 7.2: Note that these conjectures claim that there is an essential difference between adaptive and nonadaptive walks. The Theorems in this paper do not provide proof of any such difference. However, as pointed to us by Ben Morris, it seems that using the methods of evolving sets ([15]) it is possible to show that when the graph satisfies an isoperimetric inequality that implies transience (e.g. \mathbb{Z}^d for $d \geq 3$) and the environment is monotone, bounded between two constant multiples of the same conductance function and nonadaptive the RWCE is also transient. On the other hand, an adaptive example similar that in Section 6 can likely be constructed on \mathbb{Z}^d by mimicking the behaviour of excited random walk towards the middle - a walk which gets a bias towards the origin every time it visits a new vertex, for which there is a sketch of proof for recurrence [12]. It seems, however, that proving recurrence of the “excited towards the middle” RWCE is more technically involved than Theorem 6.1.

One could also consider a continuous time version of the RWCE, where the edges are equipped with Poisson clocks with rates equal to their conductance, and the walk jumps over whatever edge rings first. There seems to be an essential difference between the continuous time and discrete time models, which is that for a continuous time random walk on a graph, the stationary measure is always uniform. In fact, it follows from results of Delmotte and Deuschel [5] that for continuous time **nonadaptive** RWCE on \mathbb{Z}^d with conductances bounded above and below by a constant, heat kernel behavior is essentially the same as in \mathbb{Z}^d , and in particular the walk is recurrent if and only if $d \leq 2$. However, such heat kernel estimates, and even questions of recurrence vs. transience, are open for more general graphs (even when requiring monotonicity). We therefore ask:

Question 7.3. *Do Theorems 4.1, 4.2, 4.3 and 4.4 hold for continuous time nonadaptive RWCE on any graph?*

In fact, one could ask the same thing in the discrete time RWCE, by simply requiring all the stationary measures on the graphs G_i to be the same (e.g. by keeping the sum of the conductances at each vertex fixed).

Question 7.4. *Do Theorems 4.1, 4.2, 4.3 and 4.4 hold for nonadaptive RWCE on any graph under the condition that all G_i share the same stationary measure?*

The latter question seems closely related to the results of [1] on RWCE's on finite graphs, where it is shown that contrary to the general case where the cover time may be exponential, for a sequence of graphs with common stationary measures, the cover time is only polynomial.

Note that in the adaptive case, the “fixed” stationary measure plays no role, and in fact one can easily mimic the behavior of discrete time adaptive RWCE with bounded conductances using adaptive continuous RWCE's up to a time change — simply use the same graph, conductances and adaptive rule and change the environment immediately after the process jumps. More precisely, denoting by (Y_i, D_i) the discrete RWCE and by (X_t, C_t) the continuous time RWCE on G , and by T_i the time of the i 'th jump of X_t we get a coupling of the two processes up to time change simply by taking $(Y_i, D_i) = (X_{T_i}, C_{T_i})$.

Acknowledgments

The first draft of this paper was written in 2006, and some version circulated. We wish to thank all those who read and commented on earlier drafts. We thank Amir Dembo and Ruojun Huang for helpful observations regarding continuous time RWCE and for pointing out the reference [5]. We thank the anonymous referees for pointing out [14] and its relevance to Theorem 6.1, as well as other useful suggestions.

The research of G.A. was supported by the Israel Science Foundation grant ISF 1471/11 by a Grant from the GIF, the German-Israeli Foundation for Scientific Research and Development.

References

- [1] Chen Avin, Michal Koucký, and Zvi Lotker. How to explore a fast-changing world (cover time of a simple random walk on evolving graphs). In *Automata, languages and programming*, pages 121–132. Springer, 2008.
- [2] I. Benjamini and D.B. Wilson. Excited random walk. *Electron. Comm. Probab*, 8(9):86–92, 2003.
- [3] Sourav Chatterjee. A new approach to strong embeddings. *Probability Theory and Related Fields*, 152(1-2):231–264, 2012.
- [4] Burgess Davis. Reinforced random walk. *Probability Theory and Related Fields*, 84(2):203–229, 1990.
- [5] T Delmotte and J-D Deuschel. On estimating the derivatives of symmetric diffusions in stationary random environment, with applications to ϕ interface model. *Probability theory and related fields*, 133(3):358–390, 2005.
- [6] Amir Dembo, Ruojun Huang, and Vladas Sidoravicius. Monotone interaction of walk and graph: recurrence versus transience. *Electronic Communications in Probability*, 19:1–12, 2014.
- [7] Amir Dembo, Ruojun Huang, and Vladas Sidoravicius. Walking within growing domains: recurrence versus transience. *Electron. J. Probab*, 19(106):1–20, 2014.

- [8] Rick Durrett, Harry Kesten, and Vlada Limic. Once edge-reinforced random walk on a tree. *Probability theory and related fields*, 122(4):567–592, 2002.
- [9] Uwe Einmahl. Extensions of results of komlós, major, and tusnády to the multivariate case. *Journal of multivariate analysis*, 28(1):20–68, 1989.
- [10] Harry Kesten. Hitting probabilities of random walks on zd. *Stochastic Processes and their Applications*, 25:165–184, 1987.
- [11] János Komlós, Péter Major, and Gábor Tusnády. An approximation of partial sums of independent rv’s, and the sample df. ii. *Probability Theory and Related Fields*, 34(1):33–58, 1976.
- [12] G. Kozma. Excited towards the middle. <http://www.wisdom.weizmann.ac.il/~gadyk/site/center.pdf>.
- [13] Gregory Francis Lawler. *A self-avoiding random walk*. PhD thesis, Princeton., 1979.
- [14] Mikhail Menshikov and Serguei Popov. On range and local time of many-dimensional submartingales. *Journal of theoretical probability*, 27(2):601–617, 2014.
- [15] Ben Morris and Yuval Peres. Evolving sets, mixing and heat kernel bounds. *Probability Theory and Related Fields*, 133(2):245–266, 2005.
- [16] Robin Pemantle. A survey of random processes with reinforcement. *Probab. Surv*, 4(0):1–79, 2007.
- [17] Frank Spitzer. *Principles of random walk*, volume 34. Springer Science & Business Media, 2001.
- [18] Bálint Tóth. The” true” self-avoiding walk with bond repulsion on z: limit theorems. *The Annals of Probability*, pages 1523–1556, 1995.
- [19] MR Vervoort. Reinforced random walks. *preparation, draft version available on http://staff. science.wa. nl/vervoort*, 2002.
- [20] Andrei Yu Zaitsev. Multidimensional version of the results of komlós, major and tusnády for vectors with finite exponential moments. *ESAIM: Probability and Statistics*, 2:41–108, 1998.
- [21] M.P.W. Zerner. Multi-excited random walks on integers. *Probability Theory and Related Fields*, 133(1):98–122, 2005.

Appendix: second proof of Theorem 6.1

From a “calulatory” point of view, [2] reduces to the fact that simple random walk starting from $(0, 0)$ has a probability of $n^{-1/4}$ to avoid hitting the right half line for the first n steps (Kesten’s lemma: see [10]). This $n^{-1/4}$ factor manifests itself in the fact that, finally, they prove a $n^{3/4}$ drift (up to logarithmic factors). In their settings the probabilities for going up or down never change — the effect of the drift is only to move weight around between the left and right probabilities. We do not know how to mimic this particular detail in the settings of monotone adaptive conductances so we will need to work without it, and this would complicate the geometric settings somewhat. Below we work out our replacement for Kesten’s lemma. Hence for a while we will only develop properties of simple random walk. The impatient can jump to lemma 7.13 to see how this is used. Below $\varepsilon \in (0, \frac{1}{2})$ is some parameter that will be kept fixed throughout. The notation $x \approx y$ denotes that $cX \leq y \leq Cx$ for some absolute constants $c, C > 0$.

Lemma 7.5. *Let E_1 be the event that a random walk starting from 0 will avoid hitting the point $(-1, 0)$ for the first $\lceil n^\varepsilon \rceil$ steps. Then*

$$\mathbb{P}(E_1) \approx \frac{1}{\log n}.$$

This is a well known fact. See e.g. [17].

Lemma 7.6. *Let W be Brownian motion starting from 0. Let K be an infinite cone with opening $\theta \in [0, \pi]$ and tip v , and assume $|v| \leq 2d(0, K)$. Let $t > 4|v|^2$ and denote $\mu = \sqrt{t}/|v|$. Then*

$$\mathbb{P}(W[0, t] \cap K = \emptyset) \geq \frac{c\mu^{-\pi/(2\pi-\theta)}}{\sqrt{\log \mu}} \quad (5)$$

$$\mathbb{P}(\{W[0, t] \cap K = \emptyset\} \cap |W(t)_2| < \delta\sqrt{t}) \leq C\delta\mu^{-\pi(2\pi-\theta)}\sqrt{\log \mu}. \quad (6)$$

for any $\delta < 1$.

We remark that both $\sqrt{\log}$ factors above can be removed without much difficulty. See some additional blurbs on this in the remark on page 20 below.

Proof. Denote $\xi = \pi/(2\pi - \theta)$. Let T_r be the stopping time of W on $\partial B(0, r)$. Applying the map $z \rightarrow z^\xi$ that maps the cone to a half-space, and using conformal invariance (and a few calculations) we get that

$$\mathbb{P}(W[0, T_r] \cap K = \emptyset) \approx \left(\frac{|v|}{r}\right)^\xi.$$

On the other hand, $\mathbb{P}(t > T_{C\sqrt{t\log \mu}}) \leq \mu^{-1}$ for some C sufficiently large, so

$$\mathbb{P}(W[0, t] \cap K = \emptyset) \geq \mathbb{P}(W[0, T_{C\sqrt{t\log \mu}}] \cap K = \emptyset) - \mu^{-1} \geq c \left(\frac{1}{\mu\sqrt{\log \mu}}\right)^\xi.$$

For the other part, first notice that $\mathbb{P}(\frac{1}{2}t < T_{c\sqrt{t\log \mu}}) \leq \mu^{-1}$ which gives similarly that $\mathbb{P}(W[0, \frac{1}{2}t] \cap K = \emptyset) \leq c(\sqrt{\log \mu}/\mu)^\xi$. After $\frac{1}{2}t$ we have that for any $x \in \mathbb{R}^2$ that Brownian motion W starting from x has probability $\leq C\delta$ to be in the strip $\{(x, y) : |y| < \delta\sqrt{t}\}$. These two facts prove (6). \blacksquare

Lemma 7.7. *For $m \in (n^{2\varepsilon}, n]$ let $E_2(m, n)$ be the event that a random walk R starting from 0, satisfies*

1. E_1
2. $R[\lceil n^\varepsilon \rceil, m] \cap F = \emptyset$ where F is the funnel

$$F = \{(x, y) : x \geq -1, |y| \leq \log^3 n \sqrt{x+2}\}. \quad (7)$$

Then

$$\begin{aligned} \mathbb{P}(E_2(m, n)) &\geq \left(\frac{n^\varepsilon}{m}\right)^{1/4+o(1)}, \\ \mathbb{P}(E_2(m, n) \cap \{|R(m)_2| < \delta\sqrt{m}\}) &\leq \delta \left(\frac{n^\varepsilon}{m}\right)^{1/4+o(1)} \end{aligned}$$

if only $\delta\sqrt{m} \geq 1$.

Here and below $o(1)$ stands for an entry that goes to 0 as $n \rightarrow \infty$ uniformly in $m > n^{2\varepsilon}$.

Proof. Denote $v = R(\lceil n^\varepsilon \rceil)$. The first ingredient is Hungarian coupling [9, Theorem 4], see also [3, 20, 11], which gives that we can couple random walk starting from v to Brownian motion W also starting from v such that with probability $\geq 1 - n^{-10}$ we have $|R(t) - W(t)| \leq C_1 \log^2 t$. We therefore find two cones K^\pm satisfying $K^- + B(0, C_1 \log^2 n) \subset F$ and $F + B(0, C_1 \log^2 n) \subset K^+$. Specifically we choose

$$K^- = \{(x, 0) : x \geq C_1 \log^2 n\}$$

$$K^+ = \left\{ (x, y) : x \geq -n^{\varepsilon/4}, |y| \leq \frac{\log^3 n}{n^{\varepsilon/8}} (x + n^{\varepsilon/4}) \right\}$$

and the inclusion conditions will be satisfied for n sufficiently large.

Next we want to estimate the distance of v from K^+ . With probability $> 1 - C \log^{-2} n$ we have that $d(v, K^+) > n^{\varepsilon/2} \log^{-2} n$. To see this fix some $\lambda = 1, 2, \dots$ and examine the annulus $A := n^{\varepsilon/2} (B(0, \lambda) \setminus B(0, \lambda - 1))$. For every $w \in A$ one has that $\mathbb{P}(v = w) \leq C n^{-\varepsilon} e^{-\lambda^2}$ while the inflated cone $(K^+ + B(0, n^{\varepsilon/2} \log^{-2} n)) \cap A$ contains $\leq C n^{\varepsilon/2} (\lambda n^{(3/8)\varepsilon} \log^2 n + n^{\varepsilon/2} \log^{-2} n) \leq C \lambda n^\varepsilon \log^{-2} n$ points. Summing over λ we get the estimate for $d(v, K^+)$. Comparing to the probability of E_1 we get for n sufficiently large

$$\mathbb{P}(E_1 \cap \{d(v, K^+) > n^{\varepsilon/2} \log^{-2} n\}) \approx \frac{C}{\log n}. \quad (8)$$

Now we may invoke lemma 7.6 and get that, assuming $d(v, K^+) > n^{\varepsilon/2} \log^{-2} n$,

$$\begin{aligned} \mathbb{P}(R[\lceil n^\varepsilon \rceil, m] \cap F = \emptyset) &\geq \mathbb{P}(W[0, m - \lceil n^\varepsilon \rceil] \cap K^- = \emptyset) \\ &\stackrel{(5)}{\geq} \frac{c}{\sqrt{\log n}} \left(\frac{n^{\varepsilon/2} \log^{-2} n}{\sqrt{m - \lceil n^\varepsilon \rceil}} \right)^{1/4} \geq \left(\frac{n^\varepsilon}{m} \right)^{1/4 + o(1)} \end{aligned}$$

and

$$\begin{aligned} &\mathbb{P}(\{R[\lceil n^\varepsilon \rceil, m] \cap F = \emptyset\} \cap \{|R(m)_2| \leq \delta \sqrt{m}\}) \leq \\ &\leq \mathbb{P}(\{W[0, m - \lceil n^\varepsilon \rceil] \cap K^+ = \emptyset\} \cap \\ &\quad \cap \{|W(m - \lceil n^\varepsilon \rceil)_2| \leq \delta \sqrt{m} + C_1 \log^2 n\}) \leq \\ &\stackrel{(6)}{\leq} C \left(\delta + \frac{C_1 \log^2 n}{\sqrt{m}} \right) \sqrt{\log n} \left(\frac{n^{\varepsilon/2} \log^{-2} n}{\sqrt{m - \lceil n^\varepsilon \rceil}} \right)^{\pi / (2\pi - n^{-\varepsilon/8} \log^3 n)} \leq \\ &\leq \delta \left(\frac{n^\varepsilon}{m} \right)^{1/4 + o(1)} \end{aligned}$$

Where in the last inequality we used $\delta \sqrt{m} \geq 1$ to bound $C_1 m^{-1/2} \log^2 n \leq \delta \log^2 n$ and then this log factors can be folded into the $o(1)$ in the exponent like all the other log-s (including the one from E_1). Notice also that we didn't write the negligible probability for the coupling to fail, but it does not affect the result for n sufficiently large. \blacksquare

Lemma 7.8. *Let $E_3(m, n)$, $m \geq n^{2\varepsilon}$ be the event that a random walk R starting from 0 satisfies that*

1. $R[m - \lceil n^\varepsilon \rceil, m]$ avoids $(-1, 0) + R(m)$; and
2. $R[0, m - \lceil n^\varepsilon \rceil]$ avoids $F + R(m)$ where F is the funnel defined in (7).

Then

$$\mathbb{P}(E_3) \geq \left(\frac{n^\varepsilon}{m}\right)^{1/4+o(1)}, \quad (9)$$

$$\mathbb{P}(E_3 \cap \{|R(m)_2| < \delta\sqrt{m}\}) \leq \delta \left(\frac{n^\varepsilon}{m}\right)^{1/4+o(1)}. \quad (10)$$

for any $m \in [n^{2\varepsilon}, n]$.

Proof. This follows immediately from lemma 7.7 and time reversal symmetry. \blacksquare

Following [2] we will call m satisfying $E_3(m, n)$ “tan points” (imagine the sun being at the right infinity, then $R(m)$ gets a tan without (almost) any previous point blocking a whole “tanning funnel”).

Lemma 7.9. *Let $m_1 < m_2$ and $m_2 - m_1 > n^{2\varepsilon}$. Then*

$$\mathbb{P}(E_3(m_1, n) \cap E_3(m_2, n)) \leq \mathbb{P}(E_3(m_1, n))\mathbb{P}(E_3(m_2 - m_1, n)).$$

Proof. One only needs to notice that it is easier for $R(m_2)$ to be a tan point *with respect to the walk starting from $R(m_1)$* then to be a regular tan point. In other words, if $S(i) := R(m_1 + i) - R(m_1)$ then S is a random walk starting from 0; and if E^* is the event that $m_2 - m_1$ is a tan point for S ; then $E_3(m_2, n) \subset E^*$. \blacksquare

Lemma 7.10. *With probability $> 1 - Cn^{-2}$ there are at least $n^{3/4-(7/4)\varepsilon+o(1)}$ n^ε -separated tan points up to time n .*

Proof. Let $h = \lceil n^{1/2} \log^{-2} n \rceil$ and $l = \lfloor h^2 \log^{-1} n / \lceil n^{2\varepsilon} \rceil \rfloor$. Let T_i be the stopping times on the double line $\{(x, \pm ih) : x \in \mathbb{R}\}$. For all $i \in \mathbb{N}$ and $j = l, l+1, \dots, 2l$ let $Y_{i,j}$ be the event that $T_i + j \lceil n^{2\varepsilon} \rceil$ is a tan point with respect to T_i . Define $X_i := \#\{j : Y_{i,j}\}$. The first step is to show that

$$\mathbb{P}(X_i > n^{3/4-(7/4)\varepsilon+o(1)}) > c. \quad (11)$$

We use second moment methods. First by (9) we have

$$\mathbb{E}(X_i) \geq (l+1) \cdot \left(\frac{n^\varepsilon}{2l \lceil n^{2\varepsilon} \rceil}\right)^{1/4+o(1)} \geq n^{3/4-(7/4)\varepsilon+o(1)}.$$

For the second moment write

$$\mathbb{E}(X_i^2) = \sum_j \mathbb{P}(Y_j) + \sum_{j < k} 2\mathbb{P}(Y_j \cap Y_k)$$

and by lemma 7.9

$$\leq \mathbb{E}X_i + \sum_{j < k} 2\mathbb{P}(Y_j)\mathbb{P}(Y_{k-j}) \leq \mathbb{E}X_i + 2(\mathbb{E}X_i)^2.$$

By the well known inequality $\mathbb{P}(X \geq \frac{1}{2}\mathbb{E}X) \geq (\mathbb{E}X)^2/4\mathbb{E}(X^2)$ we get for n sufficiently large

$$\mathbb{P}(X_i > n^{3/4-(7/4)\varepsilon+o(1)}) \geq \frac{1}{12}.$$

Next we define $Y_{i,j}^*$ to be the event

$$Y_{i,j} \cap \left\{ |R(T_i + j \lceil n^{2\varepsilon} \rceil)_2| > ih + n^{1/4} \log^4 n \right\}.$$

And $X_i^* = \#\{j : Y_{i,j}^*\}$. We shall now estimate X_i^* under the assumption that $R(T_i)_2 = ih$ (rather than $-ih$) — the other case is symmetric. Examine the event

$$B_{i,j} = Y_{i,j} \cap \left\{ |R(T_i + j \lceil n^{2\varepsilon} \rceil)_2 - ih| \leq n^{1/4} \log^4 n \right\}.$$

By (10) we have (remember the definitions of l and h) that

$$\mathbb{P}(B_{i,j}) \leq \frac{n^{1/4} \log^4 n}{\sqrt{j n^{2\varepsilon}}} \left(\frac{n^\varepsilon}{j \lceil n^{2\varepsilon} \rceil} \right)^{1/4+o(1)} \leq n^{-1/2+\varepsilon/4+o(1)}$$

and summing over j we get $\mathbb{E}(\#\{j : B_{i,j}\}) \leq n^{1/2-(7/4)\varepsilon+o(1)}$. Estimating with Markov's inequality we see that the $B_{i,j}$ are negligible and then

$$\mathbb{P}(\#\{Y_{i,j} \setminus B_{i,j}\} > n^{3/4-(7/4)\varepsilon+o(1)}) > c.$$

Now $Y_{i,j} \setminus B_{i,j}$ is equal to $Y_{i,j}^* \cup \{\text{its symmetric image}\}$. Therefore we get

$$\mathbb{P}(X_i^* > N) = \mathbb{P}(\#\{j : Y_{i,j} \setminus (B_{i,j} \cup Y_{i,j}^*)\} > N) \quad \forall N$$

And hence $\mathbb{P}(X_i^* > N) \geq \frac{1}{2} \mathbb{P}(\#\{Y_{i,j} \setminus B_{i,j}\} > 2N)$.

Finally we define the event

$$G_i = \{X_i^* > n^{3/4-(7/4)\varepsilon+o(1)}\} \cap \{T_{i+1} - T_i > h^2 \log^{-1} n\}.$$

Then since $\mathbb{P}(T_{i+1} - T_i \leq h^2 \log^{-1} n) < e^{-c \log^2 n}$ we get that $\mathbb{P}(G_i) > c$.

However, G_i is independent of $R(T_i)$, including of whether it is in the line $\mathbb{R} \times \{ih\}$ or $\mathbb{R} \times \{-ih\}$, since $Y_{i,j}^*$ and the rest of the conditions are invariant with respect to translations in the x direction and reflections through the x axis. Therefore (since G_i does not examine the walk beyond T_{i+1}) the G_i are independent events. Further, with probability $> 1 - n^{-2}$ we have $\max_{m \leq n} |R(m)_2| \geq c\sqrt{n/\log n}$ and then there are at least $c \log^{3/2} n$ different i -s for which $T_{i+1} < n$. This shows that with probability $> 1 - Cn^{-2}$ at least one of the G_i -s occurred. Further, with the same probability we may also assume

$$\max_{m \leq n} |R(m)_1| \leq \sqrt{n \log n}. \tag{12}$$

This finishes the lemma. Indeed,

$$(12) \cap Y_{i,j}^* \Rightarrow T_i + j \lceil n^{2\varepsilon} \rceil \text{ is a tan point}$$

since the funnel $F + R(T_i + j \lceil n^{2\varepsilon} \rceil)$ intersects the band $\{(x, y) : |y| \leq ih\}$ only for $|x| > c\sqrt{n} \log^2 n$ and R does not go so far. Hence the $T_i + j \lceil n^{2\varepsilon} \rceil$ -s for which $Y_{i,j}^*$ occurred are n^ε -separated tan points and the lemma is proved. \blacksquare

Remark 7.11: Lemma 7.9 alleviates most of the agony usually associated with second moment methods. However it is by no means necessary. There are at least two additional paths one might take to prove the result i.e. lemma 7.10:

- It is not very difficult to get rid of all the log factors we have so lavishly neglected and show explicitly that $\mathbb{P}(E_3) \approx \frac{1}{\log n} \left(\frac{n^\varepsilon}{m}\right)^{1/4}$ — the $1/\log n$ comes from E_1 and is the only log that represents a real phenomenon. Further one can show that $\mathbb{P}(E_3(m, n) \cap \{R(m)_2 > \sqrt{m}\}) \approx \frac{1}{\log n} \left(\frac{n^\varepsilon}{m}\right)^{1/4}$ which would allow to estimate X_i^* without going through the symmetry argument.
- Alternatively, if the second moment methods only show that $\mathbb{P}(G_i) > c \log^{-10} n$ one can simply take $h = \sqrt{n} \log^{-12} n$. This will give $\log^{11+1/2} n$ possible i -s and one of them would satisfy E_i .

This concludes what we need to know about simple random walk. The next step is to couple MAW and SRW. We shall do so in the natural way: if the MAW is in a vertex whose left neighbor was visited in the past (NV-vertex), make the MAW and the SRW walk together. Otherwise, do as follows:

- With probability $\frac{1}{7}$ they both walk to the left
- With probability $\frac{1}{4}$ they both walk up, another $\frac{1}{4}$ for right, and another for down.
- With probability $\frac{1}{28}$ the SRW walks left and the MAW walk up, etc.

Denoting by R the SRW and by E the MAW we get that $E(n) - R(n)$ changes only when E is in a non-NV vertex. $(E(n) - R(n))_1$ only increases while $(E(n) - R(n))_2$ performs a random walk at these times.

Lemma 7.12. *Let R and E be an SRW and an MAW coupled as above. Let $D(k, l) = E(l) - R(l) - E(k) + R(k)$. Let n be some number. Then*

$$\mathbb{P}\left(\exists k < l < n : |D(k, l)_2| \geq C \log n \sqrt{D(k, l)_1 + 1}\right) \leq n^{-1}$$

for some C sufficiently large.

Proof. Fix some $k < l$. For an integer K let $N(K)$ be the event that exactly K non-NV vertices were visited by E between k and l . It is easy to see that

$$\mathbb{P}(N(K), D(k, l)_1 + 1 < \frac{1}{\lambda} K) \leq C e^{-c\lambda} \quad \forall \lambda > 0, \forall K \quad (13)$$

so for some C_1 sufficiently large, setting $\lambda = C_1 \log(nK)$ will ensure that the probability is $\leq \frac{1}{10} (nK)^{-3}$. Denote this event by $B_1(K)$. Next we note that

$$\mathbb{P}(N(K), |D(k, l)_2| > \lambda \sqrt{K}) \leq C e^{-c\lambda^2} \quad \forall \lambda > 0 \quad (14)$$

and setting $\lambda = C_2 \sqrt{\log(nK)}$ for some C_2 sufficiently large will ensure that the probability is $\leq \frac{1}{10} (nK)^{-3}$. Denote this event by $B_2(K)$. We get that

$$\mathbb{P}\left(\bigcup_{K=1}^{\infty} N(K) \cup B_1(K) \cup B_2(K)\right) \leq \frac{1}{n^3}.$$

However, if this event did not happen then (13) gives that the number of non-NV vertices is smaller than $C \log n (D(k, l)_1 + 1)$ and with (14) we get $|D(k, l)_2| \leq C \log n \sqrt{D(k, l)_1 + 1}$. Summing over k and l proves the lemma. ■

Lemma 7.13. *Let R and E be an SRW and an MAW coupled as above. Assume the event of lemma 7.12 did not happen. Let $m < n$ be a tan point of R . Then at least one of $[m - n^\varepsilon, m]$ is a non-NV point of E .*

Proof. If all of $[m - n^\varepsilon, m - 1]$ were NV points of E , then $E(l) - R(l)$ did not change throughout this time. Hence the first condition in the definition of a tan point, that $R(l)$ did not visit the left neighbor of $R(m)$, ensures that $E(l)$ did not visit the left neighbor of $E(m)$. So this period is secured. Examine now the time $[0, m - n^\varepsilon]$. If for some $l \in [0, m - n^\varepsilon]$ we have that $E(l)$ is the left neighbor of $E(m)$, then $R(l) = \text{left neighbor of } R(m) + D(l, m)$. But we assumed (this is the event of lemma 7.12) that $|D(k, l)_2| \leq C \log n \sqrt{D(k, l)_1 + 1}$. Hence (if n is sufficiently large), $R(l) \in R(m) + F$, in contradiction to the second condition in the definition of a tan point. ■

Proof of Theorem 6.1. Couple the MAW E to a SRW R as above. Examine the first n steps of both. By lemma 7.10 there are (with probability $> 1 - C/n$) $n^{3/4+o(1)}$ tan points $m_i < n$ which are separated i.e. $|m_i - m_j| > n^\varepsilon$ for all $i \neq j$. By lemma 7.13 this shows that there are at least so many visits of E to non-NV vertices. This shows that with probability $> 1 - C/n$ that $(E(n) - R(n))_1 > n^{3/4+o(1)}$. This shows that with probability 1, $E(2^n) - R(2^n) > 2^{3n/4+o(n)}$. Since $(E - R)_1$ is monotone we get $E(n) - R(n) > n^{3/4+o(1)}$ with probability 1. Hence E is transient. ■