On a partial theta function and its spectrum

Vladimir Petrov Kostov

Université de Nice, Laboratoire de Mathématiques, Parc Valrose, 06108 Nice Cedex 2, France, e-mail: kostov@math.unice.fr

Abstract

The bivariate series $\theta(q, x) := \sum_{j=0}^{\infty} q^{j(j+1)/2} x^j$ defines a partial theta function. For fixed $q(|q| < 1)$, $\theta(q,.)$ is an entire function. For $q \in (-1,0)$ the function $\theta(q,.)$ has infinitely many negative and infinitely many positive real zeros. There exists a sequence $\{\bar{q}_i\}$ of values of q tending to -1^+ such that $\theta(\bar{q}_k,.)$ has a double real zero \bar{y}_k (the rest of its real zeros being simple). For k odd (resp. for k even) $\theta(\bar{q}_k,.)$ has a local minimum at \bar{y}_k and \bar{y}_k is the rightmost of the real negative zeros of $\theta(\bar{q}_k,.)$ (resp. $\theta(\bar{q}_k,.)$) has a local maximum at \bar{y}_k and for k sufficiently large \bar{y}_k is the second from the left of the real negative zeros of $\theta(\bar{q}_k,.)$. For k sufficiently large one has $-1 < \bar{q}_{k+1} < \bar{q}_k < 0$. One has $\bar{q}_k = 1 - (\pi/8k) + o(1/k)$ and $|\bar{y}_k| \to e^{\pi/2} = 4.810477382...$

AMS classification: 26A06

Keywords: partial theta function; spectrum

1 Introduction

The bivariate series $\theta(q, x) := \sum_{j=0}^{\infty} q^{j(j+1)/2} x^j$ defines an entire function in x for every fixed q from the open unit disk. This function is called a *partial theta function* because $\theta(q^2, x/q)$ $\sum_{j=0}^{\infty} q^{j^2} x^j$ whereas the Jacobi theta function is defined by the same series, but when summation is performed from $-\infty$ to ∞ (i.e. when summation is not partial).

There are several domains in which the function θ finds applications: in the theory of (mock) modular forms (see [\[3\]](#page-12-0)), in statistical physics and combinatorics (see [\[18\]](#page-13-0)), in asymptotic analysis (see $[2]$) and in Ramanujan type q-series (see $[19]$). Recently it has been considered in the context of problems about hyperbolic polynomials (i.e. real polynomials having all their zeros real, see [\[4\]](#page-12-2), [\[16\]](#page-13-2), [\[5\]](#page-12-3), [\[15\]](#page-13-3), [\[6\]](#page-12-4), [\[13\]](#page-13-4) and [\[9\]](#page-12-5)). These problems have been studied earlier by Hardy, Petrovitch and Hutchinson (see [\[4\]](#page-12-2), [\[5\]](#page-12-3) and [\[16\]](#page-13-2)). For more information about θ , see also [\[1\]](#page-12-6).

For $q \in \mathbb{C}$, $|q| \leq 0.108$, the function $\theta(q,.)$ has no multiple zeros, see [\[11\]](#page-12-7). For $q \in [0,1)$ the function θ has been studied in [\[13\]](#page-13-4), [\[8\]](#page-12-8), [\[9\]](#page-12-5) and [\[10\]](#page-12-9). The results are summarized in the following theorem:

Theorem 1. (1) For any $q \in (0,1)$ the function $\theta(q,.)$ has infinitely many negative zeros.

(2) There exists a sequence of values of q (denoted by $0 < \tilde{q}_1 < \tilde{q}_2 < \cdots$) tending to 1^{-} such that $\theta(\tilde{q}_k,.)$ has a double negative zero y_k which is the rightmost of its real zeros and which is a local minimum of $\theta(q,.)$. One has $\tilde{q}_1 = 0.3092493386...$

(3) For the remaining values of $q \in [0,1)$ the function $\theta(q,.)$ has no multiple real zero.

(4) For $q \in (\tilde{q}_k, \tilde{q}_{k+1})$ (we set $\tilde{q}_0 = 0$) the function $\theta(q,.)$ has exactly k complex conjugate pairs of zeros counted with multiplicity.

(5) One has $\tilde{q}_k = 1 - (\pi/2k) + o(1/k)$ and $y_k \to -e^{\pi} = -23.1407 \ldots$

Definition 2. A value of $q \in \mathbb{C}$, $|q| < 1$, is said to belong to the *spectrum* of θ if $\theta(q,.)$ has a multiple zero.

In the present paper we consider the function θ in the case when $q \in (-1, 0]$. In order to use the results about the case $q \in [0, 1)$ one can notice the following fact. For $q \in (-1, 0]$ set $v := -q$. Then

$$
\theta(q, x) = \theta(-v, x) = \theta(v^4, -x^2/v) - vx\theta(v^4, -vx^2) . \tag{1}
$$

We prove the analog of the above theorem. The following three theorems are proved respectively in Sections [3,](#page-5-0) [4](#page-5-1) and [5.](#page-11-0)

Theorem 3. For any $q \in (-1,0)$ the function $\theta(q,.)$ has infinitely many negative and infinitely many positive real zeros.

Theorem 4. (1) There exists a sequence of values of q (denoted by \bar{q}_j) tending to -1^+ such that $\theta(\bar{q}_k,.)$ has a double real zero \bar{y}_k (the rest of its real zeros being simple). For the remaining values of $q \in (-1,0)$ the function $\theta(q,.)$ has no multiple real zero.

(2) For k odd (resp. for k even) one has $\bar{y}_k < 0$, $\theta(\bar{q}_k,.)$ has a local minimum at \bar{y}_k and \bar{y}_k is the rightmost of the real negative zeros of $\theta(\bar{q}_k,.)$ (resp. $\bar{y}_k > 0$, $\theta(\bar{q}_k,.)$ has a local maximum at \bar{y}_k and for k sufficiently large \bar{y}_k is the leftmost but one (second from the left) of the real negative zeros of $\theta(\bar{q}_k, .)$).

(3) For k sufficiently large one has $-1 < \bar{q}_{k+1} < \bar{q}_k < 0$.

(4) For k sufficiently large and for $q \in (\bar{q}_{k+1}, \bar{q}_k)$ the function $\theta(q,.)$ has exactly k complex conjugate pairs of zeros counted with multiplicity.

Remark 5. Numerical experience confirms the conjecture that parts (2), (3) and (4) of the theorem are true for any $k \in \mathbb{N}$. Proposition [14](#page-6-0) clarifies part (2) of the theorem.

Theorem 6. One has $\bar{q}_k = 1 - (\pi/8k) + o(1/k)$ and $|\bar{y}_k| \to e^{\pi/2} = 4.810477382...$

Remarks 7. (1) Theorems [1](#page-0-0) and [4](#page-1-0) do not tell whether there are values of $q \in (-1, 1)$ for which $\theta(q,.)$ has a multiple complex conjugate pair of zeros.

(2) It would be interesting to know whether the sequences $\{y_k\}$ and $\{\bar{y}_{2k-1}\}$ are monotone decreasing and $\{\bar{y}_{2k}\}\$ is monotone increasing. This is true for at least the five first terms of each sequence.

(3) It would be interesting to know whether there are complex non-real values of q of the open unit disk belonging to the spectrum of θ and (as suggested by A. Sokal) whether $|\tilde{q}_1|$ is the smallest of the moduli of the spectral values.

(4) The following statement is formulated and proved in [\[7\]](#page-12-10):

The sum of the series $\sum_{j=0}^{\infty} q^{j(j+1)/2} x^j$ (considered for $q \in (0,1)$ and $x \in \mathbb{C}$) tends to $1/(1-x)$ (for x fixed and as $q \rightarrow 1^-$) exactly when x belongs to the interior of the closed Jordan curve ${e^{|s|+is}, s \in [-\pi, \pi]}$.

This statement and equation [\(1\)](#page-1-1) imply that as $q \to -1^+$, $\theta(q, x) \to (1-x)/(1+x^2)$ for $x \in (-e^{\pi/2}, e^{\pi/2}), e^{\pi/2} = 4.810477381...$ Notice that the radius of convergence of the Taylor series at 0 of the function $(1-x)/(1+x^2)$ equals 1.

(5) On Fig. [1](#page-3-0) and [2](#page-4-0) we show the graphs of $\theta(\bar{q}_k,.)$ for $k = 1, ..., 8$. The ones for $k = 1, 2$, 5 and 6 are shown in black, the others are drawn in grey. One can notice by looking at Fig. [2](#page-4-0) that for $x \in [-2.5, 2.5]$ the graphs of $\theta(\bar{q}_k,.)$ for $k \geq 5$ are hardly distinguishable from the one of $(1-x)/(1+x^2)$.

(6) The approximative values of \bar{q}_k and \bar{y}_k for $k = 1, 2, \ldots 8$ are:

2 Some facts about θ

This section contains properties of the function θ , known or proved in [\[9\]](#page-12-5). When a property is valid for all q from the unit disk or for all $q \in (-1,1)$, we write $\theta(q, x)$. When a property holds true only for $q \in [0, 1)$ or only for $q \in (-1, 0]$, we write $\theta(v, x)$ or $\theta(-v, x)$, where $v \in [0, 1)$.

Theorem 8. (1) The function θ satisfies the following functional equation:

$$
\theta(q, x) = 1 + qx\theta(q, qx) \tag{2}
$$

and the following differential equation:

$$
2q\partial\theta/\partial q = 2x\partial\theta/\partial x + x^2\partial^2\theta/\partial x^2 = x\partial^2(x\theta)/\partial x^2
$$
\n(3)

(2) For $k \in \mathbb{N}$ one has $\theta(v, -v^{-k}) \in (0, v^{k}).$

(3) In the following two situations the two conditions $sgn(\theta(v, -v^{-k-1/2})) = (-1)^k$ and $|\theta(v, -v^{-k-1/2})| > 1$ hold true:

(i) For $k \in \mathbb{N}$ and $v > 0$ small enough;

(ii) For any $v \in (0,1)$ fixed and for $k \in \mathbb{N}$ large enough.

The real entire function $\psi(z)$ is said to belong to the Laguerre-Pólya class $\mathcal{L} - \mathcal{P}$ if it can be represented as

$$
\psi(x) = cx^m e^{-\alpha x^2 + \beta x} \prod_{k=1}^{\omega} (1 + x/x_k) e^{-x/x_k},
$$

where ω is a natural number or infinity, c, β and x_k are real, $\alpha \geq 0$, m is a nonnegative integer and $\sum x_k^{-2} < \infty$. Similarly, the real entire function $\psi_*(x)$ is a function of type I in the Laguerre-Pólya class, written $\psi_* \in \mathcal{L} - \mathcal{PI}$, if $\psi_*(x)$ or $\psi_*(-x)$ can be represented in the form

$$
\psi_*(x) = cx^m e^{\sigma x} \prod_{k=1}^{\omega} (1 + x/x_k),\tag{4}
$$

where c and σ are real, $\sigma \geq 0$, m is a nonnegative integer, $x_k > 0$, and $\sum 1/x_k < \infty$. It is clear that $\mathcal{L} - \mathcal{PI} \subset \mathcal{L} - \mathcal{P}$. The functions in $\mathcal{L} - \mathcal{P}$, and only these, are uniform limits, on compact subsets of \mathbb{C} , of hyperbolic polynomials (see, for example, Levin [\[14,](#page-13-5) Chapter 8]). Similarly, $\psi \in \mathcal{L} - \mathcal{PI}$ if and only if ψ is a uniform limit on the compact sets of the complex plane of polynomials whose zeros are real and are either all positive, or all negative. Thus, the classes $\mathcal{L} - \mathcal{P}$ and $\mathcal{L} - \mathcal{PI}$ are closed under differentiation; that is, if $\psi \in \mathcal{L} - \mathcal{P}$, then $\psi^{(\nu)} \in \mathcal{L} - \mathcal{P}$ for every $\nu \in \mathbb{N}$ and similarly, if $\psi \in \mathcal{L} - \mathcal{PI}$, then $\psi^{(\nu)} \in \mathcal{L} - \mathcal{PI}$. Pólya and Schur [\[17\]](#page-13-6) proved that if

$$
\psi(x) = \sum_{k=0}^{\infty} \gamma_k \frac{x^k}{k!}
$$
\n(5)

Figure 1: The graphs of the functions $\theta(\bar q_k,.)$ for $k=1,\,2,\,3$ and 4.

Figure 2: The graphs of the functions $\theta(\bar q_k,.)$ for $k=5,\,6,\,7$ and 8.

belongs to $\mathcal{L} - \mathcal{P}$ and its Maclaurin coefficients $\gamma_k = \psi^{(k)}(0)$ are all nonnegative, then $\psi \in$ $\mathcal{L} - \mathcal{PI}.$

The following theorem is the basic result contained in [\[12\]](#page-13-7):

Theorem 9. (1) For any fixed $q \in \mathbb{C}^*$, $|q| < 1$, and for k sufficiently large, the function $\theta(q,.)$ has a zero ζ_k close to $-q^{-k}$ (in the sense that $|\zeta_k + q^{-k}| \to 0$ as $k \to \infty$). These are all but finitely-many of the zeros of θ .

(2) For any $q \in \mathbb{C}^*$, $|q| < 1$, one has $\theta(q, x) = \prod_k(1 + x/x_k)$, where $-x_k$ are the zeros of θ counted with multiplicity.

(3) For $q \in (\tilde{q}_i, \tilde{q}_{i+1}]$ the function $\theta(q,.)$ is a product of a degree 2j real polynomial without real roots and a function of the Laguerre-Pólya class $\mathcal{L} - \mathcal{PI}$. Their respective forms are $\prod_{k=1}^{2j} (1+x/\eta_k)$ and $\prod_k (1+x/\xi_k)$, where $-\eta_k$ and $-\xi_k$ are the complex and the real zeros of θ counted with multiplicity.

(4) For any fixed $q \in \mathbb{C}^*$, $|q| < 1$, the function $\theta(q,.)$ has at most finitely-many multiple zeros.

(5) For any $q \in (-1,0)$ the function $\theta(q,.)$ is a product of the form $R(q,.)\Lambda(q,.)$, where $R = \prod_{k=1}^{2j} (1 + x/\tilde{\eta}_k)$ is a real polynomial with constant term 1 and without real zeros and $\Lambda = \prod_k (1+x/\tilde\xi_k)$, $\tilde\xi_k \in \mathbb R^*$, is a function of the Laguerre-Pólya class $\mathcal L - \mathcal P$. One has $\tilde\xi_k\tilde\xi_{k+1} < 0$. The sequence $\{|\xi_k|\}$ is monotone increasing for k large enough.

3 Proof of Theorem [3](#page-1-2)

One can use equation [\(1\)](#page-1-1). By part (3) of Theorem [8](#page-2-0) with v^4 for v, for each $v \in (0,1)$ fixed and for k large enough, if $-x^2/v = -(v^4)^{-k-1/2}$ (i.e. if $|x| = v^{-2k-1/2}$), then $|\theta(v^4, -x^2/v)| >$ 1 and $sgn(\theta(v^4, -x^2/v)) = (-1)^k$. At the same time part (2) of Theorem [8](#page-2-0) implies that for $-vx^2 = -(v^4)^{-k}$ (i.e. again for $|x| = v^{-2k-1/2}$) one has $\theta(v^4, -vx^2) \in (0, v^{4k})$ hence $|vx\theta(v^4, -vx^2)| < v^{2k+1/2} < 1$. This means that for $v \in (0,1)$ fixed and for k large enough the equality $sgn(\theta(v^4, -x^2/v)) = (-1)^k$ holds, i.e. there is a zero of θ on each interval of the form $(-v^{-2k-1/2}, -v^{-2k+1/2})$ and $(v^{-2k+1/2}, v^{-2k-1/2})$.

4 Proof of Theorem [4](#page-1-0)

4.1 Properties of the zeros of θ

The present subsection contains some preliminary information about the zeros of θ .

Lemma 10. For $q \in [-0.108, 0)$ all zeros of $\theta(q,.)$ are real and distinct.

Proof. Indeed, it is shown in [\[11\]](#page-12-7) that for $q \in \mathbb{C}$, $|q| \leq 0.108$, the zeros of θ are of the form $-q^{-j}\Delta_j$, $\Delta_j \in [0.2118, 1.7882]$. This implies (see [\[11\]](#page-12-7)) that the moduli of all zeros are distinct for $|q| \leq 0.108$. When q is real, then as all coefficients of θ are real, each of its zeros is either real or belongs to a complex conjugate pair. The moduli of the zeros being distinct the zeros are all real and distinct. \Box

Notation 11. We denote by $0 < x_1 < x_3 < \cdots$ the positive and by $\cdots < x_4 < x_2 < 0$ the negative zeros of θ . For $q \in [-0.108, 0)$ this notation is in line with the fact that x_j is close to $-q^{-j}$.

Remarks 12. (1) The quantities Δ_i are constructed in [\[11\]](#page-12-7) as convergent Taylor series in q of the form $1 + O(q)$.

(2) For $q \in (-1,0)$ the function $\theta(q,.)$ has no zeros in $[-1,0)$. Indeed, this follows from

$$
\theta = 1 + q^3 x^2 + q^{10} x^4 + \dots + qx(1 + q^5 x^2 + q^{14} x^4 + \dots).
$$

Each of the two series is sign-alternating, with positive first term and with decreasing moduli of its terms for $q \in (-1,0), x \in [-1,0)$. Hence their sums are positive; as $qx > 0$, one has $\theta > 0$.

Lemma 13. For $q \in [-0.108, 0]$ the real zeros of θ and their products with q are arranged on the real line as shown on Fig. [3.](#page-6-1)

Proof. The lemma follows from equation [\(2\)](#page-2-1). Indeed,

$$
0 = \theta(q, x_{4k+2}) = 1 + qx_{4k+2}\theta(q, qx_{4k+2}) \quad , \quad x_{4k+2} < 0 \quad \text{and} \quad q < 0 \quad \text{hence} \quad \theta(q, qx_{4k+2}) < 0 \; .
$$

For small values of q the quantity qx_{4k+2} is close to $-q^{-4k-1}$ (see Lemma [10](#page-5-2) and part (1) of Remarks [12\)](#page-6-2), i.e. close to x_{4k+1} and as $\theta(q, qx_{4k+2}) < 0$, one must have $x_{4k+1} < qx_{4k+2} < x_{4k+3}$. By continuity these inequalities hold for all $q < 0$ for which the zeros $x_{4k+1} < x_{4k+3}$ are real and distinct.

In the same way one can justify the disposition of the other points of the form qx_j w.r.t. the \Box points x_j .

Figure 3: The real zeros of θ for $q \in (-1,0)$.

Proposition 14. The function $\theta(q,.)$ with $q \in (-1,0)$ has a zero in the interval $(0,-1/q)$. More precisely, one has $\theta(q, -1/q) < 0$.

Proof. Setting as above $v = -q$ one gets

$$
\theta(-v, 1/v) = -v\Phi(v) + v^3\Psi(v) \quad , \text{ where } \Phi(v) = \sum_{j=0}^{\infty} (-1)^j v^{2j^2+3j} \quad , \quad \Psi(v) = \sum_{j=0}^{\infty} (-1)^j v^{2j^2+5j} \ .
$$

Further we use some results of [\[9\]](#page-12-5). Consider the functions $\varphi_k(\tau) := \sum_{j=0}^{\infty} (-1)^j \tau^{kj+j(j-1)/2} =$ $\theta(\tau, -\tau^{k-1})$ and $\xi_k(\tau) := 1/(1+\tau^k), \tau \in [0,1), k > 0.$

Proposition 15. (1) The functions φ_k are real analytic on [0,1]; when $k > 0$, then $\varphi_k < \xi_k$; when $k > 1$, then $\varphi_k > \xi_{k-1}$; one has $\lim_{\tau \to 1^-} \varphi_k(\tau) = 1/2$.

(2) Consider the function φ_k as a function of the two variables (k, τ) . One has $\partial \varphi_k/\partial k > 0$ for $k > 0$.

- (3) For $q \in (0,1)$, $x \in (-q^{-1}, \infty)$ one has $\partial \theta / \partial x > 0$.
- (4) For $q \in (0,1)$, $x \in (-q^{-1/2}, \infty)$ one has $\theta > 1/2$.

Before proving Proposition [15](#page-7-0) we finish the proof of Proposition [14.](#page-6-0) Part (2) of Proposition [15](#page-7-0) implies $\varphi_{3/4}(v^4) > \varphi_{1/4}(v^4)$. As $-v\Phi(v) = \varphi_{1/4}(v^4) - 1$ and $-v^3\Psi(v) = \varphi_{3/4}(v^4) - 1$, this means that $\theta(-v, 1/v) < 0$, i.e. $\theta(q, -1/q) < 0$. \Box

Proof of Proposition [15:](#page-7-0) Part (1) of the proposition is proved in [\[9\]](#page-12-5).

To prove part (2) observe that

$$
\partial \varphi_k / \partial k = (-\log \tau) \sum_{j=1}^{\infty} (-1)^{j-1} j \tau^{kj+j(j-1)/2}
$$

$$
= (-\log \tau) \tau^k \sum_{j=1}^{\infty} (\varphi_{k+j} - \tau^{k+j} \varphi_{k+j+1})
$$

As $\varphi_{k+j}-\tau^{k+j}\varphi_{k+j+1}=2\varphi_{k+j}-1$ and (see part (1) of the proposition) as $\varphi_{k+j}(\tau)>\xi_{k+j-1}(\tau)\geq$ 1/2, each difference $\varphi_{k+j} - \tau^{k+j} \varphi_{k+j+1}$ is positive on [0, 1). The factors $-\log \tau$ and τ^k are also positive.

For $x \in (-q^{-1}, 0]$ part (3) follows from part (2) and from $\varphi_k(\tau) = \theta(\tau, -\tau^{k-1})$. Indeed, one can represent x as $-\tau^{k-1}$ for some $k > 0$; for fixed τ , as x increases, k also increases. One has

$$
0 < \partial \varphi_k / \partial k = (-\log \tau) \partial \theta / \partial x |_{x = -\tau^{k-1}} \text{ and } -\log \tau > 0.
$$

For $x > 0$ part (3) results from all coefficients of $\theta(v, x)$ considered as a series in x being positive.

For $x \in (-q^{-1/2}, 0]$ part (4) follows from part (3). Indeed, consider the function $\psi :=$ $1+2\sum_{j=1}^{\infty}(-1)^{j}\tau^{j^2}, \tau \in [0,1).$ This function is positive valued, decreasing and tending to 0 as $\tau \to 1^-$, see [\[8\]](#page-12-8). As $0 < \psi(\tau^{1/2})/2 = \varphi_{1/2}(\tau) - 1/2$, for $k \ge 1/2$ part (2) of Proposition [15](#page-7-0) implies

$$
\theta(\tau, -\tau^{k-1}) = \varphi_k(\tau) \ge \varphi_{1/2}(\tau) > 1/2
$$
.

For $x = 0$ one has $\theta = 1$ hence for $x > 0$ part (4) follows from part (2).

 \Box

The following lemma follows immediately from the result of V. Katsnelson cited in part (4) of Remarks [7](#page-1-3) and from Proposition [14:](#page-6-0)

Lemma 16. For any $\varepsilon > 0$ sufficiently small there exists $\delta > 0$ such that for $q \in (-1, -1 + \delta]$ the function $\theta(q,.)$ has a single real zero in the interval $(-e^{\pi/2} + \varepsilon, e^{\pi/2} - \varepsilon)$. This zero is simple and positive.

4.2 How do the zeros of θ coalesce?

Further we describe the way multiple zeros are formed when q decreases from 0 to -1 .

Definition 17. We say that the phenomenon A happens before the phenomenon B if A takes place for $q = q_1$, B takes place for $q = q_2$ and $-1 < q_2 < q_1 < 0$. By phenomena we mean that certain zeros of θ or another function coalesce.

Notation 18. We denote by $x_j \prec x_k$ the following statement: The zeros x_k and x_{k+2} of θ can coalesce only after x_j and x_{j+2} have coalesced.

Lemma 19. One has x_{4k+2} $\prec x_{4k+3}$, x_{4k+2} $\prec x_{4k+1}$, x_{4k+3} $\prec x_{4k+4}$ and x_{4k+5} $\prec x_{4k+4}$, $k \in \mathbb{N} \cup 0$.

Proof. The statements follow respectively from

$$
qx_{4k+5} < x_{4k+4} < x_{4k+2} < qx_{4k+3} \quad , \quad x_{4k+1} < qx_{4k+2} < qx_{4k+4} < x_{4k+3} \quad ,
$$
\n
$$
qx_{4k+4} < x_{4k+3} < x_{4k+5} < qx_{4k+6} \quad \text{and} \quad x_{4k+6} < qx_{4k+7} < qx_{4k+5} < x_{4k+4} \quad .
$$

Lemma 20. One has $x_{4k+2} \prec x_{4k+6}$, $k \in \mathbb{N} \cup 0$.

Proof. Indeed, equation [\(2\)](#page-2-1) implies the following one:

$$
\theta(q,x) = 1 + qx + q^3x^2 + q^6x^3 + q^{10}x^4\theta(q,q^4x) \tag{6}
$$

 \Box

For $x = x_{4k+2}/q^4$ one obtains

$$
\theta(q, x_{4k+2}/q^4) = 1 + x_{4k+2}/q^3 + x_{4k+2}^2/q^5 + x_{4k+2}^3/q^6 = (1 + x_{4k+2}^2/q^5) + (x_{4k+2}/q^3 + x_{4k+2}^3/q^6).
$$

Each of the two sums is negative due to $q \in (-1,0)$, $x_{4k+2} < -1$ (see part (2) of Remarks [12\)](#page-6-2). For small values of |q| one has $x_{4k+2}/q^4 \in (x_{4k+8}, x_{4k+6})$ because $x_j = -q^{-j}(1 + O(q))$ and $\theta(q, x_{4k+2}/q^4) < 0$. By continuity this holds true for all $q \in (-1,0)$ for which the zeros x_{4k+8} , x_{4k+6}, x_{4k+4} and x_{4k+2} are real and distinct. Hence if x_{4k+2} and x_{4k+4} have not coalesced, then x_{4k+6} and x_{4k+8} are real and distinct. \Box

Remarks 21. [\(1\)](#page-1-1) Recall that for $q \in (-1, 0)$ we set $v := -q$ and that equation (1) holds true.

(2) Equation [\(3\)](#page-2-2) implies that the values of θ at its local maxima decrease and its values at local minima increase as q decreases in $(-1, 0)$. Indeed, at a critical point one has $\partial \theta / \partial x = 0$, so $2q\partial\theta/\partial q = x^2\partial^2\theta/\partial x^2$. At a minimum one has $\partial^2\theta/\partial x^2 \geq 0$, so $\partial\theta/\partial q \leq 0$ and the value of θ increases as q decreases; similarly for a maximum.

(3) In particular, this means that θ can only lose real zeros, but not acquire such as q decreases on $(-1, 0)$. Indeed, the zeros of θ depend continuously on q. If at some point of the real axis a new zero of even multiplicity appears, then it cannot be a maximum because the critical value must decrease and it cannot be a minimum because its value must increase.

(4) To treat the cases of odd multiplicities of the zeros it suffices to differentiate both sides of equation [\(3\)](#page-2-2) w.r.t. x. For example, a simple zero x_0 of θ cannot become a triple one because

$$
2q\partial/\partial q(\partial\theta/\partial x) = 2\partial\theta/\partial x + 4x\partial^2\theta/\partial x^2 + x^2\partial^3\theta/\partial x^3
$$

which means that as $\partial\theta/\partial x|_{x=x_0} = \partial^2\theta/\partial x^2|_{x=x_0} = 0$, then either $\partial^3\theta/\partial x^3|_{x=x_0} > 0$ hence in a neighbourhood of x_0 one has $\partial\theta/\partial x \geq 0$ and $\partial/\partial q(\partial\theta/\partial x) < 0$, or $\partial^3\theta/\partial x^3|_{x=x_0} < 0$ hence $\partial\theta/\partial x \leq 0$ and $\partial/\partial q(\partial\theta/\partial x) > 0$ (in a neighbourhood of x_0), so in both cases the triple zero bifurcates into a simple one and a complex pair as q decreases. The case of a zero of multiplicity $2m + 1$, $m \in \mathbb{N}$, is treated by analogy.

(5) In equation [\(1\)](#page-1-1) the first argument (i.e. v^4) of both functions $\theta(v^4, -x^2/v)$ and $\theta(v^4, -vx^2)$ is the same, so when one of them has a double zero, then they both have double zeros. If the double zero of the first one is at $x = a$, then the one of the second is at $x = a/v$.

Figure 4: The graphs of ψ_1 (solid line) and ψ_2 (dashed line).

Proposition 22. For any $k \in \mathbb{N} \cup 0$ there exists $q_k^* \in (-1,0)$ such that for $q = q_k^*$ the zeros x_{4k+2} and x_{4k+4} coalesce.

Notation 23. We denote by ψ_1 and ψ_2 the functions $\theta(v^4, -x^2/v)$ and $-vx\theta(v^4, -vx^2)$. By $y_{\pm k}$ and $z_{\pm k}$ we denote their real zeros for $v^4 \in (0, 0.108]$, their moduli increasing with $k \in \mathbb{N}$, $y_k > 0$, $y_{-k} < 0$, $z_k > 0$, $z_{-k} < 0$. We set $z_0 = 0$.

Proof of Proposition [22:](#page-9-0) On Fig. [4](#page-9-1) we show for $v^4 \in (0, 0.108]$ how the graphs of the functions ψ_1 and ψ_2 (drawn in solid and dashed line respectively) look like, the former being even and the latter odd, see part (1) of Remarks [21.](#page-8-0) The signs + and − indicate places, where it is certain that their sum $\theta(-v, x)$ is positive or negative. It is positive (resp. negative) if both functions are of this sign. It is positive near the origin because $\psi_1|_{x=0} > 0$ while $\psi_2|_{x=0} = 0$.

For values of v close to 0 the zeros $y_{\pm k}$ of ψ_1 (resp. the zeros $z_{\pm k}$ of $\psi_2(x)$ are close to the numbers $\pm v^{-(4k-1)/2}$ (resp. $\pm v^{-(4k+1)/2}$), $k \in \mathbb{N}$, see part (1) of Remarks [12.](#page-6-2) From these remarks follows also that for small positive values of v the zeros of $\theta(-v, x)$ are close to the numbers $-v^{-k}$, $k \in \mathbb{N}$. Hence on the negative half-axis x one obtains the following arrangement of these numbers:

$$
\cdots < -v^{-11/2} < -v^{-9/2} < -v^{-4} < -v^{-7/2} < -v^{-5/2} < -v^{-2} < -v^{-3/2} < 0.
$$
\n
$$
y_{-5} \qquad z_{-3} \qquad x_4 \qquad y_{-3} \qquad z_{-1} \qquad x_2 \qquad y_{-1}
$$

Hence for $v^4 \in (0, 0.108]$ between a zone marked by a + and one marked by a - there is exactly one zero of $\theta(-v, .)$.

As v increases from 0 to 1, it takes countably-many values at each of which one of the functions ψ_1 and ψ_2 (in turn) has a double zero and when the value is passed, this double zero becomes a conjugate pair, see Theorem [1.](#page-0-0) Hence on the negative real half-line the regions, where the corresponding function ψ_1 or ψ_2 is negative, disappear one by one starting from the right. As two consecutive changes of sign of $\theta(-v,.)$ are lost, $\theta(-v,.)$ has a couple of consecutive real negative zeros replaced by a complex conjugate pair. The quantity of these losses being countable implies the proposition. \Box

Proposition 24. Any interval of the form $[\gamma, 0], \gamma \in (-1, 0)$, contains at most finitely many spectral values of q.

Proof. Indeed, if $\gamma \ge -0.108$, the interval contains no spectral value of q, see Lemma [10;](#page-5-2) all zeros of θ are real and distinct and the graphs of the functions $\psi_{1,2}$ look as shown on Fig. [4.](#page-9-1)

Suppose that $\gamma < -0.108$. When q decreases in [γ , 0], for each of the functions this happens at most finitely many times that it has a double zero which then gives birth to a complex conjugate pair. This is always the zero which is closest to 0, see part (2) of Theorem [1.](#page-0-0)

Hence the presentation of the graphs of $\psi_{1,2}$ changes only on some closed interval containing 0, but outside it the zones marked by $+$ and $-$ continue to exist (but their borders change continuously) and the simple zeros of θ that are to be found between two such consecutive zones of opposite signs are still to be found there. Besides, no new real zeros appear, see part (3) of Remarks [21.](#page-8-0)

Therefore there exists $s_0 \in \mathbb{N}$ such that when q decreases from 0 to γ , the zeros x_{s_0}, x_{s_0+1} , x_{s_0+2} , ... remain simple and depend continuously on q. Hence only the rest of the zeros (i.e. x_1, \ldots, x_{s_0-1} can participate in the bifurcations. \Box

Proposition 25. For $q \in (-1,0)$ the function $\theta(q,.)$ can have only simple and double real zeros. Positive double zeros are local maxima and negative double zeros are local minima.

Proof. Equality [\(2\)](#page-2-1) implies the following one: $\theta(q, x/q^2) = 1 + x/q + (x^2/q)\theta(q, x)$. Set $x = x_{4k+2}$. Hence $x_{4k+2} < -1$, see part (2) of Remarks [12.](#page-6-2) As $\theta(q, x_{4k+2}) = 0$ and $1 + x_{4k+2}/q > 0$, this implies $\theta(q, x_{4k+2}/q^2) > 0$. In the same way $\theta(q, x_{4k+4}/q^2) > 0$.

For q close to 0 the numbers x_{4k+2}/q^2 and x_{4k+4}/q^2 are close respectively to x_{4k+4} and x_{4k+6} , see part (1) of Remarks [12.](#page-6-2) Hence for such values of q one has

$$
x_{4k+6} < x_{4k+4}/q^2 < x_{4k+2}/q^2 < x_{4k+4} \tag{7}
$$

This string of inequalities holds true (by continuity) for q belonging to any interval of the form $(a, 0), a \in (-1, 0)$, for any q of which the zeros x_{4k+6}, x_{4k+4} and x_{4k+2} are real and distinct.

Equation [\(7\)](#page-10-0) implies that x_{4k+6} and x_{4k+4} cannot coalesce if x_{4k+4} and x_{4k+2} are real (but not necessarily distinct). Hence when for the first time negative zeros of θ coalesce, this happens with exactly two zeros, and the double zero is a local minimum of θ .

For positive zeros one obtains in the same way the string of inequalities

$$
x_{4k+5} < x_{4k+3}/q^2 < x_{4k+5}/q^2 < x_{4k+7} \ .
$$

Indeed, one has $1 + x_{4k+3}/q < 0$ because $x_{4k+3} > 1$ (see Proposition [14\)](#page-6-0) and $q \in (0,1)$. Hence $\theta(q, x_{4k+3}/q^2)$ < 0 and in the same way $\theta(q, x_{4k+5}/q^2)$ < 0. Thus x_{4k+5} and x_{4k+7} cannot coalesce if x_{4k+3} and x_{4k+5} are real (but not necessarily distinct). Hence when for the first time positive zeros of θ coalesce, this happens with exactly two zeros, and the double zero is a local maximum of θ .

After a confluence of zeros takes place, one can give new indices to the remaining real zeros so that the indices of consecutive zeros differ by $2 (x_{2s+2} < x_{2s} < 0$ and $0 < x_{2s+1} < x_{2s+3})$. After this for the next confluence the reasoning is the same. \Box

Lemma 26. For $k \in \mathbb{N}$ sufficiently large one has $x_{4k+3} \prec x_{4k+6}$.

Proof. Suppose that $x_{4k+3} \geq 3$. Then

$$
\theta(q, x_{4k+3}/q^3) = 1 + x_{4k+3}/q^2 + x_{4k+3}^2/q^3 + (x_{4k+3}^3/q^3)\theta(q, x_{4k+3}) = 1 + x_{4k+3}/q^2 + x_{4k+3}^2/q^3.
$$

For $x_{4k+3} \geq 3$ and $q \in (-1,0)$ the right-hand side is negative. In the same way $\theta(q, x_{4k+5}/q^3)$ 0. We prove below that

$$
x_{4k+8} < x_{4k+5}/q^3 < x_{4k+3}/q^3 < x_{4k+6} \tag{8}
$$

Hence the zeros x_{4k+8} and x_{4k+6} cannot coalesce before x_{4k+5} and x_{4k+3} have coalesced. To prove the string of inequalities [\(8\)](#page-11-1) observe that for q close to 0 the numbers x_{4k+8} and x_{4k+5}/q^3 are close to one another $(x_{4k+3}/q^3$ and x_{4k+6} as well, see part (1) of Remarks [12\)](#page-6-2) which implies [\(8\)](#page-11-1). By continuity, as long as $x_{4k+3} \geq 3$ and $q \in (-1,0)$, the string of inequalities holds true also for q not necessarily close to 0.

The result of V. Katsnelson (see part (4) of Remarks [7\)](#page-1-3) implies that there exists $a \in (-1,0)$ such that for $q \in (-1, a]$ the function $\theta(q, x)$ has no zeros in $[-3, 3]$ except the one which is simple and close to 1 (see Proposition [14\)](#page-6-0). Hence for $q \in (-1, a]$ the condition $x_{4k+3} \geq 3$ is fulfilled, if the zero x_{4k+3} has been real and simple for $q \in (a, 0)$. On the other hand for $q \in [a, 0)$ only finitely many real zeros of θ have coalesced, and only finitely many have belonged to the interval [−3, 3] for some value of q, see Proposition [24.](#page-9-2) Therefore there exists $k_0 \in \mathbb{N}$ such that for $k \geq k_0$ one has $x_{4k+3} \geq 3$. \Box

4.3 Completion of the proof of Theorem [4](#page-1-0)

Proposition [25](#page-10-1) and Remarks [21](#page-8-0) show that $\theta(q,.)$ has no real zero of multiplicity higher that 2. Lemma [20](#page-8-1) implies the string of inequalities $-1 < \cdots < \bar{q}_{2l+2} < \bar{q}_{2l} < \cdots < 0$. For k sufficiently large one has $-1 < \cdots < \bar{q}_{k+1} < \bar{q}_k < \cdots < 0$. This follows from Lemmas [19](#page-8-2) and [26.](#page-10-2) It results from Proposition [24](#page-9-2) and from the above inequalities that the set of spectral values has −1 as unique accumulation point. This proves part (3) of the theorem.

Part (2) of the theorem results from Proposition [25.](#page-10-1)

Part (4) follows from Remarks [21.](#page-8-0) These remarks show that real zeros can be only lost and that no new real zeros are born.

5 Proof of Theorem [6](#page-1-4)

Recall that $\psi_1 = \theta(v^4, -x^2/v)$ and $\psi_2 = -vx\theta(v^4, -vx^2)$, see Notation [23.](#page-9-3) Recall that the spectral values \tilde{q}_k of q for $\theta(q, x)$, $q \in (0, 1)$ satisfy the asymptotic relation $\tilde{q}_k = 1 - \pi/2k + o(1/k)$. Hence the values of v for which the function $\theta(v^4, x)$ has a double zero are of the form

$$
\tilde{v}_k = (\tilde{q}_k)^{1/4} = 1 - \pi/8k + o(1/k)
$$

and the functions $\psi_{1,2}$ have double zeros for $v = \tilde{v}_k$.

Consider three consecutive values of k the first of which is odd – k_0 , $k_0 + 1$ and $k_0 + 2$. Set $v := \tilde{v}_{k_0}$. Denote by $a < b < 0$ the double negative zeros of the functions $\psi_{1,2}|_{v=\tilde{v}_{k_0}}$. These zeros are local minima and on the whole interval $[a, b]$ one has $\theta > 0$. The values of θ at local minima increase (see part (2) of Remarks [21\)](#page-8-0), therefore the double zero of $\theta(\bar{q}_{k_0},.)$ is obtained for some $|q| < |\tilde{v}_{k_0}|$, i.e. before the functions $\psi_{1,2}|_{v=\tilde{v}_{k_0}}$ have double zeros. This follows from equality [\(1\)](#page-1-1) in which both summands in the right-hand side have local minima (recall that as k_0 is odd, the double zero of θ is negative, so $x < 0$ in equality [\(1\)](#page-1-1)). Hence

$$
|\bar{q}_{k_0}| < |\tilde{v}_{k_0}| = 1 - \pi/8k_0 + o(1/k_0) \tag{9}
$$

In the same way

$$
|\bar{q}_{k_0+2}| < |\tilde{v}_{k_0+2}| = 1 - \pi/8(k_0+2) + o(1/(k_0+2)). \tag{10}
$$

In the case of $k_0 + 1$ the function $\psi_1|_{v=\tilde{v}_{k_0+1}}$ has a local minimum while $\psi_2|_{v=\tilde{v}_{k_0+1}}$ has a local maximum (because $k_0 + 1$ is even, the double zero of θ is positive, so $x > 0$ in equality [\(1\)](#page-1-1) $x > 0$). As θ has a local maximum and as the values of θ at local maxima decrease (see part (2) of Remarks [21\)](#page-8-0), the double zero of $\theta(\bar{q}_{k_0+1},.)$ is obtained for some $|q| > |\tilde{v}_{k_0+1}|$, i.e. after the functions $\psi_{1,2}|_{v=\tilde{v}_{k_0+1}}$ have double zeros. Therefore

$$
|\bar{q}_{k_0+1}| > |\tilde{v}_{k_0+1}| = 1 - \pi/8(k_0+1) + o(1/(k_0+1)) . \tag{11}
$$

When k_0 is sufficiently large one has $|\bar{q}_{k_0}| < |\bar{q}_{k_0+1}| < |\bar{q}_{k_0+2}|$ (this follows from part (3) of Theorem [4\)](#page-1-0). Using equations [\(10\)](#page-11-2) and [\(11\)](#page-12-11) one gets

$$
1 - \pi/8(k_0 + 1) + o(1/(k_0 + 1)) < |\bar{q}_{k_0+1}| < |\bar{q}_{k_0+2}| < 1 - \pi/8(k_0 + 2) + o(1/(k_0 + 2))
$$

Hence $|\bar{q}_{k_0+1}| = 1 - \pi/8(k_0+1) + o(1/(k_0+1))$ and $|\bar{q}_{k_0+2}| = 1 - \pi/8(k_0+2) + o(1/(k_0+2)).$ This implies the statement of Theorem [6.](#page-1-4)

References

- [1] G. E. Andrews, B. C. Berndt, Ramanujan's lost notebook. Part II. Springer, NY, 2009.
- [2] B. C. Berndt, B. Kim, Asymptotic expansions of certain partial theta functions. Proc. Amer. Math. Soc. 139 (2011), no. 11, 3779–3788.
- [3] K. Bringmann, A. Folsom, R. C. Rhoades, Partial theta functions and mock modular forms as q-hypergeometric series, Ramanujan J. 29 (2012), no. 1-3, 295-310, <http://arxiv.org/abs/1109.6560>
- [4] G. H. Hardy, On the zeros of a class of integral functions, Messenger of Mathematics, 34 (1904), 97–101.
- [5] J. I. Hutchinson, On a remarkable class of entire functions, Trans. Amer. Math. Soc. 25 (1923), pp. 325–332.
- [6] O.M. Katkova, T. Lobova and A.M. Vishnyakova, On power series having sections with only real zeros. Comput. Methods Funct. Theory 3 (2003), no. 2, 425–441.
- [7] B. Katsnelson, On summation of the Taylor series for the function $1/(1-z)$ by the theta summation method, [arXiv:1402.4629v](http://arxiv.org/abs/1402.4629)1 [math.CA].
- [8] V.P. Kostov, About a partial theta function, Comptes Rendus Acad. Sci. Bulgare 66, No 5 (2013) 629-634.
- [9] V.P. Kostov, On the zeros of a partial theta function, Bull. Sci. Math. 137, No. 8 (2013) 1018-1030.
- [10] V.P. Kostov, Asymptotics of the spectrum of partial theta function, Revista Mat. Complut. 27, No. 2 (2014) 677-684, DOI: 10.1007/s13163-013-0133-3.
- [11] V.P. Kostov, On the spectrum of a partial theta function, Proc. Royal Soc. Edinb. A (to appear).
- [12] V.P. Kostov, A property of a partial theta function, Comptes Rendus Acad. Sci. Bulgare (to appear).
- [13] V.P. Kostov and B. Shapiro, Hardy-Petrovitch-Hutchinson's problem and partial theta function, Duke Math. J. 162, No. 5 (2013) 825-861, [arXiv:1106.6262v](http://arxiv.org/abs/1106.6262)1[math.CA].
- [14] B. Ja. Levin, Zeros of Entire Functions, AMS, Providence, Rhode Island 1964.
- [15] I. V. Ostrovskii, On zero distribution of sections and tails of power series, Israel Math. Conf. Proceedings, 15 (2001), 297–310.
- [16] M. Petrovitch, Une classe remarquable de séries entières, Atti del IV Congresso Internationale dei Matematici, Rome (Ser. 1), 2 (1908), 36–43.
- [17] G. Pólya and J. Schur, Über zwei Arten von Faktorenfolgen in der Theorie der algebraischen Gleichungen, J. Reine Angew. Math. 144 (1914) 89–113.
- [18] A. Sokal, The leading root of the partial theta function, Adv. Math. 229 (2012), no. 5, 2603-2621, [arXiv:1106.1003.](http://arxiv.org/abs/1106.1003)
- [19] S. O. Warnaar, Partial theta functions. I. Beyond the lost notebook, Proc. London Math. Soc. (3) 87 (2003), no. 2, 363–395.