

Random block coordinate descent methods for linearly constrained optimization over networks

I. Necoara, Yu. Nesterov and F. Glineur

Received: March 2013 / Updated: November 2014

Abstract In this paper we develop random block coordinate gradient descent methods for minimizing large scale linearly constrained separable convex problems over networks. Since we have coupled constraints in the problem, we devise an algorithm that updates in parallel $\tau \geq 2$ (block) components per iteration. Moreover, for this method the computations can be performed in a distributed fashion according to the structure of the network. However, its complexity per iteration is usually cheaper than of the full gradient method when the number of nodes N in the network is large. We prove that for this method we obtain in expectation an ϵ -accurate solution in at most $\mathcal{O}(\frac{N}{\tau\epsilon})$ iterations and thus the convergence rate depends linearly on the number of (block) components τ to be updated. For strongly convex functions the new method converges linearly. We also focus on how to choose the probabilities to make the randomized algorithm to converge as fast as possible and we arrive at solving a sparse SDP. Finally, we describe several applications that fit in our framework, in particular the convex feasibility problem. Numerically, we show that the parallel coordinate descent method with $\tau > 2$ accelerates on its basic counterpart corresponding to $\tau = 2$.

1 Introduction

The performance of a network composed of interconnected subsystems can be improved if the traditionally separated subsystems are optimized together. Recently, coordinate descent methods have emerged as a powerful tool for solving large data network problems: e.g. resource allocation [7, 17], coordination in multi-agent systems [5, 7, 18], estimation in sensor networks or distributed control [10], image

I. Necoara is with Automatic Control and Systems Engineering Department, University Politehnica Bucharest, 060042 Bucharest, Romania. E-mail: ion.necoara@acse.pub.ro.

Yu. Nesterov and F. Glineur are with Center for Operations Research and Econometrics, Catholic University of Louvain, Louvain-la-Neuve, B-1348, Belgium. E-mail: {[Yurii.Nesterov](mailto:Yurii.Nesterov@uclouvain.be), [Francois.Glineur](mailto:Francois.Glineur@uclouvain.be)}@uclouvain.be.

The present paper is a generalization of our paper: I. Necoara, Yu. Nesterov and F. Glineur, A random coordinate descent method on large-scale optimization problems with linear constraints, Tech. Rep., Univ. Politehnica Bucharest, 1-25, July 2011 (ICCOPT 2013, Lisbon).

processing [3, 4, 19] and other areas [6, 13, 15]. The problems we consider in this paper have the following features: *the size of data* is big so that usual methods based on whole gradient computations are prohibitive. Moreover *the incomplete structure of information* (e.g. the data are distributed over the nodes of the network, so that at a given time we need to work only with the data available then) may also be an obstacle for whole gradient computations. In this case, an appropriate way to approach these problems is through coordinate descent methods. These methods were among the first optimization methods studied in literature but until recently they haven't received much attention.

The main differences in all variants of coordinate descent methods consist in the criterion of choosing at each iteration the coordinate over which we minimize the objective function and the complexity of this choice. Two classical criteria used often in these algorithms are the cyclic and the greedy coordinate descent search, which significantly differs by the amount of computations required to choose the appropriate index. For cyclic coordinate search estimates on the rate of convergence were given recently in [2], while for the greedy coordinate search (e.g. Gauss-Southwell rule) the convergence rate is given e.g. in [16]. One paper related to our work is [1], where a 2-coordinate greedy descent method is developed for minimizing a smooth function subject to a single linear equality constraint and additional bound constraints on the decision variables. Another interesting approach is based on random choice rule, where the coordinate search is random. Recent complexity results on random coordinate descent methods for smooth convex objective functions were obtained in [7, 12]. The extension to composite convex objective functions was given e.g. in [8, 14]. These methods are inherently serial. Recently, parallel and distributed implementations of coordinate descent methods were also analyzed e.g. in [6, 9, 10, 15].

Contributions: In this paper we develop random block coordinate gradient descent methods suited for large optimization problems in networks where the information cannot be gathered centrally, but rather it is distributed over the network. Moreover, in our paper we focus on optimization problems with linearly coupled constraints (i.e. the constraint set is coupled). Due to the coupling in the constraints we introduce a $\tau \geq 2$ block variant of random coordinate gradient descent method, that involves at each iteration the closed form solution of an optimization problem only with respect to τ block variables while keeping all the other variables fixed. Our approach allows us to analyze in the same framework several methods: full gradient, serial random coordinate descent and any parallel random coordinate descent method in between. For this method we obtain for the expected values of the objective function a convergence rate $\mathcal{O}(\frac{N}{\tau^k})$, where k is the iteration counter and N is the number of nodes in the network. Thus, the theoretical speedup in terms of the number of iterations needed to approximately solve the problem, as compared to the basic method corresponding to $\tau = 2$, is an expression depending on the number of components τ to be updated (processors) and for a complete network the speedup is equal to τ (number of components updated). This result also shows that the speedup achieved by our method on the class of separable problems with coupling constraints is the same as for separable problems without coupling constraints. For strongly convex functions we prove that the new method converges linearly. We also focus on how to choose the probabilities to make the randomized algorithm to converge as fast as possible and we arrive at solving sparse SDPs. While the most obvious benefit of randomization is that it can lead

to faster algorithms, either in worst case complexity analysis and/or numerical implementation, there are also other benefits of our algorithm that are at least as important: e.g., the use of randomization leads to a simpler algorithm that is easier to analyze, produces a more robust output and can often be organized to exploit modern computational architectures (e.g distributed and parallel computers).

Contents: The paper is organized as follows. In Section 2 we introduce our optimization model and assumptions. In Section 3 we propose a random block coordinate descent algorithm and derive the convergence rate in expectation. Section 4 provides means to choose optimally the probability distribution. In Section 5 we discuss possible applications and we conclude with some preliminary numerical results in Section 6.

Notation: We work in the space \mathbb{R}^N composed by column vectors. For $x, y \in \mathbb{R}^N$ denote the standard Euclidian inner product $\langle x, y \rangle = x^T y$ and the Euclidian norm $\|x\| = \langle x, x \rangle^{1/2}$. For symmetric matrices X, Y we consider the inner product $\langle X, Y \rangle = \text{trace}(XY)$. We use the same notation $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ for spaces of different dimension. We define the partition of the identity matrix: $I_N = [e_1 \cdots e_N]$, where $e_i \in \mathbb{R}^N$. Then, for any $x \in \mathbb{R}^N$ we write $x = \sum_i e_i x_i$. Moreover, D_x denotes the diagonal matrix with the entries x on the diagonal and $x^{-1} = [x_1^{-1} \cdots x_N^{-1}]^T$. We denote with $e \in \mathbb{R}^N$ the N -dimensional vector with all entries equal to one. For a positive semidefinite matrix $W \in \mathbb{R}^{N \times N}$ we consider the following order on its eigenvalues $0 \leq \lambda_1 \leq \cdots \leq \lambda_N$ and $\|x\|_W^2 = x^T W x$.

2 Problem formulation

We consider large data network optimization problems where each agent in the network is associated with a local variable so that their sum is fixed and we need to minimize a separable convex objective function:

$$\begin{aligned} f^* &= \min_{x_i \in \mathbb{R}} f_1(x_1) + \cdots + f_N(x_N) \\ \text{s.t.} & \quad x_1 + \cdots + x_N = 0. \end{aligned} \tag{1}$$

For convenience, we will focus on scalar convex functions $f_i : \mathbb{R} \rightarrow \mathbb{R}$, i.e. $x_i \in \mathbb{R}$, in the optimization model (1). However, our results can be easily extended to the block case, when $x_i \in \mathbb{R}^n$, with $n \geq 1$, using the Kronecker product. Moreover, constraints of the form $\alpha_1 x_1 + \cdots + \alpha_N x_N = b$, where $x_i \in \mathbb{R}^n$ and $\alpha_i \in \mathbb{R}$, can be easily handled in our framework by a change of coordinates.

Optimization problems with linearly coupled constraints (1) arise in many areas such as resource allocation [7, 17], coordination in multi-agent systems [5, 7, 18], image processing [3, 4, 10, 19] and other areas [6, 13, 15]. For problem (1) we associate a network composed of several nodes $[N] = \{1, \cdots, N\}$ that can exchange information according to a communication graph $\mathcal{G} = ([N], E)$, where E denotes the set of edges, i.e. $(i, j) \in E \subseteq [N] \times [N]$ models that node j sends information to node i . We assume that the graph \mathcal{G} is undirected and connected. For an integer $\tau \geq 2$, we also define with \mathcal{P}_τ the set of paths of τ vertices in the graph. Note that we have at most $\binom{\tau}{N}$ paths of τ vertices in a graph. The local information structure imposed by the graph \mathcal{G} should be considered as part of the problem formulation.

Our goal is to devise a distributed algorithm that iteratively solves the convex problem (1) by passing the estimate of the optimizer only between neighboring nodes along paths of τ vertices. There is great interest in designing such distributed and parallel algorithms, since centralized algorithms scale poorly with the number of nodes and are less resilient to failure of the central node. We use the notation:

$$x = [x_1 \cdots x_N]^T \quad \text{and} \quad f(x) = f_1(x_1) + \cdots + f_N(x_N).$$

Let us define the extended subspace $S \subseteq \mathbb{R}^N$ and its orthogonal complement $T \subseteq \mathbb{R}^N$:

$$S = \left\{ x : \sum_{i=1}^N x_i = 0 \right\}, \quad T = \{ u : u_1 = \cdots = u_N \}.$$

The basic assumption considered in this paper is:

Assumption 2.1 *We assume that each function f_i is convex and has Lipschitz continuous gradient with constants $L_i > 0$, i.e. the following inequality holds:*

$$\|\nabla f_i(x_i) - \nabla f_i(y_i)\| \leq L_i \|x_i - y_i\| \quad \forall x_i, y_i \in \mathbb{R}. \quad (2)$$

From the Lipschitz property of the gradient (2), the following inequality holds for all $x_i, d_i \in \mathbb{R}$ [11]:

$$f_i(x_i + d_i) \leq f_i(x_i) + \langle \nabla f_i(x_i), d_i \rangle + \frac{L_i}{2} \|d_i\|^2. \quad (3)$$

We denote with X^* the set of optimal solutions for problem (1). Note that x^* is optimal solution for (1) if and only if:

$$\sum_{i=1}^N x_i^* = 0, \quad \nabla f_i(x_i^*) = \nabla f_j(x_j^*) \quad \forall i \neq j \in [N].$$

3 Random coordinate descent algorithms

In this section we devise randomized block coordinate gradient descent algorithms for solving the separable convex problem (1) and analyze their convergence. Since we have coupled constraints in the problem, the algorithm has to update in parallel $\tau \geq 2$ components per iteration. Usually, the algorithm can be accelerated by parallelization, i.e. by using more than one pair of coordinates per iteration. Our approach allows us to analyze in the same framework several methods: full gradient ($\tau = N$), serial random coordinate descent ($\tau = 2$) and any parallel random coordinate descent method in between ($< 2\tau < N$). Let us fix $N \geq \tau \geq 2$ and we denote with $\mathcal{N} \in \mathcal{P}_\tau$ a path of τ vertices in the connected undirected graph \mathcal{G} . We also assume available a probability distribution $p_{\mathcal{N}}$ over the set \mathcal{P}_τ of paths of τ vertices in the graph \mathcal{G} . Then, we can derive a randomized τ coordinate descent algorithm where we update at each iteration only τ coordinates in the vector x . Let us define $\mathcal{N} = (i_1, \cdots, i_\tau) \in \mathcal{P}_\tau$, with $i_l \in [N]$, $s_{\mathcal{N}} = [s_{i_1} \cdots s_{i_\tau}]^T \in \mathbb{R}^\tau$, $L_{\mathcal{N}} = [L_{i_1} \cdots L_{i_\tau}]^T \in \mathbb{R}^\tau$ and $\nabla f_{\mathcal{N}} = [\nabla f_{i_1} \cdots \nabla f_{i_\tau}]^T \in \mathbb{R}^\tau$. Under assumption (2) the following inequality holds:

$$f(x + \sum_{i \in \mathcal{N}} e_i s_i) \leq f(x) + \langle \nabla f_{\mathcal{N}}(x), s_{\mathcal{N}} \rangle + \frac{1}{2} \|s_{\mathcal{N}}\|_{D_{L_{\mathcal{N}}}}^2. \quad (4)$$

Based on the inequality (4) we can devise a general randomized τ coordinate descent algorithm for problem (1), let us call it RCD_τ . Given an x in the feasible set S , we choose the coordinate τ -tuple $\mathcal{N} \in \mathcal{P}_\tau$ with probability $p_\mathcal{N}$. Let the next iterate be chosen as follows:

$$x^\dagger = x + \sum_{i \in \mathcal{N}} e_i d_i,$$

i.e. we update τ components in the vector x , where the direction $d_\mathcal{N}$ is determined by requiring that the next iterate x^\dagger to be also feasible for (1) and minimizing the right hand side in (4), i.e.:

$$d_\mathcal{N} = \arg \min_{s_\mathcal{N}: \sum_{i \in \mathcal{N}} s_i = 0} f(x) + \langle \nabla f_\mathcal{N}(x), s_\mathcal{N} \rangle + \frac{1}{2} \|s_\mathcal{N}\|_{D_{L_\mathcal{N}}}^2$$

or explicitly, in closed form:

$$d_i = \frac{1}{L_i} \frac{\sum_{j \in \mathcal{N}} \frac{1}{L_j} (\nabla f_j(x_j) - \nabla f_i(x_i))}{\sum_{j \in \mathcal{N}} \frac{1}{L_j}} \quad \forall i \in \mathcal{N}.$$

In conclusion, we obtain the following randomized τ coordinate gradient descent method:

Algorithm RCD_τ

1. choose τ -tuple $\mathcal{N}_k = (i_1^k, \dots, i_\tau^k)$ with probability $p_\mathcal{N}$
2. set $x_i^{k+1} = x_i^k \quad \forall i \notin \mathcal{N}_k$ and $x_i^{k+1} = x_i^k + d_i^k \quad \forall i \in \mathcal{N}_k$.

Clearly, algorithm RCD_τ is distributed since only neighboring nodes along a path in the graph need to communicate at each iteration. Further, at each iteration only τ components of x are updated, so that our method has low complexity per iteration. Finally, in our algorithm we maintain feasibility at each iteration, i.e. $x_1^k + \dots + x_N^k = 0$ for all $k \geq 0$. The random choice of coordinates makes the algorithm adequate for parallel and distributed implementations and thus more flexible than greedy coordinate descent methods [1, 16]. In particular, for $\tau = 2$, we choose one pair of connected nodes $(i_k, j_k) \in E$ with some given probability $p_{i_k j_k}$ and obtain the following basic iteration:

$$x^{k+1} = x^k + \frac{1}{L_{i_k} + L_{j_k}} (e_{i_k} - e_{j_k}) \left(\nabla f_{j_k}(x_{j_k}^k) - \nabla f_{i_k}(x_{i_k}^k) \right).$$

Note that our algorithm belongs to the class of center-free methods (in [17] the term center-free refers to the absence of a coordinator) with the following iteration:

$$x_i^{k+1} = x_i^k + \sum_{j \in \mathcal{N}} w_{ij}^k \left(\nabla f_j(x_j^k) - \nabla f_i(x_i^k) \right) \quad \forall i \in [N], \quad (5)$$

with appropriate weights w_{ij}^k . Based on the inequality (4) and the optimality conditions for the subproblem corresponding to $d_\mathcal{N}$, the following decrease in the

objective function values can be derived:

$$\begin{aligned} f(x^+) &\leq f(x) - \sum_{i \in \mathcal{N}} \frac{\left| \sum_{j \in \mathcal{N}} \frac{1}{L_j} (\nabla f_j(x_j) - \nabla f_i(x_i)) \right|^2}{2L_i \left(\sum_{j \in \mathcal{N}} \frac{1}{L_j} \right)^2} \\ &= f(x) - \frac{1}{2} \nabla f(x)^T G_{\mathcal{N}} \nabla f(x), \end{aligned}$$

where the matrix $G_{\mathcal{N}}$ is defined as follows:

$$G_{\mathcal{N}} = D_{L_{\mathcal{N}}}^{-1} - \frac{1}{\sum_{i \in \mathcal{N}} 1/L_i} L_{\mathcal{N}}^{-1} (L_{\mathcal{N}}^{-1})^T, \quad (6)$$

where, with an abuse of notation, $L_{\mathcal{N}} \in \mathbb{R}^N$ denotes the vector with components zero outside the index set \mathcal{N} and components L_i for $i \in \mathcal{N}$. Therefore, taking the expectation over the random τ -tuple $\mathcal{N} \in \mathcal{P}_{\tau}$, we obtain the following inequality:

$$E[f(x^+) | x] \leq f(x) - \frac{1}{2} \nabla f(x)^T G_{\tau} \nabla f(x), \quad (7)$$

where $G_{\tau} = \sum_{\mathcal{N} \in \mathcal{P}_{\tau}} p_{\mathcal{N}} G_{\mathcal{N}}$ and can be interpreted as a weighted Laplacian for the graph \mathcal{G} . From the decrease in the objective function values given above, it follows immediately that the matrix G_{τ} is positive semidefinite and has an eigenvalue $\lambda_1(G_{\tau}) = 0$ with the corresponding eigenvector $e \in T$. Since the graph is connected, it also follows that the eigenvalue $\lambda_1(G_{\tau}) = 0$ is simple, i.e. $\lambda_2(G_{\tau}) > 0$. On the extended subspace S we now define a norm that will be used subsequently for measuring distances in this subspace. We define the primal ‘‘norm’’ induced by the positive semidefinite matrix G_{τ} as:

$$\|u\|_{G_{\tau}} = \sqrt{u^T G_{\tau} u} \quad \forall u \in \mathbb{R}^N.$$

Note that $\|u\|_{G_{\tau}} = 0$ for all $u \in T$ and $\|u\|_{G_{\tau}} > 0$ for all $u \in \mathbb{R}^n \setminus T$. On the subspace S we introduce its extended dual norm:

$$\|x\|_{G_{\tau}}^* = \max_{u \in \mathbb{R}^N: \|u\|_{G_{\tau}} \leq 1} \langle x, u \rangle \quad \forall x \in S.$$

Using the definition of conjugate norms, the Cauchy-Schwartz inequality holds:

$$\langle u, x \rangle \leq \|u\|_{G_{\tau}} \cdot \|x\|_{G_{\tau}}^* \quad \forall x \in S, u \in \mathbb{R}^N.$$

Let us define the average value: $\hat{u} = \frac{1}{N} \sum_{i=1}^N u_i$. Then, the dual norm can be computed for any $x \in S$ as follows:

$$\begin{aligned} \|x\|_{G_{\tau}}^* &= \max_{u \in \mathbb{R}^N: \langle G_{\tau} u, u \rangle \leq 1} \langle x, u \rangle = \max_{u: \langle G_{\tau}(u - \hat{u}e), u - \hat{u}e \rangle \leq 1} \langle x, u - \hat{u}e \rangle \\ &= \max_{u: \langle G_{\tau} u, u \rangle \leq 1, \sum_{i=1}^N u_i = 0} \langle x, u \rangle = \max_{u: \langle G_{\tau} u, u \rangle \leq 1, e^T u = 0} \langle x, u \rangle \\ &= \max_{u: \langle G_{\tau} u, u \rangle \leq 1, u^T e e^T u \leq 0} \langle x, u \rangle \\ &= \min_{\nu, \mu \geq 0} \max_{u \in \mathbb{R}^N} [\langle x, u \rangle + \mu(1 - \langle G_{\tau} u, u \rangle) - \nu \langle e e^T u, u \rangle] \\ &= \min_{\nu, \mu \geq 0} \mu + \langle (\mu G_{\tau} + \nu e e^T)^{-1} x, x \rangle = \min_{\nu \geq 0} \min_{\mu \geq 0} \left[\mu + \frac{1}{\mu} \langle (G_{\tau} + \frac{\nu}{\mu} e e^T)^{-1} x, x \rangle \right] \\ &= \min_{\zeta \geq 0} \sqrt{\langle (G_{\tau} + \zeta e e^T)^{-1} x, x \rangle}. \end{aligned}$$

In conclusion, we obtain an extended dual norm that is well defined on subspace S :

$$\|x\|_{G_\tau}^* = \min_{\zeta \geq 0} \sqrt{\langle (G_\tau + \zeta ee^T)^{-1} x, x \rangle} \quad \forall x \in S. \quad (8)$$

Using the eigenvalue decomposition of the positive semidefinite matrix $G_\tau = \Xi \text{diag}(0, \lambda_2, \dots, \lambda_N) \Xi^T$, where λ_i are its positive eigenvalues and $\Xi = [e \ \xi_2 \ \dots \ \xi_N]$ such that $\langle e, \xi_i \rangle = 0$ for all i , then:

$$(G_\tau + \zeta ee^T)^{-1} = \Xi \text{diag}(\zeta \|e\|^2, \lambda_2, \dots, \lambda_N)^{-1} \Xi^T.$$

From (8) it follows immediately that our defined norm has the following closed form expression:

$$\|x\|_{G_\tau}^* = \sqrt{x^T G_\tau^+ x} \quad \forall x \in S, \quad (9)$$

where $G_\tau^+ = \Xi \text{diag}(0, \frac{1}{\lambda_2}, \dots, \frac{1}{\lambda_N}) \Xi^T$ denotes the pseudoinverse of the matrix G_τ .

3.1 Convergence rate: smooth case

In order to estimate the rate of convergence of our algorithm in the smooth case (Assumption 2.1) we introduce the following distance that takes into account that our algorithm is a descent method:

$$\mathcal{R}(x^0) = \max_{\{x \in S: f(x) \leq f(x^0)\}} \min_{x^* \in X^*} \|x - x^*\|_{G_\tau}^*,$$

which measures the size of the level set of f given by x^0 . We assume that this distance is finite for the initial iterate x^0 . After k iterations of the algorithm, we generate a random output $(x^k, f(x^k))$, which depends on the observed implementation of random variable:

$$\eta^k = (\mathcal{N}_0, \dots, \mathcal{N}_k).$$

Let us define the expected value of the objective function w.r.t. η^k :

$$\phi_k = E \left[f(x^k) \right].$$

We now prove the main result of this section, i.e. sublinear convergence in mean for the smooth convex case:

Theorem 3.1 *Let Assumption 2.1 hold for the optimization problem (1) and the sequence $(x^k)_{k \geq 0}$ be generated by algorithm RCD_τ . Then, we have the following sublinear rate of convergence for the expected values of the objective function:*

$$\phi_k - f^* \leq \frac{2\mathcal{R}^2(x^0)}{k}. \quad (10)$$

Proof Recall that all our iterates are feasible, i.e. $x^k \in S$. From convexity of f and the definition of the norm $\|\cdot\|_{G_\tau}$ on the subspace S , we get:

$$\begin{aligned} f(x^l) - f^* &\leq \langle \nabla f(x^l), x^l - x^* \rangle \leq \|x^l - x^*\|_{G_\tau}^* \|\nabla f(x^l)\|_{G_\tau} \\ &\leq \mathcal{R}(x^0) \cdot \|\nabla f(x^l)\|_{G_\tau} \quad \forall l \geq 0. \end{aligned}$$

Combining this inequality with (7), we obtain:

$$f(x^l) - E \left[f(x^{l+1}) \mid x^l \right] \geq \frac{(f(x^l) - f^*)^2}{2\mathcal{R}^2(x^0)},$$

or equivalently

$$E \left[f(x^{l+1}) \mid x^l \right] - f^* \leq f(x^l) - f^* - \frac{(f(x^l) - f^*)^2}{2\mathcal{R}^2(x^0)}.$$

Taking the expectation of both sides of this inequality in η_{l-1} and denoting $\Delta_l = \phi_l - f^*$ leads to:

$$\Delta_{l+1} \leq \Delta_l - \frac{\Delta_l^2}{2\mathcal{R}^2(x^0)}.$$

Dividing both sides of this inequality with $\Delta_l \Delta_{l+1}$ and taking into account that $\Delta_{l+1} \leq \Delta_l$ (see (7)), we obtain:

$$\frac{1}{\Delta_l} \leq \frac{1}{\Delta_{l+1}} - \frac{1}{2\mathcal{R}^2(x^0)} \quad \forall l \geq 0.$$

Adding these inequalities from $l = 0, \dots, k-1$ we get that $0 \leq \frac{1}{\Delta_0} \leq \frac{1}{\Delta_k} - \frac{k}{2\mathcal{R}^2(x^0)}$ from which we obtain the statement (10) of the theorem. \square

Theorem 3.1 shows that for smooth convex problem (1) algorithm RCD_τ has sub-linear rate of convergence in expectation but with a low complexity per iteration. More specifically, the complexity per iteration is $\mathcal{O}(\tau n_f + \tau)$, where n_f is the maximum cost of computing the gradient of each function f_i and τ is the cost of updating x^+ . We assume that the cost of choosing randomly a τ -tuple of indices \mathcal{N} for a given probability distribution is negligible (e.g. for $\tau = 2$ the cost is $\ln N$).

3.2 Convergence rate: strongly convex case

Additionally to the assumption of Lipschitz continuous gradient for each function f_i (see Assumption (2.1)), we now assume that the function f is also strongly convex with respect to the extended norm $\|\cdot\|_{G_\tau}^*$ with convexity parameter σ_{G_τ} on the subspace S . More precisely, the objective function f satisfies for all $x, y \in S$:

$$f(x) \geq f(y) + \langle \nabla f(y), x - y \rangle + \frac{\sigma_{G_\tau}}{2} (\|x - y\|_{G_\tau}^*)^2. \quad (11)$$

We now derive linear convergence estimates for algorithm RCD_τ under the additional strong convexity assumption:

Theorem 3.2 *Let Assumption (2.1) hold and additionally we assume f to be also σ_{G_τ} -strongly convex function with respect to norm $\|\cdot\|_{G_\tau}^*$ (see (11)). Then, for the sequence $(x^k)_{k \geq 0}$ generated by algorithm RCD_τ we have the following linear estimate for the convergence rate in expectation:*

$$\phi_k - f^* \leq (1 - \sigma_{G_\tau})^k \left(f(x^0) - f^* \right). \quad (12)$$

Proof From (7) we have:

$$2 \left(f(x^k) - E \left[f(x^{k+1}) \mid x^k \right] \right) \geq \|\nabla f(x^k)\|_{G_\tau}^2.$$

On the other hand, minimizing both sides of inequality (11) over $x \in S$ we have:

$$\|\nabla f(y)\|_{G_\tau}^2 \geq 2\sigma_{G_\tau}(f(y) - f^*) \quad \forall y \in S$$

and for $y = x^k$ we get:

$$\|\nabla f(x^k)\|_{G_\tau}^2 \geq 2\sigma_{G_\tau} \left(f(x^k) - f^* \right).$$

Combining the first inequality with the last one, and taking expectation in η_{k-1} in both sides, we prove the statement of the theorem. \square

We notice that if f_i 's are strongly convex functions with respect to the Euclidian norm, with convexity parameter σ_i , i.e.:

$$f_i(x_i) \geq f_i(y_i) + \langle \nabla f_i(y_i), x_i - y_i \rangle + \frac{\sigma_i}{2} |x_i - y_i|^2 \quad \forall x_i, y_i, \quad i \in [N],$$

then the whole function $f = \sum_i f_i$ is also strongly convex w.r.t. the extended norm induced by the positive definite matrix D_σ , where $\sigma = [\sigma_1 \cdots \sigma_N]^T$, i.e.:

$$f(x) \geq f(y) + \langle \nabla f(y), x - y \rangle + \frac{1}{2} \|x - y\|_{D_\sigma}^2 \quad \forall x, y.$$

Note that in this extended norm $\|\cdot\|_{D_\sigma}$ the strongly convex parameter of the function f is equal to 1. It follows immediately that the function f is also strongly convex with respect to the norm $\|\cdot\|_{G_\tau}^*$ with the strongly convex parameter σ_{G_τ} satisfying:

$$\sigma_{G_\tau} D_\sigma^{-1} \preceq G_\tau + \zeta e e^T,$$

for some $\zeta \geq 0$. In conclusion, the strong convexity parameter σ_{G_τ} needs to satisfy the following LMI:

$$\sigma_{G_\tau} I_N \preceq D_\sigma^{1/2} (G_\tau + \zeta e e^T) D_\sigma^{1/2}. \quad (13)$$

Finally, we should notice that we can also easily derive results showing that the problem is approximately solved with high probability in both situations, smooth and/or strongly convex case, see e.g. [7] for details.

3.3 How the number of updated blocks τ enters into the convergence rates

Note that matrix G_τ depends directly on the number of components τ to be updated and therefore, $R(x^0)$ is also depending on τ . Moreover, the convergence rate can be explicitly expressed in terms of τ for some specific choices for probabilities and for the graph \mathcal{G} . In particular, let us assume a complete graph \mathcal{G} and that we know some constants $R_i > 0$ such that for any x satisfying $f(x) \leq f(x^0)$ there exists an $x^* \in X^*$ such that:

$$|x_i - x_i^*| \leq R_i \quad \forall i \in [N],$$

and recall that $L = [L_1 \cdots L_N]^T$ and D_L is the diagonal matrix with entries on the diagonal given by the vector L . Moreover, let us consider probabilities depending on the Lipschitz constants L_i for any path $\mathcal{N} \in \mathcal{P}_\tau$ of τ vertices in the complete graph \mathcal{G} , defined as:

$$p_{\mathcal{N}}^L = \frac{\sum_{i \in \mathcal{N}} 1/L_i}{\sum_{\mathcal{N} \in \mathcal{P}_\tau} \sum_{i \in \mathcal{N}} 1/L_i}. \quad (14)$$

Theorem 3.3 *Under Assumption 2.1 and for the choice of probabilities (14) on a complete graph \mathcal{G} , the following sublinear convergence rate for algorithm RCD_τ in the expected values of the objective function is obtained:*

$$\phi_k - f^* \leq \frac{N-1}{\tau-1} \cdot \frac{2 \sum_i L_i R_i^2}{k}.$$

Proof Using the definition of the indicator function $1_{\mathcal{N}}$, we can see that:

$$\begin{aligned} \Sigma_\tau^{-1} &= \sum_{\mathcal{N} \in \mathcal{P}_\tau} \sum_{i \in \mathcal{N}} \frac{1}{L_i} = \sum_{j=1}^{\binom{\tau}{N}} \sum_{i \in \mathcal{N}_j} \frac{1}{L_i} = \sum_{j=1}^{\binom{\tau}{N}} \sum_{i=1}^N 1_{\mathcal{N}_j}(i) \frac{1}{L_i} \\ &= \sum_{i=1}^N \frac{1}{L_i} \left(\sum_{j=1}^{\binom{\tau}{N}} 1_{\mathcal{N}_j}(i) \right) = \sum_{i=1}^N \frac{1}{L_i} \binom{\tau-1}{N-1}. \end{aligned}$$

Thus, using $G_\tau = \sum_{\mathcal{N} \in \mathcal{P}_\tau} p_{\mathcal{N}} G_{\mathcal{N}}$ and the expression of $G_{\mathcal{N}}$ given in (6), we can derive that matrix G_τ has the following expression:

$$\begin{aligned} G_\tau &= \frac{1}{\Sigma_\tau^{-1}} \sum_{\mathcal{N} \in \mathcal{P}_\tau} \left[\sum_{i \in \mathcal{N}} \frac{1}{L_i} D_{L_{\mathcal{N}}}^{-1} - L_{\mathcal{N}}^{-1} (L_{\mathcal{N}}^{-1})^T \right] \\ &= \frac{1}{\Sigma_\tau^{-1}} \sum_{j=1}^{\binom{\tau}{N}} \left[\sum_{i \in \mathcal{N}_j} \frac{1}{L_i} D_{L_{\mathcal{N}_j}}^{-1} - L_{\mathcal{N}_j}^{-1} (L_{\mathcal{N}_j}^{-1})^T \right] \\ &= \frac{1}{\Sigma_\tau^{-1}} \binom{\tau-2}{N-2} \left[\sum_{i=1}^N \frac{1}{L_i} D_L^{-1} - L^{-1} (L^{-1})^T \right] \\ &= \frac{\tau-1}{N-1} \left[D_L^{-1} - \frac{1}{e^T L^{-1}} L^{-1} (L^{-1})^T \right], \end{aligned}$$

Using the previous expression for G_τ in (9), we get:

$$(\|x\|_{G_\tau}^*)^2 = \frac{N-1}{\tau-1} \sum_{i \in [N]} L_i |x_i|^2 \quad \forall x \in S. \quad (15)$$

Using the definition for R_i and the expression (15) for the norm on S , we obtain:

$$\mathcal{R}^2(x^0) \leq \mathcal{R}^2(R) = \frac{N-1}{\tau-1} \sum_{i \in [N]} L_i R_i^2,$$

where $R = [R_1 \cdots R_N]^T$. Using this expression for $\mathcal{R}(x^0)$ in Theorem 3.1 we get the statement of the theorem. \square

Note that for $\tau = N$ we recover the convergence rate of the full gradient method, while for $\tau = 2$ we get the convergence rate of the basic random coordinate descent method. Thus, the theoretical speedup of the parallel algorithm RCD_τ in terms of the number of iterations needed to approximately solve the problem, as compared to the basic random coordinate descent method, is equal in this case to τ - the number of components to be updated (number of processors available). This result also shows that the speedup achieved by our method on the class of separable problems with coupling constraints is the same as for separable problems without coupling constraints.

For the strongly convex case, we note that combining the Lipschitz inequality (3) with the strong convex inequality (11) we get:

$$\sum_{i \in [N]} L_i |x_i - y_i|^2 \geq \sigma_{G_\tau} (\|x - y\|_{G_\tau}^*)^2 \quad \forall x, y \in S.$$

Now, if we consider e.g. a complete graph and the probabilities given in (14), then using the expression for the norm $\|\cdot\|_{G_\tau}$ given in (15) we obtain $\sigma_{G_\tau} \leq \frac{\tau-1}{N-1}$. Thus, in the strongly convex case the linear convergence rate in expectation (12) can be also expressed in terms of τ .

4 Design of optimal probabilities

We have several choices for the probabilities $(p_{\mathcal{N}})_{\mathcal{N} \in \mathcal{P}_\tau}$ corresponding to paths of τ vertices in the complete graph \mathcal{G} , which the randomized coordinate descent algorithm RCD_τ depends on. For example, we can choose probabilities dependent on the Lipschitz constants L_i :

$$p_{\mathcal{N}}^\alpha = \frac{\sum_{i \in \mathcal{N}} L_i^\alpha}{\Sigma_\tau^\alpha}, \quad \Sigma_\tau^\alpha = \sum_{\mathcal{N} \in \mathcal{P}_\tau} \sum_{i \in \mathcal{N}} L_i^\alpha, \quad \alpha \in \mathbb{R}. \quad (16)$$

Note that for $\alpha = 0$ we recover the uniform probabilities. Finally, we can design optimal probabilities from the convergence rate of the method. From the definition of the constants R_i it follows that:

$$\mathcal{R}(x^0) \leq \mathcal{R}(R) = \max_{x \in S, |x_i| \leq R_i} \|x\|_{G_\tau}^*, \quad \text{with } G_\tau = \sum_{\mathcal{N} \in \mathcal{P}_\tau} p_{\mathcal{N}} G_{\mathcal{N}}.$$

We have the freedom to choose the matrix G_τ that depends linearly on the probabilities p_N . For the probabilities $(p_N)_{N \in \mathcal{P}_\tau}$ corresponding to paths of τ vertices in the complete graph \mathcal{G} we define the following set of matrices:

$$\mathcal{M} = \left\{ G_\tau : G_\tau = \sum_{N \in \mathcal{P}_\tau} p_N G_N, \quad G_N = D_{L_N}^{-1} - \frac{1}{\sum_{i \in N} 1/L_i} L_N^{-1} (L_N^{-1})^T \right\}.$$

Therefore, we search for the probabilities p_N that are the optimal solution of the following optimization problem:

$$\mathcal{R}^*(x^0) = \min_{p_N} \mathcal{R}(x^0) \leq \min_{p_N} \mathcal{R}(R) = \min_{G_\tau \in \mathcal{M}} \max_{x \in S, \|x_i\| \leq R_i} \|x\|_{G_\tau}^*.$$

Let us define $R = [R_1 \cdots R_N]^T$ and $\nu = [\nu_1 \cdots \nu_N]^T$. In the next theorem we derive an easily computed upper bound on $\mathcal{R}^*(x_0)$ and we provide a way to suboptimally select the probabilities p_N :

Theorem 4.1 *Let Assumption 2.1 hold. Then, a suboptimal choice of probabilities $(p_N)_{N \in \mathcal{P}_\tau}$ can be obtained as a solution of the following SDP problem whose optimal value is an upper bound on $\mathcal{R}^*(x_0)$, i.e.:*

$$(\mathcal{R}^*(x_0))^2 \leq \min_{G_\tau \in \mathcal{M}, \zeta \geq 0, \nu \geq 0} \left\{ \langle \nu, R^2 \rangle : \begin{bmatrix} G_\tau + \zeta ee^T & I_N \\ I_N & D_\nu \end{bmatrix} \succeq 0 \right\}. \quad (17)$$

Proof Using the definition of $\mathcal{R}(R)$ and of the norm $\|\cdot\|_{G_\tau}^*$ we get:

$$\begin{aligned} \min_{p_N} (\mathcal{R}(R))^2 &\leq \min_{G_\tau \in \mathcal{M}} \max_{x \in S, \|x_i\| \leq R_i} (\|x\|_{G_\tau}^*)^2 \\ &= \min_{G_\tau \in \mathcal{M}} \max_{x \in S, \|x_i\| \leq R_i} \min_{\zeta \geq 0} \langle (G_\tau + \zeta ee^T)^{-1} x, x \rangle \\ &= \min_{G_\tau \in \mathcal{M}, \zeta \geq 0} \max_{x \in S, \|x_i\| \leq R_i} \langle (G_\tau + \zeta ee^T)^{-1} x, x \rangle \\ &= \min_{G_\tau \in \mathcal{M}, \zeta \geq 0} \max_{x \in S, \|x_i\| \leq R_i} \langle (G_\tau + \zeta ee^T)^{-1}, xx^T \rangle \\ &= \min_{G_\tau \in \mathcal{M}, \zeta \geq 0} \max_{X \in \mathcal{X}} \langle (G_\tau + \zeta ee^T)^{-1}, X \rangle, \end{aligned}$$

where $\mathcal{X} = \{X : X \succeq 0, \text{rank} X = 1, \langle ee^T, X \rangle = 0, \langle X, E_{ii} \rangle \leq R_i^2 \ \forall i\}$ and $E_{ii} = e_i e_i^T$. Using the well-known relaxation from the SDP literature, we have:

$$\min_{p_N} (\mathcal{R}(R))^2 \leq \min_{G_\tau \in \mathcal{M}, \zeta \geq 0} \max_{X \in \mathcal{X}_r} \langle (G_\tau + \zeta ee^T)^{-1}, X \rangle,$$

where $\mathcal{X}_r = \{X : X \succeq 0, \langle ee^T, X \rangle = 0, \langle X, E_{ii} \rangle \leq R_i^2 \ \forall i\}$, i.e. we have removed the rank constraint: $\text{rank} X = 1$. Then, the right hand side of the previous optimization problem can be reformulated equivalently, using Lagrange multipliers, as follows:

$$\begin{aligned} &\min_{G_\tau \in \mathcal{M}, \zeta \geq 0} \max_{X \in \mathcal{X}_r} \langle (G_\tau + \zeta ee^T)^{-1}, X \rangle \\ &= \min_{G_\tau \in \mathcal{M}, \zeta, \nu \geq 0, Z \geq 0, \theta \in \mathbb{R}} \max_{X \in \mathbb{R}^{N \times N}} \left[\langle (G_\tau + \zeta ee^T)^{-1} + Z + \theta ee^T, X \rangle \right. \\ &\quad \left. + \sum_{i=1}^N \nu_i (R_i^2 - \langle X, E_{ii} \rangle) \right]. \end{aligned}$$

where $\nu = [\nu_1 \cdots \nu_N]^T$. Rearranging the terms, we can write the previous convex problem equivalently:

$$\begin{aligned} & \min_{G_\tau \in \mathcal{M}, \theta \in \mathbb{R}, Z \succeq 0, \zeta, \nu \geq 0} \left[\sum_i \nu_i R_i^2 \right. \\ & \quad \left. + \max_{X \in \mathbb{R}^{N \times N}} \langle (G_\tau + \zeta ee^T)^{-1} + Z + \theta ee^T - \sum_i \nu_i E_{ii}, X \rangle \right] \\ & = \min_{(G_\tau, Z, \zeta, \nu, \theta) \in \mathcal{F}} \sum_i \nu_i R_i^2, \end{aligned}$$

where the feasible set is described as: $\mathcal{F} = \{(G_\tau, Z, \zeta, \nu, \theta) : G_\tau \in \mathcal{M}, \theta \in \mathbb{R}, Z \succeq 0, \zeta, \nu \geq 0, (G_\tau + \zeta ee^T)^{-1} + Z + \theta ee^T - \sum_i \nu_i E_{ii} = 0\}$. Moreover, since $Z \succeq 0$, the feasible set can be rewritten as:

$$\{(G_\tau, \zeta, \nu, \theta) : G_\tau \in \mathcal{M}, \theta \in \mathbb{R}, \zeta, \nu \geq 0, \sum_i \nu_i E_{ii} - (G_\tau + \zeta ee^T)^{-1} - \theta ee^T \succeq 0\}.$$

We observe that we can take $\theta = 0$ and then we get the feasible set:

$$\{(G_\tau, \zeta, \nu) : G_\tau \in \mathcal{M}, \zeta, \nu \geq 0, G_\tau + \zeta ee^T \succeq D_\nu^{-1}\}.$$

In conclusion, we obtain the following SDP:

$$\min_{P_N} (\mathcal{R}(R))^2 \leq \min_{G_\tau \in \mathcal{M}, \zeta, \nu \geq 0, G_\tau + \zeta ee^T \succeq D_\nu^{-1}} \langle \nu, R^2 \rangle.$$

Finally, the SDP (17) is obtained from Schur complement formula applied to the previous optimization problem. \square

Since we assume a connected graph \mathcal{G} , we have that $\lambda_1(G_\tau) = 0$ is simple and consequently $\lambda_2(G_\tau) > 0$. Then, the following equivalence holds:

$$G_\tau + t \frac{ee^T}{\|e\|^2} \succeq tI_N \quad \text{if and only if} \quad t \leq \lambda_2(G_\tau), \quad (18)$$

since the spectrum of the matrix $G_\tau + \zeta ee^T$ is $\{\zeta \|e\|^2, \lambda_2(G_\tau), \dots, \lambda_N(G_\tau)\}$. It follows that $\zeta = \frac{t}{\|e\|^2}$, $\nu_i = \frac{1}{t}$ for all i , and G_τ such that $t \leq \lambda_2(G_\tau)$ is feasible for the SDP problem (17). We conclude that:

$$\begin{aligned} \left(\mathcal{R}^*(x^0) \right)^2 & \leq \min_{G_\tau \in \mathcal{M}, \zeta, \nu \geq 0, G_\tau + \zeta ee^T \succeq D_\nu^{-1}} \langle \nu, R^2 \rangle \\ & \leq \min_{G_\tau \in \mathcal{M}, t \leq \lambda_2(G_\tau)} \sum_{i=1}^N R_i^2 \frac{1}{t} \leq \frac{\sum_i R_i^2}{\lambda_2(G_\tau)} \quad \forall G_\tau \in \mathcal{M}. \end{aligned} \quad (19)$$

Then, according to Theorem 3.1 we obtain the following upper bound on the rate of convergence for the expected values of the objective function in the smooth convex case:

$$\phi_k - f^* \leq \frac{2 \sum_{i=1}^N R_i^2}{\lambda_2(G_\tau) \cdot k} \quad \forall G_\tau \in \mathcal{M}. \quad (20)$$

From the convergence rate for algorithm RCD_τ given in (20) it follows that we can choose the probabilities such that we maximize the second eigenvalue of G_τ :

$$\max_{G_\tau \in \mathcal{M}} \lambda_2(G_\tau).$$

In conclusion, in order to find some suboptimal probabilities $(p_N)_{N \in \mathcal{P}_\tau}$, we can solve the following simpler SDP problem than the one given in (17):

Corollary 4.1 *From (18) we get for the smooth case that a suboptimal choice of probabilities $(p_N)_{N \in \mathcal{P}_\tau}$ can be obtained as a solution of the following SDP problem:*

$$p_N^* = \arg \max_{t, G_\tau \in \mathcal{M}} \left\{ t : G_\tau \succeq t \left(I_N - \frac{ee^T}{\|e\|^2} \right) \right\}. \quad (21)$$

Note that the matrices on both sides of the LMI from (21) have the common eigenvalue zero associated to the eigenvector e , so that this LMI has empty interior which can cause problems for some classes of interior point methods. We can overcome this problem by replacing the LMI constraint in (21) with the following equivalent LMI:

$$G_\tau + \frac{ee^T}{\|e\|^2} \succeq t \left(I_N - \frac{ee^T}{\|e\|^2} \right).$$

Finally, when the functions f_i are σ_i -strongly convex, from Theorem 3.2 and the LMI (13) it follows that in order to get a better convergence rate we need to search for σ_{G_τ} as large as possible. Therefore, we get the following result:

Corollary 4.2 *For the strongly convex case the optimal probabilities are chosen as the solution of the following SDP problem:*

$$p_N^* = \arg \max_{\sigma_{G_\tau}, \zeta \geq 0, G_\tau \in \mathcal{M}} \left\{ \sigma_{G_\tau} : \sigma_{G_\tau} I_N \preceq D_\sigma^{1/2} (G_\tau + \zeta ee^T) D_\sigma^{1/2} \right\}. \quad (22)$$

In [17], the authors propose a (center-free) distributed scaled gradient method in the form (5) to solve the separable optimization problem (1) with strongly convex objective function, where at each iteration the full gradient needs to be computed. A similar rate of convergence is obtained as in Theorem 3.2 under the Lipschitz and strong convexity assumption on f , where the weights are designed by solving an SDP in the form (22). Our randomized algorithm also belongs to this class of methods and for $\tau = N$ we recover a version of the method in [17]. Moreover, our convergence analysis covers the smooth case, i.e. without the strong convexity assumption.

5 Applications

Problem (1) arises in many real applications, e.g. image processing [3, 4, 19], resource allocation [7, 17] and coordination in multi-agent systems [7, 18]. For example, we can interpret (1) as N agents exchanging n goods to minimize a total cost, where the constraint $\sum_i x_i = 0$ is the equilibrium or market clearing constraint. In this context $[x_i]_j \geq 0$ means that agent i receives $[x_i]_j$ of good j from exchange and $[x_i]_j < 0$ means that agent i contributes $|[x_i]_j|$ of good j to exchange. It can be also viewed as the distributed dynamic energy management problem: N devices

exchange power in time periods $t = 1, \dots, n$. Furthermore, $x_i \in \mathbb{R}^n$ is the power flow profile for device i and $f_i(x_i)$ is the cost of profile x_i (and usually encodes constraints). In this application the constraint $\sum_i x_i = 0$ represents the energy balance (in each time period).

Problem (1) can also be seen as the dual corresponding to an optimization of a sum of convex functions. Consider the following primal convex optimization problem that arises in many engineering applications:

$$g^* = \min_{v \in \bigcap_{i=1}^N Q_i} g_1(v) + \dots + g_N(v), \quad (23)$$

where g_i are all σ_i -strongly convex functions and $Q_i \subseteq \mathbb{R}^n$ are convex sets. Denote with v^* the unique optimal solution of problem (23). This problem can be reformulated as:

$$\min_{u_i \in Q_i, u_i = v \ \forall i \in [N]} g_1(u_1) + \dots + g_N(u_N).$$

Let us define $u = [u_1^T \dots u_N^T]^T$ and $g(u) = g_1(u_1) + \dots + g_N(u_N)$. By duality, using the Lagrange multipliers x_i for the constraints $u_i = v$, we obtain the equivalent convex problem (1), where $f_i(x_i) = \tilde{g}_i^*(x_i)$ and \tilde{g}_i^* is the convex conjugate of the function $\tilde{g}_i = g_i + 1_{Q_i}$, i.e.

$$f_i(x_i) = \max_{u_i \in Q_i} \langle x_i, u_i \rangle - g_i(u_i) \quad \forall i. \quad (24)$$

Further we have $f^* + g^* = 0$. Note that if g_i is σ_i -strongly convex, then the convex conjugate f_i is well-defined and has Lipschitz continuous gradient with constants $L_i = \frac{1}{\sigma_i}$ (see [11]), so that Assumption 2.1 holds. A particular application is the problem of finding the projection of a point v_0 in the intersection of the convex sets $\bigcap_{i=1}^N Q_i \subseteq \mathbb{R}^n$. This problem can be written as an optimization problem in the form:

$$\min_{v \in \bigcap_{i=1}^N Q_i} p_1 \|v - v_0\|^2 + \dots + p_N \|v - v_0\|^2,$$

where $p_i > 0$ such that $\sum_i p_i = 1$. This is a particular case of the separable problem (23). Note that since the functions $g_i(v) = p_i \|v - v_0\|^2$ are strongly convex, then f_i have Lipschitz continuous gradient with constants $L_i = 1/p_i$ for all $i \in [N]$.

We now show how we can recover an approximate primal solution for the primal problem (23) by solving the corresponding dual problem (1) with algorithm RCD_τ . Let us define for any dual variable x_i the primal variable:

$$u_i(x_i) = \arg \min_{u_i \in Q_i} g_i(u_i) - \langle x_i, u_i \rangle \quad \forall i \in [N].$$

Let us define $\hat{v}^* = e \otimes v^*$, where \otimes is the Kronecker product, and the norm $\|u\|_{D_\sigma}^2 = \sum_i \sigma_i \|u_i\|^2$. Moreover, let $\sigma_{\min} = \min_i \sigma_i$ and λ_N the largest eigenvalue of G_τ . Furthermore, for simplicity of the presentation we consider the initial starting point $x^0 = 0$. Then, we can derive convergence estimates on primal infeasibility and suboptimality for (23).

Theorem 5.1 *For the convex optimization problem (23) we assume that all functions g_i are σ_i -strongly convex. Let x^k be the sequence generated by algorithm RCD_τ for solving the corresponding dual problem (1) and the primal sequence $u^k = u(x^k)$. Then, we have the following convergence estimates in the expected values on primal infeasibility and suboptimality:*

$$E \left[\|u^k - \hat{v}^*\|_{D_\sigma}^2 \right] \leq \frac{4R^2(x^0)}{k} \quad \text{and} \quad E \left[|g(u^k) - g^*| \right] \leq \frac{4R^2(x^0)\lambda_N}{\sigma_{\min}\sqrt{k}}.$$

Proof Since all the functions g_i are σ_i -strongly convex, then the objective function $\sum_{i=1}^N g_i(u_i) - \langle x_i, u_i \rangle$ is also 1-strongly convex in the variable u w.r.t. the norm $\|u\|_{D_\sigma}^2 = \sum_i \sigma_i \|u_i\|^2$. Using this property and the expression of $u_i(x_i)$, we obtain the following inequalities:

$$\begin{aligned} \frac{1}{2} \|u(x) - \hat{v}^*\|_{D_\sigma}^2 &= \sum_{i=1}^N \frac{\sigma_i}{2} \|u_i(x_i) - v^*\|^2 \\ &\leq \left(\sum_{i=1}^N g_i(v^*) - \langle x_i, v^* \rangle \right) - \left(\sum_{i=1}^N g_i(u_i(x_i)) - \langle x_i, u_i(x_i) \rangle \right) \\ &= (-f^*) - (-f(x)) = f(x) - f^* \quad \forall x \in S. \end{aligned} \quad (25)$$

Now, let us consider for x the sequence x^k generated by algorithm RCD_τ and let $u^k = u(x^k)$. We note that $u_i^{k+1} = u_i^k$ for all $i \in [N] \setminus \mathcal{N}_k$ and $u_i^{k+1} = u_i(x_i^{k+1})$ for all $i \in \mathcal{N}_k$. Taking expectation over the entire history η^k and using Theorem 3.1, we get an estimate on primal infeasibility:

$$E[\|u^k - \hat{v}^*\|_{D_\sigma}^2] \leq 2E[f(x^k) - f^*] = 2(\phi_k - f^*) \leq \frac{4R^2(x^0)}{k}.$$

Moreover, for deriving estimates on primal suboptimality, we first observe:

$$\|u\|_{G_\tau} \leq \frac{\lambda_N}{\sigma_{\min}} \|u\|_{D_\sigma} \quad \forall u,$$

and combining with (25) we get:

$$\|u(x) - \hat{v}^*\|_{G_\tau} \leq \frac{\lambda_N \sqrt{2(f(x) - f^*)}}{\sigma_{\min}}. \quad (26)$$

For the left hand side suboptimality, we proceed as follows:

$$\begin{aligned} f(x^*) &= \langle x^*, \hat{v}^* \rangle - g(\hat{v}^*) \\ &= \max_{u_i \in Q_i} \langle x^*, u \rangle - g(u) \geq \langle x^*, u(x) \rangle - g(u(x)), \end{aligned}$$

which leads to the following relation:

$$\begin{aligned} g(u(x)) - g^* &\geq \langle x^*, u(x) - \hat{v}^* \rangle \geq -\|x^*\|_{G_\tau}^* \|\hat{v}^* - u(x)\|_{G_\tau} \\ &\stackrel{(26)}{\geq} -\|x^*\|_{G_\tau}^* \frac{\lambda_N \sqrt{2(f(x) - f^*)}}{\sigma_{\min}} \quad \forall x \in S. \end{aligned} \quad (27)$$

Secondly, from the definition of the dual function, we have:

$$f(x) = \langle x, u(x) \rangle - g(u(x)) \quad \forall x \in S.$$

Subtracting $f^* = f(x^*)$ from both sides and using the complementarity condition $\langle x^*, u(x^*) \rangle = 0$, where $u(x^*) = \hat{v}^*$, we get the following relations:

$$\begin{aligned}
g(u(x)) - g^* &= \langle x, u(x) \rangle - f(x) + f^* \\
&= f(x^*) - f(x) + \langle x - x^*, u(x^*) \rangle + \langle x, u(x) - u(x^*) \rangle \\
&= f(x^*) + \langle x - x^*, \nabla f(x^*) \rangle - f(x) + \langle x, u(x) - u(x^*) \rangle \\
&\leq \langle x, u(x) - u(x^*) \rangle \leq \|x\|_{G_\tau}^* \|\hat{v}^* - u(x)\|_{G_\tau} \\
&\leq (\|x - x^*\|_{G_\tau}^* + \|x^*\|_{G_\tau}^*) \|u(x) - \hat{v}^*\|_{G_\tau} \\
&\stackrel{(26)}{\leq} (\|x - x^*\|_{G_\tau}^* + \|x^*\|_{G_\tau}^*) \frac{\lambda_N \sqrt{2(f(x) - f^*)}}{\sigma_{\min}},
\end{aligned}$$

valid for all $x \in S$ and $x^* \in X^*$, where in the first inequality we used convexity of the function f and the relation $\nabla f(x) = u(x)$, and in the second inequality the Cauchy-Schwartz inequality. Now, using the definition of $R(x^0)$ and that $x^0 = 0$, replacing x with the sequence x^k in the previous derivations and taking the expectation over the entire history η^k , we obtain a bound on primal suboptimality:

$$\begin{aligned}
E[\|g(u^k) - g^*\|] &\leq E \left[\left(\|x^k - x^*\|_{G_\tau}^* + \|x^*\|_{G_\tau}^* \right) \frac{\lambda_N \sqrt{2(f(x^k) - f^*)}}{\sigma_{\min}} \right] \\
&\leq \frac{2R(x^0)\lambda_N}{\sigma_{\min}} \sqrt{2E[f(x^k) - f^*]} \leq \frac{4R^2(x^0)\lambda_N}{\sigma_{\min}\sqrt{k}},
\end{aligned}$$

which gives us a convergence estimate for primal suboptimality for problem (23). \square

In conclusion, the expected values of the distance between the primal generated points $u_i^k \in Q_i$ and the unique optimal point $v^* \in \cap_i Q_i$ of (23), i.e. $E[\|u_i^k - v^*\|]$, is less than $\mathcal{O}(\frac{1}{\sqrt{k}})$. Similar convergence rates for other projection algorithms for solving convex feasibility problems have been derived in the literature, see e.g. [3,4].

6 Numerical experiments

In this section we report some preliminary numerical results on solving the optimization problem (1), where the functions f_i are taken as in paper [17]:

$$f_i(x_i) = \frac{1}{2}a_i(x_i - c_i)^2 + \log(1 + \exp(b_i(x_i - d_i))) \quad \forall i \in [N], \quad (28)$$

where the coefficients $a_i \geq 0, b_i, c_i$ and d_i are generated randomly with uniform distributions on $[-15, 15]$. The second derivatives of these functions have the following expressions:

$$f_i''(x_i) = a_i + \frac{b_i^2 \exp(b_i(x_i - d_i))}{(1 + \exp(b_i(x_i - d_i)))^2},$$

which have the following lower and upper bounds:

$$\sigma_i = a_i \quad \text{and} \quad L_i = a_i + \frac{1}{4}b_i^2.$$


We assume that the sum of the variables is fixed to zero, i.e.:

$$\sum_{i=1}^N x_i = 0.$$

In all our numerical tests we consider a complete graph and Lipschitz dependent probabilities as in (14) (if not specified otherwise).

In the first set of experiments, we solve a single randomly generated problem with $N = 10^4$ nodes for $\tau = 2, 4$ and 7 cores in parallel using MPI. The Fig. 1 displays the evolution of $f(x^k) - f^*$ along normalized iterations k/N of algorithm RCD_τ . From the plot we can observe that increasing the number of cores reduces substantially the number of full iterations k/N .

Fig. 1 Typical performance of algorithm RCD_τ for different numbers of updated variables (processors) τ : evolution of $f(x^k) - f^*$ along normalized iterations k/N , with $\tau = 2, 4$ and 7.



Then, we tested algorithm RCD_2 , i.e. $\tau = 2$, and thus at each iteration we choose a pair of nodes in the graph $(i, j) \in E$ with probability p_{ij} and then update only the components i and j of x as follows:

$$x_i^+ = x_i + \frac{1}{L_i + L_j} (\nabla f_j(x_j) - \nabla f_i(x_i)), \quad x_j^+ = x_j + \frac{1}{L_i + L_j} (\nabla f_i(x_i) - \nabla f_j(x_j)).$$

We consider three choices of the probabilities in Fig. 2 for algorithm RCD_2 : uniform probability, probabilities depending on Lipschitz constants as given in (14) and optimal probabilities obtained from solving the SDP (21). As we expected, the method based on choosing the optimal probabilities has the fastest convergence.

Finally, we compare our algorithm RCD_2 , i.e. $\tau = 2$, for two choices for the probabilities p_{ij} (uniform and Lipschitz dependent probabilities (14)) with the full gradient method and the center-free gradient method with Metropolis weights proposed in [17]. The global Lipschitz constant in the full gradient is taken as $L_{\max} = \max_i L_i$. Note that the computations of the local Lipschitz constant L_i required by algorithm RCD_τ can be done locally in each node for the corresponding function f_i , while computing the global Lipschitz constant L_{\max} for projected

Fig. 2 Evolution of $f(x^k) - f^*$ along full iterations for algorithm RCD_2 for different choices for the probability: uniform probabilities, Lipschitz dependent probabilities as given in (14) and optimal probabilities obtained from solving the SDP problem (21).





Fig. 3 Evolution of $f(x^k) - f^*$ along full iterations for the methods: projected gradient, center-free gradient method with Metropolis weights [17] and algorithm RCD_τ with uniform and Lipschitz dependent probabilities for $\tau = 2$.



gradient method on problems of very large dimension is difficult. We have also implemented the center-free gradient method with Metropolis weights from [17]:

$$x_i^+ = x_i + \sum_{j \in \mathcal{N}_i} w_{ij} (\nabla f_j(x_j) - \nabla f_i(x_i)) \quad \forall i \in [N],$$

where \mathcal{N}_i are the neighbors of node i in the graph and the weights satisfy the relation: $\sum_{j \in \mathcal{N}_i} w_{ij} = 0$. In Fig. 3 we plot the evolution of $f(x^k) - f^*$ along full iterations k/N for the following methods: the center-free gradient algorithm from [17] with the Metropolis weights, the full projected gradient algorithm and the RCD_2 algorithm with uniform and Lipschitz dependent probabilities. We clearly see that the best accuracy is achieved by the RCD_2 algorithm with Lipschitz dependent probabilities.

7 Conclusions

In this paper we have derived parallel random coordinate descent methods for minimizing linearly constrained convex problems over networks. Since we have coupled constraints in the problem, we have devised an algorithm that updates in parallel $\tau \geq 2$ (block) components per iteration. We have proved that for this method we obtain in expectation an ϵ -accurate solution in at most $\mathcal{O}(\frac{N}{\tau\epsilon})$ iterations and thus the convergence rate depends linearly on the number of (block) components to be updated. Preliminary numerical results show that the parallel coordinate descent method with $\tau > 2$ accelerates on its basic counterpart corresponding to $\tau = 2$. For strongly convex functions the new method converges linearly. We have also provided SDP formulations that enable us to choose the probabilities in an optimal fashion.

References

1. A. Beck, *The 2-coordinate descent method for solving double-sided simplex constrained minimization problems*, Journal of Optimization Theory and Applications, 162(3), 892–919, 2014.
2. A. Beck and L. Tetruashvili, *On the convergence of block coordinate descent type methods*, SIAM Journal on Optimization, 23(4), 2037–2060, 2013.
3. H. Bauschke, J.M. Borwein, *On projection algorithms for solving convex feasibility problems*, SIAM Review, 38(3), 367–426, 1996.
4. P. L. Combettes, *The convex feasibility problem in image recovery*, in Advances in Imaging and Electron Physics (P. Hawkes Ed.), 95, 155–270, Academic Press, 1996.
5. H. Ishii, R. Tempo and E. Bai, *A web aggregation approach for distributed randomized pagerank algorithms*, IEEE Transactions Automatic Control, 57, 2703–2717, 2012.
6. Ji Liu and S. Wright, *Asynchronous stochastic coordinate descent: parallelism and convergence properties*, SIAM Journal on Optimization, 25(1), 2014.
7. I. Necoara, *Random coordinate descent algorithms for multi-agent convex optimization over networks*, IEEE Transactions Automatic Control, 58(8), 2001–2012, 2013.
8. I. Necoara and A. Patrascu, *A random coordinate descent algorithm for optimization problems with composite objective function and linear coupled constraints*, Computational Optimization and Applications, 57(2), 307–337, 2014.
9. I. Necoara and D. Clipici, *Parallel coordinate descent methods for composite minimization: convergence analysis and error bounds*, SIAM Journal on Optimization, 1–29, 2016 (<http://arxiv.org/abs/1312.5302>).
10. I. Necoara and D. Clipici, *Efficient parallel coordinate descent algorithm for convex optimization problems with separable constraints: application to distributed MPC*, Journal of Process Control, 23(3), 243–253, 2013
11. Y. Nesterov, *Introductory Lectures on Convex Optimization: A Basic Course*, Boston, Kluwer, 2004.
12. Y. Nesterov, *Efficiency of coordinate descent methods on huge-scale optimization problems*, SIAM Journal on Optimization, 22(2), 341–362, 2012.
13. Z. Qin, K. Scheinberg and D. Goldfarb, *Efficient block-coordinate descent algorithms for the group lasso*, Mathematical Programming Computation, 5(2), 143–169, 2013.
14. P. Richtarik and M. Takac, *Iteration complexity of randomized block-coordinate descent methods for minimizing a composite function*, Mathematical Programming, 144(1-2), 1–38, 2014.
15. P. Richtarik and M. Takac, *Parallel coordinate descent methods for big data optimization*, Mathematical Programming, 1–52, 2015.
16. P. Tseng and S. Yun, *A Block-coordinate gradient descent method for linearly constrained nonsmooth separable optimization*, Journal of Optimization Theory and Applications, 140, 513–535, 2009.
17. L. Xiao and S. Boyd, *Optimal scaling of a gradient method for distributed resource allocation*, Journal of Optimization Theory and Applications, 129(3), 469–488, 2006.

-
18. K. You and L. Xie, *Network topology and communication data rate for consensusability of discrete-time multi-agent systems*, IEEE Transactions Automatic Control, 56(10), 2262–2275, 2011.
 19. S. Wright, *Accelerated block coordinate relaxation for regularized optimization*, SIAM Journal on Optimization, 22(1), 159–186, 2012.