Erdős-Feller-Kolmogorov-Petrowsky law of the iterated logarithm for self-normalized martingales: a game-theoretic approach

Takeyuki Sasai, Kenshi Miyabe[†] and Akimichi Takemura^{*}

April, 2015

Abstract

We prove an Erdős–Feller–Kolmogorov–Petrowsky law of the iterated logarithm for self-normalized martingales. Our proof is given in the framework of the game-theoretic probability of Shafer and Vovk. As many other game-theoretic proofs, our proof is self-contained and explicit.

Keywords and phrases: Bayesian strategy, constant-proportion betting strategy, lower class, upper class, self-normalized processes.

1 Main Result

Let S_n be a martingale with respect to a filtration $\{\mathcal{F}_n\}_{n=0}^{\infty}$ and let $x_n = S_n - S_{n-1}$ be the martingale difference. On some regularity conditions on the growth of $|x_n|$, various versions of the law of the iterated logarithm (LIL) have been given in literature. In particular the Erdős–Feller–Kolmogorov–Petrowsky law of the iterated logarithm (EFKP-LIL [16, Chapter 5.2]) is an important extension of LIL. Erdős [6] proved EFKP-LIL for symmetric Bernoulli random variables. EFKP-LIL has been generalized by Feller [7] for bounded and independent random variables and [8] (see also Bai [1]) for the i.i.d. case. Further, EFKP-LIL has been generalized for martingales by Strassen [19], Jain, Jogdeo and Stout [10], Philipp and Stout [15], Einmahl and Mason [5] and Berkes, Hörmann and Weber [2]. In particular, Einmahl and Mason [5] proved a martingale analogue of Feller's result in [7], just as Stout [18] obtained a martingale analogue of Kolmogorov's result in [11].

For self-normalized processes, EFKP-LIL was derived by [9, 3] in the i.i.d. case. However EFKP-LIL has not been derived in the martingale case, even though de la Peña, Klass and Lai [4] obtained the usual LIL. The purpose of this paper is to prove EFKP-LIL for self-normalized martingales. For a positive non-decreasing continuous function $\psi(\lambda)$ let

$$I(\psi) := \int_{1}^{\infty} \psi(\lambda) e^{-\psi(\lambda)^{2}/2} \frac{d\lambda}{\lambda}.$$
 (1)

We state our main theorem.

^{*}Graduate School of Information Science and Technology, University of Tokyo

[†]School of Science and Technology, Meiji University

Theorem 1.1. Let S_n , $n = 1, 2, ..., be a martingale with <math>S_0 = 0$ and $x_n = S_n - S_{n-1}$ be a martingale difference with respect to a filtration $\{\mathcal{F}_n\}_{n=0}^{\infty}$ such that

$$|x_n| \leq c_n \ a.s.$$

for some \mathcal{F}_{n-1} -measurable random variable c_n . Let

$$A_n^2 := \sum_{i=1}^n x_i^2 \ge 0$$

and let ψ be a positive non-decreasing continuous function.

If $I(\psi) < \infty$, then

$$P\left(S_n < A_n \psi(A_n^2) \ a.a. \mid \lim A_n = \infty, \limsup c_n \frac{\psi(A_n^2)^3}{A_n} < \infty\right) = 1.$$
(2)

If $I(\psi) = \infty$, then

$$P\left(S_n \ge A_n \psi(A_n^2) \ i.o. \mid \lim A_n = \infty, \limsup c_n \frac{\psi(A_n^2)^3}{A_n} < \infty\right) = 1.$$
(3)

This theorem is a self-normalization of the result in Einmahl and Mason [5] and a generalization of the result in de la Peña, Klass and Lai [4]. The order of growth $A_n/(\psi(A_n^2))^3$ for c_n is currently the best known order for EFKP-LIL even in the independent case ([2]). We call (2) the *validity* and (3) the *sharpness* of EFKP-LIL.

In (2) and (3), we are not assuming that the conditioning events happen with probability one. We can state (2) equivalently as

$$P\left(\lim A_n = \infty, \limsup c_n \frac{\psi(A_n^2)^3}{A_n} < \infty, S_n \ge A_n \psi(A_n^2) \ i.o.\right) = 0.$$
(4)

For our proof we adopt the framework of game-theoretic probability by Shafer and Vovk [17]. In a game-theoretic approach, for proving (2), we explicitly construct a non-negative martingale diverging to infinity on the event of (4).

We use the following notation throughout the paper

$$\ln_k n := \underbrace{\ln \ln \ldots \ln}_{k \text{ times}} n.$$

We also fix a small positive δ for the rest of this paper, e.g., $\delta = 0.01$. For our proof, as is often seen in the upper-lower class theory (cf. Feller [8, Lemma 1]), we can restrict our attention to ψ such that

$$\psi^{L}(n) \le \psi(n) \le \psi^{U}(n)$$
 for all sufficiently large n , (5)

where

$$\psi^{L}(n) := \sqrt{2 \ln_2 n + 3 \ln_3 n}, \quad \psi^{U}(n) := \sqrt{2 \ln_2 n + 4 \ln_3 n}.$$

Here *L* means the lower class and *U* means the upper class. It can be verified that $I(\psi^U) < \infty$ and $I(\psi^L) = \infty$.

The rest of this paper is organized as follows. In Section 2 we give a game-theoretic statement corresponding to our main theorem. In Section 3 we give a proof of the validity and in Section 4 we give a proof of the sharpness.

2 Preliminaries on Game-Theoretic Probability

In order to state a game-theoretic version of Theorem 1.1, consider the following simplified predictably unbounded forecasting game (SPUFG, Section 5.1 of [17]) with the initial capital $\alpha > 0$.

SIMPLIFIED PREDICTABLY UNBOUNDED FORECASTING GAME **Players:** Forecaster, Skeptic, Reality

Protocol:

 $\mathcal{K}_{0} := \alpha.$ FOR n = 1, 2, ...Forecaster announces $c_{n} \ge 0.$ Skeptic announces $M_{n} \in \mathbb{R}.$ Reality announces $x_{n} \in [-c_{n}, c_{n}].$ $\mathcal{K}_{n} := \mathcal{K}_{n-1} + M_{n}x_{n}.$

Collateral Duties: Skeptic must keep \mathcal{K}_n non-negative. Reality must keep \mathcal{K}_n from tending to infinity.

Usually α is taken to be 1, but in Section 4 we use $\alpha \neq 1$ for notational simplicity.

We prove the following theorem, which implies Theorem 1.1 by Chapter 8 of [17].

Theorem 2.1. Consider SPUFG. Let ψ be a positive non-decreasing continuous function. If $I(\psi) < \infty$, Skeptic can force

$$A_n^2 \to \infty \text{ and } \limsup c_n \frac{\psi(A_n^2)^3}{A_n} < \infty \implies S_n < A_n \psi(A_n^2) \text{ a.a.}$$
 (6)

and if $I(\psi) = \infty$, Skeptic can force

$$A_n^2 \to \infty \text{ and } \limsup c_n \frac{\psi(A_n^2)^3}{A_n} < \infty \implies S_n \ge A_n \psi(A_n^2) \text{ i.o.}$$
 (7)

We use the same line of arguments as in [14] and Chapter 5 of Shafer and Vovk [17]. We employ a Bayesian mixture of constant-proportion betting strategies. Here we give basic properties of constantproportion betting strategies.

A constant-proportion betting strategy with betting proportion $\gamma > 0$ sets

$$M_n = \gamma \mathcal{K}_{n-1}.$$

However, \mathcal{K}_n becomes negative if $\gamma x_n < -1$. For simplicity we consider applying the strategy ("keep the account open") as long as $\gamma c_n \leq \delta$ and sets $M_n = 0$ once $\gamma c_n > \delta$ happens ("freeze the account"). Define a stopping time

$$\sigma_{\gamma} := \min\{n \mid \gamma c_n > \delta\}. \tag{8}$$

Note the monotonicity of σ_{γ} , i.e., $\sigma_{\gamma'} \ge \sigma_{\gamma}$ if $\gamma' \le \gamma$. We denote the capital process of the constantproportion betting strategy with this stopping time by \mathcal{K}_n^{γ} . With the initial capital of $\mathcal{K}_0^{\gamma} = \alpha$, the value of \mathcal{K}_n^{γ} is written as

$$\mathcal{K}_n^{\gamma} = \alpha \prod_{i=1}^{\min(n,\sigma_{\gamma}-1)} (1+\gamma x_i).$$

By

$$t - \frac{t^2}{2} - t^2 \times |t| \le \ln(1+t) \le t - \frac{t^2}{2} + t^2 \times |t|$$

for $|t| \leq \delta$, taking the logarithm of $\prod_{i=1}^{n} (1 + \gamma x_i)$, for $n < \sigma_{\gamma}$, we have

$$\gamma S_n - \frac{\gamma^2 A_n^2}{2} - \gamma^3 A_n^2 \bar{c}_n \le \ln\left(\mathcal{K}_n^{\gamma}/\alpha\right) \le \gamma S_n - \frac{\gamma^2 A_n^2}{2} + \gamma^3 A_n^2 \bar{c}_n$$

and

$$e^{-\gamma^{3}A_{n}^{2}\bar{c}_{n}}e^{\gamma S_{n}-\gamma^{2}A_{n}^{2}/2} \leq \mathcal{K}_{n}^{\gamma}/\alpha \leq e^{\gamma^{3}A_{n}^{2}\bar{c}_{n}}e^{\gamma S_{n}-\gamma^{2}A_{n}^{2}/2},$$
(9)

where

$$\bar{c}_n := \max_{1 \le i \le n} c_i$$

We also set up some notation for expressing the condition in (6) and (7). An infinite sequence of Forecaster's and Reality's announces $\omega = (c_1, x_1, c_2, x_2, ...)$ is called a *path* and the set of paths $\Omega = \{\omega\}$ is called the sample space. Define a subset $\Omega_{<\infty}$ of Ω as

$$\Omega_{<\infty} := \left\{ \omega \mid A_n^2 \to \infty, \limsup_n c_n \frac{\psi(A_n^2)^3}{A_n} < \infty \right\}.$$

For an arbitrary path $\omega \in \Omega_{<\infty}$ we have

$$\exists C(\omega) < \infty, \exists n_1(\omega), \forall n > n_1(\omega), c_n < C(\omega) \frac{A_n}{\psi(A_n^2)^3}, \ \psi(A_n^2) \ge 1.$$
(10)

The last inequality holds by the lower bound in (5).

3 Validity

We prove the validity in (6) of Theorem 2.1. In this section we let $\alpha = 1$. We discretize the integral in (1) as

$$\sum_{k=1}^{\infty} \frac{\psi(k)}{k} e^{-\psi(k)^2/2} < \infty.$$
(11)

Since $xe^{-x^2/2}$ is decreasing for $x \ge 1$, the function $\lambda \mapsto \frac{\psi(\lambda)}{\lambda}e^{-\psi(\lambda)^2/2}$ is decreasing for λ such that $\psi(\lambda) \ge 1$ and convergences of the integral in (1) and the sum in (11) are equivalent.

The convergence of the infinite series in (11) implies the existence of a non-decreasing sequence of positive reals a_k diverging to infinity ($a_k \uparrow \infty$), such that the series multiplied term by term by a_k is still convergent:

$$Z:=\sum_{k=1}^{\infty}a_k\frac{\psi(k)}{k}e^{-\psi(k)^2/2}<\infty.$$

This is easily seen by dividing the infinite series into blocks of sums less than or equal to $1/2^k$ and multiplying the *k*-th block by *k* (see also [13, Lemma 4.15]).

For $k \ge 1$ let

$$p_k := \frac{1}{Z} a_k \frac{\psi(k)}{k} e^{-\psi(k)^2/2}$$

and consider the capital process of a countable mixture of constant-proportion strategies

$$\mathcal{K}_n := \sum_{k=1}^{\infty} p_k \mathcal{K}_n^{\gamma_k}, \quad \text{where} \quad \gamma_k := \frac{\psi(k)}{\sqrt{k}}.$$
 (12)

Note that \mathcal{K}_n is never negative. By the upper bound in (5), as $k \to \infty$ we have

$$\gamma_k \le \frac{\psi^U(k)}{\sqrt{k}} = \sqrt{\frac{2\ln_2 k + 4\ln_3 k}{k}} \to 0.$$
 (13)

We show that $\limsup_n \mathcal{K}_n = \infty$ if a path $\omega \in \Omega_{<\infty}$ satisfies $S_n \ge A_n \psi(A_n^2)$ *i.o.* We bound $Z\mathcal{K}_n$ as

$$Z\mathcal{K}_n \ge \sum_{k=\lfloor A_n^2 - A_n^2/\psi(A_n^2)\rfloor}^{\lfloor A_n^2 \rfloor} p_k \mathcal{K}_n^{\gamma_k}.$$
(14)

At this point we check that all accounts on the right-hand side of (14) are open for sufficiently large *n* and the lower bound in (9) can be applied to each term of (14) for $\omega \in \Omega_{<\infty}$. We have the following two lemmas.

Lemma 3.1. Let $\omega \in \Omega_{<\infty}$. Let $C = C(\omega)$ in (10). For sufficiently large n

$$\bar{c}_n = \max_{1 \le i \le n} c_i < (1+\delta)C \frac{A_n}{\psi(A_n^2)^3}.$$
(15)

Proof. Note that the first $n_1(\omega)$ *c*'s i.e., $c_1, \ldots, c_{n_1(\omega)}$, do not matter since $\lim_{n\to\infty} A_n/\psi(A_n^2)^3 = \infty$. For $l > n_1(\omega)$, by (10) we have

$$c_l \le C \frac{A_l}{\psi(A_l^2)^3} \le C A_l$$

Hence c_l such that $A_l \leq A_n/\psi(A_n^2)^3$ do not matter in \bar{c}_n .

For c_l such that $A_l > A_n/\psi(A_n^2)^3$ we have

$$c_l \le C \frac{A_l}{\psi(A_n^2/\psi(A_n^2)^6)^3} \le C \frac{A_n}{\psi(A_n^2/\psi(A_n^2)^6)^3} = C \frac{A_n}{\psi(A_n^2)^3} \frac{\psi(A_n^2)^3}{\psi(A_n^2/\psi(A_n^2)^6)^3}$$

But by (5), both $\psi(A_n^2)$ and $\psi(A_n^2/\psi(A_n^2)^6)$ are of the order $\sqrt{2 \ln_2 A_n^2} (1+o(1))$ and $\psi(A_n^2)/\psi(A_n^2/\psi(A_n^2)^6) \to 1$ as $n \to \infty$. Hence (15) holds.

Lemma 3.2. Let $\omega \in \Omega_{<\infty}$. For sufficiently large n, $\sigma_{\gamma_k} > n$ for all $k = \lfloor A_n^2 - A_n^2/\psi(A_n^2) \rfloor, \dots, \lfloor A_n^2 \rfloor$.

Proof. By the monotonicity of ψ , we have $\gamma_k \leq \psi(A_n^2) / \sqrt{\lfloor A_n^2 - A_n^2 / \psi(A_n^2) \rfloor}$ for $k = \lfloor A_n^2 - A_n^2 / \psi(A_n^2) \rfloor, \dots, \lfloor A_n^2 \rfloor$. Then by the monotonicity of σ_{γ} , it suffices to show

$$\frac{\psi(A_n^2)}{\sqrt{[A_n^2 - A_n^2/\psi(A_n^2)]}} \bar{c}_n \le \delta$$

for sufficiently large n. By (15), the left-hand side is bounded from above by

$$\frac{\psi(A_n^2)}{\sqrt{[A_n^2 - A_n^2/\psi(A_n^2)]}} \times (1+\delta)C\frac{A_n}{\psi(A_n^2)^3} = (1+\delta)C\frac{A_n}{\sqrt{[A_n^2 - A_n^2/\psi(A_n^2)]}}\frac{1}{\psi(A_n^2)^2}.$$

But this converges to 0 as $n \to \infty$.

By Lemma 3.2 and the lower bound in (9), for sufficiently large *n*, we have

$$\mathcal{K}_n^{\gamma_k} \ge e^{-\gamma_k^3 A_n^2 \bar{c}_n} e^{\gamma_k S_n - \gamma_k^2 A_n^2/2}, \quad k = \lfloor A_n^2 - A_n^2/\psi(A_n^2) \rfloor, \dots, \lfloor A_n^2 \rfloor$$

and $Z\mathcal{K}_n$ can be evaluated from below as

$$Z\mathcal{K}_{n} \geq Z \sum_{k=\lfloor A_{n}^{2} - A_{n}^{2}/\psi(A_{n}^{2}) \rfloor}^{\lfloor A_{n}^{2} \rfloor} p_{k} \exp(\gamma_{k}S_{n} - \frac{\gamma_{k}^{2}A_{n}^{2}}{2} - \gamma_{k}^{3}A_{n}^{2}\bar{c}_{n})$$

$$= \sum_{k=\lfloor A_{n}^{2} - A_{n}^{2}/\psi(A_{n}^{2}) \rfloor}^{\lfloor A_{n}^{2} \rfloor} a_{k} \frac{\psi(k)}{k} \exp(-\frac{\psi(k)^{2}}{2} + \gamma_{k}S_{n} - \frac{\gamma_{k}^{2}A_{n}^{2}}{2} - \gamma_{k}^{3}A_{n}^{2}\bar{c}_{n})$$

Now we assume that $S_n \ge A_n \psi(A_n^2)$ *i.o.* for the path $\omega \in \Omega_{<\infty}$. Then for sufficiently large *n* such that $S_n \ge A_n \psi(A_n^2), \psi(A_n^2)/(\psi(A_n^2)-1) \le 1+\delta$ and $A_n/(\lfloor A_n^2 - A_n^2/\psi(A_n^2) \rfloor)^{1/2} \le 1+\delta$, we evaluate the exponent part by (9) as

$$-\frac{\psi(k)^{2}}{2} + \gamma_{k}S_{n} - \frac{\gamma_{k}^{2}A_{n}^{2}}{2} \ge -\frac{\psi(k)^{2}}{2} + A_{n}\psi(A_{n}^{2})\frac{\psi(k)}{\sqrt{k}} - \frac{\psi(k)^{2}}{k}\frac{A_{n}^{2}}{2}$$
$$= \psi(k)\left(-\frac{1}{2}\left(1 + \frac{A_{n}^{2}}{k}\right)\psi(k) + \sqrt{\frac{A_{n}^{2}}{k}}\psi(A_{n}^{2})\right)$$
$$\ge -\frac{\psi(A_{n}^{2})^{2}}{2}\left(\sqrt{\frac{A_{n}^{2}}{k}} - 1\right)^{2} \ge -\frac{\psi(A_{n}^{2})^{2}}{2}\left(\frac{A_{n}^{2}}{k} - 1\right)^{2}$$
$$\ge -\frac{1}{2}\left(\frac{\psi(A_{n}^{2})}{\psi(A_{n}^{2}) - 1}\right)^{2} \ge -\frac{1}{2} - 2\delta$$

and by Lemma 3.1

$$\begin{split} \gamma_k^3 A_n^2 \bar{c}_n &\leq \frac{\psi(A_n^2)^3}{\left(\lfloor A_n^2 - A_n^2/\psi(A_n^2) \rfloor\right)^{3/2}} A_n^2 (1+\delta) C \frac{A_n}{\psi(A_n^2)^3} \\ &\leq (1+\delta) C \left(\frac{A_n}{\left(\lfloor A_n^2 - A_n^2/\psi(A_n^2) \rfloor\right)^{1/2}}\right)^3 \\ &\leq C(1+\delta)^4. \end{split}$$
(16)

For sufficiently large n, we have

$$\psi(A_n^2) \le \psi^U(A_n^2) < \psi^U(2k) = \sqrt{2\ln_2 2k} + 4\ln_3 2k < 2\sqrt{2\ln_2 k} + 3\ln_2 k = 2\psi^L(k) \le 2\psi(k).$$

Thus by (16),

$$Z\mathcal{K}_{n} \geq \sum_{k=\lfloor A_{n}^{2} - A_{n}^{2}/\psi(A_{n}^{2}) \rfloor}^{\lfloor A_{n}^{2} \rfloor} a_{k} \frac{\psi(k)}{k} \exp\left(-\frac{1}{2} - 2\delta - C(1+\delta)^{4}\right)$$
$$\geq a_{\lfloor A_{n}^{2} - A_{n}^{2}/\psi(A_{n}^{2}) \rfloor} \frac{\psi(A_{n}^{2})}{2A_{n}^{2}} \sum_{k=\lfloor A_{n}^{2} - A_{n}^{2}/\psi(A_{n}^{2}) \rfloor}^{\lfloor A_{n}^{2} \rfloor} \exp\left(-\frac{1}{2} - 2\delta - C(1+\delta)^{4}\right)$$

$$\geq a_{\lfloor A_n^2 - A_n^2/\psi(A_n^2) \rfloor} \frac{\psi(A_n^2)}{2A_n^2} \left(\frac{A_n^2}{\psi(A_n^2)} - 1 \right) \exp\left(-\frac{1}{2} - 2\delta - C(1+\delta)^4 \right)$$
$$= a_{\lfloor A_n^2 - A_n^2/\psi(A_n^2) \rfloor} \left(\frac{1}{2} - \frac{\psi(A_n^2)}{2A_n^2} \right) \exp\left(-\frac{1}{2} - 2\delta - C(1+\delta)^4 \right).$$

Since $a_{|A_n^2 - A_n^2/\psi(A_n^2)|} \to \infty$ as $n \to \infty$, we have shown

$$\omega \in \Omega_{<\infty}, S_n \ge A_n \psi(A_n^2) \text{ i.o. } \Rightarrow \limsup_{n \to \infty} \mathcal{K}_n = \infty.$$

4 Sharpness

We prove the sharpness in (7) of Theorem 2.1. As in Section 4.2 of [17] and in [13], in order to prove the sharpness, it suffices to show the following proposition.

Proposition 4.1. Consider SPUFG. Let ψ be a positive non-decreasing continuous function. If $I(\psi) = \infty$, then for each C > 0, Skeptic can force

$$A_n^2 \to \infty$$
, $\limsup_n c_n \frac{\psi(A_n^2)^3}{A_n} \le C \implies S_n \ge A_n \psi(A_n^2)$ i.o. (17)

Once we prove this proposition, we can take the mixture over C = 1, 2, ... Then the sharpness follows, because for each $\omega \in \Omega_{<\infty}$, there exists $C(\omega)$ satisfying (10). We denote

$$\Omega_{C} := \left\{ \omega \in \Omega \mid A_{n}^{2} \to \infty, \limsup_{n} c_{n} \frac{\psi(A_{n}^{2})^{3}}{A_{n}} < (1 - \delta)C \right\},$$

$$\Omega_{0} := \left\{ \omega \in \Omega \mid \lim_{n \to \infty} A_{n}^{2} < \infty \right\},$$

$$\Omega_{=\infty} := \left\{ \omega \in \Omega \mid A_{n}^{2} \to \infty, \limsup_{n} c_{n} \frac{\psi(A_{n}^{2})^{3}}{A_{n}} = \infty \right\}.$$

We divide our proof of Proposition 4.1 into several subsections. For notational simplicity we use the initial capital of $\alpha = 1 - 2/e = (e - 2)/e$ in this section. In Sections 4.1 and 4.2 we only consider γ and n with $n < \sigma_{\gamma}$. As in Lemma 3.2 for the validity, this condition will be satisfied for sufficiently small γ and relevant n.

4.1 Uniform mixture of constant-proportion betting strategies

We consider a continuous uniform mixture of constant-proportion strategies with the betting proportion $u\gamma$, $2/e \le u \le 1$. This is a Bayesian strategy, a similar one to which has been considered in [12].

Define

$$\mathcal{L}_n^{\gamma} := \int_{2/e}^1 \prod_{i=1}^{\min(n,\sigma_{\gamma}-1)} (1+u\gamma x_i) du, \qquad \mathcal{L}_0^{\gamma} = \alpha = 1-e/2.$$

At round $n < \sigma_{\gamma}$ this strategy bets $M_n = \int_{2/e}^{1} u\gamma \prod_{i=1}^{n-1} (1 + u\gamma x_i) du$. Then by induction on $n < \sigma_{\gamma}$ the capital process is indeed written as

$$\mathcal{L}_{n}^{\gamma} = \mathcal{L}_{n-1}^{\gamma} + M_{n}x_{n} = \int_{2/e}^{1} \prod_{i=1}^{n-1} (1 + u\gamma x_{i}) du + x_{n} \int_{2/e}^{1} u\gamma \prod_{i=1}^{n-1} (1 + u\gamma x_{i}) du$$

$$=\int_{2/e}^{1}\prod_{i=1}^{n}(1+u\gamma x_{i})du.$$

Applying (9), we have

$$e^{-\gamma^{3}A_{n}^{2}\bar{c}_{n}}\int_{2/e}^{1}e^{u\gamma S_{n}-u^{2}\gamma^{2}A_{n}^{2}/2}du\leq \mathcal{L}_{n}^{\gamma}\leq e^{\gamma^{3}A_{n}^{2}\bar{c}_{n}}\int_{2/e}^{1}e^{u\gamma S_{n}-u^{2}\gamma^{2}A_{n}^{2}/2}du,$$

for $n < \sigma_{\gamma}$. We further bound the integral in the following lemma.

Lemma 4.2. For $n < \sigma_{\gamma}$,

$$\begin{pmatrix} e^{\gamma^3 A_n^2 \bar{c}_n} e^{2\gamma (S_n/e - \gamma A_n^2/e^2)} & \text{if } S_n \le 2\gamma A_n^2/e, \\
\end{cases}$$
(18)

$$\mathcal{L}_{n}^{\gamma} \leq \left\{ e^{\gamma^{3}A_{n}^{2}\tilde{c}_{n}} \min\left\{ e^{S_{n}^{2}/(2A_{n}^{2})} \frac{\sqrt{2\pi}}{\gamma A_{n}}, e^{\gamma S_{n}/2} \right\} \qquad if \quad 2\gamma A_{n}^{2}/e < S_{n} < \gamma A_{n}^{2}, \tag{19}$$

$$\left\{ e^{\gamma^3 A_n^2 \bar{c}_n} \min\left\{ e^{S_n^2/(2A_n^2)} \frac{\sqrt{2\pi}}{\gamma A_n}, e^{\gamma S_n - \gamma^2 A_n^2/2} \right\} \quad if \quad S_n \ge \gamma A_n^2.$$

$$(20)$$

Proof. Completing the square we have

$$-\frac{1}{2}u^{2}\gamma^{2}A_{n}^{2}+u\gamma S_{n}=-\frac{\gamma^{2}A_{n}^{2}}{2}\left(u-\frac{S_{n}}{\gamma A_{n}^{2}}\right)^{2}+\frac{S_{n}^{2}}{2A_{n}^{2}}$$

Hence by the change of variables

$$v = \gamma A_n \left(u - \frac{S_n}{\gamma A_n^2} \right), \qquad du = \frac{dv}{\gamma A_n},$$

we obtain

$$\int_{2/e}^{1} e^{u\gamma S_n - u^2 \gamma^2 A_n^2/2} du = e^{S_n^2/(2A_n^2)} \int_{2/e}^{1} \exp\left(-\frac{\gamma^2 A_n^2}{2} \left(u - \frac{S_n}{\gamma A_n^2}\right)^2\right) du$$
$$= e^{S_n^2/(2A_n^2)} \frac{1}{\gamma A_n} \int_{2\gamma A_n/e - S_n/A_n}^{\gamma A_n - S_n/A_n} e^{-v^2/2} dv.$$

Then for all cases we can bound \mathcal{L}_n^{γ} from above as

$$\mathcal{L}_n^{\gamma} \le e^{\gamma^3 A_n^2 \bar{c}_n + S_n^2 / (2A_n^2)} \frac{\sqrt{2\pi}}{\gamma A_n}.$$
(21)

Without change of variables, we can also bound the integral $\int_{2/e}^{1} g(u) du$, $g(u) := e^{u\gamma S_n - u^2\gamma^2 A_n^2/2}$, directly as

$$\int_{2/e}^{1} g(u)du \leq \max_{2/e \leq u \leq 1} g(u).$$

Note that

$$g(2/e) = e^{2\gamma(S_n/e - \gamma A_n/e^2)}, \quad g(1) = e^{\gamma S_n - \gamma^2 A_n^2/2}.$$
 (22)

We now consider the following three cases.

- **Case 1** $S_n \le 2\gamma A_n^2/e$. In this case $S_n/(\gamma A_n^2) \le 2/e$ and by the unimodality of g(u) we have $\max_{2/e \le u \le 1} g(u) = g(2/e)$. Hence (18) follows from (22).
- **Case 2** $2\gamma A_n^2/e < S_n < \gamma A_n^2$. In this case $\max_{2/e \le u \le 1} g(u) = g(S_n/(\gamma A_n^2)) = e^{S_n^2/(2A_n^2)}$ and $\mathcal{L}_n^{\gamma} \le e^{\gamma^3 A_n^2 \bar{c}_n} e^{S_n^2/(2A_n^2)}$. Furthermore in this case $S_n^2 < \gamma A_n^2 S_n$ implies $S_n^2/(2A_n^2) < \gamma S_n/2$ and we also have

$$\mathcal{L}_n^{\gamma} \le e^{\gamma^3 A_n^2 \bar{c}_n} e^{\gamma S_n/2}.$$
(23)

By (21) and (23), we have (19).

Case 3 $S_n \ge \gamma A_n^2$. Then $S_n/(\gamma A_n^2) \ge 1$ and $\max_{2/e \le u \le 1} g(u) = g(1)$. Hence

$$\mathcal{L}_{n}^{\gamma} \leq e^{\gamma^{3} A_{n}^{2} \bar{c}_{n}} e^{\gamma S_{n} - \gamma^{2} A_{n}^{2}/2}.$$
(24)

By (21) and (24), we have (20).

4.2 Buying a process and selling a process

Next we consider the following capital process.

$$\mathcal{Q}_n^{\gamma} := 2\mathcal{L}_n^{\gamma} - \mathcal{K}_n^{\gamma e}. \tag{25}$$

This capital process consists of buying two units of \mathcal{L}_n^{γ} and selling one unit of $\mathcal{K}_n^{\gamma e}$. This combination of selling and buying is essential in the game-theoretic proof of LIL in Chapter 5 of [17] and [14]. However, unlike Chapter 5 of [17] and [14], where a combination of *three* capital processes is used, we only combine *two* capital processes.

We want to bound Q_n^{γ} from above.

|C|

Lemma 4.3. Let

$$C_1 := 2e^{\gamma^3 A_n^2 \bar{c}_n} \exp\left(\frac{(2e-1)((1+e^3)\gamma^3 A_n^2 \bar{c}_n + \ln 2)}{(e-1)^2}\right).$$
(26)

Then for $n < \sigma_{\gamma e}$,

$$\begin{pmatrix} C_1 & \text{if } S_n \le \gamma A_n^2/e, \\ (\sqrt{2}) \end{pmatrix}$$
 (27)

$$Q_n^{\gamma} \leq \left\{ 2e^{\gamma^3 A_n^2 \bar{c}_n} \min\left\{ e^{S_n^2/(2A_n^2)} \frac{\sqrt{2\pi}}{\gamma A_n}, e^{\gamma S_n} \right\} \quad if \quad \gamma A_n^2/e < S_n < e\gamma A_n^2, \tag{28}$$

$$if \quad S_n \ge e\gamma A_n^2. \tag{29}$$

Remark 4.4. In this lemma, C_1 depends on \bar{c}_n , γ and A_n through $\gamma^3 A_n^2 \bar{c}_n$. However from Section 4.5 on, we evaluate $\gamma^3 A_n^2 \bar{c}_n$ from above by a constant. Hence, C_1 can be also taken to be a constant (cf. (50)) not depending on γ and A_n . Also note that the interval for S_n in (28) is larger than the interval in (19).

Proof. We bound $Q_n^{\gamma} = 2\mathcal{L}_n^{\gamma} - \mathcal{K}_n^{\gamma e}$ from above in the following three cases:

(i)
$$S_n \le \gamma A_n^2 / e$$
, (ii) $\gamma A_n^2 / e < S_n < e \gamma A_n^2$, (iii) $S_n \ge e \gamma A_n^2$,

Case (i) In this case $S_n/e - \gamma A_n^2/e^2 \le 0$. Hence (27) follows from (18) and $Q_n^{\gamma} \le 2\mathcal{L}_n^{\gamma}$.

Case (ii) We again use $Q_n^{\gamma} \le 2\mathcal{L}_n^{\gamma}$. If $\gamma A_n^2/e < S_n \le 2\gamma A_n^2/e$, then

$$\frac{S_n}{e} - \frac{\gamma A_n^2}{e^2} \le \frac{\gamma A_n^2}{e^2} \le \frac{S_n}{e}$$

and $\mathcal{L}_{n}^{\gamma} \leq e^{\gamma^{3}A_{n}^{2}\bar{c}_{n}}e^{2\gamma S_{n}/e} \leq e^{\gamma^{3}A_{n}^{2}\bar{c}_{n}}e^{\gamma S_{n}}$ from (18). Otherwise (28) follows from (19) and (20). **Case (iii)** Since $S_{n} \geq eA_{n}^{2}\gamma > A_{n}^{2}\gamma$, by (24) we have $\mathcal{L}_{n}^{\gamma} \leq e^{\gamma^{3}A_{n}^{2}\bar{c}_{n}}e^{\gamma S_{n}-\gamma^{2}A_{n}^{2}/2}$ and

$$\begin{aligned} Q_n^{\gamma} &\leq 2\mathcal{L}_n^{\gamma} - \mathcal{K}_n^{\gamma e} \leq 2e^{\gamma^3 A_n^2 \bar{c}_n} e^{\gamma S_n - \gamma^2 A_n^2/2} - e^{-\gamma^3 e^3 A_n^2 \bar{c}_n} e^{\gamma e S_n - \gamma^2 e^2 A_n^2/2} \\ &= 2e^{\gamma^3 A_n^2 \bar{c}_n} e^{\gamma S_n - \gamma^2 A_n^2/2} \left(1 - \frac{1}{2} e^{-(1+e^3)\gamma^3 A_n^2 \bar{c}_n} e^{\gamma(e-1)S_n - (e^2-1)\gamma^2 A_n^2/2} \right) \end{aligned}$$

Hence if the right-hand side is non-positive we have $Q_n^{\gamma} \leq 0$:

$$S_n \ge eA_n^2 \gamma \text{ and } -(1+e^3)\gamma^3 A_n^2 \bar{c}_n - \ln 2 + \gamma(e-1)S_n - \frac{1}{2}(e^2-1)\gamma^2 A_n^2 \ge 0$$

 $\Rightarrow Q_n^{\gamma} \le 0.$ (30)

Otherwise, write $B_n := (1 + e^3)\gamma^3 A_n^2 \bar{c}_n + \ln 2$ and consider the case

$$\gamma(e-1)S_n - \frac{1}{2}(e^2 - 1)\gamma^2 A_n^2 \le B_n$$

Dividing this by e - 1 and also considering $S_n \ge eA_n^2 \gamma$, we have

$$\gamma S_n - \frac{1}{2}(e+1)\gamma^2 A_n^2 \le \frac{B_n}{e-1},$$
(31)

$$-S_n + eA_n^2 \gamma \le 0. \tag{32}$$

 $\gamma \times (32) + (31)$ gives

$$\frac{1}{2}(e-1)\gamma^2 A_n^2 \le \frac{B_n}{e-1} \quad \text{or} \quad \frac{1}{2}\gamma^2 A_n^2 \le \frac{B_n}{(e-1)^2}.$$

Then by (31)

$$\gamma S_n - \frac{1}{2} \gamma^2 A_n^2 \le \frac{B_n}{e-1} + \frac{e}{2} \gamma^2 A_n^2 \le \frac{B_n}{e-1} + \frac{eB_n}{(e-1)^2} = \frac{(2e-1)B_n}{(e-1)^2}.$$

Hence just using $Q_n^{\gamma} \leq 2\mathcal{L}_n^{\gamma}$ and (24) in this case, we obtain

$$Q_n^{\gamma} \le 2e^{\gamma^3 A_n^2 \bar{c}_n} \exp\left(\frac{(2e-1)((1+e^3)\gamma^3 A_n^2 \bar{c}_n + \ln 2)}{(e-1)^2}\right) = C_1.$$
(33)

This also covers (30) and we have (33) for the whole case (iii).

_	-	-	

4.3 Change of time scale and dividing the rounds into cycles

For proving the sharpness we consider the change of time scale from λ to k:

$$\lambda = e^{5k\ln k} = k^{5k}.$$

By taking the derivative of $\ln \lambda = 5k \ln k$, we have $d\lambda/\lambda = 5(\ln k+1)dk$. Since $\ln k$ is dominant in $(\ln k+1)$, the integrability condition is written as

$$\int_{1}^{\infty} \psi(\lambda) e^{-\psi(\lambda)^{2}/2} \frac{d\lambda}{\lambda} = \infty \iff \int_{1}^{\infty} (\ln k) \psi(e^{5k \ln k}) e^{-\psi(e^{5k \ln k})^{2}/2} dk = \infty$$

Let $f(x) := \psi(e^{5x \ln x})e^{-\psi(e^{5x \ln x})^2/2}$. Since $xe^{-x^2/2}$ is decreasing for $x \ge 1$, the function f(x) is decreasing for x such that $\psi(e^{5x \ln x}) \ge 1$. Thus, for sufficiently large k and x such that $k \le x \le k + 1$, we have

$$\frac{1}{2}\ln(k+1)f(k+1) \le \ln kf(x+1) \le \ln xf(x) \le \ln(k+1)f(x) \le 2\ln kf(k).$$

Hence, we have

$$\int_{1}^{\infty} (\ln k) \psi(e^{5k \ln k}) e^{-\psi(e^{5k \ln k})^2/2} dk = \infty \quad \Leftrightarrow \quad \sum_{k=1}^{\infty} (\ln k) \psi(e^{5k \ln k}) e^{-\psi(e^{5k \ln k})^2/2} = \infty$$

Then, it suffices to show (17) if $\sum_{k=1}^{\infty} (\ln k) \psi(e^{5k \ln k}) e^{-\psi(e^{5k \ln k})^2/2} = \infty$.

As in Chapter 5 of [17] and [14], we divide the time axis into "cycles". However, unlike in Chapter 5 of [17] and [14], our cycles are based on stopping times. Let

$$n_k := k^{5k}, \quad k = 1, 2, \dots,$$
 (34)

and define a family of stopping times

$$\tau_k := \min\left\{n \mid A_n^2 \ge n_k\right\}. \tag{35}$$

We define the *k*-th cycle by $[\tau_k, \tau_{k+1}], k \ge 1$. Note that τ_k is finite for all *k* if and only if $A_n^2 \to \infty$. Betting strategy for the *k*-th cycle is based on the following betting proportion:

$$\gamma_k := \frac{\psi(n_{k+1})}{\sqrt{n_{k+1}}} k^2.$$
(36)

Note that γ_k in (36) is slightly different from (12).

For the rest of this section, we check the growth of various quantities along the cycles. Let $\omega \in \Omega_C$. For sufficiently large *n*,

$$|x_n| \le c_n \le C \frac{A_n}{\psi(A_n^2)^3}.$$
(37)

Furthermore $A_n^2 = A_{n-1}^2 + x_n^2$. This allows us to bound x_n^2 and A_n^2 in terms of A_{n-1}^2 . By squaring (37) we have

$$x_n^2 \le C^2 \frac{A_{n-1}^2}{\psi(A_n^2)^6 - C^2}$$
(38)

and

$$A_n^2 = A_{n-1}^2 + x_n^2 \le A_{n-1}^2 (1 + \frac{C^2}{\psi(A_n^2)^6 - C^2}) = A_{n-1}^2 \frac{\psi(A_n^2)^6}{\psi(A_n^2)^6 - C^2}.$$
(39)

Since $\psi(A_n^2)^6/(\psi(A_n^2)^6 - C^2) \to 1$ as $n \to \infty$, we have

$$\lim_{n \to \infty} \frac{A_n^2}{A_{n-1}^2} = 1$$

Note that $A_{\tau_k-1}^2 < n_k \le A_{\tau_k}^2$ by the definition of τ_k . Hence for $\omega \in \Omega_C$ we also have

$$\lim_{k \to \infty} \frac{A_{\tau_k}^2}{n_k} = 1.$$
(40)

The limits in the following lemma will be useful for our argument.

Lemma 4.5. For $\omega \in \Omega_C$

$$\lim_{k \to \infty} \frac{\psi^U(n_k)}{\psi(n_{k+1})} = 1, \quad \lim_{k \to \infty} \frac{k^5 A_{\tau_k}^2}{n_{k+1}} = e^{-5}, \quad \lim_{k \to \infty} \gamma_k A_{\tau_k} \psi(n_{k+1}) = 0.$$
(41)

Proof. All of $\psi^U(n_k)$, $\psi^U(n_{k+1})$, $\psi^L(n_k)$, $\psi^L(n_{k+1})$, $\psi(n_{k+1})$, $\psi(n_{k+1}/k^4)$ are of the order

$$\sqrt{2\ln\ln e^{5k\ln k}}(1+o(1)) = \sqrt{2\ln k}(1+o(1))$$
(42)

as $k \to \infty$ and the first equality holds by (5). The second equality holds by (40) and

$$\lim_{k \to \infty} \frac{k^5 n_k}{n_{k+1}} = \lim_{k \to \infty} \frac{k^{5(k+1)}}{(k+1)^{5(k+1)}} = \lim_{k \to \infty} \left(1 - \frac{1}{k+1}\right)^{5(k+1)} = e^{-5}.$$

Then $A_{\tau_k}^2/n_{k+1} = (1 + o(1))n_k/n_{k+1} = O(k^{-5})$ and the third equality holds by

$$\gamma_k A_{\tau_k} \psi(n_{k+1}) \le \psi(n_{k+1})^2 k^2 ((1+\delta)n_k/n_{k+1})^{1/2} \to 0 \qquad (k \to \infty).$$

4.4 Stopping times for aborting and sequential freezing for each cycle

In (48) of the next section we will introduce another capital process $\mathcal{M}_n^{\gamma_k,k}$, which will be employed in each cycle. Here we introduce some stopping times for aborting the cycle and for sequential freezing of accounts in $\mathcal{M}_n^{\gamma_k,k}$.

We say that we *abort* the *k*-th cycle, when we freeze all accounts in the *k*-th cycle and wait for the (k + 1)-st cycle. There are two cases for aborting the *k*-th cycle. The first case is when some c_n is too large for $\omega \in \Omega_C$. Define

$$\sigma_{k,C} := \min\left\{n \ge \tau_k \mid c_n \psi(A_{\tau_k}^2)^3 > (1+\delta)CA_{n-1}\right\}.$$
(43)

We will abort the *k*-th cycle if $\sigma_{k,C} < \tau_{k+1}$. Note that for $\omega \in \Omega_C$, there exists $k_1(\omega)$ such that

$$\sigma_{k,C} = \infty, \text{ for } k \ge k_1(\omega).$$
 (44)

Another case is when S_n is too large. Define

$$v_k := \min\{n \ge \tau_k \mid A_n \psi(A_n^2) < S_n\}.$$
(45)

If $v_k < \tau_{k+1}$, then Skeptic is happy to abort the *k*-th cycle, because he wants to force $S_n \ge A_n \psi(A_n^2)$ *i.o.* The above two stopping times will be used in the final construction of a dynamic strategy in Section 4.6. For each cycle, we define another family of stopping times indexed by $w = 1, ..., \lceil \ln k \rceil$, by

$$\tau_{k,w} := \min\left\{n \mid A_n^2 \ge e^{2(w+2)} \frac{n_{k+1}}{k^4}\right\}.$$
(46)

for sequential freezing of accounts of $\mathcal{M}_n^{\gamma_k,k}$ in (48). We have $\tau_k \leq \tau_{k,w}$ for $k \geq 1$ and $w \geq 1$, because

$$\frac{n_{k+1}}{k^4} = \frac{(k+1)^{5(k+1)}}{k^4} > k^{5k} = n_k$$

Lemma 4.6. Let $\omega \in \Omega_C$. $\tau_{k, \lceil \ln k \rceil} \leq \tau_{k+1}$ for sufficiently large k.

Proof. By $A_{\tau_{k,w}-1}^2 \le e^{2(w+2)} n_{k+1}/k^4$ and by (38), for sufficiently large k we have

$$x_{\tau_{k,w}}^2 \le (1+\delta)C^2 \frac{A_{\tau_{k,w}-1}^2}{\psi(A_{\tau_k}^2)^6} \le \frac{(1+\delta)C^2}{\psi(A_{\tau_k}^2)^6} \times \frac{e^{2(w+2)}n_{k+1}}{k^4}$$

and

$$A_{\tau_{k,w}}^2 \le A_{\tau_{k,w}-1}^2 + x_{\tau_{k,w}}^2 \le (1+\delta)e^{2(w+2)}\frac{n_{k+1}}{k^4}.$$
(47)

Then

$$A_{\tau_{k,\lceil \ln k\rceil}}^2 \le (1+\delta) \left(e^{2(\ln k+2)} \frac{n_{k+1}}{k^4} \right) = (1+\delta) e^4 \frac{n_{k+1}}{k^2} \le n_{k+1} \le A_{\tau_{k+1}}^2.$$

We also compare $\tau_{k,w}$ to $\sigma_{\gamma_k e^{-w+1}}$ defined in (8). This is needed for applying the bounds derived in previous sections to $\mathcal{M}_n^{\gamma_k,k}$ in the next section.

Lemma 4.7. Let $\omega \in \Omega_C$. $\tau_{k,w} \leq \sigma_{\gamma_k e^{-w+1}}$ for sufficiently large k.

Proof. By (47) and by Lemma 3.1, for sufficiently large k

$$\gamma_k e^{-w+1} \bar{c}_{\tau_{k,w}} \le \frac{\psi(n_{k+1})}{\sqrt{n_{k+1}}} k^2 e^{-w+1} \times (1+\delta)^2 C \frac{e^{w+2} \sqrt{n_{k+1}}}{k^2 \psi(A_{\tau_k}^2)^3} \le (1+\delta)^2 C e^3 \frac{\psi(n_{k+1})}{\psi(A_{\tau_k}^2)^3} \le \delta,$$

because $\psi(n_{k+1})/\psi(A_{\tau_k}^2)^3 \to 0$ as $k \to \infty$ by (42).

4.5 Further discrete mixture of processes for each cycle with sequential freezing

We introduce another discrete mixture of capital process for the k-th cycle. Define

$$\mathcal{M}_{n}^{\gamma_{k},k} := \frac{1}{\lceil \ln k \rceil} \sum_{w=1}^{\lceil \ln k \rceil} \mathcal{Q}_{\min(n,\tau_{k,w})}^{\gamma_{k}e^{-w}} = \frac{1}{\lceil \ln k \rceil} \sum_{w=1}^{\lceil \ln k \rceil} (2\mathcal{L}_{\min(n,\tau_{k,w})}^{\gamma_{k}e^{-w}} - \mathcal{K}_{\min(n,\tau_{k,w})}^{\gamma_{k}e^{-w+1}}).$$
(48)

Note that the *w*-th account in the sum of $\mathcal{M}_n^{\gamma_k,k}$ is frozen at the stopping time $\tau_{k,w}$. This is needed since the bound for c_n is growing even during the *k*-th cycle.

In order to bound $\mathcal{M}_n^{\gamma_k,k}$, we first bound C_1 in (26) for each *w* in the sum of (48) by a constant independent of *n*. Note that we only need to consider $n \le \tau_{k,w}$ for the *w*-th account.

Lemma 4.8. Let $\omega \in \Omega_C$. $(\gamma_k e^{-w})^3 A_n^2 \bar{c}_n$ and hence C_1 are bounded from above by

$$(\gamma_k e^{-w})^3 A_n^2 \bar{c}_n \le (1+\delta)^5 C e^6, \tag{49}$$

$$C_1 \le 2e^{(1+\delta)^5 Ce^6} \exp\left(\frac{(2e-1)((1+\delta)^5 Ce^6(1+e^3) + \ln 2)}{(e-1)^2}\right) =: \bar{C}_1,$$
(50)

for sufficiently large k.

Proof. By (42), for sufficiently large k

$$\frac{\psi(n_{k+1})}{\psi(A_{\tau_{k,\nu}}^2)} \le \frac{\psi(n_{k+1})}{\psi(n_k)} \le 1 + \delta.$$

$$(51)$$

Thus

$$\begin{split} \gamma_{k}^{3} e^{-3w} A_{\min(n,\tau_{k,w})}^{2} \bar{c}_{\min(n,\tau_{k,w})} &\leq \gamma_{k}^{3} e^{-3w} \times A_{\tau_{k,w}}^{2} \times \bar{c}_{\min(n,\tau_{k,w})} \\ &\leq \frac{\psi(n_{k+1})^{3}}{n_{k+1}^{3/2}} k^{6} e^{-3w} \times A_{\tau_{k,w}}^{2} \times (1+\delta) C \frac{A_{\tau_{k,w}}}{\psi(A_{\tau_{k}}^{2})^{3}} \\ &\leq (1+\delta) C \frac{\psi(n_{k+1})^{3}}{\psi(A_{\tau_{k}}^{2})^{3}} k^{6} e^{-3w} \frac{A_{\tau_{k,w}}^{3}}{n_{k+1}^{3/2}} \leq (1+\delta)^{5} C e^{6}. \end{split}$$

Lemma 4.9. Let $\omega \in \Omega_C$. For sufficiently large k,

$$\mathcal{M}_{n}^{\gamma_{k},k} \leq \bar{C}_{1} + \frac{2}{\lceil \ln k \rceil} e^{(1+\delta)^{5}Ce^{\delta}} \max_{\gamma \in [\gamma_{k}/k,\gamma_{k}]} \left(\min\{e^{S_{n}^{2}/(2n)} \frac{\sqrt{2\pi}}{\gamma A_{n}}, e^{\gamma S_{n}}\} \right), \quad n \in [\tau_{k}, \tau_{k+1}],$$
(52)

where \bar{C}_1 is given by the right-hand side of (50).

Proof. We have $|\gamma_k e^{-w} \bar{c}_{\min(n,\tau_{k,w})}| \le |\gamma_k e^{-w+1} \bar{c}_{\min(n,\tau_{k,w})}| \le \delta$ by Lemma 4.7. Then we can complete the proof of (52) by Lemma 4.3 and Lemma 4.7 because the length of the interval

$$\left\{w \mid \frac{S_n}{ne} < \gamma e^{-w} < \frac{S_n e}{n}\right\}$$

is equal to 2.

As in Chapter 5 of Shafer and Vovk [17], we use $\mathcal{M}_n^{\gamma_k,k}$ in the following form.

$$\mathcal{N}_{n}^{\gamma_{k},D} := \alpha + \frac{1}{D} \lceil \ln k \rceil \psi(n_{k+1}) e^{-\psi(n_{k+1})^{2}/2} (\alpha - \mathcal{M}_{n-\tau_{k}}^{\gamma_{k},k}), \quad \alpha = 1 - \frac{2}{e}, \ D = \frac{24\sqrt{2\pi}e^{(1+\delta)^{5}e^{6}C} + 4\bar{C}_{1}}{\alpha}.$$
 (53)

Here we give a specific value of *D* for definiteness, but from the proof below it will be clear that any sufficiently large *D* can be used. Since the strategy for $\mathcal{M}_{n-\tau_k}^{\gamma_k,k}$ is applied only to x_n 's in the cycle, $\alpha = \mathcal{N}_{\tau_k}^{\gamma_k,D} = \mathcal{M}_0^{\gamma_k}$. Concerning $\mathcal{N}_n^{\gamma_k,D}$ we prove the following two propositions.

Proposition 4.10. Let $\omega \in \Omega_C$. Suppose that

$$-A_n \psi^U(A_n^2) \le S_n \le A_n \psi(A_n^2), \qquad \forall n \in [\tau_k, \tau_{k+1}].$$
(54)

and $\tau_{k+1} < \sigma_{k,C}$. Then for sufficiently large k

$$\mathcal{N}_{n}^{\gamma_{k},D} \geq \frac{\alpha}{2}, \qquad \forall n \in [\tau_{k}, \tau_{k+1}],$$
(55)

and

$$\mathcal{N}_{\tau_{k+1}}^{\gamma_{k},D} \ge \alpha \left(1 + \frac{1-\delta}{D} \lceil \ln k \rceil \psi(n_{k+1}) e^{-\psi(n_{k+1})^{2}/2} \right).$$
(56)

Proof. In our proof we denote $t = n - \tau_k$, $S_t = S_n - S_{\tau_k}$ and $A_t^2 = A_n^2 - A_{\tau_k}^2$ for $n > \tau_k$. For proving (55), we use (52) for S_t . We bound $\mathcal{M}_t^{\gamma_k,k}$ from above. By the term $\frac{2}{\lceil \ln k \rceil}$ on the right-hand side of (52), it suffices to show

$$S_{t} \leq A_{\tau_{k}}\psi^{U}(A_{\tau_{k}}^{2}) + \sqrt{A_{\tau_{k}}^{2} + A_{t}^{2}\psi(A_{\tau_{k}}^{2} + A_{t}^{2})}$$

$$\Rightarrow \psi(n_{k+1})e^{-\psi(n_{k+1})^{2}/2}2e^{(1+\delta)^{5}e^{\delta}C}\min\{e^{S_{t}^{2}/(2A_{t}^{2})}\frac{\sqrt{2\pi}}{\gamma A_{t}}, e^{\gamma S_{t}}\} \leq \frac{D\alpha}{4}, \forall \gamma \in [\gamma_{k}/k, \gamma_{k}], \forall t \in [0, \tau_{k+1} - \tau_{k}]$$

for sufficient large k. Let

$$c_1 = \frac{9}{(1+2\delta)^2}$$
 s.t. $\frac{1}{2} - \frac{1}{\sqrt{c_1}} - \delta > 0.$ (57)

We distinguish two cases:

(a)
$$A_t^2 \leq \frac{\psi(n_{k+1})^2}{c_1 \gamma^2}$$
, (b) $\frac{\psi(n_{k+1})^2}{c_1 \gamma^2} < A_t^2 \leq A_{\tau_{k+1}}^2 - A_{\tau_k}^2$.

For case (a), $A_{\tau_k}\psi^U(A_{\tau_k}^2) \le (1 + \delta)A_{\tau_k}\psi(n_{k+1})$ by the first equality in Lemma 4.5 for sufficiently large k. Also $\psi(A_{\tau_k}^2 + A_t^2) \le \psi(n_{k+1})$. Hence in this case

$$\gamma S_t \leq \left((1+\delta)\gamma A_{\tau_k} + \sqrt{\gamma^2 A_{\tau_k}^2 + \psi(n_{k+1})^2/c_1} \right) \psi(n_{k+1}).$$

Then for $\gamma \leq \gamma_k$ by the third equality in Lemma 4.5

$$\gamma S_t \le \left((1+\delta)\gamma_k A_{\tau_k} + \sqrt{\gamma_k^2 A_{\tau_k}^2 + \psi(n_{k+1})^2 / c_1} \right) \psi(n_{k+1}) = \psi(n_{k+1})^2 \left(\frac{1}{\sqrt{c_1}} + \delta \right)$$
(58)

for sufficiently large k. Since

$$\psi(n_{k+1})e^{-\psi(n_{k+1})^2/2}2e^{(1+\delta)^5}e^{\delta C}e^{\gamma S_t} \le \psi(n_{k+1})\exp\left(-\psi(n_{k+1})^2(\frac{1}{2}-\frac{1}{\sqrt{c_1}}-\delta)\right)2e^{(1+\delta)^5}e^{\delta C} \to 0 \quad (k\to\infty),$$

we have $\mathcal{N}_n^{\gamma_k,D} \ge \alpha/2$ uniformly in $\gamma \in [\gamma_k/k, \gamma_k]$.

For case (b), $\psi(n_{k+1})/\sqrt{c_1} < \gamma A_t$ and $S_t \le \left((1+\delta)A_{\tau_k} + \sqrt{A_{\tau_k}^2 + A_t^2}\right)\psi(n_{k+1})$. Hence

$$\psi(n_{k+1})e^{-\psi(n_{k+1})^2/2} \times 2e^{(1+\delta)^5}e^{6C}e^{S_t^2/(2A_t^2)}\frac{\sqrt{2\pi}}{\gamma A_t}$$

$$\leq \psi(n_{k+1})e^{-\psi(n_{k+1})^{2}/2} \times \frac{2e^{(1+\delta)^{5}e^{\delta}C}\sqrt{2\pi}\sqrt{c_{1}}}{\psi(n_{k+1})} \exp\left(\frac{\left((1+\delta)A_{\tau_{k}} + \sqrt{A_{\tau_{k}}^{2} + A_{t}^{2}}\right)^{2}}{2A_{t}^{2}}\psi(n_{k+1})^{2}\right)$$
$$= 2e^{(1+\delta)^{5}e^{\delta}C}\sqrt{2\pi}\sqrt{c_{1}}\exp\left(\frac{(1+(1+\delta)^{2})A_{\tau_{k}}^{2} + 2(1+\delta)A_{\tau_{k}}\sqrt{A_{\tau_{k}}^{2} + A_{t}^{2}}}{2A_{t}^{2}}\psi(n_{k+1})^{2}\right).$$
(59)

For $\gamma \leq \gamma_k$,

$$\frac{\psi(n_{k+1})^2}{c_1\gamma^2} < A_t^2 \implies \frac{A_{\tau_k}^2}{A_t^2}\psi(n_{k+1})^2 < c_1\gamma^2 A_{\tau_k}^2 \le c_1\gamma_k^2 A_{\tau_k}^2 = c_1\frac{A_{\tau_k}^2}{n_{k+1}}k^4\psi(n_{k+1})^2 = O(k^{-1}\ln k)$$

Hence $\psi(n_{k+1})^2 A_{\tau_k}^2 / A_t^2 \to 0$ as $k \to \infty$. Similarly $\psi(n_{k+1})^2 A_{\tau_k} / A_t \to 0$ as $k \to \infty$, because $\psi(n_{k+1})^2 A_{\tau_k} / A_t = O(k^{-1/2} (\ln k)^{3/2})$. Therefore the right-hand side of (59) is bounded from above by $2e^{(1+\delta)^5}e^{\delta C} \sqrt{2\pi} \sqrt{c_1}(1+\delta)$ for sufficiently large k and

$$\psi(n_{k+1})e^{-\psi(n_{k+1})^2/2} \times 2e^{(1+\delta)^5 e^6 C} e^{S_t^2/(2A_t^2)} \frac{\sqrt{2\pi}}{\gamma A_t} \le \frac{D\alpha}{4},$$

with the choice of *D* in (53) and c_1 in (57). This proves (55).

Now we prove (56). We focus on the *w*-th account when $n \ge \tau_{k,w}$. Recall that in this proof we have been denoting $A_t^2 = A_n^2 - A_{\tau_k}^2$. Similarly we denote $A_{\tau_{k,w}}^2$ instead of $A_{\tau_{k,w}}^2 - A_{\tau_k}^2$. Thus

$$e^{2(w+2)}\frac{n_{k+1}}{k^4} - A_{\tau_k}^2 \le A_{\tau_{k,w}}^2.$$
(60)

We will show that $\limsup_{k\to\infty} \mathcal{M}_{\tau_{k+1}-\tau_k}^{\gamma_k,k} \leq 0$, if

$$S_{\tau_{k,w}} \le A_{\tau_k} \psi(A_{\tau_k}^2) + A_{\tau_{k,w}} \psi(A_{\tau_{k,w}}^2) \le \psi(n_{k+1}) \left\{ A_{\tau_k} + A_{\tau_{k,w}} \right\} \le 2\psi(n_{k+1}) A_{\tau_{k,w}}.$$
(61)

We evaluate

$$\mathcal{L}_{\tau_{k,w}}^{\gamma_k e^{-w},k} := \int_{2/e}^{1} \exp\left(u\gamma_k e^{-w}S_{\tau_{k,w}} - u^2\gamma_k^2 e^{-2w}A_{\tau_{k,w}}^2/2\right) du$$

from above. Because $u\gamma_k e^{-w}S_{\tau_{k,w}} - u^2\gamma_k^2 e^{-2w}A_{\tau_{k,w}}^2/2$ is maximized at $u = S_{\tau_{k,w}}/(\gamma_k e^{-w}A_{\tau_{k,w}}^2)$ and

$$\frac{S_{\tau_{k,w}}}{\gamma_k e^{-w} A_{\tau_{k,w}}^2} \le \frac{2\psi(n_{k+1})A_{\tau_{k,w}}}{(\psi(n_{k+1})k^2/\sqrt{n_{k+1}})e^{-w} A_{\tau_{k,w}}^2} \le \frac{2\sqrt{n_{k+1}}}{k^2 e^{-w} A_{\tau_{k,w}}} \le \frac{2}{e^2} \le \frac{2}{e},$$

the integrand in $\mathcal{L}_{\tau_{k,w}}^{\gamma_k e^{-w},k}$ is maximized at 2/e and we have

$$\mathcal{L}_{\tau_{k,w}}^{\gamma_{k}e^{-w},k} \leq \exp\left(\frac{2}{e}\gamma_{k}e^{-w}S_{\tau_{k,w}} - \frac{2\gamma_{k}^{2}e^{-2w}A_{\tau_{k,w}}^{2}}{e^{2}}\right)$$

By (60) and (61), for sufficiently large k,

$$\frac{2}{e}\gamma_k e^{-w} S_{\tau_{k,w}} - \frac{2\gamma_k^2 e^{-2w} A_{\tau_{k,w}}^2}{e^2} \le \frac{4\gamma_k \psi(n_{k+1}) A_{\tau_{k,w}}}{e^{w+1}} - \frac{2\gamma_k^2 A_{\tau_{k,w}}^2}{e^{2(w+1)}}$$

$$= \frac{\psi(n_{k+1})^2 k^2 A_{\tau_{k,w}}}{\sqrt{n_{k+1}} e^w} \left(\frac{4}{e} - \frac{2k^2 A_{\tau_{k,w}}}{e^2 \sqrt{n_{k+1}} e^w}\right)$$

$$\leq \frac{\psi(n_{k+1})^2 k^2 A_{\tau_{k,w}}}{\sqrt{n_{k+1}} e^w} \left(\frac{4}{e} - \frac{2}{e^2} \sqrt{e^4 - \frac{(1+\delta)k^4 n_k}{n_{k+1} e^{2w}}}\right)$$

$$\leq -\psi(n_{k+1})^2 \frac{k^2}{\sqrt{n_{k+1}} e^w} \times \frac{\sqrt{n_{k+1}} e^{w+2}}{k^2} \times \frac{1}{2}$$

$$= -\frac{e^2 \psi(n_{k+1})^2}{2}.$$

The last inequality holds because $\lim_{k\to\infty} k^4 n_k / n_{k+1} = 0$ and 4/e - 2 < -1/2. Hence $\mathcal{L}_{\tau_{k,w}}^{\gamma_k e^{-w}, k} \to 0$ uniformly in $1 \le w \le \lceil \ln k \rceil$. This implies $\limsup_{k\to\infty} \mathcal{M}_{\tau_{k+1}-\tau_k}^{\gamma_k, k} \le 0$.

Proposition 4.11. Let $\omega \in \Omega_C$. Suppose that $v_k \leq \min(\tau_{k+1}, \sigma_{k,C})$ and

$$-A_n\psi^U(A_n^2) \leq S_n, \ \forall n \in [\tau_k, \nu_k].$$

Then for sufficiently large k

$$\mathcal{N}_{\nu_k}^{\gamma_k,D} \geq \frac{\alpha}{2}.$$

Proof. As in the proof of the previous lemma, we denote $t = n - \tau_k$, $S_t = S_n - S_{\tau_k}$ and $A_t^2 = A_n^2 - A_{\tau_k}^2$. We distinguish two cases:

(a)
$$A_{\nu_k}^2 \leq \frac{\psi(n_{k+1})^2}{c_1 \gamma^2}$$
, (b) $\frac{\psi(n_{k+1})^2}{c_1 \gamma^2} < A_{\nu_k}^2 \leq A_{\tau_{k+1}}^2 - A_{\tau_k}^2$.

For case (a), for sufficiently large k and for any $\gamma \le \gamma_k$, as in (58),

$$\begin{split} \gamma S_{\nu_{k}} &\leq \gamma \left(S_{\nu_{k}-1} + c_{\nu_{k}} \right) \leq \gamma \left(\left((1+\delta)A_{\tau_{k}} + \sqrt{A_{\tau_{k}}^{2} + A_{\nu_{k}-1}^{2}} \right) \psi(n_{k+1}) + (1+\delta)C \frac{\sqrt{A_{\tau_{k}}^{2} + A_{\nu_{k}-1}^{2}}}{\psi(A_{\tau_{k}}^{2})^{3}} \right) \\ &\leq \psi(n_{k+1})^{2} \left(\frac{1}{\sqrt{c_{1}}} + \delta \right) \end{split}$$

and

$$\psi(n_{k+1})e^{-\psi(n_{k+1})^2/2}2e^{(1+\delta)^5e^6C}e^{\gamma S_{\nu_k}}\to 0 \quad (k\to\infty).$$

Hence $\mathcal{N}_{\nu_k}^{\gamma_k,D} \geq \alpha/2$ uniformly in $\gamma \in [\gamma_k/k, \gamma_k]$.

For case (b), S_{ν_k} can be evaluated as

$$S_{\nu_{k}} \leq S_{\nu_{k}-1} + c_{\nu_{k}} \leq S_{\nu_{k}-1} + (1+\delta)C\frac{\sqrt{A_{\tau_{k}}^{2} + A_{\nu_{k}-1}^{2}}}{\psi(A_{\tau_{k}}^{2})^{3}}$$
$$\leq \left((1+\delta)A_{\tau_{k}} + \sqrt{A_{\tau_{k}}^{2} + A_{\nu_{k}}^{2}}\right)\psi(n_{k+1}) + (1+\delta)C\frac{\sqrt{A_{\tau_{k}}^{2} + A_{\nu_{k}}^{2}}}{\psi(A_{\tau_{k}}^{2})^{3}}$$

$$\leq \left((1+\delta)A_{\tau_{k}} + \sqrt{A_{\tau_{k}}^{2} + A_{\nu_{k}}^{2}} \left(1 + \frac{(1+\delta)C}{\psi(A_{\tau_{k}}^{2})^{3}\psi(n_{k+1})} \right) \right) \psi(n_{k+1})$$

by (51). Put

$$q_k^2 := \frac{A_{\tau_k}^2}{A_{\nu_k}^2} \le \frac{c_1 \gamma_k^2}{\psi(n_{k+1})^2}, \qquad s_k := \frac{(1+\delta)C}{\psi(A_{\tau_k}^2)^3 \psi(n_{k+1})},$$

so that $\lim_{k} q_k \psi(n_{k+1})^2 = 0$ and $\lim_{k} s_k \psi(n_{k+1})^2 = 0$. Then for sufficiently large k

$$\begin{aligned} \frac{S_{\nu_k}^2}{2A_{\nu_k}^2} &\leq \left((1+\delta)^2 \frac{q_k^2}{2} + (1+\delta)(1+s_k)q_k \sqrt{1+q_k^2} + (1+s_k)^2 \left(\frac{1}{2} + \frac{q_k^2}{2}\right) \right) \psi(n_{k+1})^2 \\ &\leq \frac{\psi(n_{k+1})^2}{2} + \delta. \end{aligned}$$

Then

$$\psi(n_{k+1})e^{-\psi(n_{k+1})^2/2} \times 2e^{(1+\delta)^5 e^6 C} e^{S_{\nu_k}^2/(2A_{\nu_k}^2)} \frac{\sqrt{2\pi}}{\gamma A_{\nu_k}} \le 2e^{(1+\delta)^5 e^6 C+\delta} \sqrt{2\pi c_1} e^{\delta} \le \frac{D\alpha}{4}.$$

4.6 Dynamic strategy forcing the sharpness

Finally, we prove Proposition 4.1. We assume that by the validity result, Skeptic already employs a strategy forcing $S_n \ge -A_n \psi^U(A_n^2) a.a.$ for $\omega \in \Omega_C$. In addition to this strategy, based on Proposition 4.10, consider the following strategy.

Start with initial capital $\mathcal{K}_0 = \alpha$. Set k = 1.

Do the followings repeatedly:

- 1. Apply the strategy in Proposition 4.10 for $n \in [\tau_k, \tau_{k+1}]$.
 - If $\tau_{k+1} < \min(\sigma_{k,C}, \nu_k)$, then go to 2. Otherwise go to 3.
- 2. Let k = k + 1. Go to 1.
- 3. Wait until $\exists k'$ such that $-\sqrt{\tau_{k'}}\psi^U(\tau_{k'}) \leq S_{\tau_{k'}} \leq \sqrt{\tau_{k'}}\psi(\tau_{k'})$. Set k = k' and go to 1.

By this strategy Skeptic keeps his capital non-negative for every path ω . For $\omega \in \Omega_0$, $\tau_k = \infty$ for some *k* and Skeptic stays in Step 1 forever. For $\omega \in \Omega_{=\infty}$, Step 3 is performed infinite number of times, but the overshoot of $|x_n|$ in Step 3 does not make Skeptic bankrupt by Proposition 4.11. Now consider $\omega \in \Omega_C$. Since Skeptic already employs a strategy forcing $S_n \ge -A_n \psi^U(A_n^2) a.a.$, the lower bound in (54) violated only finite number of times. By $\omega \in \Omega_C$, $n \ge \sigma_{k,C}$ is happens only finite number of times. Hence if $S_n \le A_n \psi(A_n^2) a.a.$, then Step 3 is performed only finite number of times and there exists k_0 such that only Step 2 is repeated for all $k \ge k_0$. Now for each iteration of Step 2, Skeptic multiplies his capital at least by

$$1 + \frac{1-\delta}{D} \lceil \ln k \rceil \psi(n_{k+1}) e^{-\psi(n_{k+1})^2/2}$$

Then

$$\frac{1-\delta}{D}\sum_{k=k_0}^{\infty} \lceil \ln k \rceil \psi(n_{k+1}) e^{-\psi(n_{k+1})^2/2} \le \prod_{k=k_0}^{\infty} \left(1 + \frac{1-\delta}{D} \lceil \ln k \rceil \psi(n_{k+1}) e^{-\psi(n_{k+1})^2/2} \right).$$

Since the left-hand side diverges to infinity, the above strategy forces the sharpness.

References

- [1] Z. D. Bai. A theorem of Feller revisited. Ann. Probab., 17(1):385–395, 1989.
- [2] I. Berkes, S. Hörmann, and M. Weber. Upper-lower class tests for weighted i.i.d. sequences and martingales. J. Theoret. Probab., 23(2):428–446, 2010.
- [3] M. Csörgő, B. Szyszkowicz, and Q. Wang. Darling-Erdős theorem for self-normalized sums. Ann. Probab., 31(2):676–692, 2003.
- [4] V. H. de la Peña, M. J. Klass, and T. L. Lai. Self-normalized processes: exponential inequalities, moment bounds and iterated logarithm laws. *Ann. Probab.*, 32(3A):1902–1933, 2004.
- [5] U. Einmahl and D. M. Mason. Some results on the almost sure behavior of martingales. In *Limit theorems in probability and statistics (Pécs, 1989)*, volume 57 of *Colloq. Math. Soc. János Bolyai*, pages 185–195. North-Holland, Amsterdam, 1990.
- [6] P. Erdős. On the law of the iterated logarithm. Ann. of Math. (2), 43:419–436, 1942.
- [7] W. Feller. The general form of the so-called law of the iterated logarithm. *Trans. Amer. Math. Soc.*, 54:373–402, 1943.
- [8] W. Feller. The law of the iterated logarithm for identically distributed random variables. *Ann. of Math.* (2), 47:631–638, 1946.
- [9] P. S. Griffin and J. D. Kuelbs. Some extensions of the LIL via self-normalizations. *Ann. Probab.*, 19(1):380–395, 1991.
- [10] N. C. Jain, K. Jogdeo, and W. F. Stout. Upper and lower functions for martingales and mixing processes. Ann. Probab., 3:119–145, 1975.
- [11] A. Kolmogoroff. Über das Gesetz des iterierten Logarithmus. Math. Ann., 101(1):126–135, 1929.
- [12] M. Kumon, A. Takemura, and K. Takeuchi. Capital process and optimality properties of a Bayesian skeptic in coin-tossing games. *Stoch. Anal. Appl.*, 26(6):1161–1180, 2008.
- [13] K. Miyabe and A. Takemura. Convergence of random series and the rate of convergence of the strong law of large numbers in game-theoretic probability. *Stochastic Process. Appl.*, 122(1):1–30, 2012.
- [14] K. Miyabe and A. Takemura. The law of the iterated logarithm in game-theoretic probability with quadratic and stronger hedges. *Stochastic Process. Appl.*, 123(8):3132–3152, 2013.
- [15] W. Philipp and W. F. Stout. Invariance principles for martingales and sums of independent random variables. *Math. Z.*, 192(2):253–264, 1986.
- [16] P. Révész. Random Walk in Random and Non-Random Environments. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, third edition, 2013.
- [17] G. Shafer and V. Vovk. *Probability and Finance: It's only a game!* Wiley Series in Probability and Statistics. Financial Engineering Section. Wiley-Interscience, New York, 2001.

- [18] W. F. Stout. A martingale analogue of Kolmogorov's law of the iterated logarithm. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, 15:279–290, 1970.
- [19] V. Strassen. Almost sure behavior of sums of independent random variables and martingales. In Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. II: Contributions to Probability Theory, Part 1, pages 315–343. Univ. California Press, Berkeley, Calif., 1967.