# TOWARD A CONJECTURE OF TAN AND TU ON FIBERED GENERAL TYPE SURFACES

A. HUITRADO-MORA, M. CASTAÑEDA-SALAZAR AND A. G. ZAMORA

ABSTRACT. Given a semistable non-isotrivial fibered surface  $f:X\to\mathbb{P}^1$  it was conjectured by Tan and Tu that if X is of general type, then f admits at leats 7 singular fibers. In this paper we prove this conjecture in several particular cases, i.e. assuming f is obtained from blowing-up the base locus of a transversal pencil on an exceptional minimal surface S or assuming that f is obtained as the blow-up of the base locus of a transversal and adjoint pencil on a minimal surface.

## 1. Notation

We work on the complex field number  $\mathbb{C}$ . All the considered varieties will be assumed irreducible and projective. Through the paper we shall use the following notation:

- . X will be a general type surface and S its minimal model.  $\pi:X\to S$  will be the associated chain of blowing-downs.
- .  $f: X \to \mathbb{P}^1$  will be a semi-stable, non-isotrivial fibration, and F the general fibre. We set g = the genus of F. By C we will denote the image of F under  $\pi$  and

$$\Lambda: S \dashrightarrow \mathbb{P}^1$$
.

the pencil induced by f. We denote by s the number of singular fibers.

- . We shall say that  $\Lambda$  is transversal if its general member  $C \in \Lambda$  is non-singular and intersects transversally any other general member  $C' \in \Lambda$ .
- . We'll freely use the standard notation in surfaces' theory. In particular  $q=h^1(X,\mathcal{O}_X)$  will be the irregularity of X and  $p_g=h^0(X,K_X)$  its geometric genus. By e(X) will be denote the topological Euler characteristic.
- Given divisors  $D_1$  and  $D_2$  in an algebraic surface we denote as usual  $D_1 \equiv D_2$  for the numeric equivalence and  $D_1 \sim D_2$  for the linear one. Most of the time we will be working on regular surfaces and in this case we indistinctly use both symbols.
- . The number m will be:

$$m := K_S^2 - K_X^2 = e(X) - e(S).$$

Date: 03/25/16.

2000 Mathematics Subject Classification. 14D06, 14J25.

Key words and phrases. Algebraic Surfaces, Fibrations, Singular Fibres.

This paper was partially supported by CONACyT Grant 257079. First and second authors were supported by Conacyt Doctoral Scholarships. Third author was partially supported by Conacyt Grant 265621 for Academic Sabbatical Year.

Note that in general  $m \leq C^2$ . Adjunction Formula gives the important inequality  $C.K_S + m \le 2(g-1)$  with equality holding if  $\Lambda$  is transversal.

## 2. Introduction

Let  $f: X \to \mathbb{P}^1$  be a non-isotrivial semistable fibered surface. It is a classical result that such a fibration admits a certain number of singular fibers (in contrast to the case when the base of the fibration is not rational or elliptic [1]). In the seminal paper [2] it was proved that this number s must be at least 4. Subsequently the bound have been sharpened to  $s \geq 5$  if  $g \geq 2$  and  $s \geq 6$  if the surface X is not birationally ruled ([11] [12], [13]). It was conjectured by Tan and Tu, in a preprint previous to [13] that this bound must raise to 7 if X is of general type (Tan-Tu conjecture for what follows). They also proved the conjecture for genus  $2 \leq g \leq 4$  and characterized fibrations of genus 5 with s=6 on a general type surface as those obtained from blowing up the base locus of a transversal pencil on a Horikawa surface. Using this characterization the proof of the conjecture for q = 5 was completed in [15].

Roughly speaking the proof of these bounds are based, in case  $g \geq 2$  on the canonical class inequality:

$$K_f^2 < 2(g-1)(2g_B - 2 + s),$$

for any non iso-trivial semi-stable fibration  $f: X \to B$  of genus  $g \geq 2$ . Here  $K_f$  is the relatively canonical divisor,  $K_f := K_X - f^*K_B$ , which turns out to be  $K_X(-2F)$  if  $B=\mathbb{P}^1$ . The bounds for s are obtained from the positivity properties of  $K_f$  and  $K_f(-F)$ .

Unfortunately, this approach is useless for proving Tan-Tu conjecture, since in this case the only relevant information the inequality provides is that  $K_X^2 < 0$  if s=6. However for most of the cases a fibered surface (of general type or not) satisfies  $K_X^2 < 0$ . Indeed, such a surface is obtained by blowing up the base locus of some pencil  $\Lambda$  on a minimal surface S.

In this paper we deal with Tan-Tu conjecture in some particular cases. We first impose on the minimal model S of X the conditions of being exceptional in the sense that either  $K_S^2 = 2$ ,  $p_g = 3$  or  $K_S^2 = 1$  and  $p_g = 2$  (see [1], VII.8) and assuming that f is obtained as the blowing-up of the base locus of a transversal pencil  $\Lambda$  in S. In these cases we are able to prove the conjecture using explicit descriptions of these surfaces as double coverings of rational surfaces, by means of the canonical or bi-canonical map. This is the content of Section 3, Theorems 4.1 and 4.2. In this sense there is some hope of extending the result to a wider class of surfaces.

Next, in section 5 we prove the conjecture assuming that the pencil  $\Lambda$  is adjoint, i.e.  $C = B + K_S$  with B a big and nef divisor in S and  $K_S^2 \geq 3$  and for  $K_S^2 \leq 2$ assuming not only that  $\Lambda$  is adjoint, but also transversal (Theorem 5.3). The case  $K_S^2 = p_q = 1$  is the subtler and is stated and proved in Proposition 5.2.

We list below the cases in which Tan-Tu conjecture have been proved in this article:

- . If  $\chi(\mathcal{O}_S) = 1$  with no extra assumption on S or  $\Lambda$ .
- . If either  $K_S^2=1,\,p_g=2$  or  $K_S^2=2,\,p_g=3$  assuming  $\Lambda$  is transversal. . If  $\Lambda$  is transversal and adjoint and  $K_S^2\leq 2$  or if  $\Lambda$  is merely adjoint and  $K_S^2 \ge 3$ .

# 3. Some general facts and results

The following inequality will be systematically used: given a semi-stable non-isotrivial fibration of genus  $g \ge 2$ ,  $f: X \to B$ , for any integer  $e \ge 2$ :

$$\frac{1}{3}e^2(K_X^2 - 2(g-1)(6(g_B - 1) + s - s/e)) \le e_f.$$

Original formulation involves the number of (-2) vertical curves in X, but for our purposes this version will be sufficient. The proof can be founded in [12] and is based on successive changes of the base B of the fibration. We call this Tan's inequality. In particular if  $B = \mathbb{P}^1$  (our interest's case) and s = 6 we obtain:

(1) 
$$\frac{1}{3}e(K_X^2e + 12(g-1)) \le e_f.$$

Useful forms of this inequality are collected in the following:

**Lemma 3.1.** Let  $f: X \to \mathbb{P}^1$  be semistable, non-isotrivial of genus  $g \geq 2$ . If s = 6, then evaluating (1) we obtain:

i) 
$$K_S^2 + C.K_S \le 3\chi(\mathcal{O}_S) \text{ if } e = 3,$$

ii) 
$$19K_S^2 + 18C \cdot K_S < m + 36\chi(\mathcal{O}_S) \text{ if } e = 4.$$

iii) 
$$7K_S^2 + 6C.K_S < m + 9\chi(O_S) \text{ if } e = 5.$$

*Proof.* Evaluate (1) at the indicated value of e and substitute:

$$m + C.K_S \le 2(g - 1),$$

$$e(X) = 12\chi(\mathcal{O}_X) - K_X^2$$
 (Noether's Formula),

$$K_X^2 = K_S^2 - m,$$

and

$$e_f = 4(g-1) + e(X)$$
 (because  $f$  is semistable) .

We start by sharpening the bound for m obtained in [13], inequality (3.2) (compare with the proof of Theorem 2.1(4) in [13]).

**Lemma 3.2.** Assume  $f: X \to \mathbb{P}^1$  is a semistable fibration and the minimal model S of X is a general type surface, then:

$$m \leq C^2 \leq \frac{4(g-1) + K_S^2 - \sqrt{8(g-1)K_S^2 + (K_S^2)^2}}{2}.$$

*Proof.* From Index Hodge Theorem applied to  $K_S$  and C we get:

$$mK_S^2 \le C^2 K_S^2 \le (C.K_S)^2$$
.

Adjoint formula gives  $C.K_S \leq 2(g-1) - m$ , therefore

$$0 \le m^2 - (4(g-1) + K_S^2)m + 4(g-1)^2.$$

Consider the right hand term of the previous inequality as a polynomial in m. Its discriminant turns out to be

$$\Delta = 8(g-1)K_S^2 + (K_S^2)^2.$$

Thus its roots are:

$$m_{\pm} = \frac{4(g-1) + K_S^2 \pm \sqrt{\Delta}}{2}.$$

It follows that either  $m \leq m_-$  or  $m \geq m_+$ . We claim that  $m \leq m_-$  is the only possible case. Indeed, if

$$\frac{4(g-1) + K_S^2 + \sqrt{\Delta}}{2} \le m,$$

then  $C.K_S \leq 2(g-1) - m \leq 0$  that is impossible.

As a consequence of Lemma 3.2 we obtain our first general fact concerning the number s:

**Proposition 3.3.** Let S be of general type. If s = 6, then

$$K_S^2 + \sqrt{8(g-1)K_S^2 + (K_S^2)^2} \le 6\chi(\mathcal{O}_S).$$

In particular, if  $g \ge 6$  and  $\chi(O_S) = 1$ , then  $s \ge 7$ .

*Proof.* Assume s = 6, by Tan's inequality:

$$\frac{1}{3}e(K_X^2e + 12(g-1)) \le e_f,$$

for any natural number  $e \ge 2$ . Evaluating in e = 3 we obtain:

$$3K_X^2 + 12(g-1) \le e_f.$$

Since f is semistable  $e(X) = -4(g-1) + e_f$  and (2) becomes:

(3) 
$$3K_X^2 + 12(g-1) \le e_f = e(X) + 4(g-1).$$

By definition  $m=K_S^2-K_X^2=e(X)-e(S)$ . This, combined with Noether Formula  $e(S)=12\chi(\mathfrak{O}_S)-K_S^2$  leads to:

$$K_S^2 + 2(g-1) \le 3\chi(\mathcal{O}_S) + m.$$

The desired inequality follows after applying Lemma 3.2.

The importance of this Proposition is that given a family of general type surfaces with given invariants  $K_S^2$  and  $\chi(\mathcal{O}_S)$  only a finite numbers of values of g must be discharged in order to conclude that a fibered surface birational to S has at least 7 singular fibers. This principle will be illustrated in the next section.

Note that in Proposition 3.3 the hypothesis of being S of general type is essential ([2], Example 2).

## 4. Fibrations obtained from exceptional surfaces

Surfaces satisfying either  $K_S^2 = 2$  and  $p_g = 3$  or  $K_S^2 = 1$  and  $p_g = 2$  are called exceptional because of the behavior of the tri-canonical map (Theorem VII 8.3 in [1]). In this section we study fibrations in these surfaces by means of the canonical and bi-canonical map, respectively.

**Theorem 4.1.** Assume S satisfies  $K_S^2 = 2$  and  $p_g = 3$ . Let  $f: X \to \mathbb{P}^1$  be obtained as the blowing-up of a transversal pencil  $\Lambda$  in S. Then,  $s \geq 7$ .

*Proof.* By Debarre's Inequality ([6]) we know that q = 0 and therefore  $\chi(\mathcal{O}_S) = 4$ . Assume s = 6, by Proposition 3.3 it is sufficient to consider:

$$6 \le g \le 31.$$

In this case the canonical map  $\phi_{K_S}$  defines a 2 : 1 covering:

$$\phi_{K_S}: S \to \mathbb{P}^2,$$

ramified along  $R \equiv 4K_S$  ([8]). Consider the restriction  $\phi := \phi_{K_S}|_C$ .

First, we consider the case  $\phi$  is a 2:1 covering. Denote by  $G \subset \mathbb{P}^2$  the image of C under  $\phi$  and d for its degree. We have  $C = \phi^*G$  and therefore:

$$m = C^2 = (\phi^* G)^2 = 2G^2 = 2d^2.$$

On the other hand, taking into account that  $\phi^*H = K_S$ , with H a hyperplane section (i.e. the divisor associated with  $\mathcal{O}_{\mathbb{P}^2}(1)$ ) we obtain:

$$d = \frac{C.K_S}{2}$$
 and  $m = \frac{(C.K_S)^2}{2}$ .

From this we get  $2m = (2(g-1) - m)^2$ . The possible values of g satisfying such a relation in the range  $6 \le g \le 31$  are: g = 7, 13, 21, 31. The corresponding values of m and  $C.K_S$  are listed below:

-			
ĺ	g-1	$C.K_S$	m
ĺ	6	4	8
ĺ	12	6	18
ĺ	20	8	32
ſ	30	10	50

If g - 1 = 20 or 30 we use the inequalities of Lemma 3.1 ii), in order to obtain a contradiction.

Assume g - 1 = 6. In this case the number of singular points in the fibers of f will be:

$$e_f = e(X) + 4(g - 1)$$

$$= -K_X^2 + 12\chi(\mathcal{O}_X) + 4(g - 1)$$

$$= 6 + 48 + 24 = 78.$$

Since s=6 there exists at least a singular fiber  $F_0$  of f containing  $\sigma_0=13=78/6$  singular points. But then, denoting  $F_0=F_1+\ldots+F_l$  for the decomposition into irreducible components:

$$6 = (g - 1) = \sum_{i=1}^{l} (g_i - 1) + \sigma_0$$
$$\geq \sum_{i=1}^{l} (g_i - 1) + 13,$$

with  $g_i$  standing for the geometric genus of  $F_i$  and  $\sigma_0$  for the number of singular points of  $F_0$ . We have:

$$\sum_{i=1}^{l} (g_i - 1) \le -7.$$

In particular,  $F_0$  have at least 7 irreducible rational components. Being  $\phi$  a 2:1 covering and C be applied under  $\phi$  to a plane curve G of degree  $d=C.K_S/2=2$  we have that the number l of irreducible components of  $F_0$  is at most 4. In this way we get a contradiction with the assumption s=6.

The case g-1=12 follows after similar considerations, this time tanking into account that the covering  $\phi$  sends C onto a curve G of degree 3.

Consider now the case  $\phi$  restricted to C is 1 : 1. In this case there exists a curve C' such that

$$C + C' = \phi^* G$$
.

Moreover, since the ramifications of  $\phi$  over C occurs exactly on the intersections of C and C' we have  $C.C' = C.R = C.4K_S$ . Similarly, we conclude that  $C.K_S = C'.K_S$ . Also, we have that:

$$2C.K_S = \phi^*G.K_S = 2d,$$

i.e.  $d = C.K_S$ .

It follows that  $\phi^*G = \phi^*((C.K_S)H) = (C.K_S)K_S$ . From

$$C + C' = (C.K_S)K_S,$$

we obtain, after intersecting with C:

$$4K_S.C = C.C' = (C.K_S)^2 - m.$$

The possibilities for such a relationship are listed below:

(g - 1)	$C.K_S$	m
5	5	5
9	6	12
14	7	21
20	8	32
27	9	45

Using Lemma 3.1 we obtain a contradiction in the following cases: g - 1 = 20 and 27 when evaluating at e = 4, and g - 1 = 5, 9 when evaluating at e = 5.

Finally, we must analyze the case g - 1 = 14. In this case,

$$e_f = 4(g-1) - K_X^2 + 12\chi(\mathfrak{O}_X)$$
  
= 123.

Assuming s = 6 there must to exist a singular fiber  $F_0$  with its number of nodes  $\sigma_0 \ge 21$ . Assume  $F_0 = F_1 + ... + F_l$  is its decomposition into irreducible components. Denoting by  $g_i$  the geometric genus of  $F_i$  we have:

$$14 = \sum_{i=1}^{l} (g_i - 1) + \sigma_0 \ge \sum_{i=1}^{l} (g_i - 1) + 21.$$

From this we get  $l \geq 7$ . Now, the image of  $F_0$  in  $\mathbb{P}^2$  (under the composition  $\phi \circ \pi$ ) is a degree 7 curve  $G_0$ . It follows that l = 7,  $G_0 = L_1 + ... + L_7$  is the sum of seven lines  $L_i$ , and  $C_0 := \pi(F_0)$  must be the sum of seven irreducible components  $C_0 = C_1 + ... + C_7$  ( $C_i = \pi(F_i)$ ). We have, moreover,  $\phi^* L_i = C_i + C_i' \equiv K_S$ ,  $C_i^2 = C_i'^2 = -3$  and  $C_i C_i' = 4$ .

A simple cohomological computation shows that  $h^0(\mathcal{O}_S(C_{i_1}+C_{i_2}+C_{i_3})=1$  for any indexes  $i_1,i_2,i_3\in\{1,...,7\}$  and  $h^0(\mathcal{O}_S(C_1+...+C_4))=2$ . This, together with  $(C_1+...+C_4)^2=0$  means that  $|C_1+...+C_4|$  is a base point free pencil. Call  $\Delta:=C_1+...+C_4$ . We have,  $C_i'.\Delta=0$  for i=5,6,7, thus  $C_5'+C_6'+C_7'$ , being connected, must be a vertical divisor with respect to  $|\Delta|$ . We conclude the existence of an effective divisor D' such that  $\Delta\sim C_5'+C_6'+C_7'+D'$ . It is easy to deduce that D' is a rational (-3)-curve and  $\phi(D')$  is a line in  $\mathbb{P}^2$ .

From this relation and

$$(C_1 + C_1') + \dots + (C_7 + C_7') \sim 7K_S,$$

it follows that:

$$C \sim C_1 + ... + C_7 \sim 3K_S + D'.$$

This gives a contradiction, as the image  $\phi(C)$  is a degree 7 curve and the image  $\phi(3K_S+D')$  is a degree 4 curve.

**Theorem 4.2.** Assume  $f: X \to \mathbb{P}^1$  is obtained as a blow up of a transversal pencil  $\Lambda$  on a minimal surface S with  $K_S^2 = 1$  and  $p_q = 2$ . Then  $s \ge 7$ .

*Proof.* By Proposition 3.3 we must consider only the values  $6 \le g \le 37$ .

Considering the classical Horikawa's construction and notation ([9]), let  $\Pi : \bar{S} \to S$  be the blowing up centered in the base point p of  $|K_S|$ , denote by E its exceptional divisor and consider the ramified double covering:

$$\phi_2: \bar{S} \to \mathbb{F}_2.$$

The map  $\phi_2$  is given as follow: the bicanonical map of S determines a double cover on the singular quadric  $Q \subset \mathbb{P}^3$ , the singular point being the image of p.  $\phi_2$  is the induced map on  $\bar{S}$  after considering the desingularization  $\mathbb{F}_2$  of the quadric. The locus branch of  $\phi_2$  is the divisor  $B = 6\Delta + 10\Gamma$ , with  $\Delta$  and  $\Gamma$  denoting respectively the class of the (-2)-section and the class of the fiber in  $\mathbb{F}_2$  of the structural morphism and the ramification divisor R is  $5K_S + E$ . Here and in what follows given D any divisor in S we just write D for the divisor  $\Pi^*D$  in  $\bar{S}$ .

Denote by  $\bar{\Lambda}$  the induced pencil on  $\bar{S}$ . Depending on whether p is a base point of  $\Lambda$  or not we have

(4) 
$$\bar{C} = \begin{cases} \Pi^* C & \text{if} \quad p \notin \Lambda \\ \Pi^* C - E & \text{if} \quad p \in \Lambda. \end{cases}$$

Let G be the image of  $\bar{C}$  under  $\phi_2$ . If we denote  $G = a\Delta + b\Gamma$  and considering that  $\phi^*\Delta = 2E, \ \phi^*\Gamma = K_S - E$  we have

(5) 
$$\phi^* G = bK_S + (2a - b)E.$$

Let be  $\phi := \phi_2|_{\bar{C}} : \bar{C} \to G$  with  $\deg \phi = n = 1$  or 2.

We analyze the two cases n=1 or 2 and within each one the subcases  $p\in\Lambda$  or not.

Case 1  $\phi$  is 2:1. In this case  $\phi^*G = \bar{C}$ .

Assume first that  $p \notin \Lambda$ .

By (4) and (5) we have that 2a - b = 0, using this  $\bar{C}^2 = m = b^2$ . On the other hand  $m = bC.K_S$ , therefore  $b = C.K_S$ . The next table shows the possible values of m,  $C.K_S$  and g - 1.

$C.K_S$	a	m	g-1
4	2	16	10
6	3	36	21
8	4	64	36

Assume g-1=10. In this case the number of singular points in the fibers of f will be:

$$e_f = 12\chi(\mathcal{O}_X) - K_X^2 + 4(g-1)$$
  
=  $36 + 15 + 4(10) = 91$ 

Since s = 6 there exists at least a singular fiber  $F_0$  of f containing 16 singular points. Let  $F_0 = F_1 + \ldots + F_l$  be the decomposition into irreducible components, then:

(6) 
$$10 = g - 1 \ge \sum_{i=1}^{l} (g_i - 1) + \sigma_0$$
$$\ge \sum_{i=1}^{l} (g_i - 1) + 16$$

where  $g_i$  denotes the geometric genus of  $F_i$ . This imply that  $l \geq 6$ . Moreover there are at least 6 of this components that are rational curves and are mapping onto rational components of  $G_0$ .

From [7] (Corollary V.5.18) we know that the possible irreducible curves in  $\mathbb{F}_2$  are:  $\Gamma$ ,  $\Delta$  and  $\alpha\Delta + \beta\Gamma$  with  $\alpha > 0$ ,  $\beta \geq 2a$ . Let  $G_0 = G_1 + \ldots + G_s$  be the decomposition into irreducible components.

Note that even being  $\Delta$  a rational curve, is not a possibility for any of the  $G_i$ 's, that because if  $G_1 = \Delta$  then

$$\bar{C}_1 = \phi^* G_0 = 2E$$

that contradicts the semistability of f.

With respect to the components  $G_i = a_i \Delta + b_i \Gamma$ ,  $a_i > 0$ ,  $b_i \ge 2a$ ,  $G_i$  is rational if and only if  $a_i = 1$  and  $b_i \ge 2$ . Since  $a = \sum_{i=1}^s a_i$  and  $b = \sum_{i=1}^s b_i$  the only possible

decomposition is  $G_0 = G_1 + G_2$  with  $G_i = \Delta + 2\Gamma$ , i = 1, 2, i.e. we can't have the 6 needed rational components.

If g-1=21,36 we use similar arguments with  $\sigma_0=26,41$  respectively. In both cases there exist at least 5 rational components and this is not possible because a<5.

Now consider the case  $p \in \Lambda$ .

By (4) and (5) b = 2a + 1, so  $m - 1 = \bar{C}^2 = 4a(a + 1)$  and  $m = b^2$ . Moreover,

$$\bar{C}.K_{\bar{S}} = C.K_S + 1 = b + 1$$

Therefore  $m = (C.K_S)^2$ . Keeping in mind the previous notation, we get the next possible values

$C.K_S$	a	m	g-1	$e_f$	$\sigma_0$	$l \ge$
3	1	9	6	68	12	6
5	2	25	15	120	20	5
7	3	49	28	196	33	5

Observe that l is, as before, the minimal number of rational components in  $G_0$ , so analogous to the case  $p \notin \Lambda$ , all possibilities in the table can't occur because of a < l.

Case  $2 \phi$  is 1:1. In this case there exists a divisor  $\bar{C}'$  such that  $\phi^*G = \bar{C} + \bar{C}'$ . Denote as before by  $F_0$  a singular fiber of f,  $C_0$  its image under  $\pi$ . If  $F_0 = F_1 + \ldots + F_k$  is the decomposition of  $F_0$  into irreducible components, we denote by  $C_0 = C_1 + \ldots + C_k$  the corresponding decomposition for  $C_0$  and by  $\bar{C}_0 = \bar{C}_1 + \ldots + \bar{C}_k$  the corresponding curves and decomposition in  $\bar{S}$  and by  $G_0 = G_1 + \ldots + G_k$  their images in  $\mathbb{F}_2$ .

We begin by stating the following:

**Lemma 4.3.** In the previous situation, let  $G_1$  be any irreducible component of  $G_0$ , then neither  $G_1 \sim \Delta$ , nor  $G_1 \sim \Gamma$  nor  $G_1 \sim \Delta + 2\Gamma$ . In particular, if  $G_0 \sim a\Delta + b\Gamma$ , then the number of rational irreducible components of  $G_0$  is least or equal than a.

*Proof.* Let  $G_1$  be equivalent to  $\Delta$ . Then  $\phi_2^*(G_1) \sim 2E$ . Note that there exists a divisor  $\bar{C}_1'$  such that  $\bar{C}_1 + \bar{C}_1' = \phi_2^*(G_1)$ . This implies  $\bar{C}_1 = E$ , which is impossible by the definition of  $\bar{C}$ .

Now, assume  $G_1 \equiv \Gamma$ , then  $\phi^* G_1 \equiv K_S - E = C_1 + C'_1$ . From this, intersecting with  $K_S$ , and using that  $K_S$  is nef we obtain that either  $C_1.K_S = 0$  or  $K_S.C'_1 = 0$ . It follows that there exists a curve D on S with  $K_S.D = 0$ , which contradicts that ampleness of  $|K_S|$ .

Finally, suppose that  $G_1 = \Delta + 2\Gamma$ , in this case  $\phi^* G_1 \equiv 2K_S$ .

Note that for any decomposition  $2K_S = A + A'$  we must have that both, A and A' must be irreducible and equivalent to  $K_S$  and A.A' = 1. Indeed, assume A irreducible, that from  $2K_S = A + A'$  with easy it follows that  $A.K_S = A'.K_S = 1$ . Therefore,  $A^2 = 1$ , because  $2K_S$  is 1-connected,  $A \equiv K_S$  follows from HIT.

One stablished this fact, just note that if  $2K_S = C_1 + C_1'$ , then  $\phi: C_1 \to G_1$  can not be 1:1, because  $C_1$  is a genus 2 curve.

The last assertion follows from the fact that the only irreducible rational curves on  $\mathbb{F}_2$  are equivalent to either  $\Delta$ ,  $\Gamma$  or  $\Delta + b\Gamma$  with  $b \geq 2$ .

Continuing the proof of the Theorem, assume first that  $p \notin \Lambda$ . We have the commutative diagram:

$$\overline{S} \xrightarrow{\phi_2} \mathbb{F}_2 ,$$

$$\Pi \downarrow \qquad \qquad \downarrow$$

$$S \xrightarrow{\phi_{2K}} Q$$

therefore  $\phi_{2K}(\Pi(\bar{C}'))$  contains the singular point of Q and from this it follows that  $\bar{C}' = \Pi^*C'$ , for some effective divisor C' in S. By (5) b = 2a and  $\bar{C} + \bar{C}' \sim bK_S$ . Therefore,

$$C.K_S + C'.K_S = (\bar{C} + \bar{C}')K_{\bar{S}} = (bK_S)(K_S + E) = b.$$

The ramifications of  $\phi_2$  occurring on  $\bar{C}$  are given by the intersections of  $\bar{C}$  and  $\bar{C}'$ , so we have  $\bar{C}.\bar{C}'=\bar{C}.R=\bar{C}'.R$ . In particular, the right hand term of the previous equation implies that  $C.K_S=C'.K_S$  and  $b=2C.K_S$ .

Moreover,

$$bC.K_S = \phi^* G.\bar{C} = (\bar{C} + \bar{C}').\bar{C} = m + \bar{C}.\bar{C}' = m + 5C.K_S.$$

We conclude that  $m = C.K_S(2C.K_S - 5)$  and we get the possible values (keeping in mind the previous notation introduced for  $\sigma_0$  and l):

b	$C.K_S$	m	g-1
8	4	12	8
10	5	25	15
12	6	42	24
14	7	63	35

The values g - 1 = 24, 35 are impossible because of the Hodge Index Theorem.

If g-1=8 we have that  $e_f=79$  and then there must exists a singular fiber  $F_0$  with at least 14 singular points. It follows that  $G_0$  has 6 or more rational components. Using that a=4 and Lemma 4.3, we get a contradiction. The case g-1=15 follows after similar considerations.

It remains to analyze the case  $p \in \Lambda$ .

As in the previous case, like  $\bar{C}=C-E$  also  $\bar{C}'=C'-E$ . From this we get 2a-b=-2. Therefore we have:

$$\phi_2^* G = C + C' - 2E = bK_S - 2E.$$

Moreover,  $\bar{C}.R = \bar{C}'.R$  and then  $C.K_S + 1 = \bar{C}.K_{\bar{S}} = \bar{C}'.K_{\bar{S}}$ We get the next formulas

$$\phi_2^* G. K_{\bar{S}} = b + 2 = 2C. K_S + 2$$

and therefore

$$m + 5C.K_S = \phi_2^*G.\bar{C} = (bK_S - 2E).(C - E) = bC.K_S - 2$$

We conclude that  $b = 2C.K_S$  and  $m = 2(C.K_S)^2 - 5C.K_S - 2$ . The table of possible values is:

$C.K_S$	b	a	m	g-1
4	8	3	10	7
5	10	4	23	14
6	12	5	40	23
7	14	6	61	34

If g-1=23,34 we get a contradiction by Hodge Index Theorem. If g-1=7, then  $e_f=73$ , therefore there exists a singular fiber  $F_0$  with at least 13 singular points and at least 6 rational components. Taking in consideration that a=3 and Lemma 4.3 we obtain a contradiction. The case g-1=14 is similar.

## 5. The adjoint case

In this section we consider fibrations  $f: X \to \mathbb{P}^1$  satisfying the property that C is an adjoint linear system, i.e.,  $C \equiv B + K_S$  with B a big and nef divisor. The typical example for bearing in mind is  $C \equiv nK_S$ , i.e. the fibration f is obtained after blowing up the base locus of a generic pencil of curves  $\Lambda \subset |nK_S|$ .

We collect, for further use, some general elemental facts in the following:

**Lemma 5.1.** Assume  $C \equiv B + K_S$  with B a big and nef divisor, denoting by  $g_B$  the arithmetic genus of B, we have:

i) 
$$2(g_B - 1) = B^2 + B.K_S.$$

ii) 
$$m = (q_B - 1) + (q - 1).$$

iii) 
$$(g-1) = (g_B - 1) + B.K_S + K_S^2.$$

iv)  $2 \leq B.K_S$ , and if  $g \geq 5$ , then

$$g+1 \leq m$$
.

*Proof.* Assertions i)-iii) follow immediately from adjunction formula. As for iv), note that it is enough to prove that  $2 \leq B.K_S$ , because then using i)  $g_B - 1 \geq 2$  and the desired inequality follows from ii). Now, if  $B.K_S = 1$ , by Index Hodge Theorem  $B^2 = K_S^2 = 1$  and applying from  $m = B^2 + 2B.K_S + K_S^2$  we have m = 4. On the other hand, from i) and ii)  $m = 1 + (g - 1) = g \geq 5$ .

We start by studding the case  $K_S^2 = 1$ , which is similar in nature to Theorems 4.1 and 4.2:

**Proposition 5.2.** Let  $f: X \to \mathbb{P}^1$  be a fibration obtained as the blowing up of the base locus of a transversal and adjoint pencil  $\Lambda$  on a minimal surface with  $K_S^2 = 1$ . Then,  $s \geq 7$ .

*Proof.* By Noether Inequality  $p_g \leq 2$ , and by Proposition 3.3 and Theorem 4.2, we can assume  $p_g = 1$ . It is well known that for a surface with such invariants the bicanonical map  $\phi_{2K_S}$  defines a 4:1 morphism onto  $\mathbb{P}^2$  ([5], [10], [14]):

$$\phi_{2K_S}: S \to \mathbb{P}^2,$$

ramified along a divisor  $R \equiv 7K_S$ . We consider, as before, the restriction of this map to C:



In this case n is a divisor of 4,  $G \subset \mathbb{P}^2$  is the image of C and j denotes its normalization. Denote by d the degree of G.

If we assume s=6 we only need, according to Proposition 3.3 to consider  $6 \le g \le 16$ .

We start by analyzing the case n=4: in this case we have  $C=\phi^*G$  and therefore:

$$m = C^2 = (\phi^* G)^2 = 4G^2 = 4d^2.$$

On the other hand, taking in account that  $\phi^* H \equiv 2K_S$ , with H hyperplane section (i.e the divisor associated with  $\mathcal{O}_{\mathbb{P}^2}(1)$ ) we obtain:

$$C.K_S = \phi^* G.K_S = \phi^* (dH).K_S = 2d.$$

Adjunction formula gives 2(g-1) = 2d(2d+1). The only value of g that satisfies the relation in the range  $6 \le g \le 16$  is g-1=10 with d=2 and m=16. Evaluating in Tan's inequality for e=4 (Lemma 3.1 ii))we obtain a contradiction.

If n < 4, then there exists an effective divisor C' > 0 such that:

$$C + C' \equiv \phi_{2K_S}^* G \equiv 2dK_S.$$

Note that:

$$C'.K_S = 2d - C.K_S$$

and

$$C'^2 = 4d^2 - 4dC.K_S + C^2.$$

Moreover, since  $h^1(C) = h^2(C) = 0$ ,

$$h^0(C) = \frac{C^2 - C.K_S}{2} + 2.$$

Next, the only possibility for  $h^2(C') = h^0(K_S - C') \neq 0$  is  $C' \equiv K_S$ , because  $p_q = 1$ . This would imply  $C \equiv K_S$ , which is impossible.

It is easy to prove that  $H^0(C) \simeq H^0(C')$  and by Riemann-Roch we get:

$$h^0(C) = h^0(C') \ge \frac{C'^2 - C' \cdot K_S}{2} + 2,$$

and substituting the values of  $C'^2$  and  $C'.K_S$ :

$$h^0(C) \ge h^0(C) + \frac{4d^2 - (4d - 2)C.K_S - 2d}{2}.$$

This implies,

$$2d^2 - (2d - 1)C.K_S - d \le 0$$

that is equivalent to  $d \leq C.K_S$ .

Now, assume n = 1. The linear system  $|2K_S| |_C$  defines a base point free linear system on C and the associated map a 1:1 cover onto a plane degree d curve.

Thus, we have  $d = 2C.K_S$  and we obtain a contradiction with the just obtained bound  $d \leq C.K_S$ .

The case n=2 remains to be analyzed: in this case we have  $C.K_S=d$ . Note that the intersections of C and C' gives place to ramifications points of  $\phi_{2K_S}$ . Therefore:

$$2dC.K_S - m = C.C' < C.R = 7C.K_S.$$

From this we get:

$$d(2d - 7) \le m \le 2d^2 - 2.$$

It follows that  $d \leq 5$ . In general we have that

(7) 
$$e_f = e(X) + 4(g-1)$$
$$= 24 - (1-m) + 4(g-1),$$
$$= 23 - d + 6(g-1).$$

Thus, assuming s=6 and  $d\leq 3$ , there must exits a singular fiber  $F_0$  of f having at least (g-1)+4 nodes. Call  $\sigma_0$  the number of nodes of  $F_0$ . Note that the number of nodes of  $C_0=\pi(F_0)$  is also  $\sigma_0$ . Then,  $\sigma_0\geq 9$ . On the other hand, the plane curve  $G_0=\phi_{2K_S}(C_0)$ , being of degree  $d\leq 3$  admits at most 3 nodes. In this way we get the contradiction  $\sigma_0\leq 6$ .

Similar argumentations lead to contradictions for the cases d=4,5. Indeed, if d=4, then  $\sigma_0 \geq g+3$ . We have:

$$g-1 = \sum_{i=1}^{l} (g_i - 1) + \sigma_0 \ge -l + g + 3,$$

with  $g_i$  standing for the geometric genus of the components  $F_i$  of  $F_0$ . It follows that  $l \geq 4$  and therefore  $F_0$  (and in consequence  $C_0$ ) has at least 4 rational components. From this it follows that  $G_0$  has at least 2 irreducible rational components. Taking in account that  $G_0$  is a degree 4 curve we have that, either  $G_0$  contains a line as an irreducible component or it is the product of two irreducible conics. If  $G_0$  is the product of two irreducible conics then it has only 4 nodes and we get, as before a contradiction, in any other case, if L is an irreducible component of  $G_0$  and  $C_0 = C_1 + \ldots + C_l$ , then

$$\phi_{2K_S}^* L = C_i + C_i',$$

for  $C_i$  some rational components of  $C_0$ . But then:

$$2K_S \equiv C_i + C_i',$$

and it follows ([4] Lemma 1, page 181) that  $C_i = \Delta$ , the only effective divisor in  $|K_S|$ . This give a contradiction, since  $\Delta$  is a curve of geometric genus 2.

Finally the case d=5 follows after similar considerations. In this case  $C_0$  admits at least 3 irreducible rational components and  $G_0$  at least 2 irreducible rational components. The only subtle case to be treated careful being the possibility that  $G_0=Q+E$ , with Q an irreducible conic and E a singular irreducible cubic. But in this case either Q or E must satisfies that it pull back under  $\phi_{2K_S}$  is the sum of two irreducible components  $C_i+C_j$  of  $C_0$ .

Note that  $\phi_{2K_S}^*E = C_i + C_j$  is impossible, because then

$$6K_S \equiv C_i + C_i$$

and then  $K_S(C_i + C_i) = 6$ , that contradicts  $K_S(C_0) = 5$ . On the other hand

$$4K_S = \phi_{2K_S}^* Q = C_i + C_j$$

implies that  $C_0$  has exactly 3 irreducible components:  $C_0=C_1+C_2+C_3$  . Suppose i=1 and j=2, then  $C_3.K_S=1$  and

$$\phi_{2K_S}: C_3 \to E,$$

must be a 2:1 map onto a degree cubic and we obtain the contradiction

$$2K_S.C_3 = 6.$$

Finally we have:

**Theorem 5.3.** Let  $f: X \to \mathbb{P}^1$  a semi-stable non-isotrivial fibration obtained as the blow-up of the base locus of an adjoint pencil  $\Lambda$  on the minimal surface S. Then if  $K_S^2 \geq 3$  the number s of singular fibers of f is at least 7.If  $K_S^2 \leq 2$  and  $\Lambda$  is also transversal, then  $s \geq 7$ .

*Proof.* We assume s=6 and  $2 \le K_S^2$ . From Noether's inequality:

$$p_g \le \frac{K_S^2}{2} + 2,$$

we have:

$$\chi(\mathcal{O}_S) \le \frac{K_S^2}{2} + 3.$$

Applying Lemma 3.3:

$$K_S^2 + \sqrt{8K_S^2(g-1) + (K_S^2)^2} \le 3K_S^2 + 18,$$

which implies

$$8K_S^2(g-1)+(K_S^2)^2 \leq 4(K_S^2)^2+72K_S^2+18^2.$$

Substituting  $2(g-1) \ge C.K_S + m = B.K_S + K_S^2 + m$ , we get:

$$m \leq \frac{72 - 4B.K_S}{4} + \frac{18^2 - (K_S^2)^2}{4K_S^2}.$$

Now, combine the previous bound for m with Lemma 3.1 ii), in order to deduce:

$$19K_S^2 + 19B.K_S \le 18 + \frac{18^2 - (K_S^2)^2}{4K_S^2} + 108.$$

Using again the adjoint hypothesis, this amount to:

$$\frac{(K_S^2)}{4} + 19C.K_S \le 166,$$

that is,

$$C.K_S < 8.$$

Now, use Hodge Index Theorem:

$$m \le \frac{(C.K_S)^2}{K_S^2} \le \frac{64}{K_S^2},$$

and apply one more time Lemma 3.1 ii):

$$K_S^2 + 18C.K_S \le \frac{64}{K_S^2} + 108,$$

$$19K_S^2 + 18B.K_S \le \frac{64}{K_S^2} + 108.$$

Finally, using  $2 \leq B.K_S$ , (Lemma 5.1 iv)) we arrive to:

$$19K_S^2 \le \frac{64}{K_S^2} + 72.$$

This implies  $K_S^2 \leq 4$ . Thus, only the cases  $K_S^2 = 2, 3, 4$  remains to be discharged. This is easy and essentially is a reproduction of the previous argument.

For instance, for case  $K_S^2 = 2$  we have by Lemma 3.2 and Proposition 3.3 that  $g \le 16$  and  $m \le 26$ , moreover, assuming  $\Lambda$  is transversal, we can apply Theorem 4.1 and assume that  $\chi(\mathcal{O}_S) \le 3$ . Evaluating Tan's Inequality at e = 5 (Lemma 3.1 iii)) we get:

$$7K_S^2 + 6C.K_S \le m + 9\chi(\mathcal{O}_S),$$

that, under our fixed values becomes  $C.K_S \leq 6$ . Using Hodge Index Theorem we obtain  $m \leq 18$ . Evaluating again Tan's Inequality at e=4 we have  $C.K_S \leq 4$  and  $m \leq 8$ . Finally, we evaluate once again Tan's Inequality at e=5 and get the final contradiction  $C.K_S \leq 3$  and  $g+1 \leq m \leq 4$ .

Cases  $K_S^2 = 3,4$  are quite analogous, only that in these cases we don't need Theorem 4.1, and in consequence the transversality hypothesis can be avoided.  $\Box$ 

## References

- [1] Barth, W., Peters, C., Van de Ven, A. Compact Complex Surfaces Springer Verlag (1984).
- [2] Beauville, A.: Le nombre minimum de fibers singulieres d'un courbe stable sur P¹. Asterisque, 86, (1981), 97-108.
- [3] Beauville A.: Surfaces algébriques complexes. Asterisque 54, (1978).
- [4] Bombieri, E. Canonical models of surfaces of general type. Publ. Math. IHES, 42, (1973), 171-219.
- [5] Catanese, F. Surfaces with K2 = pg = 1 and their period mapping. In Algebraic geometry (Proc. Summer Meeting, Univ. Copenhagen, 1978), Springer LNM, 732 (1979), 1-29.
- [6] Debarre, O. Inégalités numériques pour les surfaces de type général. Bull. Soc. Math. France 110 (3), (1982), 319-346.
- [7] Hartshorne, R. Algebraic Geometry. Springer-Verlag GTM, (1997).
- [8] Horikawa, E. Algebraic surfaces of general type with small  $c_1^2$ . I. Annals of Mathematics, 104, (1976), 357-387.
- [9] Horikawa, E. Algebraic surfaces of general type with small c<sub>1</sub><sup>2</sup> II, Inventiones Math. 37 (1976), 121-155.
- [10] Kunev, V. Thesis for Masters Degree. Sofia University, (1976).
- [11] Liu, K. Remarks on the Geometry of Moduli Spaces Proceedings of the AMS 124, (3), (1996), 689-695.
- [12] Tan S-L: The minimal number of singular fibers of a semi stable curve over P<sup>1</sup>. J. Algebraic Geom., 4, (1995), 591-596.

- [13] Tan S-L , Tu Y. , Zamora A. G. On complex surfaces with 5 or 6 semistable singular fibers over  $\mathbb{P}^1$  . Math. Zeitschrift 249 (2005), 427-438.
- [14] Todorov, A. N. Surfaces of General type with  $p_g=1$  and (K,K)=1. I Annales scientifiques de l'E.N.S. 13 (1980), 1-21.
- [15] Zamora, A. G. Genus 5 general type P<sup>1</sup>-semistable curves have at least 7 singular points. Note di Matematica 32 (2) (2012), 1-4.

CENTRO DE CIENCIAS MATEMÁTICAS,

Universidad Nacional Autónoma de México, Campus Morelia.

Apartado Postal 61-3, Santa María, 58089.

Morelia, Michoacán, México.

 $E ext{-}mail\ address:$  huitrado at matmor.unam.mx

CENTRO DE CIENCIAS MATEMÁTICAS,

Universidad Nacional Autónoma de México, Campus Morelia.

Apartado Postal 61-3, Santa María, 58089.

Morelia, Michoacán, México.

E-mail address: mcastaneda at matmor.unam.mx

U. A. MATEMÁTICAS, U.DE ZACATECAS

CAMINO A LA BUFA Y CALZADA SOLIDARIDAD, C.P. 98000

Zacatecas, Zac. México

 $E ext{-}mail\ address:}$  alexiszamora06 at gmail.com