ON SCHAUDER ESTIMATES FOR A CLASS OF NONLOCAL FULLY NONLINEAR PARABOLIC EQUATIONS

HONGJIE DONG AND HONG ZHANG

ABSTRACT. We obtain Schauder estimates for a class of concave fully nonlinear nonlocal parabolic equations of order $\sigma \in (0,2)$ with rough and non-symmetric kernels. We also prove that the solution to a translation invariant equation with merely bounded data is C^{σ} in x variable and Λ^1 in t variable, where Λ^1 is the Zygmund space. From these results, we can derive the corresponding results for nonlocal elliptic equations with rough and non-symmetric kernels, which are new even in this case.

1. Introduction

This paper is devoted to the study of Schauder estimates for a class of concave fully nonlinear nonlocal parabolic equations. There is a vast literature on Schauder estimates for classical elliptic and parabolic equations, for instance, see [14, 19, 4]. Since the work by Caffarelli and Silvestre [1, 3, 2], nonlocal equations, which naturally arise from models in physics, engineering, and finance that involve long range interactions (for instance, see [10]), attract an increasing level of interest recently. An example of nonlocal operators, which is associated with pure jump processes (see, for instance, [20]), is the following

$$L_{a}u = \int_{\mathbb{R}^{d}} \left(u(t, x+y) - u(t, x) - y^{T} D u(t, x) \right) K_{a}(t, x, y) \, dy \quad \text{for } \sigma \in (1, 2),$$

$$L_{a}u = \int_{\mathbb{R}^{d}} \left(u(t, x+y) - u(t, x) - y^{T} D u(t, x) \chi_{B_{1}} \right) K_{a}(t, x, y) \, dy \quad \text{for } \sigma = 1$$

$$\text{with} \quad \int_{S_{r}} y K_{a}(t, x, y) \, ds = 0 \quad \forall r > 0,$$

$$L_{a}u = \int_{\mathbb{R}^{d}} \left(u(t, x+y) - u(t, x) \right) K_{a}(t, x, y) \, dy \quad \text{for } \sigma \in (0, 1),$$

$$(1.1)$$

where

$$K_a \in \mathcal{L}_0 := \left\{ K : \frac{\lambda(2-\sigma)}{|y|^{d+\sigma}} \le K(t,x,y) \le \frac{\Lambda(2-\sigma)}{|y|^{d+\sigma}} \right\}$$

for some ellipticity constants $0 < \lambda \le \Lambda$, with no regularity assumption imposed with respect to the y variable. This type of nonlocal operators was first considered by Komatsu [17], Mikulevičius and Pragarauskas [20, 21], and later by Dong and Kim [12, 11], and Schwab and Silvestre [24], to name a few. In particular, the condition (1.1) appeared in all these references except for [24], where a similar cancellation condition was imposed. Notice that this class of operators is scaling invariant.

1

 $[\]rm H.$ Dong and $\rm H.$ Zhang were partially supported by the NSF under agreements DMS-1056737 and DMS-1600593.

The fully nonlinear nonlocal parabolic equation that we are interested in is of the form

$$u_t = \inf_{a \in \mathcal{A}} (L_a u + f_a), \tag{1.2}$$

where $K_a \in \mathcal{L}_0$ for $a \in \mathcal{A}$ and \mathcal{A} is an index set. For fully nonlinear second-order equations with $f_a \equiv 0$, the celebrated $C^{2,\alpha}$ estimate was established independently by Evans [13] and Krylov [18] in early nineteen-eighties. Nonhomogeneous second-order equations were considered a bit later by Safonov [23]. Recently, Caffarelli and Silvestre [2] investigated the nonlocal version of Evans-Krylov theorem with translation invariant and symmetric kernels, i.e., $K_a(x,y) = K_a(y) = K_a(-y)$, satisfying additional regularity assumptions

$$[K_a]_{C^2(\mathbb{R}^d \setminus B_\rho)} \le \Lambda(2-\sigma)\rho^{-d-\sigma-2}.$$
(1.3)

More recently, their result was extended to nonhomogeneous fully nonlinear elliptic equations by Jin and Xiong [15] by using a recursive Evans-Krylov theorem. At almost the same time, Serra [26] removed the regularity assumption (1.3) and proved the Evans-Krylov theorem and Schauder estimates with symmetric kernels. His proof relies on a Liouville type theorem and a blow-up analysis. In this paper, we do not assume that the kernels are symmetric, which is certainly more general than the kernels considered in [2, 15, 26]. Specifically, when the kernels are symmetric, (1.1) is satisfied automatically, and

$$L_a u = \frac{1}{2} \int_{\mathbb{R}^d} (u(x+y) + u(x-y) - 2u(x)) K_a(y) \, dy,$$

which is the form of the operators considered in [2, 15, 26].

For equations with non-symmetric kernels, Dong and Kim [11, 12] proved L_p and Schauder estimates for linear elliptic equations. Chang-Lara and Dávila [7, 8] considered nonlocal parabolic equations with non-symmetric kernels and critical drift, and proved the corresponding C^{α} and $C^{1,\alpha}$ estimate. Recently in [5], they proved a version of the Evans-Krylov theorem for concave nonlocal parabolic equations with critical drift, where they assumed the kernels to be non-symmetric but translation invariant and smooth (1.3). We also mention that Schauder estimates for linear nonlocal parabolic equations were studied in [16, 21].

The objective of this paper is twofold. First we extend the previous results in [26, 5, 15, 16] to include concave nonlocal parabolic equations with non-symmetric rough kernels. More specifically, for any small α , if f_a and $K_a(t, x, y)$ are C^{α} in x and $C^{\alpha/\sigma}$ in t, then we have the following $C^{1+\alpha/\sigma,\sigma+\alpha}$ a priori estimate of any smooth solution u to (1.2) in $(-1,0) \times B_1$.

Theorem 1.1. Let $\sigma \in (0,2)$, $0 < \lambda \leq \Lambda < \infty$, and A be an index set. There is a constant $\hat{\alpha} \in (0,1)$ depending on d, σ, λ , and Λ so that the following holds. Let $\alpha \in (0,\hat{\alpha})$ such that $\sigma + \alpha$ is not an integer. Assume $K_a \in \mathcal{L}_0$ and satisfies (1.1) when $\sigma = 1$, and

$$|K_a(t, x, y) - K_a(t', x', y)| \le A(|x - x'|^{\alpha} + |t - t'|^{\alpha/\sigma}) \frac{\Lambda(2 - \sigma)}{|y|^{d + \sigma}},$$
 (1.4)

where $A \ge 0$ is a constant. Suppose $u \in C^{1+\alpha/\sigma,\sigma+\alpha}(Q_1) \cap C^{\alpha/\sigma,\alpha}((-1,0) \times \mathbb{R}^d)$ is a solution of

$$u_t = \inf_{a \in \mathcal{A}} (L_a u + f_a) \quad in \ Q_1, \tag{1.5}$$

where $f_a \in C^{\alpha/\sigma,\alpha}(Q_1)$ satisfying

$$C_0 := \sup_{a \in \mathcal{A}} [f_a]_{\alpha/\sigma,\alpha;Q_1} < \infty, \quad \sup_{(t,x) \in Q_1} \left| \inf_{a \in \mathcal{A}} f_a(t,x) \right| < \infty.$$

Then,

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{1/2}} \le C||u||_{\alpha/\sigma,\alpha;(-1,0)\times\mathbb{R}^d} + CC_0,$$
 (1.6)

where C > 0 is a constant depending only on $d, \lambda, \Lambda, \alpha, A$, and σ , and is uniformly bounded as $\sigma \to 2$.

In the theorem above, $\|\cdot\|_{\alpha/\sigma,\alpha;\Omega}$ is the Hölder norm of order α/σ in t and α in x with underlying domain Ω . We used Q_r to denote the parabolic cylinder with radius r centered at the origin. For precise definitions, see Section 2. As pointed out in [26], the $C^{\alpha/\sigma,\alpha}$ Hölder norm of u on the right-hand side of (1.6) is necessary and cannot be replaced by the L_{∞} norm or any lower-order Hölder norm of u. We also note that by keeping track of the constants in the proofs below, in the symmetric case, if $\sigma \in [\sigma_0, 2)$ for some $\sigma_0 \in (0, 1)$, then the constant C in (1.6) depends on σ_0 , not σ . In the non-symmetric case, if $0 < \sigma_0 \le \sigma \le \sigma_1 < 1$ (or $1 < \sigma_2 \le \sigma < 2$), then the constant C depends on σ_0 and σ_1 (or σ_2), not σ . In particular, C does not blow up as σ approaches 2.

Roughly speaking, the proof of Theorem 1.1 can be divided into three steps. First we prove a Liouville type theorem for solutions in $(-\infty,0)\times\mathbb{R}^d$. For the classical PDEs, we generally apply interpolation and iteration to obtain $C^{1,\alpha}$ and $C^{2,\alpha}$ estimates. One notable feature of nonlocal operators is that the boundary data is prescribed on the complement of the domain where the equation is satisfied, which makes it difficult to implement these techniques. However, if we assume that (1.2) is satisfied in $(-\infty,0)\times\mathbb{R}^d$, then we do not need to handle boundary data any more, which is the advantage of considering an equation satisfied in the whole space. Second, we prove the a priori estimate for equations with translation invariant kernels by combining the Liouville theorem and a blow-up analysis. Particularly in this step, the extension from symmetric kernels to non-symmetric kernels is non-trivial. A key idea in the classical Evans-Krylov theorem for $F(D^2u) = 0$ is that, since the function F is concave, any second directional derivative D_{ee}^2u is a subsolution. It is relatively easy to adapt this idea to the nonlocal equations with symmetric kernels due to the appearance of centered second order difference in the definition of the operator. For nonsymmetric kernels, some new ideas are required to obtain a similar subsolution as in the symmetric case. Moreover, the dependence of the t variable also makes the proof more involved. Finally, we implement a more or less standard perturbation argument to treat the general case.

The second objective of this paper is to consider the end-point situation when $\alpha=0$. For second-order elliptic equations, even the Poisson equation $\Delta u=f$, when f is merely bounded, it is well known that u may fail to be $C^{1,1}$. However, this is not the case for nonlocal equations. When $\sigma \neq 1$ and the kernels are independent of t and x, we prove a priori C^{σ} estimate in the x variable and Λ^1 estimate in the t variable when f_a is merely bounded and measurable, where Λ^1 is the Zygmund space. When $\sigma=1$, we obtain a priori Λ^1 estimate in both t and x. We assume that the solution is smooth because the spaces in which estimates are obtained are not fine enough for the nonlocal operators to be defined pointwise.

Theorem 1.2. (i) Let $\sigma \neq 1$. Assume that u is a smooth solution to (1.2) in $(-\infty,0) \times \mathbb{R}^d$ with K_a independent of t and x. When $\sigma \in (1,2)$, we also assume

that Du is $C^{(\sigma-1)/\sigma}$ in t. Then there exists a constant C depending on d, λ, Λ , and σ such that for $\sigma > 1$,

$$[u]_{\Lambda^1}^t + [u]_{\sigma}^* + [Du]_{\frac{\sigma-1}{\sigma}}^t \le C \sup_a ||f_a||_{L_{\infty}};$$

and for $\sigma < 1$,

$$[u]_{\Lambda^1}^t + [u]_{\sigma}^* \le C \sup_a ||f_a||_{L_{\infty}}.$$

(ii) Let $\sigma = 1$. Assume that u is a smooth solution to (1.2) in $(-\infty, 0) \times \mathbb{R}^d$ with K_a independent of t and x. Then there exists a constant C depending on d, λ, Λ , and σ such that

$$[u]_{\Lambda^1} \leq C \sup_a ||f_a||_{L_\infty}.$$

Here all the norms are taken in $\mathbb{R}_0^{d+1} := (-\infty, 0) \times \mathbb{R}^d$.

In the theorem above, $[\cdot]^*$, $[\cdot]^t_{\alpha}$, $[\cdot]^t_{\Lambda^1}$, and $[\cdot]_{\Lambda^1}$ are the Hölder semi-norms in x, t, the Zygmund semi-norm in t, and the Zygmund semi-norm with respect to (t,x), respectively. See the precise definitions in Section 2.

We localize Theorem 1.2 to obtain the following corollary.

Corollary 1.3. (i) Let $\sigma \neq 1$. Assume that u is a smooth solution to (1.2) with K_a independent of t and x. Then for $\sigma > 1$,

$$[u]_{\sigma;Q_{1/2}}^* + [u]_{\Lambda^1(Q_{1/2})}^t + [Du]_{\frac{\sigma-1}{\sigma};Q_{1/2}}^t \le C \bigg(\sup_{a \in \mathcal{A}} \|f_a\|_{L_{\infty}(Q_1)} + \|u\|_{L_{\infty}((-1,0) \times \mathbb{R}^d)} \bigg);$$

and for $\sigma < 1$,

$$[u]_{\sigma;Q_{1/2}}^* + [u]_{\Lambda^1(Q_{1/2})}^t \le C \Big(\sup_{a \in \mathcal{A}} ||f_a||_{L_{\infty}(Q_1)} + ||u||_{L_{\infty}((-1,0) \times \mathbb{R}^d)} \Big).$$

(ii) Let $\sigma = 1$. Assume that u is a smooth solution to (1.5) with K_a independent of t and x. Then we have

$$[u]_{\Lambda^1;Q_{1/2}} \le C \Big(\sup_{a \in \mathcal{A}} \|f_a\|_{L_{\infty}(Q_1)} + \|u\|_{L_{\infty}((-1,0) \times \mathbb{R}^d)} \Big).$$

To our best knowledge, such result is new even for nonlocal elliptic equations with symmetric kernels. A similar result was obtained very recently by Mou [22] for elliptic equations with symmetric, smooth kernels and Dini continuous data. With merely bounded and measurable data, the best estimates known in the literature are the C^{β} regularity of u in t for $\beta < 1$ and the C^{γ} regularity of u in x for $\gamma < \min\{\sigma, 1 + \alpha\}$, where $\alpha > 0$ is a small constant. See [25, 9]. Because $\Lambda^1 \subseteq C^{\beta}$ for any $\beta < 1$, Theorem 1.2 improves these results and is optimal even in the linear case. See Remark 1.4 below.

The proof of Theorem 1.2 is based on a perturbation type argument using Campanato's approach. We first refine the estimate in Theorem 1.1 when the operator is translation invariant. In particular, we replace $||u||_{\alpha/\sigma,\alpha;(-1,0)\times\mathbb{R}^d}$ on the right-hand side of (1.6) by

$$[u]_{\alpha/\sigma,\alpha;(-1,0)\times B_2} + \sum_{j=2}^{\infty} 2^{-j\sigma} [u]_{\alpha/\sigma,\alpha;(-1,0)\times (B_{2^j}\setminus B_{2^{j-1}})}.$$
 (1.7)

The advantage of the replacement will be explained below. Another important ingredient in the proof is the fact that, for example, when $\sigma > 1$,

$$[u]_{\Lambda^{1}(\mathbb{R}_{0}^{d+1})}^{t} + [u]_{\sigma;\mathbb{R}_{0}^{d+1}}^{*} + [Du]_{\frac{\sigma-1}{\sigma};\mathbb{R}_{0}^{d+1}}^{t} \leq C \sup_{r>0} \sup_{(t,x)\in\mathbb{R}_{0}^{d+1}} E[u;Q_{r}(t,x)], \qquad (1.8)$$

where

$$E[u; Q_r(t, x)] := \inf_{p \in \mathcal{P}_1} r^{-\sigma} ||u - p||_{L_{\infty}(Q_r(t, x))},$$

 \mathcal{P}_1 is the set of linear functions in (t, x), and $Q_r(t, x)$ is the parabolic cylinder with center (t, x); see (2.1). Therefore, instead of directly estimating

$$[u]_{\Lambda^1}^t + [u]_{\sigma}^* + [Du]_{\frac{\sigma-1}{\sigma}}^t,$$

we estimate $E[u; Q_r(t, x)]$ for any fixed r and (t, x). It is worth noting that in view of the proofs of Lemmas 5.2, 5.4, and 5.5, the two quantities on the left and right hand sides of (1.8) are actually equivalent.

More specifically, without loss of generality, we set (t, x) = (0, 0) and let v_K solve the homogeneous equation

$$\begin{cases} \partial_t v_K = \inf_{a \in \mathcal{A}} L_a v_K & \text{in } Q_{2R} \\ v_K = g_K := \max\{-K, \min\{u - p, K\}\} & \text{in } (-(2R)^{\sigma}, 0) \times B_{2R}^c \end{cases}$$

where K is a large constant, p is a carefully chosen linear function, and R>2r is a constant to be determined. Now we apply Theorem 1.1 to v_K and control $[v_K]_{1+\alpha/\sigma,\alpha+\sigma;Q_{R/2}}$ by using scaling argument and replacing $||v_K||_{\alpha/\sigma,\alpha}$ by (1.7). It is easily seen that in each cylindrical domain $(-R^{\sigma},0)\times(B_{2^jR}\setminus B_{2^{j-1}R})$, the Hölder norm of g_K is bounded and independent of K, but globally it depends on K and goes to infinity as $K\to\infty$. This is also the advantage of decomposing the domain into annuli. We then set q_K to be the first-order Taylor expansion of v_K and estimate

$$||u - p - q_K||_{L_{\infty}(Q_r)} \le ||u - p - v_K||_{L_{\infty}(Q_r)} + ||v_K - q_K||_{L_{\infty}(Q_r)},$$

where the first term is bounded by CR^{σ} due to the Aleksandrov-Bakelman-Pucci estimate and second term is controlled by $[v_K]_{1+\alpha/\sigma,\alpha+\sigma;Q_r}$. Finally, we are able to obtain

$$r^{-\sigma} \| u - p - q_K \|_{L_{\infty}(Q_r)} \le C(r/R)^{\alpha} \left([u]_{\sigma}^* + [u]_{\Lambda^1}^t + [Du]_{\frac{\sigma-1}{\sigma}}^t \right) + C(R/r)^{\sigma} \| f \|_{L_{\infty}}.$$

By setting R = Mr, using (1.8), and taking M sufficiently large, the terms involving u on the right-hand side above are absorbed in the left-hand side.

Remark 1.4. We give two simple examples which indicate that the estimates in Theorem 1.2 (and thus in corollary 1.3) are optimal even in the linear case. Set d=1. Let $f=f(t,x)=\chi_{\{t<-M|x|^{\sigma}\}}$ for some constant M>0 and u be a solution to the equation

$$u_t + (-\Delta)^{\sigma/2} u = f$$
 in $(-\infty, 0) \times \mathbb{R}^d$.

Then by the explicit representation of solutions, it is easily seen that for sufficiently large M,

$$\lim_{t \to 0} u_t(t,0) = -\infty.$$

Thus u cannot be Lipschitz in t in this case. Next we set $\sigma = 1$, $g = g(t, x) = \chi_{\{t < 0, x > t\}}$, and v be a solution to

$$v_t + (-\Delta)^{1/2}u = g$$
 in $(-\infty, 0) \times \mathbb{R}^d$.

Again by the explicit representation of solutions,

$$\lim_{x \to 0} v_x(0, x) = \infty.$$

Therefore, v cannot be Lipschitz in x in this case.

By viewing solutions to elliptic equations as steady state solutions to parabolic equations, from Theorems 1.1, 1.2, and Corollary 1.3, we obtain the corresponding results for nonlocal elliptic equations with nonsymmetric and rough kernels.

The organization of this paper is as follows. In the next section, we introduce some notation and preliminary results that are necessary in the proof of our main results. We prove the Liouville theorem in Section 3 and Theorem 1.1 in Section 4. In Section 5, we apply Theorem 1.1 to prove Theorem 1.2.

Remark added after the proof. After we finished the paper, we learnt that Chang-Lara and Kriventsov [9] also established a Schauder estimates for fully non-linear nonlocal parabolic equations with rough kernels, by using a different method. In their paper, it is assumed that the kernels are time-independent and symmetric. They also obtained a C^{β} estimate of u in t for $\beta < 1$ when the data is bounded and measurable.

2. Notation and preliminary results

In this section, we introduce some notation which will be used throughout this paper and some preliminary results which are useful in our proof. We use $B_r(x)$ to denote the Euclidean ball in \mathbb{R}^d with center x and radius r. The parabolic cylinder $Q_r(t,x)$ is defined as follows

$$Q_r(t,x) = (t - r^{\sigma}, t) \times B_r(x). \tag{2.1}$$

We simply use Q_r to denote $Q_r(0,0)$ and $\mathbb{R}_0^{d+1} := (-\infty,0) \times \mathbb{R}^d$. Let $\Omega \subset \mathbb{R}^{d+1}$ and we define the Hölder semi-norm as follows: for any $\alpha, \beta \in (0,1]$, and function f,

$$[f]_{\beta,\alpha;\Omega} = \sup \Big\{ \frac{|f(t,x) - f(s,y)|}{\max(|x-y|^{\alpha},|t-s|^{\beta})} : (t,x), (s,y) \in \Omega, (t,x) \neq (s,y) \Big\}.$$

We denote

$$||f||_{\beta,\alpha;\Omega} = ||f||_{L_{\infty}(\Omega)} + [f]_{\beta,\alpha;\Omega}.$$

For any nonnegative integers m and n,

$$\|f\|_{n+\beta,m+\alpha;\Omega} = \|f\|_{L_\infty(\Omega)} + [D^m f]_{\beta,\alpha;\Omega} + [\partial_t^n f]_{\beta,\alpha;\Omega}.$$

The spaces corresponding to $\|\cdot\|_{\alpha,\beta;\Omega}$ and $\|\cdot\|_{m+\alpha,n+\beta;\Omega}$ are denoted by $C^{\alpha,\beta}(\Omega)$ and $C^{m+\alpha,n+\beta}(\Omega)$, respectively. Next, for any $\alpha,\beta\in(0,1]$, we define the Hölder semi-norms only with respect to x or t

$$[f]_{\alpha;\Omega}^* = \sup \left\{ \frac{|f(t,x) - f(t,y)|}{|x - y|^{\alpha}} : (t,x), (t,y) \in \Omega, x \neq y \right\},$$
$$[f]_{\beta;\Omega}^t = \sup \left\{ \frac{|f(t,x) - f(s,x)|}{|t - s|^{\beta}} : (t,x), (s,x) \in \Omega, t \neq s \right\}.$$

When $\sigma = k + \alpha$ with some integer $k \geq 1$,

$$[f]_{\sigma:\Omega}^* = [D^k f]_{\alpha:\Omega}^*$$

For $\alpha \in (0,2)$, we define the Lipschitz-Zygmund semi-norm and norm by

$$[u]_{\Lambda^{\alpha}} = \sup_{|h|>0} |h|^{-\alpha} ||u(\cdot + h) + u(\cdot - h) - 2u(\cdot)||_{L_{\infty}},$$

$$||u||_{\Lambda^{\alpha}} = ||u||_{L_{\infty}} + [u]_{\Lambda^{\alpha}}.$$

We say $u \in \Lambda^{\alpha}$ if $||u||_{\Lambda^{\alpha}} < \infty$.

For simplicity of notation, we denote

$$\delta u(t, x, y) = \begin{cases} u(t, x + y) - u(t, x) - y^T D u(t, x) & \text{for } \sigma \in (1, 2), \\ u(t, x + y) - u(t, x) - y^T D u(t, x) \chi_{B_1} & \text{for } \sigma = 1, \\ u(t, x + y) - u(t, x) & \text{for } \sigma \in (0, 1). \end{cases}$$

The Pucci extremal operator is defined as follows: for $\sigma \neq 1$

$$\mathcal{M}^{+}u(t,x) = \int_{\mathbb{R}^{d}} \left(\Lambda \delta u(t,x,y)^{+} - \lambda \delta u(t,x,y)^{-}\right) \frac{2-\sigma}{|y|^{d+\sigma}} dy,$$
$$\mathcal{M}^{-}u(t,x) = \int_{\mathbb{R}^{d}} \left(\lambda \delta u(t,x,y)^{+} - \Lambda \delta u(t,x,y)^{-}\right) \frac{2-\sigma}{|y|^{d+\sigma}} dy.$$

When $\sigma = 1$, the extremal operator cannot be written out explicitly, due to the condition (1.1). Nevertheless, we do not use exact representation directly and define the extremal operator by

$$\mathcal{M}^+ u = \sup_a L_a u$$
 and $\mathcal{M}^- u = \inf_a L_a u$,

where the infimum (or supremum) is taken with respect to all L_a 's with kernels K_a satisfying (1.1).

We recall the weak Harnack inequality of [24, Theorem 6.1].

Proposition 2.1. Assume that $0 < \sigma_0 \le \sigma < 2$ and C > 0 is a constant. Let u be a function such that

$$u_t - \mathcal{M}^- u \ge -C$$
 in Q_1 , $u \ge 0$ in $(-1,0) \times \mathbb{R}^d$.

Then there are constants $C_1 > 0$ and $\varepsilon_1 \in (0,1)$ depending only on $\sigma_0, \lambda, \Lambda$, and d, such that

$$\left(\int_{(-1,-2^{-\sigma})\times B_{1/4}} u^{\varepsilon_1} dx dt\right)^{1/\varepsilon_1} \le C_1 \left(\inf_{Q_{1/4}} u + C\right).$$

From Proposition 2.1, we obtain the following corollary for $\sigma \in (1,2)$, the proof of which is provided in the appendix.

Corollary 2.2. Let $\sigma_2 \in (1,2)$, $\sigma \in (\sigma_2,2)$, C>0 be constants, and u satisfy

$$u_t - \mathcal{M}^- u \ge -C$$
 in Q_{2r} , $u \ge 0$ in $(-(2r)^{\sigma}, 0) \times \mathbb{R}^d$.

Let ε_1 be the constant in Proposition 2.1. For any $r, \delta \in (0,1)$, denote $\tilde{Q}_{\delta r} = (-r^{\sigma}, -(\delta r)^{\sigma}) \times B_r$. Then we have

$$r^{-(d+\sigma)/\varepsilon_1} \Big(\int_{\tilde{Q}_{\delta r}} u^{\varepsilon_1} \, dx \, dt \Big)^{1/\varepsilon_1} \le C_2 \Big(\inf_{Q_{\delta r/2}} u + C r^{\sigma} \Big),$$

where $C_2 > 0$ is a constant depending only on δ , σ_2 , λ , Λ , and d.

We state the following local boundedness estimate from [6, Corollary 6.2].

Proposition 2.3. Let $\Omega \subset \mathbb{R}^d$, $t_1 < t_2$, and u satisfy

$$u_t - \mathcal{M}^+ u \le 0$$
 in $(t_1, t_2] \times \Omega$.

Then for any $(t'_1, t_2] \times \Omega' \subset \subset (t_1, t_2] \times \Omega$,

$$\sup_{\Omega' \times (t_1', t_2]} u^+ \le C(2 - \sigma) \int_{t_1}^{t_2} \int_{\mathbb{R}^d} \frac{u^+}{1 + |x|^{d + \sigma}} \, dx \, dt,$$

where C depends on Ω , Ω' , t_1 , t_2 , and t'_1 .

Let us point out that the kernels considered in [6] are more general than our kernels. Specifically, Chang-Lara and Dávila considered when $\sigma \in [1, 2)$

$$Lu = (2 - \sigma) \int_{\mathbb{R}^d} \hat{\delta}u(x, y) K(y) \, dy + b \cdot Du(x),$$

where $\hat{\delta}u(x,y) = u(x+y) - u(x) - Du(x)y\chi_{B_1}$, $K(y) \in \mathcal{L}_0$, and for some $\beta > 0$,

$$\sup_{r \in (0,1)} r^{\sigma - 1} \left| b + (2 - \sigma) \int_{B_1 \setminus B_r} y K(y) \, dy \right| \le \beta. \tag{2.2}$$

Note that for $\sigma > 1$, since

$$\delta u(x,y) = \hat{\delta}u(x,y) - Du(x)y\chi_{B_1^c},$$

we can rewrite our operator and get

$$b = -(2 - \sigma) \int_{B_1^c} y K(y) \, dy.$$

Obviously, $|b| \leq C$, where C depends d, σ , and Λ , and it is easy to check that (2.2) holds for b and K above.

The next proposition is [24, Theorem 7.1].

Proposition 2.4. Let $0 < \sigma_0 \le \sigma < 2$ and u satisfy in Q_1

$$u_t - \mathcal{M}^+ u \le C_0$$
 and $u_t - \mathcal{M}^- u \ge -C_0$.

Then there are constants $\gamma \in (0,1)$ and C>0 only depending on d, σ_0, λ , and Λ such that

$$[u]_{\gamma/\sigma,\gamma;Q_{1/2}} \le C ||u||_{L_{\infty}((-1,0);L_{1}(\omega_{\sigma}))} + CC_{0}.$$

Here

$$||u||_{L_{\infty}((-1,0);L_{1}(\omega_{\sigma}))} = (2-\sigma) \sup_{t \in (0,1)} \int_{\mathbb{R}^{d}} \frac{|u(t,x)|}{1+|x|^{d+\sigma}} dx.$$

Note that we replaced $||u||_{L_{\infty}((-1,0)\times\mathbb{R}^d)}$ by $||u||_{L_{\infty}((-1,0);L_1(\omega_{\sigma}))}$, which follows from a simple localization argument. See, for instance, [6, Corollary 7.1]. In the sequel, we always assume $\gamma < \sigma$.

We finish this section by proving the following global Hölder estimate.

Lemma 2.5. Let u satisfy in Q_1

$$u_t - \mathcal{M}^+ u \leq C_0$$
 and $u_t - \mathcal{M}^- u \geq -C_0$.

where C_0 is a constant and $u \equiv 0$ in $\mathbb{R}_0^{d+1} \setminus Q_1$. Then there exists a constant $\alpha \in (0,1)$ depending on d, λ, Λ , and σ (uniformly as $\sigma \to 2$), so that

$$[u]_{\alpha/\sigma,\alpha;Q_1} \leq CC_0,$$

where C depends on d, λ, Λ , and σ , which is uniformly bounded as $\sigma \to 2$.

Proof. Thanks to the interior Hölder estimate Proposition 2.4, it suffices to prove the estimate near the parabolic boundary of Q_1 . We consider the lateral boundary and bottom separately. Define $\phi : \mathbb{R}^d \to \mathbb{R}^+$ as

$$\phi(x) = \begin{cases} x_d^{\beta} & \text{for } x_d > 0 \\ 0 & \text{for } x_d \le 0 \end{cases},$$

where $\beta \in (0, 1)$. We claim that for sufficiently small $\beta \in (0, \sigma_0)$ depending on d, λ , Λ , and σ_0 , we have

$$\mathcal{M}^+\phi(x) < -\hat{C}x_d^{\beta-\sigma}$$
 in $\{x_d > 0\},$

where \hat{C} depends on d, λ, Λ , and σ_0 . By scaling, it is obvious that

$$\mathcal{M}^+\phi(x) = x_d^{\beta-\sigma}\mathcal{M}^+\phi(e),$$

where e = (0, 0, ..., 0, 1). Therefore, we only need to estimate $\mathcal{M}^+\phi(e)$. Case 1: $\sigma > 1$. By definition,

 $\mathcal{M}^+\phi(e)$

$$= (2 - \sigma)x_d^{\beta - \sigma} \left(\int_{\{y_d > -1\}} + \int_{\{y_d < -1\}} \right) \left(\Lambda(\delta \phi(e, y))^+ - \lambda(\delta \phi(e, y))^- \right) \frac{1}{|y|^{d + \sigma}} dy$$

=: $(2 - \sigma)x_d^{\beta - \sigma} (I_1 + I_2)$.

When $y_d > -1$, by concavity, it follows that

$$\phi(e+y) = (1+y_d)^{\beta} < 1 + \beta y_d \text{ and } \delta\phi(e,y) \le 0.$$

Therefore,

$$I_{1} \leq \int_{\{|y_{d}|<1\}} \lambda \delta \phi(e, y) \frac{1}{|y|^{d+\sigma}} dy$$

$$= \lambda \int_{y_{d} \in (0, 1)} \left((1 + y_{d})^{\beta} + (1 - y_{d})^{\beta} - 2 \right) \frac{1}{|y|^{d+\sigma}} dy.$$

Notice that for any $s \in (-1, 1)$,

$$(1+s)^{\beta} < 1 + \beta s + \frac{\beta(\beta-1)}{2}s^2 + \frac{\beta(\beta-1)(\beta-2)}{6}s^3,$$

which implies that for $y_d \in (0,1)$

$$(1+y_d)^{\beta} + (1-y_d)^{\beta} - 2 < \beta(\beta-1)y_d^2.$$

Therefore,

$$I_1 < \lambda \beta(\beta - 1) \int_{y_d \in (0,1)} \frac{y_d^2}{|y|^{d+\sigma}} dy = C_1 \frac{\beta(\beta - 1)}{2 - \sigma},$$

where C_1 depends on d and λ .

Now we turn to I₂. Since $\phi(e+y)=0$ when $y_d<-1$, we have

$$I_{2} = \int_{\{y_{d}<-1\}} \frac{\Lambda(-\beta y_{d}-1)^{+} - \lambda(-\beta y_{d}-1)^{-}}{|y|^{d+\sigma}} dy$$

$$\leq \int_{\{y_{d}<-1/\beta\}} \frac{\Lambda(-\beta y_{d}-1)}{|y|^{d+\sigma}} dy = C_{2}\beta^{\sigma},$$

where C_2 depends on Λ, d , and σ , and is uniformly bounded as $\sigma \to 2$. Thanks to the estimates of I_1 and I_2 above, it follows that

$$\mathcal{M}^+\phi(e) \le C_1\beta(\beta-1) + C_2\beta^{\sigma}.$$

By choosing β sufficiently small depending on λ , Λ , d, and σ (but uniformly as $\sigma \to 2$) so that

$$C_1(\beta - 1) + C_2\beta^{\sigma - 1} \le -C_1/2,$$

the claim is proved.

Case 2: $\sigma < 1$. Let I_1 and I_2 be defined as before. Since $\phi(x) = 0$ for $x_d < 0$, we get

$$I_2 = -(2 - \sigma) \int_{\{y_d < -1\}} \frac{\lambda}{|y|^{d+\sigma}} dy = -C_3,$$
 (2.3)

where $C_3 > 0$ depends on σ, λ , and d. For I_1 , we have

$$I_{1} = (2 - \sigma) \int_{\{y_{d} > 0\}} \frac{\Lambda((1 + y_{d})^{\beta} - 1)}{|y|^{d + \sigma}} dy - (2 - \sigma) \int_{y_{d} \in (-1, 0)} \frac{\lambda(1 - (1 + y_{d})^{\beta})}{|y|^{d + \sigma}} dy$$

$$\leq (2 - \sigma) \Lambda \int_{\{y_{d} > 0\}} \frac{(1 + y_{d})^{\beta} - 1}{|y|^{d + \sigma}} dy \to 0 \quad \text{as} \quad \beta \to 0$$

by the monotone convergence theorem. Therefore, we can choose β small depending on Λ , λ , d, and σ so that

$$\mathcal{M}^+ \phi(e) \le -C_3/2.$$

The claim is proved.

Case 3: $\sigma = 1$. In this case, we still have (2.3). For I_1 , we notice that integrand in the region $\{-1 < y_d < 0\} \cup \{|y| < 1\}$ is negative, and

$$\int_{\{y_d > 0\} \cap \{|y| > 1\}} \frac{(1 + y_d)^{\beta} - 1}{|y|^{d + \sigma}} \, dy \to 0 \quad \text{as} \quad \beta \to 0$$

by the monotone convergence theorem. Thus the claim follows as well.

Now we are ready to consider u near the lateral boundary. By a translation and rotation of the coordinates, we replace the ball B_1 by $B_1(e)$ and estimate u near the origin. Define the barrier function $\psi(t,x) = \frac{C_0}{2^{\beta-\sigma}\hat{C}}\phi(x)$. Obviously,

$$\partial_t \psi(t,x) - \mathcal{M}^+ \psi(t,x) = -\frac{C_0}{2\beta - \sigma \hat{C}} \mathcal{M}^+ \phi(x) > \frac{C_0}{2\beta - \sigma} x_d^{\beta - \sigma} > C_0$$

when $x \in B_1(e)$. On the other hand, $\psi \geq 0$ in $\mathbb{R} \times \mathbb{R}^d$. Since

$$u_t - \mathcal{M}^+ u \le C_0$$

and $u \equiv 0$ outside Q_1 , by the comparison principle,

$$u(t,x) \le \psi(t,x) = \frac{C_0}{2^{\beta-\sigma}\hat{C}} x_d^{\beta}.$$

By considering -u instead of u, we have $u \ge -\frac{C_0}{2^{\beta-\sigma}\hat{C}}x_d^{\beta}$. Hence, around the origin $|u| \le C|x|^{\beta}$. By rotation of the coordinate, we obtain the estimate near the lateral boundary.

For the bottom, let $\tilde{\phi} = C_0(t+1)$ so that $\tilde{\phi}(-1) = 0$ and $\tilde{\phi}'(t) = C_0$. This yields that

$$\partial_t \tilde{\phi} - \mathcal{M}^+ \phi = C_0.$$

Moreover, $\tilde{\phi} \geq 0$ in $(-1,0) \times \mathbb{R}^d$. By the comparison principle again, $u \leq \phi$ in Q_1 . In particular, near the bottom $u \leq C_0(t+1)$, which further implies $|u| \leq C_0(t+1)$ by symmetry.

Combining the estimates of lateral boundary and bottom with the interior Hölder estimate, we prove the lemma. \Box

3. A LIOUVILLE THEOREM

The aim of this section is to prove the following Liouville theorem for the fully nonlinear parabolic nonlocal equation with non-symmetric kernels. The elliptic version for symmetric kernels was established in [26].

Theorem 3.1. Let $\sigma \in (0,2)$. There is a constant $\hat{\alpha} \in (0,1/2)$ depending on d, λ , Λ , and σ (but is uniform as $\sigma \to 2$) such that the following statement holds. Let $\alpha \in (0,\hat{\alpha})$ be such that $[\sigma + \hat{\alpha}] < \sigma + \alpha$ and suppose that $u \in C^{1+\frac{\alpha}{\sigma},\sigma+\alpha}_{loc}(\mathbb{R}^{d+1}_0)$ satisfies the following properties:

(i) For any $\beta \in [0, \sigma + \alpha]$ and $R \geq 1$, we have

$$[u]_{\beta/\sigma,\beta;Q_R} \le N_0 R^{\sigma+\alpha-\beta}; \tag{3.1}$$

(ii) For any $(s,h) \in \mathbb{R}_0^{d+1}$, we have

$$\partial_t (u(\cdot + s, \cdot + h) - u) - \mathcal{M}^- (u(\cdot + s, \cdot + h) - u) \ge 0, \tag{3.2}$$

$$\partial_t (u(\cdot + s, \cdot + h) - u) - \mathcal{M}^+ (u(\cdot + s, \cdot + h) - u) \le 0; \tag{3.3}$$

(iii) If $\sigma > 1$, for any nonnegative measure μ in \mathbb{R}^d with compact support,

$$\partial_t u_\mu - \mathcal{M}^+ u_\mu \le 0$$
, where $u_\mu(t, x) = \int_{\mathbb{R}^d} \delta u(t, x, h) \, d\mu(h)$.

Then u is a polynomial of degree ν in x and 1 in t, where ν is the integer part of $\sigma + \alpha$.

Remark 3.2. As in [26], it is possible to relax Condition (i) in Theorem 3.1 by assuming that (3.1) is satisfied for any $\beta \in [0, \sigma + \alpha']$ and $R \ge 1$, where $\alpha' \in (0, \alpha)$ is a constant satisfying $\sigma + \alpha' > \nu$. A simple computation reveals that in the case we also require that $\sigma > 1 + \alpha - \alpha'$ when $\sigma > 1$ and $\nu = 1$; and $\sigma > \alpha - \alpha'$ when $\sigma < 1$ and $\nu = 0$.

To prove Theorem 3.1, we first present a few lemmas. Define

$$P(t,x) = \int_{\mathbb{R}^d} \left(\delta u(t,x,y) - \delta u(0,0,y) \right)^+ \frac{2-\sigma}{|y|^{d+\sigma}} dy,$$

$$N(t,x) = \int_{\mathbb{R}^d} \left(\delta u(t,x,y) - \delta u(0,0,y) \right)^- \frac{2-\sigma}{|y|^{d+\sigma}} dy.$$

Lemma 3.3. Let $\hat{\alpha} \in (0, 1/2)$ be a constant satisfying $\hat{\alpha} < \sigma/2$. Under the conditions (i) and (ii) of Theorem 3.1, for any $\kappa \geq 2$ and $l \in \mathbb{N} \cup \{0\}$, we have

$$\sup_{Q_{\kappa^l}} \left(P + N + |u_t - u_t(0, 0)| \right) \le C N_0 \kappa^{\alpha l} \le C N_0 \kappa^{\hat{\alpha} l}$$

$$\tag{3.4}$$

and

$$[u_t]_{\gamma/\sigma,\gamma;Q_{1/2}} \le CN_0 \kappa^{\hat{\alpha}},\tag{3.5}$$

where C depends only on d, λ, Λ , and σ , and is uniformly bounded as $\sigma \to 2$.

Proof. We first estimate P and N assuming that $\nu = 2$. Fix $(t, x), (t', x') \in Q_1$ and set $l = |x - x'| + |t - t'|^{1/\sigma}$. By Condition (i), when |y| < l,

$$|\delta u(t, x, y) - \delta u(t', x', y)|$$

$$= \left| \int_0^1 y \left[Du(t, x + sy) - Du(t, x) - \left(Du(t', x' + sy) - Du(t', x') \right) \right] ds \right|$$

$$\leq C|y|^2 l^{\sigma + \alpha - 2} [u]_{1 + \alpha/\sigma, \sigma + \alpha; Q_2} \leq CN_0 |y|^2 l^{\sigma + \alpha - 2}. \tag{3.6}$$

Similarly, when $|y| \ge l$,

$$|\delta u(t, x, y) - \delta u(t', x', y)| \le C|y|^{\sigma + \alpha - 1} l[u]_{1 + \alpha/\sigma, \alpha + \sigma; Q_{1 + |y|}} \le CN_0 l|y|^{\sigma + \alpha - 1}. \quad (3.7)$$

Combining (3.6) and (3.7), we have

$$\int_{\mathbb{R}^{d}} |\delta u(t, x, y) - \delta u(t', x', y)| \frac{2 - \sigma}{|y|^{d + \sigma}} dy$$

$$\leq C N_{0} l^{\sigma + \alpha - 2} \int_{B_{l}} \frac{(2 - \sigma)|y|^{2}}{|y|^{d + \sigma}} dy + C N_{0} l \int_{\mathbb{R}^{d} \setminus B_{l}} \frac{(2 - \sigma)|y|^{\sigma + \alpha - 1}}{|y|^{d + \sigma}} dy$$

$$\leq C N_{0} l^{\alpha}. \tag{3.8}$$

Hence $P, N \in C^{\alpha,\alpha/\sigma}(Q_1)$. Because P(0,0) = N(0,0) = 0, we have

$$P(t,x) + N(t,x) \le CN_0$$
 in Q_1 .

By modifying the estimate above, we can prove the same estimate for P when $\nu=0$ or 1.

We then use a scaling argument. Define $\hat{u}(t,x) = \eta^{-\alpha-\sigma}u(\eta^{\sigma}t,\eta x)$ for any $\eta > 1$. It is easily seen that \hat{u} satisfies all the conditions in this lemma. Hence, we know that

$$\int_{\mathbb{R}^d} \frac{(2-\sigma) \left| \delta \hat{u}(t,x,y) - \delta \hat{u}(0,0,y) \right|}{|y|^{d+\sigma}} \, dy \le C N_0 \quad \text{in } Q_1.$$

Therefore,

$$P(\eta^{\sigma}t, \eta x) + N(\eta^{\sigma}t, \eta x) \le CN_0\eta^{\alpha}$$
 in Q_1 ,

which together with (3.2) and (3.3) implies (3.4).

To prove (3.5), we take h=0 in (3.2) and (3.3), and then multiply them by 1/s. By letting $s \to 0$, we know that u_t as well as $u_t - u_t(0,0)$ are sub and super-solutions at the same time. By Proposition 2.4, we obtain that $u_t \in C^{\gamma/\sigma,\gamma}(Q_{1/2})$ for some $\gamma > 0$ depending on d, λ, Λ , and σ_0 , and

$$[u_t]_{\gamma/\sigma,\gamma;Q_{1/2}} = [u_t - u_t(0,0)]_{\gamma/\sigma,\gamma;Q_{1/2}} \le C \sup_{t \in (-1,0)} \int_{\mathbb{R}^d} \frac{|u_t - u_t(0,0)|}{1 + |x|^{\sigma+d}} \, dx.$$

Using (3.4), for any $t \in (-1, 0)$,

$$\int_{\mathbb{R}^{d}} \frac{|u_{t} - u_{t}(0,0)|}{1 + |x|^{d+\sigma}} dx dt
\leq \int_{B_{1}} \frac{|u_{t} - u_{t}(0,0)|}{1 + |x|^{d+\sigma}} dx dt + \sum_{i=0}^{\infty} \int_{B_{\kappa^{i+1}} \setminus B_{\kappa^{i}}} \frac{|u_{t} - u_{t}(0,0)|}{1 + |x|^{d+\sigma}} dx dt
\leq CN_{0} \int_{B_{1}} \frac{1}{1 + |x|^{d+\sigma}} dx dt + CN_{0} \sum_{i=0}^{\infty} \int_{B_{\kappa^{i+1}} \setminus B_{\kappa^{i}}} \frac{\kappa^{\hat{\alpha}(i+1)}}{1 + |x|^{d+\sigma}} dx dt
\leq CN_{0} + CN_{0} \sum_{i=0}^{\infty} \kappa^{\hat{\alpha}(i+1)} \int_{\kappa^{i}}^{\kappa^{i+1}} \frac{r^{d-1}}{1 + r^{d+\sigma}} dr
\leq CN_{0} + CN_{0} \frac{\kappa^{\hat{\alpha}}(1 - \kappa^{-\sigma})}{1 - \kappa^{\hat{\alpha} - \sigma}} \leq CN_{0} \kappa^{\hat{\alpha}},$$

where C depends only on d, λ, Λ , and σ_0 . Here we used the fact $\hat{\alpha} < \sigma/2$ and $\kappa \ge 2$ in the last inequality. Therefore, the lemma is proved.

Dividing u by a CN_0 , where C is the constant in (3.4), and using Lemma 3.3, we have that for any $\kappa \geq 2$ and $l \in \mathbb{N} \cup \{0\}$,

$$\sup_{Q_{\kappa^l}} P \le \kappa^{\hat{\alpha}l}, \quad \sup_{Q_{\kappa^l}} N \le \kappa^{\hat{\alpha}l}. \tag{3.9}$$

We are going to prove inductively that there exists a sufficiently large $\kappa \geq 2$ and sufficiently small $\hat{\alpha} \in (0, 1/2)$ such that

$$\sup_{Q_{\kappa^{-l}}} P \leq \kappa^{-\hat{\alpha}l}, \quad \sup_{Q_{\kappa^{-l}}} N \leq \kappa^{-\hat{\alpha}l} \quad \text{for any } l \in \mathbb{N}.$$

For a fixed $r \in (0,1)$, assume that P attains its maximum in $\overline{Q_r}$ at (t_0,x_0) . Denote

$$\mathbb{A} := \{ y : \delta u(t_0, x_0, y) - \delta u(0, 0, y) > 0 \}.$$

Then

$$P(t_0, x_0) = \int_{\mathbb{A}} \left(\delta u(t_0, x_0, y) - \delta u(0, 0, y) \right) \frac{2 - \sigma}{|y|^{d + \sigma}} dy,$$

$$N(t_0, x_0) = \int_{\mathbb{R}^d \setminus \mathbb{A}} \left(\delta u(t_0, x_0, y) - \delta u(0, 0, y) \right) \frac{2 - \sigma}{|y|^{d + \sigma}} dy.$$

We define

$$v(t,x) = \int_{\mathbb{A}} \left(\delta u(t,x,y) - \delta u(0,0,y) \right) \frac{2-\sigma}{|y|^{d+\sigma}} \, dy.$$

Notice that $v \leq P$, and in particular $v \leq 1$ in Q_1 . Moreover, $P(t_0, x_0) = v(t_0, x_0)$. We denote $\mathfrak{v} = (1 - v)^+$.

Lemma 3.4. Suppose that $\hat{\alpha} \in (0, 1/2)$ satisfying $\hat{\alpha} < \sigma/2$ and $\kappa \geq 2$. Then we have

$$\mathfrak{v}_t - \mathcal{M}^- \mathfrak{v} \ge -C(\kappa^{\hat{\alpha}} - 1) \quad in \ Q_{3/4}, \tag{3.10}$$

where C is a positive constant depending only on d, λ , Λ , and σ , and is uniformly bounded as $\sigma \to 2$.

Proof. Since $v \leq 1$ in Q_1 , for any $(t,x) \in Q_1$ we have $\mathfrak{v}(t,x) = 1 - v(t,x)$, thus

$$\begin{split} \delta \mathfrak{v}(t,x,y) &= \mathfrak{v}(t,x+y) - \mathfrak{v}(t,x) - D \mathfrak{v}(t,x) y \\ &= (1-v)^+(t,x+y) - (1-v)(t,x) + D v(t,x) y \\ &= (v-1)^+(t,x+y) - \delta v(t,x,y). \end{split}$$

Therefore, we have

$$(\delta v(t, x, y))^{-} \ge (\delta v(t, x, y))^{+} - (v - 1)^{+}(t, x + y),$$

$$(\delta v(t, x, y))^{+} \le (\delta v(t, x, y))^{-} + (v - 1)^{+}(t, x + y).$$

These imply

$$\mathfrak{v}_{t} - \mathcal{M}^{-}\mathfrak{v} = -v_{t} - (2 - \sigma) \int_{\mathbb{R}^{d}} \frac{\lambda(\delta \mathfrak{v}(t, x, y))^{+} - \Lambda(\delta \mathfrak{v}(t, x, y))^{-}}{|y|^{d+\sigma}} dy$$

$$\geq -v_{t} + \mathcal{M}^{+}v - (2 - \sigma)(\Lambda + \lambda) \int_{\mathbb{R}^{d}} \frac{(v - 1)^{+}(t, x + y)}{|y|^{d+\sigma}} dy. \tag{3.11}$$

From Condition (iii) and an approximation (see, for instance, the proof of Lemma 4.4 below), v satisfies

$$v_t - \mathcal{M}^+ v \le 0. \tag{3.12}$$

On the other hand, we have

$$\int_{\mathbb{R}^d} \frac{(v-1)^+(t,x+y)}{|y|^{d+\sigma}} \, dy \le \int_{\mathbb{R}^d} \frac{(P-1)^+(t,x+y)}{|y|^{d+\sigma}} \, dy.$$

Since P satisfies (3.9), for $(t, x) \in Q_{3/4}$, we have $P(t, x + y) \le 1$ when $y \in B_{1/4}$, and thus the right-hand side above is equal to

$$\sum_{i=0}^{\infty} \int_{B_{\kappa^{i+1}-3/4} \setminus B_{\kappa^{i}-3/4}} \frac{(P-1)^{+}(t,x+y)}{|y|^{d+\sigma}} \, dy,$$

which by (3.9) is bounded by

$$\begin{split} &\sum_{i=0}^{\infty} \int_{B_{\kappa^{i+1}-3/4} \backslash B_{\kappa^{i}-3/4}} \frac{\kappa^{(i+1)\hat{\alpha}} - 1}{|y|^{d+\sigma}} \, dy \\ &\leq C \sum_{i=0}^{\infty} (\kappa^{(i+1)\hat{\alpha}} - 1) \kappa^{-i\sigma} = C \Big(\frac{\kappa^{\hat{\alpha}}}{1 - \kappa^{\hat{\alpha}-\sigma}} - \frac{1}{1 - \kappa^{-\sigma}} \Big) \leq C \Big(\kappa^{\hat{\alpha}} - 1 \Big), \end{split} \tag{3.13}$$

where C only depends on d, λ, Λ , and σ_0 . Here we used the fact $\hat{\alpha} < \sigma/2$ and $\kappa \ge 2$ in the last inequality. Combining (3.12)-(3.13) with (3.11), we prove the lemma.

Let $\hat{\theta} = \lambda/(4\Lambda)$ and γ be the constant in Proposition 2.4. For any $r_1 > 0$, define the set

$$\mathcal{D}_{r_1} = \{ (t, x) \in Q_{r_1} : v \ge 1 - \hat{\theta} \}.$$

Lemma 3.5. Suppose that $\hat{\alpha} \in (0, 1/2)$ satisfying $\hat{\alpha} < \sigma/2$ and $\kappa \ge 2$. There exist some $\eta \in (0, 1)$ sufficiently close to 1 and $c \in (0, 1)$ sufficiently small, both depending only on d, λ , Λ , and σ (and is uniform as $\sigma \to 2$), such that for $r_1 = c\kappa^{-\hat{\alpha}/\gamma}$,

$$|\mathcal{D}_{r_1}| < \eta |Q_{r_1}|. \tag{3.14}$$

Proof. By contradiction we assume that $|\mathcal{D}_{r_1}| > \eta |Q_{r_1}|$, and consider

$$w := \int_{\mathbb{R}^{d \setminus \Delta}} \left(\delta u(t, x, y) - \delta u(0, 0, y) \right) \frac{2 - \sigma}{|y|^{d + \sigma}} \, dy.$$

By Condition (iii) and an approximation argument, w is a subsolution, i.e.,

$$w_t - \mathcal{M}^+ w \le 0 \quad \text{in} \quad \mathbb{R}_0^{d+1}. \tag{3.15}$$

From (3.2) and (3.3), we know that in \mathbb{R}_0^{d+1} ,

$$\frac{\lambda}{\Lambda}P - \frac{u_t - u_t(0,0)}{\Lambda} \le N \le \frac{\Lambda}{\lambda}P - \frac{u_t - u_t(0,0)}{\lambda},\tag{3.16}$$

implying that in $Q_{1/2}$

$$\frac{\lambda}{\Lambda} P - \frac{[u_t]_{\gamma/\sigma,\gamma;Q_{1/2}}}{\Lambda} \left(|x|^{\gamma} + |t|^{\gamma/\sigma} \right) \le N \le \frac{\Lambda}{\lambda} P + \frac{[u_t]_{\gamma/\sigma,\gamma;Q_{1/2}}}{\lambda} \left(|x|^{\gamma} + |t|^{\gamma/\sigma} \right).$$

From (3.5), (3.9), and the above inequality, we obtain

$$w = P - v - N \le (1 - \lambda/\Lambda)P - v + C\kappa^{\hat{\alpha}}(|x|^{\gamma} + |t|^{\gamma/\sigma})$$

$$\le 1 - \lambda/\Lambda - (1 - \hat{\theta}) + C\kappa^{\hat{\alpha}}(|x|^{\gamma} + |t|^{\gamma/\sigma})$$

$$\le -\lambda/\Lambda + \hat{\theta} + C\kappa^{\hat{\alpha}}(|x|^{\gamma} + |t|^{\gamma/\sigma}) \quad \text{in } \mathcal{D}_{r_1},$$

where C only depends on d, λ, Λ , and σ , and is uniformly bounded as $\sigma \to 2$. Now we choose c sufficiently small depending only on d, λ, Λ , and σ (uniformly as $\sigma \to 2$), such that

$$-\lambda/\Lambda + \hat{\theta} + C\kappa^{\hat{\alpha}} (|x|^{\gamma} + |t|^{\gamma/\sigma})$$

$$\leq -3\hat{\theta} + C\kappa^{\hat{\alpha}} (c\kappa^{-\hat{\alpha}/\gamma})^{\gamma} \leq -3\hat{\theta} + Cc^{\gamma} \leq -\hat{\theta} \quad \text{in } Q_{r_1}, \tag{3.17}$$

which implies $w \leq -\hat{\theta}$ in \mathcal{D}_{r_1} . Since w is a subsolution (3.15), it follows immediately that for any $\varepsilon \in (0, r_1)$, $\bar{w}_{\varepsilon}(t, x) := (w + \hat{\theta})^+(\varepsilon^{\sigma}t, \varepsilon x)$ is a subsolution as well. Moreover,

$$\left| \left\{ \bar{w}_{\varepsilon} \le 0 \right\} \cap Q_{r_1/\varepsilon} \right| > \eta \left| Q_{r_1/\varepsilon} \right|. \tag{3.18}$$

We estimate \bar{w}_{ε} by applying Proposition 2.3 with $t_1 = -1$, $t_2 = 0$, and $\Omega = \mathbb{R}^d$

$$\bar{w}_{\varepsilon}(0,0) \leq C \int_{-1}^{0} \int_{\mathbb{R}^{d}} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt
= C \int_{-1}^{0} \int_{B_{r_{1}/\varepsilon}} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt + C \int_{-1}^{0} \int_{B_{r_{1}/\varepsilon}^{c}} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt.$$
(3.19)

We first consider the second term on the right-hand side of the inequality above

$$\int_{-1}^{0} \int_{B_{r_{1}/\varepsilon}^{c}} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt$$

$$\leq \int_{-1}^{0} \int_{B_{r_{1}/\varepsilon}^{c}} \frac{\hat{\theta}}{1 + |x|^{d+\sigma}} dx dt + \int_{-1}^{0} \int_{B_{r_{1}/\varepsilon}^{c}} \frac{|w|(\varepsilon^{\sigma}t, \varepsilon x)}{1 + |x|^{d+\sigma}} dx dt$$

$$\leq C \hat{\theta}(\varepsilon/r_{1})^{\sigma} + \int_{-\varepsilon^{\sigma}}^{0} \int_{B_{\varepsilon}^{c}} \frac{|w|(t, x)}{\varepsilon^{d+\sigma} + |x|^{d+\sigma}} dx dt. \tag{3.20}$$

Since $|w| \leq \max\{P, N\}$, from (3.9), for any $l \geq 0$,

$$\sup_{Q_{wl}} |w| \le \kappa^{\hat{\alpha}l}.$$

Therefore,

$$\int_{-\varepsilon^{\sigma}}^{0} \int_{B_{r_{1}}^{c}} \frac{|w|(t,x)}{\varepsilon^{d+\sigma} + |x|^{d+\sigma}} dx dt
\leq \int_{-\varepsilon^{\sigma}}^{0} \int_{B_{1} \setminus B_{r_{1}}} \frac{|w|}{\varepsilon^{d+\sigma} + |x|^{d+\sigma}} dx dt + \sum_{i=0}^{\infty} \int_{-\varepsilon^{\sigma}}^{0} \int_{B_{\kappa^{i+1}} \setminus B_{\kappa^{i}}} \frac{|w|}{|x|^{d+\sigma}} dx dt
\leq C((\varepsilon/r_{1})^{\sigma} + \varepsilon^{\sigma} \kappa^{\hat{\alpha}}),$$
(3.21)

where C only depends on d, λ, Λ , and σ , and is uniformly bounded as $\sigma \to 2$. We combine (3.20) and (3.21) to obtain that

$$C \int_{-1}^{0} \int_{B_{r_1/\varepsilon}^{c}} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt \le C((\varepsilon/r_1)^{\sigma} + \varepsilon^{\sigma} \kappa^{\hat{\alpha}}).$$

Recall that $r_1 = c\kappa^{-\hat{\alpha}/\gamma}$. For the right-hand side of the inequality above, we want to choose ε sufficiently small such that $C\varepsilon^{\sigma}(r_1^{-\sigma} + \kappa^{\hat{\alpha}}) \leq \hat{\theta}/4$. Indeed, since $c \in (0,1)$

and $\sigma > \gamma$, we have $r_1^{-\sigma} \ge \kappa^{\hat{\alpha}}$. It is sufficient to fix ε such that $2C\varepsilon^{\sigma}r_1^{-\sigma} = \hat{\theta}/4$, i.e.,

$$\varepsilon/r_1 = \left(\hat{\theta}/(8C)\right)^{1/\sigma} := c_1,$$

where c_1 only depends on d, λ, Λ , and σ , and is uniformly bounded as $\sigma \to 2$. In other words, by taking $\varepsilon = c_1 r_1$ we have

$$C \int_{-1}^{0} \int_{B_{r_1/\varepsilon}^c} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt \le \hat{\theta}/4.$$

$$(3.22)$$

Next we estimate the first term on the right-hand side of (3.19) using (3.18):

$$C \int_{-1}^{0} \int_{B_{1/c_{1}}} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt$$

$$= C \int_{\left((-1,0) \times B_{1/c_{1}}\right) \cap \{\bar{w}_{\varepsilon} > 0\}} \frac{\bar{w}_{\varepsilon}}{1 + |x|^{d+\sigma}} dx dt$$

$$\leq C(1 - \eta) |Q_{1/c_{1}}| \sup_{(-1,0) \times B_{1/c_{1}}} \bar{w}_{\varepsilon}$$

$$\leq C(1 - \eta) |Q_{1/c_{1}}| \sup_{(-1,0) \times B_{r_{1}}} (w + \hat{\theta})^{+} \leq \hat{\theta}/4$$
(3.23)

upon taking η sufficiently close to 1 depending only on d, λ, Λ , and σ (uniformly as $\sigma \to 2$).

Combining (3.23) and (3.22) with (3.19), we have $\bar{w}_{\varepsilon}(0,0) \leq \hat{\theta}/2$ indicating that $w(0,0) \leq -\hat{\theta}/2$, which contradicts with w(0,0) = 0 by the definition of w. Therefore, the lemma is proved.

Now we are ready to prove Theorem 3.1.

Proof of Theorem 3.1. Below we first elaborate on the case when $\sigma > 1$. At the end, we briefly discuss the case when $\sigma = 1$. The proof of the case when $\sigma < 1$ is omitted due to similarity to the first case.

Let η and c be the constants in Lemma 3.5 and $r_1 = c\kappa^{-\hat{\alpha}/\gamma}$. From (3.14),

$$|\{1-v \ge \hat{\theta}\} \cap Q_{r_1}| \ge (1-\eta)|Q_{r_1}|.$$

Recall that $\tilde{Q}_{\delta r_1} = (-r_1^{\sigma}, -(\delta r_1)^{\sigma}) \times B_{r_1}$. For δ sufficiently small depending on η , we have

$$\left|\left\{1-v\geq\hat{\theta}\right\}\cap\tilde{Q}_{\delta r_1}\right|\geq\frac{1-\eta}{2}\left|\tilde{Q}_{\delta r_1}\right|.$$

We set $r = \delta r_1/2$ and apply the weak Harnack inequality Corollary 2.2 to $(1-v)^+$ in $Q_{3/4}$ with (3.10) to obtain

$$\inf_{Q_{r}} (1 - v) + C(\kappa^{\hat{\alpha}} - 1) r_{1}^{\sigma}
\geq C(\delta) \| (1 - v)^{+} \|_{L_{\varepsilon_{1}}(\tilde{Q}_{\delta r_{1}})} r_{1}^{-(d + \sigma)/\varepsilon_{1}}
\geq C(\delta) \hat{\theta} ((1 - \eta) |\tilde{Q}_{\delta r_{1}}|/2)^{1/\varepsilon_{1}} r_{1}^{-(d + \sigma)/\varepsilon_{1}}
= C(\delta) \hat{\theta} ((1 - \eta) (1 - \delta^{\sigma})/2)^{1/\varepsilon_{1}} =: 2\theta,$$
(3.24)

where θ is a small constant depending only on d, λ , Λ , and σ , and is uniformly bounded as $\sigma \to 2$. We fix $\kappa = 4/(c\delta)^2$, where c is chosen according to (3.17), which guarantees that

$$\kappa^{-1} < c\delta/2 \cdot \kappa^{-\hat{\alpha}/\gamma}$$

for any $\hat{\alpha} \in (0, \gamma/2)$. Since $r = \delta r_1/2$ and $r_1 = c\kappa^{-\hat{\alpha}/\gamma}$, the inequality above can be written as $\kappa^{-1} < r$.

Next, we choose $\hat{\alpha}_1 = \log(1 + \theta/C)/\log \kappa$ such that for any $\hat{\alpha} \in (0, \hat{\alpha}_1)$,

$$C(\kappa^{\hat{\alpha}} - 1)r_1^{\sigma} < C(\kappa^{\hat{\alpha}} - 1) < \theta.$$

Therefore, from (3.24) and the inequality above we obtain that $\sup_{Q_r} v \leq 1 - \theta$. Since $\kappa^{-1} < r$,

$$\sup_{Q_{\kappa^{-1}}} P \le \sup_{Q_r} P = P(t_0, x_0) = v(t_0, x_0) = \sup_{Q_r} v \le 1 - \theta.$$
 (3.25)

Similarly,

$$\sup_{Q_{\kappa^{-1}}} N \le 1 - \theta. \tag{3.26}$$

Set

$$\hat{\alpha} = \min \{ -\log(1-\theta)/\log \kappa, \hat{\alpha}_1, \gamma/2 \}.$$

Then (3.25) and (3.26) imply

$$\sup_{Q_{\kappa^{-1}}} \kappa^{\hat{\alpha}} P \leq 1, \quad \sup_{Q_{\kappa^{-1}}} \kappa^{\hat{\alpha}} N \leq 1. \tag{3.27}$$

Let

$$\tilde{P}(t,x) = \kappa^{\hat{\alpha}} P(\kappa^{-1} x, \kappa^{-\sigma} t), \quad \tilde{N}(t,x) = \kappa^{\hat{\alpha}} N(\kappa^{-1} x, \kappa^{-\sigma} t),$$

and

$$\tilde{u}(t,x) = \kappa^{\sigma + \hat{\alpha}} u(\kappa^{-1} x, \kappa^{-\sigma} t).$$

From (3.16), we have

$$\frac{\lambda}{\Lambda}\tilde{P} - \frac{\tilde{u}_t - \tilde{u}_t(0,0)}{\Lambda} \le \tilde{N} \le \frac{\Lambda}{\lambda}\tilde{P} - \frac{\tilde{u}_t - \tilde{u}_t(0,0)}{\lambda}.$$

Since $\kappa \geq 2$ and $\hat{\alpha} \leq \gamma$, we get

$$[\tilde{u}_t]_{\gamma/\sigma,\gamma;Q_{1/2}} \le \kappa^{\hat{\alpha}-\gamma} [u_t]_{\gamma/\sigma,\gamma;Q_{1/2}} \le [u_t]_{\gamma/\sigma,\gamma;Q_{1/2}}.$$

On the other hand, for any $l \geq 0$,

$$\sup_{Q_{\kappa^l}} \tilde{P} = \kappa^{\hat{\alpha}} \sup_{Q_{\kappa^{l-1}}} P \leq \kappa^{\hat{\alpha}} \kappa^{(l-1)\hat{\alpha}} = \kappa^{l\hat{\alpha}}, \quad \sup_{Q_{\kappa^l}} \tilde{N} \leq \kappa^{l\hat{\alpha}}.$$

Therefore, \tilde{P} and \tilde{N} satisfy all the conditions that P and N satisfied. Applying (3.27) to \tilde{P} and \tilde{N} , we have

$$\sup_{Q_{\kappa^{-1}}} \tilde{P} \leq \kappa^{-\hat{\alpha}}, \quad \sup_{Q_{\kappa^{-1}}} \tilde{N} \leq \kappa^{-\hat{\alpha}},$$

which further implies that

$$\sup_{Q_{\kappa^{-2}}} P \leq \kappa^{-2\hat{\alpha}}, \quad \sup_{Q_{\kappa^{-2}}} N \leq \kappa^{-2\hat{\alpha}}.$$

By induction, for any $l \in \mathbb{N}$,

$$\sup_{Q_{\kappa^{-l}}} P \leq \kappa^{-l\hat{\alpha}}, \quad \sup_{Q_{\kappa^{-l}}} N \leq \kappa^{-l\hat{\alpha}}.$$

Therefore, we have in Q_1

$$P(t,x) \le C(|x|^{\hat{\alpha}} + |t|^{\hat{\alpha}/\sigma}), \quad N(t,x) \le C(|x|^{\hat{\alpha}} + |t|^{\hat{\alpha}/\sigma}).$$

Since for any $\eta \geq 1$, $\hat{u}(t,x) = \eta^{-\sigma-\alpha}u(\eta^{\sigma}t,\eta x)$ satisfies the same condition as u, replacing u by \hat{u} in the definition of P and denoting it as $P_{\hat{u}}$, we obtain

$$P_{\hat{u}}(t,x) \le C(|x|^{\hat{\alpha}} + |t|^{\hat{\alpha}/\sigma})$$
 in Q_1 .

Returning to P, we have

$$\eta^{-\alpha}P(\eta^{\sigma}t,\eta x) \le C(|x|^{\hat{\alpha}} + |t|^{\hat{\alpha}/\sigma})$$

in Q_1 , which further implies that

$$\sup_{Q_n} \frac{P(t,x)}{|x|^{\hat{\alpha}} + |t|^{\hat{\alpha}/\sigma}} \le C\eta^{\alpha - \hat{\alpha}}.$$

Let $\eta \to \infty$ yields

$$\sup_{(t,x)\in\mathbb{R}_0^{d+1}} \frac{P(t,x)}{|x|^{\hat{\alpha}} + |t|^{\hat{\alpha}/\sigma}} = 0,$$

which gives P = 0. Similarly, N = 0.

From the definition of P and N, we have

$$u(t, x + y) - u(t, x) - Du(t, x)y = u(0, y) - u(0, 0) - Du(0, 0)y.$$

Taking derivative in y, we have, for any $t \in (-\infty, 0)$ and $x, y \in \mathbb{R}^d$

$$Du(t, x + y) - Du(t, x) = Du(0, y) - Du(0, 0),$$

which implies for fixed t, u is a polynomial in x of order at most two. Using Condition (i) with $\beta=0$, we infer that this order is at most ν . Condition (ii), together with P=N=0, yields $u_t=c$ for some constant c. The proof is completed for $\sigma>1$.

Finally, we sketch the proof for $\sigma=1$. From Condition (ii), we know that $u(\cdot+s,\cdot+h)-u$, and thus Du and u_t are both sub and supersolutions, and are in $C^{\gamma/\sigma,\gamma}(Q_{1/2})$. By Proposition 2.4, we have

$$[u_t]_{\gamma/\sigma,\gamma;Q_{1/2}} + [Du]_{\gamma/\sigma,\gamma;Q_{1/2}} \le C \sup_{t \in [0,1]} \int_{\mathbb{R}^d} \frac{|Du| + |u_t|}{1 + |x|^{d+1}} \, dx.$$

From Condition (i), the right-hand side of the inequality above is less than

$$\int_{B_1} \frac{1}{1+|x|^{1+d}} \, dx + \sum_{j=1}^{\infty} \int_{B_{2^j} \setminus B_{2^{j-1}}} \frac{C2^{j\alpha}}{1+|x|^{1+d}} \, dx \le \frac{C}{1-2^{\alpha-1}}$$

for any $\alpha \in (0,1)$. By taking $\hat{\alpha} \leq \gamma$ and scaling as before, we can prove that

$$[u]_{1+\gamma,1+\gamma;Q_R} \le CR^{\alpha-\gamma},$$

i.e., u must be a linear function by sending $R \to \infty$. The theorem is proved. \square

4. Schauder estimate for nonlocal parabolic equations

In this section, we prove Theorem 1.1 by applying the Liouville theorem, a blow-up analysis, and a localization procedure. In the rest of the paper, we do not specify the domain associated with the norm when it is $\mathbb{R}_0^{d+1} = (-\infty, 0) \times \mathbb{R}^d$.

4.1. Equations with translation invariant kernels. In this subsection, we consider equations with translation invariant kernels, i.e., K = K(y). The main result of the subsection is the following theorem.

Theorem 4.1. Let $\sigma \in (0,2)$ be a constant and A be an index set. There exists a constant $\hat{\alpha} > 0$ depending on d, λ , Λ , and σ (uniformly as $\sigma \to 2$), such that given $0 < \alpha' < \alpha < \hat{\alpha}$ satisfying $[\sigma + \alpha] < \sigma + \alpha' < \sigma + \alpha$ the following holds. Let $u \in C^{1+\alpha'/\sigma,\alpha'+\sigma}((-1,0) \times \mathbb{R}^d) \cap C^{1+\alpha/\sigma,\alpha+\sigma}(Q_1)$ satisfy

$$u_t = \inf_{a \in \mathcal{A}} (L_a u + f_a)$$
 in Q_1 ,

where $L_a \in \mathcal{L}_0(\sigma, \lambda, \Lambda)$ with $K_a = K_a(y)$ for any $a \in \mathcal{A}$. Assume that

$$\sup_{(t,x)\in Q_1} \left| \inf_{a\in \mathcal{A}} f_a(t,x) \right| < \infty.$$

Then

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{1/2}} \leq C[u]_{1+\alpha'/\sigma,\alpha'+\sigma;(-1,0)\times\mathbb{R}^d} + C\sup_{a\in\mathcal{A}}[f_a]_{\alpha/\sigma,\alpha;Q_1},$$

where C only depends on $d, \lambda, \Lambda, \sigma, \alpha$, and α' , and is uniformly bounded as $\sigma \to 2$.

We denote

$$Q^{k} = \left(-1 + 2^{-(k+1)\sigma}/(1 - 2^{-\sigma}), 0\right) \times B_{1-2^{-k}}(0)$$
(4.1)

for all sufficiently large integers k such that $2^{-(k+1)\sigma} < 1 - 2^{-\sigma}$. We shall prove a stronger result:

$$\sup_{k} 2^{-k(\alpha - \alpha')} [u]_{1+\alpha/\sigma, \alpha+\sigma; Q^{k}}
\leq C[u]_{1+\alpha'/\sigma, \alpha'+\sigma; (-1,0) \times \mathbb{R}^{d}} + C \sup_{a \in \mathcal{A}} [f_{a}]_{\alpha/\sigma, \alpha; Q_{1}}.$$
(4.2)

The conclusion of the theorem is a particular case for k large only depending on σ (uniformly as $\sigma \to 2$) so that $Q_{1/2} \subset Q^k$. Since we assume that $u \in C^{1+\alpha/\sigma,\alpha+\sigma}(Q_1)$, there exists an integer k such that

$$2^{-k(\alpha-\alpha')}[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^k} = \sup_l 2^{-l(\alpha-\alpha')}[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^l}.$$

Next, we prove (4.2) by contradiction. Assume that we can find solutions u_j and index sets A_j such that

$$\begin{split} \partial_t u_j &= \inf_{a \in \mathcal{A}_j} (L_a u_j + f_a) \quad \text{in} \quad Q_1, \quad \sup_{(t,x) \in Q_1} \left| \inf_{a \in \mathcal{A}_j} f_a(t,x) \right| < \infty, \\ [u_j]_{1+\alpha'/\sigma,\sigma+\alpha';(-1,0) \times \mathbb{R}^d} &+ \sup_{a \in \mathcal{A}_j} [f_a]_{\alpha/\sigma,\alpha;Q_1} \le 1, \\ \text{and} \quad \sup_k 2^{-k(\alpha-\alpha')} [u_j]_{1+\alpha/\sigma,\sigma+\alpha;Q^k} \ge j, \end{split} \tag{4.3}$$

where for any $a \in \mathcal{A}_k$, $L_a \in \mathcal{L}_0$ with $K_a = K_a(y)$. As explained above, for each j there exists an integer k_j so that

$$2^{-k_j(\alpha-\alpha')}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}} = \sup_k 2^{-k(\alpha-\alpha')}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^k}.$$

Lemma 4.2. For any $j \ge 1$, we have

$$[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}} \leq \sup_{r>0} \sup_{(t,x)\in Q^{k_j}} r^{-(\alpha-\alpha')} [u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_r(t,x)\cap\{t>-1\}}$$

$$\leq 4^{\alpha-\alpha'} [u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}. \tag{4.4}$$

Moreover, we can find $(t_j, x_j) \in Q^{k_j}$ and r_j such that

$$\frac{1}{2}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}} \le r_j^{-(\alpha-\alpha')}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_{r_j}(t_j,x_j)\cap\{t>-1\}}$$
(4.5)

and

$$2^{k_j}r_j \to 0 \quad as \ j \to \infty. \tag{4.6}$$

Proof. The first inequality in (4.4) follows from the fact that for any $(t,x), (s,y) \in Q^{k_j}$ with $t \geq s$, we have $(s,y) \in Q_r(t,x) \cap \{t > -1\}$, where $r = \max(|x-y|, |t-s|^{1/\sigma})$. See, for instance, Claim 3.2 of [26]. For the second inequality, if $r \leq 2^{-(k_j+1)}$, for any $(t,x) \in Q^{k_j}$, we have $Q_r(t,x) \subset Q^{k_j+1}$ and

$$r^{-(\alpha-\alpha')}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_r(t,x)\cap\{t>-1\}}$$

$$\leq 2^{\alpha-\alpha'}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j+1}} \leq 4^{\alpha-\alpha'}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}},$$

where the last inequality is due to the choice of k_j . On the other hand, if $r > 2^{-(k_j+1)}$, for any $(t,x) \in Q^{k_j}$,

$$r^{-(\alpha-\alpha')}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_r(t,x)\cap\{t>-1\}} \\ \leq 2^{(k_j+1)(\alpha-\alpha')}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;(-1,0)\times\mathbb{R}^d} \leq 2^{\alpha-\alpha'}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}},$$

where the last inequality follows from (4.3). Thus, we obtain the second inequality in (4.4).

Due to (4.4), we can find $(t_j, x_j) \in Q^{k_j}$ and r_j such that (4.5) is satisfied and thus by (4.3),

$$(2^{k_j} r_j)^{\alpha - \alpha'} \le \frac{2[u_j]_{1 + \alpha'/\sigma, \alpha' + \sigma; Q_{r_j}(t_j, x_j) \cap \{t > -1\}}}{2^{-k_j(\alpha - \alpha')} [u_j]_{1 + \alpha/\sigma, \alpha + \sigma; Q^{k_j}}} \to 0 \quad \text{as } j \to \infty,$$

which further implies (4.6). The lemma is proved.

Let T_j be the Taylor expansion of u_j at $X_j = (t_j, x_j)$ of order $\nu = [\sigma + \alpha]$ in x and 1 in t. Now we consider the blow-up sequence

$$v_{j}(t,x) = \frac{u_{j}(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) - T_{j}(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x)}{r_{j}^{\sigma+\alpha}[u_{j}]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_{j}}}}.$$

Here (t_j, x_j) and r_j are from Lemma 4.2. Note that v_j is well defined on $(-R_j^{\sigma}, 0) \times \mathbb{R}^d$, where by Lemma 4.2,

$$R_j := 2^{-(k_j+1)} r_j^{-1} \to \infty \quad \text{as} \quad j \to \infty.$$

Observe that from (4.5) and (4.6), for sufficiently large j such that $r_j \leq 2^{-(k_j+1)}$,

$$[v_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_1} = \frac{r_j^{\sigma+\alpha'}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_{r_j}(t_j,x_j)}}{r_j^{\sigma+\alpha}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_{r_j}(t_j,x_j)}} \ge 1/2.$$
(4.7)

Lemma 4.3. For any R > 0 and $\beta \in [0, \sigma + \alpha']$, we have

$$[v_j]_{\beta/\sigma,\beta;Q_R\cap\{t>-R_i^\sigma\}} \le CR^{\sigma+\alpha-\beta},\tag{4.8}$$

where C depends only on α and α' . Moreover, for any $0 < R < R_j$ and $\beta \in [0, \sigma + \alpha]$, we have

$$[v_i]_{\beta/\sigma,\beta:Q_R} \le CR^{\sigma+\alpha-\beta},$$
 (4.9)

where C depends only on α and α' . Thus, we can find $v \in C^{1+\alpha/\sigma,\sigma+\alpha}(\mathbb{R}_0^{d+1})$ such that v satisfies (4.9) for any R > 0 and $\beta \in [0, \sigma + \alpha]$, and along a subsequence $v_j \to v$ in $C^{\beta/\sigma,\beta}$ locally uniformly for any $\beta \in [0, \sigma + \alpha)$.

We remark that (4.8) will be used below to prove that v satisfies Condition (iii) in Theorem 3.1, and (4.9) will be used to show that v satisfies Condition (i).

Proof of Lemma 4.3. For any R > 0 and $\beta \in [0, \sigma + \alpha']$,

$$\begin{split} &[v_j]_{\beta/\sigma,\beta;Q_R\cap\{t>-R_j^\sigma\}} \\ &= \frac{[u_j(t_j+r_j^\sigma\cdot,x_j+r_j\cdot)-T_j(t_j+r_j^\sigma\cdot,x_j+r_j\cdot)]_{\beta/\sigma,\beta;Q_R\cap\{t>-R_j^\sigma\}}}{r_j^{\sigma+\alpha}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}} \\ &\leq \frac{r_j^\beta[u_j-T_j]_{\beta/\sigma,\beta;Q_{Rr_j}(t_j,x_j)\cap\{t>-1\}}}{r_j^{\sigma+\alpha}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}} \\ &\leq \frac{r_j^\beta(Rr_j)^{\sigma+\alpha'-\beta}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_{Rr_j}(t_j,x_j)\cap\{t>-1\}}}{r_j^{\sigma+\alpha}[u_j]_{1+\alpha'/\sigma,\alpha'+\sigma;Q^{k_j}}} \leq CR^{\sigma+\alpha-\beta}, \end{split}$$

where we used (4.4) in the last inequality.

For any $R < R_j$, by the choice of k_j we have

$$\begin{split} &[v_j]_{1+\alpha/\sigma,\alpha+\sigma;Q_R} = \frac{[u_j(t_j + r_j^\sigma \cdot, x_j + r_j \cdot)]_{1+\alpha/\sigma,\alpha+\sigma;Q_R}}{r_j^{\sigma+\alpha}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}} \\ &= \frac{[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q_{Rr_j}(t_j,x_j)}}{[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}} \leq \frac{[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j+1}}}{[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}} \leq 2^{\alpha-\alpha'}. \end{split}$$

Using the interpolation inequality, we reach (4.9). The last statement of the lemma follows from the Arzela-Ascoli theorem and the Cauchy diagonal method. \Box

Lemma 4.4. The function v in Lemma 4.3 satisfies the conditions in Theorem 3.1.

Proof. By Lemma 4.3, Condition (i) is satisfied. Next we verify Condition (iii) for $\sigma \in (1,2)$. For any measure μ with compact support and $\delta \in (0,1)$, we define

$$V_j(t,x) = \int_{\mathbb{R}^d} v_j(t,x+h) - v_j(t,x) - \frac{v_j(t,x) - v_j(t,x-\delta h)}{\delta} d\mu(h).$$

Since T_j is linear in t, from the definition of v_j , we have

$$\begin{split} \partial_t V_j(t,x) &= \frac{r_j^{-\alpha}}{[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}} \\ &\cdot \int_{\mathbb{R}^d} \left[\partial_t u_j(t_j + r_j^{\sigma}t, x_j + r_j(x+h)) - \partial_t u_j(t_j + r_j^{\sigma}t, x_j + r_jx) \right. \\ &\left. - \frac{\partial_t u_j(t_j + r_j^{\sigma}t, x_j + r_jx) - \partial_t u_j(t_j + r_j^{\sigma}t, x_j + r_j(x-\delta h))}{\delta} \right] d\mu(h), \end{split}$$

which is equal to

$$\frac{r_{j}^{-\alpha}}{[u_{j}]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_{j}}}} \left[\int_{\mathbb{R}^{d}} \partial_{t} u_{j}(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}(x+h)) d\mu(h) \right]
+ \int_{\mathbb{R}^{d}} \frac{\partial_{t} u_{j}(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}(x-\delta h))}{\delta} d\mu(h)
- (1+1/\delta) \|\mu\|_{L_{1}} \partial_{t} u_{j}(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) \right].$$
(4.10)

For any $a \in \mathcal{A}_j$, define $\hat{K}_a(y) = r_j^{d+\sigma} K_a(r_j y)$, which satisfies

$$\frac{\lambda(2-\sigma)}{|y|^{d+\sigma}} \le \hat{K}_a(y) \le \frac{\Lambda(2-\sigma)}{|y|^{d+\sigma}},$$

and \hat{L}_a be the corresponding operator with kernel \hat{K}_a . Clearly,

$$\begin{split} \hat{L}_{a}V_{j} &= \int_{\mathbb{R}^{d}} \left[\hat{L}_{a}(v_{j}(t,x+h) - v_{j}(t,x)) - \frac{\hat{L}_{a}(v_{j}(t,x) - v_{j}(t,x-\delta h))}{\delta} \right] d\mu(h) \\ &= \frac{r_{j}^{-\alpha}}{\left[u_{j}\right]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_{j}}}} \int_{\mathbb{R}^{d}} \left\{ (L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x + r_{j}h) - (L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) - \frac{(L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) - (L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) - (L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) - (L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}(x - \delta h))}{\delta} \right\} d\mu(h), \end{split}$$

where in the second equality, we used the definitions of \hat{L}_a , v_j and the fact that T_j is at most second-order in x variable, so that for $\sigma > 1$ and any $y \in \mathbb{R}^d$

$$\delta T_j(t, x + h, y) - \delta T_j(t, x, y) = 0.$$

Therefore, for any $(t, x) \in (-R_i^{\sigma}, 0) \times \mathbb{R}^d$,

$$\sup_{a \in \mathcal{A}_{j}} \hat{L}_{a}(V_{j}) = \frac{r_{j}^{-\alpha}}{[u_{j}]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_{j}}}} \sup_{a \in \mathcal{A}_{j}} \left\{ \int_{\mathbb{R}^{d}} \left((L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x + r_{j}h) + \frac{(L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}(x - \delta h))}{\delta} \right) d\mu(h) - (1 + \frac{1}{\delta})(L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) \right\}$$

$$= \frac{r_j^{-\alpha}}{[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}} \sup_{a \in \mathcal{A}_j} \left\{ \int_{\mathbb{R}^d} (L_a u_j)(t_j + r_j^{\sigma} t, x_j + r_j x + r_j h) \right.$$

$$+ f_a(t_j + r_j^{\sigma} r, x_j + r_j x + r_j h) + \frac{1}{\delta} \left((L_a u_j)(t_j + r_j^{\sigma} t, x_j + r_j (x - \delta h)) \right.$$

$$+ f_a(t_j + r_j^{\sigma} t, x_j + r_j (x - \delta h)) \right) d\mu(h)$$

$$- (1 + \frac{1}{\delta}) ((L_a u_j)(t_j + r_j^{\sigma} t, x_j + r_j x) + f_a(t_j + r_j^{\sigma} t, x_j + r_j x)) \cdot \|\mu\|_{L_1}$$

$$- \int_{\mathbb{R}^d} \left(f_a(t_j + r_j^{\sigma} t, x_j + r_j x + r_j h) - f_a(t_j + r_j^{\sigma} t, x_j + r_j x) \right.$$

$$+ \frac{1}{\delta} (f_a(t_j + r_j^{\sigma} t, x_j + r_j (x - \delta h)) - f_a(t_j + r_j^{\sigma} t, x_j + r_j x) \right) d\mu(h) \right\}.$$

Note that for sufficiently large j such that $\max((-t)^{1/\sigma}, |x| + |h|) \leq R_j$ whenever $h \in \text{supp}\mu$, we have

$$|f_a(t_j + r_j^{\sigma}t, x_j + r_j x + r_j h) - f_a(t_j + r_j^{\sigma}t, x_j + r_j x)| \le [f_a]_{\alpha/\sigma, \alpha; Q_1} |r_j h|^{\alpha},$$

$$|f_a(t_j + r_j^{\sigma}t, x_j + r_j (x - \delta h)) - f_a(t_j + r_j^{\sigma}t, x_j + r_j x)| \le [f_a]_{\alpha/\sigma, \alpha; Q_1} |\delta r_j h|^{\alpha}.$$

Therefore, by the inequality

$$\sup\{f+g-h\} \ge \inf f + \inf g - \inf h,$$

we have that for $(t,x) \in \mathbb{R}_0^{d+1}$ and $h \in \operatorname{supp} \mu$ so that $\max((-t)^{1/\sigma}, |x| + |h|) \leq R_j$,

$$\sup_{a \in \mathcal{A}_j} \hat{L}_a(V_j)(t, x)$$

$$\geq \frac{r_{j}^{-\alpha}}{[u_{j}]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_{j}}}} \Big[\inf_{a \in \mathcal{A}_{j}} \int_{\mathbb{R}^{d}} (L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x + r_{j}h) \\ + f_{a}(t_{j} + r_{j}^{\sigma}r, x_{j} + r_{j}x + r_{j}h) d\mu(h) \\ + \inf_{a \in \mathcal{A}_{j}} \int_{\mathbb{R}^{d}} \frac{1}{\delta} \Big((L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}(x - \delta h)) \\ + f_{a}(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}(x - \delta h)) \Big) d\mu(h) \\ - \frac{\delta + 1}{\delta} \|\mu\|_{L_{1}} \inf_{a} \Big[(L_{a}u_{j})(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) + f_{a}(t_{j} + r_{j}^{\sigma}t, x_{j} + r_{j}x) \Big] \\ - (1 + \delta^{\alpha - 1}) r_{j}^{\alpha} \sup_{a \in \mathcal{A}_{j}} \Big[f_{a} \Big]_{\alpha/\sigma,\alpha;Q_{1}} \int_{\mathbb{R}^{d}} |h|^{\alpha} d\mu(h) \Big].$$

$$(4.11)$$

Since each u_j satisfies

$$\partial_t u_j = \inf_{a \in \mathcal{A}_j} (L_a u_j + f_a) \quad \text{in } Q_1, \tag{4.12}$$

it follows from (4.10) and (4.11) that in any bounded subset of \mathbb{R}_0^{d+1} , for sufficiently large j,

$$\partial_t V_j - \sup_{a \in \mathcal{A}_j} \hat{L}_a V_j \le \frac{1 + \delta^{\alpha - 1}}{[u_j]_{1 + \alpha/\sigma, \alpha + \sigma; Q^{k_j}}} \sup_{a \in \mathcal{A}_j} [f_a]_{\alpha/\sigma, \alpha} \int_{\mathbb{R}^d} |h|^{\alpha} d\mu(h). \tag{4.13}$$

We denote

$$V(t,x) := \int_{\mathbb{R}^d} \left[v(t,x+h) - v(t,x) - \frac{v(t,x) - v(t,x-\delta h)}{\delta} \right] d\mu(h).$$

For fixed $(t, x) \in \mathbb{R}_0^{d+1}$, by (4.8) in Lemma 4.3 and using the fact that μ has compact support, we have

$$\lim_{j} \partial_t V_j(t, x) = \partial_t \lim_{j} V_j(t, x) = \partial_t V(t, x), \tag{4.14}$$

for $|y| \leq 1$,

$$|\delta V_j(t, x, y)| \le C|y|^{\sigma + \alpha'}, \quad |\delta V(t, x, y)| \le C|y|^{\sigma + \alpha'}, \tag{4.15}$$

and for |y| > 1,

$$|V_i(t,y)|, |V(t,y)| \le C|y|^{\alpha},$$
 (4.16)

where C depends on μ . Clearly,

$$\sup_{a \in \mathcal{A}_j} |L_a(V_j - V)(t, x)| \le C \int_{\mathbb{R}^d} |\delta(V_j - V)(t, x, y)| |y|^{-d - \sigma} \, dy.$$

It follows from Lemma 4.3 that $\delta(V_j - V) \to 0$ locally uniformly. Therefore, by (4.15), (4.16), and the dominated convergence theorem, we have

$$\lim_{j} \sup_{a \in \mathcal{A}_{j}} |(L_{a}(V_{j} - V)(t, x))| = 0,$$

i.e.,

$$\lim_{j} \sup_{a \in \mathcal{A}_k} L_a V_j(t, x) = \sup_{a \in \mathcal{A}_k} L_a V(t, x). \tag{4.17}$$

Since μ has compact support, by Lemma 4.2 and (4.3), we have $R_i \to \infty$ and

$$[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}} \ge 2^{-k_j(\alpha-\alpha')}[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}} \to \infty.$$

For fixed $\delta \in (0,1)$, we send j to infinity to get from (4.13), (4.14), and (4.17) that

$$\partial_t V - \mathcal{M}^+ V \leq 0$$
 in \mathbb{R}_0^{d+1} .

By sending δ to 0 and using the dominated convergence theorem, we conclude that

$$\int_{\mathbb{R}^d} \left(v(t, x+h) - v(t, x) - h^T D v(t, x) \right) d\mu(h)$$

is a subsolution as well. Therefore, for $\sigma > 1$, v satisfies Condition (iii).

It remains to verify that v satisfies Condition (ii). Clearly, for fixed $(t, x), (s, h) \in \mathbb{R}_0^{d+1}$, when j is sufficiently large,

$$\partial_t (v_j(t+s,x+h) - v_j(t,x)) = r_j^{-\alpha} [u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}^{-1} \cdot (\partial_t u_j(t_j + r_j^{\sigma}(t+s), x_j + r_j(x+h)) - \partial_t u_j(t_j + r_j^{\sigma}t, x_j + r_jx)).$$
(4.18)

On the other hand,

$$\mathcal{M}^{-}(v_{j}(t+s,x+h)-v_{j}(t,x)) = r_{j}^{-\alpha}[u_{j}]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_{j}}}^{-1} \cdot \mathcal{M}^{-}(u_{j}(t_{j}+r_{j}^{\sigma}(t+s),x_{j}+r_{j}(x+h)) - u_{j}(t_{j}+r_{j}^{\sigma}t,x_{j}+r_{j}x)).$$
(4.19)

Combining (4.12), (4.18), and (4.19), we obtain that for j sufficiently large,

$$\partial_t (v_j(t+s,x+h) - v_j(t,x)) - \mathcal{M}^- (v_j(t+s,x+h) - v_j(t,x))$$

$$\geq -[u_j]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_j}}^{-1} \sup_a [f_a]_{\alpha/\sigma,\alpha;Q_1}.$$

By sending j to infinity, we get for any $(t, x) \in \mathbb{R}_0^{d+1}$,

$$\partial_t \big(v(t+s,x+h) - v(t,x) \big) - \mathcal{M}^- \big(v(t+s,x+h) - v(t,x) \big) \ge 0.$$

Similarly,

$$\partial_t \big(v(t+s,x+h) - v(t,x) \big) - \mathcal{M}^+ \big(v(t+s,x+h) - v(t,x) \big) \le 0.$$

The lemma is proved.

Now we are ready to finish

Proof of Theorem 4.1. By Lemma 4.4 and Theorem 3.1, v is a polynomial of order ν in x and 1 in t. Since at the origin v_j along with its first derivative in t and up to ν -th order derivatives in x are 0, by Lemma 4.3 the same is true for v. Therefore, $v \equiv 0$. This gives us a contradiction with (4.7) and Lemma 4.3. The proof is completed.

4.2. Equations with (t,x)-dependent kernels. In this subsection, we consider the case that kernels also depend on (t,x) and Hölder continuous in (t,x), i.e., there exists A>0 such that for any $a\in\mathcal{A}$, (1.4) is satisfied. We only prove Theorem 1.1 in the case when $\sigma+\alpha>2$ and the proof of the cases $\sigma+\alpha<2$ is similar and actually simpler. Below we divide the proof into several steps.

Let η be a nonnegative smooth cutoff function with $\eta \equiv 1$ in Q_1 and vanishes outside $(-(5/4)^{\sigma}, (5/4)^{\sigma}) \times B_{5/4}$. Set $v := \eta u$ and note that in Q_1 ,

$$v_{t} = \eta u_{t} + \eta_{t} u = \inf_{a \in \mathcal{A}} (\eta L_{a} u + \eta f_{a} + \eta_{t} u)$$
$$= \inf_{a \in \mathcal{A}} (L_{a}^{0} v + (L_{a} - L_{a}^{0}) v + \eta L_{a} u - L_{a} v + \eta f_{a} + \eta_{t} u)$$

where

$$L_a^0 v = \int_{\mathbb{R}^d} \delta v(t, x, y) K_a(0, 0, y) \, dy.$$

We further define

$$h_a := \eta L_a u - L_a v$$

$$= \int_{\mathbb{R}^d} ((\eta(t, x) - \eta(t, x + y)) u(t, x + y) + y^T D \eta(t, x) u(t, x)) K_a(t, x, y) dy$$

and

$$g_a := (L_a - L_a^0)v = \int_{\mathbb{R}^d} \delta v(t, x, y) \big(K_a(t, x, y) - K_a(0, 0, y) \big) dy.$$

Here in order to apply the argument of freezing the coefficients, we subtracted and added $K_a(0,0,y)$ in the formula above.

Lemma 4.5. Assume that $u \in C^{1+\alpha/\sigma,\sigma+\alpha}(Q_{11/8}) \cap C^{\alpha/\sigma,\alpha}((-(11/8)^{\sigma},0) \times \mathbb{R}^d)$. Let h_a and g_a be functions defined above. Then for any $\alpha \in \mathcal{A}$, we have

$$[g_a]_{\alpha/\sigma,\alpha:Q_1} \le CA([v]_{1+\alpha/\sigma,\alpha+\sigma} + [v]_{\alpha/\sigma,\alpha}), \tag{4.20}$$

$$[h_a]_{\alpha/\sigma,\alpha;Q_1} \le C(A+1) (\|u\|_{\alpha/\sigma,\alpha;(-(11/8)^\sigma,0)\times\mathbb{R}^d} + \|D^2 u\|_{L_\infty(Q_{11/8})}). \tag{4.21}$$

Moreover,

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{1/2}} \le C\Big([u]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_{5/4}} + C_0 + A[u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{5/4}} + (A+1)(\|u\|_{\alpha/\sigma,\alpha;(-(11/8)^{\sigma},0)\times\mathbb{R}^d} + \|D^2u\|_{L_{\infty}(Q_{11/8})})\Big). \tag{4.22}$$

Here the constant C depends only on d, λ , Λ , α , and σ , and is uniformly bounded as $\sigma \to 2$.

Proof. For $(t, x), (t', x') \in Q_1$, set $l = \max(|x - x'|, |t - t'|^{1/\sigma})$. Without loss of generality, we may assume that $l \le 1/4$.

Estimates of g_a : From the definition and the triangle inequality,

$$\begin{aligned} & \left| g_{a}(t,x) - g_{a}(t',x') \right| \\ & = \left| \int_{\mathbb{R}^{d}} \delta v(t,x,y) \left(K_{a}(t,x,y) - K_{a}(0,0,y) \right) dy \right| \\ & - \int_{\mathbb{R}^{d}} \delta v(t',x',y) \left(K_{a}(t',x',y) - K_{a}(0,0,y) \right) dy \right| \\ & \leq \left| \int_{\mathbb{R}^{d}} \left(\delta v(t,x,y) - \delta v(t',x',y) \right) \left(K_{a}(t,x,y) - K_{a}(0,0,y) \right) dy \right| \\ & + \left| \int_{\mathbb{R}^{d}} \delta v(t',x',y) \left(K_{a}(t,x,y) - K_{a}(t',x',y) \right) dy \right| \\ & =: I + II. \end{aligned}$$

Then we estimate I and II separately. First, similar to (3.8), I is less than

$$\begin{split} &\int_{B_l} \left| \left(\delta v(t,x,y) - \delta v(t',x',y) \right) \left(K_a(t,x,y) - K_a(0,0,y) \right) \right| dy \\ &+ \int_{\mathbb{R}^d \setminus B_l} \left| \left(\delta v(t,x,y) - \delta v(t',x',y) \right) \left(K_a(t,x,y) - K_a(0,0,y) \right) \right| dy := \mathbf{I}_1 + \mathbf{I}_2. \end{split}$$

Applying (3.6), we have

$$\begin{split} & \mathrm{I}_{1} \leq C[v]_{1+\alpha/\sigma,\alpha+\sigma;Q_{5/4}} \int_{B_{l}} l^{\alpha+\sigma-2} |y|^{2} \big(K_{a}(t,x,y) - K_{a}(0,0,y) \big) \, dy \\ & \leq C A(2-\sigma)[v]_{1+\alpha/\sigma,\alpha+\sigma} l^{\alpha+\sigma-2} \big(|x|^{\alpha} + |t|^{\alpha/\sigma} \big) \int_{B_{l}} |y|^{2} |y|^{-d-\sigma} \, dy \\ & = C A l^{\alpha}[v]_{1+\alpha/\sigma,\alpha+\sigma}. \end{split}$$

For I_2 , we have

$$I_{2} \leq C[v]_{1+\alpha/\sigma,\alpha+\sigma} l \int_{\mathbb{R}^{d} \setminus B_{l}} |y|^{\sigma+\alpha-1} |K_{a}(t,x,y) - K_{a}(0,0,y)| dy$$

$$\leq CA(2-\sigma)[v]_{1+\alpha/\sigma,\alpha+\sigma} l^{\alpha} (|x|^{\alpha} + |t|^{\alpha/\sigma}) \leq CAl^{\alpha}[v]_{1+\alpha/\sigma,\alpha+\sigma}.$$

Next, we bound

$$II \leq \int_{\mathbb{R}^d \setminus B_1} \left([v]_{\alpha/\sigma,\alpha} |y|^{\alpha} + \|Dv\|_{L_{\infty}} |y| \right) |K_a(t,x,y) - K_a(t',x',y)| \, dy
+ \int_{B_1} \|D^2 v\|_{L_{\infty}} |y|^2 |K_a(t,x,y) - K_a(t',x',y)| \, dy
\leq CAl^{\alpha} \left([v]_{\alpha/\sigma,\alpha} + \|Dv\|_{\infty} + \|D^2 v\|_{L_{\infty}} \right).$$

Combining the estimates of I, II, and the interpolation inequality, we get

$$|g_a(t,x) - g_a(t',x')| \le CAl^{\alpha}([v]_{1+\alpha/\sigma,\alpha+\sigma} + [v]_{\alpha/\sigma,\alpha}),$$

which implies (4.20).

Estimates of h_a : For simplicity of notation, we denote

$$\xi(t, x, y) = (\eta(t, x) - \eta(t, x + y))u(t, x + y) + y^{T}D\eta(t, x)u(t, x).$$
(4.23)

By the Leibniz rule, we have

$$\xi(t, x, y) = y^{T} \int_{0}^{1} \left(D\eta(t, x) u(t, x) - D\eta(t, x + sy) u(t, x + y) \right) ds$$

$$= -\int_{0}^{1} \int_{0}^{1} \left(u(t, x) s y^{T} D^{2} \eta(t, x + s'sy) y + y^{T} D\eta(t, x + sy) y^{T} Du(t, x + s'y) \right) ds' ds, \tag{4.24}$$

which implies that when $|y| \leq 1/8$,

$$|\xi(t,x,y)| \le C|y|^2 (||u||_{L_{\infty}(Q_{11/8})} + ||Du||_{L_{\infty}(Q_{11/8})}).$$
 (4.25)

On the other hand, clearly when $|y| \ge 1/8$,

$$|\xi(t,x,y)| \le C(||u||_{L_{\infty}((-1,0)\times\mathbb{R}^d)} + |y|||u||_{L_{\infty}(Q_1)}).$$
 (4.26)

Note that

$$\left| h_a(t,x) - h_a(t',x') \right| \le \int_{\mathbb{R}^d} \left| \xi(t,x,y) - \xi(t',x',y) \right| K_a(t,x,y) \, dy
+ \int_{\mathbb{R}^d} \left| \xi(t',x',y) \right| \left| K_a(t,x,y) - K_a(t',x',y) \right| \, dy =: \text{III} + \text{IV}.$$
(4.27)

Estimate of III: By (4.24) when $|y| \le 1/8$, we have

$$\begin{split} \left| \xi(t,x,y) - \xi(t',x',y) \right| \\ &= \left| \int_0^1 \int_0^1 s \left(u(t',x') y^T D^2 \eta(t',x'+ss'y) y - u(t,x) y^T D^2 \eta(t,x+ss'y) y \right) dx \, ds' \\ &+ \int_0^1 \int_0^1 \left[y^T D \eta(t',x'+sy) y^T D u(t',x'+s'y) \\ &- y^T D \eta(t,x+sy) y^T D u(t,x+s'y) \right] ds \, ds' \right| \\ &\leq C |y|^2 l^{\alpha} \left([u]_{\alpha/\sigma,\alpha;Q_1} + \|u\|_{L_{\infty}(Q_1)} \right) + C |y|^2 l \left(\|D^2 u\|_{L_{\infty}(Q_{11/8})} + \|D u\|_{L_{\infty}(Q_{11/8})} \right) \\ &\leq C |y|^2 l^{\alpha} \left(\|u\|_{L_{\infty}(Q_{11/8})} + \|D^2 u\|_{L_{\infty}(Q_{11/8})} \right), \end{split}$$

where we used the interpolation inequalities in the last inequality. On the other hand, when |y| > 1/8,

$$\left| y^T D\eta(t,x) u(t,x) - y^T D\eta(t',x') u(t',x') \right| \le |y| l^{\alpha} ||u||_{\alpha/\sigma,\alpha;Q_1}$$

and

$$\begin{split} & \left| \left(\eta(t,x) - \eta(t,x+y) \right) u(t,x+y) - \left(\eta(t',x') - \eta(t',x'+y) \right) u(t',x'+y) \right| \\ &= \left| u(t,x+y) \left(\eta(t,x) - \eta(t',x') - \eta(t,x+y) + \eta(t',x'+y) \right) \right. \\ & \left. + \left(\eta(t',x') - \eta(t',x'+y) \right) \left(u(t,x+y) - u(t',x'+y) \right) \right| \\ &\leq C(l \|u\|_{L_{\infty}((-1,0) \times \mathbb{R}^d)} + l^{\alpha} \|u\|_{\alpha/\sigma,\alpha;(-1,0) \times \mathbb{R}^d}), \end{split}$$

which imply that when |y| > 1/8,

$$\begin{aligned} & |\xi(t, x, y) - \xi(t', x', y)| \\ & \leq C(l||u||_{L_{\infty}((-1, 0) \times \mathbb{R}^d)} + l^{\alpha}||u||_{\alpha/\sigma, \alpha; (-1, 0) \times \mathbb{R}^d}) + C|y|l^{\alpha}||u||_{\alpha/\sigma, \alpha; Q_1}. \end{aligned}$$

Now with the above estimates, we obtain

$$\begin{split} & \text{III} \leq \int_{B_{1/8}} C|y|^2 l^{\alpha} \big(\|u\|_{L_{\infty}(Q_{11/8})} + \|D^2 u\|_{L_{\infty}(Q_{11/8})} \big) K_a(t, x, y) \, dy \\ & + C l^{\alpha} \|u\|_{\alpha/\sigma, \alpha; Q_1} \int_{B_{1/8}^c} |y| K_a(t, x, y) \, dy \\ & + C \big(l\|u\|_{L_{\infty}((-1, 0) \times \mathbb{R}^d)} + l^{\alpha} \|u\|_{\alpha/\sigma, \alpha; (-1, 0) \times \mathbb{R}^d} \big) \int_{B_{1/8}^c} K_a(t, x, y) \, dy \\ & \leq C l^{\alpha} \big(\|D^2 u\|_{L_{\infty}(Q_{11/8})} + \|u\|_{\alpha/\sigma, \alpha; (-(11/8)^{\sigma}, 0) \times \mathbb{R}^d} \big). \end{split}$$

Estimate of IV: By (4.25) and (4.26), we have

$$IV \le Cl^{\alpha} A(\|u\|_{L_{\infty}((-(11/8)^{\sigma},0)\times\mathbb{R}^d)} + \|Du\|_{L_{\infty}(Q_{11/8})}).$$

The estimates of III and IV with the interpolation inequalities give (4.21). Now we apply Theorem 4.1 to v with the estimates of g_a and h_a in Lemma 4.5 to obtain

$$[v]_{1+\alpha/\sigma,\alpha+\sigma;Q_{1/2}} \le C\Big([v]_{1+\alpha'/\sigma,\alpha'+\sigma} + A[v]_{1+\alpha/\sigma,\alpha+\sigma} + A[v]_{\alpha/\sigma,\alpha} + (A+1)(\|u\|_{\alpha/\sigma,\alpha;(-(11/8)^{\sigma},0)\times\mathbb{R}^d} + \|D^2u\|_{L_{\infty}(Q_{11/8})}) + \sup_{\sigma} [\eta f_a]_{\alpha/\sigma,\alpha;Q_1}\Big).$$

Since $\eta \equiv 1$ in Q_1 and has compact support in $(-(5/4)^{\sigma}, (5/4)^{\sigma}) \times B_{5/4}$, we get (4.21). The lemma is proved.

Proof of Theorem 1.1. We first use a scaling argument. For any $\varepsilon > 0$, set $\hat{u}(t, x) := \varepsilon^{-\sigma} u(\varepsilon^{\sigma} t, \varepsilon x)$. Since u satisfies (1.5), we have

$$\hat{u}_t(t,x) = \inf_{a} \left\{ \int_{\mathbb{R}^d} \delta \hat{u}(t,x,y) K_a^{\varepsilon}(t,x,y) \, dy + f_a(\varepsilon^{\sigma}t,\varepsilon x) \right\} \quad \text{in } Q_{1/\varepsilon},$$

where

$$K_a^{\varepsilon}(t, x, y) = \varepsilon^{d+\sigma} K_a(\varepsilon^{\sigma} t, \varepsilon x, \varepsilon y).$$

Clearly,

$$\left| K_a^{\varepsilon}(t,x,y) - K_a^{\varepsilon}(t',x',y) \right| \le A(2-\sigma)\varepsilon^{\alpha} \left(|x-x'|^{\alpha} + |t-t'|^{\alpha/\sigma} \right) \frac{\Lambda}{|y|^{d+\sigma}}.$$

Then we apply (4.22) to \hat{u} and get

$$[\hat{u}]_{1+\alpha/\sigma,\alpha+\sigma;Q_{1/2}} \leq C \Big([\hat{u}]_{1+\alpha'/\sigma,\alpha+\sigma;Q_{5/4}} + C_0 \varepsilon^{\alpha} + A \varepsilon^{\alpha} [\hat{u}]_{1+\alpha/\sigma,\alpha+\sigma;Q_{5/4}} + (A \varepsilon^{\alpha} + 1) \Big(\|\hat{u}\|_{\alpha/\sigma,\alpha;(-(11/8)^{\sigma},0) \times \mathbb{R}^d} + \|D^2 \hat{u}\|_{L_{\infty}(Q_{11/8})} \Big) \Big).$$

Returning back to u, we have

$$[u]_{1+\alpha/\sigma,\sigma+\alpha;Q_{\varepsilon/2}} \leq C \Big(\varepsilon^{\alpha'-\alpha} [u]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_{5\varepsilon/4}} + C_0 + A \varepsilon^{\alpha} [u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{5\varepsilon/4}} + (A \varepsilon^{\alpha} + 1) \Big(\varepsilon^{-\sigma-\alpha} \|u\|_{\alpha/\sigma,\alpha;(-(11\varepsilon/8)\sigma,0)\times\mathbb{R}^d} + \varepsilon^{2-\sigma-\alpha} \|D^2 u\|_{L_{\infty}(Q_{11\varepsilon/8})} \Big) \Big).$$

By a translation of the coordinates, the inequality above holds for any $(t, x) \in Q_1$ for sufficiently small $\varepsilon > 0$

$$[u]_{1+\alpha/\sigma,\sigma+\alpha;Q_{\varepsilon/2}(t,x)}$$

$$\leq C \Big(\varepsilon^{\alpha'-\alpha}[u]_{1+\alpha'/\sigma,\alpha'+\sigma;Q_{5\varepsilon/4}(t,x)} + C_0 + A\varepsilon^{\alpha}[u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{5\varepsilon/4}(t,x)} + (A\varepsilon^{\alpha}+1)\Big(\varepsilon^{-\sigma-\alpha}\|u\|_{\alpha/\sigma,\alpha;(t-(11\varepsilon/8)^{\sigma},t)\times\mathbb{R}^d} + \varepsilon^{2-\sigma-\alpha}\|D^2u\|_{L_{\infty}(Q_{11\varepsilon/8}(t,x))}\Big)\Big).$$

$$(4.28)$$

Let Q^k be defined in (4.1). It is obvious that Q^k monotonically increases to Q_1 . Then for any $(t,x), (s,y) \in Q^k$ such that $t \geq s$, we set $l := \max(|t-s|^{1/\sigma}, |x-y|)$. When $l \geq \varepsilon/2$,

$$\frac{|D^{2}u(t,x) - D^{2}u(s,y)|}{l^{\sigma+\alpha-2}} + \frac{|u_{t}(t,x) - u_{t}(s,y)|}{l^{\sigma+\alpha-2}} \le 2^{\sigma+\alpha-1} \varepsilon^{2-\sigma-\alpha} (||u_{t}||_{L_{\infty}(Q^{k})} + ||D^{2}u||_{L_{\infty}(Q^{k})});$$

when $l < \varepsilon/2$,

$$\frac{|D^2u(t,x)-D^2u(s,y)|}{l^{\sigma+\alpha-2}}+\frac{|u_t(t,x)-u_t(s,y)|}{l^{\sigma+\alpha-2}}\leq 2[u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{\varepsilon/2}(t,x)}\cdot$$

Now we choose $\varepsilon = 2^{-k-2}$ so that for any $(t,x) \in Q^k$, $Q_{11\varepsilon/8}(t,x) \subset Q^{k+1}$ and $(t-(11\varepsilon/8)^{\sigma},t) \subset (-1,0)$. Combining the two inequalities above with (4.28), we obtain

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k}} \leq 2^{(k+3)(\sigma+\alpha-2)+1} (\|u_{t}\|_{L_{\infty}(Q^{k})} + \|D^{2}u\|_{L_{\infty}(Q^{k})})$$

$$+ C \Big(2^{(k+2)(\alpha-\alpha')} [u]_{1+\alpha'/\sigma,\alpha'+\sigma;Q^{k+1}} + C_{0} + 2^{-(k+2)\alpha} A[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k+1}}$$

$$+ (2^{-(k+2)\alpha}A + 1) \Big(2^{(k+2)(\sigma+\alpha)} \|u\|_{\alpha/\sigma,\alpha;(-1,0)\times\mathbb{R}^{d}}$$

$$+ 2^{(k+2)(\sigma+\alpha-2)} \|D^{2}u\|_{L_{\infty}(Q^{k+1})}\Big)\Big). \tag{4.29}$$

By the interpolation inequalities

$$\begin{split} &[u]_{1+\alpha'/\sigma,\alpha'+\sigma;Q^{k+1}} \leq 2^{-2(k+2)(\alpha-\alpha')}[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k+1}} + C2^{2(k+2)(\sigma+\alpha')}\|u\|_{L_{\infty}}, \\ &\|D^2u\|_{L_{\infty}(Q^{k+1})} + \|u_t\|_{L_{\infty}(Q^{k+1})} \\ &\leq 2^{-2(k+1)(\sigma+\alpha-2)}[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k+1}} + C2^{4(k+1)}\|u\|_{L_{\infty}(Q^{k+1})}, \end{split}$$

we reorganize the right-hand side of (4.29) to get

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^k} \le C\Big(\Big(2^{-(k+1)(\sigma+\alpha-2)} + 2^{-(k+2)(\alpha-\alpha')} \Big) [u]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k+1}} + 2^{5k(\sigma+\alpha)} ||u||_{\alpha/\sigma,\alpha;(-1,0)\times\mathbb{R}^d} + C_0 \Big),$$

where C depends on A. Obviously, $\sigma + \alpha < 3$ and there exists a constant k_0 depends on d, σ_0 , α , λ , Λ , and A such that $Q_{1/2} \subset Q^{k_0}$ and for any $k \geq k_0$,

$$C(2^{-(k+1)(\sigma+\alpha-2)} + 2^{-2(k+1)(\alpha-\alpha')}) < 2^{-16}.$$

Therefore, we have for any $k \ge k_0$,

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^k} \le 2^{-16} [u]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k+1}} + C2^{15k} ||u||_{\alpha/\sigma,\alpha} + CC_0.$$

We multiply both sides above by $2^{-16(k-k_0)}$ and then sum from $k = k_0$ to infinity and obtain that

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q^{k_0}} \le C2^{15k_0} ||u||_{\alpha/\sigma,\alpha;(-1,0)\times\mathbb{R}^d} + CC_0.$$

In particular,

$$[u]_{1+\alpha/\sigma,\alpha+\sigma;Q_{1/2}} \le C(\|u\|_{\alpha/\sigma,\alpha;(-1,0)\times\mathbb{R}^d} + C_0).$$

The proof is completed.

4.3. An improved estimate. By a more careful analysis, we obtain the following corollary when the kernels depend only on y.

Corollary 4.6. Let $\sigma \in (0,2)$ and $0 < \lambda \leq \Lambda$. Assume that for any $a \in \mathcal{A}$, K_a only depends on y. There is a constant $\hat{\alpha} \in (0,1)$ depending on d, σ , λ , and Λ (uniformly as $\sigma \to 2$) so that the following holds. Let $\alpha \in (0,\hat{\alpha})$. Suppose $u \in C^{1+\alpha/\sigma,\sigma+\alpha}(Q_1) \cap C^{\alpha/\sigma,\alpha}_{loc}([-1,0] \times \mathbb{R}^d)$ is a solution of

$$u_t = \inf_{a \in A} (L_a u + f_a)$$
 in Q_1 .

Then,

$$[u]_{1+\alpha/\sigma,\alpha+\sigma,Q_{1/2}}$$

$$\leq C[u]_{\alpha/\sigma,\alpha;(-1,0)\times B_2} + C\sum_{j=2}^{\infty} 2^{-j\sigma}[u]_{\alpha/\sigma,\alpha;(-1,0)\times (B_{2^j}\setminus B_{2^{j-1}})} + CC_0,$$
(4.30)

where $C_0 = \sup_a [f_a]_{\alpha/\sigma,\alpha;Q_1}$ and C > 0 depends only on d, λ , Λ , α , and σ , and is uniformly bounded as $\sigma \to 2$.

Proof. Since the proof is quite similar to the proof of Theorem 1.1, we only provide a sketch here. By a standard scaling and covering argument, we may assume that $u \in C^{1+\alpha/\sigma,\sigma+\alpha}(Q_2) \cap C^{\alpha/\sigma,\alpha}_{loc}([-2,0] \times \mathbb{R}^d)$ and the equation is satisfied in Q_2 . Let η be a cutoff function such that $\eta \in C^{\infty}_0((-2^{\sigma},2^{\sigma}) \times B_2)$ and $\eta \equiv 1$ in $Q_{5/4}$. Let $v = \eta u$, which satisfies

$$v_t = \inf_a (L_a v + h_a + \eta f_a + \eta_t u),$$

where

$$h_a = \int_{\mathbb{R}^d} \xi(t, x, y) K_a(y) \, dy$$

and ξ is defined in (4.23). It is sufficient to estimate $[h_a]_{\alpha/\sigma,\alpha;Q_1}$. Since K_a only depends on y, it follows that

$$|h_a(t,x) - h_a(t',x')| = III,$$

where III is defined in (4.27). The estimate is similar to the one in the proof of Lemma 4.5. For any $(t, x), (t', x') \in Q_1$, since $\eta \equiv 1$ in $Q_{5/4}, D\eta(t, x) = D\eta(t', x') = 0$. When $|y| \le 1/4$, $\xi(t, x, y) = 0$; When |y| > 1/4, we have

$$\begin{aligned} & \left| \xi(t, x, y) - \xi(t', x', y) \right| \\ &= \left| (\eta(t, x) - \eta(t, x + y)) u(t, x + y) + y^T D \eta(t, x) u(t, x) \right. \\ & \left. - (\eta(t', x') - \eta(t', x' + y)) u(t', x' + y) + y^T D \eta(t', x') u(t', x') \right| \\ &\leq \left| \eta(t, x + y) u(t, x + y) - \eta(t', x' + y) u(t', x' + y) \right| + \left| u(t, x + y) - u(t', x' + y) \right|. \end{aligned}$$

Recall that $l = \max\{|x - x'|, |t - t'|^{1/\sigma}\}$. Combining the estimate above, we obtain

$$\begin{aligned} & \text{III} = \int_{B_{1/4}^c} \left| \xi(t, x, y) - \xi(t', x', y) \right| K_a(y) \, dy \\ & \leq C l^{\alpha} \|u\|_{\alpha/\sigma, \alpha; (-1,0) \times B_2} + \int_{B_{1/4}^c} \left| u(t, x + y) - u(t', x' + y) \right| K_a(y) \, dy \\ & \leq C l^{\alpha} \|u\|_{\alpha/\sigma, \alpha; (-1,0) \times B_2} + \sum_{j=-1}^{\infty} \int_{B_{2j} \setminus B_{2j-1}} |u(t, x + y) - u(t', x' + y)| K_a(y) \, dy \\ & \leq C l^{\alpha} \|u\|_{\alpha/\sigma, \alpha; (-1,0) \times B_2} + C l^{\alpha} \sum_{j=-1}^{\infty} 2^{-j\sigma} [u]_{\alpha/\sigma, \alpha; (-1,0) \times B_{2j+1}} \\ & \leq C l^{\alpha} \left(\|u\|_{\alpha/\sigma, \alpha; (-1,0) \times B_2} + \sum_{j=-1}^{\infty} 2^{-j\sigma} [u]_{\alpha/\sigma, \alpha; (-1,0) \times (B_{2j} \setminus B_{2j-1})} \right), \end{aligned}$$

which implies that

$$[h_a]_{\alpha/\sigma,\alpha;Q_1} \le C\Big(\|u\|_{\alpha/\sigma,\alpha;(-1,0)\times B_2} + \sum_{i=1}^{\infty} 2^{-j\sigma}[u]_{\alpha/\sigma,\alpha;(-1,0)\times (B_{2^j}\setminus B_{2^{j-1}})}\Big).$$

Then we apply Theorem 1.1 to v and obtain

$$[v]_{1+\alpha/\sigma,\sigma+\alpha;Q_{1/2}} \le C\Big(\|v\|_{\alpha/\sigma,\alpha;(-1,0)\times\mathbb{R}^d} + \|u\|_{\alpha/\sigma,\alpha;(-1,0)\times B_2} + \sum_{j=1}^{\infty} 2^{-j\sigma} [u]_{\alpha/\sigma,\alpha;(-1,0)\times (B_{2j}\setminus B_{2j-1})} + C_0\Big).$$

Combining the fact that $\eta \equiv 1$ in $Q_{5/4}$ and replacing u by u - u(0,0), we reach (4.30). Therefore, the proof is completed.

5. Equations with bounded inhomogeneous terms

In this section, we present an application of Corollary 4.6 to nonlocal parabolic equations with merely bounded nonhomogeneous terms:

$$u_t = \inf_{a \in \mathcal{A}} (L_a u + f_a), \tag{5.1}$$

where $\sup_a ||f_a||_{L_\infty} < \infty$ and

$$L_a u(x) = \int_{\mathbb{R}^d} \delta u(t, x, y) K_a(y) \, dy.$$

Before proving Theorem 1.2, we first present an interpolation inequality involving the Zygmund semi-norm. The proof can be found in the appendix

Lemma 5.1. Let $\alpha \in (0,1)$ and $f \in \Lambda^1((-1,0)) \cap L_{\infty}((-1,0))$. Then we have $f \in C^{\alpha}((-1,0))$ and

$$[f]_{\alpha;(-1,0)} \le C ||f||_{L_{\infty}((-1,0))} + C[f]_{\Lambda^{1}((-1,0))},$$
 (5.2)

where C depends only on α .

In the sequel, we set

$$[u]_{\Lambda^1}^t := [u]_{\Lambda^1(\mathbb{R}_0^{d+1})}^t, \quad [u]_{\sigma}^* := [u]_{\sigma;\mathbb{R}_0^{d+1}}^*, \quad \text{and} \quad [Du]_{\frac{\sigma-1}{\sigma}}^t := [Du]_{\frac{\sigma-1}{\sigma};\mathbb{R}_0^{d+1}}^t.$$

Let η be a smooth even nonnegative function in \mathbb{R} with unit integral and vanishing outside (-1,1). For R>0, we define the mollification of u with respect to t as

$$u^{(R)}(t,x) = \int_{\mathbb{R}} \left(2u(t - R^{\sigma}s, x) - u(t - 2R^{\sigma}s, x) \right) \eta(s - 1) \, ds.$$

The following lemmas will also be used in our proof. We present their proofs in the appendix.

Lemma 5.2. Let $\beta \in (0,1]$ and R > 0. Then we have

$$[\partial_t u^{(R)}]_{\beta;\mathbb{R}_0^{d+1}}^t \le C(\beta) R^{-\beta\sigma} [u]_{\Lambda^1}^t. \tag{5.3}$$

Lemma 5.3. Let $\sigma \in (1,2)$, $\alpha \in (0,1)$, and R > 0 be constants. Assume that u defined on \mathbb{R}_0^{d+1} is C^{σ} in x, Λ^1 in t, and Du is $C^{(\sigma-1)/\sigma}$ in t. Let p = p(t,x) be the first-order Taylor expansion of $u^{(R)}$ at the origin. Then for any integer $j \geq 0$, we have

$$\begin{aligned} &[u-p]^*_{\alpha;(-R^{\sigma},0)\times B_{2^{j}R}} \\ &\leq C(2^{j}R)^{\sigma-\alpha}[u]^*_{\sigma} + C2^{(1-\alpha)j}R^{\sigma-\alpha}[Du]^t_{\frac{\sigma-1}{\sigma}} + C2^{-j\alpha}R^{\sigma-\alpha}[u]^t_{\Lambda^{1}} \end{aligned} \tag{5.4}$$

and

$$\begin{split} [u-p]^t_{\alpha/\sigma;(-R^{\sigma},0)\times B_{2^jR}} &\leq C2^{j(\sigma-\alpha/2)}R^{\sigma-\alpha}[u]^*_{\sigma} + C2^{j(\sigma+1-\alpha)/2}R^{\sigma-\alpha}[Du]^t_{\frac{\sigma-1}{\sigma}} \\ &\quad + C2^{j(\sigma-\alpha/2)}R^{\sigma-\alpha}[u]^t_{\Lambda^1}, \end{split} \tag{5.5}$$

where C > 0 is a constant depending only on d, σ , and α .

In the case when $\sigma = 1$, we define $u^{(R)}$ differently. Let $\zeta \in C_0^{\infty}(B_1)$ be a radial nonnegative function with unit integral. For R > 0, we define

$$u^{(R)}(t,x) = \int_{\mathbb{R}^{d+1}} \left(2u(t - Rs, x - Ry) - u(t - 2Rs, x - Ry) \right) \eta(s - 1) \zeta(y) \, dy \, ds.$$

Lemma 5.4. Let $\alpha \in (0,1)$, and R > 0 be constants. Assume that u defined on \mathbb{R}_0^{d+1} is Λ^1 in (t,x). Let p = p(t,x) be the first-order Taylor expansion of $u^{(R)}$ at the origin. Then for any integer $j \geq 0$, we have

$$[u-p]_{\alpha,\alpha;(-R,0)\times B_{2^{j}R}} \le C2^{j(1-\alpha/2)}R^{1-\alpha}[u]_{\Lambda^{1}}, \tag{5.6}$$

where C > 0 is a constant depending only on d and α .

Define \mathcal{P}_0 to be the set of first-order polynomials of t, and \mathcal{P}_1 to be the set of first-order polynomials of (t, x).

Lemma 5.5. (i) When $\sigma \in (0,1)$, we have

$$[u]_{\Lambda^{1}}^{t} + [u]_{\sigma}^{*} \leq C \sup_{r>0} \sup_{(t,x) \in \mathbb{R}_{0}^{d+1}} r^{-\sigma} \inf_{p \in \mathcal{P}_{0}} ||u - p||_{L_{\infty}(Q_{r}(t,x))},$$

where C > 0 is a constant depending only on d and σ .

(ii) When $\sigma \in (1,2)$, we have

$$[u]_{\Lambda^{1}}^{t} + [u]_{\sigma}^{*} + [Du]_{\frac{\sigma-1}{\sigma}}^{t} \leq C \sup_{r>0} \sup_{(t,x)\in\mathbb{R}_{0}^{d+1}} r^{-\sigma} \inf_{p\in\mathcal{P}_{1}} \|u-p\|_{L_{\infty}(Q_{r}(t,x))},$$

where C > 0 is a constant depending only on d and σ .

(iii) We have

$$[u]_{\Lambda^1} \le C \sup_{r>0} \sup_{(t,x) \in \mathbb{R}_0^{d+1}} r^{-1} \inf_{p \in \mathcal{P}_1} \|u - p\|_{L_{\infty}(Q_r(t,x))},$$

where C > 0 is a constant depending only on d.

Proof. The estimates of $[u]_{\sigma}^*$ and $[Du]_{\frac{t}{\sigma}-1}^t$ are standard. See, for instance, [19, Section 3.3]. We only consider $[u]_{\Lambda^1}^t$. For any polynomial p which is linear in t, by the triangle inequality,

$$\begin{aligned} & \left| u(t+s,x) + u(t-s,x) - 2u(t,x) \right| \\ & = \left| u(t+s,x) - p(t+s,x) + u(t-s,x) - p(t-s,x) - 2(u(t,x) - p(t,x)) \right| \\ & \leq 4 \|u - p\|_{L_{\infty}(Q_r(t+s,x))}, \end{aligned}$$

where $r^{\sigma} = 2s$. Since p is arbitrary, the inequality above implies that

$$|u(t+s,x) + u(t-s,x) - 2u(t,x)| \le 8sr^{-\sigma} \inf_{p} ||u-p||_{L_{\infty}(Q_r(t+s,x))}.$$

Similarly, we can prove Assertion (iii). The lemma is proved.

Proof of Theorem 1.2. We only treat the case when $\sigma \geq 1$. For the case when $\sigma < 1$, the proof is almost the same with minor modifications.

First we assume $\sigma > 1$. Let $\hat{\alpha}$ be the constant in Corollary 4.6 and $\alpha \in (0, \hat{\alpha})$ be such that $\alpha + \sigma \leq 2$. Let R > 0 be a constant, p be defined as in Lemma 5.3, and $c_0 = \partial_t p$. Let $K \geq 2||u - p||_{L_{\infty}(Q_{2R})}$ be a constant to be specified later and denote

$$g_K = \max \big(\min(u - p, K), -K \big).$$

Clearly, $g_K \in C^{\alpha/\sigma,\alpha}$ and any $C^{\alpha/\sigma,\alpha}$ norm (or semi-norm) of g_K is less than or equal to that of u-p.

Let v_K be the solution to

$$\begin{cases} \partial_t v_K = \inf_a (L_a v_K - c_0) & \text{in } Q_{2R}, \\ v_K = g_K & \text{in } \mathbb{R}_0^{d+1} \setminus Q_{2R}. \end{cases}$$
 (5.7)

Noting that g_k is Hölder continuous in both t and x, the solvability follows from Theorem 1.1 and a regularization argument; see [2, 26]. We apply Corollary 4.6 to v_K with a scaling to get

$$[v_K]_{1+\alpha/\sigma,\alpha+\sigma;Q_{R/2}}$$

$$\leq C \Big(R^{-\sigma}[v_K]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times B_{2R}} + \sum_{j=2}^{\infty} 2^{-j\sigma} R^{-\sigma}[v_K]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times (B_{2^{j_R}}\setminus B_{2^{j-1}R})} \Big)$$

$$\leq C \Big(R^{-\sigma}[v_K]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times B_{2R}} \Big)$$

$$+\sum_{i=2}^{\infty} 2^{-j\sigma} R^{-\sigma} [u-p]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times(B_{2^{j}R}\setminus B_{2^{j-1}R})} \Big), \tag{5.8}$$

where in the last equality we used the fact that $v_K = g_K$ in $(-(2R)^{\sigma}, 0) \times B_{2R}^c$ and the Hölder norm of g_K is less than the Hölder norm of u - p.

By Lemma 5.3,

$$\begin{split} [u-p]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times B_{2^{j}R}} &\leq C2^{j(\sigma-\alpha/2)}R^{\sigma-\alpha}[u]_{\Lambda^{1}}^{t} \\ &+ C2^{j(\sigma-\alpha/2)}R^{\sigma-\alpha}[u]_{\sigma}^{*} + C2^{j(\sigma+1-\alpha)/2}R^{\sigma-\alpha}[Du]_{\frac{\sigma-1}{2}}^{t}, \end{split} \tag{5.9}$$

which together with (5.8) gives

$$[v_K]_{1+\alpha/\sigma,\alpha+\sigma;Q_{R/2}}$$

$$\leq C \left(R^{-\sigma}[v_K]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times B_{2R}} + R^{-\alpha}[u]_{\Lambda^1}^t + R^{-\alpha}[u]_{\sigma}^* + R^{-\alpha}[Du]_{\frac{\sigma-1}{2}}^t \right). \quad (5.10)$$

Next we estimate $w_K := g_K - v_K$, which is equal to $u - p - v_K$ in Q_{2R} by the choice of K. By (5.1) and (5.7), w_K satisfies

$$\begin{cases} \partial_t w_K \le \mathcal{M}^+ w_K + h_K + \sup_{a \in \mathcal{A}} \|f_a\|_{L_\infty} & \text{in } Q_{2R}, \\ \partial_t w_K \ge \mathcal{M}^- w_K + \hat{h}_K - \sup_{a \in \mathcal{A}} \|f_a\|_{L_\infty} & \text{in } Q_{2R}, \\ w_K = 0 & \text{in } \mathbb{R}_0^{d+1} \setminus Q_{2R}, \end{cases}$$

where

$$h_K := \mathcal{M}^+(u - p - g_K), \quad \hat{h}_K := \mathcal{M}^-(u - p - g_K).$$

By the dominated convergence theorem, it is not hard to see that

$$||h_K||_{L_{\infty}(Q_{2R})}, ||\hat{h}_K||_{L_{\infty}(Q_{2R})} \to 0 \text{ as } K \to \infty.$$

We then fix K large enough so that

$$||h_K||_{L_{\infty}(Q_{2R})} + ||\hat{h}_K||_{L_{\infty}(Q_{2R})} \le \sup_{a \in A} ||f_a||_{L_{\infty}}.$$

From Lemma 2.5, we have

$$||w_K||_{L_{\infty}(Q_{2R})} \le CR^{\sigma} \sup_{a \in \mathcal{A}} ||f_a||_{L_{\infty}}, \quad [w_K]_{\alpha/\sigma,\alpha;Q_{2R}} \le CR^{\sigma-\alpha} \sup_{a \in \mathcal{A}} ||f_a||_{L_{\infty}}, \quad (5.11)$$

where C depends on d, σ, λ , and Λ .

Now let q_K be the first-order Taylor expansion of v_K at the origin. Then by (5.10), for any $r \in (0, R/2)$,

$$||u - p - q_K||_{L_{\infty}(Q_r)} \le ||u - p - v_K||_{L_{\infty}(Q_r)} + ||v_K - q_K||_{L_{\infty}(Q_r)}$$

$$\le ||u - p - v_K||_{L_{\infty}(Q_r)} + Cr^{\sigma + \alpha}R^{-\sigma}[v_K]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times B_{2R}}$$

$$+ Cr^{\sigma + \alpha}R^{-\alpha}([u]_{\Lambda^1}^t + [u]_{\sigma}^* + [Du]_{\frac{\sigma-1}{\sigma}}^t).$$
(5.12)

Since $w_K = u - p - v_K$ in Q_{2R} , we plug (5.9) with j = 0, and (5.11) to (5.12) and obtain

$$||u - p - q_K||_{L_{\infty}(Q_r)} \le CR^{\sigma} \sup_{a \in \mathcal{A}} ||f_a||_{L_{\infty}} + Cr^{\sigma + \alpha}R^{-\alpha}([u]_{\Lambda^1}^t + [u]_{\sigma}^* + [Du]_{\frac{\sigma - 1}{\sigma}}^t).$$

Dividing both sides of the inequality above by r^{σ} , we have

$$r^{-\sigma} \| u - p - q_K \|_{L_{\infty}(Q_r)}$$

$$\leq C(R/r)^{\sigma} \sup_{a \in \mathcal{A}} \| f_a \|_{L_{\infty}} + C(r/R)^{\alpha} ([u]_{\Lambda^1}^t + [u]_{\sigma}^* + [Du]_{\frac{\sigma-1}{\sigma}}^t).$$

Set r = R/M, where $M \ge 2$ is a constant to be determined. Note that the center of the cylinder can be replaced by any point (t, x) in \mathbb{R}_0^{d+1} , i.e.,

$$r^{-\sigma} \| u - p - q_K \|_{L_{\infty}(Q_r(t,x))}$$

$$\leq C M^{\sigma} \sup_{a \in \mathcal{A}} \| f_a \|_{L_{\infty}} + C M^{-\alpha} ([u]_{\Lambda^1}^t + [u]_{\sigma}^* + [Du]_{\frac{\sigma-1}{\sigma}}^t),$$
(5.13)

which together with Lemma 5.5 implies

$$[u]_{\Lambda^{1}}^{t} + [u]_{\sigma}^{*} + [Du]_{\frac{\sigma-1}{\sigma}}^{t}$$

$$\leq C \sup_{r>0} \sup_{(t,x)\in\mathbb{R}_{0}^{d+1}} r^{-\sigma} \inf_{p\in\mathcal{P}_{1}} ||u-p||_{L_{\infty}(Q_{r}(t,x))}$$

$$\leq CM^{\sigma} \sup_{a\in A} ||f_{a}||_{L_{\infty}} + CM^{-\alpha} ([u]_{\Lambda^{1}}^{t} + [u]_{\sigma}^{*} + [Du]_{\frac{\sigma-1}{\sigma}}^{t}). \tag{5.14}$$

By taking M sufficiently large in (5.14) so that $CM^{-\alpha} < 1/2$, we obtain

$$[u]_{\Lambda^1}^t + [u]_{\sigma}^* + [Du]_{\frac{\sigma-1}{\sigma}}^t \le C \sup_{a \in \mathcal{A}} ||f_a||_{L_{\infty}}.$$

In the case when $\sigma = 1$, by Lemma 5.4 and (5.8),

$$[v_K]_{1+\alpha,1+\alpha;Q_{R/2}} \le C(R^{-1}[v_K]_{\alpha/\sigma,\alpha;(-R^{\sigma},0)\times B_{2R}} + R^{-\alpha}[u]_{\Lambda^1}).$$

Then by the same proof, similar to (5.13), we get

$$r^{-1} \| u - p - q_K \|_{L_{\infty}(Q_r(t,x))} \le CM \sup_{a \in \mathcal{A}} \| f_a \|_{L_{\infty}} + CM^{-\alpha} [u]_{\Lambda^1},$$

which together with Lemma 5.5 gives

$$[u]_{\Lambda^1} \le C \sup_{a \in \mathcal{A}} ||f_a||_{L_{\infty}}$$

by taking M sufficiently large. The theorem is proved.

Next, we provide a sketched proof of Corollary 1.3.

Proof of Corollary 1.3. We only consider the case $\sigma > 1$ and divide the proof into three steps.

Step 1. For $k=1,2,\ldots$, denote $Q^k:=Q_{1-2^{-k}}$. Let $\eta_k\in C_0^\infty(Q^{k+1})$ be a sequence of nonnegative smooth cutoff functions satisfying $\eta\equiv 1$ in Q^k , $|\eta|\leq 1$ in Q^{k+1} , $\|\partial_t^j D^i \eta_k\|_{L_\infty}\leq C2^{k(i+j)}$ for each $i,j\geq 0$. Set $v_k:=u\eta_k$ to be a smooth function and notice that in \mathbb{R}_0^{d+1} ,

$$\partial_t v_k = \eta_k \partial_t u + \partial_t \eta_k u = \inf_{a \in \mathcal{A}} (\eta_k L_a u + \eta_k f_a + \partial_t \eta_k u)$$
$$= \inf_{a \in \mathcal{A}} (L_a v_k + h_{ka} + \eta_k f_a + \partial_t \eta_k u),$$

where

$$h_{ka} = \eta_k L_a u - L_a v_k = \int_{\mathbb{R}^d} \xi_k(t, x, y) K_a(y) \, dy,$$

and

$$\xi_k(t, x, y) = u(t, x + y)(\eta_k(t, x + y) - \eta_k(t, x)) - y^T D\eta_k(t, x)u(t, x).$$

Obviously, we have

$$\sup_{a \in \mathcal{A}} \|\eta_k f_a\|_{L_{\infty}} \le \sup_{a \in \mathcal{A}} \|f_a\|_{L_{\infty}(Q_1)} \quad \text{and} \quad \|\partial_t \eta_k u\|_{L_{\infty}} \le C2^k \|u\|_{L_{\infty}(Q_1)}.$$

Step 2. We estimate the L_{∞} norm of h_{ka} . By the fundamental theorem of calculus,

$$\xi_k(t, x, y) = y^T \int_0^1 u(t, x + y) D\eta_k(t, x + sy) - u(t, x) D\eta_k(t, x) ds.$$

For $|y| \ge 2^{-k-3}$,

$$|\xi_k(t, x, y)| \le C2^k |y| ||u||_{L_{\infty}((-1, 0) \times \mathbb{R}^d)}.$$

For $|y| < 2^{-k-3}$, we can further write

$$\xi_k(t, x, y) = y^T \int_0^1 (u(t, x + y) - u(t, x)) D\eta_k(t, x + sy) + u(t, x) (D\eta_k(t, x + sy) - D\eta_k(t, x)) ds,$$

where the second term on the right-hand side is bounded by $C2^{2k}|y|^2|u(t,x)|$. To estimate the first term, we consider two cases: when $|x| \ge 1 - 2^{-k-2}$, because $|y| < 2^{-k-3}$, $\xi_k(t,x,y) \equiv 0$; when $|x| < 1 - 2^{-k-2}$, we have

$$\left| y^T \int_0^1 (u(t, x+y) - u(t, x)) D\eta_k(t, x+sy) \, ds \right| \le C2^k |y|^2 ||Du||_{L_{\infty}(Q^{k+3})}.$$

Hence for $|y| < 2^{-k-3}$,

$$|\xi_k(t,x,y)| \le C|y|^2 (2^{2k}|u(t,x)| + 2^k ||Du||_{L_{\infty}(Q^{k+3})}).$$

Combining with the case when $|y| > 2^{-k-3}$, we see that

$$||h_{ka}||_{L_{\infty}} \le C2^{\sigma k} ||u||_{L_{\infty}((-1,0)\times\mathbb{R}^d)} + C2^{(\sigma-1)k} ||Du||_{L_{\infty}(Q^{k+3})}.$$

Step 3. We apply Theorem 1.2 to v_k and use the bounds in the previous steps to obtain

$$[v_k]_{\Lambda^1}^t + [v_k]_{\sigma}^* + [v_k]_{\frac{\sigma-1}{\sigma}}^t \le C \sup_{a \in \mathcal{A}} ||f_a||_{L_{\infty}(Q_1)} + C2^k ||u||_{L_{\infty}(Q_1)} + C2^{\sigma k} ||u||_{L_{\infty}((-1,0) \times \mathbb{R}^d)} + C2^{(\sigma-1)k} ||Du||_{L_{\infty}(Q^{k+3})},$$

where C depends on d, λ , Λ , and σ , but independent of k. Since $\eta_k \equiv 1$ in Q^k , it follows that

$$[u]_{\Lambda^{1}(Q^{k})}^{t} + [u]_{\sigma;Q^{k}}^{*} + [Du]_{\frac{\sigma-1}{\sigma};Q^{k}}^{t}$$

$$\leq C2^{2k} ||u||_{L_{\infty}((-1,0)\times\mathbb{R}^{d})} + C2^{k} ||Du||_{L_{\infty}(Q^{k+3})} + C\sup_{a\in\mathcal{A}} ||f_{a}||_{L_{\infty}(Q_{1})}.$$
 (5.15)

By the interpolation inequality, for any $\varepsilon \in (0,1)$

$$||Du||_{L_{\infty}(Q^{k+3})} \le \varepsilon[u]_{\sigma;Q^{k+3}}^* + C\varepsilon^{-\frac{1}{\sigma-1}}||u||_{L_{\infty}(Q^{k+3})}.$$
 (5.16)

Set $N=1/(\sigma-1)$ and notice N>1. Combining (5.15) and (5.16) with $\varepsilon=C^{-1}2^{-k-10N}$, we obtain

$$[u]_{\Lambda^{1}(Q^{k})}^{t} + [u]_{\sigma;Q^{k}}^{*} + [Du]_{\frac{\sigma-1}{\sigma};Q^{k}}^{t} \le C2^{2k+(k+10N)N} ||u||_{L_{\infty}((-1,0)\times\mathbb{R}^{d})} + 2^{-10N} [u]_{\sigma;Q^{k+3}}^{*} + C \sup_{a\in\mathcal{A}} ||f_{a}||_{L_{\infty}(Q_{1})}.$$

Then we multiply 2^{-3kN} to both sides of the inequality above and get

$$\begin{split} &2^{-3kN}([u]_{\Lambda^{1}(Q^{k})}^{t}+[u]_{\sigma;Q^{k}}^{*}+[Du]_{\sigma;Q^{k}}^{t})\\ &\leq C2^{2(1-N)k}\|u\|_{L_{\infty}((-1,0)\times\mathbb{R}^{d})}+2^{-3N(k+3)-1}[u]_{\sigma;Q^{k+3}}^{*}+C2^{-3kN}\sup_{a}\|f_{a}\|_{L_{\infty}(Q_{1})}. \end{split}$$

We sum up the both sides of the inequality above and obtain

$$\begin{split} &\sum_{k=1}^{\infty} 2^{-3kN} ([u]_{\Lambda^{1}(Q^{k})}^{t} + [u]_{\sigma;Q^{k}}^{x} + [Du]_{\frac{\sigma-1}{\sigma};Q^{k}}^{t}) \\ &\leq C \sum_{k=1}^{\infty} 2^{2(1-N)k} \|u\|_{L_{\infty}((-1,0)\times\mathbb{R}^{d})} + C \sum_{k=1}^{\infty} 2^{-3kN} \sup_{a} \|f_{a}\|_{L_{\infty}(Q_{1})} \\ &+ \frac{1}{2} \sum_{k=4}^{\infty} 2^{-3kN} ([u]_{\Lambda^{1}(Q^{k})}^{t} + [u]_{\sigma;Q^{k}}^{*} + [Du]_{\frac{\sigma-1}{\sigma};Q^{k}}^{t}), \end{split}$$

which further implies that

$$\sum_{k=1}^{\infty} 2^{-3kN} ([u]_{\Lambda^{1}(Q^{k})}^{t} + [u]_{\sigma;Q^{k}}^{*} + [Du]_{\frac{\sigma-1}{\sigma};Q^{k}}^{t})$$

$$\leq C \|u\|_{L_{\infty}((-1,0)\times\mathbb{R}^{d})} + C \sup_{a} \|f_{a}\|_{L_{\infty}(Q_{1})},$$

where C depends on d, λ , Λ , and σ . In particular, by taking k=1, the corollary is proved.

APPENDIX

In the appendix, we first provide a sketch of the proof of Corollary 2.2.

Proof. By a scaling argument, we assume that r=1. Let $k \geq 1$ be a constant to be determined later. Set $\hat{\delta} = \delta/k$. Let $(t_0, x_0) \in \overline{Q_{\delta/2}}$ be such that $u(t_0, x_0) = \inf_{Q_{\delta/2}} u$. Since $\sigma \in (1, 2)$, we have $2^{-\sigma} \leq 1 - 4^{-\sigma}$. By a scaling and translation of the coordinates, we apply Proposition 2.1 to u in $Q_{\delta}(t_0, x_0)$ and obtain

$$\hat{\delta}^{-(\sigma+d)/\varepsilon} \|u\|_{L_{\varepsilon}(\hat{Q}_1)} \le C_1 \Big(\inf_{Q_{\hat{\delta}}(t_0, x_0)} u + C\hat{\delta}^{\sigma} \Big),$$

where $\hat{Q}_1 = Q_{\hat{\delta}}(t_1, x_0)$ and $t_1 = t_0 - (4^{\sigma} - 1)\hat{\delta}^{\sigma}$. For any $x_1 \in B_{\hat{\delta}/2}(x_0)$,

$$\begin{split} &\|u\|_{L_{\varepsilon}(\hat{Q}_1)} \geq \|u\|_{L_{\varepsilon}((t_1-\delta^{\sigma},t_1)\times B_{\hat{\delta}/2}(x_1))} \\ &\geq C_2 \hat{\delta}^{(\sigma+d)/\varepsilon} \inf_{(t_1-\delta^{\sigma},t_1)\times B_{\hat{\delta}/2}(x_1)} u \geq C_2 \hat{\delta}^{(\sigma+d)/\varepsilon} \inf_{Q_{\hat{\delta}}(t_1,x_1)} u, \end{split}$$

where $C_2 > 0$ depending only on d. Therefore,

$$\inf_{Q_{\hat{\delta}}(t_1,x_1)} u \le C_1/C_2 \Big(\inf_{Q_{\hat{\delta}}(t_0,x_0)} u + C\hat{\delta}^{\sigma}\Big).$$

Applying Proposition 2.1 again, we have

$$\hat{\delta}^{-(\sigma+d)/\varepsilon} \|u\|_{L_{\varepsilon}(\hat{Q}_{2})} \le C_{1} \Big(\inf_{Q_{\hat{\varepsilon}}(t_{1},x_{1})} u + C\hat{\delta}^{\sigma} \Big),$$

where $\hat{Q}_2 = Q_{\hat{\delta}}(t_2, x_1)$ and $t_2 = t_0 - 2(4^{\sigma} - 1)\hat{\delta}^{\sigma}$, and for any $x_2 \in B_{\hat{\delta}/2}(x_1)$,

$$\inf_{Q_{\hat{\delta}}(t_2, x_2)} u \le C_1 / C_2 \Big(\inf_{Q_{\hat{\delta}}(t_1, x_1)} u + C \hat{\delta}^{\sigma} \Big).$$

By induction, or any $x_{n-1} \in B_{(n-1)\hat{\delta}/2}(x_0) \cap B_1$,

$$\hat{\delta}^{-(\sigma+d)/\varepsilon} \|u\|_{L_{\varepsilon}(\hat{Q}_n)} \le C_3 \left(\inf_{Q_{\hat{\delta}}(t_0, x_0)} u + C \hat{\delta}^{\sigma} \right)$$

$$\le C_3 \left(u(t_0, x_0) + C \hat{\delta}^{\sigma} \right) = C_3 \left(\inf_{Q_{\delta/2}} u + C \hat{\delta}^{\sigma} \right),$$

where $\hat{Q}_n = Q_{\hat{\delta}}(t_n, x_{n-1})$, $t_n = t_0 - n(4^{\sigma} - 1)\hat{\delta}^{\sigma}$, and C_3 is a constant depending only on λ , Λ , d, and n. Notice that $|x_0| \leq \delta/2$, $t_0 \in [-(\delta/2)^{\sigma}, 0]$, and $\sigma > 1$. We can choose $k \geq 1$ in a suitable range depending only on σ_2 and δ , and then $n \leq [2k/\delta] + 1$, such that \hat{Q}_n runs through $(-\delta^{\sigma}, -\delta^{\sigma} + (4^{\sigma} - 1)\hat{\delta}^{\sigma}) \times B_1$. Finally, by applying Proposition 2.1 again and using a simple covering argument, we prove the corollary.

Finally, we give the proofs of Lemmas 5.1, 5.2, and 5.3.

Proof of Lemma 5.1. By mollification, it suffices to prove (5.2) assuming that $f \in C^{\alpha}((-1,0))$. Let $x,y \in (-1,0), y < x$, and h := x - y. When $h \ge 1/3$,

$$\frac{|f(x) - f(y)|}{h^{\alpha}} \le 2 \cdot 3^{\alpha} ||f||_{L_{\infty}((-1,0))}.$$

When h < 1/3, either x < -1/3 or y > -2/3. If x < -1/3, then $2x - y \in (x, 0)$ and

$$\frac{|f(x) - f(y)|}{h^{\alpha}} \le \frac{1}{2} \frac{|f(2x - y) + f(y) - 2f(x)|}{h^{\alpha}} + \frac{1}{2} \frac{|f(2x - y) - f(y)|}{h^{\alpha}}$$
$$\le \frac{3^{\alpha - 1}}{2} [f]_{\Lambda^{1}((-1,0))} + \frac{1}{2^{1 - \alpha}} [f]_{\alpha;(-1,0)}.$$

The case when y > -2/3 is similar. Therefore,

$$[f]_{\alpha;(-1,0)} \le 2 \cdot 3^{\alpha} ||f||_{L_{\infty}((-1,0))} + \frac{1}{2^{1-\alpha}} [f]_{\alpha;(-1,0)} + \frac{3^{\alpha-1}}{2} [f]_{\Lambda^{1}((-1,0))},$$
 which yields (5.2).

Proof of Lemma 5.2. First we consider the case when $\beta = 1$. Integrating by part and noting that η'' is an even function and $\int \eta'' = 0$, we obtain

$$\begin{split} \left| \partial_t^2 u^{(R)} \right| &= \left| \int_{\mathbb{R}} \left(2 \partial_t^2 u(t - R^{\sigma} s, x) - \partial_t^2 u(t - 2 R^{\sigma} s, x) \right) \eta(s - 1) \, ds \right| \\ &= \left| \int_{\mathbb{R}} R^{-2\sigma} \left(2 u(t - R^{\sigma} s, x) - \frac{1}{4} u(t - 2 R^{\sigma} s, x) \right) \eta''(s - 1) \, ds \right| \\ &= \left| \int_{\mathbb{R}} R^{-2\sigma} \left(u(t - R^{\sigma} s, x) + u(t - R^{\sigma} (2 - s), x) - 2 u(t - R^{\sigma}, x) \right) - \frac{1}{8} \left(u(t - 2 R^{\sigma} s, x) + u(t - 2 R^{\sigma} (2 - s), x) - 2 u(t - 2 R^{\sigma}, x) \right) \right) \eta''(s - 1) \, ds \\ &\leq C R^{-\sigma} [u]_{\Lambda^1}^t. \end{split}$$

For $\beta \in (0,1)$, when $r \geq R^{\sigma}$,

$$r^{-1-\beta}|u^{(R)}(t,x)+u^{(R)}(t-2r,x)-2u^{(R)}(t-r,x)|\leq r^{-\beta}[u]_{\Lambda^1}^t\leq R^{-\beta\sigma}[u]_{\Lambda^1}^t.$$

When $r \in (0, R^{\sigma})$, by (5.3) with $\beta = 1$,

$$\begin{split} r^{-1-\beta}|u^{(R)}(t,x) + u^{(R)}(t-2r,x) - 2u^{(R)}(t-r,x)| &\leq r^{1-\beta}[\partial_t u^{(R)}]_{1;\mathbb{R}_0^{d+1}}^t \\ &\leq C r^{1-\beta} R^{-\sigma}[u]_{\Lambda^1}^t \leq C R^{-\beta\sigma}[u]_{\Lambda^1}^t. \end{split}$$

From the above two inequalities, we immediately get (5.3).

Proof of Lemma 5.3. We first estimate the Hölder semi-norm in x. By the interpolation inequality,

$$[u-p]_{\alpha;(-R^{\sigma},0)\times B_{2^{j}R}}^{*}$$

$$\leq (2^{j}R)^{-\alpha}\|u-p\|_{L_{\infty}((-R^{\sigma},0)\times B_{2^{j}R})} + (2^{j}R)^{\sigma-\alpha}[u-p]_{\sigma;(-R^{\sigma},0)\times B_{2^{j}R}}^{*}.$$
(5.17)

Because p is linear,

$$[u-p]^*_{\sigma;(-R^{\sigma},0)\times B_{2j_R}} = [u]^*_{\sigma;(-R^{\sigma},0)\times B_{2j_R}}.$$
(5.18)

Since η has unit integral, we have

$$|u^{(R)}(t,x) - u(t,x)| = \left| \int_{\mathbb{R}} \left(2u(t - R^{\sigma}s, x) - u(t - 2R^{\sigma}s, x) - u(t, x) \right) \eta(s - 1) \, ds \right| \le CR^{\sigma} [u]_{\Lambda^{1}}^{t}.$$
(5.19)

Furthermore, for any $(t, x) \in (-R^{\sigma}, 0) \times B_{2^{j}R}$,

$$\begin{aligned} & \left| u^{(R)}(t,x) - p(t,x) \right| = \left| u^{(R)}(t,x) - u^{(R)}(0,0) - \partial_t u^{(R)}(0,0)t - x^T D u^{(R)}(0,0) \right| \\ & \leq \left| u^{(R)}(t,x) - u^{(R)}(t,0) - x^T D u^{(R)}(t,0) \right| \\ & + \left| u^{(R)}(t,0) - u^{(R)}(0,0) - \partial_t u^{(R)}(0,0)t \right| + \left| x^T D u^{(R)}(0,0) - x^T D u^{(R)}(t,0) \right| \\ & \leq (2^j R)^{\sigma} [u]_{\sigma}^* + R^{2\sigma} \left\| \partial_t^2 u^{(R)} \right\|_{L_{\infty}(-R^{\sigma},0) \times B_{2^j R}} + C 2^j R^{\sigma} \left[D u^{(R)} \right]_{\frac{\sigma-1}{\sigma}}^t. \end{aligned}$$

Using Lemma 5.2, we have

$$\left\| u^{(R)} - p \right\|_{L_{\infty}((-R^{\sigma},0) \times B_{2^{j}R})} \leq C(2^{j}R)^{\sigma} [u]_{\sigma}^{*} + C2^{j}R^{\sigma} [Du]_{\frac{\sigma-1}{\sigma}}^{t} + CR^{\sigma} [u]_{\Lambda^{1}}^{t},$$

which together with (5.19) implies that

$$||u - p||_{L_{\infty}((-R^{\sigma}, 0) \times B_{2^{j}R})} \le C(2^{j}R)^{\sigma}[u]_{\sigma}^{*} + C2^{j}R^{\sigma}[Du]_{\frac{\sigma-1}{\sigma}}^{t} + CR^{\sigma}[u]_{\Lambda^{1}}^{t}.$$
 (5.20)

We plug (5.18) and (5.20) in (5.17) and get (5.4).

Next we estimate the Hölder semi-norm in t. Obviously,

$$[u-p]_{\alpha/\sigma;(-R^{\sigma},0)\times B_{2^{j}R}}^{t} \leq [u-p]_{\alpha/\sigma;(-2^{j\sigma/2}R^{\sigma},0)\times B_{2^{j}R}}^{t}.$$

From Lemma 5.1 and scaling, we have

$$\begin{split} &[u-p]_{\alpha/\sigma;(-2^{j\sigma/2}R^{\sigma},0)\times B_{2^{j}R}}^{t} \\ &\leq C(2^{j/2}R)^{-\alpha}\|u-p\|_{L_{\infty}((-2^{j\sigma/2}R^{\sigma},0)\times B_{2^{j}R})} + C(2^{j/2}R)^{\sigma-\alpha}[u-p]_{\Lambda^{1}}^{t} \\ &\leq C(2^{j/2}R)^{-\alpha}\|u-p\|_{L_{\infty}((-2^{j\sigma/2}R^{\sigma},0)\times B_{2^{j}R})} + C(2^{j/2}R)^{\sigma-\alpha}[u]_{\Lambda^{1}}^{t}. \end{split}$$

We follow the proof of (5.20) to estimate

$$\begin{split} &\|u-p\|_{L_{\infty}((-2^{j\sigma/2}R^{\sigma},0)\times B_{2^{j}R})} \\ &\leq \|u-u^{(R)}\|_{L_{\infty}((-2^{j\sigma/2}R^{\sigma},0)\times B_{2^{j}R})} + \|u^{(R)}-p\|_{L_{\infty}((-2^{j\sigma/2}R^{\sigma},0)\times B_{2^{j}R})} \\ &\leq CR^{\sigma}[u]_{\Lambda^{1}}^{t} + C(2^{j}R)^{\sigma}[u]_{\sigma}^{*} + C2^{j\sigma}R^{\sigma}[u]_{\Lambda^{1}}^{t} + C2^{j(\sigma+1)/2}R^{\sigma}[Du]_{\sigma-1}^{t}. \end{split}$$

Therefore, we reach (5.5). The lemma is proved.

Proof of Lemma 5.4. We modify the proof of Lemma 5.3. Similar to Lemma 5.1, we have

$$[u-p]_{\alpha,\alpha;(-R,0)\times B_{2^{j}R}} \leq [u-p]_{\alpha,\alpha;Q_{2^{j}R}}$$

$$\leq (2^{j}R)^{-\alpha} ||u-p||_{L_{\infty}(Q_{2^{j}R})} + (2^{j}R)^{1-\alpha} [u-p]_{\Lambda_{1};Q_{2^{j}R}}$$

$$\leq (2^{j}R)^{-\alpha} ||u-p||_{L_{\infty}(Q_{2^{j}R})} + (2^{j}R)^{1-\alpha} [u]_{\Lambda_{1}}.$$
(5.21)

Since η and ζ have unit integral and ζ is radial, we have

$$\begin{aligned} & \left| u^{(R)}(t,x) - u(t,x) \right| \\ &= \left| \int_{\mathbb{R}^{d+1}} \left(2u(t-Rs,x-Ry) - u(t-2Rs,x-Ry) - u(t,x) \right) \eta(s-1) \zeta(y) \, dy \, ds \right| \\ &= \left| \int_{\mathbb{R}^{d+1}} \left(2u(t-Rs,x-Ry) - u(t-2Rs,x-Ry) - u(t,x-Ry) \right) \right. \\ &+ \frac{1}{2} \left(u(t,x+Ry) + u(t,x-Ry) - 2u(t,x) \right) \eta(s-1) \zeta(y) \, dy \, ds \right| \\ &\leq CR[u]_{\Lambda^{1}}. \end{aligned} \tag{5.22}$$

For any $(t, x) \in Q_{2^j R}$, similar to Lemma 5.2 with $\beta = \alpha/2$,

$$\left|u^{(R)}(t,x)-p(t,x)\right| \leq (2^{j}R)^{1+\alpha/2}[u^{(R)}]_{1+\alpha/2,1+\alpha/2;Q_{2^{j}R}} \leq 2^{j(1+\alpha/2)}R[u]_{\Lambda^{1}},$$

which together with (5.22) implies that

$$||u - p||_{L_{\infty}((-2^{j}R,0) \times B_{2^{j}R})} \le 2^{j(1+\alpha/2)}R[u]_{\Lambda^{1}}.$$
 (5.23)

We plug (5.23) in (5.21) and get (5.6). The lemma is proved.

ACKNOWLEDGMENTS

The authors would like to thank the referees for their careful review as well as many valuable comments and suggestions.

REFERENCES

- [1] Luis Caffarelli and Luis Silvestre. Regularity theory for fully nonlinear integro-differential equations. Comm. Pure Appl. Math., 62(5):597–638, 2009.
- [2] Luis Caffarelli and Luis Silvestre. The Evans-Krylov theorem for nonlocal fully nonlinear equations. Ann. of Math. (2), 174(2):1163–1187, 2011.
- [3] Luis Caffarelli and Luis Silvestre. Regularity results for nonlocal equations by approximation. Arch. Ration. Mech. Anal., 200(1):59–88, 2011.
- [4] Luis A. Caffarelli and Xavier Cabré. Fully nonlinear elliptic equations, volume 43 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, BI, 1995.
- 5] Chang Lara, Hctor and Dvila, Gonzalo. $C^{\sigma,\alpha}$ estimates for concave, non-local parabolic equations with critical drift. J. Integral Equations Appl. 28 (2016), no. 3, 373–394.
- [6] H. Chang-Lara and G. Davila. Hölder estimates for non-local parabolic equations with critical drift. J. Differential Equations, 260(5):4237–4284, 2016.

- [7] Héctor Chang-Lara and Gonzalo Dávila. Regularity for solutions of non local parabolic equations. Calc. Var. Partial Differential Equations, 49(1-2):139–172, 2014.
- [8] Héctor Chang-Lara and Gonzalo Dávila. Regularity for solutions of nonlocal parabolic equations II. J. Differential Equations, 256(1):130-156, 2014.
- [9] Héctor A. Chang-Lara, Dennis Kriventsov. Further Time Regularity for Non-Local, Fully Non-Linear Parabolic Equations. Comm. Pure Appl. Math. http://dx.doi.org/10.1002/cpa.21671
- [10] Rama Cont and Peter Tankov. Financial modelling with jump processes. Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [11] Hongjie Dong and Doyoon Kim. On L_p -estimates for a class of non-local elliptic equations. J. Funct. Anal., 262(3):1166–1199, 2012.
- [12] Hongjie Dong and Doyoon Kim. Schauder estimates for a class of non-local elliptic equations. Discrete Contin. Dyn. Syst., 33(6):2319–2347, 2013.
- [13] Lawrence C. Evans. Classical solutions of fully nonlinear, convex, second-order elliptic equations. Comm. Pure Appl. Math., 35(3):333–363, 1982.
- [14] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [15] Tianling Jin and Jingang Xiong. Schauder estimates for nonlocal fully nonlinear equations. Ann. Inst. H. Poincaré Anal. Non Linéaire, 33 (2016), no. 5, 1375–1407.
- [16] Tianling Jin and Jingang Xiong. Schauder estimates for solutions of linear parabolic integrodifferential equations. Discrete Contin. Dyn. Syst., 35(12):5977–5998, 2015.
- [17] Takashi Komatsu. Pseudodifferential operators and Markov processes. J. Math. Soc. Japan, 36(3):387–418, 1984.
- [18] N. V. Krylov. Boundedly inhomogeneous elliptic and parabolic equations in a domain. Izv. Akad. Nauk SSSR Ser. Mat., 47(1):75–108, 1983.
- [19] N. V. Krylov. Lectures on elliptic and parabolic equations in Hölder spaces, volume 12 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 1996.
- [20] R. Mikulevičius and H. Pragarauskas. On the Cauchy problem for certain integro-differential operators in Sobolev and Hölder spaces. *Liet. Mat. Rink.*, 32(2):299–331, 1992.
- [21] R. Mikulevicius and H. Pragarauskas. On the Cauchy problem for integro-differential operators in Hölder classes and the uniqueness of the martingale problem. *Potential Anal.*, 40(4):539–563, 2014.
- [22] Chenchen Mou. Interior regularity for nonlocal fully nonlinear equations with Dini continuous terms. J. Differential Equations, 260(11):7892–7922, 2016.
- [23] M. V. Safonov. Classical solution of second-order nonlinear elliptic equations. Izv. Akad. Nauk SSSR Ser. Mat., 52(6):1272–1287, 1328, 1988.
- [24] R. W. Schwab and L. Silvestre. Regularity for parabolic integro-differential equations with very irregular kernels. Anal. PDE 9 (2016), no. 3, 727–772.
- [25] Joaquim Serra. Regularity for fully nonlinear nonlocal parabolic equations with rough kernels. Calc. Var. Partial Differential Equations, 54(1):615–629, 2015.
- [26] Joaquim Serra. $C^{\sigma+\alpha}$ regularity for concave nonlocal fully nonlinear elliptic equations with rough kernels. Calc. Var. Partial Differential Equations, 54(4):3571–3601, 2015.
- (H. Dong) Division of Applied Mathematics, Brown University, 182 George Street, Providence, RI 02912, USA

E-mail address: Hongjie_Dong@brown.edu

(H. Zhang) Division of Applied Mathematics, Brown University, 182 George Street, Providence, RI 02912, USA

 $E\text{-}mail\ address{:}\ \texttt{Hong_Zhang@brown.edu}$