

ASPHERICITY OF A LENGTH FOUR RELATIVE GROUP PRESENTATION

Abd Ghafur Bin Ahmad, Muna A Al-Mulla and Martin Edjvet

Abstract

We consider the relative group presentation $\mathcal{P} = \langle G, \mathbf{x} | \mathbf{r} \rangle$ where $\mathbf{x} = \{x\}$ and $\mathbf{r} = \{xg_1xg_2xg_3x^{-1}g_4\}$. We show modulo a small number of exceptional cases exactly when \mathcal{P} is aspherical. If $H = \langle g_1^{-1}g_2, g_1^{-1}g_3g_1, g_4 \rangle \leq G$ then the exceptional cases occur when H is isomorphic to one of C_5, C_6, C_8 or $C_2 \times C_4$.

Mathematics Subject Classification: 20F05, 57M05.

Keywords: relative group presentations; asphericity.

1 Introduction

A *relative group presentation* is a presentation of the form $\mathcal{P} = \langle G, \mathbf{x} | \mathbf{r} \rangle$ where G is a group, \mathbf{x} a set disjoint from G and \mathbf{r} is a set of cyclically reduced words in the free product $G * \langle \mathbf{x} \rangle$ where $\langle \mathbf{x} \rangle$ denotes the free group on \mathbf{x} . If $G(\mathcal{P})$ denotes the group defined by \mathcal{P} then $G(\mathcal{P})$ is the quotient group $G * \langle \mathbf{x} \rangle / N$ where N denotes the normal closure in $G * \langle \mathbf{x} \rangle$ of \mathbf{r} . A relative presentation is defined in [2] to be *aspherical* if every spherical picture over it contains a *dipole*. If \mathcal{P} is aspherical then statements about $G(\mathcal{P})$ can be deduced and the reader is referred to [2] for a discussion of these; in particular torsion in $G(\mathcal{P})$ can easily be described.

We will consider the case when both \mathbf{x} and \mathbf{r} consists of a single element. Thus $\mathbf{r} = \{r\}$ where $r = x^{\varepsilon_1}g_1 \dots x^{\varepsilon_k}g_k$ where $g_i \in G$, $\varepsilon_i = \pm 1$ and $g_i = 1$ implies $\varepsilon_i + \varepsilon_{i+1} \neq 0$ for $1 \leq i \leq k$, subscripts mod k . If $k \leq 3$ or if $r \in \{xg_1xg_2xg_3xg_4, xg_1xg_2xg_3xg_4xg_5\}$ then, modulo some open cases, a complete classification of when \mathcal{P} is aspherical has been obtained in [1], [2], [7] and [11]. The case $r = (xg_1)^p(xg_2)^q(xg_3)^r$ for $p, q, r > 1$ was studied in [12] and $r = x^n g_1 x^{-1} g_2$ ($n \geq 4$) was studied in [5]. The authors of [10] used results from [7] in which $r = xg_1xg_2x^{-1}g_3$ to prove asphericity for certain LOG groups. In this paper we continue the study of asphericity and consider $r = xg_1xg_2xg_3x^{-1}g_4$. Observe that $r = 1$ if and only if $x^{-1}g_2^{-1}x^{-1}g_1^{-1}x^{-1}g_4^{-1}xg_3^{-1} = 1$ so replacing x^{-1} by x it follows that we can work modulo $g_1 \leftrightarrow g_2^{-1}$ and $g_3 \leftrightarrow g_4^{-1}$. A standard approach is to make the transformation $t = xg_1$ and then consider the subgroup H of G generated by the resulting coefficients. In our case r becomes $t^2g_1^{-1}g_2tg_1^{-1}g_3g_1t^{-1}g_4$ and so $H = \langle g_1^{-1}g_2, g_1^{-1}g_3g_1, g_4 \rangle$. One then usually proceeds according to either when H is non-cyclic or when H is cyclic. (Note that $\langle g_1^{-1}g_2, g_1^{-1}g_3g_1, g_4 \rangle$ is cyclic if and only if $\langle g_2g_1^{-1}, g_2g_4^{-1}g_2^{-1}, g_3^{-1} \rangle$ is cyclic.) The latter case seems to be the more complicated – indeed the open cases referred to in the above paragraph almost all involve H being cyclic. Our results reflect this difference in difficulty. When H is non-cyclic we obtain a complete answer except for the following case (modulo $g_1 \leftrightarrow g_2^{-1}, g_3 \leftrightarrow g_4^{-1}$) in which $H \cong C_2 \times C_4$:

$$(\mathbf{E}) \quad |g_3| = 2; |g_4| = 4; g_1^{-1}g_2 = 1; g_1^{-1}g_3g_1g_4 = g_4g_1^{-1}g_3g_1.$$

Theorem 1.1 *Let \mathcal{P} be the relative presentation*

$$\mathcal{P} = \langle G, x | xg_1xg_2xg_3x^{-1}g_4 \rangle,$$

where $g_i \in G$ ($1 \leq i \leq 4$), $g_3 \neq 1$, $g_4 \neq 1$ and $x \notin G$. Let $H = \langle g_1^{-1}g_2, g_1^{-1}g_3g_1, g_4 \rangle$ and assume that H is non-cyclic and the exceptional case (\mathbf{E}) does not hold. Then \mathcal{P} is aspherical if and only if (modulo $g_1 \leftrightarrow g_2^{-1}$, $g_3 \leftrightarrow g_4^{-1}$) none of the following conditions holds:

- (i) $|g_4| < \infty$ and $g_3g_1g_2^{-1} = 1$;
- (ii) $|g_1^{-1}g_2| < \infty$, $|g_3| = |g_4| = 2$ and $g_1^{-1}g_3g_1g_4 = g_2g_4^{-1}g_2^{-1}g_3^{-1} = 1$;
- (iii) $\frac{1}{|g_1^{-1}g_2|} + \frac{1}{|g_3|} + \frac{1}{|g_4|} + \frac{1}{|g_2g_4g_1^{-1}g_3^{-1}|} > 2$.

Now let H be cyclic. Before stating the theorem we make a list of exceptions (modulo $g_1 \leftrightarrow g_2^{-1}$, $g_3 \leftrightarrow g_4^{-1}$).

$$(\mathbf{E1}) \quad |g_4| = 5; g_1^{-1}g_2 = g_4^2; g_1^{-1}g_3g_1 = g_4^3.$$

$$(\mathbf{E2}) \quad |g_4| = 6; g_1^{-1}g_2 = 1; g_1^{-1}g_3g_1 = g_4^2.$$

$$(\mathbf{E3}) \quad |g_4| = 6; g_1^{-1}g_2 = 1; g_1^{-1}g_3g_1 = g_4^4.$$

$$(\mathbf{E4}) \quad |g_4| = 8; g_1^{-1}g_2 = 1; g_1^{-1}g_3g_1 = g_4^4.$$

Observe that $(\mathbf{E1})$ implies $H \cong C_5$; $(\mathbf{E2})$ and $(\mathbf{E3})$ imply $H \cong C_6$; and $(\mathbf{E4})$ implies $H \cong C_8$.

Theorem 1.2 *Let \mathcal{P} be the relative presentation*

$$\mathcal{P} = \langle G, x | xg_1xg_2xg_3x^{-1}g_4 \rangle$$

where $g_i \in G$ ($1 \leq i \leq 4$), $g_3 \neq 1$, $g_4 \neq 1$ and $x \notin G$. Let $H = \langle g_1^{-1}g_2, g_1^{-1}g_3g_1, g_4 \rangle$ be a cyclic group. Suppose that none of the exceptional conditions $(\mathbf{E1})$ – $(\mathbf{E4})$ holds. Then \mathcal{P} is aspherical if and only if either H is infinite or H is finite and (modulo $g_1 \leftrightarrow g_2^{-1}$, $g_3 \leftrightarrow g_4^{-1}$) none of the following conditions holds:

- (i) $g_3g_1g_2^{-1} = 1$;
- (ii) $g_3^{-1}g_1g_2^{-1} = g_2g_4^{-1}g_1^{-1}g_3^{-1} = 1$;
- (iii) $g_3^{-1}g_1g_2^{-1} = g_4g_2^{-1}g_1 = 1$;
- (iv) $|g_3| = 2; |g_4| = 2$;
- (v) $|g_3| = 2; g_1^{-1}g_3g_2g_4 = g_1^{-1}g_2g_4^{-2} = 1$;
- (vi) $|g_3| = 2; g_1^{-1}g_3g_2g_4 = (g_1^{-1}g_2)^2g_4^{-1} = 1$;
- (vii) $g_1^{-1}g_2 = 1; g_1^{-1}g_3g_1g_4^{\pm 1} = 1$;
- (viii) $g_1^{-1}g_2 = 1; |g_3| = 2; |g_4| = 3$;
- (ix) $g_1^{-1}g_2 = 1; 4 \leq |g_3| \leq 5; g_1^{-1}g_3^2g_1g_4$;
- (x) $g_1^{-1}g_2 = 1; |g_3| = 6; g_1^{-1}g_3^3g_1g_4$.

In Section 2 we discuss pictures and curvature; in Section 3 there are some preliminary results; Theorem 1.1 and Theorem 1.2 are proved in Sections 4 and 5.

Acknowledgement: the authors thank R M Thomas for his helpful comments concerning this paper.

2 Pictures

The definitions of this section are taken from [2]. The reader should consult [2] and [1] for more details.

A *picture* \mathbf{P} is a finite collection of pairwise disjoint discs $\{\Delta_1, \dots, \Delta_m\}$ in the interior of a disc D^2 , together with a finite collection of pairwise disjoint simple arcs $\{\alpha_1, \dots, \alpha_n\}$ embedded in the closure of $D^2 - \bigcup_{i=1}^m \Delta_i$ in such a way that each arc meets $\partial D^2 \cup \bigcup_{i=1}^m \Delta_i$ transversely in its end points. The *boundary* of \mathbf{P} is the circle ∂D^2 , denoted $\partial \mathbf{P}$. For $1 \leq i \leq m$, the *corners* of Δ_i are the closures of the connected components of $\partial \Delta_i - \bigcup_{j=1}^n \alpha_j$, where $\partial \Delta_i$ is the boundary of Δ_i . The *regions* of \mathbf{P} are the closures of the connected components of $D^2 - \left(\bigcup_{i=1}^m \Delta_i \cup \bigcup_{j=1}^n \alpha_j \right)$.

An *inner region* of \mathbf{P} is a simply connected region of \mathbf{P} that does not meet $\partial \mathbf{P}$. The picture \mathbf{P} is *non-trivial* if $m \geq 1$, is *connected* if $\bigcup_{i=1}^m \Delta_i \cup \bigcup_{j=1}^n \alpha_j$ is connected, and is *spherical* if it is non-trivial and if none of the arcs meets the boundary of D^2 . The number of edges in a region Δ is called the *degree* of Δ and is denoted by $d(\Delta)$. If \mathbf{P} is a spherical picture, the number of different discs to which a disc Δ_i is connected is called the *degree* of Δ_i , denoted by $\deg(\Delta_i)$.

With $\mathcal{P} = \langle G, \mathbf{x} | \mathbf{r} \rangle$ define the following labelling: each arc α_j is equipped with a normal orientation, indicated by a short arrow meeting the arc transversely, and labelled by an element of $\mathbf{x} \cup \mathbf{x}^{-1}$. Each corner of \mathbf{P} is oriented *anticlockwise* (with respect to D^2) and labelled by an element of G . If κ is a corner of a disc Δ_i of \mathbf{P} , then $W(\kappa)$ is the word obtained by reading in an anticlockwise order the labels on the arcs and corners meeting $\partial \Delta_i$ beginning with the label on the first arc we meet as we read the anticlockwise corner κ . If we cross an arc labelled x in the direction of its normal orientation, we read x , otherwise we read x^{-1} .

A picture \mathbf{P} is called a *picture over the relative presentation* \mathcal{P} if the above labelling satisfies the following conditions.

- (1) For each corner κ of \mathbf{P} , $W(\kappa) \in \mathbf{r}^*$, the set of all cyclic permutations of the members of $\mathbf{r} \cup \mathbf{r}^{-1}$ which begin with a member of \mathbf{x} .
- (2) If g_1, \dots, g_l is the sequence of corner labels encountered in a *clockwise* traversal of the boundary of an inner region Δ of \mathbf{P} , then the product $g_1 \dots g_l = 1$ in G . We say that $g_1 \dots g_l$ is the *label* of Δ .

A *dipole* in a labelled picture \mathbf{P} over \mathcal{P} consists of corners κ and κ' of \mathbf{P} together with an arc joining the two corners such that κ and κ' belong to the same region and such that if $W(\kappa) = Sg$ where $g \in G$ and S begins and ends with a member of $\mathbf{x} \cup \mathbf{x}^{-1}$, then $W(\kappa') = S^{-1}h^{-1}$. The picture \mathbf{P} is *reduced* if it does not contain a dipole. A relative

presentation \mathcal{P} is called *aspherical* if every connected spherical picture over \mathcal{P} contains a dipole.

The *star graph* \mathcal{P}^{st} of a relative presentation \mathcal{P} is a graph whose vertex set is $\mathbf{x} \cup \mathbf{x}^{-1}$ and edge set is \mathbf{r}^* . For $R \in \mathbf{r}^*$, write $R = Sg$ where $g \in G$ and S begins and ends with a member of $\mathbf{x} \cup \mathbf{x}^{-1}$. The initial and terminal functions are given as follows: $\iota(R)$ is the first symbol of S , and $\tau(R)$ is the inverse of the last symbol of S . The labelling function on the edges is defined by $\lambda(R) = g^{-1}$ and is extended to paths in the usual way. A non-empty cyclically reduced cycle (closed path) in \mathcal{P}^{st} will be called *admissible* if it has trivial label in G . In our case G embeds in $\langle G, t|r \rangle$ [8] so *every* region of a reduced picture over \mathcal{P} supports an admissible cycle in \mathcal{P}^{st} .

As described in the introduction we will consider spherical pictures over $\mathcal{P} = \langle G, t|r \rangle$ where $r = t^2 g_1^{-1} g_2 t g_1^{-1} g_3 g_1 t^{-1} g_4$. For ease of presentation we introduce the following notation: $a = 1$, $b = g_1^{-1} g_2$, $c = g_1^{-1} g_3 g_1$ and $d = g_4$ and consider $tatbtct^{-1}d$. Exception **(E)** and conditions (i)–(iii) of Theorem 1.1 can then be re-written as

$$\mathbf{(E)} \quad |c| = 2; |d| = 4; b = 1; cd = dc.$$

$$(i) \quad |d| < \infty \text{ and } cab^{-1} = 1;$$

$$(ii) \quad |a^{-1}b| < \infty, |c| = |d| = 2 \text{ and } a^{-1}cad = bd^{-1}b^{-1}c^{-1} = 1;$$

$$(iii) \quad \frac{1}{|a^{-1}b|} + \frac{1}{|c|} + \frac{1}{|d|} + \frac{1}{|bda^{-1}c^{-1}|} > 2.$$

The exceptions **(E1)**–**(E4)** and conditions (i)–(x) of Theorem 1.2 can be rewritten as

$$\mathbf{(E1)} \quad |d| = 5; b = d^2; c = d^3;$$

$$\mathbf{(E2)} \quad |d| = 6; b = 1; c = d^2;$$

$$\mathbf{(E3)} \quad |d| = 6; b = 1; c = d^4;$$

$$\mathbf{(E4)} \quad |d| = 8; b = 1; c = d^4;$$

$$(i) \quad cab^{-1} = 1;$$

$$(ii) \quad c^{-1}ab^{-1} = cadb^{-1} = 1;$$

$$(iii) \quad c^{-1}ab^{-1} = db^{-1}a = 1;$$

$$(iv) \quad |c| = |d| = 2;$$

$$(v) \quad |c| = 2; cbda^{-1} = a^{-1}bd^{-2} = 1;$$

$$(vi) \quad |c| = 2; cbda^{-1} = (a^{-1}b)^2 d^{-1} = 1;$$

$$(vii) \quad a^{-1}b = 1; cad^{\pm 1}a^{-1} = 1;$$

$$(viii) \quad a^{-1}b = 1; |c| = 2; |d| = 3;$$

$$(ix) \quad a^{-1}b = 1; 4 \leq |c| \leq 5; c^2ada^{-1} = 1;$$

$$(x) \quad a^{-1}b = 1; |c| = 6; c^3ada^{-1} = 1.$$




Figure 2.1: vertices and star graph

Let \mathbf{P} be a reduced connected spherical picture over \mathcal{P} . Then the vertices (discs) of \mathbf{P} are given by Figure 2.1(i) and (ii) where \bar{x} denotes x^{-1} for $x \in \{a, b, c, d\}$; and the star graph Γ is given by Figure 2.1(iii).

We make the following assumptions.

(A1) \mathbf{P} has a minimum number of vertices.

(A2) If $|c| = 2$ then, subject to (A1), \mathbf{P} has a maximum number of regions of degree 2 with label $c^{\pm 2}$.

Observe that (A1) implies that $c^{\varepsilon_1} w c^{\varepsilon_2}$, $d^{\varepsilon_1} w d^{\varepsilon_2}$ where $\varepsilon_1 = -\varepsilon_2 = \pm 1$ and $w = 1$ in G cannot occur as sublabels of a region. For otherwise a sequence of bridge moves [4] can be applied to produce a dipole which can then be deleted to obtain a picture with fewer vertices. Moreover if $|c| = 2$ then (A2) implies that $c^{\pm 2}$ cannot be a proper sublabel and $c^{\varepsilon} w c^{\varepsilon}$ where $\varepsilon = \pm 1$, $w = 1$ in G cannot be a sublabel of a region in \mathbf{P} . For otherwise bridge moves can be applied to increase the number of regions labelled $c^{\pm 2}$ while leaving the number of vertices unchanged.

To prove asphericity we adopt the approach of [6]. Let each corner in every region of \mathbf{P} be given an angle. The *curvature* of a vertex is defined to be 2π less the sum of the angles at that vertex. The curvature $c(\Delta)$ of a k -gonal region Δ of \mathbf{P} is the sum of all the angles of the corners of this region less $(k - 2)\pi$. Our method of associating angles is to give each corner at a vertex of degree d an angle $2\pi/d$. This way the vertices have zero curvature and we need consider only the regions. Thus if Δ is a k -gonal region of \mathbf{P} (a k -gon), denoted by $d(\Delta) = k$, and the degree of the vertices of Δ are d_i ($1 \leq i \leq k$) then

$$c(\Delta) = c(d_1, d_2, \dots, d_k) = (2 - k)\pi + 2\pi \sum_{i=1}^k (1/d_i).$$

In fact since each $d_i = 4$ ($1 \leq i \leq k$) we obtain

$$c(\Delta) = \pi(2 - k/2)$$

so if $d(\Delta) \geq 4$ then $c(\Delta) \leq 0$.

It follows from the fundamental curvature formula that $\sum c(\Delta) = 4\pi$ is where the sum is taken over all the regions Δ of \mathbf{P} . Our strategy to show asphericity will be to show that the positive curvature that exists in \mathbf{P} can be sufficiently compensated by the negative curvature. To this end, as a first step, we located the regions Δ of \mathbf{P} satisfying $c(\Delta) > 0$, that is, of positive curvature. For each such Δ we distribute all of $c(\Delta)$ to regions $\hat{\Delta}$ near Δ . For such regions $\hat{\Delta}$ of \mathbf{P} define $c^*(\hat{\Delta})$ to equal $c(\hat{\Delta})$ plus all the positive curvature $\hat{\Delta}$ receives in the distribution procedure mentioned above with the understanding that if $\hat{\Delta}$ receives no positive curvature then $c^*(\hat{\Delta}) = c(\hat{\Delta})$. Observe then that the total curvature of \mathbf{P} is at most $\sum(c^*(\hat{\Delta}))$ where the sum is taken over all regions $\hat{\Delta}$ of D that are not positively curved regions. Therefore to prove \mathcal{P} is aspherical it suffices to show that $c^*(\hat{\Delta}) \leq 0$ for each $\hat{\Delta}$.

Using the star graph Γ of Figure 2.1(iii) we can list the possible labels of regions of small degree (up to cyclic permutation and inversion).

$$d(\Delta) = 2 \Rightarrow l(\Delta) \in S_2 = \{c^2, d^2, a^{-1}b\}$$

$$d(\Delta) = 3 \Rightarrow l(\Delta) \in S_3 = \{c^3, cab^{-1}, c^{-1}ab^{-1}, d^3, db^{-1}a, d^{-1}b^{-1}a\}$$

Allowing each element in $S_2 \cup S_3$ to be either trivial or non-trivial yields 512 possibilities. This number can be reduced without any loss as follows.

1. Work modulo *T-equivalence*, that is, $a \leftrightarrow b^{-1}$, $c \leftrightarrow d^{-1}$.
2. Delete any combination that implies $c = 1$ or $d = 1$.
3. Delete any combination that yields a contradiction (for example $c^2 = 1$, $cab^{-1} = 1$, $c^{-1}ab^{-1} \neq 1$).
4. Delete any combination that yields $|d| < \infty$ and $cab^{-1} = 1$ or $|c| < \infty$ and $d^{-1}b^{-1}a = 1$ (see Lemma 3.1(i)).
5. When $H = \langle b, c, d \rangle$ is cyclic it can be assumed that H is finite (see Lemma 3.4(i)).

It can be readily verified that there remain 23 cases partitioned according to the existence in \mathbf{P} of regions of degree 2 and are listed below.

Case A: There are no regions of degree two.

- (A0) $|c| > 3$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.
- (A1) $|c| = 3$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.
- (A2) $|c| = 3$, $|d| = 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.
- (A3) $|c| > 3$, $|d| > 3$, $a^{-1}b \neq 1$, $cab^{-1} = 1$, $c^{-1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.
- (A4) $|c| > 3$, $|d| > 3$, $a^{-1}b \neq 1$, $cab^{-1} \neq 1$, $c^{-1}ab^{-1} = 1$, $d^{\pm 1}b^{-1}a \neq 1$.

- (A5) $|c| = 3, |d| > 3, a^{-1}b \neq 1, cab^{-1} = 1, c^{-1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (A6) $|c| = 3, |d| > 3, a^{-1}b \neq 1, cab^{-1} \neq 1, c^{-1}ab^{-1} = 1, d^{\pm 1}b^{-1}a \neq 1.$
- (A7) $|c| = 3, |d| > 3, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} \neq 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1.$
- (A8) $|c| = 3, |d| = 3, a^{-1}b \neq 1, cab^{-1} \neq 1, c^{-1}ab^{-1} = 1, d^{\pm 1}b^{-1}a \neq 1.$
- (A9) $|c| = 3, |d| = 3, a^{-1}b \neq 1, cab^{-1} \neq 1, c^{-1}ab^{-1} = 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1.$
- (A10) $|c| > 3, |d| > 3, a^{-1}b \neq 1, cab^{-1} \neq 1, c^{-1}ab^{-1} = 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1.$

Case B: Regions of degree two are possible.

- (B1) $|c| = 2, |d| > 3, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B2) $|c| = 2, |d| = 2, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B3) $|c| = 2, |d| = 3, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B4) $|c| = 2, |d| > 3, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} = 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B5) $|c| = 2, |d| > 3, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} \neq 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1.$
- (B6) $|c| = 2, |d| = 3, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} \neq 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1.$
- (B7) $|c| > 3, |d| > 3, a^{-1}b = 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B8) $|c| = 2, |d| > 3, a^{-1}b = 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B9) $|c| = 3, |d| > 3, a^{-1}b = 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B10) $|c| = 2, |d| = 2, a^{-1}b = 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B11) $|c| = 2, |d| = 3, a^{-1}b = 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$
- (B12) $|c| = 3, |d| = 3, a^{-1}b = 1, c^{\pm 1}ab^{-1} \neq 1, d^{\pm 1}b^{-1}a \neq 1.$

3 Preliminary results

We first exhibit spherical diagrams corresponding to some of the conditions of Theorems 1.1 and 1.2. **Note:** when drawing figures the discs (vertices) will often be represented by points; the edge arrows shown in Figure 2.1 will be omitted; and regions with label $c^{\pm 2}, d^{\pm 2}$ will be labelled simply by $c^{\pm 1}, d^{\pm 1}$.

Lemma 3.1

(a) *If any of the following conditions holds then \mathcal{P} fails to be aspherical:*

- (i) $|d| < \infty$ and $cab^{-1} = 1$;
- (ii) $|d| < \infty$ and $c^{-1}ab^{-1} = bd^{-1}a^{-1}c^{-1} = 1$;
- (iii) $|d| < \infty$ and $a^{-1}b = cad^{-1}b^{-1} = 1$.

(b) *If $bda^{-1}c^{-1} = 1$ and (f_1, f_2, f_3) is any of the following then \mathcal{P} fails to be aspherical.*

- (i) $(2, 2, < \infty)$;

- (ii) $(< \infty, 2, 2,)$;
- (iii) $(2, 3, k)$ $(3 \leq k \leq 5)$;
- (iv) $(3, 2, l)$ $(4 \leq l \leq 5)$;
- (v) $(k, 2, 3)$ $(3 \leq k \leq 5)$;

where $(f_1, f_2, f_3) = (|a^{-1}b|, |c|, |d|)$.

(c) If $a^{-1}b = 1$ and (f_1, f_2, f_3) is any of the following then \mathcal{P} fails to be aspherical.

- (i) $(2, 2, < \infty)$;
- (ii) $(2, < \infty, 2)$;
- (iii) $(2, l, 3)$ $(4 \leq l \leq 5)$;
- (iv) $(3, k, 2)$ $(3 \leq k \leq 5)$;
- (v) $(2, 3, k)$ $(3 \leq k \leq 5)$,

where $(f_1, f_2, f_3) = (|c|, |d|, |bda^{-1}c^{-1}|)$.

Proof (a)(i) If $|d| = n < \infty$ there is a spherical picture over \mathcal{P} consisting of two n -gons labelled d^n, d^{-n} together with $2n$ 3-gons labelled cab^{-1} . The sphere for $n = 5$ is shown in Figure 3.1(i).

(ii) If $|d| = n < \infty$ then there is a spherical picture over \mathcal{P} consisting of two regions with labels d^n, d^{-n} and between them two layers. Each layer has n 4-gons and n 3-gons with labels $cadb^{-1}, cba^{-1}$ and $bd^{-1}a^{-1}c^{-1}, c^{-1}ab^{-1}$. The sphere for $n = 6$ is shown in Figure 3.1(ii).

(iii) If $|d| = n < \infty$ then there is a spherical picture over \mathcal{P} consisting of a region labelled d^n , a region labelled c^{-n} and a layer of $2n$ regions between them. The layer consists of n 2-gons labelled $a^{-1}b$ and n 4-gons labelled $cad^{-1}b^{-1}$. The sphere for $n = 6$ is given by Figure 3.1(iii).

(b)(i) If $|d| = n < \infty$ then there is a spherical picture over \mathcal{P} consisting of two regions with label d^n and between them a layer consisting of n 4-gons with label $(a^{-1}b)^2$, n 2-gons with label c^{-2} and $2n$ 4-gons with label $ad^{-1}b^{-1}c$. The case $n = 5$ is shown in Figure 3.1(iv).

(ii) If $|a^{-1}b| = n < \infty$ then there is a spherical picture over \mathcal{P} consisting of two regions with label $(a^{-1}b)^n$ and between them a layer consisting of n 2-gons with label c^{-2} , n 2-gons with label d^2 and $2n$ 4-gons with label $ad^{-1}b^{-1}c$. The case $n = 3$ is shown in Figure 3.1(v).

(iii) If $k = 3$ a spherical picture is given by Figure 3.2(i) in which $A = ab^{-1}ab^{-1}$, $B = bda^{-1}c^{-1}$, $C = c^3$ and $D = d^{-3}$. If $k = 4$ a spherical picture is given by Figure 3.2(ii) in which $D = d^{-4}$. If $k = 5$ there is a spherical picture consisting of 30 A regions, 60 B regions, 20 C regions and 12 $D = d^{-5}$ regions, half of which is shown in Figure 3.2(iii).




Figure 3.1: spherical pictures

(iv) If $l = 4$ a spherical picture is given by Figure 3.3(i) in which $A = (ab^{-1})^3$, $B = bda^{-1}c^{-1}$, $C = c^2$ and $D = d^{-4}$. If $l = 5$ there is a spherical picture consisting of 20 A regions, 60 B regions, 30 C regions and 12 $D = d^{-5}$ regions, half of which is shown in Figure 3.3(ii).

(v) Spherical pictures for $k = 3$ and 4 are given in Figure 3.4(i) and (ii) in which $A = (ab^{-1})^3$ and $(ab^{-1})^4$ (respectively), $B = bda^{-1}c^{-1}$, $C = c^2$ and $D = d^{-3}$. If $k = 5$ there is a spherical picture consisting of 12 $A = (ab^{-1})^5$ regions, 60 B regions, 30 C regions and 20 D regions half of which is shown in Figure 3.4(iii).

(c)(i) If $|bda^{-1}c^{-1}| = n < \infty$ then there is a spherical picture over \mathcal{P} consisting of two $4n$ -gons each with label $(bda^{-1}c^{-1})^n$ together with n 2-gons with label c^2 , n 2-gons with label d^{-2} and $2n$ 2-gons with label ab^{-1} . The sphere for $n = 3$ is shown in Figure 3.5(i).

(ii) If $|d| = n < \infty$ then there is a sphere consisting of two n -gons with label $D = d^{-n}$ and between them a layer consisting of n 8-gons with label $B = (bda^{-1}c^{-1})^2$, $2n$ 2-gons with




Figure 3.2: spherical pictures




Figure 3.3: spherical pictures




Figure 3.4: spherical pictures

label $A = ab^{-1}$ and n 2-gons with label $C = c^2$. The sphere for $n = 4$ is shown in Figure 3.5(ii).

(iii) A spherical picture for $l = 4$ is given by Figure 3.5(iii) in which $A = ab^{-1}$, $B = (bda^{-1}c^{-1})^3$, $C = c^2$ and $D = d^{-4}$. If $l = 5$ there is a sphere consisting of 60 A regions, 20 B regions, 30 C regions and 12 $D = d^{-5}$ regions half of which is given by Figure 3.5(iv).

(iv) Spherical pictures for $k = 3$ and 4 are given by Figure 3.6(i) and (ii) in which $A = ab^{-1}$, $B = (bda^{-1}c^{-1})^2$, $C = c^3$ and $D = d^{-3}$ and d^{-4} (respectively). If $k = 5$ there is a spherical picture consisting of 60 A regions, 30 B regions, 20 C regions and 12 $D = d^{-5}$ regions half of which is given by Figure 3.6(iii).

(v) Spherical pictures for $k = 3$ and 4 are given by Figure 3.7(i) and (ii) in which $A = ab^{-1}$; $B = (bda^{-1}c^{-1})^3$ and $(bda^{-1}c^{-1})^4$ (respectively), $C = c^2$ and $D = d^{-3}$. If $k = 5$ there is a spherical picture consisting of 60 A regions, 12 $B = (bda^{-1}c^{-1})^5$ regions, 30 C regions and 20 D regions half of which is given by Figure 3.7(iii). \square

It follows from Theorem 1(2) in [1] that if $|t| < \infty$ in $G(\mathcal{P})$ then \mathcal{P} fails to be aspherical. We apply this fact in the proof of the next lemma.

Lemma 3.2 *If any of the following conditions hold then \mathcal{P} fails to be aspherical.*

- (i) $|a^{-1}b| < \infty$, $|c| = |d| = 2$ and $a^{-1}cad = bdb^{-1}c = 1$.
- (ii) $|d| < \infty$ and $c^{-1}ab^{-1} = db^{-1}a = 1$.




Figure 3.5: spherical pictures




Figure 3.6: spherical pictures






Figure 3.7: spherical pictures



(i)



(ii)

Figure 3.8: spherical pictures

- (iii) $|d| < \infty$ and $a^{-1}b = cada^{-1} = 1$.
- (iv) $c^2 = cbda^{-1} = a^{-1}bd^{-2} = 1$.
- (v) $c^2 = cbda^{-1} = (a^{-1}b)^2d^{-1} = 1$.
- (vi) $a^{-1}b = c^2 = d^3 = 1$ and $cada^{-1}cad^{-1}a^{-1} = 1$.
- (vii) $a^{-1}b = c^2ada^{-1} = 1$ and $4 \leq |c| \leq 5$.
- (viii) $a^{-1}b = c^3ada^{-1} = 1$ and $|c| = 6$.

Proof

- (i) It is enough to show that the group $G = \langle b, d, t \mid d^2 = b^k = 1, bd = db, t^2btdt^{-1}d = 1 \rangle$ has order $2k(3^{2k} - 1)$. Now $G = \langle d, t \mid d^2, t^{-2}d^{-1}td^{-1}t^{-1}dtdt^{-1}dt^2d^{-1}, (t^3dt^{-1}d)^k \rangle$ and $G/G' = \langle d, t \mid d^2 = t^{2k} = [d, t] = 1 \rangle$. Let \mathcal{K} denote the covering 2-complex associated with G' [3]. Then \mathcal{K} has edges $t_{0j}, t_{1j}, d_{j0}, d_{j1}$ ($1 \leq j \leq 2k$) and 2-cells $d_{j0}d_{j1}, t_{0j}d_{1-j}t_{1j}^{-1}d_{2-j}, t_{i1}t_{i2} \dots t_{i2k}$ where $1 \leq i \leq 2$ and $1 \leq j \leq 2k$ and the d subscripts are mod $2k$. Collapsing the maximal subtree whose edges are d_{j0} ($1 \leq j \leq 2k$), t_{0l} ($2 \leq l \leq 2k$) and using the lifts of d^2 shows that $G' = \langle t_{01}, t_{1j} \mid 1 \leq j \leq 2k \rangle$. Using the lifts of the second relator it is easily shown that $G' = \langle t_{01}, t_{11}, t_{12} \rangle$ where $t_{11}t_{01}^{-1} = t_{12}^{-3^{2k-1}}$ and, finally, using the lift of the third relator $(t^3dt^{-1}d)^k$ one can show that $G' = \langle t_{12} \mid t_{12}^r \rangle$ where $r = \frac{1}{2}(3^{2k} - 1)$. We omit the details.
- (ii) It is enough to show that $G = \langle d, x \mid d^k, t^2dtd^{-1}t^{-1}d \rangle$ has order $2k(1 + 4 + 4^2 + \dots + 4^{k-1})$. Now $G = \langle u, t \mid (ut^{-2})^k, tut^{-1}u^{-2} \rangle$ and $G/G' = \langle u, t \mid u = t^{2k} = 1 \rangle$. Let \mathcal{L} denote the covering complex associated with G' . Then \mathcal{L} has edges t_j, u_j ($1 \leq j \leq 2k$) and 2-cells $t_1t_2 \dots t_{2k}, u_j$ ($1 \leq j \leq 2k$). Collapsing the maximal tree whose edges are t_1, \dots, t_{k-1} implies $G' = \langle t_{2k}, u_j \mid 1 \leq j \leq 2k \rangle$. The lifts of $tut^{-1}u^{-2}$ yield the relators $u_l = u_1^{2^{l-1}}$ for $2 \leq l \leq 2k$ and $t_{2k}u_1t_{2k}^{-1}u_1^{-4^k}$. The lifts of $(ut^{-2})^k$ yield the relators $t_{2k}^{-1} = \prod_{i=0}^{k-1} u_{2i+1} = \prod_{i=1}^k u_{2i}$. It follows that $G' = \langle u_1 \mid u_1^r \rangle$ where $r = 1 + 4 + 4^2 + \dots + 4^{k-1}$.
- (iii) Here $r = t^3dt^{-1}d^{-1}$ and $|d| < \infty$ implies $|t| < \infty$.
- (iv) – (v) A spherical picture for (iv), (v) is shown in Figure 3.8(i), (ii) (respectively).
- (vi) – (viii) For these cases we use GAP [9]. For (vi), $r = t^3ct^{-1}d$ together with the conditions yields $|t| \leq 12$; for (vii) $r = t^3ct^{-1}c^{-2}$ and $|c| = 4, 5$ implies $|t| \leq 8, 10$ (respectively); and for (viii) $r = t^3ct^{-1}c^{-3}$ and $|c| = 6$ implies $|t| \leq 24$. \square

Lemma 3.3 *If any of the conditions (i)–(iii) of Theorem 1.1 or (i)–(x) of Theorem 1.2 holds then \mathcal{P} fails to be aspherical.*

Proof Consider Theorem 1.1. If (i) holds then \mathcal{P} fails to be aspherical by Lemma 3.1(a)(i). If (ii) holds then \mathcal{P} fails to be aspherical by Lemma 3.2(i). This leaves condition (iii). If $a^{-1}b \neq 1$ and $bda^{-1}c^{-1} \neq 1$ then (iii) does not hold; and if $a^{-1}b = bda^{-1}c^{-1} = 1$ then H is cyclic. Let $a^{-1}b = 1$. Since $(|c|, |d|, |bda^{-1}c^{-1}|)$ is T -equivalent to $(|d|, |c|, |bda^{-1}c^{-1}|)$ it can be assumed without any loss that $|c| \leq |d|$. The resulting ten cases are dealt with by Lemma 3.1(c). Let $bda^{-1}c^{-1} = 1$. Since $(|a^{-1}b|, |c|, |d|)$ is T -equivalent to $(|a^{-1}b|, |d|, |c|)$ it can again be assumed without any loss that $|c| \leq |d|$. The resulting ten cases are dealt with by Lemma 3.1(b). Now consider Theorem 1.2. If (i) holds then \mathcal{P} is aspherical by Lemma 3.1(a)(i); if (ii) holds then by Lemma 3.1(a)(ii); if (iii) holds then by Lemma 3.2(ii); if (iv) holds then by Lemma 3.2(i); if (v) holds then by Lemma 3.2(iv); if (vi) holds then by Lemma 3.2(v); if (vii) holds then by Lemmas 3.1(a)(iii) and 3.2(iii); if (viii) holds then by Lemma 3.2(vi); if (ix) holds then by Lemma 3.2(vii); and if (x) holds then by Lemma 3.2(viii). \square

A *weight function* α on the star graph Γ of Figure 2.1(iii) is a real-valued function on the set of edges of Γ . Denote the edge labelled a, b, c, d by e_a, e_b, e_c, e_d (respectively). The function α is *weakly aspherical* if the following two conditions are satisfied:

- (1) $\alpha(e_a) + \alpha(e_b) + \alpha(e_c) + \alpha(e_d) \leq 2$;
- (2) each admissible cycle in Γ has weight at least 2.

If there is a weakly aspherical weight function on Γ then \mathcal{P} is aspherical [2].

Lemma 3.4 *If any of the following conditions holds then \mathcal{P} is aspherical.*

- (i) $|c| = |d| = \infty$;
- (ii) $1 < |b| < \infty$ and $|d| = \infty$;
- (iii) $|c| < \infty, |d| < \infty$ and $|b| = \infty$.

Proof The following functions α are weakly aspherical.

- (i) $\alpha(e_a) = \alpha(e_b) = 1, \alpha(e_c) = \alpha(e_d) = 0$.
- (ii) $\alpha(e_a) = \alpha(e_b) = \frac{1}{2}, \alpha(e_c) = 1, \alpha(e_d) = 0$.
- (iii) $\alpha(e_a) = \alpha(e_b) = 0, \alpha(e_c) = \alpha(e_d) = 1$. \square

The following lemmas will be useful in later sections.

Lemma 3.5 *Let $d(\hat{\Delta}) = k$ where $\hat{\Delta}$ is a region of the spherical picture \mathbf{P} over \mathcal{P} .*

- (i) *If $\hat{\Delta}$ receives at most $\frac{\pi}{6}$ across each edge and $k \geq 6$ then $c^*(\hat{\Delta}) \leq 0$.*

- (ii) If $\hat{\Delta}$ receives at most $\frac{\pi}{6}$ across at most two-thirds of its edges, nothing across the remaining edges and $k \geq 5$ then $c^*(\hat{\Delta}) \leq 0$.
- (iii) If $\hat{\Delta}$ receives at most $\frac{\pi}{4}$ across each edge and $k \geq 8$ then $c^*(\hat{\Delta}) \leq 0$.
- (iv) If $\hat{\Delta}$ receives at most $\frac{\pi}{4}$ across at most two-thirds of its edges, nothing across the remaining edges and $k \geq 6$ then $c^*(\hat{\Delta}) \leq 0$.
- (v) If $\hat{\Delta}$ receives at most $\frac{\pi}{4}$ across at most half of its edges, nothing across the remaining edges and $k \geq 5$ then $c^*(\hat{\Delta}) \leq 0$.
- (vi) If $\hat{\Delta}$ receives at most $\frac{\pi}{2}$ across at most half of its edges, nothing across the remaining edges and $k \geq 7$ then $c^*(\hat{\Delta}) \leq 0$.
- (vii) If $\hat{\Delta}$ receives at most $\frac{\pi}{2}$ across at most three-fifths of its edges, nothing across the remaining edges and $k \geq 8$ then $c^*(\hat{\Delta}) \leq 0$.

Proof Recall that $c(\hat{\Delta}) = \pi(2 - \frac{k}{2})$. (i) $c^*(\hat{\Delta}) \leq c(\hat{\Delta}) + k \cdot \frac{\pi}{6} \leq 0$ for $k \geq 6$. (ii) $c^*(\hat{\Delta}) \leq c(\hat{\Delta}) + \frac{2k}{3} \cdot \frac{\pi}{6} \leq 0$ for $k \geq 6$. If $k = 5$ then $\hat{\Delta}$ receives at most $\frac{\pi}{6}$ across at most three edges and so $c^*(\hat{\Delta}) \leq \pi(2 - \frac{5}{2}) + 3 \cdot \frac{\pi}{6} = 0$. (iii) $c^*(\hat{\Delta}) \leq c(\Delta) + k \cdot \frac{\pi}{4} \leq 0$ for $k \geq 8$. (iv) $c^*(\hat{\Delta}) \leq c(\Delta) + \frac{2k}{3} \cdot \frac{\pi}{4} \leq 0$ for $k \geq 6$. (v) $c^*(\hat{\Delta}) \leq c(\Delta) + \frac{k}{2} \cdot \frac{\pi}{4} \leq 0$ for $k \geq 6$. If $k = 5$ then $\hat{\Delta}$ receives at most $\frac{\pi}{4}$ across at most two edges and $c^*(\hat{\Delta}) \leq \pi(2 - \frac{5}{2}) + 2 \cdot \frac{\pi}{4} = 0$. (vi) $c^*(\hat{\Delta}) \leq c(\Delta) + \frac{k}{2} \cdot \frac{\pi}{2} \leq 0$ for $k \geq 8$. If $k = 7$ then $\hat{\Delta}$ receives at most $\frac{\pi}{2}$ across at most three edges and $c^*(\hat{\Delta}) \leq \pi(2 - \frac{7}{2}) + 3 \cdot \frac{\pi}{2} = 0$. (vii) $c^*(\hat{\Delta}) \leq c(\Delta) + \frac{3k}{5} \cdot \frac{\pi}{2} \leq 0$ for $k \geq 10$. If $k = 9$ then $\hat{\Delta}$ receives at most $\frac{\pi}{2}$ across five edges and $c^*(\hat{\Delta}) \leq \pi(2 - \frac{9}{2}) + 5 \cdot \frac{\pi}{2} = 0$; if $k = 8$ then $\hat{\Delta}$ receives across at most four edges and $c^*(\hat{\Delta}) \leq \pi(2 - \frac{8}{2}) + 4 \cdot \frac{\pi}{2} = 0$. \square

Remark We will use the above lemmas as follows. Suppose that $\hat{\Delta}$ receives positive curvature across its edge e_i . If we know that it then never receives curvatures across e_{i-1} or across e_{i+1} then we can apply the “half” results; or if we know that it receives positive curvature across at most one of e_{i-1}, e_{i+1} then we can apply the “two-thirds” results.

Let $\hat{\Delta}$ be a region of \mathbf{P} and let e be an edge of $\hat{\Delta}$. If $\hat{\Delta}$ receives no curvature across e then e is called a *gap*; if at most $\frac{\pi}{6}$ then e is called a *two-thirds gap*; and if at most $\frac{\pi}{4}$ then e is called a *half gap*.

Lemma 3.6 (The Four Gaps Lemma) *If $\hat{\Delta}$ has a total of at least four gaps (in particular, four edges across which $\hat{\Delta}$ does not receive any curvature) and the most curvature that crosses any edge is $\frac{\pi}{2}$ then $c^*(\hat{\Delta}) \leq 0$.*

Proof It follows that $c^*(\hat{\Delta}) \leq \pi(2 - \frac{k}{2}) + (k - 4)\frac{\pi}{2} = 0$. \square

Checking the star graph shows that we will have the following LIST for the labels of regions of degree k where $k \in \{2, 3, 4, 5, 6, 7\}$:

If $d(\Delta) = 2$ then $l(\Delta) \in \{c^2, a^{-1}b, d^2\}$.

If $d(\Delta) = 3$ then $l(\Delta) \in \{c^3, cab^{-1}, c^{-1}ab^{-1}, db^{-1}a, d^{-1}b^{-1}a, d^3\}$.

If $d(\Delta) = 4$ then $l(\Delta) \in \{d^4, d^2a^{-1}b, d^2b^{-1}a, c^2ab^{-1}, c^2ba^{-1}, c^4, ab^{-1}ab^{-1}, d\{a^{-1}, b^{-1}\}\{c, c^{-1}\}\{a, b\}\}$.

If $d(\Delta) = 5$ then $l(\Delta) \in \{d^5, d^3a^{-1}b, d^3b^{-1}a, c^3ab^{-1}, c^3ba^{-1}, c^5, cab^{-1}ab^{-1}, cba^{-1}ba^{-1}, da^{-1}ba^{-1}b, db^{-1}ab^{-1}a, d^2\{a^{-1}, b^{-1}\}\{c, c^{-1}\}\{a, b\}, c^2\{a, b\}\{d, d^{-1}\}\{a^{-1}, b^{-1}\}\}$.

If $d(\Delta) = 6$ then $l(\Delta) \in \{d^6, d^4a^{-1}b, d^4b^{-1}a, c^4ab^{-1}, c^4ba^{-1}, c^6, ab^{-1}ab^{-1}ab^{-1}, d^2a^{-1}ba^{-1}b, d^2b^{-1}ab^{-1}a, c^2ab^{-1}ab^{-1}, c^2ba^{-1}ba^{-1}, d^3\{a^{-1}, b^{-1}\}\{c, c^{-1}\}\{a, b\}, d^2\{a^{-1}, b^{-1}\}\{c^2, c^{-2}\}\{a, b\}, d\{a^{-1}, b^{-1}\}\{c^3, c^{-3}\}\{a, b\}, c\{ab^{-1}, ba^{-1}\}\{c, c^{-1}\}\{ab^{-1}, ba^{-1}\}, d\{a^{-1}b, b^{-1}a\}\{d, d^{-1}\}\{a^{-1}b, b^{-1}a\}, c\{ab^{-1}a, ba^{-1}b\}\{d, d^{-1}\}\{a^{-1}, b^{-1}\}, c\{a, b\}\{d, d^{-1}\}\{a^{-1}ba^{-1}, b^{-1}ab^{-1}\}\}$.

Where, for example $d^2\{a^{-1}, b^{-1}\}\{c, c^{-1}\}\{a, b\}$ yields the eight labels $d^2a^{-1}c^{\pm 1}a$, $d^2a^{-1}c^{\pm 1}b$, $d^2b^{-1}c^{\pm 1}a$, $d^2b^{-1}c^{\pm 1}b$.

We will use the above LIST throughout the following sections often without explicit reference.

4 Proof of Case A

In this section we prove Theorems 1.1 and 1.2 for Case A, that is, we make the following assumptions:

$$|c| > 2, |d| > 2 \text{ and } a^{-1}b \neq 1.$$

This implies that $d(\Delta) \geq 3$ for each region Δ of the spherical diagram \mathcal{P} . If $d(\Delta) = 3$ then we will fix the names of the fifteen neighbouring regions Δ_i ($1 \leq i \leq 15$) of Δ as shown in Figure 4.1(i) and we use this notation throughout the section. We treat each of the cases (A0)–(A10) in turn.

(A0) $|c| > 3$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

In this case $d(\Delta) > 3$ for all regions Δ . Since the degree of each vertex equals 4 it follows that $c(\Delta) = (2 - d(\Delta))\pi + d(\Delta)\frac{2\pi}{4} \leq 0$ and so \mathcal{P} is aspherical.

(A1) $|c| = 3$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 3$ then $c(\Delta) = c(4, 4, 4) = \frac{\pi}{2}$, $l(\Delta) = c^3$ and Δ is given by Figure 4.1(ii). Observe that if $d(\Delta_i) = 4$ for $i \in \{1, 3, 5\}$ then $l(\Delta_i) = bd\omega$ and so $l(\Delta_i) \in \{bd^2a^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}\}$. (See the LIST of Section 3.) But $bdb^{-1}c^{\pm 1}$ implies $|d| = |c|$, a contradiction. Observe further that at most one of bd^2a^{-1} , $bda^{-1}c$, $bda^{-1}c^{-1}$ equals 1 otherwise there is a contradiction to $|c| = 3$ or $|d| > 3$. This leaves the following cases: (i) $bd^2a^{-1} \neq 1$; $bda^{-1}c^{\pm 1} \neq 1$; (ii) $bd^2a^{-1} = 1$; (iii) $bda^{-1}c = 1$; (iv) $bda^{-1}c^{-1} = 1$.

Consider case (i). Here $d(\Delta_i) > 4$ for $i \in \{1, 3, 5\}$ of Figure 4.1(ii), so add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each of $c(\Delta_i)$ across the bd edge as shown. Observe from Figure 4.1(ii) that Δ_1 , say, does




Figure 4.1: the region Δ and curvature distribution for Cases (A1,2,6,7,8) (B9,12)

not receive any curvature from Δ_2 or Δ_6 . Therefore if $\hat{\Delta}$ receives positive curvature, since $d(\hat{\Delta}) \geq 5$, it follows that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(ii).

Now consider case (ii), $bd^2a^{-1} = 1$. If $d(\Delta_i) > 4$ for at least two of the Δ_i where $i \in \{1, 3, 5\}$, say Δ_1 and Δ_3 , then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to each of $c(\Delta_1)$ and $c(\Delta_3)$ across the bd edge as in Figure 4.1(iii). Otherwise by symmetry it can be assumed without any loss of generality that $d(\Delta_1) = d(\Delta_3) = 4$. Then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ firstly to $c(\Delta_3)$ and then on to $c(\Delta_2)$ across the bd and ca edges as shown in Figure 4.1(iv). If $d(\Delta_5) > 4$ then add the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_5)$ across the bd edge; otherwise $d(\Delta_5) = 4$ and similarly add the $\frac{\pi}{4}$ firstly to $c(\Delta_5)$ then on to $c(\Delta_4)$. Now observe that in Figure 4.1(iii) Δ_1 does not receive positive curvature from Δ_2 or Δ_6 and in Figure 4.1(iv) that Δ_2 does not receive positive curvature from Δ_1 or Δ_9 . It follows that if $\hat{\Delta}$ receives positive curvature then $\hat{\Delta}$ receives curvature across at most half of its edges and so if $d(\hat{\Delta}) \geq 5$, it follows that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(v). This leaves the case when $d(\hat{\Delta}) = 4$ and $l(\hat{\Delta}) = cad^{-1}\omega$. Therefore $l(\hat{\Delta}) \in \{cad^{-1}b^{-1}, cad^{-1}a^{-1}\}$. In each case $l(\hat{\Delta})$ together with $bd^2a^{-1} = c^3 = 1$ implies $|d| = 3$, a contradiction.

Now consider (iii), $bda^{-1}c = 1$. As before, if $d(\Delta_1) > 4$ and $d(\Delta_3) > 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to each of $c(\Delta_1)$ and $c(\Delta_3)$ across the bd edge as in Figure 4.1(iii). If $d(\Delta_1) = d(\Delta_3) = 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ firstly to $c(\Delta_1)$ and then on to $c(\Delta_2)$ across the bd and ad edges as shown in Figure 4.2(i). The remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ is distributed either to Δ_5 when $d(\Delta_5) > 4$ or to Δ_6 via Δ_5 when $d(\Delta_5) = 4$. Now observe that in Figure 4.1(iii) Δ_1 does




Figure 4.2: curvature distribution for Cases (A1) and (A7)

not receive positive curvature from Δ_2 or Δ_6 ; and in Figure 4.2(i) that Δ_2 does not receive positive curvature from Δ_3 or Δ_8 . It follows that if the region $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and therefore if $d(\hat{\Delta}) \geq 5$ then $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(v). This leaves the case when $d(\hat{\Delta}) = 4$ and $l(\hat{\Delta}) = b^{-1}adw$. Therefore $l(\hat{\Delta}) = b^{-1}ad^2$ and H is cyclic. If this occurs then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_8)$ as shown in Figure 4.2(ii). If $d(\Delta_8) = 4$ then $l(\Delta_8) \in \{cad^{\pm 1}b^{-1}, cad^{\pm 1}a^{-1}, cab^{-1}c\}$ which, together with $b^{-1}add$, contradicts the (A1) assumptions. Also note that in Figure 4.2(ii), Δ_8 does not receive positive curvature from Δ_7 or from Δ_{16} and so if $\hat{\Delta}$ receives positive curvature it does so across at most half its edges and $c^*(\hat{\Delta}) \leq 0$.

Finally consider case (iv), $bda^{-1}c^{-1} = 1$. First assume that $b^2 \neq 1$. Again if $d(\Delta_1) > 4$ and $d(\Delta_3) > 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to each of $c(\Delta_1)$ and $c(\Delta_3)$ across the bd edge as in Figure 4.1(iii). If $d(\Delta_1) = d(\Delta_3) = 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ firstly to $c(\Delta_1)$ and then on to $c(\Delta_2)$ across the bd and ab^{-1} edges as shown in Figure 4.2(iii) and add the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to Δ_5 or Δ_6 as in the above. Now observe that in Figure 4.1(iii) Δ_1 does not receive positive curvature from Δ_2 or Δ_6 and in Figure 4.2(iii) that Δ_2 does not receive positive curvature from Δ_3 or Δ_8 . Thus if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges so $d(\hat{\Delta}) \geq 5$ implies $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(v). This leaves the case when $d(\hat{\Delta}) = 4$ and $l(\hat{\Delta}) = b^{-1}ab^{-1}\omega$. Therefore $l(\hat{\Delta}) = b^{-1}ab^{-1}a$ and there is a contradiction to $b^2 \neq 1$. Now assume that $b^2 = 1$. The distribution in this case is different in that we add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_j)$ for $j \in \{1, 3, 5\}$ as in Figure 4.1(ii) across the bd edge. If $d(\Delta_1) = 4$, say, then distribute the $\frac{\pi}{6}$ further on to $c(\Delta_7)$ across the d^{-2} edge




Figure 4.3: curvature distribution for Cases (A2,8) and (B3,6,11,12)

as shown in Figure 4.2(iv). Therefore if $\hat{\Delta}$ receives positive curvature and if $d(\hat{\Delta}) \geq 6$, it follows that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(i). This leaves the case when $4 \leq d(\hat{\Delta}) \leq 5$ and $l(\hat{\Delta}) \in \{bd\omega, d^{-2}\omega\}$. If $d(\hat{\Delta}) = 4$ then $l(\hat{\Delta}) = d^{-2}\omega$ and $l(\hat{\Delta}) \in \{d^{-4}, d^{-2}a^{-1}b, d^{-2}b^{-1}a\}$ therefore $d^4 = 1$ and there is a sphere by Lemma 3.1(b)(iii). So let $d(\hat{\Delta}) = 5$. If $\hat{\Delta}$ receives curvature across at most three edges then $c^*(\hat{\Delta}) \leq \pi(2 - \frac{5}{2}) + 3 \cdot \frac{\pi}{6} = 0$. Checking (the LIST) now shows that $c^*(\hat{\Delta}) \leq 0$ except when $l(\hat{\Delta}) = d^{-5}$ and in this case we obtain a sphere by Lemma 3.1(b)(iii).

In conclusion \mathcal{P} fails to be aspherical in this case if and only if $b^2 = bda^{-1}c^{-1} = 1$ and $|d| \in \{4, 5\}$. Note that if these conditions hold then H is non-cyclic for otherwise we would obtain $d^6 = 1$, a contradiction.

(A2) $|c| = 3$, $|d| = 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 3$ then Δ is given by Figures 4.1(ii) and 4.3(i). If $d(\Delta) = 4$ and $l(\Delta) \in \{bd\omega, ca\omega\}$ then $l(\Delta) \in S = \{bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}, cad^{\pm 1}a^{-1}, cad^{\pm 1}b^{-1}\}$ otherwise there is a contradiction to one of the assumptions. (Throughout this case unless otherwise stated this means one of the **(A2)** assumptions.)

The cases to be considered are (where for example case (ii) means $bda^{-1}c$ is the only member of S to equal 1): (i) $bda^{-1}c^{\pm 1} \neq 1$, $bdb^{-1}c^{\pm 1} \neq 1$, $cad^{\pm 1}a^{-1} \neq 1$, $cadb^{-1} \neq 1$; (ii) $bda^{-1}c = 1$; (iii) $bda^{-1}c^{-1} = 1$; (iv) $bdb^{-1}c = 1$; (v) $bdb^{-1}c^{-1} = 1$; (vi) $cada^{-1} = 1$; (vii) $cad^{-1}a^{-1} = 1$; (viii) $cadb^{-1} = 1$; (ix) $bdb^{-1}c = 1$, $cada^{-1} = 1$; (x) $bdb^{-1}c = 1$, $cad^{-1}a^{-1} = 1$; (xi) $bdb^{-1}c^{-1} = 1$, $cada^{-1} = 1$; (xii) $bdb^{-1}c^{-1} = 1$, $cad^{-1}a^{-1} = 1$.




Figure 4.4: curvature distribution for Case (A2)

Note that any other combination of these conditions gives a contradiction to one of the assumptions. Moreover, (ii) is T-equivalent to (viii); (iv) is T-equivalent to (vi); (v) is T-equivalent to (vii); and (x) is T-equivalent to (xi). So it remains to consider (i), (ii), (iii), (iv), (v), (ix), (x) and (xii).

Now let $c(\Delta) > 0$ and so $l(\Delta) \in \{c^3, d^3\}$. In cases (i), (ii), (iv) and (v) add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ for $i \in \{1, 3, 5\}$ as shown in Figures 4.1(ii) and 4.3(i). If $d(\Delta_i) > 4$ then no further distribution takes place. Suppose without any loss of generality that $d(\Delta_1) = 4$. This cannot happen in case (i); in case (ii) Δ_1 is given by Figure 4.3(ii) and so add the $\frac{\pi}{6}$ from $c(\Delta)$ to $c(\Delta_7)$ across the bd and bd^{-1} edges noting that $d(\Delta_7) > 4$ otherwise $l(\Delta_7) \in \{bd^{-2}a^{-1}, bd^{-1}a^{-1}c^{\pm 1}, bd^{-1}b^{-1}c^{\pm 1}\}$ which contradicts one of the assumptions; in case (iv) Δ_1 is given by Figure 4.3(iii) and so add the $\frac{\pi}{6}$ from $c(\Delta)$ to $c(\Delta_2)$ across the bd and ad edges noting that $d(\Delta_2) > 4$ otherwise $l(\Delta_2) \in \{ad^2b^{-1}, ada^{-1}c^{\pm 1}, ad^{-1}b^{-1}c^{\pm 1}\}$ which contradicts one of the assumptions or yields case (ix) or (x); and in case (v) Δ is given by Figure 4.3(iv) and so add the $\frac{\pi}{6}$ from $c(\Delta)$ to $c(\Delta_7)$ across the bd and $d^{-1}a^{-1}$ edges noting $d(\Delta_7) > 4$ otherwise $l(\Delta_7) \in \{d^{-2}a^{-1}b, d^{-1}a^{-1}c^{\pm 1}a, d^{-1}a^{-1}c^{\pm 1}b\}$ which contradicts one of the assumptions or yields case (xi) or (xii). Therefore if the region $\hat{\Delta}$ receives positive curvature then it receives $\frac{\pi}{6}$ across each edge and so if $d(\hat{\Delta}) \geq 6$ then $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(i). This leaves the case when $d(\hat{\Delta}) = 5$. After checking for vertex labels that contain the sublabeled (bd) , (ca) , (ad) and (bd^{-1}) corresponding to the edges crossed in Figures 4.1(ii) and 4.3(i)–(iv) we obtain $c^*(\hat{\Delta}) \leq \pi(2 - \frac{5}{2}) + 3 \cdot \frac{\pi}{6} = 0$. This completes cases (i), (ii), (iv) and (v).

Consider case (iii), $bda^{-1}c^{-1} = 1$. If $b^2 = 1$ then we obtain a sphere by Lemma 3.1(b)(iii). Note also that H is non-cyclic in this case otherwise we obtain $b = 1$, a contradiction. Suppose then that $b^2 \neq 1$. First let $l(\Delta) = c^3$. Add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ for $i \in \{1, 3, 5\}$ as in Figure 4.1(ii). If $d(\Delta_i) > 4$ then no further distribution takes place. Suppose that $d(\Delta_1) = 4$. Then add $\frac{\pi}{6}$ from $c(\Delta)$ to $c(\Delta_6)$ across the bd and ab^{-1} edges as shown in Figure 4.4(i) noting that $d(\Delta_6) > 4$, otherwise $l(\Delta_6) \in \{b^{-1}ab^{-1}a, b^{-1}ad^{\pm 2}\}$ which contradicts $b^2 \neq 1$ or one of the assumptions; and if $d(\Delta_3) = 4$ or $d(\Delta_5) = 4$ in Figure 4.1(ii) then similarly add $\frac{\pi}{6}$ to Δ_2 or Δ_4 . Secondly, let $l(\Delta) = d^3$. Add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ for $i \in \{1, 3, 5\}$ as in Figure 4.3(i). If $d(\Delta_i) > 4$ then no further distribution takes place.




Figure 4.5: curvature distribution for Case (A2)

Suppose without any loss of generality that $d(\Delta_1) = 4$. Then add $\frac{\pi}{6}$ from $c(\Delta)$ to $c(\Delta_2)$ across the ca and ba^{-1} edges as shown in Figure 4.4(ii), noting $d(\Delta_2) > 4$, otherwise $l(\Delta_2) \in \{ba^{-1}ba^{-1}, ba^{-1}c^{\pm 2}\}$ which contradicts $b^2 \neq 1$ or one of the assumptions. If $d(\Delta_3) = 4$ or $d(\Delta_5) = 4$ then similarly add $\frac{\pi}{6}$ to Δ_4 or Δ_6 . If $\hat{\Delta}$ receives positive curvature and $d(\hat{\Delta}) \geq 6$, it follows by Lemma 3.5(i) that $c^*(\hat{\Delta}) \leq 0$. It remains to check $d(\hat{\Delta}) = 5$. After checking for vertex labels that contain the sublabels $(bd), (ca), (b^{-1}a)$ and (ba^{-1}) corresponding to the edges crossed in Figures 4.1(ii), 4.3(i) and 4.4(i)–(ii) we obtain $c^*(\hat{\Delta}) \leq \pi(2 - \frac{5}{2}) + 3 \cdot \frac{\pi}{6} = 0$ or $l(\hat{\Delta}) \in \{bda^{-1}ba^{-1}, cab^{-1}ab^{-1}\}$ and this contradicts one of the assumptions. Therefore $c^*(\hat{\Delta}) \leq 0$.

Consider case (ix), $bdb^{-1}c = 1$ and $cada^{-1} = 1$. If $d(\Delta_i) > 4$ for at least two of Δ_i where $i \in \{1, 3, 5\}$, say Δ_1 and Δ_3 , then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to each of $c(\Delta_1)$ and $c(\Delta_3)$ across the bd and ca edges as shown in Figures 4.1(iii) and 4.4(iii). By symmetry it can be assumed that $d(\Delta_1) = d(\Delta_3) = 4$. The two possibilities are given in Figures 4.4(iv) and 4.5(i) and in both cases add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_2)$ across the bd, ad or ca, bd^{-1} edges as shown. If $d(\Delta_5) > 4$ then add the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_5)$; or if $d(\Delta_5) = 4$ then apply the above to Δ_1 and Δ_5 to distribute the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ similarly to $c(\Delta_6)$. Now observe that if Δ_1 receives positive curvature from Δ then it does not receive positive curvature from Δ_2 ; and if Δ_2 receives positive curvature from Δ (as in Figures 4.4(iv) and 4.5(i)) then it does not receive positive curvature from Δ_3 . It follows that if the region $\hat{\Delta}$ receives positive curvature then it does so across at most two-thirds of its edges and therefore if $\hat{\Delta}$ receives positive curvature and if $d(\hat{\Delta}) \geq 6$ then $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(iv). Note that $d(\Delta_2) > 4$ in Figures 4.4(iv) and 4.5(i) otherwise $l(\Delta_2) \in \{c^{-1}ada^{-1}, c^{-1}adb^{-1}, cbd^{-1}a^{-1}, cbd^{-1}b^{-1}\}$ which contradicts one of the assumptions. So there remains the case $d(\hat{\Delta}) = 5$ and $l(\hat{\Delta}) \in \{bd\omega, ca\omega, c^{-1}ad\omega, cbd^{-1}\omega\}$. Checking shows that in each case $c^*(\hat{\Delta}) \leq \pi(2 - \frac{5}{2}) + 2 \cdot \frac{\pi}{4} = 0$.

Consider (x), $bdb^{-1}c = 1$ and $cad^{-1}a^{-1} = 1$. First consider $l(\Delta) = d^3$. If at least two of the Δ_i where $i \in \{1, 3, 5\}$ have degree greater than four, say Δ_1 and Δ_3 , then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_1)$ and $c(\Delta_3)$ as shown in Figure 4.4(iii). So assume otherwise and without any loss




Figure 4.6: curvature distribution for Case (A2)

of generality let $d(\Delta_1) = d(\Delta_3) = 4$ as shown in Figure 4.5(ii) where $d(\Delta_2) > 4$ otherwise $l(\Delta_2) = a^{-1}bd^{-2}$ which contradicts $d^{-1}b^{-1}a \neq 1$. So add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_2)$ as shown in Figure 4.5(ii). If $d(\Delta_5) > 4$ in Figure 4.5(ii) add the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_5)$ otherwise use the same argument as above for Δ_1 and Δ_5 and add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_6)$. Now consider $l(\Delta) = c^3$. If at least two of the Δ_i where $i \in \{1, 3, 5\}$ have degree > 4 , say, Δ_1 and Δ_3 , then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_1)$ and $c(\Delta_3)$ as in Figure 4.1(iii). Suppose exactly two of the Δ_i have degree $= 4$, say, Δ_1 and Δ_3 . Add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_5)$. If $d(\Delta_2) > 4$ then add the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_2)$ as in Figure 4.4(iv). If $d(\Delta_2) = 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_{10})$ as in Figure 4.5(iii). If now $d(\Delta_{10}) = 4$ then $l(\Delta_{10}) = ba^{-1}ba^{-1}$ and so add the $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_9)$ as in Figure 4.6(i). Observe that $l(\Delta_9) = ad^{-1}d^{-1}w$ forces $d(\Delta_9) > 4$ otherwise there is a contradiction to $d^{-1}b^{-1}a \neq 1$. Finally suppose that $d(\Delta_i) = 4$ for $i \in \{1, 3, 5\}$. Then $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ is added to either Δ_2 , Δ_{10} or Δ_9 exactly as above; similarly $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ is added to Δ_6 , Δ_7 or Δ_{15} as shown in Figure 4.6(i).

Now observe that in Figures 4.4(iii) and 4.1(iii) Δ_1 does not receive positive curvature from Δ_2 ; in Figures 4.5(ii) and 4.4(iv) Δ_2 does not receive positive curvature from Δ_3 ; in Figure 4.5(iii) Δ_{10} does not receive positive curvature from Δ_{11} ; and in Figure 4.6(i) Δ_9 does not receive positive curvature from Δ_2 . Observe that if $\hat{\Delta}$ receives positive curvature then $d(\hat{\Delta}) \geq 5$. It follows from Lemma 3.5(iv) that if $d(\hat{\Delta}) \geq 6$ then $c^*(\hat{\Delta}) \leq 0$ so let $d(\hat{\Delta}) = 5$. If $\hat{\Delta}$ receives across at most two edges then $c^*(\hat{\Delta}) \leq 0$ so it remains to check if $\hat{\Delta}$ receives curvature from more than two edges. From the above we see that positive curvature is transferred across (ca) , (bd) , (bd^{-1}) , (ad) , (ba^{-1}) , (ad^{-1}) -edges. The only two labels that contain more than two such sublabeled and do not yield a contradiction are $a^{-1}ccad$ and $a^{-1}cadd$ as shown in Figures 4.6(ii)–(iii). Let $l(\Delta) = a^{-1}ccad = 1$ as in Figure 4.6(ii). Here Δ receives nothing from $\hat{\Delta}_1$ or $\hat{\Delta}_5$. If $d(\hat{\Delta}_2) > 3$ then Δ receives nothing from $\hat{\Delta}_2$ and so $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}_2) = 3$ then $d(\hat{\Delta}_3) > 3$ and Δ receives nothing from $\hat{\Delta}_3$ via $\hat{\Delta}_4$ as in Figure 4.4(iv) and again $c^*(\hat{\Delta}) \leq 0$. Let $l(\Delta) = a^{-1}cadd = 1$ as in Figure 4.6(iii). Here Δ receives nothing from $\hat{\Delta}_4$ or $\hat{\Delta}_5$. If $d(\hat{\Delta}_1) > 3$ then Δ receives nothing from $\hat{\Delta}_1$




Figure 4.7: curvature distribution for Cases (A2) and (A3)

and so $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}_1) = 3$ then $d(\hat{\Delta}_2) > 3$ and Δ receives nothing from $\hat{\Delta}_2$ via $\hat{\Delta}_3$ and again $c^*(\hat{\Delta}) \leq 0$.

Finally consider case (xii), $bdb^{-1}c^{-1} = cad^{-1}a^{-1} = 1$. Then $c = d$ and so $bdb^{-1}d^2 = 1$. If now $b^2 = 1$ then we obtain $bdbd^2 = 1$ and $H = \langle bd \rangle$ is cyclic. Assume first that H is non-cyclic so, in particular, $|b| > 2$. Add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ for $i \in \{1, 3, 5\}$ as in Figures 4.1(ii) and 4.3(i) across the bd and ca edges. If, say, $d(\Delta_1) > 4$ then no further distribution takes place. If $d(\Delta_1) = 4$ then the $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ is added to $c(\Delta_2)$ if $l(\Delta) = c^3$ across the bd and ab^{-1} edges, or to $c(\Delta_6)$ if $l(\Delta) = d^3$ across the ca and $a^{-1}b$ edges as shown in Figures 4.7(i)–(ii). Observe that $d(\Delta_2) > 4$ and $d(\Delta_6) > 4$. If $\hat{\Delta}$ receives positive curvature and $d(\hat{\Delta}) \geq 6$, it follows by Lemma 3.5(i) that $c^*(\hat{\Delta}) \leq 0$. It remains to check $d(\hat{\Delta}) = 5$. After checking for vertex labels that contain the sublabeled $(bd), (ca), (ab^{-1})$ and $(a^{-1}b)$ corresponding to the edges crossed in Figures 4.1(ii), 4.3(i) and 4.7(i)–(ii) it follows either that $\hat{\Delta}$ receives at most $3 \cdot \frac{\pi}{6}$ and so $c^*(\hat{\Delta}) \leq 0$ or $l(\hat{\Delta}) \in \{bda^{-1}ba^{-1}, cab^{-1}ab^{-1}\}$ which in each case yields a contradiction to H non-cyclic. Now assume that H is cyclic. If at least two of the Δ_i where $i \in \{1, 3, 5\}$ have degree greater than four, say Δ_1 and Δ_3 , then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to each of $c(\Delta_1)$ and $c(\Delta_3)$ as shown in Figures 4.1(iii) and 4.4(iii). By symmetry assume then that $d(\Delta_1) = d(\Delta_3) = 4$. The two possibilities are in Figures 4.7(iii) and 4.5(ii) and in each case add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_2)$ as shown. If $d(\Delta_5) > 4$ then add the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_5)$; or if $d(\Delta_5) = 4$ then similarly distribute the remaining $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ via Δ_5 to Δ_4 or Δ_6 . Now observe that if Δ_1 receives positive curvature from Δ it does not receive curvature from Δ_2 ; and if Δ_2 receives positive curvature from Δ it does not receive positive curvature from Δ_1 in Figure 4.7(iii) or from Δ_3 in Figure 4.5(ii).




Figure 4.8: curvature distribution for Cases (A4), (A6) and (A8)

It follows that if $d(\hat{\Delta}) \geq 6$ then $c^*(\Delta) \leq 0$ by Lemma 3.5(iv). Now $d(\Delta_2) > 4$ in Figures 4.7(iii) and 4.5(ii) so there remains the case $d(\hat{\Delta}) = 5$ and $d(\hat{\Delta}) \in \{bdw, caw, c^{-1}aw, bd^{-1}w\}$. But checking shows that in all cases $c^*(\hat{\Delta}) \leq \pi(2 - \frac{5}{2}) + 2 \cdot \frac{\pi}{4} = 0$.

In conclusion \mathcal{P} fails to be aspherical in this case if and only if $b^2 = bda^{-1}c^{-1} = 1$ (and H is non-cyclic).

(A3) $|c| > 3$, $|d| > 3$, $a^{-1}b \neq 1$, $cab^{-1} = 1$, $c^{-1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 3$ then $l(\Delta) = cab^{-1}$ and Δ is given by Figure 4.7(iv). Add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ across the d^2 edge. If Δ_5 receives no curvature across at least four edges then there are at least four *gaps* as defined in Section 3 and $c^*(\Delta_5) \leq 0$ by Lemma 3.6. This leaves the case when $l(\Delta_1) \in \{d^n, d^na^{-1}b, d^nb^{-1}a\}$. If $l(\Delta_5) = d^n$ then there is a sphere by Lemma 3.1(a)(i); and if $d^n(a^{-1}b)^{\pm 1} = 1$ then H is cyclic. Therefore \mathcal{P} is aspherical if and only if $|d| = \infty$.

(A4) $|c| > 3$, $|d| > 3$, $a^{-1}b \neq 1$, $cab^{-1} \neq 1$, $c^{-1}ab^{-1} = 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 3$ then $l(\Delta) = c^{-1}ab^{-1}$ and Δ is given by Figure 4.8(i). First assume that H is non-cyclic. Add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ across the db^{-1} edge and note that $l(\Delta_1) = db^{-1}\omega$ forces $d(\Delta_1) \geq 5$, otherwise $l(\Delta_1) \in \{db^{-1}ad, db^{-1}c^{\pm 1}a, db^{-1}c^{\pm 1}b\}$ which contradicts $|d| > 3$ or H non-cyclic. Observe also that Δ_1 does not receive curvature from Δ_2 or Δ_6 and so if $d(\Delta_1) \geq 7$, it follows that $c^*(\Delta_1) \leq 0$ by Lemma 3.5(vi); if $d(\Delta_1) = 5$ then $l(\Delta_1)$ contains at most one occurrence of $(db^{-1})^{\pm 1}$ and so $c^*(\Delta_1) \leq \pi(2 - \frac{5}{2}) + \frac{\pi}{2} = 0$; and if $d(\Delta_1) = 6$ then $l(\Delta_1)$ contains at most two occurrence of $(db^{-1})^{\pm 1}$ and so $c^*(\Delta_1) \leq \pi(2 - \frac{6}{2}) + 2 \cdot \frac{\pi}{2} = 0$. Now let H be cyclic. If $d(\Delta_1) > 4$ in Figure 4.8(i) then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$. Assume otherwise and $d(\Delta_1) = 4$. Then $l(\Delta_1) \in \{db^{-1}ad, db^{-1}c^{\pm 1}a, db^{-1}c^{\pm 1}b\}$ which yields a contradiction to the **(A4)** assumptions except when $l(\Delta_1) \in \{db^{-1}ad, db^{-1}ca\}$. If $db^{-1}ca = 1$ then there is a sphere by Lemma 3.1(a)(ii) so assume otherwise. Thus if $d(\Delta_1) = 4$ then $l(\Delta_1) = db^{-1}ad$. In this case as shown in Figure 4.8(ii) add $c(\Delta) = \frac{\pi}{2}$ firstly to $c(\Delta_1)$ then on to $c(\Delta_6)$. Observe that $d(\Delta_6) > 4$ otherwise $l(\Delta_6) \in \{da^{-1}bd, da^{-1}c^{\pm 1}a, da^{-1}cb, da^{-1}c^{-1}b\}$ which either contradicts $|c| > 3$, $|d| > 3$ or $d^{\pm 1}b^{-1}a \neq 1$, or implies the exceptional case **(E1)** when $da^{-1}c^{-1}b = 1$. Now observe that if Δ_1 receives positive curvature from Δ then it does not receive positive curvature from Δ_2 or Δ_6 in Figure 4.8(i); and if Δ_6 receives positive curvature from Δ then it does not receive positive curvature from Δ_5 or Δ_{15} in Figure




Figure 4.9: curvature distribution for Case (A6)

4.8(ii). Therefore if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges which implies that if $d(\hat{\Delta}) \geq 7$, then $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). If $d(\hat{\Delta}) = 5$ then checking shows that $l(\hat{\Delta})$ contains at most one occurrence of $(db^{-1})^{\pm 1}$ or $(da^{-1})^{\pm 1}$. It follows that $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}) = 6$ then checking shows that $l(\hat{\Delta})$ contains at most two occurrences of $(db^{-1})^{\pm 1}$ or $(da^{-1})^{\pm 1}$ and so $c^*(\hat{\Delta}) \leq 0$. In conclusion in this case \mathcal{P} fails to be aspherical if and only if $db^{-1}ca = 1$ (and so H is cyclic).

(A5) $|c| = 3$, $|d| > 3$, $a^{-1}b \neq 1$, $cab^{-1} = 1$, $c^{-1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $|d| < \infty$ then there is a sphere by Lemma 3.1(a)(i) and if $|d| = \infty$ then \mathcal{P} is aspherical by Lemma 3.4(ii).

(A6) $|c| = 3$, $|d| > 3$, $a^{-1}b \neq 1$, $cab^{-1} \neq 1$, $c^{-1}ab^{-1} = 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 3$ then Δ is given by Figure 4.1(ii) and 4.8(iii). First assume that either H is non-cyclic or H is cyclic but $b \neq d^{\pm 2}$. If $l(\Delta) = c^3$ then add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ for $i \in \{1, 3, 5\}$ across the bd edge. Note that if $d(\Delta_i) = 4$ then $l(\Delta_i) \in \{bdda^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}\}$ which contradicts $|d| > 3$ or $b \neq d^{\pm 2}$ and so $d(\Delta_i) \geq 5$. If $l(\Delta) = c^{-1}ab^{-1}$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_5)$ across the $d^{-1}a^{-1}$ edge and note that if $d(\Delta_5) = 4$ then $l(\Delta_5) \in \{d^{-1}a^{-1}bd^{-1}, d^{-1}a^{-1}c^{\pm 1}a, d^{-1}a^{-1}c^{\pm 1}b\}$ which contradicts $|d| > 3$ or $b \neq d^{\pm 2}$ and so $d(\Delta_5) \geq 5$. Observe now that Δ_1 does not receive curvature from Δ_2 or Δ_6 in Figure 4.1(ii); and Δ_5 does not receive curvature from Δ_4 or Δ_6 in Figure 4.8(iii). Therefore if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges which implies that if $d(\hat{\Delta}) \geq 7$, then $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). It remains to consider $5 \leq d(\hat{\Delta}) \leq 6$. If $d(\hat{\Delta}) = 5$ then checking shows that $l(\hat{\Delta})$ contains at most one occurrence of $(bd)^{\pm 1}$ or $(d^{-1}a^{-1})^{\pm 1}$. It follows that $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}) = 6$ then checking shows that $l(\hat{\Delta})$ contains at most two occurrences of $(bd)^{\pm 1}$ or $(d^{-1}a^{-1})^{\pm 1}$ and so $c^*(\hat{\Delta}) \leq 0$ and \mathcal{P} is aspherical.

Now let $b = d^{-2}$. This implies $c = d^2$ and $d^6 = 1$. It follows that if $d(\Delta) \in \{4, 5\}$ then $l(\Delta) \in \{bdda^{-1}, ccab^{-1}, cab^{-1}ab^{-1}, dda^{-1}c^{-1}a, ddb^{-1}ca, ddb^{-1}c^{-1}b\}$. If $l(\Delta) = c^{-1}ab^{-1}$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_1)$ and to $c(\Delta_5)$ as shown in Figure 4.8(iv). Let $l(\Delta) = c^3$. If at least two of $d(\Delta_i) > 4$, say $i = 1, 3$, then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_1)$ and to $c(\Delta_3)$ as shown in Figure 4.1(iii); if say $d(\Delta_1) = d(\Delta_3) = 4$ and $d(\Delta_5) > 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_5)$ and $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_2)$ via Δ_3 as shown in Figure 4.1(iv); and if $d(\Delta_i) = 4$ for $i \in \{1, 3, 5\}$ then add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each of $c(\Delta_j)$ for $j \in \{2, 4, 6\}$ similarly. Therefore if $\hat{\Delta}$ receives




Figure 4.10: curvature distribution for Cases (A6), (A7), (B5) and (B6)

positive curvature then $d(\hat{\Delta}) \geq 5$. Observe that Δ_1 does not receive positive curvature from Δ_2 in Figure 4.1(iii); Δ_1, Δ_5 does not receive positive curvature from Δ_6, Δ_4 respectively in Figure 4.8(iv); and Δ_2 does not receive positive curvature from Δ_9 in Figure 4.1(iv). It follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most two-thirds of its edges. Therefore $d(\hat{\Delta}) \geq 6$ implies $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(iv). Since there are at most two occurrences of $(ad)^{\pm 1}$, $(db^{-1})^{\pm 1}$, $(bd)^{\pm 1}$, $(cad^{-1})^{\pm 1}$ in each of $cab^{-1}ab^{-1}$, $dda^{-1}c^{-1}a$, $ddb^{-1}ca$, $ddb^{-1}c^{-1}b$ it follows that if $d(\hat{\Delta}) = 5$ then again $c^*(\hat{\Delta}) \leq 0$.

Finally let $b = d^2$. This implies $c = d^{-2}$ and $d^6 = 1$. It follows that if $d(\Delta) \in \{4, 5\}$ then $l(\Delta) \in \{d^2b^{-1}a, c^2ab^{-1}, bd^2b^{-1}c, bd^2a^{-1}c^{-1}, d^2a^{-1}ca, cab^{-1}ab^{-1}\}$. Let $c(\Delta) = c^3$. If at least one of Δ_i where $i \in \{1, 3, 5\}$ has degree > 5 , say $d(\Delta_1) > 5$, then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ as shown in Figure 4.9(i); or if $d(\Delta_i) = 5$ for $i \in \{1, 3, 5\}$ then add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each $c(\Delta_i)$ as shown in Figure 4.9(ii) in which $l(\Delta_i) \in \{bd^2b^{-1}c, bd^2a^{-1}c^{-1}\}$. Let $c(\Delta) = c^{-1}ab^{-1}$. If $d(\Delta_3) > 4$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_3)$ as in Figure 4.8(v); if $d(\Delta_3) = 4$ and $d(\Delta_4) > 5$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_4)$ via Δ_3 as in Figure 4.9(iii); and if $d(\Delta_3) = 4$ and $d(\Delta_4) = 5$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_5)$ as in Figure 4.9(iv). Finally if $d(\hat{\Delta}) = 5$, $l(\hat{\Delta}) = bd^2b^{-1}c$ and $\hat{\Delta}$ receives positive curvature from two neighbouring regions then add $\frac{\pi}{2}$ from $c^*(\hat{\Delta})$ to $c(\hat{\Delta}_1)$ as shown in Figure 4.10(i). Now observe that Δ_1 does not receive positive curvature from Δ_6 in Figure 4.9(i); Δ_1 , for example, does not receive from Δ_6 or Δ_2 in Figure 4.9(ii); Δ_3

does not receive from Δ_4 in Figure 4.8(v); Δ_4 does not receive from Δ_5 in Figure 4.9(iii); Δ_5 does not receive from Δ_4 or Δ_6 in Figure 4.9(iv); and $\hat{\Delta}_1$ does not receive from $\hat{\Delta}_2$ or $\hat{\Delta}_3$ in Figure 4.10(i). It follows that if $\hat{\Delta}$ receives positive curvature across two consecutive edges then $d(\hat{\Delta}) > 5$ and $\hat{\Delta}$ is given by Figure 4.10(ii) in which $\hat{\Delta}$ does not receive from $\hat{\Delta}_1$ or $\hat{\Delta}_2$. Since two sublabels of the form $(c^{-1}bd)^{\pm 1}$ must be separated by a sublabel of length at least 2 it follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most $\frac{3}{5}$ of its edges. Therefore if $d(\hat{\Delta}) \geq 8$ then $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vii); and if $l(\hat{\Delta})$ does not involve $(c^{-1}bd)^{\pm 1}$ then Lemma 3.5(vi) applies and $c^*(\hat{\Delta}) \leq 0$ for $d(\hat{\Delta}) \geq 7$. If $d(\hat{\Delta}) = 5$ then either $c^*(\hat{\Delta}) \leq 0$ or $\hat{\Delta}$ is given by Figure 4.10(i) and $c^*(\hat{\Delta}) - \frac{\pi}{2} \leq 0$. If $l(\hat{\Delta})$ does not involve $(c^{-1}bd)^{\pm 1}$ it remains to consider $d(\hat{\Delta}) = 6$. But checking shows that each candidate for $l(\hat{\Delta})$ contains at most two occurrences of $(bd)^{\pm 1}$, $(b^{-1}c)^{\pm 1}$ or $(cad)^{\pm 1}$ so $c^*(\hat{\Delta}) \leq 0$. Finally if $l(\hat{\Delta}) = c^{-1}bdw$ and $d(\hat{\Delta}) = 6$ then $l(\hat{\Delta}) = c^{-1}bd^2b^{-1}c^{-1}$ and $c^*(\hat{\Delta}) \leq 0$; or if $d(\hat{\Delta}) = 7$ then $l(\hat{\Delta}) \in \{c^{-1}bd^4b^{-1}, c^{-1}bda^{-1}bdb^{-1}, c^{-1}bda^{-1}bd^{-1}a^{-1}, c^{-1}bdb^{-1}ad^{-1}b^{-1}\}$ and $c^*(\hat{\Delta}) \leq c(\hat{\Delta}) + 3(\frac{\pi}{2}) = 0$. In conclusion \mathcal{P} is aspherical for this case.

(A7) $|c| = 3$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $db^{-1}a = 1$, $d^{-1}b^{-1}a \neq 1$.

If $d(\Delta) = 3$ then $l(\Delta) \in \{c^3, db^{-1}a\}$ and Δ is given by Figures 4.1(ii) and 4.10(iii). Note that if $l(\Delta_i) = bdw$, $c^{-1}aw$ and $d(\Delta_i) = 4$ then $l(\Delta_i) = bda^{-1}c^{\pm 1}$, $c^{-1}ad^{-1}b^{-1}$ (respectively) otherwise there is a contradiction to $|d| > 3$. First assume that $bda^{-1}c^{\pm 1} \neq 1$. If $l(\Delta) = c^3$ then add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ for $i \in \{1, 3, 5\}$ across the bd edge as shown in Figure 4.1(ii); and if $l(\Delta) = db^{-1}a$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_5)$ as in Figure 4.10(iii). Observe then that Δ_1 does not receive positive curvature from Δ_2 or Δ_6 in Figure 4.1(ii); and Δ_5 does not receive from Δ_4 or Δ_6 in Figure 4.10(iii). It follows that $c^*(\hat{\Delta}) \leq 0$ for $d(\hat{\Delta}) \geq 7$ by Lemma 3.5(vi). If $d(\hat{\Delta}) = 5$ then checking shows that either $l(\hat{\Delta})$ contains at most one occurrence of $(bd)^{\pm 1}$ or $(c^{-1}a)^{\pm 1}$ and so $c^*(\hat{\Delta}) \leq 0$ or $l(\hat{\Delta}) = c^{-1}ad^{-2}b^{-1}$. If $d(\hat{\Delta}) = 6$ then checking shows that $l(\hat{\Delta})$ contains at most two occurrences of $(bd)^{\pm 1}$ or $(c^{-1}a)^{\pm 1}$ and so $c^*(\hat{\Delta}) \leq 0$. Thus if $c^*(\hat{\Delta}) > 0$ then $\hat{\Delta}$ is given by Figure 4.10(iv) and add $c^*(\hat{\Delta}) = \frac{\pi}{6}$ to $c(\hat{\Delta}_1)$ as shown. Now $\hat{\Delta}_1$ does not receive positive curvature from $\hat{\Delta}_2$ or $\hat{\Delta}_3$ in Figure 4.10(iv) and the above statements still hold for Δ_i of Figure 4.1(ii) and Δ_5 of Figure 4.10(iii). Therefore if $\hat{\Delta}$ receives positive curvature it does so across at most half of its edges and so $c^*(\hat{\Delta}) \leq 0$ for $d(\hat{\Delta}) \geq 7$. Let $d(\hat{\Delta}) < 7$. If $l(\hat{\Delta})$ does not involve $a^{-1}b$ then checking shows that $c^*(\hat{\Delta}) \leq 0$ so assume $l(\hat{\Delta}) = a^{-1}bw$. But if now $d(\hat{\Delta}) < 6$ then $l(\hat{\Delta})$ together with $c^{-1}ad^{-2}b^{-1}$ contradicts $|d| = 9$; and if $d(\hat{\Delta}) = 6$ then the only labels with more than two occurrences of $(a^{-1}b)^{\pm 1}$, $(bd)^{\pm 1}$ or $(c^{-1}a)^{\pm 1}$ that do not yield a contradiction are $a^{-1}ba^{-1}cbd$, $a^{-1}bda^{-1}cb$ and $a^{-1}bd^{-1}b^{-1}ad$ in which case $c^*(\hat{\Delta}) \leq c(\Delta) + 2(\frac{\pi}{6}) + \frac{\pi}{2} < 0$. If $bda^{-1}c = 1$ then, since $db^{-1}a \leftrightarrow c^{-1}ab^{-1}$ and $bda^{-1}c \leftrightarrow a^{-1}c^{-1}bd^{-1}$, there is a sphere by Lemma 3.1(a)(ii). This leaves the case $bda^{-1}c^{-1} = 1$ which implies $d^6 = 1$ and $c = d^2$. If $l(\Delta) = db^{-1}a$ then again add $c(\Delta) = \frac{\pi}{2}$ for $c(\Delta_5)$ as in Figure 4.10(iii). Let $l(\Delta) = c^3$. If at least two $d(\Delta_i) \geq 5$, say Δ_1 and Δ_3 , then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_1)$ and to $c(\Delta_3)$ as in Figure 4.1(iii). If, say, $d(\Delta_1) = d(\Delta_3) = 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_2)$ via Δ_1 as in Figure 4.2(iii) and $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_5)$ if $d(\Delta_5) \geq 5$ or to $c(\Delta_6)$ via Δ_5 if $d(\Delta_5) = 4$ also. Observe that Δ_5 does not receive positive curvature from Δ_4 or Δ_6 in Figure 4.10(iii); Δ_1 does not receive any from Δ_6 in Figure 4.1(iii); and Δ_2 does not receive any from Δ_3 in




Figure 4.11: curvature distribution for Cases (A7) and (A8)

Figure 4.2(iii). It follows from Lemma 3.5(v) that if $d(\hat{\Delta}) \geq 7$ then $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}) = 6$ then checking the allowable labels shows that $l(\hat{\Delta})$ involves at most two occurrences of $(bd)^{\pm 1}$, $(c^{-1}a)^{\pm 1}$ or $(b^{-1}ab)^{\pm 1}$ and so $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}) = 5$ then $c^*(\hat{\Delta}) \leq 0$ except when $\hat{\Delta}$ is given by Figure 4.11(i) in which case add $c^*(\hat{\Delta}) = \frac{\pi}{4}$ to $c(\hat{\Delta}_1)$ across the bd edge as shown; and note that $\hat{\Delta}_1$ does not receive positive curvature from $\hat{\Delta}_2$ (and so none of the above is affected). If $c^*(\hat{\Delta}_1) \leq 0$ then we are done, otherwise $l(\hat{\Delta}_1) \in \{bda^{-1}c^{-1}, bda^{-1}c^2\}$. Let $l(\hat{\Delta}_1) = bda^{-1}c^{-1}$. Then add $c^*(\hat{\Delta}) = \frac{\pi}{4}$ to $\hat{\Delta}_2$ via $\hat{\Delta}_1$ as shown in Figure 4.11(ii) noting that $\hat{\Delta}_2$ does not receive positive curvature from the region Δ . It follows that if $c^*(\hat{\Delta}) > 0$ then $l(\hat{\Delta}) = b^{-1}ad^{-1}w$ and $d(\hat{\Delta}) \leq 6$. But this implies that $l(\hat{\Delta}) \in \{b^{-1}ad^{-1}a^{-1}bd, b^{-1}ad^{-1}a^{-1}ca\}$ and $c^*(\hat{\Delta}) \leq 0$. Finally if $\hat{\Delta}_1$ is given by Figure 4.11(iii) then add $c^*(\hat{\Delta}_1) = \frac{\pi}{4}$ to $\hat{\Delta}_3$ as shown and repeat the above argument. This procedure must terminate at a region $\hat{\Delta}_k$, say, where either $c^*(\hat{\Delta}_k) \leq 0$ or $d(\hat{\Delta}_k) = 4$ as in Figure 4.11(ii), in which case $c^*(\hat{\Delta}_k) = \frac{\pi}{4}$ is added to $c(\hat{\Delta}_{k+1})$ where $l(\hat{\Delta}_k) = b^{-1}ad^{-1}w$ and so $c^*(\hat{\Delta}_k) \leq 0$. In conclusion \mathcal{P} is aspherical if and only if $bda^{-1}c \neq 1$. (Note that if $bda^{-1}c = 1$ then H is cyclic.)

(A8) $|c| = 3$, $|d| = 3$, $a^{-1}b \neq 1$, $cab^{-1} \neq 1$, $c^{-1}ab^{-1} = 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 3$ then $l(\Delta) \in \{c^3, d^3, c^{-1}ab^{-1}\}$ and Δ is given by Figures 4.1(ii), 4.3(i) and 4.8(iii). If $l(\Delta) = c^3$ then add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ for $i \in \{1, 3, 5\}$ across the bd edge as shown in Figure 4.1(ii). Note that if $d(\Delta_i) = 4$ then $l(\Delta_i) \in \{bdda^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}\}$ which contradicts $|d| = 3$ or $d^{\pm 1}b^{-1}a \neq 1$ and so $d(\Delta_i) \geq 5$. If $l(\Delta) = c^{-1}ab^{-1}$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_5)$ across the $d^{-1}a^{-1}$ edge as shown in Figure 4.8(iii) and note that

if $d(\Delta_5) = 4$ then $l(\Delta_5) \in \{d^{-1}a^{-1}bd^{-1}, d^{-1}a^{-1}c^{\pm 1}a, d^{-1}a^{-1}c^{\pm 1}b\}$ which again contradicts $|d| = 3$ or $d^{\pm 1}b^{-1}a \neq 1$ and so $d(\Delta_5) \geq 5$. If $l(\Delta) = d^3$ then there are three subcases. First assume that $d(\Delta_i) \geq 5$ for $i \in \{1, 3, 5\}$. Then add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each $c(\Delta_i)$ across the ca edge as shown in Figures 4.3(i). Now suppose that at least two of the Δ_i have degree four, without loss of generality Δ_1 and Δ_3 . Then $l(\Delta_1) = l(\Delta_3) = cab^{-1}c$ (otherwise we will have a contradiction to $d^{\pm 1}b^{-1}a \neq 1$) which in turn forces $l(\Delta_2) = c^{-1}bd\omega$ and so $d(\Delta_2) > 4$ otherwise $l(\Delta_2) \in \{c^{-1}bda^{-1}, c^{-1}bdb^{-1}\}$ which contradicts $|d| = 3$ or $d^{\pm 1}b^{-1}a \neq 1$. In this case add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_2)$ and $c(\Delta_5)$ across the ca and bd edges as shown in Figure 4.11(iv) and if $d(\Delta_5) = 4$ then in the same way add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to $c(\Delta_6)$. This leaves the case where exactly one of Δ_i has degree four and without loss assume $d(\Delta_1) = 4$. Then add $\frac{1}{2}c(\Delta) = \frac{\pi}{4}$ to each of $c(\Delta_3)$ and $c(\Delta_5)$ as shown in Figure 4.11(v).

Observe that in Figure 4.1(ii) Δ_1 does not receive any positive curvature from Δ_2 or Δ_6 ; in Figure 4.3(i) Δ_1 does not receive positive curvature from Δ_2 ; in Figure 4.8(iii) Δ_5 does not receive any positive curvature from Δ_6 ; in Figure 4.11(iv) Δ_2 does not receive any positive curvature from Δ_3 or Δ_8 and Δ_5 does not receive any positive curvature from Δ_6 ; and in Figure 4.11(v) Δ_3, Δ_5 (respectively) does not receive any positive curvature from Δ_4, Δ_6 (respectively). Let $\hat{\Delta}$ receive positive curvature and first suppose that $l(\hat{\Delta})$ does not involve $(ad)^{\pm 1}$. It follows that $\hat{\Delta}$ receives positive curvature across at most half of its edges and so, since $d(\hat{\Delta}) \geq 5$, $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(v). Now assume that $l(\hat{\Delta}) = ad\omega$ and put $d(\hat{\Delta}) = k$. Since positive curvature crosses $(bd), (ca)$ and (ad) edges only it follows that if $l(\hat{\Delta})$ involves at least two occurrences of $(ad)^{\pm 1}$ then there are at least four gaps and $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.6, so assume otherwise. Moreover, $l(\hat{\Delta}) = ad\omega$ forces $\hat{\Delta}$ to have at least two gaps and so $c^*(\hat{\Delta}) \leq \pi(2 - \frac{k}{2}) + \frac{\pi}{2} + (k-3)\frac{\pi}{4}$ which implies that if $k \geq 7$ then $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}) = 5$ then $l(\hat{\Delta}) \in \{ad^3b^{-1}, adb^{-1}ab^{-1}, ad^2a^{-1}c^{\pm 1}, ad^2b^{-1}c^{\pm 1}, ada^{-1}c^{\pm 2}, adb^{-1}c^{\pm 2}\}$ which contradicts $|b| = 3$ or $d^{\pm 1}b^{-1}a \neq 1$. If $d(\hat{\Delta}) = 6$ then checking shows that $\hat{\Delta}$ receives curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$. In conclusion in this case \mathcal{P} is aspherical.

(A9) $|c| = 3, |d| = 3, a^{-1}b \neq 1, cab^{-1} \neq 1, c^{-1}ab^{-1} = 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1$. Since H is cyclic \mathcal{P} is aspherical by Lemma 3.2(ii).

(A10) $|c| > 3, |d| > 3, a^{-1}b \neq 1, cab^{-1} \neq 1, c^{-1}ab^{-1} = 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1$. Since H is cyclic, $|d| < \infty$ and so \mathcal{P} is aspherical by Lemma 3.2(ii).

It follows from **(A0)**–**(A10)** above that either \mathcal{P} is aspherical or modulo T -equivalence one of the conditions from Theorem 1.1(i), (iii) or Theorem 1.2(i), (ii), (iii) is satisfied and so Theorems 1.1 and 1.2 are proved for Case A.

5 Proof of Case B

In this section we prove Theorems 1.1 and 1.2 for Case B, that is, we make the following assumption: at least one of $c^2, d^2, a^{-1}b$ equals 1 in H .




Figure 5.1: the region Δ and curvature distribution for Cases (B1,2,3,5,6,8,10)

If $d(\Delta) = 2$ then we will fix the names of the four neighbouring regions Δ_i ($1 \leq i \leq 4$) of Δ as shown in Figure 5.1(i) and we use this notation throughout the section.

Remark Recall that if $c^2 = 1$ then $c^{\pm 2}$ cannot be a proper sublabel. This fact will be used often without explicit reference.

We treat each of the cases (B1) to (B12) in turn.

(B1) $|c| = 2$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 2$ then Δ is given by Figure 5.1(ii). Observe that if $d(\Delta_i) = 4$ for $i \in \{1, 2\}$ then $l(\Delta_i) = \{bdda^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}\}$. But $bdb^{-1}c^{\pm 1} = 1$ implies $|d| = |c|$, a contradiction. Observe further that at most one of $bd^2a^{-1}, bda^{-1}c^{\pm 1}$ equals 1 otherwise there is a contradiction to $|d| > 3$. This leaves the following cases:

- (i) $bd^2a^{-1} \neq 1$, $bda^{-1}c^{\pm 1} \neq 1$;
- (ii) $bd^2a^{-1} = 1$, $bda^{-1}c^{\pm 1} \neq 1$;
- (iii) $bda^{-1}c^{\pm 1} = 1$, $bd^2a^{-1} \neq 1$.

- (i) In this case $d(\Delta_i) > 4$ for Δ_i ($1 \leq i \leq 2$) of Figure 5.1(ii) so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to each of $c(\Delta_i)$ ($1 \leq i \leq 2$). Observe from Figure 5.1(ii) that Δ_i does not receive positive curvature from Δ_j for $j \in \{3, 4\}$. It follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies that




Figure 5.2: curvature distribution for Case (B1)

$c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). Checking (the LIST of Section 3) shows that if $d(\hat{\Delta}) = 5$ then $\hat{\Delta}$ receives positive curvature across at most one edge and so $c^*(\hat{\Delta}) \leq 0$. Also if $d(\hat{\Delta}) = 6$ then checking shows $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$.

- (ii) Suppose that $l(\Delta) = bd^2a^{-1} = 1$, $bda^{-1}c^{\pm 1} \neq 1$. If $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$ as in Figure 5.1(ii). If say $d(\Delta_2) = 4$ as in Figure 5.1(iii) then $l(\Delta_2) = bdda^{-1}$ which forces $l(\Delta_3) = ca\omega$. First assume that $cadb^{-1} \neq 1$. Then $d(\Delta_3) > 3$ and so add $\frac{\pi}{2}$ to $c(\Delta_3)$ via Δ_2 . If $d(\Delta_1) = 4$ then add $\frac{\pi}{2}$ to $c(\Delta_4)$ via Δ_1 in a similar way. Observe that Δ_1 does not receive positive curvature from Δ_3 or Δ_4 in Figure 5.1(ii); and Δ_3 does not receive positive curvature from Δ_1 or Δ_5 in Figure 5.1(iii). It follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). It remains to study $5 \leq d(\hat{\Delta}) \leq 6$. Checking shows that if $d(\hat{\Delta}) = 5$ then either the label contradicts $|c| \neq 1$ or $\hat{\Delta}$ receives positive curvature across at most one edge and so $c^*(\hat{\Delta}) \leq 0$. Also if $d(\hat{\Delta}) = 6$ then checking shows that $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$. Now assume that $cadb^{-1} = 1$, in which case $c = d^3$, $b = d^{-2}$ and $|d| = 6$. The distribution of curvature is exactly the same except when $d(\Delta_3) = 4$ in Figure 5.1(iii). In this case add $\frac{2}{3}c(\Delta) = \frac{2\pi}{3}$ to $c(\Delta_1)$ and $\frac{1}{3}c(\Delta) = \frac{\pi}{3}$ to $c(\Delta_5)$ via Δ_3 as shown in Figure 5.1(iv). Together with the observations above (which still hold) we also have that Δ_1 does not receive positive curvature from Δ_3 , Δ_4 or Δ_7 and that Δ_5 does not receive positive curvature from Δ_9 or Δ_{10} in Figure 5.1(iv). An argument similar to those given in the proof of Lemma 3.5 now shows that if $d(\hat{\Delta}) \geq 8$ then $c^*(\hat{\Delta}) \leq 0$; and that if $l(\hat{\Delta})$ does not involve $(cbd)^{\pm 1}$ then $c^*(\hat{\Delta}) \leq 0$ for $d(\hat{\Delta}) \geq 7$ by Lemma 3.5(vi). The conditions on b , c and d imply that if $2 < d(\hat{\Delta}) < 6$ then $l(\hat{\Delta}) \in \{d^2a^{-1}b, db^{-1}c^{\pm 1}a\}$ so if $l(\hat{\Delta})$ does not involve $(cbd)^{\pm 1}$ it remains to consider $d(\hat{\Delta}) = 6$. But checking shows that $l(\hat{\Delta})$ will then either involve at most two non-adjacent occurrences of $(bd)^{\pm 1}$, $(ca)^{\pm 1}$ or




Figure 5.3: curvature distribution for Case (B1)

$(ba^{-1})^{\pm 1}$ and so $c^*(\hat{\Delta}) \leq 0$ or $l(\hat{\Delta}) = (ab^{-1})^3$ in which case $c^*(\hat{\Delta}) \leq c(\hat{\Delta}) + 3(\frac{\pi}{3}) = 0$. Finally if $l(\hat{\Delta}) = cbd\omega$ and $d(\hat{\Delta}) \leq 7$ then $l(\hat{\Delta}) \in \{cbda^{-1}ba^{-1}, cbdb^{-1}ab^{-1}, cbd^3b^{-1}\}$ and $c^*(\hat{\Delta}) \leq c(\hat{\Delta}) + \frac{2\pi}{3} + \frac{\pi}{3} = 0$.

- (iii) Now suppose that $l(\Delta) = bda^{-1}c^{\pm 1} = 1$, $bd^2a^{-1} \neq 1$. First assume that $|b| \geq 3$. Add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$ as in Figure 5.1(ii). If say $d(\Delta_2) = 4$ then $l(\Delta_2) = bda^{-1}c^{\pm 1}$. First let $l(\Delta_2) = bda^{-1}c$ as in Figure 5.2(i). This forces $l(\Delta_4) = ad\omega$ and so $d(\Delta_4) = 4$ forces $l(\Delta_4) = addb^{-1}$. But if $b = d^2$ then $c = d^3$ and there is a sphere by Lemma 3.2(iv), so it can be assumed that $d(\Delta_4) > 4$. So add $\frac{\pi}{2}$ to $c(\Delta_4)$ via Δ_2 as shown. Suppose now that $l(\Delta_2) = bda^{-1}c^{-1}$ as in Figure 5.2(ii). This forces $l(\Delta_4) = ab^{-1}\omega$ and so $d(\Delta_4) > 4$, otherwise there is a contradiction to $|b| \geq 3$. So add $\frac{\pi}{2}$ to $c(\Delta_4)$ via Δ_2 as shown. Similarly add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_3)$ if $d(\Delta_1) = 4$. Observe that Δ_1 does not receive positive curvature from Δ_3 or Δ_4 in Figure 5.1(ii); Δ_2 does not receive positive curvature from Δ_3 or Δ_4 in Figure 5.1(ii); and Δ_4 does not receive positive curvature from Δ_1 or Δ_8 in Figures 5.2(i), (ii). It follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). It remains to study $5 \leq d(\hat{\Delta}) \leq 6$.

If $|b| > 3$ then checking shows that if $d(\hat{\Delta}) = 5$ then either the label contradicts one of the **(B1)** assumptions or $\hat{\Delta}$ receives positive curvature across at most one edge and so $c^*(\hat{\Delta}) \leq 0$. Checking shows that if $d(\hat{\Delta}) = 6$ then $\hat{\Delta}$ receives positive curvature across at most two edges or $l(\hat{\Delta}) = (ab^{-1})^3$ contradicting $|b| > 3$, and so $c^*(\hat{\Delta}) \leq 0$.

Let $|b| = 3$. If H is cyclic then $bdc = 1$ implies $b = d^2$ and we obtain a sphere as before, so assume otherwise. If $|b| = 3$ and $|d| \in \{4, 5\}$ then we obtain a sphere by Lemma 3.1(b)(iv). So let $|b| = 3$, $|d| \geq 6$. Distribute curvature from Δ as above and as shown in Figure 5.2. Checking shows that if $d(\hat{\Delta}) = 5$ then either the label contradicts $|b| = 3$ or $|d| \geq 6$ or $\hat{\Delta}$ receives positive curvature across at most one edge and so $c^*(\hat{\Delta}) \leq 0$. Checking shows that if $d(\hat{\Delta}) = 6$ then $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$ except when $l(\hat{\Delta}) = ba^{-1}ba^{-1}ba^{-1}$. This case is shown in Figure 5.3(i) where $c^*(\hat{\Delta}) = \frac{\pi}{2}$ and so add $\frac{1}{3}c^*(\hat{\Delta}) = \frac{\pi}{6}$ to $c(\hat{\Delta}_i)$ for $i \in \{1, 2, 3\}$ across the edge ad^{-1} . If $d(\hat{\Delta}_i) = 4$ then $l(\hat{\Delta}_i) \in \{ad^{-1}b^{-1}c^{-1}, ad^{-1}b^{-1}c\}$. Suppose that $l(\hat{\Delta}_1) = ad^{-1}b^{-1}c^{-1}$ as in Figure 5.3(ii). Then $l(\hat{\Delta}_4) = d^2b^{-1}\omega$ and $d(\hat{\Delta}_4) > 4$ otherwise there is a contradiction to H non-cyclic. So add $\frac{\pi}{6}$ to $c(\hat{\Delta}_4)$ across the edge db^{-1} . If $l(\hat{\Delta}_1) = ad^{-1}b^{-1}c$ as in Figure 5.3(iii) then $l(\hat{\Delta}_4) = d^3\omega$ and $d(\hat{\Delta}_4) > 4$ otherwise there is a contradiction to $|d| \geq 6$. So add $\frac{\pi}{6}$ to $c(\hat{\Delta}_4)$ across the edge d^2 . Observe that if $\hat{\Delta}$ receives positive curvature then it receives $\frac{\pi}{2}$ across the edges ab^{-1} , ad or bd ; and receives $\frac{\pi}{6}$ across the edges ad^{-1} , db^{-1} or dd . Thus there is a gap preceding $c^{\pm 1}$, b , a and a gap after $c^{\pm 1}$, b^{-1} , a^{-1} and there is a two-thirds gap across the edges ad^{-1} , db^{-1} and dd . Also there is always a gap between two d 's (other than when the subword is $d^{\pm 2}$). Now since c^2 cannot be a proper sublabel it follows that if there are at least two occurrences of $c^{\pm 1}$ then we obtain four gaps. Suppose now that there is at most one occurrence of $c^{\pm 1}$. If there is exactly one occurrence of c and either no occurrences of b or no occurrences of d then H is cyclic, a contradiction; and if there




Figure 5.4: curvature distribution for Case (B2)

are no occurrences of c and exactly one occurrence of d or of b then again H is cyclic, a contradiction. So assume otherwise. It follows that $l(\hat{\Delta})$ contains at least four gaps or $l(\hat{\Delta}) \in \{c^{\pm 1}ad^{\pm 1}a^{-1}ba^{-1}, c^{\pm 1}bd^{\pm 1}a^{-1}ba^{-1}, c^{\pm 1}ad^{\pm 1}b^{-1}ab^{-1}, c^{\pm 1}bd^{\pm 1}b^{-1}ab^{-1}, d(a^{-1}b)^{\pm 1}d^{\pm 1}(a^{-1}b)^{\pm 1}\}$. But since $c = bd$ each of these labels contradicts H non-cyclic, $|d| \geq 6$ or $|b| = 3$ except when $l(\hat{\Delta}) = da^{-1}bda^{-1}b$. In this case if $c^*(\hat{\Delta}) > 0$ then it can be assumed without loss of generality that $\hat{\Delta}$ is given by Figure 5.3(iv) and $\hat{\Delta}$ receives $\frac{1}{3}c^*(\hat{\Delta}_1) = \frac{\pi}{6}$ from $c(\hat{\Delta}_1)$. This implies that $l(\hat{\Delta}_3) = a^{-1}c^{-1}bd$ and $l(\hat{\Delta}_4) = ad^{-1}d^{-1}\omega$. So add $\frac{\pi}{6}$ from $c(\hat{\Delta})$ to $c(\hat{\Delta}_4)$. Since this $\frac{\pi}{6}$ is across an ad^{-1} edge and since $l(\hat{\Delta}_4) = ad^{-2}\omega$ it follows from the above that $c^*(\hat{\Delta}_4) \leq 0$. If $l(\hat{\Delta}_2) = b^{-1}ab^{-1}ab^{-1}$ in Figure 5.3(iv) then a similar argument applies to $c(\hat{\Delta}_5)$.

Finally let $|b| = 2$. In particular, H is non-cyclic for otherwise $d^2 = 1$, a contradiction. If $|d| < \infty$ then we obtain a sphere by Lemma 3.1(b)(i) and if $|d| = \infty$ then \mathcal{P} is aspherical by Lemma 3.4(ii).

In conclusion \mathcal{P} is aspherical in this case except when H is non-cyclic, $bda^{-1}c^{\pm 1} = 1$ and either $|b| = 3$, $|d| \in \{4, 5\}$ or $|b| = 2$, $|d| < \infty$; or when H is cyclic, $b = d^2$, $c = d^3$ and $|d| = 6$.

(B2) $|c| = 2$, $|d| = 2$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If H is cyclic then $c = d$. Therefore $bd^{-1}b^{-1}c = cada^{-1} = 1$ and there is a sphere by Lemma 3.2(i). So suppose from now on that H is non-cyclic. If $d(\Delta) = 2$ then Δ is given by Figures 5.1(ii) and 5.4(i). Moreover, if $d(\Delta_i) = 4$ for $i \in \{1, 2\}$ then $l(\Delta_i) \in \{bd^2a^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}, cad^{\pm 1}a^{-1}, cad^{\pm 1}b^{-1}\}$. But $bdda^{-1} = 1$ implies that $b = 1$ which is a contradiction. This leaves the following cases:

- (i) $bda^{-1}c^{\pm 1} \neq 1$, $bdb^{-1}c^{\pm 1} \neq 1$, $cad^{\pm 1}a^{-1} \neq 1$;
- (ii) $bda^{-1}c^{\pm 1} = 1$, $bdb^{-1}c^{\pm 1} \neq 1$, $cad^{\pm 1}a^{-1} \neq 1$;
- (iii) $bdb^{-1}c^{\pm 1} = 1$, $bda^{-1}c^{\pm 1} \neq 1$, $cad^{\pm 1}a^{-1} \neq 1$;
- (iv) $cad^{\pm 1}a^{-1} = 1$, $bda^{-1}c^{\pm 1} \neq 1$, $bdb^{-1}c^{\pm 1} \neq 1$;

(v) $bdb^{-1}c^{\pm 1} = 1$, $cad^{\pm 1}a^{-1} = 1$, $bda^{-1}c^{\pm 1} \neq 1$.

(Note that any other combination implies $b = 1$.)

- (i) In this case $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ in Figures 5.1(ii) and 5.4(i) so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to each of $c(\Delta_1)$ and $c(\Delta_2)$. Observe that Δ_1 and Δ_2 do not receive positive curvature from Δ_3 or Δ_4 in Figures 5.1(ii) and 5.4(i). It follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). It remains to study $5 \leq d(\hat{\Delta}) \leq 6$. Checking shows that if $d(\hat{\Delta}) = 5$ then either the label contradicts $c^{\pm 1}ab^{-1} \neq 1$ or $\hat{\Delta}$ receives positive curvature across at most one edge and so $c^*(\hat{\Delta}) \leq 0$. Also if $d(\hat{\Delta}) = 6$ then $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$.
- (ii) In this case the labels $bda^{-1}c^{\pm 1}$ can occur. If $|b| < \infty$ then we obtain spheres by Lemma 3.1(b)(ii) and if $|b| = \infty$ then \mathcal{P} is aspherical by Lemma 3.4(iii).
- (iii) In this case the labels $bdb^{-1}c^{\pm 1}$ can occur and H is non-Abelian. If $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ as in Figures 5.1(ii) and 5.4(i) then add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. If say $d(\Delta_2) = 4$ as in Figure 5.4(ii) then $l(\Delta_3) = c^{-1}a\omega$ and so $d(\Delta_3) > 4$, otherwise there is a contradiction to $c \neq d$ or $b \neq 1$. So add $\frac{\pi}{2}$ to $c(\Delta_3)$ via Δ_2 as shown. If $d(\Delta_1) = 4$ then similarly add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_4)$ via Δ_1 . We see from Figures 5.1(ii) and 5.4(i)–(ii) that if $\hat{\Delta}$ receives positive curvature then it does so across the edges bd, ca or $c^{-1}a$. It follows that the only word of length 3 that contains no gaps is $a^{-1}c^{\pm 1}a$. Suppose that $l(\hat{\Delta}) = \omega_1 a^{-1} c^{\pm 1} a \omega_2 \omega$ where ω_1 and ω_2 have length 2 and ω has length at least 0. Then there is a gap preceding a^{-1} in $\omega_1 a^{-1}$ and after a in $a \omega_2$. Moreover, if ω_1 does not contain a gap then $\omega_1 = bd$ and if ω_2 does not contain a gap then $\omega_2 = d^{-1}b^{-1}$. It follows that if $l(\omega) > 0$ then $l(\hat{\Delta})$ contain at least 4 gaps and $c^*(\hat{\Delta}) \leq 0$. If $l(\omega) = 0$ and $l(\hat{\Delta}) = \omega_1 a^{-1} c^{\pm 1} a d^{-1} b^{-1}$ then $l(\hat{\Delta}) \in \{c^{\pm 1} b a^{-1} c^{\pm 1} a d^{-1} b^{-1}, a d^{\pm 1} a^{-1} c^{\pm 1} a d^{-1} b^{-1}\}$ and there are 4 gaps; if $l(\omega) = 0$ and $l(\hat{\Delta}) = b d a^{-1} c^{\pm 1} a \omega_2$ then $l(\hat{\Delta}) \in \{b d a^{-1} c^{\pm 1} a d^{\pm 1} a^{-1}, b d a^{-1} c^{\pm 1} a b^{-1} c^{\pm 1}\}$ and again there are 4 gaps. If now $l(\hat{\Delta}) = a^{-1} c^{\pm 1} a \omega$ and $d(\hat{\Delta}) \leq 6$ then either $l(\hat{\Delta}) = a^{-1} c^{\pm 1} a d^{\pm 2}$, a contradiction, or there are 4 gaps. Now suppose that $l(\hat{\Delta})$ does not contain the subword $a^{-1} c^{\pm 1} a$. Then $\hat{\Delta}$ receives positive curvature across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). So let $d(\hat{\Delta}) \leq 6$ and $l(\hat{\Delta}) \in \{c a \omega, b d \omega, c^{-1} a \omega\}$. If $d(\hat{\Delta}) < 6$ then checking shows that there is a contradiction to H non-Abelian; and if $d(\hat{\Delta}) = 6$ then checking shows that $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$.
- (iv) In this case the labels $cad^{\pm 1}a^{-1}$ can occur and H is non-Abelian. If $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ as in Figures 5.1(ii) and 5.4(i) then add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. If say $d(\Delta_2) = 4$ then $l(\Delta_2) = cad^{\pm 1}a^{-1}$ which forces $l(\Delta_4) = bd^{-1}\omega$ and so $d(\Delta_3) > 4$, otherwise there is a contradiction to $b \neq 1$ or $bd^{-1}b^{-1}c^{\pm 1} \neq 1$. So add $\frac{\pi}{2}$ to $c(\Delta_4)$ via Δ_2 as shown in Figure 5.4(iii). If $d(\Delta_1) = 4$ then similarly add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_3)$ via

Δ_1 . We see from Figures 5.1(ii), 5.4(i) and 5.4(iii) that if $\hat{\Delta}$ receives positive curvature then it does so across the edges bd , bd^{-1} or ca . It follows that the only word of length 3 that contains no gaps is $bd^{\pm 1}b^{-1}$. Suppose that $l(\hat{\Delta}) = \omega_1 bd^{\pm 1}b^{-1}\omega_2\omega$ where ω_1 and ω_2 have length 2 and ω has length at least 0. Then there is a gap preceding b in ω_1b and after b^{-1} in $b^{-1}\omega_2$. Moreover, if ω_1 does not contain a gap then $\omega_1 = a^{-1}c^{-1}$ and if ω_2 does not contain a gap then $\omega_2 = ca$. It follows that if $l(\omega) > 0$ then $l(\hat{\Delta})$ contain at least 4 gaps and $c^*(\hat{\Delta}) \leq 0$. If $l(\omega) = 0$ and $l(\hat{\Delta}) = \omega_1 bd^{\pm 1}b^{-1}ca$ then $l(\hat{\Delta}) \in \{d^{\pm 1}a^{-1}bdb^{-1}ca, bc^{\pm 1}bdb^{-1}ca\}$ and there are 4 gaps; if $l(\omega) = 0$ and $l(\hat{\Delta}) = a^{-1}c^{-1}bd^{\pm 1}b^{-1}\omega_2$ then $l(\hat{\Delta}) \in \{a^{-1}c^{-1}bdb^{-1}ad^{\pm 1}, a^{-1}c^{-1}bdb^{-1}c^{\pm 1}b\}$ and again there are 4 gaps. If $l(\hat{\Delta}) = bd^{\pm 1}b^{-1}\omega$ and $d(\hat{\Delta}) \leq 6$ then there are 4 gaps. Now suppose that $l(\hat{\Delta})$ does not contain the subword $bd^{\pm 1}b^{-1}$. Then $\hat{\Delta}$ receives positive curvature across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). So let $d(\hat{\Delta}) \leq 6$ and $l(\hat{\Delta}) \in \{bd\omega, bd^{-1}\omega, ca\omega\}$. If $d(\hat{\Delta}) < 6$ then checking shows that there is a contradiction to H non-Abelian; and if $d(\hat{\Delta}) = 6$ then checking shows that $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$.

- (v) In this case the labels $bdb^{-1}c^{\pm 1}$ and $cad^{\pm 1}a^{-1}$ can occur. If $|b| = \infty$ then \mathcal{P} is aspherical by Lemma 3.4(iii); and if $|b| < \infty$ then there is a sphere by Lemma 3.2(i).

In conclusion \mathcal{P} fails to be aspherical in this case either when H is cyclic or when H is non-cyclic and when $bda^{-1}c^{-1} = 1$, $|b| < \infty$ or when $bdb^{-1}c^{-1} = a^{-1}cad = 1$, $|b| < \infty$.

(B3) $|c| = 2$, $|d| = 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 2$ then Δ is given by Figure 5.1(ii). If $d(\Delta) = 3$ then Δ is given by Figure 4.3(i).

Moreover, if $d(\Delta_i) = 4$ and $l(\Delta_i) = bdw$ or caw then

$l(\Delta_i) \in \{bd^2a^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}, cad^{\pm 1}a^{-1}, cad^{\pm 1}b^{-1}\}$. But each of $bd^2a^{-1} = 1$,

$bdb^{-1}c^{\pm 1} = 1$ and $cad^{\pm 1}a^{-1} = 1$ implies a contradiction to one of the **(B3)** assumptions.

Thus we have the following cases:

- (i) $bda^{-1}c^{\pm 1} \neq 1$, $cadb^{-1} \neq 1$;
- (ii) $bda^{-1}c^{\pm 1} = 1$, $cadb^{-1} \neq 1$;
- (iii) $cadb^{-1} = 1$, $bda^{-1}c^{\pm 1} \neq 1$.

- (i) In this case $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ in Figure 5.1(ii), so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to each of $c(\Delta_1)$ and $c(\Delta_2)$. In Figure 4.3(i) $d(\Delta_1) > 4$, $d(\Delta_3) > 4$ and $d(\Delta_5) > 4$ so add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each of $c(\Delta_1)$, $c(\Delta_3)$ and $c(\Delta_5)$. Observe that Δ_1 and Δ_2 do not receive positive curvature from Δ_3 or Δ_4 in Figure 5.1(ii). Also Δ_1 , Δ_3 and Δ_5 do not receive positive curvature from Δ_m for $m \in \{2, 4, 6\}$ in Figure 4.3(i). It follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). It remains to study $5 \leq d(\hat{\Delta}) \leq 6$. Checking shows that if $d(\hat{\Delta}) = 5$ then either the label contradicts




Figure 5.5: curvature distribution for Case (B3)

$cab^{-1} \neq 1$ or $\hat{\Delta}$ receives positive curvature across at most one edge and so $c^*(\hat{\Delta}) \leq 0$. Also if $d(\hat{\Delta}) = 6$ then $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$.

- (ii) In this case the labels $bda^{-1}c^{\pm 1}$ can occur. If H is cyclic then $d = b^2$, $c = b^3$ and there is a sphere by Lemma 3.2(v), so assume that H is non-cyclic. If $|b| \in \{2, 3, 4, 5\}$ then we obtain spheres by Lemma 3.1(b)(i), (v), so assume that $|b| \geq 6$. In Figure 5.1(ii) if $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$ as shown. If say $d(\Delta_1) > 4$ and $d(\Delta_2) = 4$ as in Figure 5.5(i) then add $\frac{\pi}{2}$ to $c(\Delta_1)$. This implies that $l(\Delta_2) = bda^{-1}c^{\pm 1}$ as shown. This forces $l(\Delta_3) = b^{-1}a\omega$ and so $d(\Delta_3) > 4$, otherwise there is a contradiction to $|b| \geq 6$ so add $\frac{\pi}{2}$ to $c(\Delta_3)$ via Δ_2 as shown. If $d(\Delta_1) = d(\Delta_2) = 4$ then add $\frac{\pi}{2}$ to $c(\Delta_j)$ for $j \in \{3, 4\}$ as shown in Figure 5.5(ii). The one exception to the above is when $l(\Delta_1) = bda^{-1}\omega$ and $d(\Delta_1) > 4$. Then $d(\Delta_4) > 4$ and in this situation add the $\frac{\pi}{2}$ from $c(\Delta)$ to $c(\Delta_4)$ via Δ_1 as shown in Figure 5.5(iii). The same applies to Δ_2 . In Figure 4.3(i) if $d(\Delta_1) > 4$, $d(\Delta_3) > 4$ and $d(\Delta_5) > 4$ then add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each of $c(\Delta_1)$, $c(\Delta_2)$ and $c(\Delta_5)$. If say $d(\Delta_1) = 4$, $d(\Delta_3) > 4$ and $d(\Delta_5) > 4$ then $l(\Delta_2) = ba^{-1}\omega$ and $d(\Delta_2) > 4$ otherwise there is a contradiction to $|b| \geq 6$ so add $\frac{\pi}{6}$ to $c(\Delta_2)$, $c(\Delta_3)$ and $c(\Delta_5)$ as shown in Figure 5.6(i). Now suppose that $d(\Delta_1) = 4$ and $d(\Delta_3) = 4$. This implies that $l(\Delta_2) = l(\Delta_4) = ba^{-1}\omega$ as shown in Figure 5.6(ii). So add $\frac{\pi}{6}$ to $c(\Delta_2)$, $c(\Delta_4)$ and $c(\Delta_5)$. If $d(\Delta_1) = d(\Delta_3) = d(\Delta_5) = 4$ then similarly add $\frac{\pi}{6}$ to $c(\Delta_m)$ for $m \in \{2, 4, 6\}$.

We now see that if $\hat{\Delta}$ receives positive curvature then it receives at most $\frac{\pi}{2}$ across $(bd)^{\pm 1}$ and $(b^{-1}a)^{\pm 1}$; and it receives at most $\frac{\pi}{6}$ across $(ca)^{\pm 1}$ and $(ab^{-1})^{\pm 1}$. Thus there is always a gap immediately preceding c and d^{-1} ; and there is a gap immediately after c^{-1} and d . This implies that if there are at least four occurrences of $c^{\pm 1}$ or $d^{\pm 1}$ then $l(\hat{\Delta})$ contains at least four gaps and so $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.6. Suppose that there are at most three occurrences of $c^{\pm 1}$ or $d^{\pm 1}$ in $l(\hat{\Delta})$. Observe that in addition to the four gaps mentioned above the following sublabeled yield gaps: $(cb)^{\pm 1}$ and $(ad)^{\pm 1}$ each




Figure 5.6: curvature distribution for Case (B3)

yields a gap; $(bda^{-1})^{\pm 1}$ yields two gaps (see Figure 5.5(iii)); and $(ca)^{\pm 1}$ and $(ab^{-1})^{\pm 1}$ each yields the equivalent of a two-thirds gap. If $l(\hat{\Delta}) = (b^{-1}a)^{\pm n}$ where $n \geq 1$ then $l(\hat{\Delta})$ obtains at least four gaps since $|b| \geq 6$. If $l(\hat{\Delta}) \in \{d^{\pm 1}(b^{-1}a)^{\pm n}, (ab^{-1})^{\pm n}c^{\pm 1}\}$ then H is cyclic so it can be assumed that $l(\hat{\Delta})$ involves either two or three occurrences of $c^{\pm 1}$ or $d^{\pm 1}$. It follows that if there are three occurrences then $c^*(\hat{\Delta}) \leq 0$; or if exactly two occurrences then either $c^*(\hat{\Delta}) \leq 0$ or $l(\hat{\Delta}) \in \{d^{\pm 1}b^{-1}ada^{-1}b, d^{\pm 1}b^{-1}ada^{-1}ba^{-1}b, d^{\pm 1}b^{-1}ab^{-1}ada^{-1}b, cba^{-1}bd^{\pm 1}b^{-1}, cbd^{\pm 1}b^{-1}ab^{-1}\}$. But each of these labels forces H cyclic or $|b| < 6$ or a **(B3)** contradiction, therefore $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.6.

- (iii) In this case the label $cadb^{-1}$ can occur. First assume that H is non-cyclic. In Figure 5.1(ii) $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ otherwise there is a contradiction to $|c| = 2$ or $|d| = 3$, so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. In Figure 4.3(i) if $d(\Delta_1) > 4$, $d(\Delta_3) > 4$ and $d(\Delta_5) > 4$ add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each of $c(\Delta_1)$, $c(\Delta_2)$ and $c(\Delta_5)$. If say $d(\Delta_1) = 4$ only then add $\frac{\pi}{4}$ to $c(\Delta_3)$ and $c(\Delta_5)$. Now suppose that $d(\Delta_1) = d(\Delta_3) = 4$. This implies that their label is $cadb^{-1}$ which forces $l(\Delta_2) = cba^{-1}\omega$ as shown in Figure 5.6(iii). So add $\frac{\pi}{4}$ to $c(\Delta_2)$ via Δ_1 and to $c(\Delta_5)$. Finally if $d(\Delta_1) = d(\Delta_3) = d(\Delta_5) = 4$ then in a similar way add $\frac{\pi}{6}$ to $c(\Delta_m)$ for $m \in \{2, 4, 6\}$. Observe that Δ_1 does not receive positive curvature from Δ_3 and Δ_2 does not receive positive curvature from Δ_4 in Figure 5.1(ii). In Figure 4.3(i) Δ_1 does not receive positive curvature from Δ_2 . In Figure 5.6(iii) Δ_2 does not receive positive curvature from Δ_3 . Since $\hat{\Delta}$ receives $\frac{\pi}{2}$ across the bd edge and $\frac{\pi}{4}$ across the ca , ba^{-1} edges it follows that $\hat{\Delta}$ receives an average of $\frac{\pi}{4}$ across each of its edges, so $d(\hat{\Delta}) \geq 8$ implies that $c^*(\hat{\Delta}) \leq 0$. It remains to study $5 \leq d(\hat{\Delta}) \leq 7$. Checking shows that if $d(\hat{\Delta}) = 5$ then either the label contradicts $|d| = 3$ or H non-cyclic or $\hat{\Delta}$ receives positive curvature across at most one edge and so $c^*(\hat{\Delta}) \leq 0$, except when $l(\hat{\Delta}) = bdda^{-1}c^{-1}$ as in Figure 5.7(i). In this case $\hat{\Delta}$ receives $\frac{\pi}{2}$ from $c(\hat{\Delta}_1)$ and $\frac{\pi}{4}$ from $c(\hat{\Delta}_2)$. If $|b| > 2$ then this implies that $d(\hat{\Delta}_3) > 4$ and so add $\frac{\pi}{4}$ to $c(\hat{\Delta}_3)$ noting that this is a similar edge to the one crossed in Figure 5.6(iii) so there is no change to the above argument and $c^*(\hat{\Delta}) \leq 0$ in this




Figure 5.7: curvature distribution for Case (B3)

case. Suppose now that $|b| = 2$ and $l(\hat{\Delta}_3) = ab^{-1}ab^{-1}$ as in Figure 5.7(ii). If $d(\hat{\Delta}_4) > 5$ then add $\frac{\pi}{4}$ to $c(\hat{\Delta}_4)$ across the da^{-1} edge. If $d(\hat{\Delta}_4) = 5$ then $l(\hat{\Delta}_4) = da^{-1}c^{-1}bd$ which implies that $l(\hat{\Delta}_5) = caw$ and so if $d(\hat{\Delta}_5) > 5$ then add $\frac{\pi}{4}$ to $c(\hat{\Delta}_5)$ as in Figure 5.7(iii). If $d(\hat{\Delta}_5) \in \{4, 5\}$ then $l(\hat{\Delta}_5) \in \{cadb^{-1}, cad^{-2}b^{-1}\}$ and this forces $l(\hat{\Delta}_6) = c^{-1}ba^{-1}\omega$ and so $d(\hat{\Delta}_6) > 5$ otherwise there is a contradiction to $|c| \neq 1$. So add $\frac{\pi}{4}$ to $c(\hat{\Delta}_6)$ again as shown in Figure 5.7(iii).

Observe that $\hat{\Delta}_4$ in Figure 5.7(ii) can now receive $\frac{\pi}{4}$ from $c(\hat{\Delta})$, however it receives no positive curvature from $\hat{\Delta}_3$ or any other region across the da^{-1} edge. Moreover, it is clear from Figure 5.7(iii) that $\hat{\Delta}_5$ receives only the $\frac{\pi}{4}$ from $\hat{\Delta}_4$ across its ca edge; and $\hat{\Delta}_6$ receives only the $\frac{\pi}{4}$ from $\hat{\Delta}_5$ across its ba^{-1} edge. Finally observe that Figures 5.7(ii)–(iii) do not alter the fact that Δ_1 does not receive positive curvature from Δ_3 and Δ_2 does not receive positive curvature from Δ_4 in Figure 5.1(ii). Therefore the average positive curvature that $\hat{\Delta}$ receives across each edge is still $\frac{\pi}{4}$ and so if $d(\hat{\Delta}) \geq 8$ then $c^*(\hat{\Delta}) \leq 0$. It remains to check $6 \leq d(\hat{\Delta}) \leq 7$ for the sublabeled $(bd)^{\pm 1}(\pi/2)$ and $(ca)^{\pm 1}, (ab^{-1})^{\pm 1}, (da^{-1}c^{-1})^{\pm 1}(\pi/4)$. Checking shows that if $d(\hat{\Delta}) = 6$ then the most curvature that $\hat{\Delta}$ can receive is either $2(\frac{\pi}{2})$ or $\frac{\pi}{2} + 2(\frac{\pi}{4})$ or $4(\frac{\pi}{4})$ and so $c^*(\hat{\Delta}) \leq 0$. If $d(\hat{\Delta}) = 7$ then the most curvature received is $3(\frac{\pi}{2})$ or $2(\frac{\pi}{2}) + 2(\frac{\pi}{4})$ or $\frac{\pi}{2} + 4(\frac{\pi}{4})$ or $6(\frac{\pi}{4})$ and $c^*(\hat{\Delta}) \leq 0$ except for $l(\hat{\Delta}) = da^{-1}c^{-1}bda^{-1}b$; but this implies $cd = 1$, a contradiction.

Now let H be cyclic. Then $d = b^4$ and $c = b^3$. Again add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to each of $c(\Delta_1), c(\Delta_2)$ as in Figure 5.1(ii). In Figure 4.3(i) if say $d(\Delta_1) > 5$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and so it can be assumed that $d(\Delta_i) \leq 5$ for $i \in \{1, 3, 5\}$ in which case $l(\Delta_i) \in \{cadb^{-1}, cad^{-2}b^{-1}\}$. If say $d(\Delta_1) = 4$ then add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_6)$ via Δ_1 as shown in Figure 5.7(iv). It can be assumed then that $d(\Delta_i) = 5$ for $i \in \{1, 3, 5\}$ in which case add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to each $c(\hat{\Delta})$ via Δ_i where $i \in \{1, 3, 5\}$ as shown in Figure 5.8(i). If say $\hat{\Delta} = \hat{\Delta}_1$ and $d(\hat{\Delta}_1) = 5$ then




Figure 5.8: curvature distribution for Cases (B3) and (B5)

repeat the above, that is, add the $\frac{\pi}{6}$ from $c(\Delta)$ across another ca edge and continue in this way until $\frac{\pi}{6}$ is eventually added to a region $\hat{\Delta}_k$ where either $d(\hat{\Delta}_k) > 5$ (and so the

process terminates) or $d(\hat{\Delta}_k) = 4$ in which case the $\frac{\pi}{6}$ from $c(\Delta)$ is added to $c(\hat{\Delta}_{k+1})$ as shown in Figure 5.8(ii), where $k = 3$. If $d(\hat{\Delta}_{k+1}) > 5$ then the process terminates (and note that $l(\hat{\Delta}_{k+1}) = d^{-1}b^{-1}c^{-1}w$); otherwise $l(\hat{\Delta}_{k+1}) = d^{-1}b^{-1}c^{-1}ad^{-1}$ and the $\frac{\pi}{6}$ from $c(\Delta)$ is added to $c(\hat{\Delta}_{k+2})$ where $\hat{\Delta}_{k+2}$ is the region shown in Figure 5.8(ii) with $k = 3$. Observe that $l(\hat{\Delta}_{k+2}) = ba^{-1}c^{-1}w$ so $d(\hat{\Delta}_{k+2}) > 5$ and the process terminates. This completes the distribution of curvature that occurs. It follows that if $\hat{\Delta}$ receives positive curvature across an edge e_i say then $\hat{\Delta}$ does not receive any curvature across the adjacent edges e_{i-1} , e_{i+1} except when $\hat{\Delta}$ is given by $\hat{\Delta}_{k+1} = \hat{\Delta}_4$ in Figure 5.8(ii). Therefore if $l(\hat{\Delta})$ does not involve $(cbd)^{\pm 1}$ then Lemma 3.5(vi) applies and $c^*(\hat{\Delta}) \leq 0$ for $d(\hat{\Delta}) \geq 7$; and if $d(\hat{\Delta}) = 6$ then checking for $(bd)^{\pm 1}$, $(ca)^{\pm 1}$ and $(cba^{-1})^{\pm 1}$ shows that $\hat{\Delta}$ receives positive curvature across at most two edges and $c^*(\hat{\Delta}) \leq 0$. Finally if $l(\hat{\Delta}) = cbdw$ then we see from Figure 5.8(ii) that the maximum amount $\hat{\Delta}$ receives is on average $\frac{\pi}{3}$ across $\frac{2}{3}$ of its edges and so if $d(\hat{\Delta}) \geq 8$ then $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vii). Checking shows that if $6 \leq d(\hat{\Delta}) \leq 7$ then $l(\hat{\Delta}) \in \{cbdb^{-1}ab^{-1}, cbda^{-1}bdb^{-1}, cbda^{-1}c^{-1}ba^{-1}, cbda^{-1}cba^{-1}\}$ and so if $d(\hat{\Delta}) = 6, 7$ then $\hat{\Delta}$ receives curvature across at most 2, 3 edges (respectively) and $c^*(\hat{\Delta}) \leq 0$.

In conclusion \mathcal{P} fails to be aspherical in this case when H is non-cyclic, $bda^{-1}c^{-1} = 1$ and $|b| \in \{2, 3, 4, 5\}$; or when H is cyclic and $bda^{-1}c^{\pm 1} = 1$.

(B4) $|c| = 2$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} = 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $|d| < \infty$ then there is a sphere by Lemma 3.1(a)(i) and if $|d| = \infty$ then \mathcal{P} is aspherical by Lemma 3.4(ii).

(B5) $|c| = 2$, $|d| > 3$, $a^{-1}b \neq 1$, $c^{\pm 1}ab^{-1} \neq 1$, $db^{-1}a = 1$, $d^{-1}b^{-1}a \neq 1$.

If $d(\Delta) = 2$ then Δ is given by Figure 5.1(ii) in which case add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to each of $c(\Delta_1)$ and $c(\Delta_2)$ as shown; and if $d(\Delta) = 3$ then Δ is given by Figure 4.10(iii) in which case add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_5)$ as shown. Assume that H is non-cyclic. If say $d(\Delta_1) = 4$ in Figure 5.1(ii) then $l(\Delta_1) \in \{bd^2a^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}\}$ which contradicts $|d| > 3$ or H non-cyclic; and if $d(\Delta_5) = 4$ in Figure 4.10(iii) then $l(\Delta_5) \in \{c^{-1}ad^{\pm 1}a^{-1}, c^{-1}ad^{\pm 1}b^{-1}\}$ which contradicts $|d| > 3$ or H non-cyclic. Observe that in Figure 5.1(ii) Δ_1 say does not receive positive curvature from Δ_3 or Δ_4 ; and in Figure 4.10(iii) Δ_5 does not receive any from Δ_4 or Δ_6 . Therefore $c^*(\hat{\Delta}) \leq 0$ for $d(\hat{\Delta}) \geq 7$ by Lemma 3.5(vi) and so it remains to consider $5 \leq d(\hat{\Delta}) \leq 6$. But checking for subwords $(bd)^{\pm 1}, (c^{-1}a)^{\pm 1}$ shows that if $d(\hat{\Delta}) = 5, 6$ then $\hat{\Delta}$ receives positive curvature across 1, 2 edges respectively except when $l(\hat{\Delta}) = d^2a^{-1}cb$. But H would then be cyclic, so $c^*(\hat{\Delta}) \leq 0$.

Now assume that H is cyclic. If $bda^{-1}c = 1$ then the conditions are T -equivalent to those of Lemma 3.1(a)(ii) and there is a sphere, so assume otherwise. Then $d(\Delta_1) > 4$ in Figure 5.1(ii) and $d(\Delta_5) > 4$ in Figure 4.10(iii). The argument above now applies except when $l(\Delta_1) = bd^2a^{-1}c$ in which case add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_3)$ via Δ_1 as shown in Figure 5.8(iii). We see from Figures 4.10(iii), 5.1(ii) and 5.8(iii) that if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and so $c^*(\hat{\Delta}) \leq 0$ for $d(\hat{\Delta}) \geq 7$ by Lemma 3.5(vi), so let $4 \leq d(\hat{\Delta}) \leq 6$. Then either $l(\hat{\Delta}) = bd^2a^{-1}c^{-1}$ and $c^*(\hat{\Delta}) \leq c(\Delta) + \frac{\pi}{2} = 0$; or $d(\hat{\Delta}) = 6$, $\hat{\Delta}$ receives across at most two edges (checking for $(bd)^{\pm 1}, (c^{-1}a)^{\pm 1}, (ad)^{\pm 1}$) and again $c^*(\hat{\Delta}) \leq 0$.

In conclusion \mathcal{P} is aspherical if and only if $bda^{-1}c \neq 1$.

(B6) $|c| = 2, |d| = 3, a^{-1}b \neq 1, c^{\pm 1}ab^{-1} \neq 1, db^{-1}a = 1, d^{-1}b^{-1}a \neq 1$.

If $d(\Delta) = 2$ then Δ is given by Figure 5.1(ii). If $d(\Delta) = 3$ then Δ is given by Figures 4.3(i) and 4.10(iii). In Figure 5.1(ii) if $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. If $d(\Delta_1) > 4$ and $d(\Delta_2) = 4$ then $l(\Delta_2) \in \{bd^2a^{-1}, bda^{-1}c^{\pm 1}, bdb^{-1}c^{\pm 1}\}$ and so $l(\Delta_2) = bd^2a^{-1}$ otherwise there is a contradiction to $|d| = 3$. Therefore Δ_2 is given by Figure 5.1(iii) forcing $l(\Delta_3) = ca\omega$ and so $d(\Delta_3) > 4$, otherwise there is a contradiction to $|c| = 2$, so add $\frac{\pi}{2}$ to $c(\Delta_1)$ and to $c(\Delta_3)$ via Δ_2 . If $d(\Delta_1) = 4$ then add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_4)$ via Δ_1 . Observe in Figure 4.10(iii) that if $d(\Delta_5) = 4$ then $l(\Delta_5) \in \{c^{-1}ad^{\pm 1}a^{-1}, c^{-1}ad^{\pm 1}b^{-1}\}$ which contradicts $|c| = 2$ so add $c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_5)$ as shown. In Figure 4.3(i) if $d(\Delta_i) = 4$ where $i \in \{1, 3, 5\}$ then $l(\Delta_i) \in \{cad^{\pm 1}a^{-1}, cad^{\pm 1}b^{-1}\}$ which contradicts $|c| = 2$ so add $c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$ as shown. We see from Figures 4.3(i), 4.10(iii) and 5.1(ii)–(iii) that if $\hat{\Delta}$ receives positive curvature then it does so across the edges bd, ca or $c^{-1}a$. It follows that the only word of length 3 that contains no gaps is $a^{-1}c^{\pm 1}a$. Suppose that $l(\hat{\Delta}) = \omega_1a^{-1}c^{\pm 1}a\omega_2$ where ω_1 and ω_2 have length 2 and ω has length at least 0. Then there is a gap preceding a^{-1} in ω_1a^{-1} and after a in $a\omega_2$. Moreover, if ω_1 does not contain a gap then $\omega_1 = bd$ and if ω_2 does not contain a gap then $\omega_2 = d^{-1}b^{-1}$. It follows that if $l(\omega) > 0$ then $l(\hat{\Delta})$ contain at least 4 gaps and $c^*(\hat{\Delta}) \leq 0$. If $l(\omega) = 0$ and $l(\hat{\Delta}) = \omega_1a^{-1}c^{\pm 1}ad^{-1}b^{-1}$ then $l(\hat{\Delta}) \in \{c^{\pm 1}ba^{-1}c^{\pm 1}ad^{-1}b^{-1}, ad^{\pm 1}a^{-1}c^{\pm 1}ad^{-1}b^{-1}\}$ and there are 4 gaps; if $l(\omega) = 0$ and $l(\hat{\Delta}) = bda^{-1}c^{\pm 1}a\omega_2$ then $l(\hat{\Delta}) \in \{bda^{-1}c^{\pm 1}ad^{\pm 1}a^{-1}, bda^{-1}c^{\pm 1}ab^{-1}c^{\pm 1}\}$ and again there are 4 gaps. If $l(\hat{\Delta}) = a^{-1}c^{\pm 1}a\omega$ and $d(\hat{\Delta}) \leq 6$ then either we obtain a contradiction to $|c| = 2$




Figure 5.9: curvature distribution for Cases (B7,8,9,11,12)

or $l(\hat{\Delta}) = a^{-1}c^{\pm 1}ab^{-1}c^{\pm 1}b$ and there are 4 gaps. Now suppose that $l(\hat{\Delta})$ does not contain the subword $a^{-1}c^{\pm 1}a$. Then $\hat{\Delta}$ receives positive curvature across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). So let $d(\hat{\Delta}) \leq 6$ and

$l(\hat{\Delta}) \in \{caw, bdw, c^{-1}aw\}$. If $d(\hat{\Delta}) < 6$ then checking shows that there is a contradiction to a **(B6)** assumption or $l(\hat{\Delta}) = bda^{-1}ba^{-1}$ and there are four gaps; and if $d(\hat{\Delta}) = 6$ then checking for $c^{-1}aw$ shows that either there is a contradiction to a **(B6)** assumption or $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$. Therefore \mathcal{P} is aspherical.

(B7) $|c| > 3$, $|d| > 3$, $a^{-1}b = 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 2$ then Δ is given by Figure 5.9(i). Here $l(\Delta_1) = b^{-1}c\omega$ and $l(\Delta_2) = ad^{-1}\omega$. First assume that H is non-cyclic. This implies that $d(\Delta_1) > 4$, $d(\Delta_2) > 4$, otherwise there is a contradiction to H non-cyclic, $|c| > 3$, or $|d| > 3$. So add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. Observe that each of Δ_1 and Δ_2 does not receive positive curvature from Δ_j for $j \in \{3, 4\}$. It follows that if $\hat{\Delta}$ receives positive curvature then it does so across at most half of its edges and so $d(\hat{\Delta}) \geq 7$ implies that $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.5(vi). Checking shows that if $d(\hat{\Delta}) = 5$ then $l(\hat{\Delta})$ contradicts $|c| > 3$, $|d| > 3$ or H non-cyclic. Also if $d(\hat{\Delta}) = 6$ then $\hat{\Delta}$ receives positive curvature across at most two edges and so $c^*(\hat{\Delta}) \leq 0$.

Now assume that H is cyclic. If $c = d^{\pm 1}$ there is a sphere by Lemma 3.1(a)(iii) and Lemma 3.2(iii) so assume from now on that $c \neq d^{\pm 1}$. If $c \neq d^2$ and $d \neq c^2$ then \mathcal{P} is aspherical by the above argument. If $c = d^2$ and $d = c^2$ then $c^3 = 1$, a contradiction, so assume that $c = d^2$ and $d \neq c^2$. Then $c^*(\hat{\Delta}) \leq 0$ as above except when $l(\hat{\Delta}) = b^{-1}cad^{-2}$ and $\hat{\Delta}$ is given by Figure 5.9(ii). In this case add $c^*(\hat{\Delta}) = \frac{\pi}{2}$ to $c(\hat{\Delta}_1)$ as shown noting that $\hat{\Delta}_1$ does not receive any curvature from Δ . Therefore if a region receives positive curvature across an edge it is across a $b^{-1}c$ or ad^{-1} or $a^{-1}c^{-1}$ edge; and if across consecutive edges then the sublabel is $(da^{-1}c^{-1})^{\pm 1}$. It follows that if $l(\hat{\Delta})$ does not involve $(da^{-1}c^{-1})^{\pm 1}$ and $d(\hat{\Delta}) \geq 7$ then $c^*(\hat{\Delta}) \leq 0$. Observe that each occurrence of $da^{-1}c^{-1}$ contributes two gaps. It follows that if $l(\hat{\Delta})$ contains at least two occurrences of $(da^{-1}c^{-1})^{\pm 1}$ or if $l(\hat{\Delta})$ contains exactly one occurrence and $d(\hat{\Delta}) \geq 8$ then there are four gaps and $c^*(\hat{\Delta}) \leq 0$. But if $l(\hat{\Delta}) = da^{-1}c^{-1}w$

and $d(\hat{\Delta}) = 7$ then checking possible $l(\hat{\Delta})$ shows that again there are four gaps and $c^*(\hat{\Delta}) \leq 0$. If $5 \leq d(\hat{\Delta}_1) \leq 6$ then either there is a contradiction to one of our assumptions or $c^*(\hat{\Delta}_1) \leq 0$ except when $l(\hat{\Delta}_1) \in \{a^{-1}c^{-2}ad^{-1}, a^{-1}c^{-2}bd^{-1}, a^{-1}c^{-1}ad^2, a^{-1}c^{-1}bd^2, a^{-1}c^{-3}bd\}$. If $d = c^{-2}$ or $d = c^3$ then $|c| = 5$ and there is a sphere by Lemma 3.2(vii). This leaves $l(\hat{\Delta}_1) \in \{a^{-1}c^{-1}ad^2, a^{-1}c^{-1}bd^2\}$ and these are given by Figure 5.9(iii), (iv). Add $c^*(\hat{\Delta}_1) = \frac{\pi}{2}$ to $c(\hat{\Delta}_2)$ as shown. Repeat the argument for $\hat{\Delta}_2$ noting that as before $\hat{\Delta}_2$ does not receive positive curvature from the corresponding region Δ . This procedure will terminate at a region $\hat{\Delta}_k$, say, such that $\hat{\Delta}_i \neq \hat{\Delta}_j$ for $i \neq j$ where $1 \leq i, j \leq k$ and $c^*(\hat{\Delta}_k) \leq 0$. Repeat this for each copy of the region $\hat{\Delta}_1$ to conclude that \mathcal{P} is aspherical. If $d = c^2$ and $c \neq d^2$ then the argument is the same by symmetry.

In conclusion \mathcal{P} is aspherical in this case except when either $c = d^{\pm 1}$ or $c^5 = 1$ and $c = d^2$ or $d^5 = 1$ and $d = c^2$.

(B8) $|c| = 2$, $|d| > 3$, $a^{-1}b = 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 2$ then Δ is given by Figures 5.1(ii) and 5.9(i). In Figure 5.9(i) $l(\Delta_1) = b^{-1}c\omega$ and $l(\Delta_2) = ad^{-1}\omega$. This implies that $d(\Delta_1) > 4, d(\Delta_2) > 4$, otherwise there is a contradiction to $|d| > 3$ so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. In Figure 5.1(ii) $l(\Delta_1) = l(\Delta_2) = bd\omega$. This similarly implies that $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$, so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. Observe that in Figure 5.9(i) Δ_1 does not receive positive curvature from Δ_4 ; and Δ_2 does not receive positive curvature from Δ_4 . Observe also that if $\hat{\Delta}$ receives positive curvature then it does so across the edges $b^{-1}c$, ad^{-1} or bd . Thus there is always a gap immediately preceding c^{-1} and a ; and there is a gap after c and a^{-1} . This implies that if there are at least four occurrences of $c^{\pm 1}$ then $l(\hat{\Delta})$ contains at least four gaps and so $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.6. We will proceed according to the number of occurrences of $c^{\pm 1}$ in $l(\hat{\Delta})$. If there are no occurrences of $c^{\pm 1}$ then either $l(\hat{\Delta}) = (ab^{-1})^k$ where $|k| \geq 2$ and there are four gaps, or $l(\hat{\Delta}) = d(ab^{-1})^{k_1} \dots d(ab^{-1})^{k_m}$ where $k_i \in \mathbb{Z}$ ($1 \leq i \leq m$). But since there is always at least one gap between any two occurrences of $d^{\pm 1}$ it follows that again there are four gaps or $|d| \leq 3$, a contradiction. So $c^*(\hat{\Delta}) \leq 0$ in this case.

Assume first that H is non-cyclic. If there is exactly one occurrence of $c^{\pm 1}$ in $l(\hat{\Delta})$ then H is cyclic so suppose that there are either two or three occurrences of $c^{\pm 1}$. Then either the label contains at least four gaps or it contradicts one of the **(B8)** assumptions or one of the following cases $\hat{\Delta}_i$ ($1 \leq i \leq 9$) occurs:

- (1) $cad^{-1}b^{-1}cad^{-1}b^{-1}$;
- (2) $cad^{-1}b^{-1}cbd^{-1}b^{-1}$;
- (3) $cad^{-1}b^{-1}c^{-1}ad^{-1}b^{-1}$;
- (4) $cad^{-1}b^{-1}c^{-1}bd^{-1}b^{-1}$;
- (5) $cad^{-1}b^{-1}cbda^{-1}$;
- (6) $cad^{-1}b^{-1}cbdb^{-1}$;




Figure 5.10: curvature distribution for Case (B8)

$$(7) \quad cad^{-1}b^{-1}c^{-1}bda^{-1};$$

$$(8) \quad cad^{-1}b^{-1}c^{-1}bdb^{-1};$$

$$(9) \quad (bda^{-1}c^{-1})^3.$$

If any of (1)–(4) occurs with any of (5)–(8) or with (9) then $|d| = 2$, a contradiction. Also if any of (5)–(8) occurs with (9) then $c = d^3$ and H is cyclic, so assume otherwise.

Consider (1)–(4). These yield the relator $(cd)^2$ and it follows that $cd^k = d^{-k}c$ for $k \in \mathbb{Z}$. Moreover if $|d| < \infty$ then there is a sphere by Lemma 3.1(c)(ii) so it can be assumed that $|d| = \infty$. In case (1) $\hat{\Delta}_1$ is given by Figure 5.10(i) where, given that $c^*(\hat{\Delta}_1) > 0$, it can be assumed that $d(\Delta_1) = d(\Delta_2) = 2$ and at least one of $d(\Delta_3)$, $d(\Delta_4)$ equals 2. Add $\frac{\pi}{2}$ from $c(\hat{\Delta}_1)$ to $c(\hat{\Delta}_{10})$ as shown in Figure 5.10(i); and if $d(\Delta_3) = d(\Delta_4) = 2$ add a further $\frac{\pi}{2}$ of $c(\hat{\Delta}_1)$ to $c(\hat{\Delta}_{12})$ as shown. In cases (2)–(4) $c^*(\hat{\Delta}) \leq \frac{\pi}{2}$ where $\hat{\Delta} \in \{\hat{\Delta}_2, \hat{\Delta}_3, \hat{\Delta}_4\}$ and $\frac{\pi}{2}$ is added from $c(\hat{\Delta})$ to $c(\hat{\Delta}_{10})$ as shown in Figure 5.10(ii). Observe that $x \neq b$ in Figure 5.10(i), (ii) for otherwise c^2 would be a proper sublabeled, and so $x \in \{a, d\}$. If $x = a$ then the sublabeled ad yields a gap so let $x = d$. Then either dd yields a gap or $\hat{\Delta}_{11} \in \{\hat{\Delta}_i : 1 \leq i \leq 4\}$ and $\frac{\pi}{2}$ is added to $c(\hat{\Delta}_{10})$ from $c(\hat{\Delta}_{11})$. Continuing this way, since $|d| = \infty$, eventually we

get a sublabel ad or dd which contributes a gap. Consider $l(\hat{\Delta}_{10})$. If it contains an odd number of occurrences of c then $cd^k = d^{-k}c$ implies that $c \in \langle d \rangle$ and H is cyclic. This leaves the case when there are exactly two occurrences of c and $cd^{\alpha_1}cd^{\alpha_2} = 1$ for $\alpha_1, \alpha_2 \in \mathbb{Z} \setminus \{0\}$. If $|\alpha_1|, |\alpha_2| > 1$ then there are four gaps and $c^*(\hat{\Delta}_{10}) \leq 0$; and if $|\alpha_1| > 1, |\alpha_2| = 1$ this implies $|d| < \infty$, a contradiction.

Consider (5)–(8). These yield the relator $cdcd^{-1}$ and H is Abelian. Observe that $|d| = 4$ yields **(E)**, so assume otherwise. In each case add $c^*(\hat{\Delta}) = \frac{\pi}{2}$ to $c(\hat{\Delta}_{13})$ as shown in Figure 5.10(iii)–(vi). Observe that $\hat{\Delta}_{13}$ receives no curvature from Δ_1 or Δ_2 ; that $l(\hat{\Delta}_{13}) = adw$ implies $d(\hat{\Delta}_{13}) > 4$ otherwise there is a contradiction to $|d| > 3$; and there is still a gap between each pair of occurrences of d . If $l(\hat{\Delta}_{13})$ contains an odd number of occurrences of c then H is cyclic so it can be assumed that $l(\hat{\Delta}_{13})$ yields the relator $cd^{\beta_1}cd^{\beta_2}$. If $|\beta_1| > 1$ and $|\beta_2| > 1$ then there are four gaps and if $(\beta_1, \beta_2) \in \{(2, 1), (2, -1), (1, 1)\}$ then $|d| \leq 3$, so this leaves the case $\beta_1 = 1, \beta_2 = -1$. Again there are four gaps except when $l(\hat{\Delta}_{13}) = adb^{-1}cad^{-1}b^{-1}c$ and this is shown in Figure 5.10(vii): add $c^*(\hat{\Delta}_{13}) = \frac{\pi}{2}$ to $c(\hat{\Delta}_{14})$ and observe that $\hat{\Delta}_{14}$ does not receive positive curvature from Δ . Consider $l(\hat{\Delta}_{14}) = bddw$. If there are at least four occurrences of c then $c^*(\hat{\Delta}_4) \leq 0$; and if there is an odd number of occurrences then H is cyclic. Suppose firstly that there are no occurrences of c in $l(\hat{\Delta}_{14})$. Since $|d| \geq 5$, if there is one occurrence of b then $l(\hat{\Delta}_4) = a^{-1}bd^k$ ($k \geq 5$) and there are four gaps; and since each $(a^{-1}b)^{\pm 1}$ yields a gap and each $(bd^l)^{\pm 1}$ ($l \geq 2$) yields a gap it follows that if there are at least two occurrences of b then again $c^*(\hat{\Delta}_{14}) \leq 0$. Suppose finally that there are two occurrences of c and so $cd^{\beta_1}cd^{\beta_2} = 1$ where $\beta_1 \geq 2$ and $|\beta_2| \geq 0$. If $|\beta_2| > 1$ then there are four gaps; and if $|\beta_2| = 1$ then $\beta_1 \geq 4$, otherwise there is a contradiction to $|d| > 4$, and again there are four gaps, so $c^*(\hat{\Delta}_{14}) \leq 0$.

Finally consider case (9). In this case $\hat{\Delta}_9$ is given by Figure 5.10(viii). Suppose that $c^*(\hat{\Delta}_9) > 0$. Then it can be assumed that $d(\Delta_i) = 2$ for $1 \leq i \leq 6$ and $c^*(\hat{\Delta}) = \frac{\pi}{2}$ so add $\frac{1}{3}c^*(\hat{\Delta}) = \frac{\pi}{6}$ to $c(\hat{\Delta}_l)$ for $l \in \{15, 16, 17\}$. In this case if $|d| \in \{4, 5\}$ then we obtain a sphere by Lemma 3.1(c)(iii). Now if $|d| \geq 6$ then as shown in Figure 5.10(viii) $l(\Delta_l) = d^{-2}\omega$ and d^{-2} will contribute two-thirds of a gap. If there are now at least two occurrences of c then either $|d| < 6$ or H is cyclic, a contradiction, or there are four gaps; if there is exactly one occurrence of c then this contradicts H non-cyclic; and if there are no occurrences of c then $l(\Delta_l) = d^{k_1}(b^{-1}a)^{m_1} \dots d^{k_n}(b^{-1}a)^{m_n}$ where $m_i \in \mathbb{Z}, k_i \geq 1$. Since $k_1 + \dots + k_n \geq 6$ it follows that there are at least four gaps and $c^*(\hat{\Delta}_l) \leq 0$.

Now let H be cyclic. If $c = d^2$ or $c = d^3$ then there is a sphere by T -equivalence and Lemma 3.2(vii), (viii); and $c = d^4$ is **(E4)**, so assume otherwise. In particular, $|d| > 4$. We follow the same argument as above and so if $l(\hat{\Delta})$ contains no occurrences or at least four occurrences of c then, as before, $c^*(\hat{\Delta}) \leq 0$; and if $l(\hat{\Delta})$ contains an odd number of occurrences of c then $c = d^k$ for some $k \geq 4$ which implies there are at least four gaps and $c^*(\hat{\Delta}) \leq 0$. Suppose then that $l(\hat{\Delta})$ involves c exactly twice. Subcases (1)–(4) imply $d^2 = 1$ and (9) implies $c = d^3$, a contradiction. This leaves subcases (5)–(8).

Add $c^*(\hat{\Delta}) = \frac{\pi}{2}$ to $c(\hat{\Delta}_{13})$ as in Figure 5.10(iii)–(vi). Since there is still a gap between each pair of occurrences of d it follows from the above paragraph and the previous argument




Figure 5.11: curvature distribution for Case (B9)

that $c^*(\hat{\Delta}_{13}) \leq 0$ except when $l(\hat{\Delta}_{13}) = adb^{-1}cad^{-1}b^{-1}c$. Again add $c^*(\hat{\Delta}_{13}) = \frac{\pi}{2}$ to $c(\hat{\Delta}_{14})$ as shown in Figure 5.10(vii). If $l(\hat{\Delta}_{14}) = bddw$ involves at least three occurrences of c then, since $\hat{\Delta}_4$ does not receive positive curvature from Δ in Figure 5.10(vii), there are at least four gaps and $c^*(\hat{\Delta}_4) \leq 0$. Otherwise checking the possible labels for $l(\hat{\Delta}_4) = bddw$ shows that there are four gaps or a contradiction to $|d| > 4$ or $c \notin \{d^3, d^4\}$.

In conclusion \mathcal{P} is aspherical except when $|cd| = 2$, $|d| < \infty$ or when $|cd| = 3$, $|d| \in \{4, 5\}$ or when H is cyclic and $c = d^2$ or d^3 .

(B9) $|c| = 3$, $|d| > 3$, $a^{-1}b = 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $d(\Delta) = 2$ then Δ is given by Figure 5.9(i). If $d(\Delta) = 3$ then Δ is given by Figure 4.1(ii). In Figure 5.9(i) $l(\Delta_1) = b^{-1}c\omega$ and $l(\Delta_2) = ad^{-1}\omega$ which implies that $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ otherwise there is a contradiction to $|d| > 3$ or $|c| = 3$, so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. In Figure 4.1(ii) $l(\Delta_1) = l(\Delta_3) = l(\Delta_5) = bdw$ which implies that $d(\Delta_i) > 4$ where $i \in \{1, 3, 5\}$ otherwise there is a contradiction to $|d| > 3$ so add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_i)$. Observe that if $\hat{\Delta}$ receives positive curvature then it receives $\frac{\pi}{2}$ across the edges $b^{-1}c$, ad^{-1} or receives $\frac{\pi}{6}$ across the edge bd . Thus there is always a gap immediately preceding c^{-1} and a ; and there is a gap after c and a^{-1} . Also there is a two thirds of a gap across the edge bd . This implies that if there are at least four occurrences of $c^{\pm 1}$ then $l(\hat{\Delta})$ contains at least four gaps and so $c^*(\hat{\Delta}) \leq 0$ by Lemma 3.6. We will proceed according to the number of occurrences of $c^{\pm 1}$ in $l(\hat{\Delta})$. If there are no occurrences of $c^{\pm 1}$ then either $l(\hat{\Delta}) = (ab^{-1})^k$ where $|k| \geq 3$ and there are four gaps, or $l(\hat{\Delta}) = d(ab^{-1})^{k_1} \dots d(ab^{-1})^{k_m}$ where $k_i \in \mathbb{Z}$ ($1 \leq i \leq m$). But since there is always at least one gap between any two occurrences of $d^{\pm 1}$ it follows that again there are four gaps or $|d| \leq 3$, a contradiction. So $c^*(\hat{\Delta}) \leq 0$ in this case. On the other hand if there are between one and three occurrences of $c^{\pm 1}$ inclusive and $c^*(\hat{\Delta}) > 0$ then either there is a contradiction to one of the **(B9)**

assumptions or $c \in \{d^{\pm 2}, d^3\}$ or $l(\hat{\Delta}) = cad^{-1}b^{-1}cad^{-1}b^{-1}$. If $c = d^{\pm 2}$ then **(E2)** or **(E3)** occurs, so assume otherwise.




Figure 5.12: curvature distribution for Cases (B11) and (B12)

First suppose that $l(\hat{\Delta}) = cad^{-1}b^{-1}cad^{-1}b^{-1}$ (in particular, H is non-cyclic) as in Figure 5.11(i). If $c^*(\hat{\Delta}) > 0$ then it can be assumed that $d(\Delta_i) = 2$ for $1 \leq i \leq 4$ and $c^*(\hat{\Delta}) = \frac{\pi}{3}$, so add $\frac{1}{2}c^*(\hat{\Delta}) = \frac{\pi}{6}$ to $c(\hat{\Delta}_j)$ for $j \in \{1, 2\}$. Therefore $d^{\pm 2}$ contributes a two-thirds gap. If $|d| \in \{4, 5\}$ then we obtain a sphere by Lemma 3.1(c)(iv). Assume now that $|d| \geq 6$ and without any loss of generality assume that $\hat{\Delta}_l \neq \hat{\Delta}$ of Figure 5.11(i). If $l(\hat{\Delta}_l)$ has at least two occurrences of c then either there is a contradiction to H non-cyclic or one of the **(B9)** assumptions or there are four gaps and $c^*(\hat{\Delta}_l) \leq 0$; if there is exactly one occurrence of c then this contradicts H non-cyclic; and if there are no occurrences of c then $l(\hat{\Delta}_l) = d^{k_1}(b^{-1}a)^{m_1} \dots d^{k_n}(b^{-1}a)^{m_n}$ where $m_i \in \mathbb{Z}$, $k_i \geq 1$. Since $k_1 + \dots + k_n \geq 6$ it follows that there are at least four gaps and $c^*(\hat{\Delta}_l) \leq 0$.

Now let $c = d^3$. It follows from the above argument and checking possible vertex labels that if $\hat{\Delta}_1$ receives positive curvature then $c^*(\hat{\Delta}_1) \leq 0$ except when $l(\hat{\Delta}_1) = c^{-1}bd^3a^{-1}$ and $c^*(\hat{\Delta}) = \frac{\pi}{6}$. Add $c^*(\hat{\Delta}_1)$ to $c(\hat{\Delta}_2)$ as shown in Figure 5.11(ii) (where we note that $\hat{\Delta}_2$ receives no curvature from $\hat{\Delta}$). Checking $\{d^{-1}, a, b\}d^{-2}w$ with the understanding that there is a gap preceding the d^{-2} and each d^{-2} contributes a two-thirds gap it follows that either there is a contradiction or $c^*(\hat{\Delta}_2) \leq 0$ except when $l(\hat{\Delta}_2) = l(\hat{\Delta}_1)^{-1} = cad^{-3}b^{-1}$. If $\hat{\Delta}_2$ is given by Figure 5.11(iii) then $c^*(\hat{\Delta}_2) \leq c(\hat{\Delta}_2) + \frac{\pi}{2} + 2(\frac{\pi}{6}) < 0$, so let $\hat{\Delta}_2$ be given by Figure 5.11(iv). Observe that the neighbour Δ_1 has label $c^{-2}bw$ and no longer has degree 3 and that if either $d(\Delta_2) > 2$ or $d(\Delta_3) > 2$ then $c^*(\hat{\Delta}) < 0$ and so Figure 5.11(iv) shows the only possibility for $c^*(\hat{\Delta}_2) > 0$. Now repeat the argument for $\hat{\Delta}_2$ and $\hat{\Delta}_3$. The conditions on the neighbouring regions Δ_1 , Δ_2 and Δ_3 ensure that this procedure will terminate at a region $\hat{\Delta}_k$, say, such that $\hat{\Delta}_i \neq \hat{\Delta}_j$ for $i \neq j$ where $1 \leq i, j \leq k$ and $c^*(\hat{\Delta}_k) \leq 0$. Repeat this for each copy of the region $\hat{\Delta}_1$ to conclude \mathcal{P} is aspherical.

In conclusion \mathcal{P} is aspherical except when H is non-cyclic, $|c^{-1}d| = 2$ and $|d| \in \{4, 5\}$.

(B10) $|c| = 2$, $|d| = 2$, $a^{-1}b = 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If $|cd| < \infty$ then we obtain a sphere by Lemma 3.1(c)(i), so assume otherwise (in particular,

H is non-cyclic). Observe that if a label involves c and is not $c^{\pm 2}$ then it yields the relation $(cd)^k = 1$ and so it follows that if $c(\Delta) > 0$ then Δ is given by Figure 5.1(ii) in which $l(\Delta_1) = l(\Delta_2) = bdw$. Add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$ as shown. This completes the distribution of curvature. Consider Δ_1 of Figure 5.1(ii). Since positive curvature only crosses the edge $(bd)^{\pm 1}$ it follows that if $l(\Delta_1)$ involves at least four occurrences of d then $c^*(\Delta_1) \leq 0$. This leaves the case when $l(\Delta_1)$ involves exactly two occurrences of d . But d^2 cannot be a sublabel since there would then be a region with label caw , and so $l(\Delta_1) = bd\{a^{-1}b, b^{-1}a\}da^{-1}$ has four gaps and $c^*(\Delta_1) \leq 0$. In conclusion \mathcal{P} is aspherical in this case if and only if $|cd| = \infty$.

(B11) $|c| = 2$, $|d| = 3$, $a^{-1}b = 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If H is cyclic then there is a sphere by Lemma 3.2(vi), so assume otherwise. If $d(\Delta) = 2$ then Δ is given by Figures 5.1(ii) and 5.9(i); and if $d(\Delta) = 3$ then Δ is given by Figure 4.3(i). In Figure 5.9(i) $l(\Delta_1) = b^{-1}c\omega$ and $l(\Delta_2) = ad^{-1}\omega$. This implies that $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ otherwise there is a contradiction to $|d| = 3$ so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. In Figure 5.1(ii) $l(\Delta_1) = l(\Delta_2) = bdw$. This implies that $d(\Delta_1) > 4$ and $d(\Delta_2) > 4$ otherwise there is a contradiction to $|d| = 3$ so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. In Figure 4.3(i) $l(\Delta_1) = l(\Delta_3) = l(\Delta_5) = ca\omega$. This implies that $d(\Delta_1) > 4$, $d(\Delta_3) > 4$ and $d(\Delta_5) > 4$ otherwise there is a contradiction to $|d| = 3$ so add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_1)$, $c(\Delta_3)$ and $c(\Delta_5)$.

Observe that if $\hat{\Delta}$ receives positive curvature then it does so across the edges $b^{-1}c$, ad^{-1} , bd , ca and there is a two-thirds gap in ca . If there is only one occurrence of c or d then this contradicts H non-cyclic. Also if $|cd| \in \{2, 3, 4, 5\}$ then we obtain a sphere by Lemma 3.1(c)(ii),(v). So assume that $|cd| > 5$. If there are between two and five occurrences of c inclusive then the resulting relator contradicts H non-cyclic or $|cd| > 5$ except for $(cd)^2(cd^{-1})^2$ and $(cdcd^{-1})^2$. But each of these yields four gaps. Since at least six occurrences of c yields four gaps, this leaves the case when c does not occur and this implies that there is at least one gap between any two occurrences of d . It follows that there can only be exactly three occurrences of d . Checking shows there are four gaps except when the following subcases occur:

- (1) $l(\hat{\Delta}) = bddda^{-1}$;
- (2) $l(\hat{\Delta}) = bd^2a^{-1}bda^{-1}$;
- (3) $l(\hat{\Delta}) = bda^{-1}bda^{-1}bda^{-1}$.

Consider (1), this case is given by Figure 5.12(i). If $d(\Delta_2) > 2$ then $c^*(\hat{\Delta}) \leq 0$. Assume otherwise and add $c^*(\hat{\Delta}) = \frac{\pi}{2}$ to $c(\hat{\Delta}_1)$ as shown. Observe that $l(\hat{\Delta}_1) = cax\omega$ and $x \in \{b^{-1}, d, d^{-1}\}$. If $x \in \{b^{-1}, d\}$ then $\hat{\Delta}_1$ cannot receive positive curvature across the edges ab^{-1} or ad . Suppose that $x = d^{-1}$. Then $\hat{\Delta}_1$ cannot receive positive curvature across the edge ad^{-1} otherwise $l(\Delta_1) \neq c^2$, a contradiction. So there will be a gap between c

and the next occurrence of d and the previous argument does not change. It follows that $c^*(\hat{\Delta}_1) \leq 0$.

Consider (2), this case is given by Figure 5.12(ii). If $d(\Delta_2) > 2$ or $d(\Delta_4) > 2$ then $c^*(\hat{\Delta}) \leq 0$. Assume otherwise and add $c^*(\hat{\Delta}) = \frac{\pi}{2}$ to $c(\hat{\Delta}_2)$ as shown. Observe that $l(\hat{\Delta}_2) = cax\omega$ and the same argument as for (1) can be applied to deduce that $c^*(\hat{\Delta}_2) \leq 0$.

Consider (3), this case is given by Figure 5.12(iii). If $d(\Delta_2) > 2$ or $d(\Delta_4) > 2$ or $d(\Delta_6) > 2$ then $c^*(\hat{\Delta}) \leq 0$. Assume otherwise and add $\frac{1}{3}c^*(\hat{\Delta}) = \frac{\pi}{6}$ to $c(\hat{\Delta}_i)$ for $i \in \{3, 4, 5\}$ as shown. Consider $l(\hat{\Delta}_3) = ad^{-1}x\omega$ where $x \in \{b^{-1}, a^{-1}, d^{-1}\}$. If $x = b^{-1}$ then $l(\hat{\Delta}_6) = c^{-2}bd\omega$, a contradiction. If $x = a^{-1}$ then exactly as above $l(\hat{\Delta}_3)$ either contains four gaps or there is a contradiction to either H non-cyclic, $|cd| > 5$ or one of the **(B11)** assumptions and so $c^*(\hat{\Delta}_3) \leq 0$. Finally let $x = d^{-1}$. If $l(\hat{\Delta}_3) \notin \{ad^{-3}b^{-1}, ad^{-2}b^{-1}ad^{-1}b^{-1}\}$ then again as above $c^*(\hat{\Delta}_3) \leq 0$ so assume otherwise. But if $l(\hat{\Delta}_3) = ad^{-3}b^{-1}$ then we are back in subcase (1) and if $l(\hat{\Delta}_3) = ad^{-2}b^{-1}ad^{-1}b^{-1}$ then we are back in subcase (2), the only difference being that $\hat{\Delta}$ in Figures 5.12(i), (ii) this time receives $\pi/6$ from Δ_2 rather than $\pi/2$.

In conclusion \mathcal{P} is aspherical except when H is non-cyclic and $|cd| \in \{2, 3, 4, 5\}$ or when H is cyclic.

(B12) $|c| = 3$, $|d| = 3$, $a^{-1}b = 1$, $c^{\pm 1}ab^{-1} \neq 1$, $d^{\pm 1}b^{-1}a \neq 1$.

If H is cyclic then $c = d^{\pm 1}$ and so there is a sphere by Lemma 3.1(a)(iii) or Lemma 3.2(iii), so assume otherwise. If $d(\Delta) = 2$ then Δ is given by Figure 5.9(i); and if $d(\Delta) = 3$ then Δ is given by Figures 4.1(ii) and 4.3(i). In Figure 5.9(i) $l(\Delta_1) = b^{-1}c\omega$ and $l(\Delta_2) = ad^{-1}\omega$. This implies that $d(\Delta_1) > 4$, $d(\Delta_2) > 4$, otherwise there is a contradiction to H non-cyclic or $|c| = 3$ or $|d| = 3$ so add $\frac{1}{2}c(\Delta) = \frac{\pi}{2}$ to $c(\Delta_1)$ and $c(\Delta_2)$. In Figure 4.1(ii) $l(\Delta_1) = l(\Delta_3) = l(\Delta_5) = bd\omega$. This implies that $d(\Delta_1) > 4$, $d(\Delta_3) > 4$ and $d(\Delta_5) > 4$ otherwise there is a contradiction to H non-cyclic or $|c| = 3$ or $|d| = 3$ so add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_1)$, $c(\Delta_3)$ and $c(\Delta_5)$. In Figure 4.3(i) $l(\Delta_1) = l(\Delta_3) = l(\Delta_5) = ca\omega$. This implies that $d(\Delta_1) > 4$, $d(\Delta_3) > 4$ and $d(\Delta_5) > 4$ otherwise there is a contradiction to H non-cyclic or $|c| = 3$ or $|d| = 3$ so add $\frac{1}{3}c(\Delta) = \frac{\pi}{6}$ to $c(\Delta_1)$, $c(\Delta_3)$ and $c(\Delta_5)$.

Observe that if $\hat{\Delta}$ receives positive curvature then it does so across the edges $b^{-1}c$, ad^{-1} , bd , ca and there is a two-thirds gap in bd and ca . If there is only one occurrence of c or d in $l(\hat{\Delta})$ then this contradicts H non-cyclic so assume otherwise. Also if $|c^{-1}d| = 2$ then we obtain a sphere as by Lemma 3.1(c)(iv). So assume that $|c^{-1}d| > 2$. Checking now shows that there is at least one gap between any two occurrences of c or between any two occurrences of d ; and there is at least a two-thirds gap between any two occurrences of c and d . It follows that if there are at least four occurrences of c or of d then $c^*(\hat{\Delta}) \leq 0$. Suppose there are three occurrences of c . If there are three occurrences of d then there are at least $6 \cdot \frac{2}{3} = 4$ gaps; and if there are two occurrences of d then there are four gaps or $l(\hat{\Delta})$ yields the relator $c^2d^{-1}cd^{-1}$ which forces H to be cyclic. Suppose there are two occurrences of c . If there are three occurrences of d then there are four gaps or $l(\hat{\Delta})$ yields the relator $c^{-1}dc^{-1}d^2$ which forces H to be cyclic; and if there are two occurrences of d then $l(\hat{\Delta})$ forces $|c^{-1}d| = 2$ or H cyclic or is one of $c\{a, b\}d\{a^{-1}, b\}c\{a, b\}d\{a^{-1}, b^{-1}\}$,

$c\{a, b\}d\{a^{-1}, b^{-1}\}c^{-1}\{a, b\}d^{-1}\{a^{-1}, b^{-1}\}$ and there are four gaps. This leaves the case when c does not occur. Checking $l(\hat{\Delta})$ shows that there is a contradiction to $|d| = 3$ or there are at least four gaps except when $l(\hat{\Delta}) = bddda^{-1}$ as shown in Figure 5.12(iv). In this case add $c^*(\hat{\Delta}) = \frac{\pi}{6}$ to $c(\hat{\Delta}_1)$ as shown. Since $\frac{\pi}{6}$ is distributed across ca it follows that $c^*(\hat{\Delta}_1) \leq 0$ as above.

In conclusion \mathcal{P} is aspherical except when H is cyclic or when H is non-cyclic and $|cd| = 2$.

It follows from (B1)–(B12) that either \mathcal{P} is aspherical or modulo T -equivalence one of the conditions of Theorem 1.1 (i)–(iii) or Theorem 1.2 (i), (ii), (iv)–(x) is satisfied and so Theorems 1.1 and 1.2 are proved for Case B.

References

- [1] Y G Baik, W A Bogley and S J Pride, On the asphericity of length four relative group presentations, *Int. J. Algebra Comput.* **7** (1997), 277–312.
- [2] W A Bogley and S J Pride, Aspherical relative presentations, *Proc. Edinburgh Math. Soc.* **35** (1992), 1–39.
- [3] D E Cohen, *Combinatorial group theory: a topological approach*, LMS Student Texts **14** (CUP) (1989).
- [4] D J Collins and J Huebschmann, Spherical diagrams and identities among relations, *Math. Ann.* **261** (1982), 155–183.
- [5] P J Davidson, On the asphericity of a family of relative group presentations, *Int. J. Algebra Comput.* **19** (2009), 159–185.
- [6] M Edjvet, Equations over groups and a theorem of Higman, Neumann and Neumann, *Proc. London Math. Soc.* **62** (1991), 563–589.
- [7] M Edjvet, On the asphericity of one-relator relative presentations, *Proc. Roy. Soc. Edinburgh Sec.* **124** (1994), 713–728.
- [8] M Edjvet and J Howie, The solution of length four equations over groups, *Trans. Amer. Math. Soc.* **326** (1991), 345–369.
- [9] The GAP Group: GAP – Groups, Algorithms and Programming; Version 4.6.4, 2013. <http://www.gap-system.org>.
- [10] N D Gilbert and J Howie, LOG groups and cyclically presented groups, *J. Algebra* **174** (1995), 118–131.
- [11] J Howie and V Metaftsis, On the asphericity of length five relative group presentations, *Proc. London Math. Soc.* **82** (2001), 173–194.

- [12] V Metaftsis, On the asphericity of relative group presentations of arbitrary length, Int. J. Algebra Comput. **13** (2003), 323–339.

Abd Ghafur Bin Ahmad
School of Mathematical Sciences
Faculty of Sciences and Technology
Universiti Kebangsaan Malaysia
43600 UKM, Bangi
Selangor Darul Ehsan, Malaysia
ghafur@ukm.my

Muna A Al-Mulla¹
Department of Mathematics and Statistics
King Faisal University
PO Box 400
Alahssa 31982, Kingdom of Saudi Arabia
malmulla@fku.edu.sa

Martin Edjvet
School of Mathematical Sciences
The University of Nottingham
University Park, Nottingham NG7 2RD, UK
martin.edjvet@nottingham.ac.uk

¹Research supported by Ministry of Education, Kingdom of Saudi Arabia