

August 30, 2018

## CONSTRUCTION OF RIGHT GYROGROUP STRUCTURES ON A NON-ABELIAN GROUP

AKHILESH CHANDRA YADAV

ABSTRACT. In this article we gave the notion of particular type of functions on a group termed as class assigned functions . Using class assigned functions, we have constructed several right gyrogroup structures on a given non-abelian group.

**Keywords:** Right transversals, Right loops, Gyrotransversals.  
**MSC:** 20N05.

### 1. INTRODUCTION

A gyrogroup is a grouplike structure that is defined in [3] along with a weaker structure called a left gyrogroup. In [4], Foguel and Ungar proved that any given group can be turned into a left gyrogroup which is, in turn, a gyrogroup if and only if the given group is central by a 2 – Engel group. The importance of left gyrogroups stems from the facts that (i) any gyrotransversal groupoid is a left gyrogroup, and (ii) any left gyrogroup is a twisted subgroup in a specified group. Let  $K$  be a group and let  $G$  be the inner product of the group  $K$  by  $Inn(K)$ , where  $Inn(K)$  is the inner automorphism group of  $K$  whose generic element  $\alpha_k$  denotes conjugation by  $k \in K$ . Then, the diagonal transversal  $D$  generated by  $K$  (in  $G$ ) is a subset of  $G$  given by

$$D = \{(k; \alpha_k) \mid k \in K\}.$$

Then the diagonal transversal  $D$  is a gyrotransversal (Theorem 3.2, [4]). Indeed this operation induces a binary operation  $\odot$  on  $K$  so that the groupoid  $(K, \odot)$  is a left gyrogroup (see, [4]). The operation  $\odot$  on  $K$  is given by  $x \odot y = x^2 y x^{-1}$ . It is convenient for us to study right transversals and right conjugation action. Thus, in our terminology,  $(K, \odot)$  is a right gyrogroup, where  $x \odot y = y^{-1} x y^2$ . In this article, we construct several right gyrogroup structures on a non-abelian group.

### 2. PRELIMINARIES

Let  $S$  be a right transversal to the subgroup  $H$  in a group  $G$ . Let  $g : S \rightarrow H$  be a map with  $g(e) = 1$ , where  $e$  denotes the identity of  $S$  with respect to transversal operation and 1 denotes the identity of

group  $H$ . Consider the right transversal  $S_g = \{g(x)x \mid x \in S\}$  to  $H$ . Then

**Proposition 2.1.** [2] *The right quasigroup  $(S_g, o')$  is isomorphic to the right quasigroup  $(S, o_g)$  where  $o_g$  is the operation given by  $x o_g y = x\theta g(y)oy$ .*

The operation  $o_g$  is known as the deformed operation on  $S$ .

**Definition 2.2.** [2] *Let  $(S, o)$  be a magma together with a right identity  $e$  and together with a right inverse  $a'$  for every element  $a \in S$  with respect to  $e$  in the sense that  $aoa' = e$ . Then  $(S, o)$  will be called a right gyrogroup if*

- (1) *for any  $x, y, z \in S$  there exists a unique element  $x\theta f(y, z) \in S$  such that*

$$(2.1) \quad (xoy)oz = x\theta f(y, z)o(yoz),$$

- (2) *the map  $f(y, z) : S \rightarrow S$  given by  $f(y, z)(x) = x\theta f(y, z)$  is an automorphism of  $(S, o)$  and*  
 (3) *for all  $y \in S$ ,*

$$(2.2) \quad f(y, y') = I_S.$$

**Definition 2.3.** [2] *A right transversal  $S$  to a subgroup  $H$  with identity  $e$  in a group  $G$  is called a gyrotransversal if  $S = S^{-1}$  and  $h^{-1}xh \in S, \forall x \in S$  and  $h \in H$ .*

**Proposition 2.4** ([2], Representation theorem for right gyrogroups). *A right loop  $(S, o)$  is a right gyrogroup if and only if it is a gyrotransversal to  $G_S$  in its group extension  $G_S S$ .*

In [2], we have shown the following:

**Lemma 2.5.** [2] *Let  $S$  be a gyrotransversal to a subgroup  $H$  in a group  $G$  and  $g$  be a map from  $S$  to  $H$  such that  $g(e) = e$ . Then the transversal  $S_g = \{g(x)x \mid x \in S\}$  is a gyrotransversal if and only if*

$$(2.3) \quad g(x^{-1}) = g(x)^{-1}$$

*and  $g$  is equivariant in the sense that*

$$(2.4) \quad g(h^{-1}xh) = h^{-1}g(x)h$$

*for all  $x \in S$  and  $h \in H$ .*

## 3. CONSTRUCTION OF RIGHT GYROGROUPS

**Definition 3.1.** Let  $G$  be a given non-abelian group. Define a binary operation  $\cdot$  on  $G \times G$  as

$$(a, x) \cdot (b, y) = (ab[b^{-1}xb, y^{-1}], b^{-1}xboy),$$

where  $xoy = y^{-1}xy^2$ . Then  $(G \times G, \cdot)$  is a group. The identity of the group  $G \times G$  is  $(1, 1)$ , where  $1$  denotes the identity of group  $G$  and the inverse of  $(a, x)$  is  $(a^{-1}, a^{-1}x^{-1}a)$ , for each  $(a, x)$  in  $G \times G$ .

One may easily observe that the group  $G \times G$  contains  $G \times \{1\} (\cong G)$  as a subgroup and  $\{1\} \times G$  as a gyrotransversal to the subgroup  $G \times \{1\}$  in the group  $G \times G([1, 2])$ .

**Proposition 3.2.** Let  $n \in \mathbb{N}$  and  $g$  be a map from a group  $G$  into itself with  $g(1) = 1$  and  $g(x) = x^n$  for each  $x \in G$ . Then  $S_g = \{(g(x), x) \mid x \in G\}$  is a gyrotransversal to the subgroup  $H (= G \times \{1\})$  in  $G \times G$ .

PROOF: Since  $(x^{-1})^n = (x^n)^{-1}$  and  $(h^{-1}xh)^n = h^{-1}x^nh$  for all  $x, h \in G$ . Thus the result follows from Lemma 2.5.  $\square$

Let  $G$  be a group. Let  $R$  be a relation on  $G$  given by  $aRb$  if either  $a = b$  or  $a = b^{-1}$  or  $a \sim b$  or  $a \sim b^{-1}$ , where  $\sim$  is a conjugacy relation on group  $G$ . Then  $R$  is an equivalence relation on  $G$ . The equivalence class of  $w \in G$  is denoted by  $[w]_R$ .

**Definition 3.3.** Let  $G$  be a group. A function  $k : G \rightarrow \mathbb{N} \cup \{0\}$  is called a class assigned function on  $G$  if  $k([1]_R) = 0$  and  $k$  is constant on each class  $[w]_R$ .

**Proposition 3.4.** Let  $k$  be a class assigned function on a free group  $F$ . Then it determines a function  $g : F \rightarrow F$  given by  $g(w) = w^{k(w)}$  so that the corresponding transversal  $S_g = \{(g(w), w) \mid w \in F\}$  to the subgroup  $H = F \times \{1\}$  in  $F \times F$  is a gyrotransversal.

PROOF: By definition of  $k$ ,  $k(w) = k(w^{-1}) = k(u^{-1}wu)$ , for all  $u, w$  in  $F$ . Therefore  $g(w^{-1}) = g(w)^{-1}$  and  $g(u^{-1}wu) = (u^{-1}wu)^{k(u^{-1}wu)} = u^{-1}w^{k(u^{-1}wu)}u = u^{-1}w^{k(w)}u = u^{-1}g(w)u$ . By Lemma 2.5,  $S_g$  is a gyrotransversal.  $\square$

**Proposition 3.5.** Let  $g : F \rightarrow F$  be a map satisfying the property that

- (i).  $g(1) = 1$ ,
- (ii).  $g(w^{-1}) = g(w)^{-1}$ ,
- (iii).  $g(u^{-1}wu) = u^{-1}g(w)u$ , for all  $u, w \in F$ .

Then  $g$  determines a class assigned function  $k$  on free group  $F$  such that  $g(w) = w^{k(w)}$ .

PROOF: For  $w \in F \setminus \{1\}$ , (iii) gives  $g(w) = w^{-1}g(w)w$ . Thus  $g(w)$  belongs to the centralizer of  $w$ . But  $F$  is a free group, therefore  $g(w) = w^{k(w)}$ , for unique  $k(w) \in \mathbb{N} \cup \{0\}$ . Then (ii) and (iii) implies that  $k(w^{-1}) = k(w)$  and  $k(u^{-1}wu) = k(w)$  for all  $u, w \in F$ . Define  $k(1) = 0$ , thus  $k(w)$  determines a class assigned function  $k$ .  $\square$

**Corollary 3.6.** *There is a bijective correspondence between the set of all class assigned functions and the set of all maps  $g$  described as in the Prop. (3.5). In particular, there is a bijective correspondence between the set of all class assigned functions and the set of all gyrotransversals to the subgroup  $H$  in  $F \times F$ .*

**Theorem 3.7.** *Let  $G$  be a non-abelian group and  $k$  be a class assigned function on  $G$ . Then it determines a function  $g : G \rightarrow G$  given by  $g(w) = w^{k(w)}$  such that  $S_g$  is a gyrotransversal to  $G \times \{1\}$  in  $G \times G$ . Define a binary operation  $o_k$  on  $G$  by*

$$(3.1) \quad x o_k y = y^{-k(y)} x y^{k(y)+1}.$$

*Then  $(G, o_k)$  is a right gyrogroup.*

PROOF: By the property of class assigned function  $k$ ,  $g$  satisfies  $g(x^{-1}) = g(x)^{-1}$  and  $g(h^{-1}xh) = h^{-1}g(x)h$  for all  $x, h$  in  $G$ . Thus  $S_g$  is a gyrotransversal to  $G \times \{e\}$  in the group  $G \times G$  and so deformed operation  $o_k$  on  $G$  is given by Equation 3.1. Observed that  $y^{-1}o_k y = e = y o_k y^{-1}$ . By a little computation the map  $f(y, z)(x)$  determined by the equation  $(xoy)oz = f(y, z)(x)o(yoz)$  is  $f(y, z)(x) = x\theta [y^{k(y)}(yz)^{-k(y o_k z)}z^{k(z)}]$ , where  $x\theta a = a^{-1}xa$  for all  $x, a \in G$ . Clearly  $f(y, z)$  is an inner automorphism of  $(G, o_k)$ . Also  $f(y, y^{-1})(x) = x = f(y^{-1}, y)(x)$  and so  $f(y, y^{-1}) = I_G$ . Thus,  $(G, o_k)$  is a right gyrogroup.  $\square$

**Remark 3.8.** *We do not know if its converse true.*

**Example 3.2.** *Take  $G = S_3$ , the symmetric group of degree 3. Let  $k$  be a class assigned function on  $S_3$ .*

*If  $k(x) = 0$  for all  $x$  in  $S_3$ . Then the corresponding right gyroroup  $(S_3, o_k)$  is a group  $S_3$  itself.*

*If  $k(I) = 0, k(x) = 1$  for all  $x \in S_3 \setminus \{I\}$ . Then the group generated by automorphisms  $f(y, z)$  is isomorphic to  $[S_3, S_3] = A_3$ .*

*If  $k(I) = 0, k((12))=k((2\ 3))=k((1\ 3))=1, k((1\ 2\ 3))=k((1\ 3\ 2))=2$ . Then  $f((12), (132))(x) = x\theta(13)$ ,  $f((12), (123))(x) = x\theta(23)$  and  $f((23), (132))(x) = x\theta(12)$ . Thus,  $(S_3, o_k)$  is a right gyrogroup for which the group generated by automorphisms  $f(y, z)$  is isomorphic to  $S_3$ .*

The above right gyrogroups are non-isomorphic. Note that the right gyrogroups determined by different  $k$  may be isomorphic. For, let  $k_1$  and  $k_2$  be class assigned maps given by  $k_1(x) = 0$  for all  $x \in S_3$  and  $k_2(I) = 0, k_2(x) = 6$  for all  $x \in S_3 \setminus \{I\}$  respectively. Clearly they are distinct but the corresponding right gyrogroups are the group  $S_3$ .

**Corollary 3.9.** For  $n \geq 3$ , there are

$$\prod_{\sum_{i=1}^k n_i \leq n}^{n_i > 1} [n_1, \dots, n_k]$$

gyrotransversals to the subgroup  $S_n$  in the generalized product of group  $S_n$ . Here  $[n_1, \dots, n_k]$  denotes the least common multiple of  $n_1, \dots, n_k$ .

PROOF: Let  $p, q \in S_n$ . Since  $p$  and  $p^{-1}$  both have same cycle structure and  $p, q$  are conjugates to each other if and only if they have same cycle structures. Therefore for each  $w \in S_n$ , the equivalence class  $[w]_R$  is same as the conjugacy class of  $w$ . Since the class assigned function  $k(w)$  has  $|w|$  choices on each class  $[w]_R$  and the choices are independent on these classes. Therefore the number of class assigned functions  $k$  will be  $\prod |w|$ , where the product runs over each classes with  $w \neq 0$ . Let  $n_1, n_2, \dots, n_k$  be natural numbers greater than 1 such that  $n_1 + n_2 + \dots + n_k \leq n$ . Then order of permutation with cycle structure  $(n_1, n_2, \dots, n_k)$  is  $[n_1, n_2, \dots, n_k]$ , the least common multiple of  $n_1, n_2, \dots, n_k$ . Thus, by above proposition the number of gyrotransversals to  $S_n$  in the generalized product  $S_n \times S_n$  will be

$$\prod_{\sum_{i=1}^k n_i \leq n}^{n_i > 1} [n_1, \dots, n_k]$$

□

## REFERENCES

- [1] R. Lal, *Transversals in groups*, Journal of algebra, **181**, 70-81(1996).
- [2] R. Lal and A. C. Yadav, *Topological right gyrogroups and gyrotransversals*, Communication in Algebra, 41:9, 3559-3575 (2013).
- [3] T. Foguel, A.A. Ungar, *Involutory decomposition of groups into twisted subgroups and subgroups*, J. Group theory, **3**, 27-46 (2000).
- [4] T. Foguel, A.A. Ungar, *Gyrogroups and the decomposition of groups into twisted subgroups and subgroups*, Pac. J. of Maths. Vol. 197, No. 1, (2001).