

A NOTE ON THE AFFINE-INVARIANT PLANK PROBLEM

GREGORY R. CHAMBERS AND LAWRENCE MOUILLÉ

ABSTRACT. Suppose that C is a bounded, convex subset of \mathbb{R}^n , and that P_1, \dots, P_k are planks which cover C in respective directions v_1, \dots, v_k and with widths w_1, \dots, w_k . In 1951, Bang conjectured that

$$\sum_{i=1}^k \frac{w_i}{w_{v_i}(C)} \geq 1,$$

generalizing a previous conjecture of Tarski. Here, $w_{v_i}(C)$ is the width of C in the direction v_i .

In this note we give a short proof of this conjecture under the assumption that, for every m with $1 \leq m \leq k$, $C \setminus \bigcup_{i=1}^m P_i$ is a convex set. In addition, we prove that if the projection of C onto the vector space spanned by the normal vectors of the planks has dimension d , then

$$\sum_{i=1}^k \frac{w_i}{w_{v_i}(C)} \geq \frac{1}{d}.$$

A *plank* P is the closed, connected region between two parallel hyperplanes in \mathbb{R}^n . The distance between these supporting hyperplanes is called the *width* of P . We define the *direction* of P to be the one-dimensional subspace v of \mathbb{R}^n which is perpendicular to the supporting hyperplanes of P . In 1932, Tarski proved in [6] that if P_1, \dots, P_k are planks with widths w_1, \dots, w_k which cover a unit ball in \mathbb{R}^2 , then

$$\sum_{i=1}^k w_i \geq 2.$$

He asked if an analogous result was true for every bounded, convex subset C of \mathbb{R}^n . For any one-dimensional subspace v of \mathbb{R}^n , we define the width of C in the direction v , $w_v(C)$, as the width of the smallest plank of direction v which covers C . In other words, if we choose hyperplanes perpendicular to v , which do not intersect the interior of C , and which are minimally separated, then $w_v(C)$ is the distance between those hyperplanes. Tarski's question can be thus stated as follows: If P_1, \dots, P_k are planks which cover C with widths w_1, \dots, w_k ,

then does

$$\sum_{i=1}^k w_i \geq \inf_{v \in G(n,1)} w_v(C)$$

hold? Here, $G(n,1)$ is the Grassmannian of one-dimensional linear subspaces in \mathbb{R}^n .

This conjecture was resolved in 1951 by Bang in [3]. In the same article, Bang posed the following well-known affine-invariant generalization of Tarski's problem:

Conjecture 1 (Bang [3]). Suppose that C is a convex, bounded subset of \mathbb{R}^n , and that P_1, \dots, P_k are planks in directions v_1, \dots, v_k and of widths w_1, \dots, w_k which cover C . Then

$$\sum_{i=1}^k \frac{w_i}{w_{v_i}(C)} \geq 1.$$

If $w_{v_i}(C) = 0$ for some i , then we define the corresponding ratio as $+\infty$, and so the inequality is trivially true.

There have been many articles published on this conjecture. For a survey of these results, as well as other variations of this problem, see [4]. In this note, we prove Conjecture 1 under an additional assumption:

Theorem 1. *Suppose that C is a convex, bounded subset of \mathbb{R}^n , and that P_1, \dots, P_k are planks which cover C . Furthermore, assume that*

$$C \setminus \bigcup_{i=1}^m P_i$$

is convex for every m with $1 \leq m \leq k$. Then Conjecture 1 is true for C and P_1, \dots, P_k .

Now, recall that the dimension of a convex set $C \subseteq \mathbb{R}^n$ is defined to be the dimension of the unique affine subspace of \mathbb{R}^n containing C in which C has non-empty interior. We also prove the following theorem, which gives a lower bound on the sum of relative widths that is not sharp, but does hold for general configurations of planks.

Theorem 2. *Suppose that C is a convex, bounded subset of \mathbb{R}^n , and that P_1, \dots, P_k are planks in directions v_1, \dots, v_k and of widths w_1, \dots, w_k which cover C . Define V to be the span of the directions v_1, \dots, v_k , and define $\pi_V(C)$ to be the orthogonal projection of C onto V . Then*

$$\sum_{i=1}^k \frac{w_i}{w_{v_i}(C)} \geq \frac{1}{\dim \pi_V(C)}.$$

1. PROOF OF THEOREM 1

The proof of Theorem 1 follows immediately from repeatedly applying the following proposition. We note that the idea of dilating a convex set about a point on the boundary was also used by Alexander in [1] to prove different results.

Proposition 3. *Suppose that C is a bounded, convex subset of \mathbb{R}^n , and that P_1, \dots, P_k are planks which cover C . Furthermore, suppose that P_m has the property that $C \setminus P_m$ is a convex set for some $1 \leq m \leq k$. If Conjecture 1 is true for $C \setminus P_m$ and $P_1, \dots, P_{m-1}, P_{m+1}, \dots, P_k$, then it is true for C and P_1, \dots, P_k .*

Proof. Define $X = C \setminus P_m$. We first show that we may make the following assumptions:

- (1) $m = 1$.
- (2) $w_{v_i}(C) > 0$ for all $1 \leq i \leq k$.
- (3) X is not empty.
- (4) $w_{v_i}(X) > 0$ for all $1 \leq i \leq k$.

We can clearly make the first assumption without loss of generality. If $w_{v_i}(C) = 0$ for some i , then $\frac{w_i}{w_{v_i}(C)} = +\infty > 1$, completing the proof. If X is empty, then $\frac{w_1}{w_{v_1}(C)} \geq 1$, finishing the proof. If $w_{v_i}(X) = 0$ for some i , then since X contains a point which is not in P_1 , since P_1 is closed, and since C is convex, $w_{v_i}(C) = 0$, contradicting the second assumption.

Suppose X and C satisfy all of the above assumptions. We will prove that, for every $v \in G(n, 1)$,

$$(*) \quad \frac{w_v(X)}{w_v(C)} \geq \frac{w_{v_1}(X)}{w_{v_1}(C)} \geq 1 - \frac{w_1}{w_{v_1}(C)}.$$

The second inequality is immediate, since $w_{v_1}(X) \geq w_{v_1}(C) - w_1$. We prove the first inequality, beginning with the statement of some elementary facts about the widths of convex sets. Here, v is any element of $G(n, 1)$, and K_1, K_2 , and K are any bounded convex sets in \mathbb{R}^n .

- (1) If K_1 and K_2 are translates of each other, then $w_v(K_1) = w_v(K_2)$.
- (2) If $K_1 = cK_2$ for some $c \in \mathbb{R}$, then $w_v(K_1) = |c|w_v(K_2)$.
- (3) If $K_1 \subset K_2$, then $w_v(K_1) \leq w_v(K_2)$.
- (4) $w_v(K) = w_v(\bar{K})$

Our proof proceeds by considering two cases based on how $w_{v_1}(X)$ compares to $w_{v_1}(C)$.

Case 1: If $w_{v_1}(X) = w_{v_1}(C)$, then $\overline{X} = \overline{C}$, and so from fact (4) we have that $w_v(X) = w_v(C)$ for all $v \in G(n, 1)$. As a result, $\frac{w_v(X)}{w_v(C)} \geq 1 = \frac{w_{v_1}(X)}{w_{v_1}(C)}$.

Case 2: If $w_{v_1}(C) > w_{v_1}(X)$, then one of the supporting hyperplanes of X in the direction v_1 lies in P_1 , and the other lies outside of it. Let H_1 be the first hyperplane, and let H_2 be the second; we have that $H_1 \cap P_1 = \emptyset$, and that $H_2 \subset P_1$. Since H_1 and H_2 are supporting hyperplanes, there exists a point $p \in H_1 \cap \partial C$. From fact (1), we may assume that p is the origin. Let P^* be the plank between H_1 and H_2 . From the definition of X , and the above assumptions about H_1 and H_2 ,

$$P^* \cap \overline{C} = \overline{X}.$$

In particular, if V is the unit vector perpendicular to H_1 which points into P^* , then $x \in \overline{X}$ if and only if $0 \leq V \cdot x \leq w_{v_1}(X)$.

Let $\rho = \frac{w_{v_1}(X)}{w_{v_1}(C)}$, and consider ρC . We claim that $\rho C \subset \overline{X}$. From the above inequality, we need only show that

$$0 \leq V \cdot x \leq w_{v_1}(X)$$

for every $x \in \rho C$. For every such x , there is a $y \in C$ such that $x = \rho y$. We then have that

$$0 \leq y \cdot V \leq w_{v_1}(C),$$

and so

$$0 = 0\rho \leq x \cdot V \leq \rho w_{v_1}(C) = w_{v_1}(X).$$

Since $\rho C \subset \overline{X}$, by facts (2), (3), and (4), $\rho w_v(C) \leq w_v(X)$ for every $v \in G(n, 1)$. We obtain our result by rearranging this inequality. This proof is shown in Figure 1.

Now, suppose that Conjecture 1 holds for X . We then have that

$$\sum_{i=2}^k \frac{w_i}{w_{v_i}(X)} \geq 1,$$


and so by multiplying both sides by $\frac{w_{v_1}(X)}{w_{v_1}(C)}$ and by using (*),

$$\sum_{i=2}^k \frac{w_i}{w_{v_i}(C)} \geq \frac{w_{v^*}(X)}{w_{v^*}(C)}.$$

Adding the contribution from P_1 to both sides and using (*) again, we have that

$$\sum_{i=1}^k \frac{w_i}{w_{v_i}(C)} \geq 1. \quad \square$$

FIGURE 1. Proof of Proposition 3.



Remark. We mention a noteworthy cover of a convex set by planks which cannot be ordered as in Theorem 1. As Hunter observed in [5], there is a configuration of three planks P_1, P_2 , and P_3 which cover the equilateral triangle T of side length 1 such that no plank's removal results in a convex set, and such that

$$\frac{w_1}{w_{v_1}(T)} + \frac{w_2}{w_{v_2}(T)} + \frac{w_3}{w_{v_3}(T)} = 1.$$

The configuration is as follows: Suppose that the barycenter of the equilateral triangle is the origin, and that one edge is parallel to the x -axis. Each of our three planks has width $1/3$, and they are centered on the lines passing through the origin with tangent vectors $(0, 1)$, $(\sqrt{3}, -1)$, and $(-\sqrt{3}, -1)$. In particular, we cannot hope to prove if a configuration of planks covering a convex body has the sum of their relative widths equal to 1, then they must satisfy the hypotheses of Theorem 1.

2. PROOF OF THEOREM 2

Proof. Suppose that C is a convex, bounded subset of \mathbb{R}^n , and that P_1, \dots, P_k are planks in directions v_1, \dots, v_k and of widths w_1, \dots, w_k which cover C . Define V to be the span of the directions v_1, \dots, v_k , and define $\pi_V : \mathbb{R}^n \rightarrow V$ to be the orthogonal projection onto V . We wish to show that the sum of the relative widths of C is bounded below by $\frac{1}{\dim \pi_V(C)}$.

Notice because C is convex in \mathbb{R}^n , we have that $\pi_V(C)$ is convex in V . Furthermore, the projections $\pi_V(P_1), \dots, \pi_V(P_k)$ are planks in V that cover $\pi_V(C)$. Given $i \in \{1, \dots, k\}$, the plank $\pi_V(P_i)$ has width and direction equal to that of P_i , namely w_i and v_i , respectively. Also, the relative widths of $\pi_V(C)$ in V are equal to those of C in \mathbb{R}^n , meaning $w_{v_i}(\pi_V(C)) = w_{v_i}(C)$ for each i . Thus, it suffices to show that the sum of relative widths of $\pi_V(C)$ is bounded below by $\frac{1}{\dim \pi_V(C)}$.

Define m to be the dimension of $\pi_V(C)$. In particular, because $\pi_V(C)$ is convex in V , there exists a unique m -dimensional affine subspace $A \subseteq$

$V \subseteq \mathbb{R}^n$ containing $\pi_V(C)$ in which $\pi_V(C)$ has non-empty interior. Let C', P'_1, \dots, P'_k denote the intersections of $\pi_V(C), \pi_V(P_1), \dots, \pi_V(P_k)$ with A . As in the proof of Proposition 3, we may assume the relative width $w_{v_i}(C') = w_{v_i}(\pi_V(C)) > 0$ for all i . It follows that each direction v_i is not orthogonal to A , and hence P'_1, \dots, P'_k are planks in A . For each i , define v'_i and w'_i to be the direction and width of P'_i in A , respectively. By a simple similar triangles argument, it is easy to see that the relative widths of C' with respect to $\pi_V(P_i)$ and P'_i are equal. In other words, for each i ,

$$\frac{w_i}{w_{v_i}(C')} = \frac{w'_i}{w_{v'_i}(C')}.$$

Thus, because $C' = \pi_V(C)$ and $\dim \pi_V(C) = m$, it suffices to prove that the sum of the relative widths of C' with respect to the planks P'_1, \dots, P'_k is bounded below by $\frac{1}{m}$.

Now, recall that the John ellipsoid for a given d -dimensional convex set K is the unique ellipsoid contained in K with maximal d -dimensional volume (see [2]). In addition, if B_1^d , the unit ball centered at the origin in \mathbb{R}^d , is the John ellipsoid for K , then K is contained in B_d^d , the ball of radius d centered at the origin in \mathbb{R}^d .

Since A and C' are m -dimensional and the sum of relative widths is independent of affine transformations, we can apply an affine transformation to \mathbb{R}^n so that the image of A under the transformation is $\mathbb{R}^m \times \{0\} \subseteq \mathbb{R}^n$ and the image of C' has the unit ball B_1^m as its inscribed John ellipsoid. Thus, the image of C' under this affine transformation is contained in the ball B_m^m of radius m . Because applying this affine transformation does not affect the sum of the relative widths, we will use " C', P'_1, \dots, P'_k " to refer to their respective images under this transformation.

Observe that, since the planks P'_1, \dots, P'_k cover C' , they also cover B_1^m . By Bang's solution to the original Tarski plank problem [3], it follows that the total width of all the planks, $w'_1 + \dots + w'_k$, is no less than 2 (twice the radius of the ball). Because the ball B_m^m contains C' , each width $w_{v'_i}(C')$ is no more than $2m$. Therefore,

$$\sum_{i=1}^k \frac{w_i}{w_{v_i}(C')} = \sum_{i=1}^k \frac{w'_i}{w_{v'_i}(C')} \geq \frac{1}{2m} \sum_{i=1}^k w'_i \geq \frac{1}{m}. \quad \square$$

Acknowledgments The authors would like to thank Marianna Cörnyei and Almut Burchard for useful conversations with regard to this problem. The first author was supported in part by NSF grant DMS-1906543.

REFERENCES

1. Ralph Alexander, *A problem about lines and ovals*, The American Mathematical Monthly **75** (1968), no. 5, 482–487.
2. Keith Ball, *An elementary introduction to modern convex geometry*, Flavors of Geometry (1997), 1–58.
3. Thøger Bang, *A solution of the “Plank Problem”*, Proceedings of the American Mathematical Society **2** (1951), no. 6, 990–993.
4. Károly Bezdek, *Geometry — intuitive, discrete, and convex: A tribute to László Fejes Tóth*, ch. Tarski’s Plank Problem Revisited, pp. 45–64, Springer Berlin Heidelberg, Berlin, Heidelberg, 2013.
5. Henry F. Hunter, *Some special cases of Bang’s inequality*, Proceedings of the American Mathematical Society **117** (1993), 819 – 821.
6. Alfred Tarski, *Further remarks about the degree of equivalence of polygons (in Polish)*, Parametr **2** (1932).

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TX
Email address: gchambers@rice.edu

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TX
Email address: mouille@rice.edu