

**From Monge to Higgs:
a survey of distance computations in
noncommutative geometry**

Pierre Martinetti

ABSTRACT. This is a review of explicit computations of Connes distance in noncommutative geometry, covering finite dimensional spectral triples, almost-commutative geometries, and spectral triples on the algebra of compact operators. Several applications to physics are covered, like the metric interpretation of the Higgs field, and the comparison of Connes distance with the minimal length that emerges in various models of quantum spacetime. Links with other areas of mathematics are studied, in particular the horizontal distance in sub-Riemannian geometry. The interpretation of Connes distance as a non-commutative version of the Monge-Kantorovich metric in optimal transport is also discussed.

CONTENTS

1. Introduction	2
2. Distances on the space of states of an algebra	4
3. Finite dimensional algebras	10
4. Product of geometries and the Higgs field	18
5. Sub-Riemannian geometry from gauge fluctuation of the metric	23
6. Compact operators: Moyal plane vs quantum space	33
7. Discussion	38
Notations	44
References	45

*Proceedings of the workshop “Noncommutative Geometry and Optimal Transport”,
Besançon november 2014.*

1. Introduction

The distance formula in noncommutative geometry has been introduced by Connes at the end of the 80's [13]. Given a so-called *spectral triple* $(\mathcal{A}, \mathcal{H}, D)$, that is an involutive algebra \mathcal{A} acting on a Hilbert space \mathcal{H} via a representation π , and an operator D on \mathcal{H} such that the commutator $[D, \pi(a)]$ is bounded for any a in \mathcal{A} , one defines on the space $\mathcal{S}(\mathcal{A})$ of states of \mathcal{A} the (possibly infinite) distance

$$(1.1) \quad d(\varphi, \varphi') = \sup_{a \in \mathcal{A}} \{ |\varphi(a) - \varphi'(a)|, \|[D, \pi(a)]\| \leq 1 \} \quad \forall \varphi, \varphi' \in \mathcal{S}(\mathcal{A}).$$

For $\mathcal{A} = C_0^\infty(\mathcal{M})$ the commutative algebra of smooth functions vanishing at infinity on a locally compact and complete manifold \mathcal{M} , acting on the Hilbert space \mathcal{H} of square integrable differential forms and D the signature operator, this distance computed between pure states gives back the geodesic distance on \mathcal{M} . In this sense, eq. (1.1) is a generalization of Riemannian geodesic distance that makes sense in a noncommutative setting, and provides an original tool to study the geometry of the space of states on an algebra. Besides its mathematical interest, Connes distance also has an intriguing echo in physics, for it yields a metric interpretation for the Higgs field [14], the missing piece of the *Standard Model of Fundamental Interactions* recently discovered by the Large Hadronic Collider at CERN.

In the 90's, Rieffel [55] noticed that (1.1) was a noncommutative version of the Wasserstein distance of order 1 in the theory of optimal transport (the modern version of Monge *déblais et remblais* problem). More exactly, this is a noncommutative generalization of Kantorovich dual formula of the Wasserstein distance [36]. Formula (1.1), which we call *spectral distance* in this survey, thus offers an unexpected connection between an ancient mathematical problem and the most recent discovery in high energy physics.

The meaning of this connection is far from clear. Yet, Rieffel's observation suggests that the spectral distance may provide an interesting starting point for a theory of optimal transport in noncommutative geometry, as well as a possible interpretation of the Higgs field as a cost function on spacetime. More specifically, one may wonder

- What remains of the duality Wasserstein (minimizing a cost)/Kantorovich (maximizing a profit) in the noncommutative setting? Is there some "noncommutative cost" that one is minimizing while computing the supremum in the distance formula?
- May the noncommutative geometry point of view on the Wasserstein distance help to solve some problems in optimal transport? Vice-versa, can one use results of optimal transport to address relevant issues in noncommutative geometry?
- Is such a generalization of the Riemannian distance truly interesting for physics?

These questions were at the origin of the mini-workshop *Optimal transport and noncommutative geometry* held in Besançon in november 2014, and whose present text is part of the proceedings. Both optimal transport and noncommutative geometry are active areas of research, but with little intersection. In addition, the metric aspect of noncommutative geometry is a part of the theory that has been relatively little studied so far [15]. Nevertheless several results - including explicit

computations - have been obtained in the recent years, and links with other areas of geometry (like sub-Riemannian geometry) have been discovered.

This survey aims at providing an account of the metric aspect of noncommutative geometry, readable by non experts. The questions listed above will serve as a guideline (they are discussed in a more systematic way in the last section of the paper), but our point of view is rather to emphasize explicit calculations of the spectral distance, starting with commutative examples and going further in non-commutativity: finite dimensional algebras, matrix valued functions on a manifold, compact operators. We omit the proof (that can be found in the indicated bibliography) and stress for each example some application in physics, or some relation with other part of mathematics.

More precisely, after some generalities in section 2 where we introduce formula (1.1) and discuss some of its properties, we begin our survey in section 3 with finite dimensional spectral triples. This is essentially a review of [34] with some slight generalizations to non-pure states. Depending on the finite dimensional algebra being commutative or not, one deals with distance on a graphs (§3.1 and §3.2) or on projective spaces (§3.3), like the sphere (§3.4). In section 4 we consider products of spectral triples. After general properties in §4.1 mainly taken from [21], we focus on *almost commutative geometries* in §4.2, that is the product of a manifold by a finite dimensional spectral triple. This is in this context that the Higgs field acquires a metric interpretation [8, 14] as the component of the metric in a discrete internal dimension [48], as explained in §4.3. Section 5 is entirely devoted to the relation between almost-commutative geometry and sub-Riemannian geometry. As recalled in §5.1, this relation has been pointed out in [14] but fully studied in [44, 46]: formula (1.1) yields a (possibly infinite) distance on the bundle $\mathcal{P}(\mathcal{A})$ of pure states of the algebra of matrix-valued functions on a manifold, which is finite between certain classes of leaves of the horizontal foliation of $\mathcal{P}(\mathcal{A})$. This is in contrast with the horizontal distance which, by definition, is infinite between horizontal leaves. The difference between the spectral and the horizontal distances is governed by the holonomy associated to a Dirac operator of an almost commutative geometry (§5.2). The computation of the distances is worked out in details for a simple example (bundle on a circle) in §5.3-5.6 In section 6 we consider a truly noncommutative example, that is a spectral triple on the algebra of compact operators. We view the latter first as the algebra of the Moyal plane in §6.1, then as the algebra describing some models of quantum spacetimes in §6.2. In the last section of the paper, we discuss various problems, in particular what could play the role of geodesics in a noncommutative framework (§7.1 and §7.2), and how to export Kantorovich duality to the noncommutative side (§7.3).

Although we try to cover a wide range of examples, this survey is not exhaustive. For a state of the art on the topological aspect of metric noncommutative geometry, we invite the reader to see the nice review of Latrémolière in this volume [37], or [49] for an approach oriented towards KK -theory. Among the subjects that are not treated here, let us mention applications to dynamical systems [3], fractals (see e.g. [12, 11]), as well as the pseudo-Riemannian case, e.g. in [51] and [29].

The notations are collected at the end of the text, before the bibliography.

2. Distances on the space of states of an algebra

A state φ of a complex C^* -algebra \mathcal{A} is a linear application from \mathcal{A} to \mathbb{C} which is positive (any positive element a^*a of \mathcal{A} has image a non-negative real number) and of norm 1. A similar notion exists for real algebras, although one should be careful that selfadjointness, $\varphi(x^*) = \bar{\varphi}(x)$, does not follow from positivity, as it does for unital complex algebras [31].

For any C^* -algebra the set of states $\mathcal{S}(\mathcal{A})$ is convex, and even compact in the weak-* topology in case the algebra is unital. The extremal points are the pure states $\mathcal{P}(\mathcal{A})$. By Gelfand theorem, for $\mathcal{A} = C_0(\mathcal{X})$ the algebra of continuous functions vanishing at infinity on a locally compact topological space \mathcal{X} , the pure states are in 1-to-1 correspondence with the points of \mathcal{X} , viewed as the evaluation

$$(2.1) \quad \delta_x(f) := f(x) \quad \forall x \in \mathcal{X}, f \in C_0(\mathcal{X}).$$

Taking as a rough definition of noncommutative geometry a “space whose algebra of functions \mathcal{A} is non-commutative”, pure states of \mathcal{A} thus appear as natural candidates to play the role of points in a noncommutative framework. One may prefer to focus on classes of irreducible representations rather than on pure states; this is discussed in §7.1.

2.1. Commutative case: the Monge-Kantorovich distance. In the commutative case $\mathcal{A} = C_0(\mathcal{X})$, a distance in the space of states $\mathcal{S}(\mathcal{A})$ is provided by optimal transport. Namely, given a function $c : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ called the *cost function*, the optimal transport between two probability measures μ_1, μ_2 on \mathcal{X} is

$$(2.2) \quad W(\mu_1, \mu_2) := \inf_{\rho} \int_{\mathcal{X} \times \mathcal{X}} c(x, y) d\rho(x, y)$$

where the infimum runs on all the measure ρ on $\mathcal{X} \times \mathcal{X}$ with marginals μ_1, μ_2 . When the cost function c is a distance, then $W(\mu_1, \mu_2)$ is a distance on the space of probability measures on \mathcal{X} , called the *Wasserstein* or the *Monge-Kantorovich* distance of order 1. One obtains a distance on the space of states noticing that any probability measure μ defines a state

$$(2.3) \quad \varphi(f) = \int_{\mathcal{X}} f(x) d\mu(x),$$

and any state comes in this way.

The Monge-Kantorovich distance is important in probability theory because the convergence in W always implies the weak* convergence (with convergence of moments). For \mathcal{X} compact, W actually metrizes the weak* topology on probability measures. This is not the only distance to make it, but according to Villani [59, p. 97] this is the most convenient one.

A similar definition exists for any order $p \in \mathbb{N}^*$, by considering instead of (2.2)

$$(2.4) \quad W_p(\mu_1, \mu_2) := \inf_{\rho} \left(\int_{\mathcal{X} \times \mathcal{X}} c^p(x, y) d\rho(x, y) \right)^{\frac{1}{p}}.$$

Nevertheless, in this paper we will mostly consider the distance of order 1, because in this particular case there exists a dual formulation which makes sense in a non-commutative context. Indeed, Kantorovich showed [36] that W can be equivalently

written as

$$(2.5) \quad W(\mu_1, \mu_2) := \sup_{\|f\|_{\text{Lip}} \leq 1} \int_{\mathcal{X}} f d\mu_1 - \int_{\mathcal{X}} f d\mu_2$$

where the supremum runs on all real functions which are Lipschitz with respect to the cost, that is

$$(2.6) \quad |f(x) - f(y)| \leq c(x, y) \quad \forall x, y \in \mathcal{X}.$$

As explained in §2.2 below, for $\mathcal{X} = \mathcal{M}$ a Riemannian complete manifold, the dual form (2.5) of the Wasserstein distance coincides with the spectral distance (1.1) for $\mathcal{A} = C_0^\infty(\mathcal{M})$ acting on the space of differential forms and D the signature operator.

Before entering the details, let us stress why the dual formulation of Kantorovich may be of interest for physics. Computed between pure states, $W(\delta_x, \delta_y) = c(x, y)$ gives back the cost function. In particular on a Riemannian manifold \mathcal{M} , taking as cost the geodesic distance, the Wasserstein distance (2.5) between pure states provides an algebraic formulation of the geodesic distance in terms of supremum, in contrast with the usual definition as the infimum on the length of all paths between x and y . This view on the geodesic distance does not rely on any notion ill defined in the quantum context, such as points or path between points. It only involves algebraic tools, typical from quantum physics.

Possible noncommutative generalizations of W_p for $p \geq 2$ are discussed in §7.3.

2.2. Noncommutative case: Connes spectral distance. A distance on the space $\mathcal{S}(\mathcal{A})$ of states of a non-necessarily commutative C^* -algebra \mathcal{A} has been introduced by Connes at the end of the 80's [13] in the framework of noncommutative geometry.

Assuming \mathcal{A} acts on an Hilbert space \mathcal{H} , then given an operator D on \mathcal{H} , one associates to any pair of states $\varphi, \varphi' \in \mathcal{S}(\mathcal{A})$ the quantity

$$(2.7) \quad d(\varphi, \varphi') := \sup_{a \in \text{Lip}_D(\mathcal{A})} |\varphi(a) - \varphi'(a)|$$

where the D -Lipschitz ball of \mathcal{A} is the subset of \mathcal{A} defined as

$$(2.8) \quad L_D(\mathcal{A}) := \{a \in \mathcal{A}, \|[D, a]\| \leq 1\},$$

where the norm is the operator norm on \mathcal{H} . In (2.8) as well as most of the time in the rest of the paper, we omit the symbol of representation and we identify an element a of \mathcal{A} with its representation $\pi(a)$ as bounded operator on \mathcal{H} . In these conditions (2.7) coincides with (1.1).

Eq. (2.7) is obviously invariant under the exchange of φ and φ' , and is zero if $\varphi = \varphi'$. The triangle inequality is easy to check. For two states φ, φ' that are equal everywhere but on some element a_∞ such that $[D, a_\infty]$ is unbounded, one has $d(\varphi, \varphi') = 0$ although $\varphi \neq \varphi'$. To avoid this, one requires $[D, a]$ be bounded for any $a \in \mathcal{A}$. Then (2.7) defines a distance (possibly infinite) on $\mathcal{S}(\mathcal{A})$. Asking $(D - \lambda\mathbb{I})^{-1}$ to be compact for any λ in the resolvent set of D (in case \mathcal{A} is unital, this means that D has compact resolvent), the set $(\mathcal{A}, \mathcal{H}, D)$ is called a *spectral triple* (and D a *Dirac operator*). Hence the name *spectral distance* to denote (2.7).

For $\mathcal{A} = C_0^\infty(\mathcal{M})$ the algebra of smooth functions vanishing at infinity on a locally compact complete Riemannian manifold \mathcal{M} , with multiplicative representation on the Hilbert space $\mathcal{H} = L^2(\mathcal{M}, \wedge)$ of square integrable differential forms,

$$(2.9) \quad (f\psi)(x) = f(x)\psi(x) \quad \forall x \in \mathcal{M}, \psi \in L^2(\mathcal{M}, \wedge),$$

and $D = d + d^\dagger$ the signature operator (d is the exterior derivative, d^\dagger its adjoint) then the spectral distance (2.7) computed between pure states gives back the geodesic distance on \mathcal{M} ,

$$(2.10) \quad d(\delta_x, \delta_y) = d_{\text{geo}}(x, y).$$

A similar result is obtained, in case \mathcal{M} is a spin manifold, with $\mathcal{H} = L^2(\mathcal{M}, S)$ the Hilbert space of square integrable spinors and

$$(2.11) \quad D = \not{D} := -i\gamma^\mu(\partial_\mu + \omega_\mu)$$

the usual Dirac operator, with ω_μ the spin connection and γ^μ the Dirac matrices satisfying

$$(2.12) \quad \gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = 2g^{\mu\nu}\mathbb{I}$$

where $g^{\mu\nu}$ the Riemannian metric on \mathcal{M} .

Furthermore, in [56] Rieffel noticed that for \mathcal{M} compact and for any state (pure or not), formula (2.7) was nothing but Kantorovich dual formulation (2.5) of the Wasserstein distance. This is because the norm of the commutator $[d + d^\dagger, f]$ (or $[\not{D}, f]$ in case the spin structure is taken into account) is nothing but the Lipschitz norm of f . We show in [22] that this remains true for a locally compact manifold as soon as it is complete (the latter condition guarantees that looking for the supremum on $C_0^\infty(\mathcal{M})$ or on Lipschitz functions is equivalent).

Therefore, Connes spectral distance appears as a generalization of the Wasserstein distance. More precisely, it provides a formulation of Kantorovich dual formula which makes sense also in a noncommutative context. Whether there exists a noncommutative version of the initial definition (2.2) of the Wasserstein distance as an infimum is an open question, discussed in section 7.3.

Let us stress that Connes formula makes sense in a wider context: one may look for the supremum on the Lipschitz ball $L(a) \leq 1$ for any seminorm L on \mathcal{A} , non necessarily coming from the commutator with an operator. One does not even need to work with an algebra: states and Lipschitz seminorms makes sense for *ordered unit spaces* (see [56, §11] for an extended discussion on that matter). This flexibility is useful when one focuses on topological aspects of the distance (for instance under which conditions does (2.7) metrize the weak* topology on $\mathcal{S}(\mathcal{A})$? [37]). In this review, we adopt the point of view that spectral triples provide algebras and operators D - hence seminorms L_D - that are relevant for physics as well as for other aspect of mathematics, offering thus various examples where the explicit computation of the spectral distance is worth undertaken.

It is also worth mentioning that by adding more conditions on \mathcal{A} , \mathcal{H} and D , one is able to fully characterize a Riemannian closed (spin) manifold \mathcal{M} as a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ where \mathcal{A} is commutative [16]. Focusing only on the metric aspect, one may as well be interested in characterizing a metric space in terms of algebraic datas, without the need of any smooth structure. A good reference on this topic is [61]. A general reference on the algebraic way of characterizing a smooth manifold is [52].

2.3. Isometries & projections. Before making explicit computations of the distance, let us list various definitions and easy but useful general results. In all this section, d denotes the spectral distance (2.7) associated to an arbitrary spectral triple $(\mathcal{A}, \mathcal{H}, D)$.

DEFINITION 2.1. *We call “optimal element for a pair of states $\varphi, \varphi' \in \mathcal{S}(\mathcal{A})$ ” any element a in $L_D(\mathcal{A})$ such that*

$$(2.13) \quad |\varphi(a) - \varphi'(a)| = d(\varphi, \varphi')$$

or, in case the supremum is not reached, any sequence $\{a_n \in L_D(\mathcal{A})\}$ such that

$$(2.14) \quad \lim_{n \rightarrow \infty} |\varphi(a_n) - \varphi'(a_n)| = d(\varphi, \varphi').$$

LEMMA 2.2. [34, Lem. 1] *The supremum in (2.7) can be searched equivalently on selfadjoint elements of \mathcal{A} . In case \mathcal{A} is unital, the supremum can be equivalently searched on positive elements.*

We call isometry of the state space an application $\alpha : \mathcal{S}(\mathcal{A}) \rightarrow \mathcal{S}(\mathcal{A})$ such that

$$(2.15) \quad d(\varphi, \varphi') = d(\alpha(\varphi), \alpha(\varphi')) \quad \forall \varphi, \varphi' \in \mathcal{S}(\mathcal{A}).$$

A class of isometries particularly useful for explicit computations are the lift to states of inner automorphisms of \mathcal{A} , that is

$$(2.16) \quad \alpha_u(\varphi) := \varphi \circ \alpha_u$$

where $\alpha_u := \text{Ad } u$ for some unitary $u \in \mathcal{A}$.

LEMMA 2.3. *Let u be a unitary element in \mathcal{A} that commutes with D , then α_u is an isometry of $\mathcal{S}(\mathcal{A})$. Namely*

$$(2.17) \quad d(\varphi, \varphi') = d(\varphi \circ \alpha_u, \varphi' \circ \alpha_u) \quad \forall \varphi, \varphi' \in \mathcal{S}(\mathcal{A}).$$

The proof is easy and can be found e.g. in [40, Prop. 1.29]. The result is also valid for some operator u in $\mathcal{B}(\mathcal{H})$ that is not necessarily the representation of a unitary element of \mathcal{A} . In this case one should consider only the states whose domain contains u , that is such that $\varphi \circ \alpha_u$ and $\varphi' \circ \alpha_u$ make sense.

Other useful applications are projections, that sometimes permit to reduce the search for the supremum in (2.7) to subsets of \mathcal{A} more tractable than \mathcal{A} itself.

DEFINITION 2.4. *The projection of a spectral triple $(\mathcal{A}, \mathcal{H}, D, \pi)$ (π is the representation of \mathcal{A} on \mathcal{H}) by a projection $e = e^* = e^2 \in \mathcal{B}(\mathcal{H})$ is the triple*

$$(2.18) \quad \mathcal{A}_e := \alpha_e(\mathcal{A}), \quad \mathcal{H}_e := e\mathcal{H}, \quad D_e := eDe|_{\mathcal{H}_e},$$

where $\alpha_e(a) := e\pi(a)e$ for any $a \in \mathcal{A}$.

The projected triple $(\mathcal{A}_e, \mathcal{H}_e, D_e)$ may not be a spectral triple since \mathcal{A}_e may not be an algebra, for instance when $e \notin \pi(\mathcal{A})$. Nevertheless, when \mathcal{A} is unital the set

$$(2.19) \quad \mathcal{A}_e^{\text{sa}} := \{e\pi(a)e, a = a^* \in \mathcal{A}\}$$

of selfadjoint elements of \mathcal{A}_e is an ordered unit space. Therefore, as explained at the end of §2.2, the notion of states of $\mathcal{A}_e^{\text{sa}}$ - and by extension of \mathcal{A}_e - still makes sense, with an obvious map from $\mathcal{S}(\mathcal{A}_e)$ to $\mathcal{S}(\mathcal{A})$,

$$(2.20) \quad \varphi \rightarrow \varphi \circ \alpha_e.$$

Given $\varphi, \varphi' \in \mathcal{S}(\mathcal{A}_e)$, we still call *spectral distance* the quantity

$$(2.21) \quad d_e(\varphi, \varphi') := \sup_{\text{Lip}_D(\mathcal{A}_e)} |\varphi(a) - \varphi'(a)|.$$

LEMMA 2.5. [48, Lem. 1] *Let $(\mathcal{A}, \mathcal{H}, D)$ be a unital spectral triple and e a projection in \mathcal{H} that commutes with D , then for any states φ, φ' of \mathcal{A}_e one has*

$$(2.22) \quad d_e(\varphi, \varphi') = d(\varphi \circ \alpha_e, \varphi' \circ \alpha_e).$$

Said differently, a projection that commutes with the Dirac operator behaves like an isometry. The difference between (2.22) and (2.15) is that in (2.22) the set of elements on which the supremum is searched is smaller on the l.h.s. than on the r.h.s. Notice also that the application (2.20) has no reason to be surjective.

2.4. Connected components. Given a spectral triple $(\mathcal{A}, \mathcal{H}, D)$, we denote the set of states at finite spectral distance from a state $\varphi \in \mathcal{S}(\mathcal{A})$ by

$$(2.23) \quad \text{Con}(\varphi) := \{\varphi' \in \mathcal{S}(\mathcal{A}); d(\varphi, \varphi') < +\infty\}.$$

The notation is justified because this set coincides with the connected component of φ in $\mathcal{S}(\mathcal{A})$ for the topology metrized by the spectral distance (see [22, Def. 2.1]).

PROPOSITION 2.6. *For any $\varphi \in \mathcal{S}(\mathcal{A})$, the set $\text{Con}(\varphi)$ is convex.*

PROOF. For any $\varphi_0, \varphi_1 \in \text{Con}(\varphi)$ and $s \in [0, 1]$, denote

$$(2.24) \quad \varphi_s := s\varphi_0 + (1-s)\varphi_1.$$

One easily checks that

$$(2.25) \quad d(\varphi_s, \varphi_t) = |s-t|d(\varphi_0, \varphi_1) \quad \forall s, t \in \mathbb{R}.$$

By the triangle inequality, $d(\varphi_0, \varphi_1)$ is finite. Thus the same is true for $d(\varphi_s, \varphi_t)$. In particular $d(\varphi_0, \varphi_s)$ is finite, so again by the triangle inequality $d(\varphi, \varphi_s)$ is finite for any $s \in [0, 1]$. Hence $\varphi_s \in \text{Con}(\varphi)$, showing the later is convex. \square

Restricting the connected component of a state to pure states, by prop. 2.6 one obtains a set whose convex hull is still in the connected component,

$$(2.26) \quad \overline{\text{Con}(\varphi) \cap \mathcal{P}(\mathcal{A})} \subset \text{Con}(\varphi).$$

But at this point nothing guarantees that $\text{Con}(\varphi)$ is the convex hull of its restriction to pure states. We come back to this point in section 7.3.

The following lemma is useful to characterize the connected components.

LEMMA 2.7. *For any two states φ, φ' of \mathcal{A} , the distance $d(\varphi, \varphi')$ is infinite if and only if there exists a sequence $a_n \in \mathcal{A}$ such that*

$$(2.27) \quad \lim_{n \rightarrow \infty} L_D(a_n) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \varphi(a_n) - \varphi'(a_n) = \infty.$$

In particular $d(\varphi, \varphi')$ is infinite as soon as there exists an element $a \in \mathcal{A}$ such that

$$(2.28) \quad L_D(a) = 0 \quad \text{and} \quad \varphi(a) \neq \varphi'(a).$$

PROOF. The proof that the non-finiteness of the distance is equivalent to (2.27) is easy and can be found for instance in [44, Lemma 1]. The second statement follows by considering $a_n := na$, $n \in \mathbb{N}$. \square

In the finite dimensional case, there are stronger results.

LEMMA 2.8. *For a spectral triple with finite dimensional \mathcal{A} and \mathcal{H} , the distance between two states φ, φ' is finite if and only if*

$$(2.29) \quad \varphi(a) = \varphi'(a) \quad \forall a \in \text{Ker } L_D.$$

In particular, for \mathcal{A} unital the distance is finite on the whole space of states if and only if

$$(2.30) \quad \text{Ker } L_D = \{\lambda \mathbb{1}, \lambda \in \mathbb{C}\}.$$

PROOF. For the first statement, by lemma 2.7 one just needs to show that

$$(2.31) \quad d(\varphi, \varphi') = \infty \implies \exists a \in \text{Ker } L_D \text{ such that } \varphi(a) \neq \varphi'(a).$$

Let us thus assume $d(\varphi, \varphi')$ is infinite. This means there exists a sequence $a_n \in \mathcal{A}$ satisfying (2.27). By hypothesis \mathcal{H} is isomorphic to \mathbb{C}^N for some $N \in \mathbb{N}$ and \mathcal{A} is a subalgebra of $M_N(\mathbb{C})$. The kernel of L_D is a vector subspace of $M_N(\mathbb{C})$. Let K^\perp denote its orthogonal complement in $M_N(\mathbb{C})$ and

$$(2.32) \quad \tilde{\mathcal{A}} := \mathcal{A} \cap K^\perp \simeq \mathcal{A} / \text{Ker } L_D.$$

Any a_n decomposes in a unique way as

$$(2.33) \quad a_n = \tilde{a}_n + k_n$$

where $k_n \in \text{Ker } L_D$ and $\tilde{a}_n \in \tilde{\mathcal{A}}$. On $\tilde{\mathcal{A}}$, the seminorm L_D is actually a norm. Moreover, since $L_D(a_n) = L_D(\tilde{a}_n)$ for any n , by (2.27) one gets

$$(2.34) \quad \lim_{n \rightarrow \infty} L_D(\tilde{a}_n) = \lim_{n \rightarrow \infty} L_D(a_n) = 0,$$

and all the norms on a finite dimensional vector are equivalent, so that \tilde{a}_n tends to zero in the C^* -norm of $M_N(\mathbb{C})$. Since states are continuous, this means

$$(2.35) \quad \lim_{n \rightarrow \infty} \varphi(\tilde{a}_n) - \varphi'(\tilde{a}_n) = 0,$$

hence

$$(2.36) \quad \lim_{n \rightarrow \infty} \varphi(a_n) - \varphi'(a_n) = \lim_{n \rightarrow \infty} \varphi(k_n) - \varphi'(k_n),$$

which is infinite by (2.27). This cannot be true if (2.29) holds true, since the r.h.s. of (2.36) would be zero. Therefore, for a finite dimensional spectral triple the non-finiteness of $d(\varphi, \varphi')$ implies that φ and φ' do not coincide on $\text{Ker } L_D$.

The second statement follows by noticing that for any element $a \neq \mathbb{1}$, there exist at least two states φ, φ' that do not take the same value on a . Indeed, given any non-zero $a \in \mathcal{A}$, there exists at least one state φ such that $\varphi(a) \neq 0$ [35, Theo. 4.3.4]. Assume that $\varphi(a) \neq 1$. Then the state

$$(2.37) \quad \varphi' := \frac{1}{2}\varphi + \frac{1}{2}\varphi_0$$

where φ_0 is the state that takes value 1 on each $a \in \mathcal{A}$, is such that $\varphi'(a) \neq \varphi(a)$. If $\varphi(a) = 1$, then again by [35, theo. 4.3.4] there exists at least a state φ' such that $\varphi'(a - \mathbb{1}) \neq 0$, that is $\varphi'(a) \neq 1$. \square

In a wider context (i.e. not necessarily finite dimensional and with a seminorm not necessarily coming from the commutator with a Dirac-like operator), condition (2.30) is one of the requirements of what Rieffel called a *Lip-norm* [55, 54], that is a seminorm L_D such that (2.7) metrizes the weak* topology. For a state of the art of the topological aspect of the spectral distance, we invite the reader to see the extensive contribution of Latrémolière in the present volume [37].

3. Finite dimensional algebras

To begin our survey of explicit computations of the spectral distance (2.7), let us consider finite dimensional (complex) C^* -algebras, that is finite sums of matrix algebras,

$$(3.1) \quad \mathcal{A} = \bigoplus_{i=1}^N M_{n_i}(\mathbb{C})$$

where $n_i \in \mathbb{N}$ for any $1 \leq i \leq N$. We begin by commutative examples $\mathcal{A} = \mathbb{C}^N$ in §3.1 and §3.2, then we study matrix algebras in §3.3 and §3.4.

3.1. Discrete spaces. The simplest case, that is $\mathcal{A} = \mathbb{C}^2$, is instructive although it is commutative and elementary. Making \mathcal{A} act on $\mathcal{H} = \mathbb{C}^2$ as diagonal matrices,

$$(3.2) \quad \pi(z_1, z_2) := \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix},$$

with

$$(3.3) \quad D = \begin{pmatrix} 0 & m \\ \bar{m} & 0 \end{pmatrix} \quad m \in \mathbb{C}$$

as Dirac operator (the diagonal of D commutes with the representation π and so is not relevant for the distance computation), one easily computes that the spectral distance between the two pure states

$$(3.4) \quad \delta_i(z_1, z_2) := z_i, \quad i = 1, 2$$

of \mathbb{C}^2 is

$$(3.5) \quad d(\delta_1, \delta_2) = \frac{1}{|m|}.$$

The spectral distance thus allows to equip the discrete two-point space $\{\delta_1, \delta_2\}$ with a generalization of the geodesic distance, although the usual notion of length-of-the-shortest-path no longer makes sense since there is no “points”, i.e. no pure states, between δ_1 and δ_2 . Incidentally, this raises the question of what should play the role of geodesics in noncommutative geometry: a curve in $\mathcal{S}(\mathcal{A})$, in $\mathcal{P}(\mathcal{A})$, or something else? We come back to this question in §7.1 and §7.2.

The construction above generalizes to arbitrary dimension: consider $\mathcal{A} = \mathbb{C}^N$ acting diagonally on \mathbb{C}^N , with D a $N \times N$ selfadjoint matrix with null-diagonal. For simplicity, we restrict to Dirac operators with real entries, that is

$$(3.6) \quad D_{ij} = D_{ji} \in \mathbb{R}$$

for any $i, j \in [1, N]$. One has N -pure states $\delta_i, i = 1, 2, \dots, N$ and we write the distance

$$(3.7) \quad d(i, j) := d(\delta_i, \delta_j).$$

PROPOSITION 3.1. [34, Prop. 7] *For $N = 3$, one deals with a three point space with distance*

$$(3.8) \quad d(1, 2) = \sqrt{\frac{D_{13}^2 + D_{23}^2}{D_{12}^2 D_{13}^2 + D_{12}^2 D_{23}^2 + D_{23}^2 D_{13}^2}}.$$

The other distances are obtained by cyclic permutations of the indices, and verify the triangle inequality “to the square”

$$(3.9) \quad d(1, 2)^2 + d(2, 3)^2 \geq d(1, 3)^2.$$

Formula (3.8) is invertible. That is, given three positive numbers (a, b, c) verifying (3.9), there exists a Dirac operator giving these numbers as distances.

PROPOSITION 3.2. [34, Prop. 8] *Let a, b, c three positive real numbers such that*

$$(3.10) \quad a^2 + b^2 \geq c^2, \quad b^2 + c^2 \geq a^2, \quad a^2 + c^2 \geq b^2.$$

There exists an operator D such that

$$(3.11) \quad d(1, 2) = a, \quad d(1, 3) = b, \quad d(2, 3) = c.$$

It has coefficients

$$(3.12) \quad D_{12} = \sqrt{\frac{2(b^2 + c^2 - a^2)}{(a + b + c)(-a + b + c)(a - b + c)(a + b - c)}},$$

D_{13} and D_{23} are obtained by cyclic permutations of a, b, c .

A surprising interpretation of (3.8) and (3.12) comes from electric circuits [34]. Starting with three numbers a, b, c satisfying (3.10), one defines

$$(3.13) \quad r_1 := a^2 + b^2 - c^2, \quad r_2 := a^2 + c^2 - b^2, \quad r_3 := b^2 + c^2 - a^2.$$

By (3.11), $d(1, 2)^2 = r_1 + r_2$ is the resistance between the points 1, 2 of the “star” circuit made of the three resistances r_1, r_2, r_3 (fig. 3.1), and similarly for $d(1, 3)$ and $d(2, 3)$. It is well known in electricity that the star circuit with resistance r_i is equivalent to a triangle circuit with resistance

$$(3.14) \quad R_{ij} := D_{ij}^{-2}$$

where the D_{ij} ’s are precisely given by formula (3.12). So modulo the reparametrizations (3.13) and (3.14), the passage from the distances to the coefficients of the Dirac operator is similar to the passage from the star to the triangle circuits.




FIGURE 1. Equivalent triangle and star circuits.

Unfortunately, the electric analogy no longer makes sense in higher dimension. Indeed, in order to work out $\text{Lip}_D(\mathcal{A})$, one needs to solve the characteristic polynomial of the antisymmetric matrix $[D, a]$ (assuming a is real). This is of order $\frac{N}{2}$, and is in principle not explicitly calculable as soon as $\frac{N}{2} \geq 5$, meaning there is little hope to explicitly compute the distance in a space with more than $N = 10$

points. In fact the difficulty arises much earlier, at $N = 4$. Consider $\mathcal{A} = \mathbb{C}^4$ acting diagonally on $\mathcal{H} = \mathbb{C}^4$ with D a 4×4 real symmetric matrix with entries D_{ij} . Write

$$d_1 := \frac{1}{D_{12}}, \quad d_2 := \frac{1}{D_{13}}, \quad d_3 := \frac{1}{D_{14}}, \quad d_4 := \frac{1}{D_{23}}, \quad d_5 := \frac{1}{D_{24}}, \quad d_6 := \frac{1}{D_{34}}.$$

PROPOSITION 3.3. [34, Theo. 9]

- i. On a four point space, $d(i, j)$ is the root of a polynomial of degree $\delta \leq 12$, and is not in general solvable by radicals.
- ii. However there are cases where $d(1, 2)$ is computable explicitly. For instance when $\frac{1}{d_2} = \frac{1}{d_5} = \infty$, one has

$$d(1, 2) = \begin{cases} d_1 & \text{when } d_1^2 \leq d_6^2, \\ \frac{d_1 \sqrt{(d_3^2 + d_1 d_6)^2}}{\sqrt{d_1^2 d_3^2} \sqrt{d_3^2 + d_6^2}} & \text{when } d_1 d_6 = d_3 d_4, \\ \sqrt{\frac{d_1^2 (d_3^2 + d_6^2) (d_4^2 + d_6^2)}{(d_3 d_4 - d_1 d_6)^2}} & \text{when } C \leq 0, \\ \max \left(\sqrt{\frac{d_1^2 (d_3^2 + d_4^2)}{(d_3 + d_4)^2 + (d_1 - d_6)^2}}, \sqrt{\frac{d_1^2 (d_3^2 - d_4^2)}{(d_3 - d_4)^2 + (d_1 + d_6)^2}} \right) & \text{otherwise,} \end{cases}$$

where

$$C := ((d_3 + d_4)^2 d_6 + (d_1 - d_6)(d_3 d_4 - d_6^2))((d_3 - d_4)^2 d_6 + (d_1 + d_6)(d_3 d_4 + d_6^2)).$$

As well,

$$d(1, 3) = \begin{cases} \sqrt{d_3^2 + d_6^2} & \text{when } (d_3^2 + d_6^2) \leq (d_1 d_6 - d_3 d_4)^2, \\ \sqrt{d_1^2 + d_4^2} & \text{when } (d_1^2 + d_4^2) \leq (d_1 d_6 - d_3 d_4)^2, \\ = \max \left(\frac{\sqrt{(d_1 d_3 + d_4 d_6)^2}}{\sqrt{(d_3 + d_4)^2 + (d_1 - d_6)^2}}, \frac{\sqrt{(d_1 d_3 + d_4 d_6)^2}}{\sqrt{(d_3 - d_4)^2 + (d_1 + d_6)^2}} \right) & \text{otherwise.} \end{cases}$$

The other distances are obtained by cyclic permutations.

This proposition shows that already for $N = 4$ the distance formula cannot be inverted. This means that given a set of $\frac{1}{2}N(N-1)$ real positive numbers satisfying the triangle inequality, there is no algorithm permitting to build a $N \times N$ Dirac operator D_N giving back these numbers as the spectral distance associated to the spectral triple $(\mathbb{C}^N, \mathbb{C}^N, D_N)$. However such an algorithm exists if one allows the size of the Hilbert space to increase.

PROPOSITION 3.4. [34, Prop. 13] Let $d_{ij}, 1 \leq i, j \leq N, i \neq j$, be a finite sequence of possibly infinite strictly positive numbers such that

$$(3.15) \quad d_{ij} = d_{ji} \quad \text{and} \quad d_{ij} \leq d_{ik} + d_{kj} \quad \text{for any } i, j, k.$$

Then there exists a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ with $\mathcal{A} = \mathbb{C}^N$ and $\mathcal{H} = \mathbb{C}^{\frac{3}{2}N(N-1)}$ such that the resulting distance on the set of pure states of \mathcal{A} is given by the numbers d_{ij} .

A similar construction has been proposed in [58, §2.2].

3.2. Distances on graphs. The four-point space in Prop. 3.3 suggests that for $N \geq 4$, there is little sense in trying to compute explicitly the distance in a N -point space with the most general Dirac operator. However some general properties of the distance can be worked out for arbitrary N . To this aim we let $\mathcal{A} = \mathbb{C}^N$ act as diagonal matrices on $\mathcal{H} = \mathbb{C}^N$, and we identify the N pure states of \mathbb{C}^N ,

$$(3.16) \quad \delta_i(z_1, z_2, \dots, z_N) := z_i \quad \forall (z_1, z_2, \dots, z_N) \in \mathbb{C}^N,$$

with the points $1, 2, \dots, N$ of a N -point graph. We take as a Dirac operator the incidence matrix of the graph, that is

$$(3.17) \quad D = \begin{pmatrix} 0 & D_{12} & \dots & \dots & D_{1N} \\ D_{12} & & D_{23} & & 0 \\ \vdots & D_{23} & 0 & \ddots & \vdots \\ \vdots & & \ddots & \ddots & D_{N-1,N} \\ D_{1N} & \dots & \dots & D_{N-1,N} & 0 \end{pmatrix} \quad D_{ij} \in \mathbb{R},$$

where $D_{ij} = D_{ji}$ is non-zero if and only if there is a link in the graph between the points i and j . A path γ_{ij} is a sequence of p distinct points $(i, i_2, \dots, i_{p-1}, j)$ such that

$$D_{i_k i_{k+1}} \neq 0 \quad \text{for all } k \in \{1, p-1\}.$$

Two points i, j are said connected if there exists at least one path γ_{ij} . We define the length of a path γ_{ij} as

$$L(\gamma_{ij}) := \sum_{k=1}^{p-1} \frac{1}{|D_{i_k i_{k+1}}|},$$

and the geodesic distance L_{ij} between any two connected points i, j as the length of the shortest path γ_{ij} .

PROPOSITION 3.5. [34, Prop. 4]

- i. Let D' be the operator obtained by canceling one or more lines and the corresponding columns in D , and d' the associated distance. Then $d' \geq d$.
- ii. The distance between two points i and j depends only on the matrix elements corresponding to points located on paths γ_{ij} .
- iii. The distance between any two points is finite if and only if they are connected.
- iv. For any two points i, j , one has $d(i, j) \leq L_{ij}$.

A case explicitly computable is the maximally connected graph, that is the operator D with all coefficients equal to a fixed real constant k .

PROPOSITION 3.6. [34, Prop. 5]

- i. The distance between any two points i, j is

$$(3.18) \quad d(i, j) = \frac{1}{|k|} \sqrt{\frac{2}{N}}.$$

- ii. If the link between two points i_1, i_2 - and only this link - is cut, $D_{i_1 i_2} = 0$, then

$$(3.19) \quad d(i_1, i_2) = \frac{1}{|k|} \sqrt{\frac{2}{N-2}}.$$

Examples of explicit computation of the spectral distance in lattices can be found in [4], [18], [20]. Applications to quantum gravity have been explored in [57].

3.3. Projective spaces. The space of pure states of $M_n(\mathbb{C})$, $n \in \mathbb{N}$, is the projective space $\mathbb{C}P^{n-1}$: any normalized vector $\xi \in \mathbb{C}^n$ defines the pure state

$$(3.20) \quad \omega_\xi(a) = \langle \xi, a\xi \rangle \quad \forall a \in M_n(\mathbb{C})$$

where $\langle \cdot, \cdot \rangle$ is the usual inner product on \mathbb{C}^n . Two such vectors equal up to a phase define the same state, and any pure state comes in this way. All the representations of $M_n(\mathbb{C})$ induced by these pure states via the Gelfand-Neimark-Segal construction are equivalent, that is why it is sometimes argued [39] that $M_n(\mathbb{C})$ should be considered as a 1-point space. On the contrary, we argue that the spectral distance provides the space of pure states of $M_n(\mathbb{C})$ with structure finer than the one of irreducible representations, and there is no reason to neglect it. We come back to this point in section 7.1.

We consider the spectral triple

$$(3.21) \quad \mathcal{A} = M_n(\mathbb{C}), \quad \mathcal{H} = \mathbb{C}^n, \quad D$$

where the action of \mathcal{A} on \mathcal{H} is the usual representation of matrices, while D is an arbitrary selfadjoint element of $M_n(\mathbb{C})$. There exists no explicit computation of the distance in the most general case, that is between any two states of $M_n(\mathbb{C})$ for arbitrary n . There are such computations for $n = 2$, which are the object of §3.4. For $n \geq 2$, we expose below some properties of the connected components, which are slight generalizations of unpublished results of [40]. There is also an explicit computation of the distance between any pure states of $M_n(\mathbb{C}) \oplus \mathbb{C}$ but with a particular class of operator D , presented in Prop. 3.11.

From now on we assume $n \geq 2$. To make the correspondence between $\mathbb{C}P^{n-1}$ and normalized vectors in \mathbb{C}^n explicit, it is convenient to fix as a basis of \mathcal{H} an orthonormal set of eigenvectors ψ_i of D , $i = 1, n$, so that

$$(3.22) \quad D = \text{diag} (d_1, d_2, \dots, d_n)$$

where $d_i \in \mathbb{R}$ are the eigenvalues of D (possibly null). For any eigenvector ψ_i , we call *eigenstate of D* the pure state

$$(3.23) \quad \omega_i := \langle \psi_i, \cdot \psi_i \rangle.$$

We write e_{ii} the diagonal matrix with only non-zero component the i^{th} entry that is equal to 1. Given a normalized complex n -vector ξ , we write $\xi_i := \langle \xi, \psi_i \rangle$ its components on the eigenbasis of D , and

$$(3.24) \quad \omega_\xi := \langle \xi, \cdot \xi \rangle$$

the corresponding pure state of $M_n(\mathbb{C})$. It is not difficult to characterize the pure states at finite distance from one another.

PROPOSITION 3.7. *Let ξ, ζ be normalized vectors in \mathbb{C}^n . The distance between the pure states ω_ξ and ω_ζ is finite if and only if the projections of ξ and ζ on the kernel - as well as on any eigenspace of D - are equal up to a phase. That is, for any eigenspace \mathcal{H}_J of D ($J \geq 1$ an integer), there exists a phase $\theta_J \in [0, 2\pi[$ such that*

$$(3.25) \quad \xi_i = e^{i\theta_J} \zeta_i \quad \text{for any } i \in I_J,$$

where I_J is the subset of $\{1, n\}$ such that $\mathcal{H}_J = \text{span} \{e_{ii}, i \in I_J\}$.

PROOF. By lemma 2.7, we just need to show that ω_ξ, ω_ζ coincide on the kernel of L_D if and only if (3.25) holds true. Let us first assume there is no degeneracy, that is all the eigenvalues d_i of D are distinct. Then (3.25) amounts to

$$(3.26) \quad |\xi_i| = |\zeta_i| \quad \forall i = 1, \dots, n.$$

The kernel of L_D is the set of diagonal matrices. Any two states ω_ξ, ω_ζ coincide on the kernel if and only if $\omega_\xi(e_{jj}) = |\xi_j|^2$ equals $\omega_\zeta(e_{jj}) = |\zeta_j|^2$ for any j , that is equation (3.26).

In case of degeneracy, one has

$$(3.27) \quad \text{Ker } L_D = \bigoplus_{J \in \mathbb{N}} \mathcal{B}(\mathcal{H}_J)$$

and φ, φ' coincide on each summand if and only if (3.25) holds. \square

As a corollary, one obtains that the connected component in the space of pure states of any pure state is a torus inside $\mathbb{C}P^{n-1}$.

COROLLARY 3.8. *Let ω_ξ be the pure state of $M_n(\mathbb{C})$ associated to the normalized vector $\xi \in \mathbb{C}^n$ with components ξ_i in the eigenbasis of D . Then $\text{Con}(\omega_\xi) \cap \mathcal{P}(M_n(\mathbb{C}))$ is the $k-1$ -torus*

$$(3.28) \quad U_\xi = \left\{ \left(\begin{array}{cc} \xi_i & \forall i \in I_1 \\ \xi_i e^{i\theta_2} & \forall i \in I_2 \\ \vdots & \\ \xi_i e^{i\theta_k} & \forall i \in I_k \end{array} \right), \theta_2, \dots, \theta_k \in [0, 2\pi[\right\},$$

where k is the number of distinct eigenvalues of D .

PROOF. This follows directly from Prop. 3.7. Notice that if $k = 1$, that is D is proportional to the identity, then (3.25) indicates that ω_ζ is at finite distance from ω_ξ if and only if $\xi = e^{i\theta_1} \zeta$. But this means $\omega_\xi = \omega_\zeta$, so that $\text{Con}(\omega_\xi)$ reduces to ω_ξ itself. That is why in (3.28) the phase $e^{i\theta_1}$ is factorized out. \square

3.4. The three dimensional ball. For $n = 2$, the distance can be explicitly computed. The space of pure states of $M_2(\mathbb{C})$ is the complex projective plane $\mathbb{C}P^1$, which is in 1-to-1 correspondence with the 2-sphere: to any normalized complex vector $\xi \in \mathbb{C}^2$ with components ξ_1, ξ_2 , one associates the point p_ξ of S^2 with Euclidean coordinates

$$(3.29) \quad x_\xi := 2\text{Re}(\xi_1 \bar{\xi}_2), \quad y_\xi := 2\text{Im}(\xi_1 \bar{\xi}_2) \quad \text{and} \quad z_\xi := |\xi_1|^2 - |\xi_2|^2.$$

The evaluation of ω_ξ on $a \in M_2(\mathbb{C})$ with components a_{ij} reads (see e.g. [6, §4.3])

$$\omega_\xi(a) = \frac{1+z_\xi}{2} a_{11} + \frac{1-z_\xi}{2} a_{22} + r \Re(e^{i\Xi} a_{12})$$

where

$$(3.30) \quad r e^{i\Xi} := x_\xi + i y_\xi = 2\xi_1 \bar{\xi}_2.$$

A non-pure state φ of $M_2(\mathbb{C})$ is given by a probability distribution ϕ on S^2 ,

$$(3.31) \quad \varphi(a) = \int_{S^2} \phi(\xi) \omega_\xi(a) d\xi = \frac{1+\tilde{z}_\phi}{2} a_{11} + \frac{1-\tilde{z}_\phi}{2} a_{22} + \tilde{r}_\phi \Re(e^{i\tilde{\Xi}_\phi} a_{12})$$

where $d\xi$ is the $SU(2)$ invariant measure on S^2 normalized to 1 and

$$(3.32) \quad \tilde{\mathbf{x}}_\phi := (\tilde{x}_\phi, \tilde{y}_\phi, \tilde{z}_\phi)$$

denotes the mean point of ϕ , that is

$$\tilde{x}_\phi := \int_{S^2} \phi(\xi) x_\xi d\xi$$

with similar notation for \tilde{y}_ϕ , \tilde{z}_ϕ , and $\tilde{r}_\phi e^{i\tilde{\Xi}_\phi} := \tilde{x}_\phi + i\tilde{y}_\phi$. The correspondence between a state and a mean point,

$$(3.33) \quad \varphi \longleftrightarrow \tilde{\mathbf{x}}_\phi,$$

is 1-to-1, that is $\mathcal{S}(M_2(\mathbb{C}))$ is the 3-ball. But unlike the commutative case, two distinct probability measures may have the same mean point, so that the correspondence between states and probability measures is not 1-to-1.

Let us first consider the spectral triple (3.21), that is $M_2(\mathbb{C})$ acting on \mathbb{C}^2 with D an arbitrary selfadjoint 2×2 matrix. As in §3.3, we chose as basis of \mathbb{C}^2 an orthonormal eigenbasis of D , so that the north and south poles of S^2 are the image of the eigenstates of D by (3.29).

PROPOSITION 3.9. *Assume the two eigenvalues d_1, d_2 of D are distinct (otherwise D is proportional to the identity and all the distances are infinite).*

- i. *Two points $\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}$ in the 3-ball are at finite distance iff $\tilde{z}_\phi = \tilde{z}_{\phi'}$.*
- ii. *The distance between two points $\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}$ with the same z -coordinate is proportional to the chord distance on the circle:*

$$(3.34) \quad d(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}) = \frac{1}{|d_1 - d_2|} \sqrt{(\tilde{x}_\phi - \tilde{x}_{\phi'})^2 + (y_\phi - y_{\phi'})^2}.$$

PROOF. The result has been shown for pure states in [34, Prop. 2]¹. The proof easily adapts to non-pure states as follows. If $\tilde{z}_\phi \neq \tilde{z}_{\phi'}$, then for

$$(3.35) \quad b = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \in \text{Ker } L_D$$

one has by (3.31) that $\varphi(b) \neq \varphi'(b)$, meaning the distance is infinite by lemma 2.8.

Assume $\tilde{z}_\phi = \tilde{z}_{\phi'}$. Then

$$(3.36) \quad |\varphi(a) - \varphi'(a)| = \Re(a_{12}(\tilde{x}_\phi - \tilde{x}_{\phi'} + i(\tilde{y}_\phi - \tilde{y}_{\phi'})))$$

$$(3.37) \quad \leq |a_{12}| |(\tilde{x}_\phi - \tilde{x}_{\phi'} + i(\tilde{y}_\phi - \tilde{y}_{\phi'}))|.$$

A direct calculation shows that

$$(3.38) \quad L_D(a) = |a_{12}| |d_1 - d_2|$$

so that

$$(3.39) \quad d(\varphi, \varphi') \leq \frac{1}{|d_1 - d_2|} \sqrt{(\tilde{x}_\phi - \tilde{x}_{\phi'})^2 + (\tilde{y}_\phi - \tilde{y}_{\phi'})^2}.$$

This upper bound is attained by $a = a^* \in M_2(\mathbb{C})$ with components $a_{11} = a_{22} = 0$ and $a_{12} = \frac{1}{|d_1 - d_2|} e^{-i\theta}$ with $\theta = \arg(\tilde{x}_\phi - \tilde{x}_{\phi'} + i(\tilde{y}_\phi - \tilde{y}_{\phi'}))$. \square

¹Notice the misprint of a factor 2 in the result as expressed below Prop. 2

The proposition above shows that the simplest spectral triple on $M_2(\mathbb{C})$ equips the 3-ball with a metric that slices the ball into circles at infinite distance from one another, in particular the poles of S^2 are at infinite distance from any other state. To avoid such infinities, according to lemma 2.7 one needs to reduce the kernel of the semi-norm L_D to the multiples of the identity. This can be done by changing the space of representation and the operator D . An exemple is the following spectral triple, which comes from the truncation of the spectral triple of the Moyal plane in § 6.1. Namely, one makes

$$(3.40) \quad \mathcal{A} = M_2(\mathbb{C}) \quad \text{act on} \quad \mathcal{H} = M_2(\mathbb{C}) \otimes \mathbb{C}^2 \quad \text{as} \quad m \otimes \mathbb{I}_2,$$

where the element m of the algebra $M_2(\mathbb{C})$ acts on the Hilbert space $M_2(\mathbb{C})$ by matrix multiplication. As a Dirac operator, one takes

$$(3.41) \quad D = -i\sqrt{2} \begin{pmatrix} 0_2 & [X^\dagger, \cdot] \\ -[X, \cdot] & 0_2 \end{pmatrix}$$

where the non-zero terms are the commutators with the matrix

$$(3.42) \quad X = \frac{1}{\sqrt{\theta}} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

and its adjoint. This operator is the restriction to $M_2(\mathbb{C})$ of the usual Dirac operator of the plane acting on $L^2(\mathbb{R}^2)$ (see [6] for details).

PROPOSITION 3.10. [6, Prop. 4.4]. *The spectral distance between any two states of $M_2(\mathbb{C})$, identified to points $\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}$ of the 3-ball by (3.33) is finite. More exactly,*

$$d(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}) = \sqrt{\frac{\theta}{2}} \times \begin{cases} d_{eq}(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}) & \text{if } |\tilde{z}_\phi - \tilde{z}_{\phi'}| \leq d_{eq}(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}) , \\ \frac{d_{Ec}(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'})^2}{2|\tilde{z}_\phi - \tilde{z}_{\phi'}|} & \text{if } |\tilde{z}_\phi - \tilde{z}_{\phi'}| \geq d_{eq}(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}) , \end{cases}$$

where

$$(3.43) \quad d_{Ec}(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}) = \sqrt{|\tilde{x}_\phi - \tilde{x}_{\phi'}|^2 + |\tilde{y}_\phi - \tilde{y}_{\phi'}|^2 + |\tilde{z}_\phi - \tilde{z}_{\phi'}|^2}$$

denotes the euclidean distance on B^3 while

$$(3.44) \quad d_{eq}(\tilde{\mathbf{x}}_\phi, \tilde{\mathbf{x}}_{\phi'}) = \sqrt{|\tilde{x}_\phi - \tilde{x}_{\phi'}|^2 + |\tilde{y}_\phi - \tilde{y}_{\phi'}|^2}$$

is the Euclidean distance between the projections of the points on the equatorial plane $z = 0$.

Contrary to the simplest spectral triple on $M_2(\mathbb{C})$ of proposition 3.9, with the spectral triple (3.40) inherited from the Moyal plane, the spectral distance induces on B^3 the Euclidean topology, which coincides with the weak* topology [6, §4.3].

For sake of completeness, let us mention another example of finite dimensional spectral triple that allows to orientate the 3-ball, by adding one point at finite distance from one of the pole of S^2 . This is obtained by letting

$$(3.45) \quad \mathcal{A} = M_n(\mathbb{C}) \oplus \mathbb{C} \quad \text{act on} \quad \mathcal{H} = \mathbb{C}^n \oplus \mathbb{C}$$

as

$$(3.46) \quad a = \begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix},$$

with $x \in M_n(\mathbb{C})$ and $y \in \mathbb{C}$. As a Dirac operator, one takes

$$(3.47) \quad D = \begin{pmatrix} 0_n & v \\ v^* & 0_n \end{pmatrix}.$$

where $v \in \mathbb{C}^n$.

PROPOSITION 3.11. [34, Prop. 3] *For two pure states ω_ξ, ω_ζ of $M_n(\mathbb{C})$ such that $\xi_j = e^{i\theta} \zeta_j$ for all $j \in [2, n]$, the distance is*

$$(3.48) \quad d(\omega_\xi, \omega_\zeta) = \frac{2}{\|v\|} \sqrt{1 - |\langle \xi, \zeta \rangle|^2}.$$

Furthermore, the pure state ω_c of \mathbb{C} is at infinite distance from all the pure states of $M_n(\mathbb{C})$, except ω_v for which

$$d(\omega_c, \omega_v) = \frac{1}{\|v\|}.$$

Applied to $M_2(\mathbb{C}) \oplus \mathbb{C}$, one has that the space of pure states is the disjoint union of the sphere S^2 and the point ω_c . On the sphere the condition of finitude of the distance is the same as in proposition 3.9: S^2 is sliced in circles at infinite distance from one another, while on each circle the distance is proportional to the Euclidean distance on the disk. The pure state ω_v gives the north pole of the sphere, and is at finite distance from ω_c . In this sense adding a point allows to give an orientation to the sphere, by distinguishing between the south pole at infinite distance from any other points and the north pole at finite distance from the isolated point ω_c .

4. Product of geometries and the Higgs field

We now consider the metric aspect of the product of spectral triples. Recall that a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ (with representation π) is *graded* if there exists a grading Γ of \mathcal{H} (that is a selfadjoint operator Γ such that $\Gamma^2 = \mathbb{I}$) which satisfies

$$(4.1) \quad \Gamma D = -D\Gamma, \quad [\Gamma, \pi(a)] = 0 \quad \forall a \in \mathcal{A}.$$

Given two spectral triples $T_1 = (\mathcal{A}_1, \mathcal{H}_1, D_1)$, $T_2 = (\mathcal{A}_2, \mathcal{H}_2, D_2)$ where we assume that T_1 is graded with grading Γ_1 , the product

$$(4.2) \quad \mathcal{A} = \mathcal{A}_1 \otimes \mathcal{A}_2, \quad \mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2, \quad D = D_1 \otimes \mathbb{I}_2 + \Gamma_1 \otimes D_2$$

is again a spectral triple [14] that we denote

$$(4.3) \quad T := T_1 \times T_2.$$

We are interested in the spectral distance d associated to T , and how it is related to the distance d_1, d_2 associated to T_1 and T_2 .

General results on that matter are recalled in §4.1. In §4.2 we focus on the case where T_1 is the usual spectral triple of a closed (spin) manifold described in (2.9) - (2.12) and T_2 a finite dimensional spectral triple as investigated in section 3. In that case, the product T describes a slightly noncommutative generalization of a manifold, called *almost commutative geometry*, which is important for physical applications since it is at the hearth of the description of the standard model of particle physics, as explained in §4.3. The bundle structure of the space of pure states then also opens interesting links with sub-Riemannian geometry. This is the object of section 5.

4.1. Pythagoras inequality. Till recently, the metric aspect of product of spectral triples had been studied mainly for almost commutative geometries. In particular, it came out that for the spectral triple describing the standard model of elementary particles, the distance d between pure states satisfies the Pythagoras equality with respect to the distances d_1 on the manifold and the distance d_2 of the finite dimensional spectral triple describing the gauge degrees of freedom [48]. A similar result was found for the product of the Moyal plane with the two-point space of §3.1 [43]. This raises the question whether the product (4.2) is always orthogonal in the sense of Pythagoras theorem. By this we intend that given two separable² states of $\mathcal{A} = \mathcal{A}_1 \otimes \mathcal{A}_2$,

$$(4.4) \quad \varphi := \varphi_1 \otimes \varphi_2, \quad \varphi' := \varphi'_1 \otimes \varphi'_2,$$

does one have - at least between pure states - that

$$(4.5) \quad d^2(\varphi, \varphi') \text{ equals } d_1^2(\varphi_1, \varphi'_1) + d_2^2(\varphi_2, \varphi'_2) ?$$

In [21] we proved the following Pythagoras inequalities for the product of arbitrary unital spectral triples. For a complete and pedagogical treatment on that matter, as well as some significant generalizations, we invite the reader to see the contribution of F. D'Andrea in this volume [19].

THEOREM 4.1. [21, Theo. 5] *Given the product (4.2) of two spectral triples $(\mathcal{A}_i, \mathcal{H}_i, D_i)$, $i = 1, 2$ and two separable states $\varphi = \varphi_1 \otimes \varphi_2$ and $\varphi' = \varphi'_1 \otimes \varphi'_2$ of \mathcal{A} , one has:*

$$(4.6) \quad d(\varphi, \varphi') \leq d_1(\varphi_1, \varphi'_1) + d_2(\varphi_2, \varphi'_2) .$$

Furthermore, if the spectral triples are unital, then

$$(4.7) \quad d(\varphi, \varphi') \geq \sqrt{d_1(\varphi_1, \varphi'_1)^2 + d_2(\varphi_2, \varphi'_2)^2} .$$

Combining (4.7) and (4.6) one obtains a noncommutative version of Pythagoras theorem that holds true for any separable states in the product of arbitrary unital spectral triples (it was first proven in [43, Prop. II.4] for pure states, with one of the spectral triples the two-point space \mathbb{C}^2).

COROLLARY 4.2. [21] *Let $\varphi = \varphi_1 \otimes \varphi_2$, $\varphi' = \varphi'_1 \otimes \varphi'_2$ be two separable states in the product of two unital spectral triples. Then*

$$(4.8) \quad \sqrt{d_1(\varphi_1, \varphi'_1)^2 + d_2(\varphi_2, \varphi'_2)^2} \leq d(\varphi, \varphi') \leq \sqrt{2} \sqrt{d_1(\varphi_1, \varphi'_1)^2 + d_2(\varphi_2, \varphi'_2)^2} .$$

Furthermore these inequalities are optimal, in that there exist examples that saturate the bounds.

Notice that (4.6) is not the triangle inequality

$$(4.9) \quad d(\varphi, \varphi') \leq d(\varphi_1 \otimes \varphi_2, \varphi_1 \otimes \varphi'_2) + d(\varphi_1 \otimes \varphi'_2, \varphi'_1 \otimes \varphi'_2)$$

because nothing guarantees that the distance between two states $\varphi_1 \otimes \varphi_2, \varphi_1 \otimes \varphi'_2$ that differ only on one of the algebras gives back the distance on a single spectral triple, that is $d(\varphi_1 \otimes \varphi_2, \varphi_1 \otimes \varphi'_2)$ equals $d_2(\varphi_2, \varphi'_2)$. In fact, this comes out as a corollary, initially proven in [48].

²A state $\varphi \in \mathcal{S}(\mathcal{A})$ is said separable if it decomposes as the product of two states $\varphi_1 \in \mathcal{S}(\mathcal{A}_1)$, $\varphi_2 \in \mathcal{S}(\mathcal{A}_2)$

COROLLARY 4.3. [21, Cor. 6] *Let $\varphi = \varphi_1 \otimes \varphi_2$, $\varphi' = \varphi'_1 \otimes \varphi'_2$ be two separable states in the product of two unitary spectral triples. If $\varphi_2 = \varphi'_2$, then $d(\varphi, \varphi') = d_1(\varphi_1, \varphi'_1)$, and similarly if $\varphi_1 = \varphi'_1$ then $d(\varphi, \varphi') = d_2(\varphi_2, \varphi'_2)$.*

To conclude the generalities on the product of spectral triples, let us mention an application of the projection lemma 2.5. It is not of great interest in se because of the strong conditions required, but it turns out to be extremely useful to compute the distance in the standard model of elementary particles, as explained in the next subsection. Let us consider the product (4.2) and restrict the attention to normal states for, say, the algebra \mathcal{A}_2 . To any such state φ is associated a support, namely a projection $s \in \mathcal{A}_2$ such that φ is faithful on $s\mathcal{A}_2s$. For a pure state ω , being normal implies

$$(4.10) \quad sas = \omega(a)s \quad \forall a \in \mathcal{A}_2.$$

We say that two normal pure states ω_1, ω_2 are in direct sum if

$$(4.11) \quad s_1as_2 = 0 \quad \forall a \in \mathcal{A}_2.$$

If furthermore the sum $p =: s_1 + s_2$ of their support commutes with D_2 , then the distance in the product (4.2) projects down to a two point-case $\mathcal{A}_1 \otimes \mathbb{C}^2$.

PROPOSITION 4.4. [48] *Let d be the distance associated with the product $T = T_1 \times T_2$. Let ω_2, ω'_2 be normal pure states of \mathcal{A}_2 in direct sum, and whose sum of supports p commutes with D_2 . Then for any pure states ω_1, ω'_1 of \mathcal{A}_1 one has*

$$(4.12) \quad d(\omega_1 \otimes \omega_2, \omega'_1 \otimes \omega'_2) = d_e(\omega_1 \otimes \omega_c, \omega'_1 \otimes \omega'_c)$$

where ω_c, ω'_c are the two pure states of \mathbb{C}^2 while d_e is the distance associated to the product $T_e := T_1 \times T_r$ where $T_r := (\mathcal{A}_r, \mathcal{H}_r, D_r)$ with

$$(4.13) \quad \mathcal{A}_r := \mathbb{C}^2, \quad \mathcal{H}_r := p\mathcal{H}_2, \quad D_r := pD_2p|_{\mathcal{H}_r}.$$

Note that this proposition remains true for an algebra \mathcal{A}_2 on a field other than \mathbb{C} , assuming that the notion of states is still available. For instance in the standard model one deals with real algebras.

4.2. Almost commutative geometries and fluctuation of the metric.

A slightly noncommutative generalization of a manifold is obtained by taking the product (4.2) of the spectral triple of a closed, spin manifold \mathcal{M} , that is (see (2.11))

$$(4.14) \quad T_1 = (C^\infty(\mathcal{M}), L^2(\mathcal{M}, S), \not{D}),$$

by a finite dimensional spectral triple $T_2 = (\mathcal{A}_F, \mathcal{H}_F, D_F)$. Namely one considers

$$(4.15) \quad \mathcal{A} = C^\infty(\mathcal{M}) \otimes \mathcal{A}_F, \quad \mathcal{H} = L^2(\mathcal{M}, S) \otimes \mathcal{H}_F, \quad D = \not{D} \otimes \mathbb{I}_F + \gamma^5 \otimes D_F$$

where \mathbb{I}_F is the identity operator on \mathcal{H}_F and γ^5 is the grading of $L^2(\mathcal{M}, S)$ given by the product of the Dirac matrices. The center of \mathcal{A} is infinite dimensional (as an algebra) while the noncommutative part is finite dimensional, hence the name *almost commutative geometries* often used to describe (4.15).

Because $C^\infty(\mathcal{M})$ is nuclear, the space of pure states $\mathcal{P}(\mathcal{A})$ of \mathcal{A} is [35]

$$(4.16) \quad \mathcal{P}(\mathcal{A}) \simeq \mathcal{P}(C^\infty(\mathcal{M})) \times \mathcal{P}(\mathcal{A}_F),$$

and $\mathcal{S}(\mathcal{A})$ its convex hull. $\mathcal{P}(C^\infty(\mathcal{M}))$ is homeomorphic to \mathcal{M} and $\mathcal{P}(\mathcal{A}_F)$ carries a natural action of the special unitarie group $SU(\mathcal{A}_F)$ of \mathcal{A}_F ,

$$(4.17) \quad \omega \rightarrow \omega \circ \alpha_u \quad \forall \omega \in \mathcal{P}(\mathcal{A}_F)$$

with α_u the inner automorphism of \mathcal{A}_F given by conjugate action of $u \in SU(\mathcal{A}_F)$. In other terms, $\mathcal{P}(\mathcal{A})$ is a trivial $SU(\mathcal{A}_F)$ -bundle on \mathcal{M} with fiber $\mathcal{P}(\mathcal{A}_F)$.

In the study of noncommutative algebras (or more generally noncommutative rings), there exists a notion of equivalence which is weaker than isomorphism but turns out to be very fruitful, that of Morita equivalence. Given a spectral triple $(\mathcal{A}, \mathcal{H}, D)$, there is a generic procedure to export the geometrical structure to a Morita equivalent algebra [14]. Taking advantage of the self-Morita equivalence of \mathcal{A} , this procedure yields a natural way to introduce a connection in the geometry $(\mathcal{A}, \mathcal{H}, D)$, by substituting the operator D with a *covariant Dirac operator* D_A , such that $(\mathcal{A}, \mathcal{H}, D_A)$ is still a spectral triple. Explicitly, this covariant operator is

$$(4.18) \quad D_A := D + A + JAJ^{-1},$$

where A is a selfadjoint element of the set of generalized 1-forms ³

$$(4.19) \quad \Omega_D^1 := \{a^i[D, b_i], \quad a^i, b_i \in \mathcal{A}\},$$

and J is the *real structure*. The latter is a generalization to the non-commutative setting of the charge conjugation operator on spinors. The only thing we need to know about it at the moment is that for any $a, b \in \mathcal{A}$ one has $[JaJ^{-1}, b] = 0$, so that substituting D with D_A in the distance formula yields

$$(4.20) \quad d_A(\varphi, \varphi') := \sup_{a \in \mathcal{A}} \{|\varphi(a) - \varphi'(a)|, \|[D + A, a]\| \leq 1\}.$$

There is no reason for $\|[D_A, a]\|$ to equal $\|[D, a]\|$, neither for the distance d_A computed with the covariant Dirac operator D_A to equal the distance d computed with the initial operator D . That is why one talks of a *fluctuation of the metric*.

For almost commutative geometries (4.15), a generalized 1-form in Ω_D^1 is [17]:

$$(4.21) \quad A = -i\gamma^\mu f_\mu^i \otimes m_i + \gamma^5 h^j \otimes n_j$$

where $m_i \in \mathcal{A}_F$, $h^j, f_\mu^i \in C^\infty(\mathcal{M})$, while

$$(4.22) \quad n_j \in \Omega_{D_F}^1 := \{a^i[D_F, b_i], \quad a^i, b_i \in \mathcal{A}_F\}.$$

Omitting the tensor product, a selfadjoint 1-forms $A = A^* \in \Omega_D^1$ thus decomposes as the sum

$$(4.23) \quad A = -i\gamma^\mu A_\mu + \gamma^5 H,$$

where

$$(4.24) \quad A_\mu := f_\mu^i m_i$$

is a \mathcal{A}_F -valued skew-adjoint 1-form field over \mathcal{M} , and

$$(4.25) \quad H := h^j n_j$$

is a $\Omega_{D_F}^1$ -valued selfadjoint scalar field. The part of the covariant Dirac operator D_A relevant for the distance formula is the fluctuated operator

$$(4.26) \quad \not{D} + A = \not{D} + \gamma^5 H - i\gamma^\mu A_\mu.$$

We investigate below how the distance on the bundle of pure states (4.16) is affected by the two pieces of the fluctuation: the scalar fluctuation H in §4.3, and the gauge fluctuation A_μ in §5.

³We use Einstein summation over repeated indices in alternate positions (up/down).

4.3. Two sheet models and the metric interpretation of the Higgs.

Let us consider a scalar fluctuation of the metric, namely formula (4.20) with $D + A$ given by (4.26) where

$$(4.27) \quad H \neq 0, \quad A_\mu = 0.$$

This amounts to take the product of the manifold by an internal geometry

$$(4.28) \quad T_F^x := (\mathcal{A}_F, \mathcal{H}_F, D_F(x) := D_F + H(x))$$

in which D_F is now a non-constant section of $\text{End } \mathcal{H}_F$.

Let d_{geo} , d_x , d_H denote the geodesic distance on \mathcal{M} , the spectral distance associated to the spectral triple T_F^x , and the distance (4.20) with generalized 1-form A given by (4.27). Corollary 4.3 and proposition 4.4 gives respectively

PROPOSITION 4.5. [48, Theo. 2'] *For any pure states δ_x, δ_y of $C^\infty(\mathcal{M})$ and $\omega_F, \omega'_F \in \mathcal{P}(\mathcal{A}_F)$, one has*

$$\begin{aligned} d_H(\delta_x \otimes \omega_F, \delta_x \otimes \omega'_F) &= d_x(\omega_F, \omega'_F), \\ d_H(\delta_x \otimes \omega_F, \delta_y \otimes \omega_F) &= d_{\text{geo}}(x, y). \end{aligned}$$

PROPOSITION 4.6. [48, Theo. 4'] *Let ω_1, ω_2 be two normal pure states of \mathcal{A}_F with support s_1, s_2 in direct sum, and such that the sum of their support commutes with $D_F(x)$ for all x . Then*

$$d(\delta_x \otimes \omega_1, \delta_y \otimes \omega_2) = L'((0, x), (1, y)),$$

where L' is the geodesic distance in the manifold $\mathcal{M}' := [0, 1] \times \mathcal{M}$ equipped with the metric

$$(4.29) \quad \begin{pmatrix} \|\tilde{H}(x)\|^2 & 0 \\ 0 & g^{\mu\nu}(x) \end{pmatrix}$$

in which $g^{\mu\nu}$ is the metric on \mathcal{M} and \tilde{H} is projection on $s_2 \mathcal{A}_F$ of the restriction of D_H to $s_1 \mathcal{A}_F$.




FIGURE 2. Space-time of the standard model with a pure scalar fluctuation of the metric.

Proposition 4.6 gives an intuitive picture of the spacetime of the standard model. The later is described by an almost commutative geometry (4.15) where

$$(4.30) \quad \mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}),$$

with \mathbb{H} the algebra of quaternions. \mathcal{A}_F is suitably represented over a finite dimensional vector space \mathcal{H}_F generated by elementary fermions, while D_F is a finite dimensional matrix that contains the masses of the elementary fermions together with the Cabibbo matrix and neutrinos mixing angles. We refer the reader to [9] for the most advanced version of the model pre-discovery of the Higgs, and [10], [24], [23] for enhanced version post-Higgs. Through the *spectral action* [8] the scalar fluctuation H further identifies with the Higgs field [14] (see also [28] for the first appearance of the Higgs field as a connection in a noncommutative space).

From the metric point of view, one finds that the pure states of $M_3(\mathbb{C})$ are at infinite distance from one another, whereas the states of \mathbb{C} and \mathbb{H} are in direct sum, with support the identity. Hence the model of spacetime that emerges is a two-sheet model, two copies of the manifold, one indexed by the pure state of \mathbb{C} , the other one by the pure state of \mathbb{H} (cf figure 2).

PROPOSITION 4.7. [48, Prop. 8] *The distance between the two sheets coincides with the geodesic distance in a $(\dim \mathcal{M}) + 1$ dimension manifold, and the extra-component of the metric is*

$$(4.31) \quad \left\| \tilde{H}(x) \right\|^2 = (|1 + h_1(x)|^2 + |h_2(x)|^2) m_t^2$$

where the h_i 's are the components of the Higgs field and m_t is the mass of the quark top.

5. Sub-Riemannian geometry from gauge fluctuation of the metric

In this section we study a gauge fluctuation of the metric, that is formula (4.20) with D substituted with (4.26) where

$$(5.1) \quad A_\mu \neq 0, H = 0.$$

This is a review of [44] and [46].

As recalled in §5.1, it was expected that the spectral distance on the bundle of pure states (4.16) were equal to the Carnot-Carathéodory (or horizontal) distance associated to the sub-Riemannian structure defined by the 1-form field A_μ . In fact, the link between the two distances is more intricate and interesting. As explained in §5.2, the horizontal distance is an upper bound to the spectral distance, but it has no reason to be the lowest one, unless the holonomy is trivial. In §5.3 we study the example where the base manifold \mathcal{M} is a circle. The holonomy is not trivial, and indeed the two distances are not equal. We show it by working out the connected components of both distances, in case $\mathcal{A}_F = M_2(\mathbb{C})$. This result is extended to $\mathcal{A}_F = M_n(\mathbb{C})$ with $n \geq 2$ in §5.4. The two remaining sections contain exact computations of the spectral distance: on the whole of the bundle of pure states for $\mathcal{A}_F = M_2(\mathbb{C})$ (§5.5); between two pure states on the same fiber in case $\mathcal{A}_F = M_n(\mathbb{C})$ for arbitrary n (§5.6).

Let us mention that other applications of noncommutative geometry to sub-Riemannian geometry have been investigated in [33].

5.1. Horizontal structure on the space of pure states. A gauge fluctuation (5.1) is obtained from an almost commutative geometry (4.15) by taking $D_F = 0$. In practical, we take as a finite dimensional spectral triple

$$(5.2) \quad \mathcal{A}_F = M_n(\mathbb{C}), \quad \mathcal{H}_F = M_n(\mathbb{C}), \quad D_F = 0$$

for some $n \in \mathbb{N}$, so that the almost commutative geometry we are dealing with is

$$(5.3) \quad \mathcal{A} = C^\infty(\mathcal{M}) \otimes M_n(\mathbb{C}), \quad \mathcal{H} = L^2(\mathcal{M}, S) \otimes M_n(\mathbb{C}), \quad D = \not{D} \otimes \mathbb{1}_F.$$

The vanishing of D_F implies that the scalar part H of the fluctuation in (4.25) vanishes. Since the spin connection in \not{D} commutes with the algebra, the part of the fluctuated operator (4.26) relevant in the distance formula reduces to

$$(5.4) \quad D_\mu := -i\gamma^\mu(\partial_\mu + A_\mu).$$

As explained in §3.3, the space $\mathcal{P}(M_n(\mathbb{C}))$ of pure states of $M_n(\mathbb{C})$ identifies with the projective space $\mathbb{C}P^{n-1}$. The action (4.17) of $SU(n)$ on $\mathcal{P}(M_n(\mathbb{C}))$ reads as the free action of $U(n)$ on $\mathbb{C}P^{n-1}$,

$$(5.5) \quad \xi \rightarrow u\xi \quad \forall \xi \in \mathbb{C}P^{n-1}, u \in U(n),$$

and $\mathcal{P}(\mathcal{A})$ in (4.16) is now the trivial $SU(n)$ -bundle with fiber $\mathbb{C}P^{n-1}$

$$(5.6) \quad P \xrightarrow{\pi} M.$$

We denote

$$(5.7) \quad \xi_x := (\delta_x \in \mathcal{P}(C^\infty(\mathcal{M})), \omega_\xi \in \mathcal{P}(M_n(\mathbb{C})))$$

an element of P , where δ_x is the evaluation (2.1) and ω_ξ is the pure state of $M_n(\mathbb{C})$ defined by (3.24). Its evaluation on an element of \mathcal{A}

$$(5.8) \quad a = f^i \otimes m_i, \quad f^i \in C^\infty(\mathcal{M}), m_i \in M_n(\mathbb{C})$$

reads

$$(5.9) \quad \xi_x(a) = \langle \xi, a(x)\xi \rangle,$$

where for any x in \mathcal{M} one writes

$$(5.10) \quad a(x) = f^i(x) \otimes m_i \in M_n(\mathbb{C}).$$

The gauge part A_μ of the fluctuation has value in the set of skew-adjoint elements of $M_n(\mathbb{C})$, that is the Lie algebra $\mathfrak{u}(n)$. Thus A_μ is the local form of the 1-form field associated to some Ehresmann connection Ξ on the trivial $U(n)$ -principal bundle on \mathcal{M} . By reduction to $SU(n)$ followed by a mapping to the associated bundle (5.6), one inherits from Ξ a connection on the bundle P of pure states of \mathcal{A} . This means that at any $p \in P$ the tangent space $T_p P$ splits into a vertical subspace and an horizontal subspace,

$$(5.11) \quad T_p P = V_p P \oplus H_p P \quad p \in P,$$

where HP is the kernel of the connection 1-form associated to Ξ .

A curve $t \in [0, 1] \mapsto c(t) \in P$ is horizontal when its tangent vector is everywhere horizontal, that is for any t one has

$$(5.12) \quad \dot{c}(t) \in H_{c(t)} P.$$

The horizontal (also called Carnot-Carathéodory) distance $d_h(p, q)$ is defined as the infimum on the length of the horizontal paths joining p to q ,

$$(5.13) \quad d_h(p, q) := \inf_{\dot{c}(t) \in \mathcal{H}_{c(t)}P} \int_0^1 \|\dot{c}(t)\| dt \quad \forall p, q \in P,$$

where the norm on HP is the pull back of the metric⁴

$$(5.14) \quad \|\dot{c}\| = \sqrt{g(\pi_*(\dot{c}), \pi_*(\dot{c}))}.$$

When p, q cannot be linked by any horizontal path then $d_h(p, q)$ is infinite.

To summarize, the gauge part A_μ of the covariant Dirac operator (4.26) equips the bundle P of pure states of an almost commutative geometry with two distances: the horizontal distance d_h (5.13) and the fluctuated spectral distance d_A (4.20) computed with D_μ . The rest of this section is a collection of results regarding the comparison of these two distances.

5.2. Holonomy obstruction.

DEFINITION 5.1. *A pure state at finite horizontal distance from ξ_x is said accessible, and we define*

$$(5.15) \quad \text{Acc}(\xi_x) := \{q \in P; d_h(\xi_x, q) < +\infty\}.$$

A pure state at finite spectral distance from ξ_x is said connected, and we define

$$(5.16) \quad \text{Con}(\xi_x) := \{q \in P; d_A(\xi_x, q) < +\infty\}.$$

We use the same notation as in (2.23) although here we restrict to pure states.

In the same way as the spectral distance on a manifold is bounded by the geodesic distance, for almost commutative geometry with gauge fluctuation the horizontal distance provides an upper bound to the spectral distance.

PROPOSITION 5.2. [44, Prop. 1] *For any $\xi_x, \zeta_y \in P$,*

$$(5.17) \quad d_A(\xi_x, \zeta_y) \leq d_h(\xi_x, \zeta_y) \quad \forall \xi_x, \zeta_y \in P.$$

In other terms

$$(5.18) \quad \text{Acc}(\xi_x) \subset \text{Con}(\xi_x).$$

However this upper bound is not optimal. In [14] was suggested that d_A and d_h were equal. This is true when the holonomy group reduces to the identity: then $\text{Acc}(\xi_x) = \text{Con}(\xi_x)$ coincides with the horizontal lift of \mathcal{M} passing through ξ_x . In particular, on a given fiber there is no points accessible from one another and both the spectral and the horizontal distances are infinite.

However when the holonomy is not trivial, then $\text{Acc}(\xi_x)$ has no reason to equal $\text{Con}(\xi_x)$. The obstruction comes from the number of times a minimal horizontal curve between ξ_x and $p \in \text{Acc}(\xi_x)$ - that is an horizontal curve whose length is the horizontal distance - intersects the same orbit of the holonomy group. To be more explicit, given an horizontal curve c between ξ_x and ζ_y , we call *ordered self-intersecting points at $p_0 = c(t_0)$* a set of K elements $p_1 := c(t_1), \dots, p_K := c(t_K)$ such that for any $i = 1, \dots, K$

$$(5.19) \quad \pi(p_i) = \pi(p_0), \quad d_h(p_0, p_{i+1}) > d_h(p_0, p_i).$$

⁴In all this section, π denotes the projection from P to \mathcal{M} , and not the representation of the algebra in the spectral triple.




FIGURE 3. An ordered sequence of self-intersecting points.

Assuming the spectral distance between to pure states ξ_x, ζ_y is the horizontal distance, and that there exists at least one minimal horizontal curve between ξ_x and ζ_y , then one has the following constraint on the optimal element of definition 2.1.

PROPOSITION 5.3. *Let ξ_x, ζ_y be two points in P such that $d_A(\xi_x, \zeta_y) = d_h(\xi_x, \zeta_y)$. Then for any minimal horizontal curve c between ξ_x and ζ_y one has*

$$(5.20) \quad d_A(\xi_x, c(t)) = d_h(\xi_x, c(t)).$$

Moreover there exists an optimal element $a \in \mathcal{A}$ such that for any $\xi_t := c(t)$

$$(5.21) \quad \xi_t(a) = d_h(\xi_x, c(t)) \quad \text{or} \quad \lim_{n \rightarrow \infty} \xi_t(a_n) = d_h(\xi_x, c(t)).$$

Consequently, assuming there is a minimal horizontal curve between ξ_x and ζ_y with K self-intersecting points at p_0 , proposition 5.3 puts $K + 1$ condition on the n^2 real components of the selfadjoint matrix $a(\pi(p_0))$,

$$(5.22) \quad p_i(a) = \text{Tr}(s_{p_i} a(\pi(p_0))) = d_h(\xi_x, p_i) \quad \forall i = 0, 1, \dots, K$$

where s_{p_i} denotes the support of the pure state p_i . So it is most likely that $d_A(\xi_x, \zeta_y)$ cannot equal $d_h(\xi_x, \zeta_y)$ unless there exists a minimal horizontal curve between ξ_x and ζ_y such that its projection does not self-intersect more than n^2 times. Actually, questioning the equality between d_A and d_h amounts to the following problem:

Given a minimal horizontal curve c , is there a way to deform it into another horizontal curve c' , keeping its length and its end-points fixed, such that c' has less selfintersecting points than c ?

Say differently:

Can one characterize the minimum number of selfintersecting points in a minimal horizontal curve between two given points ?

It seems that there is no known answer to these questions [50]. In some cases it might be possible indeed to reduce the number of self-intersecting points of a minimal horizontal curve by smooth deformations that keep its length constant (see [46, §2.3]). In order to escape these issues, we consider a case where there is at most one minimal horizontal curve between two points: bundles on the circle S^1 .

5.3. The counter-example of the circle. We consider (5.3) for $\mathcal{M} = S^1$. The gauge fluctuation A_μ has only one component A and we fix on \mathbb{C}^n a basis of real eigenvectors of iA such that

$$(5.23) \quad A = i \begin{pmatrix} \theta_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \theta_n \end{pmatrix},$$

where the θ_j 's are real functions on S^1 . The space of pure states of

$$(5.24) \quad \mathcal{A} = C^\infty(S^1) \otimes M_n(\mathbb{C})$$

is a $\mathbb{C}P^{n-1}$ bundle $P \xrightarrow{\pi} S^1$ on the circle. In a trivialization (π, V) , we associate to the pure state $\xi_x \in P$ with

$$(5.25) \quad V(\xi_x) = \xi = \begin{pmatrix} V_1 \\ \vdots \\ V_n \end{pmatrix} \in \mathbb{C}P^{n-1},$$

the $n-1$ -torus of $\mathbb{C}P^{n-1}$

$$(5.26) \quad T_\xi := \left\{ \begin{pmatrix} V_1 \\ e^{i\varphi_j} V_j \end{pmatrix}, \varphi_j \in \mathbb{R}, j = 2, \dots, n \right\},$$

and the n -torus of P ,

$$(5.27) \quad \mathbb{T}_\xi := S^1 \times T_\xi.$$

The set $\text{Acc}(\xi_x)$ of points in P accessible to ξ_x is the horizontal lift $c(\tau)$, $\tau \in \mathbb{R}$, of the circle with initial conditions $\pi(c(0)) = x$, $V(c(0)) = \xi$. Explicitly, one has

$$(5.28) \quad c(\tau) = (c_*(\tau), V(\tau))$$

where $c_*(\tau) := \pi(c(\tau))$ while $V(\tau)$ has components

$$(5.29) \quad V_j(\tau) = V_j e^{-i\Theta_j(\tau)} \quad \text{with} \quad \Theta_j(\tau) := \int_0^\tau \theta_j(t) dt.$$

Hence on a given fiber $\pi^{-1}(c_*(\tau))$ the set of accessible points is the sub-torus of T_ξ ,

$$(5.30) \quad H_\tau^\xi := \text{Acc}(\xi_x) \cap \pi^{-1}(c_*(\tau)) = \left\{ \begin{pmatrix} V_1(\tau) \\ e^{ik\Theta_{1j}(2\pi)} V_j(\tau) \end{pmatrix}, k \in \mathbb{Z}, j = 2, \dots, n \right\}.$$

This is at best dense in T_ξ if all the $\Theta_{1j}(2\pi)$'s are distinct and irrational. The union over all S^1 yields

$$(5.31) \quad \text{Acc}(\xi_x) = \bigcup_{\tau \in [0, 2\pi[} H_\tau^\xi \subsetneq \mathbb{T}_\xi.$$

The simplest counter-example to the equality between the horizontal and the spectral distances is given by the $n = 2$ case (i.e. $T_\xi = S^1$).

PROPOSITION 5.4. [44, Prop. 5], [46, Prop. 3.4] *For $\mathcal{A}_F = M_2(\mathbb{C})$ and a gauge fluctuation A non proportional to the identity, one has*

$$(5.32) \quad \text{Con}(\xi_x) = \mathbb{T}_\xi.$$

Thus by (5.31) one has that $\text{Acc}(\xi_x)$ is at best dense in $\text{Con}(\xi_x)$. Any element of \mathbb{T}_ξ that is not in $\text{Acc}(\xi_x)$ is at finite spectral distance from ξ_x , although it is infinitely Carnot-Carathéodory far from it. This shows that the two distances are not equal.

In this example, the discrepancy between the two distances follows from the holonomy obstruction of Prop. 5.3. The holonomy is non-trivial because the base $\mathcal{M} = S^1$ is non-simply connected. A open question is whether there is the same obstruction when the holonomy comes from the curvature of the connection.

5.4. Connected versus accessible points on the $\mathbb{C}P^{n-1}$ bundle on S^1 .

For $n > 2$, proposition 5.4 needs to be refined. $\text{Con}(\xi_x)$ is still a subset of the torus \mathbb{T}_ξ but not necessary equal to it. Viewing the torus \mathbb{T}_ξ as the subset of \mathbb{R}^n ,

$$(5.33) \quad \mathbb{T}_\xi = \{\tau \in [0, 2\pi[, \varphi_i \in [0, 2\pi[, i = 2, \dots, n\}$$

one has that $\text{Con}(\xi_x)$ is a sub-torus \mathbb{U}_ξ of \mathbb{T}_ξ ,

$$(5.34) \quad \mathbb{U}_\xi = \{\tau \in [0, 2\pi[, \varphi_i \in [0, 2\pi[, i = 2, \dots, n_c\}$$

with dimension $n_c \leq n$ given by the number of equivalence classes of the following relation.

DEFINITION 5.5. Let us fix a pure state ξ_x in P . Two directions i, j of \mathbb{T}_ξ are said *far* from each other if the components i and j of the holonomy at x are equal, and we write $\text{Far}(\cdot)$ the equivalence classes,

$$(5.35) \quad \text{Far}(i) := \{j \in [1, n] \text{ such that } \Theta_j(2\pi) = \Theta_i(2\pi) \bmod [2\pi]\}.$$

We denote n_c the numbers of such equivalence classes and we label them as

$$\text{Far}_1 = \text{Far}(1), \text{Far}_p = \text{Far}(j_p) \quad p = 2, \dots, n_c$$

where $j_p \neq 0$ is the smallest integer that does not belong to $\bigcup_{q=1}^{p-1} \text{Far}_q$. Two directions belonging to distinct equivalence classes are said *close* to each other.

The terminology comes from the following proposition, which shows that the torus-dimension of the connected components for the spectral distance is given by the number of directions close to each other. On the contrary, two directions that are not close to each other do not contribute to the connected components: from the spectral distance point of view, they are infinitely far from each other.

PROPOSITION 5.6. [46, Prop. 3.4] *Con*(ξ_x) is the n_c torus

$$(5.36) \quad \mathbb{U}_\xi := \bigcup_{\tau \in [0, 2\pi[} U_\tau^\xi$$

where $U_\tau^\xi \subset T_\xi$ is the $(n_c - 1)$ torus defined by ($V_i(\tau)$ is given in (5.29))

$$(5.37) \quad U_\tau^\xi := \left\{ \begin{pmatrix} V_i(\tau) & \forall i \in \text{Far}_1 \\ e^{i\varphi_2} V_i(\tau) & \forall i \in \text{Far}_2 \\ \dots & \\ e^{i\varphi_{n_c}} V_i(\tau) & \forall i \in \text{Far}_{n_c} \end{pmatrix}, \varphi_j \in \mathbb{R}, j \in [2, n_c] \right\}.$$

The spectral and the horizontal distances yield two distinct topologies Con and Acc on the bundle of pure states P . Obviously

$$(5.38) \quad e^{i\Theta_{1j}(2k\pi)} = e^{i\Theta_{1i}(2k\pi)} \quad \forall j \in \text{Far}(i),$$

hence $H_\tau^\xi \subset U_\tau^\xi$ fiber-wise and $\text{Acc}(\xi_x) \subset \mathbb{U}(\xi_x)$ globally, as expected from (5.18). Also obvious is the inclusion of \mathbb{U}_ξ within \mathbb{T}_ξ . To summarize the various connected components organize as follows,

$$(5.39) \quad \text{Acc}(\xi_x) \subset \text{Con}(\xi_x) = \mathbb{U}_\xi \subset \mathbb{T}_\xi \subset P,$$

or fiber-wise

$$(5.40) \quad H_\tau^\xi \subset U_\tau^\xi \subset T_\xi \subset \mathbb{C}P^{n-1}.$$

The difference between $\text{Acc}(\xi_x)$ and \mathbb{U}_ξ is governed by the irrationality of the connection, whereas the difference between \mathbb{U}_ξ and \mathbb{T}_ξ is governed by the number of close directions. More specifically

$$(5.41) \quad \mathbb{T}_\xi = \bigcup_{\zeta \in T_\xi} \text{Acc}(\zeta_x)$$

is the union of all states with equal components up to phase factors. Meanwhile

$$(5.42) \quad \mathbb{U}_\xi = \bigcup_{\zeta \in U_\xi} \text{Acc}(\zeta_x),$$

with $U_\xi = U_{\tau=0}^\xi$, is the union of all states with equal components up to phase factors, with the extra-condition that phase factors corresponding to directions far from each other must be equal.

Note that none of the distances is able to "see" between different tori $\mathbb{T}_\xi, \mathbb{T}_\eta$. However within a given \mathbb{U}_ξ the spectral distance "sees" between the horizontal components. In this sense the spectral distance keeps "better in mind" the bundle structure of the set of pure states P (see also figure 6 in §5.6). This suggests that the spectral distance could be relevant to study some transverse metric structure in a more general framework of foliation.

5.5. A low dimensional example. Having individuated the connected components of the spectral distance, we now compute the latter explicitly in two examples: on the whole of the bundle P of pure states in the low dimension case $n = 2$ below, and on a given fiber for arbitrary n in §5.6.

Identifying $\mathcal{P}(M_2(\mathbb{C})) \simeq \mathbb{C}P^{n-1}$ with the 2-sphere via (3.29), the pure state space of $C^\infty(S^1, M_2(\mathbb{C}))$ is a bundle in sphere over S^1 . The pure state ξ_x in (5.25) is mapped to the point

$$(5.43) \quad x_0 = R \cos \theta_0, \quad y_0 = R \sin \theta_0, \quad z_0 = z_\xi$$

on the fiber $\pi^{-1}(x)$, where we define

$$(5.44) \quad 2V_1\overline{V_2} =: Re^{i\theta_0}.$$

The torus T_ξ in (5.26) is mapped to the circle of radius R

$$(5.45) \quad S_R := \{x, y, z \in S^2, z = z_0\},$$

so that by Prop. 5.4 and assuming the holonomy is not trivial, the connected component

$$(5.46) \quad \text{Con}(\xi_x) = \mathbb{T}_\xi = S^1 \times S_R$$

is a 2-dimensional torus (see Figure 4).

The points accessible from ξ_x are given in (5.28) as

$$(5.47) \quad \xi_x^k := c(\tau + 2k\pi), \tau \in [0, 2\pi[, k \in \mathbb{Z}.$$

On the sphere they have coordinates

$$(5.48) \quad x_\tau^k := R \cos(\theta_0 - \theta_\tau^k), \quad y_\tau^k := R \sin(\theta_0 - \theta_\tau^k), \quad z_\tau^k := z_\xi$$

where $\theta_\tau^k := \theta(\tau + 2k\pi)$. $\text{Acc}(\xi_x)$ is discrete or dense within \mathbb{T}_ξ , depending whether $\Theta(2\pi)$ is rational or not.




FIGURE 4. The 2-torus \mathbb{T}_ξ , the accessible point ξ_x^1 and an arbitrary pure state ζ_y .

To compute the spectral distance on \mathbb{T}_ξ we use the following parametrization.

DEFINITION 5.7. *Given ξ_x in P , any pure state ζ_y in the 2-torus \mathbb{T}_ξ is in one-to-one correspondence with an equivalence class*

$$(5.49) \quad (k \in \mathbb{N}, 0 \leq \tau_0 \leq 2\pi, 0 \leq \varphi \leq 2\pi) \sim (k + \mathbb{Z}, \tau_0, \varphi - 2\mathbb{Z}\omega\pi)$$

such that

$$(5.50) \quad \tau = 2k\pi + \tau_0, \quad \omega := \frac{\Theta_1(2\pi) - \Theta_2(2\pi)}{2\pi}, \quad \zeta_y = \begin{pmatrix} V_1(\tau) \\ e^{i\varphi} V_2(\tau) \end{pmatrix}.$$

After a rather lengthy computation, one finds

PROPOSITION 5.8. [46, Prop. 4.5] *Let ξ_x be a pure state in P and $\zeta_y = (k, \tau_0, \varphi)$ a pure state in \mathbb{T}_ξ . Then either the two directions are far from each other so that $\text{Con}(\xi_x) = \text{Acc}(\xi_x)$ and*

$$(5.51) \quad d_A(\xi_x, \zeta_y) = \begin{cases} \min(\tau_0, 2\pi - \tau_0) & \text{when } \varphi = 0 \\ +\infty & \text{when } \varphi \neq 0 \end{cases};$$

or the directions are close to each other so that $\text{Con}(\xi_x) = \mathbb{T}_\xi$ and

$$(5.52) \quad d_A(\xi_x, \zeta_y) = \max_{T_\pm} H_\xi(T, \Delta)$$

where

$$H_\xi(T, \Delta) := T + z_\xi \Delta + RW_{k+1} \sqrt{(\tau_0 - T)^2 - \Delta^2} + RW_k \sqrt{(2\pi - \tau_0 - T)^2 - \Delta^2}$$

with

$$(5.53) \quad W_k := \frac{|\sin(k\omega\pi + \frac{\varphi}{2})|}{|\sin\omega\pi|};$$

and the maximum is on one of the triangles (see fig. 5.5)

$$(5.54) \quad \mathcal{T}_\pm := T \pm \Delta \leq \min(\tau_0, 2\pi - \tau_0)$$

with sign the one of z_ξ .




FIGURE 5. Unit is $\min(\tau_0, 2\pi - \tau_0)$.

For ξ an equatorial state, i.e. $z_\xi = 0$, the result greatly simplifies

PROPOSITION 5.9. [46, §4.3.1]

$$(5.55) \quad d_A(\xi_x, \zeta_y) = H_\xi(0, 0) = RW_{k+1}\tau_0 + RW_k(2\pi - \tau_0).$$

5.6. Distances on the fiber. In the general case $\mathcal{A} = C^\infty(S^1, M_n(\mathbb{C}))$ for arbitrary integer $n \in \mathbb{N}$ one can explicitly compute the spectral distance for two pure states on the same fiber. \mathbb{T}_ξ is now a n -torus and instead of (5.49) one deals with equivalence classes of $(n + 1)$ -tuples

$$(5.56) \quad (k \in \mathbb{N}, 0 \leq \tau_0 \leq 2\pi, 0 \leq \varphi_i \leq 2\pi) \sim (k + \mathbb{Z}, \tau_0, \varphi_j - 2\mathbb{Z}\omega_j\pi)$$

with

$$(5.57) \quad \omega_j := \frac{\Theta_1(2\pi) - \Theta_j(2\pi)}{2\pi} \quad \forall j \in [2, n],$$

such that ζ_y in \mathbb{T}_ξ writes

$$(5.58) \quad \zeta_y = \begin{pmatrix} V_1(\tau) \\ e^{i\varphi_j} V_j(\tau) \end{pmatrix}$$

where $\tau := 2k\pi + \tau_0$. As soon as $n > 2$ there is no longer correspondence between the fiber of P and a sphere, however in analogy with (5.44) we write

$$(5.59) \quad V_j = \sqrt{\frac{R_j}{2}} e^{i\theta_j^0}$$

where $R_j \in \mathbb{R}^+$ and $\theta_j^0 \in [0, 2\pi]$. The spectral distance on a given fiber has a simple expression. To fix notation we consider the fiber over x and we identify ξ_x to the $n + 1$ -tuple $(0, 0, \dots, 0)$.

PROPOSITION 5.10. [46, Prop. 5.2] *Given a pure state $\zeta_x = (k, 0, \varphi_j) \in \mathbb{T}_\xi$, either ζ_x does not belong to the connected component \mathbb{U}_ξ and $d(\xi_x, \zeta_x) = +\infty$ or $\zeta_x \in \mathbb{U}_\xi$ and*

$$(5.60) \quad d_A(\xi_x, \zeta_x) = \pi \operatorname{Tr} |S_k|$$

where $|S_k| = \sqrt{S_k^* S_k}$ and S_k is the matrix with components

$$(5.61) \quad S_{ij}^k := \sqrt{R_i R_j} \frac{\sin \left(k\pi(\omega_j - \omega_i) + \frac{\varphi_j - \varphi_i}{2} \right)}{\sin \pi(\omega_j - \omega_i)}.$$

In the low dimensional case $n = 2$, the connected component $\operatorname{Con}(\xi_x)$ of the spectral distance is 2-torus \mathbb{T}_ξ (see figure 4), whereas the connected component $\operatorname{Acc}(\xi_x)$ of the horizontal distance is a subset of it. Following (5.56), one parametrizes the S^1 -fiber of \mathbb{T}_ξ over x - denoted S_x - by

$$(5.62) \quad \Xi := 2k\omega\pi + \varphi \pmod{2\pi} \quad k \in \mathbb{N}, 0 \leq \varphi \leq 2\pi.$$

In this parametrization, ξ_x has coordinate $\Xi = 0$.

PROPOSITION 5.11. [44, §6.2] *For $\zeta_x \in S_x$ with coordinate $\Xi \in [0, 2\pi]$, one has*

$$(5.63) \quad d_A(\xi_x, \zeta_x) = \frac{2\pi R}{|\sin \omega\pi|} \sin \frac{\Xi}{2}.$$

It is quite interesting to note that for those points on the fiber which are accessible from ξ_x , namely $\xi_x^k = (k, 0, 0)$ or equivalently

$$(5.64) \quad \Xi = \Xi_k := 2k\omega\pi,$$

the Carnot-Carathéodory distance is $d_h(0, \Xi_k) = 2k\pi$. Hence, as soon as ω is irrational, one can find close to ξ_x in the Euclidean topology of S_x some ξ_x^k which are arbitrarily Carnot-Carathéodory-far from ξ_x . In other terms, d_h destroys the S^1 structure of the fiber. On the contrary the spectral distance preserves it since $d_A(\xi_x, \xi_y)$ is proportional to the chord distance on S^1 (see figure 6).

The chord distance already appeared in Prop. 3.9 for the finite dimensional spectral triple with $\mathcal{A} = M_2(\mathbb{C})$. This suggests that the distance in an almost commutative geometry with a gauge fluctuation may be retrieved as a spectral distance associated to the finite dimensional algebra \mathcal{A}_F only, as this happens in the product of spectral triples with the *non-fluctuated* Dirac operator (see Prop. 4.3).

Proposition 5.11 gives another example where the space of pure states equipped with the spectral distance is not a path metric space. We come back to this point in §7.1. Let us just notice that the chord distance on S^1 is smooth at the cut-locus, contrary to the Euclidean distance on the circle (cf Figure 6). A possibility to make the fiber S_x equipped with d_A a path metric space, is to interpret the chord distance as the arc length on a cardioid. We elaborated on this point in [45].




FIGURE 6. From left to right: $d_h(0, \Xi_k)$, $d_A(0, \Xi_k)$ and $d_E(0, \Xi_k)$. Vertical unit is $\frac{\pi R}{|\sin \omega \pi|}$, horizontal unit is π .

6. Compact operators: Moyal plane vs quantum space

After finite dimensional spectral triples in section 3 and almost commutative geometries in sections 4 and 5, we consider the spectral distance on a truly noncommutative algebra (that is: an infinite dimensional algebra with finite dimensional center), namely the C^* - algebra \mathcal{K} of compact operators. We first view it as the norm closure of the algebra of the Moyal plane (\mathcal{S}, \star) , in order to have a natural candidate as a Dirac operator. The main results on the spectral distance on the Moyal plane coming from [6] and [43] are exposed in §6.1.

Then we view \mathcal{K} as the algebra of functions on a quantum spacetime. This interpretation is common to various models, motivated by quantum gravity. Following the idea that at very small scale spacetime itself becomes quantum, hence the length might be quantized, a quantum length operator is defined in these models. It has a priori no links with the spectral distance but we showed [42] that between certain classes of states, including coherent states, the quantum length and the spectral distance actually capture the same metric information. This is the object of §6.2.

6.1. Spectral distance on the Moyal plane. The spectral triple of the Moyal plane [30] is

$$(6.1) \quad \mathcal{A} = (\mathcal{S}, \star), \quad \mathcal{H} = L^2(\mathbb{R}^2) \otimes \mathbb{C}^2, \quad D = -i\sigma^\mu \nabla_\mu,$$

where \mathcal{A} is a noncommutative deformation of the algebra of Schwartz functions on the plane,

$$(6.2) \quad (f \star g)(x) := \frac{1}{(\pi\theta)^2} \int d^2y d^2z f(x+y)g(x+z)e^{-2iy\Theta^{-1}z} \quad \forall f, g \in \mathcal{A},$$

induced by a symplectic form on \mathbb{R}^2 ,

$$(6.3) \quad \Theta := \theta \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \theta \in (0, 1).$$

D is the usual Dirac operator of the plane and $f \in \mathcal{A}$ acts $L_f \otimes \mathbb{I}_2$ where

$$(6.4) \quad (L_f \psi)(x) = (f \star \psi)(x) \quad \forall x \in \mathbb{R}^2$$

denotes the left-star multiplication on $L^2(\mathbb{R}^2)$.

The C^* -closure of \mathcal{A} is the algebra of compact operators. Thus the set of pure states of \mathcal{A} is the set of vector states in the irreducible faithful representation (the Schrödinger representation) of \mathcal{A} on $L^2(\mathbb{R})$, where any element of \mathcal{A} appears as an infinite dimensional matrix with rapidly decreasing coefficients. The eigenstates of the harmonic oscillator, that is the Hermite functions h_m , form a basis of $L^2(\mathbb{R})$. We denote ω_m the associated vector state

$$(6.5) \quad \omega_m = \langle h_m, \cdot h_m \rangle.$$

Another class of interesting states are the coherent states

$$(6.6) \quad \alpha_\kappa \omega_0 := \omega_0 \circ \alpha_\kappa, \quad \kappa \in \mathbb{R}^2,$$

which are the lift to the ground state ω_0 of the action of \mathbb{R}^2 on \mathcal{A} by translation, that is

$$(6.7) \quad (\alpha_\kappa f)(x) = f(x + \kappa).$$

More generally, for any state φ of \mathcal{A} we denote its translated by $\kappa \in \mathbb{R}^2$ as

$$(6.8) \quad \varphi_\kappa = \varphi \circ \alpha_\kappa.$$

The main results on the spectral distance on the Moyal plane are summarized in the following proposition.

PROPOSITION 6.1. *i. [43, Theo. 3.7] The spectral distance between any state $\varphi \in \mathcal{S}(\mathcal{A})$ of the Moyal algebra and any of its κ -translated, $\kappa \in \mathbb{C}$, is precisely the amplitude of translation*

$$(6.9) \quad d(\varphi, \alpha_\kappa \varphi) = |\kappa|.$$

ii. [6, 43] The spectral distance on the Moyal plane takes all possible values in $[0, \infty]$. In particular there are pure states at infinite distance from one another.

iii. [6, Prop. 3.6] The distance between the eigenstates ω_m of the harmonic oscillator is additive: for $m \leq n$,

$$(6.10) \quad d(\omega_m, \omega_n) = \sqrt{\frac{\theta}{2}} \sum_{k=m+1}^n \frac{1}{\sqrt{k}}.$$

Point ii. follows from i. and the fact that there exist (pure) states at infinite distance from one another (see [6, Prop. 3.10] and [7]). This an important difference with the commutative $\theta = 0$ case, where the distance between any pure states - i.e. any point - is as large as one wants, but remains finite.

The distance in the Moyal plane computed with the “harmonic Dirac operator” that appears in various physical models of quantum spacetime [32] has been investigated in [60]. One finds a multiple of the distance computed with the usual Dirac operator in (6.1), yielding a formalization of homothetic spectral triples.

6.2. Minimal length on quantum spacetime. At the Planck scale λ_P , the general relativistic picture of spacetime as a smooth manifold \mathcal{M} is expected to loose any operational meaning, due to the impossibility of simultaneously measuring with arbitrary accuracy the four spacetime coordinates x_μ . This comes as a consequence of the *principle of gravitational stability against localization* [25, 26], which states that to prevent the formation of black-hole during an arbitrarily accurate localization process, one postulates a non-zero minimal uncertainty in the *simultaneous* measurement of *all* coordinates of space-time. A way to implement

these uncertainty relations is to view the coordinates in a chart U of \mathcal{M} no more as functions $x \in U \subset \mathcal{M} \mapsto x_\mu \in \mathbb{R}$, but as quantum operators q_μ satisfying non trivial commutation relations,

$$(6.11) \quad [q_\mu, q_\nu] = i\lambda_P^2 Q_{\mu\nu},$$

where the $Q_{\mu\nu}$'s are operators whose properties depend on the model. In particular in [27, 2], the commutators $Q_{\mu\nu}$'s are central operators with selfadjoint closure, covariant under the action of the Poincaré group. In this case, the quantum coordinates Q_μ are affiliated to the algebra of compact operators \mathcal{K} , in the same way as in the commutative case the coordinates x_μ does not belong to $C_0(\mathbb{R}^n)$ but are affiliated to it. From this perspective, \mathcal{K} plays the role of algebra of noncommutative functions on the quantum plane.

A natural candidate to capture the metric information of the quantum space (6.11) is the length operator

$$(6.12) \quad L = \sqrt{\sum_{\mu} (dq_{\mu})^2} \quad \text{where} \quad dq_{\mu} := q_{\mu} \otimes 1 - 1 \otimes q_{\mu}.$$

The idea is that the minimum l_P of the spectrum of L represents the minimum value of the measurable length on a quantum space [2, 1].

Taking advantage of the double role of the algebra of compact operators as the closure of the Moyal algebra and as the algebra of functions on quantum spacetime, it is natural to wonder whether the metric information captured by the length operator L is related to the metric information captured by the spectral distance d in the Moyal plane. To this aim, given two states φ, φ' of \mathcal{K} , one needs to associate a number with the length operator L , that could then be compared with $d(\varphi, \varphi')$. Assuming that φ, φ' are in the domain of the operators Q_μ (that is $\varphi(Q_\mu)$ and $\varphi'(Q_\mu)$ make sense), the most natural choice is to consider the evaluation of the separable state $\varphi \otimes \varphi'$ on the operator L ,

$$(6.13) \quad d_L(\varphi, \varphi') := (\varphi \otimes \varphi')(L).$$

We call it the *quantum length* of the state $\varphi \otimes \varphi'$. However, to avoid the difficulties in taking the square root of an operator, it is more convenient to work with the *quantum square length*

$$(6.14) \quad \sqrt{d_L^2(\varphi, \varphi')} := \sqrt{\varphi \otimes \varphi'(L^2)}.$$

In the commutative case $q_\mu = x_\mu$, one has

$$(6.15) \quad d_L(\delta_x, \delta_y) = \sqrt{d_{L^2}(\delta_x, \delta_y)} = d(\delta_x, \delta_y) = d_{\text{geo}}(x, y).$$

All the tools introduced so far to measure a distance (spectral distance, spectrum of a length operator or of its square) all coincide with the geodesic distance. In the noncommutative case, there is no reason that the three quantities on the l.h.s. of the equation above remain equal. In particular while the spectral distance d is actually a distance between states, so that $d(\varphi, \varphi) = 0$, there is no reason that $d_{L^2}(\varphi, \varphi)$ and $d_L(\varphi, \varphi)$ vanish.

This can be checked on the set of *generalized coherent states* (see (6.6))

$$(6.16) \quad \mathcal{C} := \{\alpha_\kappa \omega_m, m \in \mathbb{N}, \kappa \in \mathbb{R}^2\}.$$

PROPOSITION 6.2. [42, 2] *The quantum square-length on \mathcal{C} is*

$$(6.17) \quad d_{L^2}(\alpha_\kappa\omega_m, \alpha_{\tilde{\kappa}}\omega_n) = 2E_m + 2E_n + |\kappa - \tilde{\kappa}|^2$$

for any $m, n \in \mathbb{N}$, $\kappa, \tilde{\kappa} \in \mathbb{R}^2$, with

$$(6.18) \quad E_m = \lambda_P^2(m + \frac{1}{2})$$

the n^{th} eigenvalue of the Hamiltonian H of the quantum harmonic oscillator. Hence the quantum square length is invariant by translation. Moreover one has

$$(6.19) \quad d_L(\alpha_\kappa\omega_m, \alpha_{\tilde{\kappa}}\omega_n) \leq \sqrt{d_{L^2}(\alpha_\kappa\omega_m, \alpha_{\tilde{\kappa}}\omega_n)}$$

with equality only when $m = n = 0$ and $\kappa = \tilde{\kappa}$.

Identifying the parameter θ in (6.3) with the square λ_P^2 of the Planck length, one has between eigenstates of the harmonic oscillator that

$$(6.20) \quad d(\omega_m, \omega_n) = \frac{\lambda_P}{\sqrt{2}} \sum_{k=m+1}^n \frac{1}{\sqrt{k}} \neq \sqrt{d_{L^2}(\omega_m, \omega_n)} = \sqrt{2E_m + 2E_n},$$

whereas between generalized coherent states

$$(6.21) \quad d(\omega_m, \alpha_\kappa\omega_m) = |\kappa| \neq \sqrt{d_{L^2}(\omega_m, \alpha_\kappa\omega_m)} = \sqrt{4E_m + |\kappa|^2}.$$

To understand the discrepancy between the quantum length and the spectral distance, one should understand how to *turn the quantum length into a true distance* that vanishes on the diagonal $\varphi' = \varphi$, or how to *give a quantum taste to the spectral distance* so that it no longer vanishes on the diagonal. As explained below, the two points of view turn out to be equivalent thanks to Pythagoras theorem of §4.1.

One “quantizes” the spectral distance by doubling the Moyal plane, that is by taking the product in the sense of (4.2) of the spectral triple (6.1) with the two point space of § 3.1, namely

$$(6.22) \quad \mathcal{A}' := \mathcal{A} \otimes \mathbb{C}^2, \quad \mathcal{H}' := \mathcal{H} \otimes \mathbb{C}^2, \quad D' := D \otimes \mathbb{I} + \Gamma \otimes D_F$$

where Γ is a grading of \mathcal{H} and

$$(6.23) \quad D_F := \begin{pmatrix} 0 & \bar{\Lambda} \\ \Lambda & 0 \end{pmatrix} \quad \text{with } \Lambda = \text{const.}$$

Pure states of \mathcal{A}' are pairs

$$(6.24) \quad \omega^i := (\omega, \delta_i), \quad \omega \in \mathcal{P}(\mathcal{A}), \quad \mathcal{P}(\mathbb{C}^2) = \{\delta_1, \delta_2\}.$$

Hence

$$(6.25) \quad \mathcal{P}(\mathcal{A}') \simeq \mathcal{P}(\mathcal{A}) \times \mathcal{P}(\mathcal{A})$$

and the geometry described by the doubled spectral triple $(\mathcal{A}', \mathcal{H}', D')$ is a *two-sheet model* - two copies of the Moyal plane. The associated distance d' is known between translated states $\varphi^i = (\varphi, \delta^i)$, $\varphi_\kappa^j = (\varphi_\kappa, \delta^j)$ non-necessarily localized on the same copy, and is given by Pythagoras theorem.

PROPOSITION 6.3. [43] *For any $\varphi \in \mathcal{S}(\mathcal{A})$ and $\kappa \in \mathbb{R}^2$, one has*

$$(6.26) \quad d'^2(\varphi^1, \varphi_\kappa^2) = d^2(\varphi, \varphi_\kappa) + d_2^2(\delta^1, \delta^2)$$

where d, d_2 are the distances on the Moyal plane and the two point space.

Rather than comparing the quantum length with the spectral distance on a single sheet, the idea is to compare the quantum square-length $\sqrt{d_{L^2}(\varphi, \tilde{\varphi})}$ with the spectral distance in the double-sheeted model $d'(\varphi^1, \tilde{\varphi}^2)$. In particular, for $\tilde{\varphi} = \varphi_\kappa$, one has from Prop. 6.3 and (3.5) that

$$(6.27) \quad d'(\varphi^1, \varphi_\kappa^2) = \sqrt{d_{L^2}(\varphi, \varphi_\kappa)}$$

if and only if on a single sheet

$$(6.28) \quad d(\varphi, \varphi_\kappa) = \sqrt{d_{L^2}(\varphi, \varphi_\kappa) - |\Lambda|^{-2}}.$$

For any $\varphi, \tilde{\varphi}$ in the domain of the length operator L , we thus define the *modified quantum length* as

$$(6.29) \quad d'_L(\varphi, \tilde{\varphi}) := \sqrt{|d_{L^2}(\varphi, \tilde{\varphi}) - \Lambda^{-2}(\varphi, \tilde{\varphi})|}$$

where

$$(6.30) \quad \Lambda^{-2}(\varphi, \tilde{\varphi}) := \sqrt{d_{L^2}(\varphi, \varphi)d_{L^2}(\tilde{\varphi}, \tilde{\varphi})}.$$

This is the correct quantity, built from the length operator L , that should be compared with the spectral distance.

PROPOSITION 6.4. [42] *On the set of generalized coherent states for a fixed $m \in \mathbb{N}$, that is*

$$(6.31) \quad \mathcal{C}(\omega_m) := \{\alpha_\kappa \omega_m, \kappa \in \mathbb{R}^2\},$$

one has

$$(6.32) \quad d_D(\omega, \tilde{\omega}) = d'_L(\omega, \tilde{\omega}) \quad \forall \omega, \tilde{\omega} \in \mathcal{C}(\omega_m).$$

On the set of all generalized coherent states (6.16), d_D coincides with d'_L asymptotically, both in the limit of large translation

$$(6.33) \quad \lim_{\kappa \rightarrow \infty} \frac{d_D(\alpha_\kappa \omega_m, \alpha_{\tilde{\kappa}} \omega_n) - d'_L(\alpha_\kappa \omega_m, \alpha_{\tilde{\kappa}} \omega_n)}{d'_L(\alpha_\kappa \omega_m, \alpha_{\tilde{\kappa}} \omega_n)} = 0, \quad \forall m, n \in \mathbb{N}, \tilde{\kappa} \in \mathbb{C},$$

and for large difference of energy

$$(6.34) \quad \lim_{n \rightarrow 0} \frac{d_D(\alpha_\kappa \omega_m, \alpha_{\tilde{\kappa}} \omega_n) - d'_L(\alpha_\kappa \omega_m, \alpha_{\tilde{\kappa}} \omega_n)}{d'_L(\alpha_\kappa \omega_m, \alpha_{\tilde{\kappa}} \omega_n)} = 0, \quad \forall m \in \mathbb{N}, \kappa, \tilde{\kappa} \in \mathbb{C}.$$

It is quite remarkable that the spectral distance d_D on a single copy of the Moyal plane coincides (exactly on the set of translated of a states, asymptotically on the set of generalized coherent states) with the “natural” quantity d'_L , vanishing on the diagonal, that one can build from the quantum length d_L . The two options “quantizing the spectral distance” by allowing the emergence of a non-zero minimal spectral distance, or “geometrizing the quantum length” by turning it into a true distance are two equivalent procedures.

7. Discussion

In §7.1 and §7.2 we gather several observations coming from the previous examples concerning the following question: what should play the role of points and geodesics in noncommutative geometry? As a concluding remark, we discuss in §7.3 Kantorovich duality in the noncommutative framework.

7.1. Points and geodesics. In the commutative case $\mathcal{A} = C(\mathcal{M})$ for \mathcal{M} a compact manifold, two pure states δ_x, δ_y provide via the GNS construction two inequivalent irreducible representations. This is no longer true in the noncommutative case. For instance all the pure states of $M_n(\mathbb{C})$ yields equivalent irreducible representations. A point of view is to consider that a “point” in noncommutative geometry should be a class of irreducible equivalent representations - that is a class of pure states - rather than a pure state. From this point of view, any spectral triple with algebra $\mathcal{A} = M_n(\mathbb{C})$ describes a one-point space. On the contrary, we argued in §3.3 that the spectral distance gives a non-trivial structure to the set of pure states of $M_n(\mathbb{C})$, regardless the unitary equivalence of the representations they induce. We do not see any good reason to wash out this structure by considering only quotients of $\mathcal{P}(\mathcal{A})$ instead of $\mathcal{P}(\mathcal{A})$ entirely.

Furthermore, several facts suggest that the purity of state might not be such a relevant concept regarding the metric aspect of noncommutative geometry. For instance Pythagoras equality holds between pure states δ_x^1, δ_y^2 in the product of a manifold by \mathbb{C}^2 , but in the case of the Moyal plane it holds between translated states $\varphi^1, \varphi_\kappa^2$, pure or not. What is important to pass from Pythagoras inequalities of theorem 4.1 to the equality is not the purity of the states, but the existence of a curve $t \rightarrow \varphi(t) \in \mathcal{S}(\mathcal{A})$ between the two considered states such that

$$(7.1) \quad d(\varphi(s), \varphi(t)) = |s - t| d(\varphi(0), \varphi(1)) \quad \forall s, t \in [0, 1].$$

In case of a manifold, such a curve is provided by the minimal geodesic between $\delta_x = \varphi(0)$ and $\delta_y = \varphi(1)$, which has value in pure states. In case of the Moyal plane, this curve is the orbit of $\varphi = \varphi(0)$ under the translation action of \mathbb{R}^2 , which lies in $\mathcal{P}(\mathcal{A})$ if φ is pure, in $\mathcal{S}(\mathcal{A})$ otherwise.

Other instances where the purity of state does not seem an adequate criteria to characterize a “point” of a noncommutative geometry are the cut-off geometries developed in [20]. There, pure states need to be approximated by non-pure states. Namely, given a spectral triple $(\mathcal{A}, \mathcal{H}, D)$, one truncates the Dirac operator via the adjoint action of a sequence of increasing projections P_N tending to \mathbb{I} ,

$$(7.2) \quad D \rightarrow D_N := P_N D P_N.$$

This has motivations from the spectral action where the Dirac operator is truncated by a cut-off energy. In case $(\mathcal{A}, \mathcal{H}, D)$ is the usual spectral triple of a manifold, substituting in the spectral distance formula the semi-norm L_D by

$$(7.3) \quad L_N(a) := ||| [D_N, a] |||$$

yields a distance between any pure states δ_x, δ_y which is infinite as soon as D_N has finite rank. To make it finite, one should truncate the pure states as well. In case $\mathcal{M} = S^1$, an explicit truncation is given by the Fejer transform of rank N , yielding non pure states of $C^\infty(\mathcal{M})$ (see [20, §5] for details).

Related to the problem of determining what points are in a noncommutative context, is the question of what should play the role of a geodesic. From a purely

metric point of view, one may take as a definition of (minimal) geodesic between two states φ, φ' a curve like (7.1) with $\varphi(0) = \varphi$, $\varphi(1) = \varphi'$. In the case of a manifold

$$(7.4) \quad \mathcal{A} = C^\infty(\mathcal{M}), \quad \varphi = \delta_x, \varphi' = \delta_y$$

there are two such curves: the usual geodesic between x and y (with t the proper length) which lies completely in $\mathcal{P}(\mathcal{A})$, and the convex combination

$$(7.5) \quad \varphi(t) = t\delta_x + (1-t)\delta_y$$

which lies in non-pure states. In case of $\mathcal{A} = M_2(\mathbb{C})$, the distance on the 2-sphere $\mathcal{P}(\mathcal{A})$ is the Euclidean distance in the 3-ball $\mathcal{S}(\mathcal{A})$, hence any curve (7.1) between two pure states necessarily goes through non-pure states. The same is true for the two point space of §3.1, the distance between the two sheets of the standard model in §4.3, and the distance on a fiber S_x with the gauge fluctuated Dirac operator in §5.6. In other terms, in all these examples the space of pure states $\mathcal{P}(\mathcal{A})$ equipped with the spectral distance is not a path metric space. This forbids to take (7.1) as a definition of a geodesic, at least as long as one imposes that the later must be a curve of pure states.

If one allows non-pure states, then a geodesic in the sense of (7.1) always exists and is given by (7.5). This does not seem a very operative definition of a geodesic; it simply shows that by the very definition of states as convex combinations of pure states, then the space of states $\mathcal{S}(\mathcal{A})$ equipped with the spectral distance is trivially always path metric. A more interesting question could be the following: is the commutative case $\mathcal{A} = C^\infty(\mathcal{M})$ the only example where the space of pure states equipped with the spectral distance is path metric ?

More understanding on these questions may come from optimal transport. As pointed out by a referee of an early version of this text, the question in this context is whether the curve of measures (a curve of states, in our terminology) is produced by the underlying measure on curves. A discussion on that matter, for the Wasserstein distance of order p though, can be found in [38], see also [59]. For the distance of order 1, one should see the appendix of [53].

7.2. Optimal elements and geodesics. Another point of view [42, 47] on the question of geodesics in noncommutative geometry could be to define a geodesic in a dual way, that is to find a substitute of the geodesic in the notion of *optimal element* introduced in definition 2.1. This makes sense because in the commutative case, the commutator norm condition

$$(7.6) \quad \|[\partial, f]\| = \sup_{x \in \mathcal{M}} \left\| \nabla f|_x \right\|_{T_x \mathcal{M}} = 1$$

characterizes the optimal element between δ_x and δ_y locally, in the sense that the constraint is carried by the gradient of f . The geodesics through x are retrieved as the curves tangent to gradient of the optimal element $f = d_{\text{geo}}(x, \cdot)$. In this sense, computing the spectral distance - that is finding an optimal element - amounts to solving the equation of the geodesics:

- eq. (7.6) plays the role of the geodesic equation;
- the optimal element $d_{\text{geo}}(x, \cdot)$ fully characterizes the geodesics through x ;
- the valuation of the optimal element on $\delta_x - \delta_y$ gives the integration of the line element on a minimal geodesic between x and y .

At the moment there is no clear translation of the above points in a non-commutative context. However, focusing on the optimal element yields interesting interpretations of the results on the Moyal plane of section 6. Recall that in the commutative case, as observed in (6.15), both the quantum length and the spectral distance coincide with the geodesic distance; and this is the same function

$$(7.7) \quad l(x_\mu) := \sqrt{\sum_{\mu} x_{\mu}^2}$$

which yields both the optimal element between two pure states $\delta_x, \delta_{\lambda x}$, $\lambda \in \mathbb{R}^+$ and - by the functional calculus - the length operator $L = l(dq_\mu)$ in (6.12). On the contrary, on the Moyal plane the quantum length and the spectral distance no longer coincide, as stressed in (6.20) and Prop. 6.4, so that one could expect the length operator not to be defined by the same function as the optimal element. This is indeed the case. To see it is convenient to work with the creation/annihilation operators

$$(7.8) \quad a := \frac{q_1 + iq_2}{\sqrt{2}}, \quad a^* := \frac{q_1 - iq_2}{\sqrt{2}},$$

as well as with their universal differentials

$$(7.9) \quad da = \frac{1}{\sqrt{2}}(dq_1 + idq_2), \quad da^* = \frac{1}{\sqrt{2}}(dq_1 - idq_2).$$

PROPOSITION 7.1. [42] *The length operator can be equivalently defined as $L = l_i(da)$, with*

$$(7.10) \quad l_1(z) := \sqrt{z\bar{z} + z\bar{z}} \quad \text{or} \quad l_2(z) := \sqrt{2(z\bar{z} - \lambda_P^2)} \quad \text{or} \quad l_3(z) := \sqrt{2(\bar{z}z + \lambda_P^2)}.$$

The optimal element between any two eigenstates of the Hamiltonian of the quantum harmonic oscillator is - up to regularization at infinity - the \star -action of the function l_0 , defined as the solution of

$$(7.11) \quad (\partial_z l_0 \star z) \star (\partial_z l_0 \star z)^* = \frac{1}{2} z^* \star z.$$

Neither $l_1(a)$ nor $l_2(a)$ or $l_3(a)$ are optimal elements between eigenstates.

In a similar way, the spectral distance between translated states φ, φ_κ being the amplitude of translation $|\kappa|$ both in the commutative and the noncommutative cases, one could expect the respective optimal elements to be related. And this is indeed the same function

$$(7.12) \quad l_\kappa(z) = \frac{ze^{-i\Xi} + \bar{z}e^{i\Xi}}{\sqrt{2}} \quad \text{with} \quad \Xi := \text{Arg } \kappa,$$

which yields the optimal element (up to regularization at infinity) both on the Euclidean plane (through the pointwise action of l_κ) and the Moyal plane (through its \star -action). For the latter, this has been shown in [43, Theo. III.9], for the former in [22, Prop. 3.2]). It is quite remarkable that the same function l_κ gives an optimal element between translated states, regardless of the commutativity of the algebra.

Let us now compare the optimal elements l_κ for translated states and l_0 for eigenstates of the harmonic oscillator. Modulo regularization at infinity, the latter

is characterized as a solution of [6, Prop. 3.7]

$$(7.13) \quad [\not\partial, L_{l_0}] = -i \begin{pmatrix} 0 & S^* \\ S & 0 \end{pmatrix}$$

where L_{l_0} denotes the \star -multiplication by l_0 defined by (7.11), while S is the shift operator (eq. (7.11) actually follows from (7.13)). In analogy with the commutative case where $[\not\partial, f] = (\not\partial f)$, we interpret $[\not\partial, L_{l_0}]$ as the derivative of the optimal element l_0 . The presence of the shift operator in this derivative suggests that the “geodesic” is somehow non smooth. A similar interpretation follows from the observation that the spectral distance (6.10)

$$(7.14) \quad d(\omega_m, \omega_n) = \lambda_P \sum_{k=m+1}^n \frac{1}{\sqrt{2k}}$$

is the middle Riemann sum approximation of the modified quantum length (6.29)

$$(7.15) \quad d'_L(\omega_m, \omega_n) = \lambda_P (\sqrt{2n+1} - \sqrt{2m+1}) = \lambda_P \int_{m+\frac{1}{2}}^{n+\frac{1}{2}} \frac{1}{\sqrt{2k}} dk.$$

In [43, 42] we interpret this result saying that the spectral distance and the quantum length are the integration of the same quantum line element

$$(7.16) \quad \lambda_P \frac{1}{\sqrt{2k}} dk$$

but along two distinct geodesics: a continuous one for the quantum length (7.15), a discrete one for the spectral distance (7.14).

Between translated states, the optimal element l_κ satisfies an equation similar to (7.13),

$$(7.17) \quad [\not\partial, L_{l_\kappa}] = -i \begin{pmatrix} 0 & e^{i\Xi} \\ e^{-i\Xi} & 0 \end{pmatrix}$$

where the shift is substituted with a term proportional to the identity. This indicates that the geodesic is “smooth”, in agreement with the analysis developed below (7.12).

In the same vein, one has

$$(7.18) \quad [\not\partial, L_{l_0}]^* [\not\partial, L_{l_0}] = \mathbb{I} - e_0$$

where e_0 is the projection on the ground state h_0 , while

$$(7.19) \quad [\not\partial, L_{l_\kappa}]^* [\not\partial, L_{l_\kappa}] = \mathbb{I}.$$

Eq. (7.19) indicates that the derivative of the optimal element l_κ is a unitary operator, whereas the derivative of l_0 is not. This comes from the fact that the set of eigenstates of the harmonic oscillator - identified to \mathbb{N} - is not a group, unlike the set of translated states. So the shift S acting on $l^2(\mathbb{N})$ is not a unitary operator.

7.3. Kantorovich duality in noncommutative geometry ? The formula (2.7) of the spectral distance is a way to export to the noncommutative setting the usual notion of Riemannian geodesic distance. Notice the change of point of view: the distance is no longer the infimum of a geometrical object (i.e. the length of the paths between points), but the supremum of an algebraic quantity (the difference of the valuation of two states).

A natural question is whether one loses any trace of the distance-as-an-infimum by passing to the noncommutative side. More specifically, is there some “noncommutative Kantorovich duality” allowing to view the spectral distance as the minimization of some “noncommutative cost” ?

<u>distance as a supremum:</u>	d_{ϑ} commutative case	→	d_D noncommutative case
	↑		
Kantorovich duality:	$d_{\vartheta} = W$		$d_D = W_D?$
	↓		↓
<u>distance as an infimum:</u>	W with cost d_{geo}		noncommutative cost ?

In this diagram, d_{ϑ} and d_D denote the spectral distances computed with the seminorms $\|[\vartheta, \cdot]\|$ and $\|[D, \cdot]\|$, while W is the Wasserstein distance and W_D its putative noncommutative generalization.

In the commutative case, the cost function is retrieved as the Monge-Kantorovich distance between pure states of $C_0(\mathcal{M})$. So in the noncommutative case, if the spectral distance were to coincide with some “Monge-Kantorovich”-like distance W_D on $\mathcal{S}(\mathcal{A})$, then the associated cost should be the spectral distance on the pure state space $\mathcal{P}(\mathcal{A})$. So given a spectral triple $(\mathcal{A}, \mathcal{H}, D)$, we aim at defining a “Monge-Kantorovich”-like distance W_D on the state space $\mathcal{S}(\mathcal{A})$, taking as a cost function the spectral distance d_D on the pure state space $\mathcal{P}(\mathcal{A})$. A first idea is to mimic formula (2.2) with $\mathcal{X} = \mathcal{P}(\mathcal{A})$, that is

$$(7.20) \quad W(\mu_1, \mu_2) = \inf_{\rho} \int_{\mathcal{P}(\mathcal{A}) \times \mathcal{P}(\mathcal{A})} d\rho \quad d_D(\omega, \tilde{\omega})$$

where μ_1, μ_2 are probability measures on $\mathcal{P}(\mathcal{A})$, $\omega, \tilde{\omega}$ are generic elements of $\mathcal{P}(\mathcal{A})$ and the infimum is on the measures ρ on $\mathcal{P}(\mathcal{A}) \times \mathcal{P}(\mathcal{A})$ with marginals μ_1, μ_2 . For this to make sense as a distance on $\mathcal{S}(\mathcal{A})$, we should restrict to states $\varphi \in \mathcal{S}(\mathcal{A})$ that are given by a probability measure on $\mathcal{P}(\mathcal{A})$. This is possible (at least) when \mathcal{A} is separable and unital: $\mathcal{S}(\mathcal{A})$ is then metrizable [5, §4.1.4] so that by Choquet theorem any state $\varphi \in \mathcal{S}(\mathcal{A})$ is given by a probability measure $\mu \in \text{Prob}(\mathcal{P}(\mathcal{A}))$. One should be careful however that the correspondence is not 1 to 1: $\mathcal{S}(\mathcal{A}) \rightarrow \text{Prob}(\mathcal{P}(\mathcal{A}))$ is injective, but two distinct probability measures μ_1, μ_2 may yield the same state φ . This is because \mathcal{A} is *not* an algebra of continuous functions on $\mathcal{P}(\mathcal{A})$ (otherwise \mathcal{A} would be commutative). Thus W_D that we are looking for should not be a distance on $\text{Prob}(\mathcal{P}(\mathcal{A}))$, but on a quotient of it, precisely given by $\mathcal{S}(\mathcal{A})$. This forbids to define W_D by formula (7.20), since by construction the latter is a distance on $\text{Prob}(\mathcal{P}(\mathcal{A}))$.

A possibility is to consider the infimum

$$(7.21) \quad \inf_{\mu_1, \mu_2} W(\mu_1, \mu_2)$$

on all the probability measures $\mu_1, \mu_2 \in \text{Prob}(\mathcal{P}(\mathcal{A}))$ such that for any $a \in \mathcal{A}$ one has

$$(7.22) \quad \varphi_1(a) = \int_{\mathcal{P}(\mathcal{A})} \omega(a) d\mu_1(\omega), \quad \varphi_2(a) = \int_{\mathcal{P}(\mathcal{A})} \omega(a) d\mu_2(\omega)$$

for two given states φ_1, φ_2 . However it is not yet clear that (7.21) is a distance on $\mathcal{S}(\mathcal{A})$.

In [41], we explored another way, consisting in viewing \mathcal{A} as an “noncommutative algebra of functions” on $\mathcal{P}(\mathcal{A})$,

$$(7.23) \quad a(\omega) := \omega(a) \quad \forall \omega \in \mathcal{P}(\mathcal{A}), a \in \mathcal{A};$$

and define the set of “ d_D -Lipschitz noncommutative functions” in analogy with (2.6) as

$$(7.24) \quad \text{Lip}_D(\mathcal{A}) := \{a \in \mathcal{A}, |a(\omega_1) - a(\omega_2)| \leq d_D(\omega_1, \omega_2) \quad \forall \omega_1, \omega_2 \in \mathcal{P}(\mathcal{A})\}.$$

By mimicking (2.7) we then defines for any $\varphi_1, \varphi_2 \in \mathcal{S}(\mathcal{A})$

$$(7.25) \quad W_D(\varphi_1, \varphi_2) := \sup_{a \in \text{Lip}_D(\mathcal{A})} |\varphi_1(a) - \varphi_2(a)|.$$

PROPOSITION 7.2. [41, Prop. 3.1] W_D is a distance, possibly infinite, on $\mathcal{S}(\mathcal{A})$. Moreover for any $\varphi_1, \varphi_2 \in \mathcal{S}(\mathcal{A})$,

$$(7.26) \quad d_D(\varphi_1, \varphi_2) \leq W_D(\varphi_1, \varphi_2).$$

The equation above is an equality on the set of convex linear combinations

$$(7.27) \quad \varphi_\lambda := \lambda \omega_1 + (1 - \lambda) \omega_2$$

of any two given pure states ω_1, ω_2 : namely for any $\lambda, \tilde{\lambda} \in [0, 1]$ one has

$$(7.28) \quad d_D(\varphi_\lambda, \varphi_{\tilde{\lambda}}) = |\lambda - \tilde{\lambda}| d_D(\omega_1, \omega_2) = W_D(\varphi_\lambda, \varphi_{\tilde{\lambda}}).$$

The difference between W_D and d_D is entirely contained in the difference between the D -Lipschitz ball (2.8) and $\text{Lip}_D(\mathcal{A})$ defined in (7.24). In the commutative case $\mathcal{A} = C_0(\mathcal{M})$, these two notions of Lipschitz functions coincide with the usual one, so that $d_\emptyset = W_\emptyset$. In the noncommutative case, they coincide on some easy low dimensional examples, like for $\mathcal{A} = M_2(\mathbb{C})$, but there are indications that this is not true in general [54, §7].

To conclude, let us mention another direction of research still largely unexplored: generalizing to the noncommutative realm the Wasserstein distance W_p of order $p \geq 2$ (2.4). The only attempt we are aware of is that of [62], where one proposes a noncommutative version of W_p based on the posets of commutative sub-algebras of a noncommutative C^* -algebra.

Notations

Given $z \in \mathbb{C}$, we denote \bar{z} its conjugate, $|z|$ its module and $\Re(z)$ its real part.

Given an involutive algebra \mathcal{A} , the adjoint of an element $a \in \mathcal{A}$ is a^* . A C^* -algebra is an associative and involutive algebra \mathcal{A} , equipped with a norm $\|\cdot\|$ in which it is complete, and such that for any $a \in \mathcal{A}$ one has

$$(7.29) \quad \|a^*a\| = \|a\|^2.$$

It is *unital* if it contains a unit, that is an element $\mathbf{1} \in \mathcal{A}$ such that

$$(7.30) \quad \mathbf{1}a = a\mathbf{1} = a \quad \forall a \in \mathcal{A}.$$

Most of the time we identify an element a of the algebra \mathcal{A} with its representation $\pi(a)$ as bounded operator on some Hilbert space \mathcal{H} . Unless otherwise specified, representations are always faithful and non-degenerate. In particular, if \mathcal{A} is unital with unit $\mathbf{1}$, this guarantees that $\pi(\mathbf{1})$ is the identity \mathbb{I} of $\mathcal{B}(\mathcal{H})$.

The adjoint action of a unitary $u \in \mathcal{A}$ is $(Adu)(a) := uau^*$, for any $a \in \mathcal{A}$.

We denote by $\mathcal{S}(\mathcal{A})$ and $\mathcal{P}(\mathcal{A})$ the space of states and of pure states of \mathcal{A} . We usually denote a state by φ , with suitable decorations $\varphi', \varphi_0, \varphi_1, \dots$ if needed. A faithful state is a state φ such state $\varphi(a^*a) = 0$ iff $a = 0$. Pure states are usually denoted by ω in the noncommutative case, and δ in the commutative case, with suitable decorations.

Given two operators A, B acting on an Hilbert space \mathcal{H} , the bracket $[A, B] = AB - BA$ is their commutator. $\mathcal{B}(\mathcal{H})$ denotes the space of bounded operators on \mathcal{H} , and \mathbb{I} is the identity operator. Unless otherwise specified, the norm $\|\cdot\|$ is the operator norm coming from the action on \mathcal{H} , that is

$$(7.31) \quad \|A\| = \sup_{\psi \in \mathcal{H}} \frac{\|A\psi\|_{\mathcal{H}}}{\|\psi\|_{\mathcal{H}}}$$

where

$$(7.32) \quad \|\psi\|_{\mathcal{H}} = \sqrt{\langle \psi, \psi \rangle}$$

is the L^2 -norm on \mathcal{H} , with $\langle \cdot, \cdot \rangle$ the inner product on \mathcal{H} . We omit the index \mathcal{H} and it should be clear from the context if one deals with the operator or the L^2 -norm.

A vector in \mathcal{H} is usually denoted by the greek letter ψ, ζ or ξ . Its components in a given orthonormal basis are the complex numbers ψ_i, ζ_i, ξ_i with $i = 1, \dots, \dim \mathcal{H}$. The dual vector is $\bar{\psi}$, with components $\bar{\psi}_i$. The canonical basis of \mathbb{C}^N is $\{e_{ij}\}_{i,j=1,\dots,N}$, that is e_{ij} is the matrix with null entries, except 1 at the i^{th} line, j^{th} column.

A spectral triple $(\mathcal{A}, \mathcal{H}, D)$ is the datum of a (non necessarily commutative) involutive algebra \mathcal{A} , acting faithfully on an Hilbert space \mathcal{H} via a representation π , together with a (non necessarily bounded) operator D on \mathcal{H} such that $D - \lambda\mathbb{I}$ is compact for any λ in the resolvent set of D , and $[D, \pi(a)]$ is a bounded operator for any $a \in \mathcal{A}$. A spectral triple is unital when the algebra \mathcal{A} is unital and the representation π is non-degenerate. It is graded if there exists a grading Γ of \mathcal{H} (that is $\Gamma = \Gamma^*$ and $\Gamma^2 = \mathbb{I}$) which commutes with any $\pi(a)$ and anticommutes with D . The D -Lipschitz ball of \mathcal{A} is

$$(7.33) \quad L_D(\mathcal{A}) := \{a \in \mathcal{A}, \|[D, a]\| \leq 1\}.$$

D denotes the (generalized) Dirac operator of an arbitrary spectral triple. \not{D} is the usual Dirac operator of a spin manifold. γ^μ are the Dirac matrices, σ^μ the Pauli matrices.

We call “distance” a function that verifies all the usual properties of a distance, except that we do not assume it is necessarily finite. d is the spectral distance (1.1), d_A the fluctuated distance defined in § 5, d_h the horizontal distance in sub-Riemannian geometry.

A path metric space is a metric space (\mathcal{X}, \tilde{d}) such that between any two points $x, y \in \mathcal{X}$ there exists a continuous curve $c : [0, 1] \rightarrow \mathcal{X}$ with $c(0) = x, c(1) = y$ and such that

$$(7.34) \quad \tilde{d}(c(s), c(t)) = |s - t| \tilde{d}(x, y) \quad \forall s, t \in [0, 1].$$

The Lipschitz norm of a function f on a Riemannian manifold \mathcal{M} with geodesic distance d_{geo} is

$$(7.35) \quad \|f\|_{\text{Lip}} := \sup_{x, y \in \mathcal{M}} \frac{|f(x) - f(y)|}{d_{\text{geo}}(x, y)}.$$

The algebra of n -dimensional complex matrices is $M_n(\mathbb{C})$. The algebra of quaternions is \mathbb{H} .

References

- [1] Giovanni Amelino-Camelia, Giulia Gubitosi, and Flavio Mercati, *Discreteness of area in non-commutative space*, Phys. Lett. B **676** (2009), 180–83.
- [2] D. Bahns, S. Doplicher, K. Fredenhagen, and G. Piacitelli, *Quantum geometry on quantum spacetime: distance, area and volume operators*, Commun. Math. Phys. **308** (2011), 567–589.
- [3] J. V. Bellissard, M. Marcolli, and K. Reihani, *Dynamical systems on spectral metric spaces*, arXiv:1008.4617v1 [math.OA] (2010).
- [4] G. Bimonte, F. Lizzi, and G. Sparano, *Distances on a lattice from noncommutative geometry*, Phys. Lett. B **341** (1994), 139–146.
- [5] O. Bratteli and D. W. Robinson, *Operator algebras and quantum statistical mechanics 1*, Springer, 1987.
- [6] Eric Cagnache, Francesco d’Andrea, Pierre Martinetti, and Jean-Christophe Wallet, *The spectral distance on Moyal plane*, J. Geom. Phys. **61** (2011), 1881–1897.
- [7] Eric Cagnache and Jean-Christophe Wallet, *Spectral distances: Results for moyal plane and noncommutative torus*, SIGMA **6** (2010), no. 026, 17 pages.
- [8] A. H. Chamseddine and A. Connes, *The spectral action principle*, Commun. Math. Phys. **186** (1996), 737–750.
- [9] A. H. Chamseddine, A. Connes, and M. Marcolli, *Gravity and the standard model with neutrino mixing*, Adv. Theor. Math. Phys. **11** (2007), 991–1089.
- [10] A. H. Chamseddine, A. Connes, and Walter van Suijlekom, *Beyond the spectral standard model: emergence of Pati-Salam unification*, JHEP **11** (2013), 132.
- [11] E. Christensen and C. Ivan, *Spectral triples for C^* -algebras and metrics on the cantor set*, J. Operator Theory **56** (2006), no. 1, 17–46.
- [12] Erik Christensen, Cristina Ivan, and Elmar Schrohe, *Spectral triples and the geometry of fractals*, Journal of Noncommutative Geometry (2011).
- [13] A. Connes, *Compact metric spaces, Fredholm modules, and hyperfiniteness*, Ergod. Th. & Dynam. Sys. **9** (1989), 207–220.
- [14] ———, *Gravity coupled with matter and the foundations of noncommutative geometry*, Commun. Math. Phys. **182** (1996), 155–176.
- [15] A. Connes and J. Lott, *The metric aspect of noncommutative geometry*, Nato ASI series B Physics **295** (1992), 53–93.
- [16] Alain Connes, *On the spectral characterization of manifolds*, J. Noncom. Geom. **7** (2013), no. 1, 1–82.

- [17] D. Testard D. Kastler, *Quantum forms of tensor products*, Commun. Math. Phys. **155** (1993), 135–142.
- [18] J. Dai and X.C. Song, *Pythagoras' theorem on a 2d-lattice from a "natural" Dirac operator and Connes' distance formula*, J. Phys. A **34** (2001), 5571–5582.
- [19] F. D'Andrea, *Pythagoras theorem in noncommutative geometry*, to be published in Comtemp. Math. arXiv 1507.08773 [math-ph] **arXiv:** (2015).
- [20] F. D'Andrea, F. Lizzi, and P. Martinetti, *Spectral geometry with a cut-off: topological and metric aspects*, J. Geom. Phys. **82** (2014), 18–45.
- [21] F. D'Andrea and P. Martinetti, *On Pythagoras theorem for products of spectral triples*, Lett. Math. Phys. **103** (2013), 469–492.
- [22] Francesco D'Andrea and Pierre Martinetti, *A view on optimal transport from noncommutative geometry*, SIGMA **6** (2010), no. 057, 24 pages.
- [23] A. Devastato, *Noncommutative geometry, grand symmetry and twisted spectral triple*, J. Phys.: Conf. Ser. **634** (2015), 012008.
- [24] A. Devastato, F. Lizzi, and P. Martinetti, *Grand Symmetry, Spectral Action and the Higgs mass*, JHEP **01** (2014), 042.
- [25] S. Doplicher, *Spacetime and fields, a quantum texture*, Proceedings 37th Karpacz Winter School of Theo. Physics (2001), 204–213.
- [26] ———, *Quantum field theory on quantum spacetime*, J. Phys.: Conf. Ser. **53** (2006), 793–798.
- [27] Sergio Doplicher, Klaus Fredenhagen, and John E. Robert, *The quantum structure of spacetime at the Planck scale and quantum fields*, Commun.Math.Phys. **172** (1995), 187–220.
- [28] M. Dubois-Violette, J. Madore, and R. Kerner, *Classical bosons in a noncommutative geometry*, Class. Quantum Grav. **6** (1989), 1709.
- [29] Nicolas Franco, *Global eikonal condition for Lorentzian distance function in noncommutative geometry*, SIGMA **6** (2010), no. 064, 11 pages.
- [30] V. Gayral, J. M. Gracia Bondia, B. Iochum, T. Schücker, and J. C. Varilly, *Moyal planes are spectral triples*, Commun. Math. Phys. **246** (2004), 569–623.
- [31] K. R. Goodearl, *Notes on real and complex c^* -algebras*, Notes on Real and Complex C^* -Algebras, 1982.
- [32] Harald Grosse and R. Wulkenhaar, *8d-spectral triple on 4d-moyal space and the vacuum of noncommutative gauge theory*, preprint arXiv 0709.0095 (2007).
- [33] S. Hasselmann, *Spectral triples on Carnot manifolds*, **arXiv 1404.5494** (2014).
- [34] Bruno Iochum, Thomas Krajewski, and Pierre Martinetti, *Distances in finite spaces from noncommutative geometry*, J. Geom. Phys. **31** (2001), 100–125.
- [35] Richard V. Kadison and John R. Ringrose, *Fundamentals of the theory of operator algebras. Volume I, Elementary theory*, Academic Press, 1983.
- [36] L. V. Kantorovich, *On the transfer of masses*, Dokl. Akad. Nauk. SSSR **37** (1942), 227–229.
- [37] F. Latrémolière, *Quantum metric space and the Gromov-Hausdorff propinquity*, to be published in Contemp. Maths. arXiv 1506.04341v2 (2015).
- [38] S. Lisini, *Characterization of absolutely continuous curves in Wasserstein spaces*, Calculus of variations and partial differential equations **28** (2007), no. 1, 85–120.
- [39] J. Madore, *An introduction to noncommutative differential geometry and its physical applications*, Cambridge University Press, 1995.
- [40] P. Martinetti, *Distances en géométrie non-commutative*, PhD thesis (2001), arXiv:math-ph/0112038v1.
- [41] ———, *Towards a Monge-Kantorovich distance in noncommutative geometry*, Zap. Nauch. Semin. POMI **411** (2013).
- [42] P. Martinetti, F. Mercati, and L. Tomassini, *Minimal length in quantum space and integrations of the line element in noncommutative geometry*, Rev. Math. Phys. **24** (2012), no. 5, 1250010–36 pages.
- [43] P. Martinetti and L. Tomassini, *Noncommutative geometry of the Moyal plane: translation isometries, Connes' distance on coherent states, Pythagoras equality*, Commun. Math. Phys. **323** (2013), no. 1, 107–141.
- [44] Pierre Martinetti, *Carnot-Carathéodory metric and gauge fluctuation in noncommutative geometry*, Commun. Math. Phys. **265** (2006), 585–616.

- [45] ———, *Smoother than a circle or how noncommutative geometry provides the torus with an egocentric metric*, proceedings of Deva intl. conf. on differential geometry and physics (Cluj university press (Roumania), ed.), 2006.
- [46] ———, *Spectral distance on the circle*, J. Func. Anal. **255** (2008), no. 1575-1612.
- [47] Pierre Martinetti and Luca Tomassini, *Length and distance on a quantum space*, Proc. of Sciences **042** (2011).
- [48] Pierre Martinetti and Raimar Wolkenhaar, *Discrete Kaluza-Klein from scalar fluctuations in noncommutative geometry*, J. Math. Phys. **43** (2002), no. 1, 182–204.
- [49] B. Mesland and A. Rennie, *Nonunital spectral triples and metric completeness in unbounded KK-theory*, (2015).
- [50] Richard Montgomery, *A tour of subriemannian geometries, their geodesics and applications*, AMS, 2002.
- [51] Valter Moretti, *Aspects of noncommutative Lorentzian geometry for globally hyperbolic space-times*, Rev. Math. Phys. **15** (2003), 1171–1217.
- [52] J. Netsruev, *Smooth manifolds and observables*, Graduate Texts in Mathematics, vol. 220, Springer - Verlag, 2003.
- [53] E. Paolini and E. Stepanov, *Decomposition of acyclic normal currents in a metric space*, J. Func. Anal. **263** (2012), no. 11, 3358–3990.
- [54] Marc Rieffel, *Compact quantum metric spaces*, Contemporary Mathematics **365** (2003).
- [55] Marc A. Rieffel, *Metrics on states from actions of compact groups*, Documenta Math. **3** (1998), 215–229.
- [56] ———, *Metric on state spaces*, Documenta Math. **4** (1999), 559–600.
- [57] Carlo Rovelli, *Lorentzian Connes distance, spectral graph distance and loop gravity*, arXiv 1408.3260 (2014).
- [58] W. van Suijlekom, *Noncommutative geometry and particle physics*, Springer, 2015.
- [59] Cédric Villani, *Optimal transport: Old and new*, Springer, 2009.
- [60] J.-C. Wallet, *Connes distance by examples: Homothetic spectral metric spaces*, Rev. Math. Phys. **24** (2012), no. 9, 1250027.
- [61] N. Weaver, *Lipschitz algebras*, World Scientific, 1999.
- [62] D. Zaev, *L_p -Wasserstein distances on state and quasi-state spaces of c^* -algebra*, (arxiv 1505.06061).

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA
E-mail address: martinetti@dima.unige.it