

# REPRESENTATION VARIETIES DETECT ESSENTIAL SURFACES

STEFAN FRIEDL, TAKAHIRO KITAYAMA, AND MATTHIAS NAGEL

ABSTRACT. Extending Culler-Shalen theory, Hara and the second author presented a way to construct certain kinds of branched surfaces in a 3-manifold from an ideal point of a curve in the  $SL_n$ -character variety. There exists an essential surface in some 3-manifold known to be not detected in the classical  $SL_2$ -theory. We prove that every connected essential surface in a 3-manifold is given by an ideal point of a rational curve in the  $SL_n$ -character variety for some  $n$ .

## 1. INTRODUCTION

In this paper we study an extension of Culler-Shalen theory for higher-dimensional representations. In their seminal work [CS] Culler and Shalen established a method to construct essential surfaces in a 3-manifold from an ideal point of a curve in the  $SL_2(\mathbb{C})$ -character variety. The method is built on a beautiful combination of the theory of incompressible surfaces in a 3-manifold, the geometry of representation varieties, and Bass-Serre theory [Se1, Se2]. We refer the reader to the exposition [Sh] for literature and related topics on Culler-Shalen theory. Hara and the second author presented an analogous extension of the Culler-Shalen method to the case of higher-dimensional representations [HK]. They showed that certain kinds of branched surfaces (possibly without any branched points) are constructed from an ideal point of a curve in the  $SL_n(\mathbb{C})$ -character variety for a general  $n$ . Such a branched surface corresponds to a nontrivial splitting of the 3-manifold group as a complex of groups [C, Ha].

The classical theory for 2-dimensional representations is not sufficient to detect all essential surfaces in Haken manifolds. Throughout the paper let  $M$  be a compact connected orientable 3-manifold. We denote by  $X_n(M)$  the  $SL_n(\mathbb{C})$ -character variety of  $\pi_1 M$ . It was discovered by Boyer and Zhang [BZ], and Motegi [Mo] that there exist infinitely many Haken manifolds  $M$ , which are even hyperbolic, such that  $X_2(M)$  has no irreducible component of positive dimension. See also [SZ] for further study on the topic. We say that *an essential surface  $S$  in  $M$  is given by an ideal point  $\chi$  of a curve in  $X_n(M)$*  if  $S$  is constructed from  $\chi$  by the Culler-Shalen method or its extension developed in [HK] as described in Subsection 2.3. Hara and the second author formulated and raised the following question [HK, Question 6.1].

---

2010 *Mathematics Subject Classification.* Primary 57N10, Secondary 57M05, 20E42.

*Key words and phrases.* 3-manifold, essential surface, character variety, Bruhat-Tits building, separable subgroup.

**Question 1.1.** *Does there exist an essential surface in some 3-manifold  $M$  not given by any ideal point of curves in  $X_2(M)$  but given by an ideal point of a curve in  $X_n(M)$  for some  $n$ ?*

The aim of this paper is to show that the extension of Culler-Shalen theory to the case of higher-dimensional representations [HK] detects all essential surfaces in Haken manifolds. The following is the main theorem of this paper, which, in particular, gives an affirmative answer to Question 1.1.

**Theorem 1.2.** *Every connected essential surface in  $M$  is given by an ideal point of a rational curve in  $X_n(M)$  for some  $n$ .*

The proof of Theorem 1.2 relies on the breakthroughs of Agol [A] and Wise [W], and the subsequent works of Przytycki and Wise [PW1, PW2] on the separability of subgroups in a 3-manifold group. For a given connected essential surface  $S$  in  $M$  there exists a non-separating lift  $T$  of  $S$  in some finite cover  $N$  of  $M$  by the separability of  $\pi_1 S$  in  $\pi_1 M$ . The non-separating surface  $T$  defines abelian representations  $\pi_1 N \rightarrow \mathrm{SL}_2(\mathbb{C})$  parameterized in  $\mathbb{C}^\times$ , which induces an affine curve  $D_T$  consisting of representations  $\pi_1 M \rightarrow \mathrm{SL}_n(\mathbb{C})$  where  $n$  is twice the degree of the cover  $N$ . The set of characters of representations in  $D_T$  is a desired rational curve in  $X_n(M)$  as in the statement of Theorem 1.2, which has a unique ideal point. Then analyzing the structure of the Bruhat-Tits building associated to the function field of  $D_T$ , we explicitly construct a PL-map from the universal cover of  $M$  to the 1-skeleton of the building. Finally, we show that the inverse image of midpoints of edges by the PL-map is isotopic to parallel copies of  $S$ .

The paper is organized as follows. In Section 2 we review the extension of the Culler-Shalen method to the case of  $\mathrm{SL}_n(\mathbb{C})$ -representations in [HK]. Here we recall some of the standard facts on  $\mathrm{SL}_n(\mathbb{C})$ -character varieties and Bruhat-Tits buildings associated to the special linear group. In Subsection 2.3, we give the precise definition of the sentence ‘*an essential surface is given by an ideal point*’. Section 3 provides a brief exposition on the separability of surface subgroups by Przytycki and Wise [PW2]. Section 4 is devoted to the proof of Theorem 1.2.

**Acknowledgment.** We wish to thank Steven Boyer and Takashi Hara for helpful conversations. The first and the third author was supported by the SFB 1085 ‘Higher Invariants’ at the Universität Regensburg funded by the Deutsche Forschungsgemeinschaft (DFG). The second author was supported by JSPS KAKENHI (No. 26800032).

## 2. $\mathrm{SL}_n$ -CULLER SHALEN THEORY

We begin with an overview of the extension of Culler-Shalen theory to the case of higher-dimensional representations in [HK].

**2.1. Character varieties.** We briefly review the  $\mathrm{SL}_n(\mathbb{C})$ -character variety of a finitely generated group. See [LM, Si1, Si2] for more details.

Let  $\pi$  be a finitely generated group. We define the following affine algebraic set

$$R_n(\pi) = \mathrm{Hom}(\pi, \mathrm{SL}_n(\mathbb{C})).$$

The algebraic group  $\mathrm{SL}_n(\mathbb{C})$  acts on the affine algebraic set  $R_n(\pi)$  by conjugation. We denote by  $X_n(\pi)$  the GIT quotient of the action [MFK]:

$$X_n(\pi) = \mathrm{Hom}(\pi, \mathrm{SL}_n(\mathbb{C})) // \mathrm{SL}_n(\mathbb{C}).$$

The affine algebraic set  $X_n(\pi)$  is called the  $\mathrm{SL}_n(\mathbb{C})$ -character variety of  $\pi$ . By definition the coordinate ring  $\mathbb{C}[X_n(\pi)]$  is isomorphic to the subring  $\mathbb{C}[R_n(\pi)]^{\mathrm{SL}_n(\mathbb{C})}$  of  $\mathbb{C}[R_n(\pi)]$  consisting of  $\mathrm{SL}_n(\mathbb{C})$ -invariant functions. Procesi [P, Theorem 1.3] showed that  $\mathbb{C}[R_n(\mathbb{C})]^{\mathrm{SL}_n(\mathbb{C})}$  is generated by *trace functions*  $I_\gamma$  for  $\gamma \in \pi$  defined by

$$I_\gamma(\rho) = \mathrm{tr} \rho(\gamma)$$

for  $\rho \in R_n(\pi)$ . Therefore  $X_n(\mathbb{C})$  is identified with the set of *characters*  $\chi_\rho$  for  $\rho \in R_n(\pi)$  defined by

$$\chi_\rho(\gamma) = \mathrm{tr} \rho(\gamma)$$

for  $\gamma \in \pi$ . For a compact connected orientable 3-manifold  $M$ , we abbreviate  $R_n(\pi_1 M)$  and  $X_n(\pi_1 M)$  with  $R_n(M)$  and  $X_n(M)$  respectively to simplify notation.

Let  $C$  be an affine variety, and denote by  $\mathbb{C}(C)$  its field of rational functions. We call  $C$  an *affine curve* if the transcendence degree of  $\mathbb{C}(C)$  over  $\mathbb{C}$  equals 1 [F, Section 6.5]. Consider an affine curve  $C$  and its projectivisation  $\overline{C}$ . The projective curve  $\overline{C}$  might not be smooth, but it has a unique smooth model, i.e., there is a smooth projective curve  $\tilde{C}$  together with a birational map  $\tilde{C} \dashrightarrow \overline{C}$  which is universal [F, Theorem 7.3]. Recall that a birational equivalence induces an isomorphism on the associated fields of rational functions [F, Proposition 6.12]. Thus their fields of rational functions all agree:  $\mathbb{C}(C) = \mathbb{C}(\overline{C}) = \mathbb{C}(\tilde{C})$ . To a point  $P$  of  $\tilde{C}$  the local ring  $\mathcal{O}_{\tilde{C}, P}$  of  $\tilde{C}$  at  $P$  is associated. As the point  $P$  is a smooth point, the ring  $\mathcal{O}_{\tilde{C}, P}$  is a discrete valuation ring, which induces a discrete valuation  $v_P$  on  $\mathbb{C}(C)$  [F, Section 7.1].

An *ideal point*  $\chi$  of an affine curve  $C$  is a point of its smooth projective model  $\tilde{C}$  corresponding to a point of  $\overline{C} \setminus C$ . We can equip the rational functions  $\mathbb{C}(C)$  with the discrete valuation  $v_\chi$  associated to an ideal point  $\chi$  as described above.

**2.2. Bruhat-Tits buildings.** Following the exposition [G], we describe the Bruhat-Tits building [BT1, BT2, IM] associated to the special linear group over a discrete valuation field. See also [AB] for more details on buildings.

Let  $F$  be a commutative field equipped with a discrete valuation  $v$  which is not necessarily complete. We denote by  $\mathcal{O}_v$  the valuation ring associated to  $v$ . The *Bruhat-Tits building* associated to  $\mathrm{SL}_n(F)$ , which is an  $(n-1)$ -dimensional simplicial complex  $B_v$ , is defined as follows: A vertex of  $B_v$  is the homothety class of a lattice in the  $n$ -dimensional vector space  $F^n$ , where a *lattice* in  $F^n$  is a free  $\mathcal{O}_v$ -submodule of full rank, and two lattices  $\Lambda$  and  $\Lambda'$  are *homothetic* if  $\Lambda = \alpha\Lambda'$  for some  $\alpha \in F^\times$ . A set of  $(m+1)$  vertices  $s_0, s_1, \dots, s_m$  forms an  $m$ -simplex in  $B_v$  if and only if there exist lattices  $\Lambda_0, \Lambda_1, \dots, \Lambda_m$  representing  $s_0, s_1, \dots, s_m$  respectively such that after relabeling indices we have the flag relation

$$\omega\Lambda_m \subsetneq \Lambda_0 \subsetneq \Lambda_1 \subsetneq \dots \subsetneq \Lambda_m,$$

where  $\omega$  is an irreducible element of  $\mathcal{O}_v$ .

The simplicial complex  $B_v$  is known to be an *Euclidean building*, and, in particular, a CAT(0)-space with respect to the standard metric. See for instance [AB, Definition 11.1] for the definition of an Euclidean building. Since  $\mathrm{SL}_n(F)$  acts on the set of lattices in  $F^n$  so that homothety classes and above flag relations are preserved,  $\mathrm{SL}_n(F)$  acts also on  $B_v$ . This action is *type-preserving*, i.e., there exists an  $\mathrm{SL}_n(F)$ -invariant map  $\tau: B_v^{(0)} \rightarrow \mathbb{Z}/n\mathbb{Z}$  such that  $\tau|_{\Delta^{(0)}}$  is a bijection for each

$(n - 1)$ -simplex  $\Delta$  in  $B_v$ . Here for a simplicial complex  $K$  we denote by  $K^{(m)}$  the  $m$ -skeleton of  $K$ . In particular, for any subgroup  $G$  of  $\mathrm{SL}_n(F)$  the quotient  $B_v/G$  is again an  $(n - 1)$ -dimensional simplicial complex.

*Remark 2.1.* In the case of  $n = 2$  the above construction is nothing but the one of the tree associated to  $\mathrm{SL}_2(F)$  in [Se1, Se2].

**2.3. An ideal point giving an essential surface.** We summarize the construction in [HK] of a certain branched surface from an ideal point of a curve in the character variety. Here we restrict our attention to the case where such a branched surface has no branched points, and is an essential surface.

Let  $C$  be a curve in  $X_n(M)$  and  $\chi$  an ideal point of  $C$ . We denote by  $t: R_n(M) \rightarrow X_n(M)$  the quotient map. There exists a curve  $D$  in  $t^{-1}(C)$  such that  $t|_D$  is not a constant map, and a regular map  $\tilde{t}|_D: \tilde{D} \rightarrow \tilde{C}$  on the smooth projective models is induced by  $t|_D$ . We take a lift  $\tilde{\chi} \in (\tilde{t}|_D)^{-1}(\{\chi\})$ , and denote by  $B_{\tilde{\chi}}$  the Bruhat-Tits building associated to  $\mathrm{SL}_n(\mathbb{C}(D))$ , where the function field  $\mathbb{C}(D)$  is equipped with the discrete valuation at  $\tilde{\chi}$ . The *tautological representation*  $\mathcal{P}: \pi_1 M \rightarrow \mathrm{SL}_n(\mathbb{C}(D))$  is defined by

$$\mathcal{P}(\gamma)(\rho) = \rho(\gamma)$$

for  $\gamma \in \pi_1 M$  and  $\rho \in D$ . Pulling back the action of  $\mathrm{SL}_n(\mathbb{C}(D))$  on  $B_{\tilde{\chi}}$  by  $\mathcal{P}$ , we obtain the action of  $\pi_1 M$  on  $B_{\tilde{\chi}}$ . Extending [CS, Theorem 2.2.1] to the case of a general  $n$ , Hara and the second author [HK, Corollary 4.5] proved that the action is *nontrivial*, i.e., for every vertex of  $B_{\tilde{\chi}}$  its stabilizer subgroup of  $\pi_1 M$  is proper.

Recall that a compact orientable properly-embedded surface  $S$  in  $M$  is called *essential* if for any component  $S_0$  of  $S$  the inclusion-induced homomorphism  $\pi_1 S_0 \rightarrow \pi_1 M$  is injective, and  $S_0$  is not boundary-parallel nor homeomorphic to the 2-sphere  $S^2$ . We say that *an essential surface  $S$  is given by an ideal point  $\chi$*  if for some lift  $\tilde{\chi}$  of  $\chi$  there exists a PL map  $f: M \rightarrow B_{\tilde{\chi}}^{(1)}/\pi_1 M$  whose inverse image of the set of midpoints of the edges in  $B_{\tilde{\chi}}$  is isotopic to some number of parallel copies of  $S$ .

When  $n = 2$ , since  $\pi_1 M$  nontrivially acts on the tree  $B_{\tilde{\chi}}$  without inversions, every ideal point  $\chi$  gives some essential surface in  $M$  [CS, Proposition 2.3.1]. In general, it follows from the proof of [HK, Theorem 4.7] that if  $n = 3$  or if  $\partial M$  is non-empty, then there exists a PL map  $f: M \rightarrow B_{\tilde{\chi}}^{(2)}/\pi_1 M$  such that  $f^{-1}(Y)$  is a certain branched surface called *essential tribranched surface* [HK, Definition 2.2], where  $Y$  is the union of edges in the first barycentric subdivision of  $B_{\tilde{\chi}}^{(2)}/\pi_1 M$  not contained in  $B_{\tilde{\chi}}^{(1)}/\pi_1 M$ . Note that an essential tribranched surface without any branched points is nothing but an essential surface in the usual sense.

*Remark 2.2.* The authors [FKN] showed that every closed 3-manifold  $M$  with  $\mathrm{rank} \pi_1 M \geq 4$  contains an essential tribranched surface.

### 3. SURFACE SUBGROUP SEPARABILITY

We recall the separability of surface subgroups in a 3-manifold group proved by Przytycki and Wise [PW2], which is a key ingredient of the proof of Theorem 1.2. A subgroup  $H$  of a group  $G$  is *separable* if  $H$  equals the intersection of finite index subgroups of  $G$  containing  $H$ .

**Theorem 3.1.** ([PW2, Theorem 1.1]) *Let  $S$  be a connected essential surface  $S$  in  $M$ . Then  $\pi_1 S$  is separable in  $\pi_1 M$ .*

Theorem 3.1 was proved by Przytycki and Wise [PW1] when  $M$  is a graph manifold, and by Wise [W] when  $M$  is a hyperbolic manifold. In fact, every finitely generated subgroup of  $\pi_1 M$  is separable when  $M$  is a hyperbolic manifold, by Wise [W] in the case where  $M$  has a non-empty boundary and by Agol [A] in the case where  $M$  is closed. See also Liu [L] for a refinement of the separability.

The following is a topological interpretation of Theorem 3.1. While it is well-known for experts, nevertheless we give a proof for the sake of completeness. See also [Sc, Lemma 1.4].

**Corollary 3.2.** *For an essential surface  $S$  in  $M$  there exists some finite cover of  $M$  where the inverse image of  $S$  contains a non-separating component.*

*Proof.* We may assume that  $S$  is connected and separating. Let  $M_+$  and  $M_-$  be the two components of the complement of  $S$ . It follows from [He, Theorem 10.5] that  $\pi_1 S$  has index at least two in  $\pi_1 M_-$  and in  $\pi_1 M_+$ . It follows from Theorem 3.1 that there exists an epimorphism  $\varphi: \pi_1 M \rightarrow G$  to a finite group such that  $\varphi(\pi_1 S) \neq \varphi(\pi_1 M_{\pm})$ . In particular, we have

$$[G : \varphi(\pi_1 S)] \geq 2[G : \varphi(\pi_1 M_{\pm})].$$

Let  $p: M_{\varphi} \rightarrow M$  be the covering corresponding to  $\text{Ker } \varphi$ . The numbers of components of  $p^{-1}(M_{\pm})$  and  $p^{-1}(S)$  are equal to  $[G : \varphi(\pi_1 M_{\pm})]$  and  $[G : \varphi(\pi_1 S)]$  respectively, and the above inequality implies

$$[G : \varphi(\pi_1 S)] \geq [G : \varphi(\pi_1 M_+)] + [G : \varphi(\pi_1 M_-)].$$

Thus the number of components of  $p^{-1}(S)$  is greater than or equal to that of its complement, which shows that some component of  $p^{-1}(S)$  is non-separating.  $\square$

#### 4. PROOF OF THE MAIN THEOREM

Now we prove the main theorem. For the readers' convenience we recall the statement.

**Theorem 4.1** (Theorem 1.2). *Every connected essential surface in  $M$  is given by an the ideal point of a rational curve in  $X_n(M)$  for some  $n$ .*

Let  $S$  be a connected essential surface in  $M$ . It follows from Corollary 3.2 that there exists a  $d$ -fold covering  $p: N \rightarrow M$  for some  $d$  such that  $p^{-1}(S)$  contains a non-separating component  $T$ . Then the proof is divided into two parts: First, we construct a rational curve  $C_T$  in  $X_{2d}(M)$  with a unique ideal point  $\chi_T$ , which is determined by  $T$ . Second, for a lift  $\tilde{\chi}_T$  of  $\chi_T$  we construct a PL map  $f: M \rightarrow B_{\tilde{\chi}_T}^{(1)}$  whose inverse image of the set of midpoints of edges in  $B_{\tilde{\chi}_T}$  is isotopic to two parallel copies of  $S$ .

**4.1. Construction of a curve.** We denote by  $\psi: \pi_1 N \rightarrow \mathbb{Z}$  the epimorphism induced by the intersection pairing with  $T$ . For each  $z \in \mathbb{C}^{\times}$  we define the representation  $\tilde{\rho}_z: \pi_1 N \rightarrow \text{SL}_2(\mathbb{C})$  to be the composition of  $\psi$  and the homomorphism  $\mathbb{Z} \rightarrow \text{SL}_2(\mathbb{C})$  which sends an integer  $k$  to the matrix

$$\begin{pmatrix} z^k & 0 \\ 0 & z^{-k} \end{pmatrix}.$$

We consider the induced representation  $\rho_z : \pi_1 M \rightarrow \text{Aut}(\mathbb{C}[\pi_1 M] \otimes_{\mathbb{C}[\pi_1 N]} \mathbb{C}^2)$  of  $\tilde{\rho}_z$ . Fixing representatives  $\gamma_1, \dots, \gamma_d \in \pi_1 M$  of the elements of  $\pi_1 M/p_*(\pi_1 N)$ , we have the decomposition

$$\mathbb{C}[\pi_1 M] \otimes_{\mathbb{C}[\pi_1 N]} \mathbb{C}^2 = \bigoplus_{i=1}^d \gamma_i \otimes \mathbb{C}^2,$$

which is naturally identified with  $\mathbb{C}^{2d}$ . Thus we regard  $\rho_z$  as a representation  $\pi_1 M \rightarrow \text{SL}_{2d}(\mathbb{C})$ . We now set

$$D_T = \{\rho_z \in R_{2d}(M) : z \in \mathbb{C}^\times\},$$

$$C_T = \{\chi_{\rho_z} \in X_{2d}(M) : z \in \mathbb{C}^\times\}.$$

**Lemma 4.2.** (1) *The set  $D_T$  is a curve in  $R_{2d}(M)$  isomorphic to  $\mathbb{C}^\times$ .*  
(2) *The set  $C_T$  is a rational curve in  $X_{2d}(M)$  with a unique ideal point.*

*Proof.* These sets of representations and characters are constructed along the following commutative diagram:

$$\begin{array}{ccccccc} \mathbb{C}^\times & \longrightarrow & R_2(\mathbb{Z}) & \longrightarrow & R_2(N) & \longrightarrow & R_{2d}(M) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \mathbb{C} & \longrightarrow & X_2(\mathbb{Z}) & \longrightarrow & X_2(N) & \longrightarrow & X_{2d}(M), \end{array}$$

where the first vertical map sends  $z \in \mathbb{C}^\times$  to  $z + z^{-1} \in \mathbb{C}$ , and the first bottom horizontal map is an isomorphism which sends  $w \in \mathbb{C}$  to the character of  $\mathbb{Z}$  whose image of  $1 \in \mathbb{Z}$  is  $w$ . The composition of the top horizontal maps is called  $\Psi$  and the composition of the bottom horizontal maps is called  $\Phi$ . Then the sets  $D_T$  and  $C_T$  coincide with the images of  $\Psi$  and  $\Phi$

We may assume  $\gamma_1 \in \pi_1 N$ , and take  $\mu \in \pi_1 N$  with  $\psi(\mu) = 1$ . Then we have

$$\rho_z(\mu) = \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix} \oplus \bigoplus_{i=2}^d \begin{pmatrix} z^{\psi(\gamma_i^{-1}\mu\gamma_i)} & 0 \\ 0 & z^{-\psi(\gamma_i^{-1}\mu\gamma_i)} \end{pmatrix},$$

$$\chi_{\rho_z}(\mu) = z + z^{-1} + \sum_{i=2}^d \left( z^{\psi(\gamma_i^{-1}\mu\gamma_i)} + z^{-\psi(\gamma_i^{-1}\mu\gamma_i)} \right).$$

Hence the restriction of the map  $R_{2d}(M) \rightarrow \mathbb{C}^2$  sending a representation  $\rho$  to the vector of the (1, 1)- and (2, 2)-entries of  $\rho(\mu)$  gives the inverse regular map  $D_T \rightarrow \mathbb{C}^\times$  of  $\Psi$ , where  $\mathbb{C}^\times$  is identified with the curve  $xy - 1$  in  $\mathbb{C}^2$ , and (1) is proved.

We deduce from the second equation above that the map  $\Phi$  is not constant. Also by fixing an affine space  $\mathbb{C}^N$  containing  $X_{2d}(M)$ , we regard  $\Phi$  as a map  $\mathbb{C} \rightarrow \mathbb{C}^N$ . We denote by  $\overline{\Phi} : \mathbb{P}^1 \rightarrow \mathbb{P}^N$  the projective extension of  $\Phi$ . Since  $\overline{\Phi}$  is not a constant map, by the completeness of the projective line  $\mathbb{P}^1$  [Mu, Section I.9, Theorem 1] the image  $\overline{C}_T$  of  $\overline{\Phi}$  is a projective curve, and by the Riemann-Hurwitz formula the curve  $\overline{C}_T$  is rational. Therefore the set  $C_T$ , which coincides with the intersection of  $\overline{C}_T$  and  $\mathbb{C}^N$ , is an affine rational curve. Since  $\Phi$  induces a surjective regular map  $\tilde{\Phi} : \mathbb{P}^1 \rightarrow \tilde{C}_T$  on the smooth projective models, the rational curve  $C_T$  has a unique ideal point corresponding to the point at infinity of  $\mathbb{P}^1$ , which proves (2).  $\square$

It is a simple matter to check that both the two ideal points of  $D_T$  corresponding to 0 and  $\infty$  are lifts of the unique ideal point  $\chi_T$  of  $C_T$ . Let  $\tilde{\chi}_T$  be the one corresponding to 0. Then as in Section 2.3 we obtain the nontrivial action  $\pi_1 M$  on the Bruhat-Tits building  $B_{\tilde{\chi}_T}$  associated to  $\text{SL}_{2d}(\mathbb{C}(D_T))$ . We identify  $\mathbb{C}(D_T)$  with

the standard function field  $\mathbb{C}(t)$  and the valuation at  $\tilde{\chi}_T$  with the lowest degree of the Laurent expansion of a rational function. Then the vector space  $\mathbb{C}(t)^{2d}$  is decomposed into

$$\mathbb{C}[\pi_1 M] \otimes_{\mathbb{C}[\pi_1 N]} \mathbb{C}(t)^2 = \bigoplus_{i=1}^d \gamma_i \otimes \mathbb{C}(t)^2,$$

where  $\pi_1 N$  acts on  $\mathbb{C}(t)^2$  by the representation  $\mathcal{Q}: \pi_1 N \rightarrow \mathrm{SL}_2(\mathbb{C}(t))$  defined by

$$\mathcal{Q}(\gamma) = \begin{pmatrix} t^{\psi(\gamma)} & 0 \\ 0 & t^{-\psi(\gamma)} \end{pmatrix}$$

for  $\gamma \in \pi_1 N$ , and the tautological representation  $\mathcal{P}: \pi_1 M \rightarrow \mathrm{SL}_{2d}(\mathbb{C}(t))$  is given by the left multiplication on  $\mathbb{C}[\pi_1 M] \otimes_{\mathbb{C}[\pi_1 N]} \mathbb{C}(t)^2$ .

**4.2. Construction of a PL-map.** We take a triangulation of  $M$  containing  $S$  as a normal surface. We may assume that the intersection of each tetrahedron with  $S$  is connected, if necessary, replacing the triangulation by an appropriate subdivision. The triangulation of  $M$  induces ones of  $N$  and the universal cover  $\tilde{M}$  of  $M$ , so that  $T$  and its inverse image  $\tilde{T}$  by the covering  $\tilde{M} \rightarrow N$  are also normal surfaces. Then we take a cellular map  $g: N \rightarrow \mathbb{R}/\mathbb{Z}$  such that  $g^{-1}([\frac{1}{2}]) = T$ , where we consider the cellular structure of  $\mathbb{R}/\mathbb{Z}$  with one vertex corresponding to  $\mathbb{Z}$ . We define  $\tilde{g}: \tilde{M} \rightarrow \mathbb{R}$  to be the  $\pi_1 N$ -equivariant lift of  $g$ , so that  $\tilde{g}^{-1}(\frac{1}{2} + \mathbb{Z}) = \tilde{T}$ .

We now define a map  $\tilde{f}^{(0)}: \tilde{M}^{(0)} \rightarrow B_{\tilde{\chi}_T}^{(0)}$  as follows. For  $s \in \tilde{M}^{(0)}$  we consider the lattice

$$\bigoplus_{i=1}^d \gamma_i \otimes \Lambda_{\tilde{g}(\gamma_i^{-1}s)}$$

in  $\mathbb{C}[\pi_1 M] \otimes_{\mathbb{C}[\pi_1 N]} \mathbb{C}(t)^2$ , where  $\Lambda_n$  is the lattice in  $\mathbb{C}(t)^2$  generated by the vectors

$$\begin{pmatrix} t^n \\ 0 \end{pmatrix} \text{ and } \begin{pmatrix} 0 \\ t^{-n} \end{pmatrix}.$$

Note that  $\tilde{g}(\gamma_i^{-1}s)$  is an integer for each  $i$  by the construction of  $\tilde{g}$ . Then we set  $\tilde{f}^{(0)}(s)$  to be the homothety class of the above lattice. In the following two lemmas we observe the key properties of  $\tilde{f}^{(0)}$ .

**Lemma 4.3.** *The map  $\tilde{f}^{(0)}$  is  $\pi_1 M$ -equivariant.*

*Proof.* For  $\gamma \in \pi_1 M$  there exist a permutation  $\sigma$  of degree  $d$  and  $\delta_i \in \pi_1 N$  such that

$$\gamma \gamma_i = \gamma_{\sigma(i)} \delta_i$$

for each  $i$ . Then

$$\begin{aligned} \bigoplus_{i=1}^d \gamma \gamma_i \otimes \Lambda_{\tilde{g}(\gamma_i^{-1}s)} &= \bigoplus_{i=1}^d \gamma_{\sigma(i)} \delta_i \otimes \Lambda_{\tilde{g}(\gamma_i^{-1}s)} &= \bigoplus_{i=1}^d \gamma_{\sigma(i)} \otimes \mathcal{Q}(\delta_i) \cdot \Lambda_{\tilde{g}(\gamma_i^{-1}s)} \\ &= \bigoplus_{i=1}^d \gamma_{\sigma(i)} \otimes \Lambda_{\tilde{g}(\gamma_i^{-1}s) + \psi(\delta_i)} &= \bigoplus_{i=1}^d \gamma_{\sigma(i)} \otimes \Lambda_{\tilde{g}(\delta_i \gamma_i^{-1}s)} \\ &= \bigoplus_{i=1}^d \gamma_{\sigma(i)} \otimes \Lambda_{\tilde{g}(\gamma_{\sigma(i)}^{-1} \gamma s)} &= \bigoplus_{i=1}^d \gamma_i \otimes \Lambda_{\tilde{g}(\gamma_i^{-1} \gamma s)} \end{aligned}$$

for  $\gamma \in \pi_1 M$  and  $s \in \tilde{M}^{(0)}$ , which implies that  $\tilde{f}^{(0)}$  is a  $\pi_1 M$ -equivariant map.  $\square$

**Lemma 4.4.** *For each tetrahedron  $\Delta$  in  $\widetilde{M}$  the set  $\tilde{f}^{(0)}(\Delta^{(0)})$  consists of one vertex if  $\gamma_i^{-1} \cdot \Delta$  does not intersect with  $\widetilde{T}$  for any  $i$ , and two vertices of distance 2 with respect to the graph metric on  $B_{\widetilde{\chi}}^{(1)}$  otherwise.*

*Proof.* If  $\gamma_i^{-1} \cdot \Delta$  does not intersect with  $\widetilde{T}$  for any  $i$ , then it follows from the choice of  $g$  that there exists some  $n_i \in \mathbb{Z}$  such that

$$\tilde{g}(\gamma_i^{-1} \cdot \Delta^{(0)}) = \{n_i\}$$

for each  $i$ , and hence we obtain

$$\tilde{f}^{(0)}(\gamma_i^{-1} \cdot \Delta^{(0)}) = \left\{ \left[ \bigoplus_{i=1}^d \gamma_i \otimes \Lambda_{n_i} \right] \right\}.$$

In the following we consider the case where  $\gamma_i^{-1} \cdot \Delta$  intersects with  $\widetilde{T}$  for  $i = i_1, \dots, i_m$ . Then the intersections of  $\gamma_i^{-1} \cdot \Delta$  with  $\widetilde{T}$  are all connected and of same type for  $k = 1, \dots, m$ , since otherwise  $p(T) = S$  implies that the intersection of some tetrahedron in  $M$  with  $S$  is not a normal disc, which contradicts the choice of the triangulation of  $M$ . Thus  $\Delta^{(0)}$  is divided into two subsets  $\Delta_+^{(0)}$  and  $\Delta_-^{(0)}$  satisfying the following:

- (1) there exists some  $n_{i_k} \in \mathbb{Z}$  such that

$$\tilde{g}(\gamma_{i_k}^{-1} \cdot \Delta_+^{(0)}) = \{n_{i_k} + 1\} \text{ and } \tilde{g}(\gamma_{i_k}^{-1} \cdot \Delta_-^{(0)}) = \{n_{i_k}\}$$

for  $k = 1, \dots, m$ ;

- (2) there exists some  $n_i \in \mathbb{Z}$  such that

$$\tilde{g}(\gamma_i^{-1} \cdot \Delta^{(0)}) = \{n_i\}$$

for  $i \neq i_1, \dots, i_m$ .

Hence we obtain

$$\tilde{f}^{(0)}(\Delta_+^{(0)}) = \{[\Lambda_+]\} \text{ and } \tilde{f}^{(0)}(\Delta_-^{(0)}) = \{[\Lambda_-]\},$$

where

$$\begin{aligned} \Lambda_+ &= \left( \bigoplus_{k=1}^m \gamma_{i_k} \otimes \Lambda_{n_{i_k}+1} \right) \oplus \left( \bigoplus_{i \neq i_1, \dots, i_m} \gamma_i \otimes \Lambda_{n_i} \right), \\ \Lambda_- &= \left( \bigoplus_{k=1}^m \gamma_{i_k} \otimes \Lambda_{n_{i_k}} \right) \oplus \left( \bigoplus_{i \neq i_1, \dots, i_m} \gamma_i \otimes \Lambda_{n_i} \right). \end{aligned}$$

Since

$$\begin{aligned} t\Lambda'_n &\subsetneq \Lambda_{n+1} \subsetneq \Lambda'_n, \\ t\Lambda'_n &\subsetneq \Lambda_n \subsetneq \Lambda'_n \end{aligned}$$

for  $n \in \mathbb{Z}$ , where  $\Lambda'_n$  is the lattice in  $\mathbb{C}(t)^2$  generated by the vectors

$$\begin{pmatrix} t^n \\ 0 \end{pmatrix} \text{ and } \begin{pmatrix} 0 \\ t^{-n-1} \end{pmatrix},$$

we have

$$\begin{aligned} t\Lambda' &\subsetneq \Lambda_+ \subsetneq \Lambda', \\ t\Lambda' &\subsetneq \Lambda_- \subsetneq \Lambda', \end{aligned}$$



where

$$\Lambda' = \left( \bigoplus_{k=1}^m \gamma_{i_k} \otimes \Lambda'_{n_{i_k}} \right) \oplus \left( \bigoplus_{i \neq i_1, \dots, i_m} \gamma_i \otimes \Lambda_{n_i} \right).$$

By the definition of the building  $B_{\tilde{\chi}}$  these relations imply that there exist edges in  $B_{\tilde{\chi}}$  connecting  $[\Lambda_+]$  and  $[\Lambda_-]$  with  $[\Lambda']$ , and hence the distance between  $[\Lambda_+]$  and  $[\Lambda_-]$  is at most 2 in  $B_{\tilde{\chi}}^{(1)}$ . We further observe that the matrix

$$\left( \bigoplus_{i=1}^n \gamma_{i_k} \otimes \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \right) \oplus \left( \bigoplus_{i \neq i_1, \dots, i_m} \gamma_i \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right)$$

in  $\mathrm{SL}_{2d}(\mathbb{C}(t))$  sends  $[\Lambda_-]$  to  $[\Lambda_+]$ . Since the action of  $\mathrm{SL}_{2d}(\mathbb{C}(t))$  on  $B_{\tilde{\chi}}$  is type-preserving, the distance between them is exactly equal to 2, and the lemma follows.  $\square$

We are now in position to construct a desired PL map  $f: M \rightarrow B_{\tilde{\chi}_T}^{(1)}/\pi_1 M$ . It follows from Lemmas 4.3 and 4.4 that  $\tilde{f}^{(0)}$  extends to a  $\pi_1 M$ -equivariant simplicial map  $\tilde{f}: \tilde{N} \rightarrow B_{\tilde{\chi}_T}^{(1)}$  with respect to the 1st barycentric subdivision of the triangulation of  $\tilde{M}$ . We define  $f: M \rightarrow B_{\tilde{\chi}_T}^{(1)}/\pi_1 M$  to be the quotient of  $\tilde{f}$  by  $\pi_1 M$ . By the construction of  $\tilde{f}$  we check at once that the inverse image of the set of midpoints of edges in  $B_{\tilde{\chi}_T}$  by  $\tilde{f}$  is isotopic to two parallel copies of  $\tilde{T}$ . Since  $\tilde{f}$  is  $\pi_1 M$ -equivariant and since  $p(T) = S$ , the inverse image of the set of midpoints of edges in  $B_{\tilde{\chi}_T}/\pi_1 M$  by  $f$  is isotopic to two parallel copies of  $S$ . Therefore  $S$  is given by  $\chi_T$ , which completes the proof.

#### REFERENCES

- [AB] P. Abramenko and K. S. Brown, *Buildings. Theory and applications*, Graduate Texts in Mathematics, 248, Springer, New York, 2008. xxii+747 pp.
- [A] I. Agol, *The virtual Haken conjecture*, With an appendix by Agol, Daniel Groves, and Jason Manning, Doc. Math. **18** (2013), 1045–1087.
- [BZ] S. Boyer and X. Zhang, *On Culler-Shalen seminorms and Dehn filling*, Ann. of Math. (2) **148** (1998), no. 3, 737–801.
- [BT1] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d'une donnée radicielle valuée*, Inst. Hautes Études Sci. Publ. Math. **60** (1984), 197–376.
- [BT2] F. Bruhat and J. Tits, *Groupes réductifs sur un corps local* (French), Inst. Hautes Études Sci. Publ. Math. **41** (1972), 5–251.
- [C] J. M. Corson, *Complexes of groups*, Proc. London Math. Soc. (3) **65** (1992), no. 1, 199–224.
- [CS] M. Culler and P. B. Shalen, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math. (2) **117** (1983), no. 1, 109–146.
- [FKN] S. Friedl, T. Kitayama and M. Nagel, *A note on the existence of essential tribranched surfaces*, arXiv:1505.01806.
- [F] W. Fulton, *Algebraic Curves*, Advanced Book Program, W.A. Benjamin Inc., Massachusetts, 1974.
- [G] P. Garrett, *Buildings and classical groups*, Chapman & Hall, London, 1997. xii+373 pp.
- [HK] T. Hara and T. Kitayama, *Character varieties of higher dimensional representations and splittings of 3-manifolds*, arXiv:1410.4295.
- [Ha] A. Haefliger, *Complexes of groups and orbihedra*, Group theory from a geometrical viewpoint (Trieste, 1990), 504–540, World Sci. Publ., River Edge, NJ, 1991.
- [He] J. Hempel, *3-Manifolds*, Ann. of Math. Studies, no. 86. Princeton University Press, Princeton, N. J., 1976.

- [IM] N. Iwahori and H. Matsumoto, *On some Bruhat decomposition and the structure of the Hecke rings of  $p$ -adic Chevalley groups*, Inst. Hautes Études Sci. Publ. Math. **25** (1965), 5–48.
- [L] Y. Liu, *A characterization of virtually embedded subsurfaces in 3-manifolds*, arXiv:1406.4674.
- [LM] A. Lubotzky and A. R. Magid, *Varieties of representations of finitely generated groups*, Mem. Amer. Math. Soc. **58** (1985), no. 336, xi+117 pp.
- [Mo] K. Motegi, *Haken manifolds and representations of their fundamental groups in  $SL(2, \mathbb{C})$* , Topology Appl. **29** (1988), no. 3, 207–212.
- [Mu] D. Mumford, *The red book of varieties and schemes. Second, expanded edition. Includes the Michigan lectures (1974) on curves and their Jacobians. With contributions by Enrico Arbarello*, Lecture Notes in Mathematics, 1358. Springer-Verlag, Berlin, 1999. x+306 pp.
- [MFK] D. Mumford, J. Fogarty and F. Kirwan, *Geometric invariant theory, Third edition*, Ergebnisse der Mathematik und ihrer Grenzgebiete (2) 34. Springer-Verlag, Berlin, 1994. xiv+292 pp.
- [P] C. Procesi, *The invariant theory of  $n \times n$  matrices*, Advances in Math. **19** (1976), no. 3, 306–381.
- [PW1] P. Przytycki and D. T. Wise, *Graph manifolds with boundary are virtually special*, J. Topol. **7** (2014), no. 2, 419–435.
- [PW2] P. Przytycki and D. T. Wise, *Separability of embedded surfaces in 3-manifolds*, Compos. Math. **150** (2014), no. 9, 1623–1630.
- [SZ] S. Schanuel and X. Zhang, *Detection of essential surfaces in 3-manifolds with  $SL_2$ -trees*, Math. Ann. **320** (2001), no. 1, 149–165.
- [Sc] P. Scott, *Subgroups of surface groups are almost geometric*, J. London Math. Soc. (2) **17** (1978), no. 3, 555–565.
- [Se1] J.-P. Serre, *Arbres, amalgames,  $SL_2$* , Avec un sommaire anglais, Rédigé avec la collaboration de Hyman Bass, Astérisque, No. 46, Société Mathématique de France, Paris, 1977, 189 pp.
- [Se2] J.-P. Serre, *Trees*, Translated from the French by John Stillwell, Springer-Verlag, Berlin-New York, 1980. ix+142 pp.
- [Sh] P. B. Shalen, *Representations of 3-manifold groups*, Handbook of geometric topology, 955–1044, North-Holland, Amsterdam, 2002.
- [Si1] A. S. Sikora,  *$SL_n$ -character varieties as spaces of graphs*, Trans. Amer. Math. Soc. **353** (2001), no. 7, 2773–2804 (electronic).
- [Si2] A. S. Sikora, *Character varieties*, Trans. Amer. Math. Soc. **364** (2012), no. 10, 5173–5208.
- [W] D. T. Wise, *The structure of groups with a quasi-convex hierarchy*, preprint (2011), available at <http://www.math.mcgill.ca/wise/papers.html>.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF REGENSBURG, GERMANY  
*E-mail address:* `sfriedl@gmail.com`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, JAPAN  
*E-mail address:* `kitayama@ms.u-tokyo.ac.jp`

UNIVERSITÉ DU QUÉBEC À MONTRÉAL, QUÉBEC, CANADA  
*E-mail address:* `nagel@cirget.ca`