

A generalized Lyapunov's inequality for a fractional boundary value problem

Amar Chidouh

Laboratory of Dynamic Systems, Houari Boumediene University, Algiers, Algeria

Delfim F. M. Torres

*Center for Research and Development in Mathematics and Applications (CIDMA),
Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal*

Abstract

We prove existence of positive solutions to a nonlinear fractional boundary value problem. Then, under some mild assumptions on the nonlinear term, we obtain a smart generalization of Lyapunov's inequality. The new results are illustrated through examples.

Key words: fractional differential equations, Lyapunov's inequality, boundary value problem, positive solutions, Guo–Krasnoselskii fixed point theorem.

2010 MSC: 26A33, 34A08.

1. Introduction

Lyapunov's inequality is an outstanding result in mathematics with many different applications – see [6, 16] and references therein. The result, as proved by Lyapunov in 1907 [11], asserts that if $q : [a, b] \rightarrow \mathbb{R}$ is a continuous function, then a necessary condition for the boundary value problem

$$\begin{cases} y'' + qy = 0, & a < t < b, \\ y(a) = y(b) = 0 \end{cases} \quad (1)$$

to have a nontrivial solution is given by

$$\int_a^b |q(s)| ds > \frac{4}{b-a}. \quad (2)$$

Email addresses: m2ma.chidouh@gmail.com (Amar Chidouh), delfim@ua.pt (Delfim F. M. Torres)

Part of first author's Ph.D., which is carried out at Houari Boumediene University. See published version at <http://dx.doi.org/10.1016/j.cam.2016.03.035>

Submitted 30/Sept/2015; revised 22/Feb/2016; accepted 31/Mar/2016 in J. Comput. Appl. Math.

Lyapunov's inequality (2) has taken many forms, including versions in the context of fractional (noninteger order) calculus, where the second-order derivative in (1) is substituted by a fractional operator of order α .

Theorem 1 (See [3]). *Consider the fractional boundary value problem*

$$\begin{cases} {}_aD^\alpha y + qy = 0, & a < t < b, \\ y(a) = y(b) = 0, \end{cases} \quad (3)$$

where ${}_aD^\alpha$ is the (left) Riemann–Liouville derivative of order $\alpha \in (1, 2]$ and $q : [a, b] \rightarrow \mathbb{R}$ is a continuous function. If (3) has a nontrivial solution, then

$$\int_a^b |q(s)| ds > \Gamma(\alpha) \left(\frac{4}{b-a} \right)^{\alpha-1}. \quad (4)$$

A Lyapunov fractional inequality (4) can also be obtained by considering the fractional derivative in (3) in the sense of Caputo instead of Riemann–Liouville [4]. More recently, Rong and Bai obtained a Lyapunov-type inequality for a fractional differential equation but with fractional boundary conditions [13]. Motivated by [7, 8, 9, 12] and the above results, as well as existence results on positive solutions [1, 2, 10, 17], which are often useful in applications, we focus here on the following boundary value problem:

$$\begin{cases} {}_aD^\alpha y + q(t)f(y) = 0, & a < t < b, \\ y(a) = y(b) = 0, \end{cases} \quad (5)$$

where ${}_aD^\alpha$ is the Riemann–Liouville derivative and $1 < \alpha \leq 2$. Our first result asserts existence of nontrivial positive solutions to problem (5) (see Theorem 8). Then, under some assumptions on the nonlinear term f , we get a generalization of inequality (4) (see Theorem 10).

The paper is organized as follows. In Section 2 we recall some notations, definitions and preliminary facts, which are used throughout the work. Our results are given in Section 3: using the Guo–Krasnoselskii fixed point theorem, we establish in Section 3.1 our existence result; then, in Section 3.2, assuming that function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is continuous, concave and nondecreasing, we generalize Lyapunov's inequalities (2) and (4).

2. Preliminaries

Let $C[a, b]$ be the Banach space of all continuous real functions defined on $[a, b]$ with the norm $\|u\| = \sup_{t \in [a, b]} |u(t)|$. By $L[a, b]$ we denote the space of all real functions, defined on $[a, b]$, which are Lebesgue integrable with the norm

$$\|u\|_L = \int_a^b |u(s)| ds.$$

The reader interested in the fractional calculus is referred to [15]. Here we just recall the definition of (left) Riemann–Liouville fractional derivative.

Definition 2. *The Riemann–Liouville fractional derivative of order $\alpha > 0$ of a function $u : [a, b] \rightarrow \mathbb{R}$ is given by*

$${}_a D^\alpha u(t) = \frac{1}{\Gamma(n - \alpha)} \frac{d^n}{dt^n} \int_a^t \frac{u(s)}{(t - s)^{\alpha - n + 1}} ds,$$

where $n = [\alpha] + 1$ and Γ denotes the Gamma function.

Definition 3. *Let X be a real Banach space. A nonempty closed convex set $P \subset X$ is called a cone if it satisfies the following two conditions:*

- (i) $x \in P, \lambda \geq 0$, implies $\lambda x \in P$;
- (ii) $x \in P, -x \in P$, implies $x = 0$.

Lemma 4 (Jensen’s inequality [14]). *Let μ be a positive measure and let Ω be a measurable set with $\mu(\Omega) = 1$. Let I be an interval and suppose that u is a real function in $L(d\mu)$ with $u(t) \in I$ for all $t \in \Omega$. If f is convex on I , then*

$$f\left(\int_{\Omega} u(t) d\mu(t)\right) \leq \int_{\Omega} (f \circ u)(t) d\mu(t). \quad (6)$$

If f is concave on I , then the inequality (6) holds with “ \leq ” substituted by “ \geq ”.

Lemma 5 (Guo–Krasnoselskii fixed point theorem [5]). *Let X be a Banach space and let $K \subset X$ be a cone. Assume Ω_1 and Ω_2 are bounded open subsets of X with $0 \in \Omega_1 \subset \overline{\Omega}_1 \subset \Omega_2$, and let $T : K \cap (\overline{\Omega}_2 \setminus \Omega_1) \rightarrow K$ be a completely continuous operator such that*

- (i) $\|Tu\| \geq \|u\|$ for any $u \in K \cap \partial\Omega_1$ and $\|Tu\| \leq \|u\|$ for any $u \in K \cap \partial\Omega_2$; or
- (ii) $\|Tu\| \leq \|u\|$ for any $u \in K \cap \partial\Omega_1$ and $\|Tu\| \geq \|u\|$ for any $u \in K \cap \partial\Omega_2$.

Then, T has a fixed point in $K \cap (\overline{\Omega}_2 \setminus \Omega_1)$.

3. Main results

Let us consider the nonlinear fractional boundary value problem (5) and give its integral representation involving a Green function, which was deduced in [3].

Lemma 6. *Function y is a solution to the boundary value problem (5) if, and only if, y satisfies the integral equation*

$$y(t) = \int_a^b G(t, s) q(s) f(y(s)) ds,$$

where

$$G(t, s) = \frac{1}{\Gamma(\alpha)} \begin{cases} \frac{(t-a)^{\alpha-1}(b-s)^{\alpha-1}}{(b-a)^{\alpha-1}} - (t-s)^{\alpha-1}, & a \leq s \leq t \leq b, \\ \frac{(t-a)^{\alpha-1}(b-s)^{\alpha-1}}{(b-a)^{\alpha-1}}, & a \leq s \leq t \leq b, \end{cases} \quad (7)$$

is the Green function associated to problem (5).

Proof. Similar to the one found in [3]. \square

Lemma 7. *The Green function G defined by (7) satisfies the following properties:*

1. $G(t, s) \geq 0$ for all $a \leq t, s \leq b$;
2. $\max_{t \in [a, b]} G(t, s) = G(s, s)$, $s \in [a, b]$;
3. $G(s, s)$ has a unique maximum given by

$$\max_{s \in [a, b]} G(s, s) = \frac{(b-a)^{\alpha-1}}{4^{\alpha-1} \Gamma(\alpha)};$$

4. there exists a positive function $\varphi \in C(a, b)$ such that

$$\min_{t \in [\frac{2a+b}{3}, \frac{2b-a}{3}]} G(t, s) \geq \varphi(s)G(s, s), \quad a < s < b.$$

Proof. The first three properties are proved in [3]. Moreover, we know that Green's function $G(t, s)$ is decreasing with respect to t for $s \leq t$ and increasing with respect to t for $t \leq s$ [3]. To prove the fourth property, we define the following functions:

$$g_1(t, s) = \frac{1}{\Gamma(\alpha)} \left[\frac{(t-a)^{\alpha-1}(b-s)^{\alpha-1}}{(b-a)^{\alpha-1}} - (t-s)^{\alpha-1} \right]$$

and

$$g_2(t, s) = \frac{1}{\Gamma(\alpha)} \left[\frac{(t-a)^{\alpha-1}(b-s)^{\alpha-1}}{(b-a)^{\alpha-1}} \right].$$

Obviously, $G(t, s) > 0$ for $t, s \in (a, b)$ and one can seek the minimum in an interval of the form $[a + \frac{b-a}{n}, b - \frac{b-a}{n}]$, where $n \geq 3$ is a natural number. For $t \in [\frac{2a+b}{3}, \frac{2b-a}{3}]$,

$$\begin{aligned} \min_{t \in [\frac{2a+b}{3}, \frac{2b-a}{3}]} G(t, s) &= \begin{cases} g_1\left(\frac{2b-a}{3}, s\right) & \text{if } s \in (a, \frac{2a+b}{3}], \\ \min \left\{ g_1\left(\frac{2b-a}{3}, s\right), g_2\left(\frac{2a+b}{3}, s\right) \right\} & \text{if } s \in [\frac{2a+b}{3}, \frac{2b-a}{3}], \\ g_2\left(\frac{2a+b}{3}, s\right) & \text{if } s \in [\frac{2b-a}{3}, b), \end{cases} \\ &= \begin{cases} g_1\left(\frac{2b-a}{3}, s\right) & \text{if } s \in (a, \lambda], \\ g_2\left(\frac{2a+b}{3}, s\right) & \text{if } s \in [\lambda, b), \end{cases} \\ &= \frac{1}{\Gamma(\alpha)} \begin{cases} \left(\frac{(2b-4a)(b-s)}{3(b-a)} \right)^{\alpha-1} - \left(\frac{2b-a}{3} - s \right)^{\alpha-1} & \text{if } s \in (a, \lambda], \\ \left(\frac{b-s}{3} \right)^{\alpha-1} & \text{if } s \in [\lambda, b), \end{cases} \end{aligned}$$

where $\frac{2a+b}{3} < \lambda < \frac{2b-a}{3}$ is the unique solution of equation

$$g_1\left(\frac{2b-a}{3}, s\right) = g_2\left(\frac{2a+b}{3}, s\right).$$

Set

$$\varphi(s) = \begin{cases} \frac{\left(\frac{(2b-4a)(b-s)}{3}\right)^{\alpha-1} - (b-a)^{\alpha-1} \left(\frac{2b-a}{3} - s\right)^{\alpha-1}}{\left((s-a)(b-s)\right)^{\alpha-1}} & \text{if } s \in (a, \lambda], \\ \left(\frac{(b-a)}{3(s-a)}\right)^{\alpha-1} & \text{if } s \in [\lambda, b). \end{cases}$$

The proof is complete. \square

Let $X = C[a, b]$ and define the operator $T : X \rightarrow X$ as follows:

$$Ty(t) = \int_a^b G(t, s)q(s)f(y(s))ds, \quad y \in X. \quad (8)$$

To prove existence of solution to the fractional boundary value problem (5) it suffices to prove that the map T has a fixed point in K .

3.1. Existence of positive solutions

To prove existence of nontrivial positive solutions to the fractional boundary value problem (5) we consider the following hypotheses:

$$(H_1) \quad f(y) \geq \gamma^* r_1 \text{ for } y \in [0, r_1],$$

$$(H_2) \quad f(y) \leq \gamma r_2 \text{ for } y \in [0, r_2],$$

where $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is continuous. In what follows we take

$$\gamma := \left(\int_a^b G(s, s)q(s)ds \right)^{-1} \quad \text{and} \quad \gamma^* := \left(\int_{\frac{2a+b}{3}}^{\frac{2b-a}{3}} G(s, s)\varphi(s)q(s)ds \right)^{-1}.$$

Theorem 8. *Let $q : [a, b] \rightarrow \mathbb{R}_+$ be a nontrivial Lebesgue integrable function. Assume that there exist two positive constants $r_2 > r_1 > 0$ such that the assumptions (H_1) and (H_2) are satisfied. Then the fractional boundary value problem (5) has at least one nontrivial positive solution y belonging to X such that $r_1 \leq \|y\| \leq r_2$.*

For the proof of Theorem 8 we use Lemma 5 with the cone K given by

$$K := \{y \in X : y(t) \geq 0, a \leq t \leq b\}.$$

Proof of Theorem 8. Using the Ascoli–Arzela theorem, we prove that $T : K \rightarrow K$ is a completely continuous operator. Let

$$\Omega_i = \{y \in K : \|y\| \leq r_i\}.$$

From (H_1) and Lemma 7, we have for $t \in [\frac{2a+b}{3}, \frac{2b-a}{3}]$ and $y \in K \cap \partial\Omega_1$ that

$$\begin{aligned} (Ty)(t) &\geq \int_a^b \min_{t \in [\frac{2a+b}{3}, \frac{2b-a}{3}]} G(t, s) q(s) f(y(s)) ds \\ &\geq \gamma^* \left(\int_a^b G(s, s) \varphi(s) q(s) ds \right) r_1 \\ &\geq \gamma^* \left(\int_{\frac{2a+b}{3}}^{\frac{2b-a}{3}} G(s, s) \varphi(s) q(s) ds \right) r_1 = \|y\|. \end{aligned}$$

Thus, $\|Ty\| \geq \|y\|$ for $y \in K \cap \partial\Omega_1$. Let us now prove that $\|Ty\| \leq \|y\|$ for all $y \in K \cap \partial\Omega_2$. From (H_2) , it follows that

$$\|Ty\| = \max_{t \in [a, b]} \int_a^b G(t, s) q(s) f(y(s)) ds \leq \gamma \left(\int_a^b G(s, s) q(s) ds \right) r_2 = \|y\|$$

for $y \in K \cap \partial\Omega_2$. Thus, from Lemma 5, we conclude that the operator T defined by (8) has a fixed point in $K \cap (\overline{\Omega}_2 \setminus \Omega_1)$. Therefore, the fractional boundary problem (5) has at least one positive solution y belonging to X such that $r_1 \leq \|y\| \leq r_2$. \square

Example 9. Consider the following fractional boundary value problem:

$$\begin{cases} {}_0D^{3/2}y + te^y = 0 & \text{if } 0 < t < 1, \\ y(0) = y(1) = 0. \end{cases} \quad (9)$$

Firstly, let us calculate the values of γ and γ^* . Here,

$$\varphi(s) = \begin{cases} \frac{\sqrt{\frac{2(1-s)}{3}} - \sqrt{\frac{2}{3} - s}}{\sqrt{s(1-s)}} & \text{if } s \in (0, \lambda), \\ \frac{1}{\sqrt{3s}} & \text{if } s \in [\lambda, 1), \end{cases}$$

where $\lambda \simeq 0.64645$. Hence, by a simple computation, we get

$$\gamma^* \simeq 26.459 \text{ and } \gamma \simeq 4.514. \quad (10)$$

We choose $r_1 = \frac{1}{27}$ and $r_2 = 1$. Then we get

1. $f(y) = e^y \geq \gamma^* r_1$ for $y \in [0, \frac{1}{27}]$;
2. $f(y) = e^y \leq \gamma r_2$ for $y \in [0, 1]$.

Therefore, from Theorem 8, problem (9) has at least one nontrivial solution y in X such that $\frac{1}{27} \leq \|y\| \leq 1$.

3.2. Generalized Lyapunov's inequality

The next result generalizes Theorem 1: choosing $f(y) = y$ in Theorem 10, inequality (11) reduces to (4). Note that $f \in C(\mathbb{R}_+, \mathbb{R}_+)$ is a concave and nondecreasing function.

Theorem 10. *Let $q : [a, b] \rightarrow \mathbb{R}$ be a real nontrivial Lebesgue integrable function. Assume that $f \in C(\mathbb{R}_+, \mathbb{R}_+)$ is a concave and nondecreasing function. If the fractional boundary value problem (5) has a nontrivial solution y , then*

$$\int_a^b |q(t)| dt > \frac{4^{\alpha-1} \Gamma(\alpha) \eta}{(b-a)^{\alpha-1} f(\eta)}, \quad (11)$$

where $\eta = \max_{t \in [a, b]} y(t)$.

Proof. We begin by using Lemma 6. We have

$$\begin{aligned} |y(t)| &\leq \int_a^b G(t, s) |q(s)| f(y(s)) ds, \\ \|y\| &\leq \int_a^b G(s, s) |q(s)| f(y(s)) ds < \frac{(b-a)^{\alpha-1}}{4^{\alpha-1} \Gamma(\alpha)} \int_a^b |q(s)| f(y(s)) ds. \end{aligned}$$

Using Jensen's inequality (6), and taking into account that f is concave and nondecreasing, we get that

$$\|y\| < \frac{(b-a)^{\alpha-1} \|q\|_L}{4^{\alpha-1} \Gamma(\alpha)} \int_a^b \frac{|q(s)| f(y(s)) ds}{\|q\|_L} < \frac{(b-a)^{\alpha-1} \|q\|_L}{4^{\alpha-1} \Gamma(\alpha)} f(\eta),$$

where $\eta = \max_{t \in [a, b]} y(t)$. Thus,

$$\int_a^b |q(s)| ds > \frac{4^{\alpha-1} \Gamma(\alpha) \eta}{(b-a)^{\alpha-1} f(\eta)}.$$

This concludes the proof. \square

Corollary 11. *Consider the fractional boundary value problem*

$$\begin{cases} {}_a D^\alpha y + q(t) f(y) = 0, & a < t < b, \\ y(a) = y(b) = 0, \end{cases}$$

where $f \in C(\mathbb{R}_+, \mathbb{R}_+)$ is concave and nondecreasing and $q \in L([a, b], \mathbb{R}_+^*)$. If there exist two positive constants $r_2 > r_1 > 0$ such that $f(y) \geq \gamma^* r_1$ for $y \in [0, r_1]$ and $f(y) \leq \gamma r_2$ for $y \in [0, r_2]$, then

$$\int_a^b q(t) dt > \frac{4^{\alpha-1} \Gamma(\alpha) r_1}{(b-a)^{\alpha-1} f(r_2)}.$$

Example 12. Consider the following fractional boundary value problem:

$$\begin{cases} {}_0D^{3/2}y + t \ln(2 + y) = 0, & 0 < t < 1, \\ y(0) = y(1) = 0. \end{cases}$$

We have that

- (i) $f(y) = \ln(2 + y) : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is continuous, concave and nondecreasing;
- (ii) $q(t) = t : [0, 1] \rightarrow \mathbb{R}_+$ is a Lebesgue integral function with $\|q\|_L = 1 > 0$.

We computed the values of γ and γ^* in (10). Choosing $r_1 = 1/40$ and $r_2 = 1$, we get

1. $f(y) = \ln(2 + y) \geq \gamma^* r_1$ for $y \in [0, 1/40]$;
2. $f(y) = \ln(2 + y) \leq \gamma r_2$ for $y \in [0, 1]$.

Therefore, from Corollary 11, we get that

$$\int_0^1 q(t) dt > \frac{4^{\alpha-1} \Gamma(\alpha) r_1}{(b-a)^{\alpha-1} f(r_2)} \simeq 4.0334 \times 10^{-2}.$$

Acknowledgments

This research was initiated while Chidouh was visiting the Department of Mathematics of University of Aveiro, Portugal, 2015. The hospitality of the host institution and the financial support of Houari Boumedienne University, Algeria, are here gratefully acknowledged. Torres was supported through CIDMA and the Portuguese Foundation for Science and Technology (FCT), within project UID/MAT/04106/2013. The authors are grateful to Rachid Bebbouchi for putting them in touch. They are also very grateful to two anonymous referees for their careful reading of the submitted manuscript and for their valuable comments and questions.

References

- [1] F. M. Atici and G. Sh. Guseinov, On the existence of positive solutions for nonlinear differential equations with periodic boundary conditions, *J. Comput. Appl. Math.* **132** (2001), no. 2, 341–356.
- [2] Z. Bai and H. Lü, Positive solutions for boundary value problem of nonlinear fractional differential equation, *J. Math. Anal. Appl.* **311** (2005), no. 2, 495–505.
- [3] R. A. C. Ferreira, A Lyapunov-type inequality for a fractional boundary value problem, *Fract. Calc. Appl. Anal.* **16** (2013), no. 4, 978–984.

- [4] R. A. C. Ferreira, On a Lyapunov-type inequality and the zeros of a certain Mittag-Leffler function, *J. Math. Anal. Appl.* **412** (2014), no. 2, 1058–1063.
- [5] D. J. Guo and V. Lakshmikantham, *Nonlinear problems in abstract cones*, Notes and Reports in Mathematics in Science and Engineering, 5, Academic Press, Boston, MA, 1988.
- [6] M. Hashizume, Minimization problem related to a Lyapunov inequality, *J. Math. Anal. Appl.* **432** (2015), no. 1, 517–530.
- [7] M. Jleli, L. Ragoub and B. Samet, A Lyapunov-type inequality for a fractional differential equation under a Robin boundary condition, *J. Funct. Spaces* **2015** (2015), Art. ID 468536, 5 pp.
- [8] M. Jleli and B. Samet, Lyapunov-type inequalities for a fractional differential equation with mixed boundary conditions, *Math. Inequal. Appl.* **18** (2015), no. 2, 443–451.
- [9] M. Jleli and B. Samet, Lyapunov-type inequalities for fractional boundary-value problems, *Electron. J. Differential Equations* **2015** (2015), no. 88, 11 pp.
- [10] N. Li and C. Wang, New existence results of positive solution for a class of nonlinear fractional differential equations, *Acta Math. Sci. Ser. B Engl. Ed.* **33** (2013), no. 3, 847–854.
- [11] A. Liapounoff, Problème général de la stabilité du mouvement, *Ann. Fac. Sci. Toulouse Sci. Math. Sci. Phys. (2)* **9** (1907), 203–474.
- [12] D. O'Regan and B. Samet, Lyapunov-type inequalities for a class of fractional differential equations, *J. Inequal. Appl.* **2015** (2015), no. 247, 10 pp.
- [13] J. Rong and C. Bai, Lyapunov-type inequality for a fractional differential equation with fractional boundary conditions, *Adv. Difference Equ.* **2015** (2015), no. 82, 10 pp.
- [14] W. Rudin, *Real and complex analysis*, third edition, McGraw-Hill, New York, 1987.
- [15] S. G. Samko, A. A. Kilbas and O. I. Marichev, *Fractional integrals and derivatives*, translated from the 1987 Russian original, Gordon and Breach, Yverdon, 1993.
- [16] T. Sun and J. Liu, Lyapunov inequality for dynamic equation with order $n + 1$ on time scales, *J. Dyn. Syst. Geom. Theor.* **13** (2015), no. 1, 95–101.
- [17] X. Zhang, L. Wang and Q. Sun, Existence of positive solutions for a class of nonlinear fractional differential equations with integral boundary conditions and a parameter, *Appl. Math. Comput.* **226** (2014), 708–718.