

AN UPPER BOUND FOR THE SIZE OF A k -UNIFORM INTERSECTING FAMILY WITH COVERING NUMBER k .

ANDRII ARMAN AND TROY RETTER

ABSTRACT. Let $r(k)$ denote the maximum number of edges in a k -uniform intersecting family with covering number k . Erdős and Lovász proved that $\lfloor k!(e-1) \rfloor \leq r(k) \leq k^k$. Frankl, Ota, and Tokushige improved the lower bound to $r(k) \geq (k/2)^{k-1}$, and Tuza improved the upper bound to $r(k) \leq (1 - e^{-1} + o(1))k^k$. We establish that $r(k) \leq (1 + o(1))k^{k-1}$.

1. INTRODUCTION

Let X be a finite set and k be a positive integer. A family of sets $\mathcal{F} \subseteq \binom{X}{k}$ is called a k -uniform hypergraph, or a k -uniform family. The hypergraph \mathcal{F} is intersecting if all $e_1, e_2 \in \mathcal{F}$ satisfy $e_1 \cap e_2 \neq \emptyset$. A set $C \subseteq X$ is called a cover of \mathcal{F} if every $f \in \mathcal{F}$ satisfies $f \cap C \neq \emptyset$. The covering number of \mathcal{F} , denoted by $\tau(\mathcal{F})$, is defined by $\tau(\mathcal{F}) := \min\{|C| : C \text{ is a cover of } \mathcal{F}\}$.

Define

$$r(k) := \max\{|\mathcal{F}| : \mathcal{F} \text{ is } k\text{-uniform, intersecting, and } \tau(\mathcal{F}) = k\},$$

where no restriction is placed upon the size of the vertex set X .

In 1975, Erdős and Lovász [2] proved that

$$\lfloor k!(e-1) \rfloor \leq r(k) \leq k^k.$$

In 1994, Tuza [3] improved the upper bound, and in 1996, Frankl, Ota, and Tokushige [1] improved the lower bound. It follows from these results that

$$\left(\frac{k}{2}\right)^{k-1} \leq r(k) \leq (1 - e^{-1} + o(1))k^k.$$

Our main result is an improved upper bound. This will be established by using the following two lemmas, which will be proved in Sections 3 and 4, respectively. The first lemma is based upon the degree of a vertex $x \in X$, denoted $d(x)$, which is the number of elements in \mathcal{F} that contain x .

Lemma 1. *Let \mathcal{F} be a k -uniform intersecting family on X with covering number k . If $x \in X$ satisfies $d(x) \geq (\log k)k^{k-2}$, then*

$$|\{f \in \mathcal{F} : f \not\ni x\}| = o(k^{k-1}).$$

The next lemma is based upon the maximum degree of a hypergraph \mathcal{F} on X , which is defined by $\Delta(\mathcal{F}) := \max\{d(x) : x \in X\}$.

Lemma 2. *Let \mathcal{F} be a k -uniform intersecting family on X with covering number k . Let $\alpha \in \mathbb{R}^+$. If $\Delta(\mathcal{F}) \leq |\mathcal{F}|/40\alpha \log k$, then for k sufficiently large*

$$|\mathcal{F}| \leq \max\{2k^{2k/3}, ek^{k-\alpha}\}.$$

Together, these two lemmas will be used to prove our main result.

Theorem 3. *The function $r(k)$ satisfies*

$$r(k) \leq (1 + o(1))k^{k-1}.$$

Proof. Let \mathcal{F} be a k -uniform intersecting family on X with covering number k . We consider two cases.

If $\Delta(\mathcal{F}) \geq (\log k)k^{k-2}$, let $x \in X$ be a vertex with $d(x) \geq (\log k)k^{k-2}$. A simple observation (which follows from Lemma 5), is that any k -uniform intersecting family \mathcal{F} with covering number k satisfies $\Delta(\mathcal{F}) \leq k^{k-1}$. From this observation and Lemma 1,

$$|\mathcal{F}| \leq d(x) + |\{f \in \mathcal{F} : f \not\ni x\}| \leq k^{k-1} + o(k^{k-1}),$$

as desired.

In the complementary case $\Delta(\mathcal{F}) < (\log k)k^{k-2}$, we proceed by contradiction. That is, assume that $\Delta(\mathcal{F}) < (\log k)k^{k-2}$ and that $|\mathcal{F}| > k^{k-1}$. For $\alpha = k/40 \log^2 k$, we have that

$$\Delta(\mathcal{F}) < (\log k)k^{k-2} \leq \frac{|\mathcal{F}|}{40\alpha \log k},$$

and hence Lemma 2 gives that for k sufficiently large

$$|\mathcal{F}| \leq \max\{2k^{2k/3}, ek^{k-\alpha}\} < k^{k-1},$$

completing the proof. □

Lemmas 1 and 2 will be established in Sections 3 and 4, respectively. The next section will introduce some notation, a pair of general lemmas, and a Guesser-Chooser game upon which the proofs of Lemmas 1 and 2 will be based.

2. PRELIMINARIES

We will use the following notation. For $\tilde{\mathcal{F}} \subseteq \mathcal{F}$ and $S \subset X$, the degree of S in $\tilde{\mathcal{F}}$, denoted by $d_{\tilde{\mathcal{F}}}(S)$, is defined by $d_{\tilde{\mathcal{F}}}(S) := |\{f \in \tilde{\mathcal{F}} : f \supseteq S\}|$. We also take $d(S) := d_{\mathcal{F}}(S)$. For integers i and j , let $[i] := \{1, 2, \dots, i\}$, let $[i, j] := [j] \setminus [i-1]$, and let $(i, j] := [j] \setminus [i]$. We write $(\log k - i)$ to stand for $(\log(k) - i)$.

We begin by establishing the following lemma.

Lemma 4. *Let \mathcal{F} be a k -uniform intersecting family on X with covering number k and let $\tilde{\mathcal{F}} \subseteq \mathcal{F}$. Let $j \in [k]$ and let S_{j-1} be any subset of X with size $|S_{j-1}| = j - 1$. Then there exists $S_j = \{s_j\} \cup S_{j-1}$ with $|S_j| = j$ such that*

$$d_{\tilde{\mathcal{F}}}(S_j) \geq k^{-1} \cdot d_{\tilde{\mathcal{F}}}(S_{j-1}).$$

Proof. Since \mathcal{F} has covering number k and $|S_{j-1}| < k$, there is an edge $f \in \mathcal{F}$ such that $f \cap S_{j-1} = \emptyset$. Because \mathcal{F} is an intersecting family,

$$\sum_{x \in f} d_{\tilde{\mathcal{F}}}(S_{j-1} \cup x) \geq d_{\tilde{\mathcal{F}}}(S_{j-1}).$$

Therefore, for some $\tilde{x} \in f$, we have $d_{\tilde{\mathcal{F}}}(S_{j-1} \cup \tilde{x}) \geq k^{-1} \cdot d_{\tilde{\mathcal{F}}}(S_{j-1})$. It suffices to take $s_j := \tilde{x}$. \square

We will also make use of the following lemma.

Lemma 5. *Let \mathcal{F} be a k -uniform intersecting family on X with covering number k . If $U \subset X$ with $|U| = u$, then $d(U) \leq k^{k-u}$.*

Proof. We induct on u . If $u = k$, then $d(U) \leq 1$. For $u < k$, choose $f \in \mathcal{F}$ such that $f \cap U = \emptyset$; such an edge exists since $\tau(\mathcal{F}) = k > u$. Making use of the fact that every edge containing U must intersect f and our inductive hypothesis,

$$d(U) \leq \sum_{x \in f} d(U \cup \{x\}) \leq k \cdot k^{k-(u+1)} = k^{k-u},$$

completing the proof. \square

For $U = \emptyset$, this yields

$$r(k) \leq k^k, \tag{1}$$

as first proved by Erdős and Lovász in [2]. We now give another proof of (1) in order to introduce some of the key ideas and notation that will be used in the proofs of Lemmas 1 and 2.

Guesser-Chooser proof of equation (1). We consider a game played between a Chooser and a Guesser. The game is played on a fixed hypergraph \mathcal{F} , which is known to both players. The Chooser selects an edge $e \in \mathcal{F}$ which is not revealed to Guesser. Guesser then asks a sequence of questions $\Omega_1, \Omega_2, \dots, \Omega_k$ to gain information about the edge e . Each question Ω_i must have a unique answer $\omega_i \in [k]$. If Guesser can always determine the edge e after asking k such questions, it follows that $|\mathcal{F}| \leq k^k$. Equivalently, this can be thought of as creating an injection from \mathcal{F} to the set of all sequences of the form $\omega_1, \omega_2, \dots, \omega_k$ where $\omega_i \in [k]$ for all $i \in [k]$.

We remark that in this game, the questions Guesser asks may depend on the answers to the previous questions, but can not depend on knowledge of the edge e that is not available to Guesser.

We now describe such a k question strategy for Guesser. Guesser first selects an arbitrary edge $e_1 \in \mathcal{F}$ and fixes an arbitrary labeling $e_1 = \{e_1^1, e_2^1, \dots, e_k^1\}$. Question Ω_1 asks for the least ω_1 such that $e_{\omega_1}^1 \in e$; indeed, since \mathcal{F} is a k -uniform intersecting family, there is a unique answer $\omega_1 \in [k]$. Hence, our first question identifies one vertex $e_{\omega_1}^1 \in e$.

More generally, question Ω_i is determined as follows. Guesser selects an edge $e_i \in \mathcal{F}$ that does not intersect $\{e_{\omega_1}^1, e_{\omega_2}^2, \dots, e_{\omega_{i-1}}^{i-1}\}$, which exists since $\tau(\mathcal{F}) = k$. Guesser then fixes an arbitrary labeling $e_i = \{e_1^i, e_2^i, \dots, e_k^i\}$. Question Ω_i asks for the least ω_i such that $e_{\omega_i}^i \in e$. Hence, after k questions are asked, Guesser has determined $e = \{e_{\omega_1}^1, e_{\omega_2}^2, \dots, e_{\omega_k}^k\}$. \square

3. PROOF OF LEMMA 1

Let \mathcal{F} be a k -uniform intersecting family on X with $\tau(\mathcal{F}) = k$. Let $x \in X$ with $d(x) \geq (\log k)k^{k-2}$. Let

$$t := \lfloor \log k \rfloor.$$

To show $|\{f \in \mathcal{F} : f \not\ni x\}| \leq k^{k-1}$, we will make use of the Guesser-Chooser game introduced in the Guesser-Chooser proof of Equation (1). Chooser will select an edge $e \in \mathcal{F}$ with $e \not\ni x$ and then Guesser will ask a sequence of k questions $\Omega_1, \Omega_2, \dots, \Omega_k$ that will yield corresponding answers $\omega_1, \omega_2, \dots, \omega_k$ with $\omega_i \in [k]$ for all $i \in [k]$. Unlike the previous proof, Guesser will now choose his questions so that the first t answers form a non-decreasing sequence, that is

$$\omega_1 \leq \omega_2 \leq \dots \leq \omega_t.$$

The key idea to our proof is that for $i \in [t]$, after having asked questions $\Omega_1, \Omega_2, \dots, \Omega_i$ and received answers $\omega_1, \omega_2, \dots, \omega_i$, Guesser will have determined

- a set $V_i \subset e$ with $|V_i| = i$,
- a set $U_i \subset X \setminus e$ with $|U_i| = \omega_i - 1$, and
- a collection of edges $\mathcal{F}_i := \{f \in \mathcal{F} : f \supseteq U_i \text{ and } f \cap V_i = \emptyset\}$ with

$$|\mathcal{F}_i| \geq (\log k - i)k^{k-\omega_i}. \quad (2)$$

We will say that the sets V_i and U_i exhibit property \mathcal{P}_i if all three of the above criteria are satisfied. Let $V_0 := \emptyset$, let $U_0 := \{x\}$, and let $\omega_0 := 2$. It follows that $|\mathcal{F}_0| = d(x) \geq (\log k)k^{k-2}$. Observe that Guesser knows that V_0 and U_0 exhibit property \mathcal{P}_0 .

Claim 6. *Let $i \in [t]$. Given sets V_{i-1} and U_{i-1} exhibiting \mathcal{P}_{i-1} , Guesser can ask a question Ω_i whose answer ω_i will determine sets V_i and U_i exhibiting property \mathcal{P}_i . Moreover, Guesser can guarantee that $\omega_i \geq \omega_{i-1}$.*

Proof. We will specify an edge $e_i = \{e_1^i, e_2^i, \dots, e_k^i\}$. Question Ω_i will then ask for the least ω_i such that $e_{\omega_i}^i \in e$.

Fix a labeling $U_{i-1} = \{u_1, \dots, u_{\omega_{i-1}-1}\}$. For $j \in [\omega_{i-1} - 1]$, take $e_j^i := u_j$. This will ensure that $\omega_i \geq \omega_{i-1}$ as desired, since $U_{i-1} \cap e = \emptyset$.

Let $S_{\omega_{i-1}}^i := U_{i-1}$. We now proceed recursively as follows: for $j \in [\omega_{i-1}, k]$, apply Lemma 4 to S_{j-1}^i with respect to \mathcal{F}_{i-1} to obtain $S_j^i = S_{j-1}^i \cup \{e_j^i\}$. For $j \in [\omega_{i-1}, k]$, this yields sets S_j^i with

$$d_{\mathcal{F}_{i-1}}(S_j^i) \geq k^{-j+\omega_{i-1}-1} d_{\mathcal{F}_{i-1}}(S_{\omega_{i-1}}^i) = k^{-j+\omega_{i-1}-1} |\mathcal{F}_{i-1}|. \quad (3)$$

By (2),

$$k^{-j+\omega_{i-1}-1} |\mathcal{F}_{i-1}| \geq (\log k - i + 1)k^{k-j-1}. \quad (4)$$

It follows from (3) and (4) that for $j \in [\omega_{i-1}, k]$,

$$d_{\mathcal{F}_{i-1}}(S_j^i) \geq (\log k - i + 1)k^{k-j-1}. \quad (5)$$

Now making use of $i \leq t$, from (5) we have that $d_{\mathcal{F}_{i-1}}(S_k^i) > 0$. From the definition of \mathcal{F}_{i-1} , it now follows that $S_k^i \cap V_{i-1} = \emptyset$. Hence, $e_j^i \notin V_{i-1}$ for all $j \in [k]$.

Having completed our construction of e_i , we now consider the answer ω_i to question Ω_i . Define

$$V_i := V_{i-1} \cup \{e_{\omega_i}^i\} \quad \text{and} \quad U_i := \{e_1^i, e_2^i, \dots, e_{\omega_i-1}^i\}.$$

Observe that \mathcal{F}_i is precisely the edges in \mathcal{F}_{i-1} that contain $U_i = S_{\omega_{i-1}}^i$ and do not contain $e_{\omega_i}^i$. Making use of (5) and Lemma 5,

$$\begin{aligned} |\mathcal{F}_i| &\geq d_{\mathcal{F}_{i-1}}(S_{\omega_{i-1}}^i) - d_{\mathcal{F}}(S_{\omega_i}^i \cup \{e_{\omega_i}^i\}) \\ &\geq (\log k - i + 1)k^{k-\omega_i} - k^{k-\omega_i} \\ &= (\log k - i)k^{k-\omega_i}. \end{aligned}$$

Thus, we have shown that V_i and U_i exhibit property \mathcal{P}_i . \square

It follows from Claim 6 that Guesser may ask questions, $\Omega_1, \Omega_2, \dots, \Omega_t$ that necessitate a non-decreasing sequence of answers $\omega_1, \omega_2, \dots, \omega_t$. Moreover, after asking these questions, Guesser will have determined $V_t \subset e$ with $|V_t| = t$. For the remaining $k - t$ questions, Guesser will no longer ask questions that necessitate a non-decreasing sequence.

Claim 7. *Let $i \in (t, k]$. Given a set $V_{i-1} \subseteq e$ with $|V_{i-1}| = i - 1$, Guesser can ask a question Ω_i whose answer ω_i will allow Guesser to determine a set $V_i \subseteq e$ with $|V_i| = i$.*

Proof. Let e_i be any edge not covered by V_{i-1} ; such an edge exists since $\tau(\mathcal{F}) > i - 1$. Arbitrarily label $e_i = \{e_1^i, e_2^i, \dots, e_k^i\}$. Question Ω_i asks for the least ω_i such that $e_{\omega_i}^i \in e$. Let $V_i := V_{i-1} \cup \{e_{\omega_i}^i\}$. \square

Hence, after k questions are asked, Guesser will have determined $e = V_k$. Since the first t answers are non-decreasing and the number of non-decreasing sequences in $[k]^t$ is $\binom{k+t-1}{t}$, this gives that

$$\begin{aligned} |\{f \in \mathcal{F} : f \not\supseteq x\}| &\leq \binom{k+t-1}{t} k^{k-t} \\ &\leq \left(\frac{e(k+t-1)}{t}\right)^t k^{k-t} \\ &\leq \left(\frac{e}{t} \left(1 + \frac{t-1}{k}\right)\right)^t k^k \\ &\leq \left(\frac{2e}{t}\right)^t k^k \leq \left(\frac{2e}{\log k - 1}\right)^{\log k - 1} k^k \leq k^{k - (1 - o(1)) \log \log k} = o(k^{k-1}). \end{aligned}$$

This completes the proof of Lemma 1.

4. PROOF OF LEMMA 2

Let \mathcal{F} be a k -uniform intersecting family on X with $\tau(\mathcal{F}) = k$. Suppose that $\Delta(\mathcal{F}) \leq |\mathcal{F}|/40\alpha \log k$. To prove Lemma 2, it suffices to prove that if $|\mathcal{F}| > 2k^{2k/3}$, then $|\mathcal{F}| \leq ek^{k-\alpha}$. Hence, we assume that $|\mathcal{F}| > 2k^{2k/3}$.

Let

$$t := 20\lfloor \alpha \log k \rfloor. \quad (6)$$

As in the proofs of Equation (1) and Theorem 1, we will make use of the Guesser and Chooser game. As before, Chooser will select an edge $e \in \mathcal{F}$ and then Guesser will ask a sequence of k questions $\Omega_1, \Omega_2, \dots, \Omega_k$ that will yield corresponding answers $\omega_1, \omega_2, \dots, \omega_k$ with $\omega_i \in [k]$ for all $i \in [k]$. Unlike the previous two proofs, Guesser will now choose his questions so that

$$\omega_i > 2k/3 \implies \omega_{i+1} > k/3 \quad \text{for all odd } i \in [t]. \quad (7)$$

Let $V_0 := \emptyset$. The following claim establishes that Guesser can ask his first t questions so that (7) is satisfied.

Claim 8. *Let $i \in [t]$ be an odd number. Given a set $V_{i-1} \subset e$ with $|V_{i-1}| = i - 1$, Guesser can ask a pair of questions Ω_i and Ω_{i+1} whose answers will determine a set $V_{i+1} \subset e$ with $|V_{i+1}| = i + 1$. Moreover, these questions can be asked so that $\omega_i > 2k/3$ implies that $\omega_{i+1} > k/3$.*

Proof. Let $i \in [t]$ be an odd number and $V_{i-1} \subset e$ with $|V_{i-1}| = i - 1$. Let

$$\mathcal{F}_{i-1} := \{f \in \mathcal{F} : f \cap V_{i-1} = \emptyset\}.$$

It follows that

$$|\mathcal{F}_{i-1}| \geq |\mathcal{F}| - \Delta(\mathcal{F}) \cdot (i - 1) \geq |\mathcal{F}|/2 \geq k^{2k/3}.$$

We now construct a testing edge $e_i = \{e_1^i, e_2^i, \dots, e_k^i\}$. We begin by specifying the first $\lfloor k/3 \rfloor$ vertices in e_i as follows. Let $S_0^i := \emptyset$. We now proceed recursively: for $j \in [\lfloor k/3 \rfloor]$, apply Lemma 4 to S_{j-1}^i with respect to \mathcal{F}_{i-1} to obtain $S_j^i = S_{j-1}^i \cup \{e_j^i\}$. This yields sets S_j^i with

$$d_{\mathcal{F}_{i-1}}(S_j^i) \geq k^{2k/3-j}. \quad (8)$$

Having specified the first $\lfloor k/3 \rfloor$ vertices in e_{i+1} , we will now work to specify the remaining vertices. To this end, let $D_{\mathcal{F}_{i-1}}(S_{\lfloor k/3 \rfloor}^i) := \{f \in \mathcal{F}_{i-1} : f \supseteq S_{\lfloor k/3 \rfloor}^i\}$. Define

$$P_i := \{x \in X \setminus S_{\lfloor k/3 \rfloor}^i : x \in f \text{ for all } f \in D_{\mathcal{F}_{i-1}}(S_{\lfloor k/3 \rfloor}^i)\}. \quad (9)$$

It follows from (8), (9), and Lemma 5 that

$$k^{2k/3 - \lfloor k/3 \rfloor} \leq d_{\mathcal{F}_{i-1}}(S_{\lfloor k/3 \rfloor}^i) = d_{\mathcal{F}_{i-1}}(S_{\lfloor k/3 \rfloor}^i \cup P_i) \leq k^{k - \lfloor k/3 \rfloor - |P_i|}. \quad (10)$$

The inequality in (10) establishes that $|P_i| \leq k/3$. We now take e_{i+1} to be any edge in $D_{\mathcal{F}_{i-1}}(S_{\lfloor k/3 \rfloor}^i)$; such an edge is guaranteed to exist since the set is non-empty by (10). Label

$$e_i = \{e_1^i, e_2^i, \dots, e_k^i\}$$

so that $\{e_1^i, e_2^i, \dots, e_{\lfloor k/3 \rfloor}^i\} = S_{\lfloor k/3 \rfloor}^i$ and $P_i \subseteq \{e_{\lfloor k/3 \rfloor + 1}^i, e_{\lfloor k/3 \rfloor + 2}^i, \dots, e_{2k/3}^i\}$.

The question Ω_i asks for the least integer ω_i such that $e_{\omega_i}^i \in e$. Let $V_i := V_{i-1} \cup \{e_{\omega_i}^i\}$. We now consider two cases depending upon the answer ω_i .

If $\omega_i > 2k/3$, then Guesser must ensure that the answer ω_{i+1} to the next question will satisfy $\omega_{i+1} \geq k/3$. Observe that since $\omega_i > 2k/3$, it follows that $S_{\lfloor k/3 \rfloor}^i \cap e = \emptyset$. Also, since $\omega_i > 2k/3$, we have $e_{\omega_i}^i \notin P_i$. Hence, by the definition of P_i (see (9)), there exists an edge $e_{i+1} \in D_{\mathcal{F}_{i-1}}(S_{\lfloor k/3 \rfloor}^i)$ with $e_{i+1} \not\supseteq e_{\omega_i}^i$. Label the vertices of this edge

$$e_{i+1} = \{e_1^{i+1}, e_2^{i+1}, \dots, e_k^{i+1}\}$$

so that $\{e_1^{i+1}, e_2^{i+1}, \dots, e_{\lfloor k/3 \rfloor}^{i+1}\} = S_{\lfloor k/3 \rfloor}^i$. It follows that $e_{i+1} \cap V_i = \emptyset$. The answer to question Ω_{i+1} (based upon the testing edge e_{i+1}) will identify a new vertex in e and necessitate an answer $\omega_{i+1} \geq k/3$.

In the complementary case $\omega_{i+1} \leq 2k/3$, the question Ω_{i+1} must identify a new vertex in e and the answer ω_{i+1} can be any integer in $[k]$. To accomplish this, Guesser takes the testing edge e_{i+1} to be any edge that does not intersect V_i ; such an edge exists since $\tau(\mathcal{F}) = k$. \square

By Claim 6, Guesser may ask questions, $\Omega_1, \Omega_2, \dots, \Omega_t$ that necessitate a sequence of answers $\omega_1, \omega_2, \dots, \omega_t$ satisfying property (7). Moreover, after asking these questions, Guesser will have determined $V_t \subset e$ with $|V_t| = t$. For the remaining $k - t$ questions, Guesser will only require that each answer is a number in $[k]$ that identifies a new vertex in e . This is possible by Claim 7.

Hence, after k questions are asked, Guesser will have determined the edge e selected by Chooser. It follows that the size of $|\mathcal{F}|$ is bounded above by the number of sequence

$\omega_1, \omega_2, \dots, \omega_k \in [k]^k$ that satisfy property (7). Because the number of ways to select a pair $\omega_i, \omega_{i+1} \in [k]$ with the condition in (7) is less than

$$k^2 - (k/3 - 2)(k/3 - 2) = (8/9)k^2 + 4k/3 - 4 < e^{-1/10}k^2$$

for k sufficiently large, it follows that

$$|\mathcal{F}| \leq (e^{-1/10}k^2)^{t/2} k^{k-t} = e^{-t/20}k^k \leq ek^{k-\alpha}$$

for k sufficiently large. This completes the proof of Lemma 2.

5. ACKNOWLEDGEMENTS

We would like to thank Peter Frankl and Vojtěch Rödl for sharing this problem with us.

REFERENCES

- [1] P. Frankl, K. Ota, and N. Tokushige, *Covers in uniform intersecting families and a counterexample to a conjecture of Lovász*, J. Combin. Theory Ser. A **74** (1996), no. 1, 33–42. [↑1](#)
- [2] P. Erdős and L. Lovász, *Problems and results on 3-chromatic hypergraphs and some related questions*, Infinite and finite sets (Colloq., Keszthely, 1973; dedicated to P. Erdős on his 60th birthday), Vol. II, 1975, pp. 609–627. Colloq. Math. Soc. János Bolyai, Vol. 10. [↑1, 2](#)
- [3] Z. Tuza, *Inequalities for minimal covering sets in set systems of given rank*, Discrete Appl. Math. **51** (1994), no. 1-2, 187–195. 2nd Twente Workshop on Graphs and Combinatorial Optimization (Enschede, 1991). [↑1](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, MANITOBA R3T 2N2, CANADA
E-mail address: `armana@myumanitoba.ca`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, GA 30322,
 USA

E-mail address: `tretter@emory.edu`