

# Fault tolerant supergraphs with automorphisms

Ashwin Ganesan\*

## Abstract

Given a graph  $Y$  on  $n$  vertices and a desired level of fault-tolerance  $k$ , an objective in fault-tolerant system design is to construct a supergraph  $X$  on  $n + k$  vertices such that the removal of any  $k$  nodes from  $X$  leaves a graph containing  $Y$ . In order to reconfigure around faults when they occur, it is also required that any two subsets of  $k$  nodes of  $X$  are in the same orbit of the action of its automorphism group. In this paper, we prove that such a supergraph must be the complete graph. This implies that it is very expensive to have an interconnection network which is  $k$ -fault-tolerant and which also supports automorphic reconfiguration. Our work resolves an open problem in the literature. The proof uses a result due to Cameron on  $k$ -homogeneous groups.

**Index terms** — interconnection networks; fault-tolerant system design; structural fault tolerance; graph automorphisms; graph theory; permutation groups.

## 1. Introduction

The interconnection network of a computing system is modeled as a graph  $X = (V, E)$  whose vertices correspond to processors and with two vertices being adjacent whenever the corresponding two processors are connected by a direct communication link (cf. [14], [16]). In order to execute an algorithm on this computing system, it is required that the architecture  $X$  contain a given *basic graph*  $Y$  as a subgraph. If some of the nodes of  $X$  become faulty, in order to continue operation it is required that the functioning part of the network still contain the basic graph  $Y$ . We assume the basic graph  $Y$  is nonempty, i.e. it contains at least one edge. Any notation or terminology on graphs used in this paper which we do not explicitly define here is standard and can be found in [1].

Let  $Y$  be a nonempty graph on  $n$  vertices. A graph  $X$  is said to be a  *$k$ -fault-tolerant realization of  $Y$*  if  $X$  can be obtained from  $Y$  by adding a set of  $k$  new vertices (called *spare nodes*) and some edges so that the resulting graph  $X$  has the property that the removal of any  $k$  vertices from  $X$  leaves a graph which still contains  $Y$  (cf. [7]). In

---

\*School of Computer Science and Engineering, Vellore Institute of Technology (VIT University), Vellore, Tamilnadu 632014, India. Email: ashwin.ganesan@gmail.com.

other words,  $X$  is a  $k$ -fault-tolerant realization of  $Y$  if  $X$  has exactly  $n+k$  vertices and  $X - W$  contains a subgraph isomorphic to  $Y$  for each  $k$ -subset  $W \subseteq V(X)$ . In this case, if any  $k$  nodes of  $X$  become faulty, the network corresponding to the nonfaulty nodes of  $X$  contains the architecture  $Y$  and hence can continue to operate. In this sense, the architecture  $X$  can tolerate up to  $k$  node failures.

A graph  $X$  is a *supergraph* of  $Y$  if it is possible to add vertices and edges to  $Y$  to obtain  $X$ , i.e. if  $V(Y) \subseteq V(X)$  and  $E(Y) \subseteq E(X)$ . A graph  $X$  is an *edge-supergraph* of  $Y$  if  $V(X) = V(Y)$  and  $E(Y) \subseteq E(X)$ . In this paper, we consider the method of constructing supergraphs  $X$  of a given graph  $Y$  such that for all  $k$ -subsets  $F \subseteq V(X)$ ,  $X - F$  contains the subgraph  $Y$ . This type of design method is called *global sparing* because the  $k$  spare nodes added to  $Y$  are associated with all of  $V(Y)$ . In local sparing, we would partition  $V(Y)$  into  $t$  subsets  $V_1, \dots, V_t$  (for some  $t \geq 2$ ) and associate  $t$  sets of spare nodes to the  $t$  subsets  $V_i$ , respectively, such that  $k_i$  spare nodes are associated with  $V_i$  ( $i = 1, \dots, t$ ) and  $k_1 + \dots + k_t = k$ . Local sparing simplifies the design and reconfiguration process, while global sparing achieves  $k$ -fault-tolerance with fewer processors.

Many authors have investigated the use of algebraic methods in interconnection networks; see the surveys [15], [10] and the references therein. The present paper investigates an open problem posed in [7] on fault-tolerant supergraphs whose automorphism group satisfies certain properties. Another research area at the interface of graph automorphisms and interconnection networks is the study of the structure of interconnection network topologies; for example, several authors have investigated the automorphism group of graphs that arise as the topology of interconnection networks [5] [8] [9] [11] [12] [17]. More recently, researchers have questioned the practicability and advantages of interconnection networks with large automorphism groups; for example, data center interconnection networks are not hyperbolic [4].

For basic definitions of terms such as groups, symmetric groups, and homomorphisms, we refer the reader to [6, Chapter 1], [3]. Let  $G$  be a group and let  $\Omega$  be a nonempty set. Suppose the map  $\mu : \Omega \times G \rightarrow \Omega$ ,  $(\alpha, x) \mapsto \alpha^x$  satisfies the following two conditions: (i)  $(\alpha^x)^y = \alpha^{xy}$ , for all  $\alpha \in \Omega$  and all  $x, y \in G$ , and (ii)  $\alpha^1 = \alpha$  for all  $\alpha \in \Omega$ , where 1 denotes the identity element of the group  $G$ . Then, we say that this map defines an *action* of  $G$  on  $\Omega$  and that  $G$  *acts* on  $\Omega$ . This action naturally induces a homomorphism from  $G$  into the symmetric group  $\text{Sym}(\Omega)$ , and so each element of  $G$  induces a permutation of  $\Omega$ . Conversely, every homomorphism from  $G$  into  $\text{Sym}(\Omega)$  induces an action of  $G$  on  $\Omega$ . The orbit of a point  $\alpha \in \Omega$  under this action is the set  $\alpha^G := \{\alpha^x : x \in G\}$ . Thus, the action of  $G$  on  $\Omega$  partitions  $\Omega$  into orbits. The action of  $G$  on  $\Omega$  is *transitive* if for all  $\alpha, \beta \in \Omega$ , there exists a  $g \in G$  such that  $\alpha^g = \beta$ ; equivalently,  $G$  acts transitively on  $\Omega$  if the action of  $G$  on  $\Omega$  has a single orbit.

Suppose  $G$  acts on  $\Omega$ . The action of  $G$  on  $\Omega$  induces an action of  $G$  on the set of all subsets of  $\Omega$  by the rule  $\Gamma^x := \{\gamma^x : \gamma \in \Gamma\}$ , for all  $\Gamma \subseteq \Omega$ . It is clear that the set  $\Omega^{\{k\}}$  of all  $k$ -subsets of  $\Omega$  is  $G$ -invariant, i.e.  $(\Omega^{\{k\}})^x = \Omega^{\{k\}}$  for all  $x \in G$ . The group  $G$  is said to be  *$k$ -homogeneous* if  $G$  acts on  $\Omega^{\{k\}}$  transitively. (What we call  $k$ -homogeneous in this paper is referred to as  $k$ -subtransitive in [7].)

A  *$k$ -tuple* of distinct elements from  $\Omega$  is an ordered subset of  $k$  distinct elements from  $\Omega$ . For example, if  $\{\delta_1, \dots, \delta_k\}$  is a  $k$ -subset of  $\Omega$ , then  $(\delta_1, \dots, \delta_k)$  is a  $k$ -tuple of

distinct elements from  $\Omega$ . Let  $\Omega^{(k)}$  denote the set of all  $k$ -tuples of distinct elements from  $\Omega$ . We say that  $G$  is  $k$ -transitive if  $G$  acts transitively on  $\Omega^{(k)}$ . Thus, the action of  $G$  on  $\Omega$  is  $k$ -transitive iff for every two  $k$ -tuples  $(\alpha_1, \dots, \alpha_k), (\beta_1, \dots, \beta_k)$  of distinct elements from  $\Omega$ , there exists a  $g \in G$  such that  $\alpha_i^g = \beta_i$  ( $i = 1, \dots, k$ ). For further details on group actions, we refer the reader to [6, Chapter 1]; an introduction to multiply transitive groups and  $k$ -homogeneous groups can be found in [18, Chapter II] and [6, Sections 2.1 and 9.4].

Let  $X = (V, E)$  be a simple, undirected graph. Let  $\text{Sym}(V)$  denote the full symmetric group acting on the vertex set  $V = V(X)$ . Then,  $\text{Sym}(V)$  acts naturally on the set  $V^{\{2\}}$  of all 2-subsets of  $V$  by the following rule: for all  $x \in \text{Sym}(V)$  and for all  $\{u, v\} \in V^{\{2\}}$ ,  $\{u, v\}^x := \{u^x, v^x\}$ . An *automorphism* of the graph  $X = (V, E)$  is a permutation  $g \in \text{Sym}(V)$  which preserves adjacency and nonadjacency. In other words,  $g \in \text{Sym}(V)$  is an automorphism of  $X$  if  $\{x, y\} \in E$  iff  $\{x, y\}^g \in E$ . The set of all automorphisms of  $X$  forms a permutation group, called the *automorphism group of  $X$* , denoted by  $\text{Aut}(X)$ . Thus,  $\text{Aut}(X) := \{g \in \text{Sym}(V) : E^g = E\}$ . A graph  $X$  is said to be *vertex-transitive* if its automorphism group  $\text{Aut}(X)$  acts transitively on the vertex set  $V(X)$ . For an introduction to automorphisms of graphs, the reader is referred to [13].

Having stated our terminology on group actions and automorphisms of graphs, we can now describe an approach for restructuring around faults in an interconnection network. This approach, called *automorphic reconfiguration*, was introduced in [7], and refers to a specific type of reconfiguration using automorphisms in which the  $k$  spare nodes are directly mapped to the set of  $k$  faulty nodes. Automorphic reconfiguration, as defined in [7], is an impractical way of designing and reconfiguring graphs (and in particular, multiprocessor networks), and was a definition given just for theoretical purposes to finally lead to the type of  $k$ -fault-tolerant supergraphs designed in [7] that are not complete graphs.

In order to achieve so-called *automorphic reconfiguration* (cf. [7, p.253]), it is required that, when  $k$  or fewer nodes of the interconnection network become faulty, there exists an automorphism of the graph  $X$  that maps the spare nodes to the faulty nodes. During this reconfiguration process, the faulty nodes are relabeled as spare nodes, and the nonfaulty nodes are relabeled as nodes of  $Y$  and contain a subgraph isomorphic to  $Y$ . In graph-theoretic terms, the interconnection topology  $X$  must satisfy the property that if  $A$  and  $B$  are any two  $k$ -subsets of  $V(X)$ , then there is an automorphism of  $X$  that maps  $A$  to  $B$ . Equivalently, the interconnection network topology  $X$  must satisfy the property that its automorphism group  $\text{Aut}(X)$  is  $k$ -homogeneous.

Thus, our objective is the following: given a basic graph  $Y$  and a desired level of fault-tolerance  $k$ , construct a graph  $X$  such that  $X$  is a  $k$ -fault-tolerant realization of  $Y$  and such that  $\text{Aut}(X)$  is  $k$ -homogeneous. In other words, given a basic graph  $Y$ , we add  $k$  spare nodes  $S$  to  $V(Y)$  and edges to get a supergraph  $X$  such that for any  $k$ -subset  $F \subseteq V(X)$  of faulty nodes, there exists an automorphism  $g \in \text{Aut}(X)$  such that  $S^g = F$  and such that the set of nonfaulty nodes contains the subgraph  $Y$ . Dutt and Hayes settled this problem for the case  $k = 2$  by proving the following result:

**Theorem 1.** [7, Theorem 2] *If  $Y$  is a nonempty graph on  $n$  vertices,  $X$  is a 2-fault-tolerant realization of  $Y$  and  $\text{Aut}(X)$  is 2-homogeneous, then  $X$  is the complete graph  $K_{n+2}$ .*

Dutt and Hayes (cf. [7, p. 253]) posed the problem of generalizing the  $k = 2$  result of Theorem 1 to arbitrary  $k$ . In this paper, we resolve this open problem (cf. Theorem 2 below).

The following is the main result of this paper:

**Theorem 2.** *Let  $k \geq 2$ . If  $Y$  is a nonempty graph on  $n$  vertices,  $X$  is a  $k$ -fault-tolerant realization of  $Y$  and  $\text{Aut}(X)$  is  $k$ -homogeneous, then  $X$  is the complete graph  $K_{n+k}$ .*

We point out that if  $Y$  has  $n$  vertices and  $X$  is a  $k$ -fault-tolerant realization of  $Y$ , then  $X$  has exactly  $n + k$  vertices. The condition that  $X$  have exactly  $n + k$  vertices can be relaxed; see Corollary 10 and the remarks preceding it. One way to state the more general version of our main result is as follows: If  $Y$  is a nonempty graph, and  $X$  is such that  $\text{Aut}(X)$  is  $k$ -homogeneous for some  $k \geq 2$  and the removal of any  $k$  nodes from  $X$  leaves a graph containing  $Y$ , then  $X$  is a complete graph. We say “a” complete graph, rather than “the” complete graph because the order of  $X$  can be arbitrary. In most of this paper, we follow the statement of the open problem in [7, p. 253] and so assume that  $X$  has exactly  $n + k$  vertices.

## 2. Preliminaries

We recall some results on permutation groups.

**Lemma 3.** *Suppose  $G$  acts on  $\Omega$ . If  $G$  is  $k$ -transitive, then  $G$  is  $k$ -homogeneous.*

*Proof:* Let  $A = \{a_1, \dots, a_k\}$  and  $B = \{b_1, \dots, b_k\}$  be  $k$ -subsets of  $\Omega$ . To show  $G$  is  $k$ -homogeneous, it suffices to show there exists  $g \in G$  such that  $A^g = B$ . By  $k$ -transitivity of  $G$ , there exists  $g \in G$  such that  $(a_1, \dots, a_k)^g = (b_1, \dots, b_k)$ , i.e. there exists  $g \in G$  such that  $a_i^g = b_i$  ( $i = 1, \dots, k$ ). Hence,  $A^g = \{a_1^g, \dots, a_k^g\} = B$ . ■

**Lemma 4.** [6, p. 35] *Suppose  $G$  acts on  $\Omega$  and  $|\Omega| = n$ . Then,  $G$  is  $k$ -homogeneous iff  $G$  is  $(n - k)$ -homogeneous.*

*Proof:* Suppose  $G$  is  $k$ -homogeneous. To show  $G$  is  $(n - k)$ -homogeneous, let  $A, B \in \Omega^{\{n-k\}}$ . It suffices to show there exists a  $g \in G$  such that  $A^g = B$ . Let  $A' := \Omega - A$ ,  $B' := \Omega - B$ . Then,  $A', B' \in \Omega^{\{k\}}$ . By hypothesis, there exists a  $g \in G$  such that  $(A')^g = B'$ . Since  $g$  acts on  $\Omega$ ,  $g$  takes the complement of  $A'$  to the complement of  $B'$ , i.e.  $A^g = B$ . The converse is proved in a similar manner. ■

**Lemma 5.** *Suppose  $G$  acts on  $\Omega$ ,  $|\Omega| = n$  and  $2 \leq k \leq n$ . If  $G$  is  $k$ -transitive, then  $G$  is  $(k - 1)$ -transitive.*

*Proof:* Let  $\Delta = \{\delta_1, \dots, \delta_{k-1}\}$  and  $\Gamma = \{\gamma_1, \dots, \gamma_{k-1}\}$  be  $k$ -subsets of  $\Omega$ . Let  $\alpha \in \Omega - \Delta$  and  $\beta \in \Omega - \Gamma$ . By  $k$ -transitivity of  $G$ , there exists a  $g \in G$  such that  $(\delta_1, \dots, \delta_{k-1}, \alpha)^g = (\gamma_1, \dots, \gamma_{k-1}, \beta)$ . Hence, there exists a  $g \in G$  which takes the tuple  $(\delta_1, \dots, \delta_{k-1})$  to the tuple  $(\gamma_1, \dots, \gamma_{k-1})$ . ■

If  $|\Omega| = n$ , we write  $S_n$  for the full symmetric group  $\text{Sym}(\Omega)$ . The group of  $n!$  permutations in  $S_n$  acts naturally on the set  $[n] := \{1, \dots, n\}$ , and it is clear that for any two  $k$ -tuples  $\alpha = (\alpha_1, \dots, \alpha_k)$ ,  $\beta = (\beta_1, \dots, \beta_k)$  ( $k \leq n$ ), there exists a  $g \in S_n$  such that  $\alpha^g = \beta$ . Hence,  $S_n$  is  $k$ -transitive for every  $k \in [n]$ . By Lemma 3,  $S_n$  is also  $k$ -homogeneous for every  $k \in [n]$ . The next result shows some formal distinction between  $k$ -homogeneity and  $k$ -transitivity by exhibiting a permutation group (subgroup of  $S_n$ ) which is 2-homogeneous and not 2-transitive.

**Lemma 6.** [6, p. 286] *Consider the permutations  $x = (12 \dots 7)$ ,  $y = (235)(476)$  in  $S_7$ . Let  $G := \langle x, y \rangle$  be the permutation group in  $S_7$  generated by  $x$  and  $y$ . Then,  $G$  is not 2-transitive and  $G$  is 2-homogeneous.*

*Proof:* Observe that  $xy = yx^2$  and so  $|G| = 21$ . If the action of a group on  $[7]$  is 2-transitive, then the group must contain at least 42 elements. Because the order of  $G$  is 21,  $G$  is not 2-transitive.

To show  $G$  is 2-homogeneous, suppose  $\{\alpha, \beta\} \in [7]^{\{2\}}$ . Then, there exists  $z \in \langle x \rangle$  such that  $\{\alpha, \beta\}^z = \{1, \gamma\}$  for some  $\gamma$ . If  $\gamma \in \{2, 3, 5\}$ , then  $\{1, \gamma\}^{y^i} = \{1, 2\}$  for some  $i$ . If  $\gamma \in \{4, 7, 6\}$ , then  $\{1, \gamma\}^{y^j} = \{1, 7\}$  for some  $j$ , and we know  $\{1, 7\}^x = \{1, 2\}$ . Thus, there exists  $g \in G$  such that  $\{\alpha, \beta\}^g = \{1, 2\}$ . Hence, the action of  $G$  on  $[7]^{\{2\}}$  has a single orbit. ■

Recall that if a group is  $k$ -transitive, then it is also  $(k-1)$ -transitive. In general, a group which is  $k$ -homogeneous is not necessarily  $(k-1)$ -homogeneous. It will follow from our main result that if a permutation group  $G$  arises as the automorphism group of a graph, then  $k$ -homogeneity of  $G$  does imply  $(k-1)$ -homogeneity of  $G$ .

The proof of the main result uses the following result due to Cameron:

**Theorem 7.** [2, Theorem 2.2] [6, Theorem 9.4A] *Let  $G$  be a permutation group acting on a set  $\Omega$ . Let  $m, k$  be integers with  $0 \leq m \leq k$  and  $m+k \leq |\Omega|$ . Then,  $G$  has at least as many orbits in  $\Omega^{\{k\}}$  as it has in  $\Omega^{\{m\}}$ .*

### 3. Main results

We first extend Theorem 1 from the 2-homogeneous case to the 3-homogeneous case. This result is actually a special case of the main result (Theorem 2). We now give an elementary proof for the 3-homogeneous case which does not use the theory of permutation groups. In fact, the condition that  $\text{Aut}(X)$  be 3-homogeneous can be replaced by the weaker condition that every subset of 3 vertices of the graph induces the same subgraph.

**Theorem 8.** *If  $X$  is a graph on 5 or more vertices containing at least one edge and if every subset of 3 vertices of  $X$  induces the same subgraph, then  $X$  is a complete graph.*

*Proof:* Let  $A = \{a, b, c\} \subseteq V(X)$  and let  $X'$  denote the induced subgraph  $X[A]$ . Because  $\text{Aut}(X)$  is 3-homogeneous, every 3-subset of  $V(X)$  induces the same subgraph  $X'$ . By hypothesis,  $X'$  contains at least one edge. We consider three cases for the structure of  $X'$ . For the first case, suppose  $X'$  is a 3-clique. Then,  $X$  is the complete graph because every subset of 3 vertices of  $X$  induces a 3-clique. For the second case, suppose  $X'$  is isomorphic to  $K_{1,2}$ . Without loss of generality, suppose  $ab, bc \in E(X)$ ,  $ac \notin E(X)$ . Let  $x, w \in V(X) - A$ . Then,  $\{a, c, w\}$  and  $\{a, c, x\}$  each induce a  $K_{1,2}$ . Since  $a$  and  $c$  are nonadjacent, it must be that  $ax, cx, aw, cw \in E(X)$ . The subgraph induced by  $\{b, x, w\}$  must also be a  $K_{1,2}$  and hence contains at least one edge. The endpoints of this edge along with vertex  $a$  induce a  $K_3$ , a contradiction. Hence, the second case is impossible.

For the third case, suppose the induced subgraph  $X'$  is isomorphic to the disjoint union of  $K_2$  and  $K_1$ . Then, the complement graph  $\overline{X'}$  is isomorphic to  $K_{1,2}$ . Since the automorphism group of a graph and of its complement are equal,  $\text{Aut}(\overline{X'})$  is also 3-homogeneous, and by the second case above,  $\overline{X'}$  is the complete graph. But this implies  $X$  is the empty graph, contradicting the fact that its induced subgraph  $X'$  contains an edge. Hence, this third case is also impossible. ■

The lower bound of 5 in Theorem 8 is tight since the automorphism group of  $C_4$  is 3-homogeneous and  $C_4$  is not a clique.

We now prove the main result.

*Proof of Theorem 2:* The case  $k = 2$  is addressed in Theorem 1, so assume  $k \geq 3$ . Let  $Y$  be a graph on  $n$  vertices. Here,  $n \geq 2$  since  $Y$  contains at least one edge. Note that  $X$  is a graph on  $n + k$  vertices. Let  $G = \text{Aut}(X)$  and let  $\Omega = V(X)$ . By hypothesis, the action of  $G$  on the set  $\Omega$  is  $k$ -homogeneous. Thus, the number of orbits of  $G$  on  $\Omega^{\{k\}}$  is 1. Since  $2 \leq n$ ,  $2 + k \leq n + k = |\Omega|$ . Also,  $2 \leq k$ . Hence, by Theorem 7, the number of orbits of  $G$  on  $\Omega^{\{2\}}$  is also 1. Equivalently, the action of  $G$  on  $\Omega$  is 2-homogeneous.

Let  $\{u, v\}$  be an edge in  $Y$ . Then  $\{u, v\}$  is an edge in  $X$ . Let  $a$  and  $b$  be distinct vertices of  $X$ . Because  $G$  is 2-homogeneous, there is an element  $g \in G$  that maps  $\{u, v\}$  to  $\{a, b\}$ . The automorphism  $g$  preserves adjacency, whence  $\{a, b\}$  is an edge of  $X$ . This proves that any two distinct vertices in  $X$  are adjacent, i.e.  $X$  is the complete graph  $K_{n+k}$ . ■

In the proof above, we essentially showed the following:

**Corollary 9.** *Let  $k \geq 2$ . Let  $X$  be a nonempty graph on  $k + 2$  vertices. If  $\text{Aut}(X)$  is  $k$ -homogeneous, then  $X$  is the complete graph  $K_{k+2}$ .*

As stated, Theorem 2 assumes that the order  $|V(X)|$  of  $X$  is exactly equal to  $n + k$ . The proof of the theorem goes through even if this condition is relaxed to  $|V(X)| \geq n + k$ :

**Corollary 10.** *Let  $X$  be a nonempty graph. If  $\text{Aut}(X)$  is  $k$ -homogeneous for some  $2 \leq k \leq |V(X)| - 2$ , then  $\text{Aut}(X)$  is 2-homogeneous and  $X$  is a complete graph.*

We resolve some further open questions in the literature. In [7, p. 252], it is stated that there is likely no characterization of  $k$ -homogeneous automorphism groups of

graphs in the literature. In [7, p. 252], it is mentioned that  $k$ -homogeneity of  $\text{Aut}(X)$  could be a “significantly weaker restriction” than  $k$ -transitivity of  $\text{Aut}(X)$ . Also, in [7, p.253], it is mentioned that  $k$ -homogeneity of  $\text{Aut}(X)$  does not necessarily imply  $i$ -homogeneity of  $\text{Aut}(X)$  for  $2 \leq i \leq k - 1$ . All these questions are resolved by the following immediate consequence of our main result Theorem 2:

**Corollary 11.** *Let  $X$  be a graph on 4 or more vertices. Suppose  $2 \leq k \leq |V(X)| - 2$ . If  $\text{Aut}(X)$  is  $k$ -homogeneous, then  $\text{Aut}(X)$  is  $i$ -homogeneous for  $i = 1, 2, \dots, k - 1$  and  $\text{Aut}(X)$  is the full symmetric group. In particular,  $k$ -homogeneity of  $\text{Aut}(X)$  is equivalent to  $k$ -transitivity of  $\text{Aut}(X)$ .*

The bounds in  $2 \leq k \leq |V(X)| - 2$  are tight - there are many families of vertex-transitive graphs for which  $\text{Aut}(X)$  is  $k$ -homogeneous for  $k = 1$  (and hence also for  $k = |V(X)| - 1$  by Lemma 4, and obviously for  $k = |V(X)|$ ) and such that  $\text{Aut}(X)$  is not  $i$ -homogeneous for  $i = 2, \dots, |V(X)| - 2$ . For example, take  $X$  to be a cycle graph.

Theorem 2 makes two assumptions: that  $X$  is a  $k$ -fault-tolerant realization of  $Y$  and that  $\text{Aut}(X)$  is  $k$ -homogeneous. The conclusion of the theorem follows mainly from the second assumption and only a weak consequence of the first assumption is used. To illustrate, we consider the following example of a graph  $X$  which is a 2-fault-tolerant realization of  $Y = Q_3$ , the 3-dimensional cube. As this example shows, the first assumption alone is not sufficient to ensure that  $X$  is the complete graph.

**Theorem 12.** *Let  $Y$  be the 3-dimensional hypercube  $Q_3$ . Let  $X$  be the graph obtained by adding to  $Y$  two spare nodes  $x_1$  and  $x_2$  and joining each  $x_i$  ( $i = 1, 2$ ) to each vertex of  $Y$ . Then,  $X$  is a 2-fault-tolerant realization of  $Y$ , but  $\text{Aut}(X)$  is not  $k$ -homogeneous for  $k = 1, 2, \dots, |V(X)| - 1$ .*

*Proof:* Observe that if any two diametrically opposite vertices of the 3-cube  $Q_3$  are removed, the resulting graph is a 6-cycle graph; see Figure1(a). Adding two new vertices to this 6-cycle graph and joining these two vertices to each vertex of the 6-cycle gives an edge-supergraph of  $Q_3$ . Hence, if a graph contains a 6-cycle, then adding two new vertices  $x_1, x_2$  to the graph and joining each  $x_i$  to each vertex of the graph gives an edge-supergraph of  $Q_3$ .

To prove that  $X$  is a 2-fault-tolerant realization of  $Y = Q_3$ , let  $F = \{f_1, f_2\}$  be a set of two faulty nodes of  $X$ . We need to show that  $X - F$  contains a subgraph isomorphic to  $Q_3$ . This is clear if  $\{f_1, f_2\} = \{x_1, x_2\}$  and also if exactly one of the faulty nodes is in  $Y$  because this node can be replaced by the non-faulty spare node. So suppose now that both faulty nodes  $f_1$  and  $f_2$  are in  $Y$ . If  $f_1$  and  $f_2$  are nonadjacent in  $Y$ , then they can be replaced by the spare nodes  $x_1$  and  $x_2$ , respectively, and  $Y - F$  contains a subgraph isomorphic to  $Q_3$ . Finally, suppose  $f_1$  and  $f_2$  are adjacent nodes of  $Y$ . Then  $Y - F$  contains a 6-cycle (see Figure 1(b)), and by the argument in the previous paragraph,  $X - F$  contains a subgraph isomorphic to  $Q_3$ .

In the graph  $X$ , the degree of vertex  $x_i$  ( $i = 1, 2$ ) is 8, and the degree of each of the remaining vertices is 4. Hence,  $X$  is not vertex-transitive (i.e.  $\text{Aut}(X)$  is not 1-homogeneous). By Lemma 4,  $\text{Aut}(X)$  is not  $(|V(X)| - 1)$ -homogeneous. If




Figure 1: (a) Removing diametrically opposite vertices from a cube  $Q_3$  gives a 6-cycle  $y_1y_2 \dots y_6$ . (b) Removing two adjacent vertices gives a graph containing a 6-cycle  $y_1y_2 \dots y_6$ .

$\text{Aut}(X)$  is  $k$ -homogeneous for some  $2 \leq k \leq |V(X)| - 2$ , then by Theorem 2  $X$  is the complete graph, a contradiction. Hence,  $\text{Aut}(X)$  is not  $k$ -homogeneous if  $k \in \{1, 2, \dots, |V(X)| - 1\}$ .  $\blacksquare$

The paper [7] designed an iterative reconfiguration technique which after  $k$  faults occur uses  $k$  different automorphisms in a repeated manner to obtain a fault-free graph isomorphic to the basic graph  $Y$  from its  $k$ -fault-tolerant supergraph  $X$  that does not require  $X$  to be a complete graph, and in fact is quite efficient in the additional edges needed in  $X$  with respect to  $Y$  (cf. [7, Theorems 5 and 6]) as well as in the switch-based implementation of  $X$ . Both Theorem 12 and the technique in [7] are evidence that  $k$ -fault-tolerance of  $X$  is not a sufficient condition for  $X$  to be complete, and further, in the case of the technique in [7] the use of automorphisms to obtain  $k$ -fault-tolerance does not require  $X$  to be complete (and thus does not require  $\text{Aut}(X)$  to be  $k$ -homogeneous).

Theorem 1 and Theorem 2 imply that it is very expensive to have an interconnection network which is  $k$ -fault-tolerant and which also supports automorphic reconfiguration (i.e. for  $\text{Aut}(X)$  of the  $k$ -fault-tolerant graph  $X$  to also be  $k$ -homogeneous) because such an interconnection network must be the complete graph.

## Acknowledgements

Thanks are due to the anonymous reviewers for helpful suggestions.

## References

- [1] B. Bollobás. *Modern Graph Theory*. Graduate Texts in Mathematics vol. 184, Springer, New York, 1998.
- [2] P. J. Cameron. Transitivity of permutation groups on unordered sets. *Mathematische Zeitschrift*, 148:127–139, 1976.
- [3] P. J. Cameron. *Permutation Groups*. London Mathematical Society Student Texts 45, Cambridge University Press, 1999.



- [4] D. Coudert and G. Ducoffe. Data center interconnection networks are not hyperbolic. *Theoretical Computer Science*, 639:72–90, 2016.
- [5] Y.-P. Deng and X.-D. Zhang. Automorphism groups of the pancake graphs. *Information Processing Letters*, 112:264–266, 2012.
- [6] J. D. Dixon and B. Mortimer. *Permutation Groups*. Graduate Texts in Mathematics vol. 163, Springer, 1996.
- [7] S. Dutt and J. P. Hayes. Designing fault-tolerant systems using graph automorphisms. *Journal of Parallel and Distributed Computing*, 12:249–268, 1991.
- [8] Y-Q. Feng. Automorphism groups of Cayley graphs on symmetric groups with generating transposition sets. *Journal of Combinatorial Theory Series B*, 96:67–72, 2006.
- [9] A. Ganesan. Automorphism groups of Cayley graphs generated by connected transposition sets. *Discrete Mathematics*, 313:2482–2485, 2013.
- [10] A. Ganesan. Cayley graphs and symmetric interconnection networks. In *Proceedings of the Pre-Conference Workshop on Algebraic and Applied Combinatorics (31st Annual Conference of the Ramanujan Mathematical Society), Trichy, Tamilnadu, India*, pages 118–170, 2016.
- [11] A. Ganesan. Edge-transitivity of Cayley graphs generated by transpositions. *Discussiones Mathematicae Graph Theory*, 36:1035–1042, 2016.
- [12] A. Ganesan. Automorphism group of the complete transposition graph. *Journal of Algebraic Combinatorics*, 42(3):793–801, November 2015.
- [13] C. Godsil and G. Royle. *Algebraic Graph Theory*. Graduate Texts in Mathematics vol. 207, Springer, New York, 2001.
- [14] J. P. Hayes. A graph model for fault-tolerant computing systems. *IEEE Transactions on Computers*, C-25(9):875–884, 1976.
- [15] M. C. Heydemann. Cayley graphs and interconnection networks. In *Graph symmetry: algebraic methods and applications (Editors: Hahn and Sabidussi)*, pages 167–226. Kluwer Academic Publishers, Dordrecht, 1997.
- [16] F. T. Leighton. *Introduction to parallel algorithms and architectures: arrays, trees, hypercubes*. Morgan Kaufmann Publishers, 1992.
- [17] M. Wan and Z. Zhang. A kind of conditional vertex-connectivity of star graphs. *Applied Mathematics Letters*, 22:264–267, 2009.
- [18] H. Wielandt. *Finite Permutation Groups*. Academic Press, 1964.