

Lovász' original lower bound: Getting tighter bounds and Reducing computational complexity

Shuchita Goyal* and Rekha Santhanam

Abstract

In this article, we give conditions on a graph under which the Lovász' original bound of the graph can be improved by increasing the topological connectivity of its neighbourhood complex. We also work out conditions under which computing the topological connectivity of hom complex of a pair of graphs can be simplified. In particular, hom complex as a covariant functor acting on a double mapping cylinder of graphs is a homotopy pushout of hom complex functor applied to its subgraphs. We give applications of this result where the computation of hom complexes is simplified. Finally, we explain why double mapping cylinder of graphs does not give a satisfactory definition of homotopy pushout in the category of graphs.

1 Introduction

For a graph G , the question of determining the chromatic number of G has been of central importance in graph theory. In the year 1955, Kneser formulated a class of graphs, popularly known as Kneser graphs, $KG_{n,k}$, and conjectured that the chromatic number of $KG_{n,k}$ is $n - 2k + 2$. This question remained open for almost 25 years until 1978 when Lovász came up with a topological space associated to a graph G , called the neighbourhood complex, $\mathcal{N}(G)$, of that graph. His proof opened a wide area of mathematics, topological combinatorics. Lovász proved the Kneser's conjecture using topological properties of $\mathcal{N}(KG_{n,k})$. In particular, he showed that $\chi(G)$ is bounded from below by the topological connectivity of $\mathcal{N}(G)$ plus 3, for every graph G . This bound is often referred to as the Lovász' original bound. Generalizing this idea, he also conjectured that for any two graphs G, T , the chromatic number of the graph G is bounded from below as per the following inequality:

$$\chi(G) \geq \chi(T) + \text{connHom}(T, G) + 1. \quad (1)$$

A graph, T , for which this inequality holds with respect to every graph G is called a test graph. In view of Kozlov's result in [12] which states that $\text{Hom}(K_2, G)$ is homotopy equivalent to $\mathcal{N}(G)$, Lovász result [14] showed that K_2 is a test graph. It has been shown that the class of complete graphs [3], odd cycles [2], bipartite graphs [15], special Kneser graphs [18], etc, are all test graphs. Lovász conjecture has been disproved by Hoory and Linial [11] who exhibited explicit graphs G and T such that the inequality (1) fails to satisfy.

In order to apply inequality (1), we need to be able to compute the connectivity of the hom complex. Some special cases have been studied [6, 7, 17]. This computation, however, in general is extremely difficult. For instance, for n greater than 3, computing whether $\text{Hom}(T, K_n)$ is path connected is a PSPACE-complete problem [4].

*The author would like to thank CSIR, India and IRCC, IIT Bombay for their financial support.

We begin this article with methods to improve Lovász' original lower bound for chromatic number by replacing the given graph G with a smaller graph. More precisely, we give conditions on a degree 2 vertex v of a graph G so that $\chi(G-v) = \chi(G)$, and the connectivity of the neighbourhood complex of $G - v$ is greater than or equal to that of the neighbourhood complex of G . We also analyse the conditions on a degree 3 vertex v for which the connectivity of the neighbourhood complex of $G - v$ is greater than or equal to the connectivity of the neighbourhood complex of G .

Next we study pushouts in graphs and their behaviour with respect to the $\text{Hom}(T, _)$ functor for any graph T . It is a quick check that in general pushouts are not preserved by the hom functor. It turns out though that double mapping cylinders of graphs are preserved under $\text{Hom}(T, _)$ functor. For a fixed homotopy test graph T , and given any m large enough, we use the properties of the double mapping cylinder to construct a graph G whose chromatic number is m and the topological connectivity of the hom complex, $\text{Hom}(T, G)$, is 0. We also compute the homotopy type of neighbourhood complexes of a certain class of graphs.

Based on our applications it is useful to know when a given graph is \times -homotopy equivalent to the double mapping cylinder of some smaller graphs. Understanding double mapping cylinder of graphs as their homotopy pushout is then vital. In Section 4, we point out why double mapping cylinder of graphs cannot be a homotopy pushout object in the category of graphs. In a following article, we study this notion further.

2 Improving Lovász' original bound

A graph G is a pair of sets $(V(G), E(G))$. The elements of $V(G)$ are called *vertices* of G , and the elements of $E(G)$, which are two element subsets of $V(G)$, are called *edges* of G . These elements need not be distinct, that is, $\{y, y\}$ can also be an edge and this makes y into a *looped* vertex. We say that two vertices x, y are adjacent if $\{x, y\} \in E(G)$ and denote the edge by xy . A *simple graph* is a graph without any looped vertex. A *complete graph* on n vertices is a simple graph where any two distinct vertices are adjacent to each other, and is denoted by K_n . For $n \in \mathbb{N}$, let I_n be the graph with vertex set $\{0, 1, \dots, n\}$ and edge set $\{ij : |i - j| \leq 1\}$.

A *homomorphism* between two graphs $f : G \rightarrow H$ is a function from $V(G)$ to $V(H)$ such that if xy is an edge in G , then $f(x)f(y)$ is an edge in H . The *chromatic number* for any simple graph G is defined to the least integer n such that there is a graph homomorphism from G into K_n .

Definition 2.1. Let $v, v' \in G$ and $v \neq v'$. Then v is said to *fold* to $v' \in G$ if every neighbour of v is also a neighbour of v' . In this case, we say that G *folds* to $G - v$. A graph G is called *stiff* if there does not exist any $v \in G$ such that G folds to $G - v$.

Definition 2.2. Let $G, H \in \mathcal{G}$ be two graphs. The (categorical) *product* of G and H is defined to be the graph $G \times H$ whose vertex set is the cartesian product, $V(G) \times V(H)$, and $(g, h)(g', h') \in E(G \times H)$ whenever $gg' \in E(G)$, $hh' \in E(H)$.

Let \mathcal{G} be the category with objects as finite undirected graphs without multiple edges and morphisms as vertex set maps that takes edges to edges.

Definition 2.3 ([13]). For two arbitrary graphs T and G , the *hom complex*, $\text{Hom}(T, G)$, is the polyhedral complex whose cells are indexed by all the functions $\eta : V(T) \rightarrow 2^{V(G)} \setminus \{\emptyset\}$ such that $\eta(x) \times \eta(y) \subset E(G)$, whenever $xy \in E(T)$. The *closure* of a cell η consists of all cells η' , satisfying $\eta'(v) \subset \eta(v)$, for all $v \in V(G)$.

Throughout this paper, we will use the notation of polytopes to mean cells of $\text{Hom}(T, G)$. For a polytope $\eta \in \text{Hom}(T, G)$, the image of η is defined to be, $\text{Im}(\eta) := \cup_{t \in V(T)} \eta(t)$. In this section, we always consider connected loopless graphs unless stated otherwise.

For a given graph G and a vertex $v \in V(G)$, we let $N_G(v)$ denote the set of neighbours of v in G , and for a set $A \subset V(G)$, $N_G(A) = \bigcup_{a \in A} N_G(v)$. Whenever there is no scope of confusion, the subscript from the notation of neighbourhood set is dropped. *Neighbourhood complex* of graph G , $\mathcal{N}(G)$, is the simplicial complex whose simplices are formed by those subsets of vertex set of G that have a common neighbour.

By definition of neighbourhood complex of a graph G , two vertices $u, v \in V(G)$ form a 1-simplex, denoted by $\{u, v\}$, in $\mathcal{N}(G)$ if and only if u and v have a common neighbour in G . That is, there exists a path of length 2 connecting u to v in G . Therefore, two vertices lie in the same path connected component of $\mathcal{N}(G)$ if and only if there exists an even length path in G connecting these two vertices. Following this idea, it is clear that all the vertices of an odd cycle of G belong to the same path connected component of $\mathcal{N}(G)$.

Lemma 2.4. [14] *A loopless connected graph G is bipartite if and only if $\mathcal{N}(G)$ is disconnected.*

Let G be any graph and $v \in V(G)$. The *degree of vertex v* is defined as the cardinality of neighbourhood set of v , $|N_G(v)|$ if v is not looped and $|N_G(v)| + 1$ otherwise. The degree of v is denoted as $d(v)$.

Theorem 2.5. *Let G be a simple non-bipartite graph with C_p , a cycle in G and $c \in C_p$ be such that $d(c) = 2$, $b \in N_G(c)$. Then $\pi_1(\mathcal{N}(G - bc), c)$ is a subgroup of $\pi_1(\mathcal{N}(G), c)$.*

In order to prove Theorem 2.5, we first establish some results about homotopy groups of simplicial complexes.

For a topological space X , $\pi_0(X)$ denotes the path connected components of X . Let X and Y be two topological spaces and $f, g : X \rightarrow Y$ be two continuous functions. The map f is said to be homotopic to g if there exists a continuous function $F : X \times [0, 1] \rightarrow Y$ satisfying $F|_{X \times \{0\}} = f$ and $F|_{X \times \{1\}} = g$. Let \mathbb{S}^n be the n -dimensional sphere, then it is the boundary of $(n + 1)$ -dimensional disc, \mathbb{D}^{n+1} . For a topological space X with a base point x , the n^{th} -homotopy group $\pi_n(X, x)$ is defined to be the collection of homotopy classes of maps from $\mathbb{S}^n \rightarrow X$. One often writes $[X, Y]$ to mean the homotopy class of maps from the space X to Y . If a continuous function $f : \mathbb{S}^n \rightarrow X$ can be extended continuously to $\tilde{f} : \mathbb{D}^{n+1} \rightarrow X$ then f is called homotopically trivial. The n^{th} -homotopy group of X is trivial if every $f : \mathbb{S}^n \rightarrow X$ is homotopically trivial. The space X is called k -connected if $\pi_n(X, x)$ is a trivial group for all $0 \leq n \leq k$ and the connectivity of X , $\text{conn}X$ is the largest k such that X is k -connected.

Let S be a geometric simplicial complex. We define a 1-chain in S is a finite sequence of 0-simplices $v_0 v_1 \dots v_s$ such that $\{v_i, v_{i+1}\}$ is a 1-simplex in S for all $i = 0, \dots, s - 1$. A 1-chain is closed if $v_0 = v_s$ and such closed 1-chains are also referred to as loops based at v_0 .

A constant loop based at v_0 is a sequence $v_0 v_1 \dots v_s$ such that $v_i = v_0$ for all $i = 0, \dots, s$. A loop based at v_0 is said to be homotopically trivial in S if it is homotopic to the constant loop based at v_0 . We define a closed 1-chain as simple if v_0 occurs exactly once, that is, $v_i \neq v_0$ for $0 < i < s$.

For a simplicial complex S , the n -skeleton, S_n , of S is the subcomplex of S that contains all k -simplices of S , for $k \leq n$.

Definition 2.6. *Let S and T be two geometric simplicial complexes. A continuous map $f : S \rightarrow T$ is called a simplicial map if for any k -simplex $\sigma = \{s_0, s_1, \dots, s_k\}$ of S , $f(\sigma)$ is a simplex $\{f(s_0), f(s_1), \dots, f(s_k)\}$ of T .*

For an n -simplex $\sigma = \{x_0, x_1, \dots, x_n\}$ of a simplicial complex X , we define a point $x \in \sigma$ to be an interior point of σ if x is not one of the x_i , for $i = 0, 1, \dots, n$. A point $x \in X$ is called an interior point of X if it is an interior point of some simplex of X .

Lemma 2.7. *Let S be a geometric simplicial complex and $\alpha : \mathbb{S}^1 \rightarrow S$ be a closed nullhomotopic 1-chain in S . If the extension $\sigma : \mathbb{D}^2 \rightarrow S$, of α , is such that $Im(\sigma) \subset S_1$, that is, σ is contained in the 1-skeleton of S , then every interior point of α is repeated at least once.*

Proof. We prove it by contradiction. Considering the 2-ball \mathbb{D}^2 as a simplicial complex, by simplicial approximation theorem, there exists a simplicial map $\sigma_s : Y \rightarrow S$, where Y denotes the n -th barycentric subdivision of \mathbb{D}^2 for a large enough n . Then, σ_s is homotopic to σ , and $\sigma_s|_{bd(Y)}$ gives a simplicial map $\alpha_s : bd(Y) \rightarrow S$ such that α_s is homotopic to α .

Since S is a simplicial complex, the intersection of any two simplices of S is again a simplex of S and a face of both the intersecting simplices. Therefore, if no 1-simplex of α_s is repeated, then at least three consecutive 1-simplices of σ_s are unrepeated. Without any loss of generality, we can assume that α_s contains a subdivision of a hollow triangle, and hence σ_s contains a subdivision of a hollow triangle. Let η be a minimal length closed 1-chain in $Im(\sigma_s)$ containing a subdivision of a hollow triangle.

Since σ_s is a simplicial map, image of any two intersecting simplices under σ_s intersects. Let A_1 be the inverse image of η under σ_s , then A_1 contains intersecting 2-simplices that form an annulus like structure (please refer to Figure 1). As a simplicial complex in itself, A_1 has a boundary that does not enclose any 2-simplex of A_1 , say α_1 . Let A_2 be the collection of 2-simplices of $Y - A_1$




Figure 1: The shaded region in (a) denotes an annulus like structure A_1 described in the proof of Lemma 2.7, with dark line illustrating the boundary α_1 . The figure in (b) shows the next iteration where A_2 is obtained from A_1 of (a), with darkest line showing the boundary α_2 .

with a facet belonging to α_1 . For any 2-simplex $\tau \in A_2$, the value of σ_s on its facet determines the value of τ , and σ_s being a simplicial map implies that $\sigma_s(\tau) \in \eta$. Note that A_2 also forms an annulus like structure with possibly few pinches along the two boundaries. Let α_2 be the boundary of A_2 away from the boundary of Y . We repeat the same argument for A_2 , and so on. Since Y is a finite simplicial complex, the process terminates after finitely many iterations with a 2-simplex with two of its facets being mapped to different 1-simplices of η , a contradiction that σ_s is a simplicial map. \square

Similar arguments can be used to prove a more general statement, that is:

Proposition 2.8. *Let S be a geometric simplicial complex and $\alpha : \mathbb{S}^n \rightarrow S$ be a nullhomotopic continuous map such that for any extension $\sigma : \mathbb{D}^{n+1} \rightarrow S$, of α , $Im(\sigma) \subset S_n$, that is, σ is contained in the n -skeleton of S , then every interior point of facets in α is repeated at least once.*

Proof of Theorem 2.5. By hypothesis, G is non-bipartite and hence $\mathcal{N}(G)$ is connected by Lemma 2.4. Let $N_G(c) = \{a, b\}$. If c belongs to a $C_4 = (a, c, b, d)$ in G , then $N_G(c) = \{a, b\} \subset N_G(d)$. Therefore c folds to d and $\mathcal{N}(G - c)$ is homotopy equivalent to $\mathcal{N}(G)$ and the result then follows.

We, therefore, assume that the vertex c does not fold in G and c does not belong to any C_4 in G . Let $N_G(a) = \{a_1, a_2, \dots, a_k, c\}$ and $N_G(b) = \{b_1, b_2, \dots, b_r, c\}$. Let A and B denote the neighbour set of a and b in G respectively. Since c is a degree 2 vertex that belongs to a cycle, its neighbours a and b also belong to the same cycle and hence $k, r \geq 1$. From now on, we drop the subscript G from N_G . Since $c \in V(G)$ does not fold in G , $(A \cap B) - \{c\} = \emptyset$.

Consider the graph $G' = G - bc$. Since G' is a subgraph of G , $\mathcal{N}(G')$ is a subcomplex of $\mathcal{N}(G)$. Let $f : \mathcal{N}(G') \rightarrow \mathcal{N}(G)$ be the inclusion map and $f_{\#} : \pi_1(\mathcal{N}(G'), c) \rightarrow \pi_1(\mathcal{N}(G), c)$ be the group homomorphism induced by f . To establish the statement, it suffices to prove that $f_{\#}$ is an injective homomorphism.

Let $\alpha \in \pi_1(\mathcal{N}(G'), c)$ be a closed simple 1-chain such that $f_{\#}(\alpha)$ is homotopically trivial in $\mathcal{N}(G)$. Then $\alpha = cv_1v_2 \dots v_t c$ is a sequence of vertices of G' such that the first and the last vertex is c and any two consecutive vertices form a 1-simplex in $\mathcal{N}(G')$ and hence have a common neighbour in G' . Since in $\mathcal{N}(G')$, the vertex c forms 1-simplex only with neighbours of a , namely, $\{a_1, a_2, \dots, a_k\}$, we can see that $v_1, v_t \in \{a_1, a_2, \dots, a_k\}$.

Using simplicial approximation, a closed 1-chain in a simplicial complex X can be interpreted as a simplicial map from a triangulation of \mathbb{S}^1 to X . Given that $\alpha : \mathbb{S}^1 \rightarrow \mathcal{N}(G)$ is homotopically trivial in $\mathcal{N}(G)$, there exists an extension $\sigma_{\alpha} : \mathbb{D}^2 \rightarrow \mathcal{N}(G)$, that is $bd(\sigma_{\alpha}) = \alpha$. In the context of simplicial complex, the subcomplex $\sigma_{\alpha} \subset \mathcal{N}(G)$ gives a triangulation of \mathbb{D}^2 . If σ_{α} is a subcomplex of $\mathcal{N}(G')$ then it gives an extension of α to \mathbb{D}^2 in $\mathcal{N}(G')$ and α is homotopically trivial in $\mathcal{N}(G')$ and there is nothing to prove. Also, if σ_{α} belongs to 1-skeleton of $\mathcal{N}(G)$, then the result follows from Proposition 2.8. So without loss of generality, assume that σ_{α} contains a 2-simplex $\{c, b_1, b_2\}$

Note that any 2-simplex of $\mathcal{N}(G)$ not containing c belongs to $\mathcal{N}(G')$. If the 2-simplex $\{c, b_1, b_2\}$ belongs to the interior of σ_{α} , then there exists a closed 1-chain η homotopic to α in $\mathcal{N}(G')$ such that boundary of σ_{η} is η , and η contains the 1-simplex $\{b_1, b_2\}$. Therefore, we assume that if a simple closed 1-chain $\alpha \subset \mathcal{N}(G')$ is homotopically trivial in $\mathcal{N}(G)$ and σ_{α} contains a 2-simplex $\{c, b_1, b_2\}$, then the 1-simplex $\{b_1, b_2\}$ belongs to α .




Figure 2: An example to illustrate the proof of the Theorem 2.5. In (a), α is shown in black as the boundary of σ_{α} . In (b), α_1 is shown in black as obtained from α such that σ_{α_1} is contained in σ_{α} . Similarly (c) records the next step. Lastly, (d) shows the final iteration, where we ought to have $\{c, b_i, a_1\}$ as a 2-simplex shown by a dotted line.

Let i be the smallest integer such that $v_i \in \{b_1, b_2, \dots, b_k\}$, so that the simple closed 1-chain $\alpha = ca_1v_2v_3 \dots v_{i-1}v_iv_{i+1} \dots v_{t-1}a_2c$ with $v_i = b_1$, say, is homotopically trivial in $\mathcal{N}(G)$ but not in $\mathcal{N}(G')$. We will show that this is not possible by exhibiting a simple closed 1-chain in σ_{α} which is not homotopically trivial in $\mathcal{N}(G)$.

Take $\alpha_1 = ca_1v_2 \dots v_{i-1}b_1c$, then α_1 is a subcomplex of σ_{α} and hence also homotopically

trivial in $\mathcal{N}(G)$. Let σ_{α_1} denote the extension of α_1 to \mathbb{D}^2 . Note that $N(A) \cap N(B) = \emptyset$, thus $\{c, a_i, b_j\}$ cannot be a 2-simplex in the neighbourhood complex of G . If $\{c, b_1, x\}$ is a 2-simplex, then $x \in \{b_1, \dots, b_k\}$. Therefore α being homotopically trivial implies that there exists $b_{i_1} \in \{b_1, \dots, b_k\}$ such that 2-simplex $\{c, b_1, b_{i_1}\} \subset \sigma_{\alpha_1}$. Now consider $\alpha_2 = ca_1v_2 \dots b_1b_{i_1}c \subset \sigma_{\alpha_1}$, then α_2 is also homotopically trivial in $\mathcal{N}(G)$. Similar argument as above gives that there exists $b_{i_2} \in \{b_1, \dots, b_k\}$ such that 2-simplex $\{c, b_{i_1}, b_{i_2}\} \subset \sigma_{\alpha_2}$. Arguing similarly, we get $\alpha_j = ca_1v_2 \dots b_1b_{i_1}b_{i_2} \dots b_{i_j}c \subset \sigma_{\alpha_{j-1}} \subset \sigma_\alpha$ such that α_j is also homotopically trivial in $\mathcal{N}(G)$. Since G is finite and hence the set $\{b_1, \dots, b_k\}$ of neighbours of b , this process terminates in a finite number of steps. During this process, all the v_i 's gets replaced by elements from $A \cup B$ and we get a closed 1-chain $\{c, a_i, b_j\}$, which is not homotopically trivial in $\mathcal{N}(G)$, and that is a contradiction. Therefore, if α is homotopically trivial in $\mathcal{N}(G)$, then it is homotopically trivial in $\mathcal{N}(G')$ also. \square

Note that G' considered in the above theorem folds to $G - c$. Since c belongs to C_n in G , $G' = G - bc$ is connected for a degree 2 vertex c of G , if $\chi(G - c) > 2$, then $\chi(G) = \chi(G - c)$.

Corollary 2.9. *For a finite connected simple non-bipartite graph, if $c \in V(G)$ belongs to a cycle in G and $d(c) = 2$. Then*

$$\text{conn}\mathcal{N}(G - c) \geq \text{conn}\mathcal{N}(G).$$

Furthermore, if $G - c$ contains an odd cycle then $\chi(G - c) = \chi(G)$. Else, $\chi(G - c) = 2$ implying $\chi(G) = 3$.

Proof. To establish the claim, we need to only show that if $\pi_k(\mathcal{N}(G))$ is trivial for $k \geq 2$, then $\pi_k(\mathcal{N}(G - bc))$ is trivial. Let $\alpha : S^n \rightarrow \mathcal{N}(G - c)$ be a simplicial map for $n \geq 2$. Then by assumption we know it extends to a simplicial map $\alpha' : \mathbb{D}^{n+1} \rightarrow \mathcal{N}(G)$. Since we assume that a, b have only one vertex c in common, we know that the 1-simplex $\{a, b\}$ is a part of $\mathcal{N}(G)$ but not a face of higher boundary in $\mathcal{N}(G)$ and does not exist in $\mathcal{N}(G - bc)$. If image of α' includes this 1-simplex $\{a, b\}$, then by Proposition 2.8 we can see that we can get a new map $\alpha'' : \mathbb{D}^{n+1} \rightarrow \mathcal{N}(G) - \{a, b\}$. Now it is possible that $N_G(b)$ is in the image of α' . Let the simplex $c \cup N_{G-c}(b)$, be in the image of α' . Since image of α does not intersect with any simplex in $c \cup N_{G-c}(b)$ containing c , we can contract the simplices containing c down to $N_{G-c}(b)$. Thus defining an $\alpha'' : D^{n+1} \rightarrow \mathcal{N}(G - c)$. Hence proved. \square

Let G be a graph, then its minimum degree, $\delta(G)$, is defined to be the minimum of degrees taken over all the vertices of G . Let $\mathcal{C}^{(2)}$ be the class of graphs whose minimum degree is 2. Then Theorem 2.5 can be applied to any non-bipartite graph $G \in \mathcal{C}^{(2)}$. Note that G does not have any hanging (degree 1) vertices. Also, since the minimum degree, $\delta(G) = 2$, every vertex belongs to some cycle in the graph. Let $v \in V(G)$ be such that $d(v) = 2$. Since v belongs to a cycle, Theorem 2.5 applies to it, and we get the graph $G - v$ such that $\pi_1(N(G - v))$ is a subgroup of $\pi_1(N(G))$, and either $G - v$ is bipartite or $\chi(G - v) = \chi(G)$.

Let $(G, c) = (G_0, c_0)$ be a connected non-bipartite graph with vertex $c_0 \in V(G_0)$ of degree two that belongs to some cycle $C_p \subset G$. Define G_1 to be the subgraph $G_0 - \{c_0\}$, where $G_0 - \{c_0\}$ denotes the subgraph of G_0 induced by the vertex set $V(G_0) - \{c_0\}$. Then by Corollary 2.9, $\text{conn}\mathcal{N}(G_1) \geq \text{conn}\mathcal{N}(G_0)$. Since G_0 is non-bipartite, $\chi(G_0) \geq 3$. If G_1 does not contain any odd cycle then it is bipartite and hence has chromatic number 2, thereby implying that the chromatic number of G_0 is 3 and we stop. On the other hand, if G_1 contains an odd cycle, then again by Corollary 2.9, $\chi(G_1) = \chi(G_0)$. In this case, if G_1 contains a degree two vertex say c_1 , then we repeat as above to obtain a smaller graph $G_2 = G_1 - \{c_1\}$; and as before if G_2 is bipartite then $2 = \chi(G_2) = \chi(G_1) - 1 = \chi(G_0) - 1$ implying that $\chi(G_0) = 3$. Otherwise $\chi(G_2) = \chi(G_1) = \chi(G_0)$.

We continue removing degree two vertices from non-bipartite graph as explained above. Since the original graph G_0 is finite, the number of iterations is also finite. Suppose n vertices are removed during this iterative process, then using Lovász' bound, we have

$$\chi(G) = \chi(G_0) = \chi(G_n) \geq \text{conn}\mathcal{N}(G_n) + 3.$$

Since G_n is a much smaller graph compared to that of G_0 , computation of $\text{conn}\mathcal{N}(G_n)$ is expected to be easier than $\text{conn}\mathcal{N}(G)$.

The result in Theorem 2.5 deals with the fundamental group of the neighbourhood complex of the subgraph $G - c$ as compared to that of G . In the cases when $\pi_1(\mathcal{N}(G))$ is non-trivial but $\pi_1(\mathcal{N}(G - c))$ is, it improves the Lovász' lower bound by at most 1. In the same vein, we prove the next result where removing degree 3 vertices can improve the bound by 2 in good cases.

Theorem 2.10. *Let G be a non-bipartite graph with a degree three non-looped vertex $v \in V(G)$. Then*

1. *The second homotopy group $\pi_2(\mathcal{N}(G - v))$ is a subgroup of $\pi_2(\mathcal{N}(G))$.*
2. *If for every pair of two neighbours x, y of v , there exists an even length path joining x and y in $G - v$, then*

$$\pi_0(\mathcal{N}(G - v)) = \pi_0(\mathcal{N}(G)).$$

Furthermore, for some $x, y \in N_G(v)$, let $P = xp_1p_2 \dots p_{2n-1}y$ be a shortest distance path from x to y in $G - v$. If $n \geq 2$, then

$$\pi_1(\mathcal{N}(G - v)) \text{ is a subgroup of } \pi_1(\mathcal{N}(G)).$$

Proof. Let the neighbours, $N_G(v)$, of v be $\{a, b, c\}$. Without loss of generality, we can assume that $N_G(a) \cap N_G(b) \cap N_G(c) = \{v\}$, otherwise if there exists $x \in N_G(a) \cap N_G(b) \cap N_G(c)$, $x \neq v$, then v folds to x . By [12, 13], $\mathcal{N}(G - v)$ is homotopy equivalent to $\mathcal{N}(G)$, and hence the result holds.

Since $G - v$ is a subgraph of G , $\mathcal{N}(G - v)$ is a subcomplex of $\mathcal{N}(G)$. Let $f : \mathcal{N}(G - v) \rightarrow \mathcal{N}(G)$ denote this inclusion, and $f_{\#} : \pi_2(\mathcal{N}(G - v)) \rightarrow \pi_2(\mathcal{N}(G))$ be the induced group homomorphism.

Let $\alpha : \mathbb{S}^2 \rightarrow \mathcal{N}(G - v)$ be a continuous map such that $f_{\#}(\alpha) = f\alpha$ is nullhomotopic. To prove the first part, it suffices to show that α is nullhomotopic. Since $f\alpha : \mathbb{S}^2 \rightarrow \mathcal{N}(G)$ is nullhomotopic, there exists an extension $\sigma : \mathbb{D}^3 \rightarrow \mathcal{N}(G)$. The 3-ball \mathbb{D}^3 carries a simplicial complex structure and $\mathcal{N}(G)$ is a simplicial complex. Therefore, by simplicial approximation theorem, there exists a simplicial map $\sigma_s : Y \rightarrow \mathcal{N}(G)$, where Y denote the n -th barycentric subdivision of \mathbb{D}^3 for some large enough n . Let α_s be the restriction of σ_s to the boundary, $bd(Y)$, of Y .

If $Im(\sigma_s)$ is contained in $\mathcal{N}(G - v)$, then α is trivial in $\mathcal{N}(G - v)$. Therefore, assume the $Im(\sigma_s)$ contains a simplex of $\mathcal{N}(G)$ that is not in $\mathcal{N}(G - v)$. Also, if $Im(\sigma_s)$ is contained in the 2-skeleton of $\mathcal{N}(G)$, then the result follows from Proposition 2.8. So, assume that $Im(\sigma_s)$ contains a 3-simplex of $\mathcal{N}(G)$ that is not in $\mathcal{N}(G - v)$.

Let $A := N_G(a)$, $B := N_G(b)$, $C := N_G(c)$. If $\{v, x, y, z\}$ forms a 3-simplex of $\mathcal{N}(G)$ that is not in $\mathcal{N}(G - v)$, then all three x, y, z belong to either A , or B , or C . Since $\sigma|_{bd(\mathbb{D}^3)} = \alpha$ is contained in $\mathcal{N}(G - v)$, v is an interior point of $Im(\sigma)$. If the 3-simplex generated by $\{v, x, y, z\}$ lies in the interior of $Im(\sigma)$, then nullhomotopicity of $f\alpha$ in $\mathcal{N}(G)$ implies that there exists an $\eta : \mathbb{S}^2 \rightarrow \mathcal{N}(G)$ such that η is homotopic to α and the 2-simplex generated by $\{x, y, z\}$ lies on the boundary of its extension $\delta : \mathbb{D}^3 \rightarrow \mathcal{N}(G)$. Therefore, we may assume that $\{x, y, z\} \in bd(\sigma) = \alpha$. We now can use similar arguments as in the proof of Theorem 2.5 to complete the proof.

To prove the second part, note that whenever two vertices of a graph can be connected by a path of even length, the two vertices belong to the same path connected component of $\mathcal{N}(G)$. Therefore if there is a path of even length for every pair of two neighbours of v , then all three neighbours of v belong to the same path connected component of $\mathcal{N}(G - v)$, and hence the result.

For the rest of the proof, let α be a closed non-nullhomotopic 1-chain in $\mathcal{N}(G - v)$ containing a, b , and c as its 0-simplices. Let $\alpha = au_1u_2 \dots u_{k_1}bv_1v_2 \dots v_{k_2}cw_1w_2 \dots w_{k_3}a$, where $k_1, k_2, k_3 \geq 0$. Since for some $x, y \in N_G(v) = \{a, b, c\}$, the distance between x and y is more than 2, at least one of k_1, k_2, k_3 is greater than 0. Suppose α is trivial in $\mathcal{N}(G)$, then $\alpha_1 = au_1u_2 \dots u_{k_1}ba$, $\alpha_2 = bv_1v_2 \dots v_{k_2}cb$, and $\alpha_3 = cw_1w_2 \dots w_{k_3}ac$ are trivial 1-chains as $\{a, b, c\}$ is a 2-simplex in $\mathcal{N}(G)$. Let a, b be such that $P = ap_1p_2 \dots p_{2n-1}b$. If $n \geq 2$, there does not exist any 2-simplex in $\mathcal{N}(G)$ containing a, b . This implies that α_1 cannot be contracted to a point in $\mathcal{N}(G)$, a contradiction. Therefore, $\pi_1(\mathcal{N}(G - v))$ is a subgroup of $\pi_1(\mathcal{N}(G))$. \square

Remark 2.11. As before, we can show that if the n^{th} homotopy group of $\mathcal{N}(G)$ for some $n \geq 4$ is trivial, then after removing this degree 3 vertex c the $\pi_n(\mathcal{N}(G - c)) = 0$ for that n . Let $\mathcal{C}^{(3)}$ denote the class of graphs with minimum degree 3. Then for any non-bipartite graph $G \in \mathcal{C}^{(3)}$ satisfying hypothesis of Theorem 2.10 (2), we can reduce the graph G and obtain a smaller subgraph G' for which the computation of $\mathcal{N}(G')$ is easier as G' is strictly smaller than G , and $\text{conn}\mathcal{N}(G')$ is at least as much as $\text{conn}\mathcal{N}(G)$.

Remark 2.12. Removing degree 2 and 3 vertices is especially useful if they are obstructing the connectivity of $\mathcal{N}(G)$ to be higher.

For instance, consider the graph $G := C_n \sqcup_{K_2} K_m$ for any $n \geq 5$ and $m \geq 4$. Then G has a degree 2 vertex (coming from the C_n), say v . It can be checked that $\text{conn}\mathcal{N}(G) = 0$, whereas $\text{conn}\mathcal{N}(G - v) = m - 3 > 0$. And at the same time, $\chi(G - v) = \chi(G)$, improving the Lovász' bound. Thus, Theorem 2.5 helps improve the lower bound on the chromatic number of G .

Similarly a simple application of Theorem 2.10 is as follows. Take pushout of $K_3 \rightarrow K_m$ to get $K_m \sqcup_{K_3} K'_m$ for $m \geq 6$. Let $a \in V(K_m), b, c \in V(K'_m)$ be such that $a, b \notin V(K_m) \cap V(K'_m)$ and $c \in V(K_m) \cap V(K'_m)$. Take a new vertex x with neighbours a, b . Now take another vertex, say v such that v is adjacent to a, c and x . Let G be this graph. We first note that $\chi(G - v) = \chi(G)$. Also $\mathcal{N}(G)$ is 0-connected, whereas the topological connectivity of $\mathcal{N}(G - v)$ is at least 2 (cf. [5]).

However, it is possible to have graphs G with $\chi(G) = 4$ but $\chi(G - v) = 3$ and characterizing such graphs is difficult. In such cases it is not clear when this method will give us better bounds for $\chi(G)$ using inequality (1). This indicates that a degree 4 version of the above theorem may not be very fruitful.

Note that Theorem 2.10 can also be applied to graphs with a degree 3 vertex having the said properties even if the minimum degree of G is two or less, that is, this theorem applies to a general class of graphs.

3 Reducing computational complexity for hom complex

Let A, B and C be graphs and $f : A \rightarrow B$ and $g : A \rightarrow C$ be graph homomorphisms. In the category of graphs, the double mapping cylinder of height n is defined as $D_n = B \sqcup_f (A \times I_n) \sqcup_g C = (B \sqcup (A \times I_n) \sqcup C) / \sim$, where \sim denotes the equivalence relation $f(a) \sim (a, n)$ and $g(a) \sim (a, 0)$, and $[x][y] \in E(D_n)$ if there exists some $x_0 \in [x], y_0 \in [y]$ such that $\{x_0, y_0\} \in E(B) \cup E(A \times I_n) \cup E(C)$. We write x to denote its equivalence class $[x]$.




Figure 3: Double mapping cylinder.

Given any two graph homomorphisms f, g , we now examine the relation between the pushout object and the double mapping cylinder of f and g .


Lemma 3.1. *Let $A, B, C \in \mathcal{G}$ be three loopless graphs with graph homomorphisms $f : A \rightarrow B$ and $g : A \rightarrow C$. Let $G \in \mathcal{G}$ denote the pushout of f and g , and D_n be their double mapping cylinder of height n . If G is a simple graph, then*

$$\chi(G) \geq \chi(D_n).$$

Proof. For a loopless graph $X \in \mathcal{G}$, a k -colouring of X is a graph homomorphism from $X \rightarrow K_k$. Let X, Y be two loopless graphs, and $h : X \rightarrow Y$ be a graph homomorphism. Then it is easy to note that a k -colouring of $h(X)$ can be extended to a k -colouring of X .


Let $\chi(G) = m$ and $c : G \rightarrow K_m$ be a proper m -colouring of G . By definition of the pushout (quotient graph) G , $f(a)$ is identified with $g(a)$ for every $a \in V(A)$, therefore $c|_{f(A)} = c|_{g(A)}$. Then $c|_{f(A)} = c|_{g(A)}$ gives a proper colouring of $f(A)$ and $g(A)$ and hence a proper colouring, say \bar{c} of A . Define $c' : D_n \rightarrow K_m$ by letting $c'|_{B \cup C} = c|_{B \cup C}$ and $c'|_{A \times \{i\}} = \bar{c}$, for $i = 1, 2, \dots, n-1$. Clearly c' is a proper colouring of D_n , hence the claim. \square

The inequality in the other direction is not true in general. For example, consider the following graphs as A, B, C .



Let $f : A \rightarrow B$ and $g : A \rightarrow C$ be graph inclusions. Then the graph on the right is the pushout $G = B \sqcup_A C$ and has $\chi(G) = 5$. On the other hand, the double mapping cylinder $D_n = B \sqcup_f (A \times I_n) \sqcup_g C$ is 4-colourable for any $n > 1$.

This gives an example where $\chi(G)$ is strictly greater than $\chi(D_n)$.



The definition of double mapping cylinder of graphs is analogous to the one for topological spaces which we recall here.

Definition 3.2. Let X, Y, Z be topological spaces with $p : X \rightarrow Y$ and $r : X \rightarrow Z$ continuous maps. The double mapping cylinder of $\{p, r\}$ is defined to be the quotient space

$$Y \bigsqcup_p (X \times [0, 1]) \bigsqcup_r Z / \sim$$

where \sim is an equivalence relation generated by $(x, 1) \sim p(x)$ and $(x, 0) \sim r(x)$. We write $Y \bigsqcup_X^h Z$ to mean the double mapping cylinder of the maps $\{p, r\}$.

Note that the double mapping cylinders give homotopy pushouts in the category of topological spaces. In particular, given two homotopy equivalent pushout diagrams, their double mapping cylinders in spaces are homotopy equivalent. We have the following proposition about hom complexes of graphs in this context.

Proposition 3.3. Let $T \in \mathcal{G}$ and consider $\text{Hom}(T, _) : \mathcal{G} \rightarrow \text{Top}$. Then for the diagram in Figure 3 and $n \geq \text{diam}(T) + 1$,


$$\begin{array}{ccc} \text{Hom}(T, A) & \xrightarrow{f_T} & \text{Hom}(T, B) \\ g_T \downarrow & & \downarrow j_1 \\ \text{Hom}(T, C) & \xrightarrow{j_2} & \text{Hom}(T, D_n) \end{array}$$

is the homotopy pushout in Top , the category of topological spaces, that is, $\text{Hom}(T, D_n)$ is the double mapping cylinder of maps $\{f_T, g_T\}$.

We can give a combinatorial proof of the above result by describing the polytopes in the hom complex, $\text{Hom}(T, D_n)$, which we omit here for brevity. Alternatively, we can prove this result using properties of pushouts of topological spaces and modifying ideas used in the proof of [16, Theorem 5.1] by Matsushita. However, he has made no reference or use of the above proposition in his article.

Here is a quick example of a graph which is a double mapping cylinder of smaller graphs and the above proposition helps us compute its hom complex.

Application 1: Given any two natural numbers $m, r \in \mathbb{N}$, we use Theorem 3.3 to construct a graph G such that $\chi(G) = m$, and $\text{connHom}(K_r, G) = 0$. First let $r = 2$ and consider the graph $D_n = B \sqcup_f (A \times I_n) \sqcup_g C$ with $B = K_m$, $A = K_2$, $C = K_m$; $m \geq 3$ and $f : A \rightarrow B$, $g : A \rightarrow C$ be inclusions. For $m = 6$, D_n is the graph:



Since f and g are inclusions, the induced maps $f_T : \text{Hom}(T, A) \rightarrow \text{Hom}(T, B)$ and $g_T : \text{Hom}(T, A) \rightarrow \text{Hom}(T, C)$ are also inclusions. By Theorem 3.3, $\text{Hom}(T, D_n) = \text{Hom}(T, K_m \sqcup_f K_2 \times I_n \sqcup_g K_m) \simeq \text{Hom}(T, B) \sqcup_{f_T} (\text{Hom}(T, A) \times I) \sqcup_{g_T} \text{Hom}(T, C)$, that is,

$$\text{Hom}(K_2, D_n) \simeq \text{Hom}(K_2, K_m) \sqcup_{f_{K_2}} (\text{Hom}(K_2, K_2) \times I) \sqcup_{g_{K_2}} \text{Hom}(K_2, K_m).$$

The complex $\text{Hom}(K_2, K_2) \times I$ is homotopy equivalent to two disjoint line segments, say L_1 and L_2 . Suppose L_1 and L_2 have initial points l_1, l_2 and final points l'_1, l'_2 respectively. Babson and Kozlov [3] proved that $\text{Hom}(K_2, K_m)$ is homotopy equivalent to a $(m-2)$ -sphere S^{m-2} . Thus for $T = K_2$, f_T maps l_1, l_2 to $\text{Hom}(K_2, B) \simeq S^{m-2}$ and g_T maps l'_1, l'_2 to $\text{Hom}(K_2, C) \simeq S^{m-2}$.

Therefore, $\text{Hom}(K_2, D_n) \simeq S^{m-2} \vee S^1 \vee S^{m-2}$. In particular, $\text{connHom}(K_2, D_n) = 0$.

For an arbitrary r and m , $r < m$, define G to be the double mapping cylinder of f, g of height more than r , where $f = g : K_r \rightarrow K_m$ are the inclusion maps. Then the above argument shows that $\text{Hom}(K_r, G)$ is homotopy equivalent to wedge of 1-dimensional spheres, \mathbb{S}^1 , and $(m-r)$ -dimensional spheres, \mathbb{S}^{m-r} . The graph G is clearly m -colourable, and $\text{connHom}(K_r, G) = 0$.

In fact, for any $r < p \leq m$ if G is the double mapping cylinder of $f : K_r \rightarrow K_m, g : K_r \rightarrow K_p$ of height more than r , then $\chi(G) = m$ and $\text{connHom}(K_r, G) = 0$. Let \mathcal{T} be the class of graphs G as defined above. Then by choosing r small and m large, \mathcal{T} gives a class of graphs for which the homotopy test graph K_r gives a weak lower bound for the chromatic number of graphs in \mathcal{T} via the inequality (1). \square

In the previous example, the graph A is chosen to be K_2 , which we recall is a test graph. We now generalise this construction for any homotopy test graph T . Let G be a graph and $v \in G$ be such that G folds to $G - v$. In view of [13], it is clear that for any graph H , $\text{connHom}(G, H) = \text{connHom}(G - v, H)$. Also, the chromatic number of G is equal to the chromatic number of $G - v$. Therefore, a graph T is a homotopy test graph if and only if its unique (upto graph isomorphism) stiff subgraph T' is a homotopy test graph. So without loss of generality let T be a stiff homotopy test graph. We further assume that T has an endomorphism¹ (other than the identity morphism). Let $m \in \mathbb{N}$ be large enough so that there exists $i : T \rightarrow K_m$, a graph inclusion. Let $n-1 = \text{diam}(T)$, and D_n be the double mapping cylinder of f, g of length n , where $f = g = i$. Since T is stiff, the identity morphism of T is an isolated point in the hom complex of $\text{Hom}(T, T)$ by Lemma 6.5 of [9]. Therefore the hom complex, $\text{Hom}(T, D_n)$, is homotopy equivalent to $X \vee \mathbb{S}^1$ for some prodsimplicial complex X implying that $\pi_1(\text{Hom}(T, D_n))$ is non-trivial, and hence $\text{connHom}(T, D_n) \leq 0$. However, the chromatic number of D_n is m . We have shown the following:

Proposition 3.4. *Let T be a homotopy test graph with a non-identity endomorphism. For any $m \geq |V(T)|$, there exists a graph G such that $\text{connHom}(T, G) = 0$, and $\chi(G) = m$.*

One can also use Theorem 3.3 for computing homotopy type of several hom complexes easily if they are \times -homotopy equivalent to double mapping cylinders of a pushout diagram of graphs.

We next give such a quick computation for a class of graphs explored by Daneshpajouh in [8].


Application 2: Let H and L be non-bipartite graphs with disjoint vertex sets and let $x \in V(H)$ and $y \in V(L)$. Daneshpajouh defined G_{H_x, L_y} to be the graph (cf. [8, Definition 1]) to be the graph with $V(G_{H_x, L_y}) = V(H) \cup V(L) \cup \{z\}$ and $E(G_{H_x, L_y}) = E(H) \cup E(L) \cup \{xz, zy\}$ where $z \notin (V(H) \cup V(L))$. He computed the homotopy type of the neighbourhood complex of this graph. Using this construction, for any given positive integers l, m , and $2 \leq p \leq q$, he has defined a graph containing a copy of $K_{l,m}$, having the chromatic number, clique number and Lovász bound as q, p , and 3, respectively.

¹An endomorphism is a morphism from the object to itself.

We compute the homotopy type of the $\mathcal{N}(G_{H_x, L_y})$ using Theorem 3.3. Let B and C be two non-trivial graphs with special vertices $b \in V(B), c \in V(C)$. For $A = K_2$ with $V(K_2) = \{x, y\}$, let $B_b = (B \sqcup K_2)/(b \sim x), C_c = (C \sqcup K_2)/(c \sim x)$ be the graphs obtained by taking wedge sum (for definition, please refer to [1]) of B with K_2 and C with K_2 respectively.

Let $f : A \rightarrow B_b, g : A \rightarrow C_c$ be graph inclusions. Then the pushout of f, g is the graph G_{B_b, C_c} . Let D_2 be the double mapping cylinder of f and g of height 2. Then D_2 is \times -homotopic to G_{B_b, C_c} via a sequence of folds, therefore by [13], $\mathcal{N}(G_{B_b, C_c})$ is homotopically equivalent to $\mathcal{N}(D_2)$. By Theorem 3.3, $\mathcal{N}(G_{B_b, C_c})$ is homotopically equivalent to the wedge $\mathcal{N}(B_b) \vee \mathcal{N}(C_c) \vee \mathbb{S}^1 \simeq \mathcal{N}(B) \vee \mathcal{N}(C) \vee \mathbb{S}^1$ as B_b and C_c folds to B and C , respectively.

Proposition 3.3 also applies in cases when the neighbourhood complex of the double mapping cylinder is more connected than that of pushout. For instance, consider the graph G to be the wedge of K_4 and K_3 at a vertex, and choose the following as A, B and C .



With respect to inclusions of A in B , and in C , let $D_n = B \bigsqcup_A^h C$, then computations show that $D_n \simeq_{\times} K_4$ and hence $\text{Hom}(K_2, D_n)$ is homotopically equivalent to a 2-sphere, \mathbb{S}^2 . Therefore $\mathcal{N}(D_n)$ is 1-connected. We can compute $\mathcal{N}(G)$ and see that its first homology is non trivial and hence is not 1-connected. Thus connectivity of $\mathcal{N}(D_n)$ is higher in this pushout. Since $\chi(G) \geq \chi(D_n)$, here the Lovasz bound is improved.

However, $\text{conn}\mathcal{N}(G)$ is not always smaller than $\text{conn}\mathcal{N}(D_n)$. For instance, in the first application where we use Prop 3.3 to compute the connectivity of a double mapping cylinder, the neighbourhood complex of the pushout graph $K_6 \coprod_{K_2} K_6$ has connectivity at least 1 (cf. [5]).

4 Is D_n a homotopy pushout in graphs?

We have already seen several properties of the double mapping cylinder D_n . In order to fully utilize these ideas, we need a way to recognize when a given pushout diagram is weakly equivalent to a double mapping cylinder.

The definition of D_n is analogous to that of double mapping cylinder in spaces. In topological spaces, the double mapping cylinder gives a homotopy pushout. Recall homotopy pushout in spaces are preserved under homotopy equivalent diagrams and we can recognize when a given pushout is a homotopy pushout (namely, if one of the maps in the pushout diagram is a cofibration).

Consider the category, \mathcal{G} , of graphs with weak equivalences as \times -homotopy equivalences. If the double mapping cylinder construction in graphs is a homotopy pushout in the category of graphs, then we can build the theory around it to get a recognizing principle.

However, the double mapping cylinder fails to preserve \times -homotopy equivalent diagrams. Let $A \xrightarrow[h_A]{} A', B \xrightarrow[h_B]{} B', C \xrightarrow[h_C]{} C'$ be \times -homotopy equivalences with \times -homotopy inverses $A' \xrightarrow[h_{A'}]{} A$

$A, B' \xrightarrow[h_{B'}]{\simeq} B, C' \xrightarrow[h_{C'}]{\simeq} C$ respectively. Let $f' : A' \rightarrow B', g' : A' \rightarrow C'$ be graph homomorphisms. There exists a canonical graph homomorphism $F : D_n \rightarrow D'_n$ which is defined as $F(b) = h_B(b), F(a, i) = (h_A(a), i)$ and $F(c) = h_C(c)$ for all $b \in V(B), a \in V(A), i \in V(I_n), c \in V(C)$.

Let $f h_{A'} \simeq_{H_B} h_{B'} f', h_{C'} g' \simeq_{H_C} g h_{A'}$ be \times -homotopies of length m and k respectively. Analogous to the case of topological spaces, we expect to define $F' : D'_{n'} \rightarrow D_n$ using the homotopies H_C and H_B . To do this in graphs, one needs to take the double mapping cylinder of $\{f', g'\}$ of length at least $m+k$ more than that of double mapping cylinder of $\{f, g\}$. Thus there is no graph homomorphism $F' : D'_{n'} \rightarrow D_n$.

Since the domain and codomain for the composite FF' are double mapping cylinders of $\{f', g'\}$ of different lengths, FF' cannot be \times -homotopic to the identity homomorphism of any graph. As mentioned earlier, D_n is not \times -homotopic to D_m , for $n \neq m$. Theorem 3.3 implies that there is a homotopy equivalence between $\text{Hom}(T, D_n)$ and $\text{Hom}(T, D_m)$ for all T . However, this is not induced by a graph homomorphism.

We then check if FF' is \times -homotopic to a graph homomorphism, 'shrink', that induces a homotopy equivalence on $\text{Hom}(T, _)$, for all T . If this turned out to be true, then we would add a few more weak equivalences to our category of graphs without compromising the structure. In the category of graphs with this new class of weak equivalences, we will stand a chance at characterizing D_n as a homotopy pushout. However, this is also false.

We define the graph homomorphism $shrink_{m+k} : D'_{m+n+k} \rightarrow D'_n$ as identity on $B' \sqcup C'$ and

$$shrink_{m+k}(a', s) = \begin{cases} (a', 0) & \text{if } 0 \leq s \leq k, \\ (a', s - k) & \text{if } k \leq s \leq k + n, \\ (a', n) & \text{if } k + n \leq s \leq k + n + m, \end{cases} \quad (2)$$

for all $a' \in V(A')$ and $s \in V(I_{m+n+k})$. The subscript $m+k$ in $shrink_{m+k}$ indicates the levels above and below n , respectively, that the map $shrink_{m+k}$ shrinks. Clearly, $shrink_{m+k}$ induces a homotopy equivalence from $\text{Hom}(T, D'_{m+n+k}) \rightarrow \text{Hom}(T, D'_n)$, for all $T \in \mathcal{G}$ and $m, n, k \geq 0$.

Let $l = \max\{m, k\}$ so that all the given homotopies are of the same length l . This can be done by repeating the values on the last level of the homotopy on the newly added levels.

For $n' = n + 2l$, $F' : D'_{l+n+l} \rightarrow D_n$ can be defined as $F'(b') = h_{B'}(b'), F'(c') = h_{C'}(c')$, for all $b' \in B'$ and $c' \in C'$, and for $a' \in A', s \in I_{l+n+l}$,

$$F'(a', s) = \begin{cases} H_C(a', s) & \text{if } 0 \leq s \leq l, \\ (h_{A'}(a'), s - l) & \text{if } l \leq s \leq l + n, \\ H_B(a', s - (n + l)) & \text{if } l + n \leq s \leq l + n + l, \end{cases}$$

Suppose $FF' \simeq_{\times} shrink_{l+l}$, then for some $r \in \mathbb{N}$ there exists a \times -homotopy $K : D'_{l+n+l} \times I_r \rightarrow D'_n$ such that $K|_{D'_{l+n+l} \times \{0\}} = FF'$ and $K|_{D'_{l+n+l} \times \{r\}} = shrink_{l+l}$. Since $FF'|_{A' \times I_l}$ is also a \times -homotopy, in general $K(a', l-1, 0) = FF'(a', l-1) = h_C H_C(a', l-1)$ need not be adjacent to $K(a'', l, 1)$, where $a' a'' \in E(A')$. Therefore, with no other assumptions, if we use only these given homotopies, then there does not exist a \times -homotopy between FF' and the $shrink_{l+l}$.

One possible approach to resolve this issue is to define an appropriate class of cofibrations with the class of weak equivalences as the \times -homotopy equivalences. Such a setup will allow us to give a better definition of homotopy pushouts in graphs. We explore this by studying model structures on the category of finite graphs with \times -homotopy equivalences in [10].

References

- [1] M. Abu-Saleem. Folding on the wedge sum of graphs and their fundamental group. *Appl. Sci.*, 12:14-19, 2010.
- [2] Eric Babson and Dmitry N. Kozlov. Topological obstructions to graph colorings. *Electron. Res. Announc. Amer. Math. Soc.*, 9:61-68 (electronic), 2003.
- [3] Eric Babson and Dmitry N. Kozlov. Complexes of graph homomorphisms. *Israel J. Math.*, 152:285-312, 2006.
- [4] Paul Bonsma and Luis Cereceda. Finding paths between graph colourings: PSPACE-completeness and superpolynomial distances. *Theoret. Comput. Sci.*, 410(50):5215-5226, 2009.
- [5] Pter Csorba. Homotopy types of box complexes of chordal graphs. *European Journal of Combinatorics*, 31(3):861 – 866, 2010.
- [6] Sonja Lj.Čukić and Dmitry N. Kozlov. Higher connectivity of graph coloring complexes. *Int. Math. Res. Not.*, (25):1543-1562, 2005.
- [7] Sonja Lj.Čukić and Dmitry N. Kozlov. The homotopy type of complexes of graph homomorphisms between cycles. *Discrete Comput. Geom.*, 36(2):313-329, 2006.
- [8] Hamid Reza Daneshpajouh. Chromatic number, Clique number, and Lovász’s bound: In a comparison. *arXiv:1709.07159*.
- [9] Anton Dochtermann. Hom complexes and homotopy theory in the category of graphs. *European J. Combin.*, 30(2):490-509, 2009.
- [10] Shuchita Goyal and Rekha Santhanam. (Lack of) Model Structures on the Category of Graphs. *arXiv:1902.09182*
- [11] Shlomo Hoory and Nathan Linial. A counterexample to a conjecture of Björner and Lovász on the χ -coloring complex. *J. Combin. Theory Ser. B*, 95(2):346-349, 2005.
- [12] Dmitry N. Kozlov. Simple homotopy types of Hom-complexes, neighborhood complexes, Lovász complexes, and atom crosscut complexes. *Topology Appl.*, 153(14):2445-2454, 2006.
- [13] Dmitry N. Kozlov. A simple proof for folds on both sides in complexes of graph homomorphisms. *Proc. Amer. Math. Soc.*, 134(5):1265-1270 (electronic), 2006.
- [14] L. Lovász. Kneser’s conjecture, chromatic number, and homotopy. *J. Combin. Theory Ser. A*, 25(3):319-324, 1978.
- [15] Takahiro Matsushita. Answers to some problems about graph coloring test graphs. *European J. Combin.*, 45:59-64, 2015.
- [16] Takahiro Matsushita. Homotopy types of the Hom complexes of graphs. *European J. Combin.*, 63:216-226, 2017.
- [17] Carsten Schultz. A short proof of $w_1^n(\text{Hom}(C_{2r+1}, K_{n+2})) = 0$ for all n and a graph colouring theorem by Babson and Kozlov. *Israel J. Math.*, 170:125-134, 2009.
- [18] Carsten Schultz. The equivariant topology of stable Kneser graphs. *J. Combin. Theory Ser. A*, 118(8):2291-2318, 2011.