

# Counting faces of graphical zonotopes

Vladimir Grujić

Faculty of Mathematics  
Belgrade University, Serbia

Mathematics Subject Classifications: 05E05, 52B05, 16T05

## Abstract

It is a classical fact that the number of vertices of the graphical zonotope  $Z_\Gamma$  is equal to the number of acyclic orientations of a graph  $\Gamma$ . We show that the  $f$ -polynomial of  $Z_\Gamma$  is obtained as the principal specialization of the  $q$ -analog of the chromatic symmetric function of  $\Gamma$ .

**Keywords:** graphical zonotope,  $f$ -vector, graphical matroid, symmetric function

## 1 Introduction

The  $f$ -polynomial of an  $n$ -dimensional polytope  $P$  is defined by  $f(P, q) = \sum_{i=0}^n f_i(P)q^i$ , where  $f_i(P)$  is the number of  $i$ -dimensional faces of  $P$ . The  $f$ -polynomial  $f(Z_\Gamma, q)$  of the graphical zonotope  $Z_\Gamma$  is a combinatorial invariant of a finite, simple graph  $\Gamma$ . The vertices of  $Z_\Gamma$  are in one-to-one correspondence with regions of the graphical hyperplane arrangement  $\mathcal{H}_\Gamma$ , which are enumerated by acyclic orientations of  $\Gamma$ .

Stanley's chromatic symmetric function  $\Psi(\Gamma) = \sum_{f \text{ proper}} \mathbf{x}_f$  of a graph  $\Gamma = (V, E)$ , introduced in [6], is the enumerator function of proper colorings  $f : V \rightarrow \mathbb{N}$ , where  $\mathbf{x}_f = x_{f(1)} \cdots x_{f(n)}$  and  $f$  is proper if there are no monochromatic edges. The chromatic polynomial  $\chi(\Gamma, d)$  of the graph  $\Gamma$ , which counts proper colorings with a finite number of colors, appears as the principal specialization

$$\chi(\Gamma, d) = \mathbf{ps}(\Psi(\Gamma))(d) = \Psi(\Gamma) |_{x_1=\dots=x_d=1, x_{d+1}=\dots=0}.$$

The number of acyclic orientations of  $\Gamma$  is determined by the value of the chromatic polynomial  $\chi(\Gamma, d)$  at  $d = -1$ , [7]

$$a(\Gamma) = (-1)^{|V|} \chi(\Gamma, -1). \quad (1)$$

There is a  $q$ -analog of the chromatic symmetric function  $\Psi_q(\Gamma)$  introduced in a wider context of the combinatorial Hopf algebra of simplicial complexes considered in [2]. It is a symmetric function over the field of rational functions

in  $q$ . The principal specialization of  $\Psi_q(\Gamma)$  is the  $q$ -analog of the chromatic polynomial  $\chi_q(\Gamma, d)$ .

The main result of this paper is the following generalization of formula (1)

**Theorem 1.1.** *Let  $\Gamma = (V, E)$  be a simple connected graph and  $\mathcal{Z}_\Gamma$  the corresponding graphical zonotope. Then the  $f$ -polynomial of  $\mathcal{Z}_\Gamma$  is given by*

$$f(\mathcal{Z}_\Gamma, q) = (-1)^{|V|} \chi_{-q}(\Gamma, -1).$$

The cancellation-free formula for the antipode in the Hopf algebra of graphs, obtained by Humpert and Martin in [3], reflects the fact that  $f(\mathcal{Z}_\Gamma, q)$  depends only on the graphical matroid  $M(\Gamma)$  associated to  $\Gamma$ . For instance, for any tree  $T_n$  the graphical matroid is the uniform matroid  $M(T_n) = U_n^n$  and the corresponding graphical zonotope is the cube  $\mathcal{Z}_{T_n} = I^{n-1}$ . Whitney's theorem from 1933 describes how two graphs with the same graphical matroid are related [9]. It can be used to find more interesting nonisomorphic graphs with the same  $f$ -polynomials of corresponding graphical zonotopes.

The paper is organized as follows. In section 2, we review the basic facts about zonotopes. In section 3, the  $q$ -analog of the chromatic symmetric function  $\Psi_q(\Gamma)$  of a graph  $\Gamma$  is introduced. Theorem 1.1 is proved in section 4. We present some examples and calculations in section 5.

## 2 Zonotopes

A zonotope  $\mathcal{Z} = \mathcal{Z}(v_1, \dots, v_m)$  is a convex polytope determined by a collection of vectors  $\{v_1, \dots, v_m\}$  in  $\mathbb{R}^n$  as the Minkowski sum of line segments

$$\mathcal{Z} = [-v_1, v_1] + \dots + [-v_m, v_m].$$

It is a projection of the  $m$ -cube  $[-1, 1]^m$  under the linear map  $\mathbf{t} \mapsto A\mathbf{t}$ ,  $\mathbf{t} \in [-1, 1]^m$ , where  $A = [v_1 \dots v_m]$  is an  $n \times m$ -matrix whose columns are vectors  $v_1, \dots, v_m$ . The zonotope  $\mathcal{Z}$  is symmetric about the origin and all its faces are translations of zonotopes.

To a collection of vectors  $\{v_1, \dots, v_m\}$  is associated a central arrangement of hyperplanes  $\mathcal{H} = \{H_{v_1}, \dots, H_{v_m}\}$ , where  $H_v$  denotes the hyperplane perpendicular to a vector  $v \in \mathbb{R}^n$ . The zonotope  $\mathcal{Z}$  and the corresponding arrangement of hyperplanes  $\mathcal{H}$  are closely related. In fact the associated fan  $\mathcal{F}_\mathcal{H}$  of the arrangement  $\mathcal{H}$  is the normal fan  $\mathcal{N}(\mathcal{Z})$  of the zonotope  $\mathcal{Z}$  (see [10, Theorem 7.16]). It follows that the face lattice of  $\mathcal{F}_\mathcal{H}$  and the reverse face lattice of  $\mathcal{Z}$  are isomorphic. In particular, vertices of  $\mathcal{Z}$  correspond to regions of  $\mathcal{H}$  and their total numbers coincide

$$f_0(\mathcal{Z}) = r(\mathcal{H}). \tag{2}$$

The faces of the zonotope  $\mathcal{Z}$  are encoded by covectors of the oriented matroid  $\mathcal{M}$  associated to the collection of vectors  $\{v_1, \dots, v_m\}$ . The covectors are sign vectors

$$\mathcal{V}^* = \{\text{sign}(v) \in \{+, -, 0\}^m \mid v \in \mathbb{R}^m\},$$

where  $\text{sign}(v)_i = \begin{cases} +, & \langle v, v_i \rangle > 0 \\ 0, & \langle v, v_i \rangle = 0 \\ -, & \langle v, v_i \rangle < 0 \end{cases}$ ,  $i = 1, \dots, m$ . The face lattice of the zonotope  $\mathcal{Z}$  is isomorphic to the lattice of covectors componentwise induced by  $+, - < 0$  on  $\mathcal{V}^*$ .

A special class of zonotopes is determined by simple graphs. To a connected graph  $\Gamma = (V, E)$ , whose vertices are enumerated by integers  $V = \{1, \dots, n\}$ , are associated the *graphical zonotope*

$$\mathcal{Z}_\Gamma = \mathcal{Z}(e_i - e_j \mid i < j, \{i, j\} \in E)$$

and the *graphical arrangement* in  $\mathbb{R}^n$

$$\mathcal{H}_\Gamma = \{H_{e_i - e_j} \mid i < j, \{i, j\} \in E\}.$$

There is a bijective correspondence between regions of  $\mathcal{H}_\Gamma$  and acyclic orientations of  $\Gamma$ , [8, Proposition 2.5], which by (2) implies

$$f_0(\mathcal{Z}_\Gamma) = r(\mathcal{H}_\Gamma) = a(\Gamma). \quad (3)$$

The arrangement  $\mathcal{H}_\Gamma$  is refined by the braid arrangement  $\mathcal{A}_{n-1}$  consisting of all hyperplanes  $H_{e_i - e_j}$ ,  $1 \leq i < j \leq n$ . Thus  $\mathcal{Z}_\Gamma$  belongs to a wider class of convex polytopes called generalized permutohedra introduced in [4]. Since arrangements  $\mathcal{H}_\Gamma$  and  $\mathcal{A}_{n-1}$  are not essential we take their quotients by the line  $l : x_1 = \dots = x_n$  and without confusing retain the same notation. Consequently  $\dim \mathcal{Z}_\Gamma = n - 1$ .




Figure 1: Permutohedron  $Pe^3$  and cube  $I^3$

**Example 2.1.** (i) The permutohedron  $Pe^{n-1}$  is represented as the graphical zonotope  $\mathcal{Z}_{K_n}$  corresponding to the complete graph  $K_n$  on  $n$  vertices (Figure 1).

(ii) The cube  $I^{n-1}$  is represented as the graphical zonotope  $\mathcal{Z}_{T_n}$  corresponding to an arbitrary tree  $T_n$  on  $n$  vertices. This shows that the graph  $\Gamma$  is not determined by the combinatorial type of the zonotope  $\mathcal{Z}_\Gamma$ .

### 3 $q$ -analog of chromatic symmetric function of graph

Stanley's chromatic symmetric function  $\Psi(\Gamma)$  can be obtained in a purely algebraic way. A combinatorial Hopf algebra  $\mathcal{H}$  is a graded, connected Hopf algebra equipped with the multiplicative linear functional  $\zeta : \mathcal{H} \rightarrow \mathbf{k}$  to the ground field  $\mathbf{k}$ . For the theory of combinatorial Hopf algebras see [1]. Consider the combinatorial Hopf algebra of graphs  $\mathcal{G}$  which is linearly generated over a field  $\mathbf{k}$  by simple finite graphs with the product defined by disjoint union  $\Gamma_1 \cdot \Gamma_2 = \Gamma_1 \sqcup \Gamma_2$  and the coproduct

$$\Delta(\Gamma) = \sum_{I \subset V} \Gamma|_I \otimes \Gamma|_{V \setminus I},$$

where  $\Gamma|_I$  denotes the induced subgraph on  $I \subset V$ . The structure of  $\mathcal{G}$  is completed by the character  $\zeta : \mathcal{G} \rightarrow \mathbf{k}$  defined to be  $\zeta(\Gamma) = 1$  for  $\Gamma$  with no edges and  $\zeta(\Gamma) = 0$  otherwise. Then it turns out that  $\Psi(\Gamma)$  is the image of the unique morphism of combinatorial Hopf algebras to symmetric functions  $\Psi : \mathcal{G} \rightarrow Sym$ , ([1, Example 4.5.]).

An important part of the structure of the Hopf algebra  $\mathcal{G}$  is the antipode  $S : \mathcal{G} \rightarrow \mathcal{G}$ . The cancellation-free formula for the antipode in terms of acyclic orientations of a graph  $\Gamma$  is obtained in [3]. We recall some basic definitions. Terminology comes from matroid theory. Given a graph  $\Gamma = (V, E)$ , for a collection of edges  $F \subset E$  denote by  $\Gamma_{V,F}$  the graph on  $V$  with the edge set  $F$ . A *flat*  $F$  of the graph  $\Gamma$  is a collection of its edges such that components of  $\Gamma_{V,F}$  are induced subgraphs. The rank  $\text{rk}(F)$  is the size of spanning forests of  $\Gamma_{V,F}$ . We have that  $|V| = \text{rk}(F) + c(F)$ , where  $c(F)$  is the number of components of  $\Gamma_{V,F}$ . By contracting edges from a flat  $F$  we obtain the graph  $\Gamma/F$ . Finally, let  $a(\Gamma)$  be the number of acyclic orientations of  $\Gamma$ . The formula of Humpert and Martin is as follows

$$S(\Gamma) = \sum_{F \in \mathcal{F}(\Gamma)} (-1)^{c(F)} a(\Gamma/F) \Gamma_{V,F}, \quad (4)$$

where the sum is over the set of flats  $\mathcal{F}(\Gamma)$ .

The following modification of the character  $\zeta$  is considered in [2] in a wider context of the combinatorial Hopf algebra of simplicial complexes. Define  $\zeta_q(\Gamma) = q^{\text{rk}(\Gamma)}$ , which determines the algebra morphism  $\zeta_q : \mathcal{G} \rightarrow \mathbf{k}(q)$ , where  $\mathbf{k}(q)$  is the field of rational functions in  $q$ . This character produces the unique morphism  $\Psi_q : \mathcal{G} \rightarrow QSym$  to quasisymmetric functions over  $\mathbf{k}(q)$ . The expansion of  $\Psi_q(\Gamma)$  in the monomial basis of quasisymmetric functions is determined by the universal formula [1, Theorem 4.1]

$$\Psi_q(\Gamma) = \sum_{\alpha \models n} (\zeta_q)_\alpha(\Gamma) M_\alpha.$$

The sum above is over all compositions of the integer  $n = |V|$  and the coefficient of the expansion corresponding to the composition  $\alpha = (a_1, \dots, a_k) \models n$  is given by

$$(\zeta_q)_\alpha(\Gamma) = \sum_{I_1 \sqcup \dots \sqcup I_k = V} q^{\text{rk}(\Gamma|_{I_1}) + \dots + \text{rk}(\Gamma|_{I_k})},$$

where the sum is over all set compositions of  $V$  of the type  $\alpha$ . The coefficients  $(\zeta_q)_\alpha(\Gamma)$  depend only on the partition corresponding to a composition  $\alpha$ , so the function  $\Psi_q(\Gamma)$  is actually symmetric and it can be expressed in the monomial basis of symmetric functions.

The invariant  $\Psi_q(\Gamma)$  is more subtle than  $\Psi(\Gamma)$ . Obviously  $\Psi_0(\Gamma)$  is the chromatic symmetric function of a graph  $\Gamma$ . It remains open to find two non-isomorphic graphs  $\Gamma_1$  and  $\Gamma_2$  with the same  $q$ -chromatic symmetric functions  $\Psi_q(\Gamma_1) = \Psi_q(\Gamma_2)$ . Let

$$\chi_q(\Gamma, d) = \mathbf{ps}(\Psi_q(\Gamma))(d)$$

be the  $q$ -analog of the chromatic polynomial  $\chi(\Gamma, d)$ . It is a consequence of a general fact for combinatorial Hopf algebras (see [1]) that

$$\chi_q(\Gamma, -1) = (\zeta_q \circ S)(\Gamma). \quad (5)$$

**Example 3.1.** Consider the graph  $\Gamma$  on four vertices with the edge set  $E = \{12, 13, 23, 34\}$ . We find that

$$\Psi_q(\Gamma) = 24m_{1,1,1,1} + (8q + 4)m_{2,1,1} + (2q^2 + 4q)m_{2,2} + (3q^2 + q)m_{3,1} + q^3m_4.$$

By principal specialization and taking into account that

$$\mathbf{ps}(m_{\lambda_1^{i_1}, \dots, \lambda_k^{i_k}})(d) = \frac{(i_1 + \dots + i_k)!}{i_1! \dots i_k!} \binom{d}{i_1 + \dots + i_k},$$

we obtain

$$\chi_q(\Gamma, d) = d(d-1)^2(d-2) + qd(d-1)(4d-5) + 4q^2d(d-1) + q^3d,$$

which by Theorem 1.1 gives

$$f(\mathcal{Z}_\Gamma, q) = 12 + 18q + 8q^2 + q^3.$$

## 4 Proof of Theorem 1.1

By applying (5) and the formula for antipode (4) we obtain

$$(-1)^{|V|} \chi_{-q}(\Gamma, -1) = (-1)^{|V|} \sum_{F \in \mathcal{F}(\Gamma)} (-1)^{c(\Gamma)} a(\Gamma/F) (-q)^{\text{rk}(F)}.$$

It follows that the statement of the theorem is equivalent to the following expression of the  $f$ -polynomial

$$f(\mathcal{Z}_\Gamma, q) = \sum_{F \in \mathcal{F}(\Gamma)} a(\Gamma/F) q^{\text{rk}(F)}. \quad (6)$$

Therefore it should be shown that components of  $f$ -vectors are determined by

$$f_k(\mathcal{Z}_\Gamma) = \sum_{\substack{F \in \mathcal{F}(\Gamma) \\ \text{rk}(F) = k}} a(\Gamma/F), \quad 0 \leq k \leq n-1. \quad (7)$$

By duality between the face lattice of  $\mathcal{Z}_\Gamma$  and the face lattice of the fan  $\mathcal{F}_{\mathcal{H}_\Gamma}$  we have

$$f_k(\mathcal{Z}_\Gamma) = f_{n-k-1}(\mathcal{F}_{\mathcal{H}_\Gamma}).$$

Let  $L(\mathcal{H}_\Gamma)$  be the intersection lattice of the graphical arrangement  $\mathcal{H}_\Gamma$ . For a subspace  $X \in L(\mathcal{H}_\Gamma)$  there is an arrangement of hyperplanes

$$\mathcal{H}_\Gamma^X = \{X \cap H \mid X \not\subseteq H, H \in \mathcal{H}_\Gamma\}$$

whose intersection lattice  $L(\mathcal{H}_\Gamma^X)$  is isomorphic to the upper cone of  $X$  in  $L(\mathcal{H}_\Gamma)$ . Since  $\mathcal{H}_\Gamma$  is central and essential we have

$$f_{n-k-1}(\mathcal{F}_{\mathcal{H}_\Gamma}) = \sum_{\substack{X \in L(\mathcal{H}_\Gamma) \\ \dim(X) = n-k-1}} r(\mathcal{H}_\Gamma^X), \quad (8)$$

where  $r(\mathcal{H}_\Gamma^X)$  is the number of regions of the arrangement  $\mathcal{H}_\Gamma^X$ , see [8, Theorem 2.6].

The intersection lattice  $L(\mathcal{H}_\Gamma)$  is isomorphic to the lattice of flats of the graphical matroid  $M(\Gamma)$ . By this isomorphism to a flat  $F$  of rank  $k$  corresponds the intersection subspace  $X^F = \cap_{\{i,j\} \in F} H_{e_i - e_j}$  of dimension  $n-k-1$ . It is easy to see that arrangements  $\mathcal{H}_\Gamma^{X^F}$  and  $\mathcal{H}_{\Gamma/F}$  coincide, which by (3) and comparing formulas (7) and (8) proves theorem.

## 5 Examples

By applying Theorem 1.1 we obtain the following interpretation of identities elaborated in [2, Propositions 17, 19].

**Example 5.1.** (i) For the permutohedron  $Pe^{n-1} = \mathcal{Z}_{K_n}$ , the  $f$ -polynomial is given by

$$f(\mathcal{Z}_{K_n}, q) = A_n(q+1),$$

where  $A_n(q) = \sum_{\pi \in S_n} q^{\text{des}(\pi)}$  is the Euler polynomial. Recall that  $\text{des}(\pi)$  is the number of descents of a permutation  $\pi \in S_n$ . It recovers the fact that the  $h$ -polynomial of the permutohedron  $Pe^{n-1}$  is the Euler polynomial  $A_n(q)$ .

(ii) For the cube  $I^{n-1} = \mathcal{Z}_{T_n}$ , where  $T_n$  is a tree on  $n$  vertices, the  $f$ -polynomial is given by

$$f(\mathcal{Z}_{T_n}, q) = (q+2)^{n-1}.$$




Figure 2: Rhombic dodecahedron  $\mathcal{Z}_{C_4}$

**Proposition 5.2.** *The  $f$ -polynomial of the graphical zonotope  $\mathcal{Z}_{C_n}$  associated to the cycle graph  $C_n$  on  $n$  vertices is given by*

$$f(\mathcal{Z}_{C_n}, q) = q^n + q^{n-1} + (q+2)^n - 2(q+1)^n.$$

*Proof.* A flat  $F \in \mathcal{F}(C_n)$  is determined by the complementary set of edges. If  $\text{rk}(F) = n - k, k > 1$  then the complementary set has  $k$  edges and  $C_n/F = C_k$ . Since  $a(C_k) = 2^k - 2, k > 1$ , by formula (7), we obtain

$$f_{n-k}(\mathcal{Z}_{C_n}) = (2^k - 2) \binom{n}{k}, 2 \leq k \leq n,$$

which leads to the required formula. □

Specially, for  $n = 4$  the resulting zonotope is the rhombic dodecahedron (see Figure 2). We have

$$f(\mathcal{Z}_{C_4}, q) = 14 + 24q + 12q^2 + q^3.$$

**Proposition 5.3.** *Let  $\Gamma = \Gamma_1 \vee_v \Gamma_2$  be the wedge of two connected graphs  $\Gamma_1$  and  $\Gamma_2$  at the common vertex  $v$ . Then*

$$f(\mathcal{Z}_\Gamma, q) = f(\mathcal{Z}_{\Gamma_1}, q)f(\mathcal{Z}_{\Gamma_2}, q).$$

*Proof.* The graphical matroids of involving graphs are related by  $M(\Gamma) = M(\Gamma_1) \oplus M(\Gamma_2)$ . For the sets of flats it holds  $\mathcal{F}(\Gamma) = \{F_1 \cup F_2 \mid F_i \in \mathcal{F}(\Gamma_i), i = 1, 2\}$ . For  $F = F_1 \cup F_2$  we have  $\Gamma/F = \Gamma_1/F_1 \vee_{[v]} \Gamma_2/F_2$ , where  $[v]$  is the component of the vertex  $v$  in  $\Gamma_{V,F}$ . Obviously  $a(\Gamma/F) = a(\Gamma_1/F_1)a(\Gamma_2/F_2)$  and  $\text{rk}(F) = \text{rk}(F_1) + \text{rk}(F_2)$ . The proposition follows from formula (6). □

The formula for cubes in Example 5.1 (ii) follows from Proposition 5.3 since any tree is a consecutive wedge of edges and  $f(I^1, q) = q + 2$ . It also allows us to restrict ourselves only to biconnected graphs. For a biconnected graph  $\Gamma$  with a disconnecting pair of vertices  $\{u, v\}$  Whitney introduced the transformation

called the *twist* around the pair  $\{u, v\}$ . This transformation does not have an affect on the graphical matroid  $M(\Gamma)$  [9].




Figure 3: Biconnected graphs related by twist transformation

**Example 5.4.** Figure 3 shows the pair of biconnected graphs on six vertices obtained one from another by the twist transformation. The corresponding zonotopes have the same  $f$ -polynomial

$$f(\mathcal{Z}_{\Gamma_1}, q) = f(\mathcal{Z}_{\Gamma_2}, q) = 126 + 348q + 358q^2 + 164q^3 + 30q^4 + q^5.$$

On the other hand their  $q$ -chromatic symmetric functions are different. One can check that corresponding coefficients by  $m_{3,1^3}$  are different

$$\begin{aligned} [m_{3,1^3}] \Psi_q(\Gamma_1) &= (11q^2 + 8q + 1) \cdot 3!, \\ [m_{3,1^3}] \Psi_q(\Gamma_2) &= (10q^2 + 10q) \cdot 3!. \end{aligned}$$

This shows that the  $q$ -analog of the chromatic symmetric function of a graph is not determined by the corresponding graphical matroid. By taking  $q = 0$  we obtain that even the chromatic symmetric functions are different since  $[m_{3,1^3}] \Psi(\Gamma_1) = 6$  and  $[m_{3,1^3}] \Psi(\Gamma_2) = 0$ .

Let us now consider Stanley's example of nonisomorphic graphs with the same chromatic symmetric functions, see [6]. We find that the  $f$ -polynomials of the corresponding graphical zonotopes differ for those graphs. From these examples we conclude that chromatic properties of a graph and the  $f$ -vector of the corresponding graphical zonotope are not related.

We have already noted that graphical zonotopes are generalized permutohedra. The  $h$ -polynomials of simple generalized permutohedra are determined in [5, Theorem 4.2]. The only simple graphical zonotopes are products of permutohedra [5, Proposition 5.2]. They are characterized by graphs whose biconnected components are complete subgraphs. Therefore Proposition 5.3 together with Example 5.1 (i) prove that the  $h$ -polynomial of a simple graphical zonotope is the product of Eulerian polynomials, the fact obtained in [5, Corollary 5.4]. Example 3.1 is of this sort and represents the hexagonal prism which is the product  $\mathcal{Z}_{K_3} \times \mathcal{Z}_{K_2}$ .



## References

- [1] M. Aguiar, N. Bergeron and F. Sottile, Combinatorial Hopf algebras and generalized Dehn-Sommerville relations, *Compositio Math.* **142** (2006), 1–30.
- [2] C. Benedetti, J. Hallam and J. Machacek, Combinatorial Hopf algebra of simplicial complexes, *SIAM J. Discrete Math.*, **30**(3) (2016), 1737–1757.
- [3] B. Humpert and J. Martin, The incidence Hopf algebra of graphs, *SIAM J. Discrete Math.*, **26** (2) (2012), 555–570.
- [4] A. Postnikov, Permutohedra, associahedra, and beyond, *Int. Math. Res. Not.* **6** (2009), 1026–1106.
- [5] A. Postnikov, V. Reiner and L. Williams, Faces of generalized permutohedra, *Documenta Math.* **13** (2008), 207–273.
- [6] R. Stanley, A symmetric function generalization of the chromatic polynomial of a graph, *Adv. Math.* **111** (1995), 166–194.
- [7] R. Stanley, Acyclic orientations of graphs, *Discrete Math.* **5** (1973), 171–178.
- [8] R. Stanley, *An introduction to hyperplane arrangements*, in: E. Miller, V. Reiner, B. Sturmfels (eds.), *Geometric Combinatorics*, IAS/Park City Mathematics Series, vol. 13, AMS, Institute for Advance Studies, 2007, 389–496.
- [9] H. Whitney, 2-isomorphic graphs, *Amer. J. Math.* **55** (1933), 245–254
- [10] G. Ziegler, *Lectures on Polytopes*, Springer-Verlag, New York, 1995.