

LATTICE STRUCTURE OF WEYL GROUPS VIA REPRESENTATION THEORY OF PREPROJECTIVE ALGEBRAS

OSAMU IYAMA, NATHAN READING, IDUN REITEN, AND HUGH THOMAS

ABSTRACT. This paper studies the combinatorics of lattice congruences of the weak order on a finite Weyl group W , using representation theory of the corresponding preprojective algebra Π . Natural bijections are constructed between important objects including join-irreducible congruences, join-irreducible (respectively, meet-irreducible) elements of W , indecomposable τ -rigid (respectively, τ^- -rigid) modules and layers of Π . The lattice-theoretically natural labeling of the Hasse quiver by join-irreducible elements of W is shown to coincide with the algebraically natural labelling by layers of Π . We show that layers of Π are nothing but bricks (or equivalently stones, or 2-spherical modules). The forcing order on join-irreducible elements of W (arising from the study of lattice congruences) is described algebraically in terms of the doubleton extension order. We give a combinatorial description of indecomposable τ^- -rigid modules for type A and D .

CONTENTS

1. Introduction	2
2. Preliminaries	6
2.1. Lattice-theoretic preliminaries	6
2.2. Weak order preliminaries	10
2.3. Algebraic preliminaries	11
2.4. Preliminaries on preprojective algebras	12
3. Homological characterizations of layers	13
4. Bijections: Theorems 1.1 and 1.3	16
5. Doubleton extension order on layer modules	20
6. Combinatorial description of indecomposable τ -rigid modules	22
6.1. Type A	22
6.2. Type D	24
References	31

Key words and phrases. preprojective algebra, Weyl group, weak order, τ -tilting theory, brick, join-irreducible element, lattice congruence.

2010 *Mathematics Subject Classification.* Primary 16G10, Secondary 06B10, 18E40, 20F55.

Osamu Iyama's work on this project was partially supported by JSPS Grant-in-Aid for Scientific Research (B) 24340004, (B) 16H03923, (C) 23540045 and (S) 15H05738. Nathan Reading's work on this project was partially supported by the National Science Foundation under grant numbers DMS-1101568 and DMS-1500949. Idun Reiten was supported by the FriNat grants 196600 and 231000 from the Research Council of Norway. Hugh Thomas's work on this project was partially supported by an NSERC Discovery Grant and the Canada Research Chairs program. The authors also gratefully acknowledge the hospitality of MSRI, Oberwolfach, Bielefeld University, and the Mittag-Leffler Institute.

1. INTRODUCTION

Let Δ be a simply laced Dynkin diagram and W the corresponding Weyl group. Once we fix an orientation Q of Δ , then the representation theory of Q categorifies the root system associated with Δ in the sense that we have Gabriel's bijection between positive roots and indecomposable representations of Q . The preprojective algebra Π of Δ unifies the representation theory of different quivers with the same underlying graph Δ , and their various aspects has been studied, e.g. [BKT, BGL, BIRS, CH, DR, GLS, Lu, KS, N]. Mizuno [Mi] showed that the support τ -tilting theory of Π categorifies the Weyl group W with the weak order in the following sense: There exists a bijection $W \ni w \mapsto I(w)$ from W to the set $\text{s}\tau\text{-tilt}\Pi$ of support τ -tilting Π -modules with the property that $v \leq w$ in the weak order on W if and only if $I(v) \geq I(w)$ in the generation order on $\text{s}\tau\text{-tilt}\Pi$. The ideal $I(w)$ was introduced in [IR, BIRS] and has been studied by several authors, e.g. [AM, A, AIRT, BKT, GLS, K, Le, Ma, ORT, SY1]. In what follows, we will overload the symbol W to denote not only the group W , but also the weak order on W .

The weak order on W is a lattice [BB]: a partial order such that meets (greatest lower bounds) and joins (least upper bounds) exist. It is enlightening to take a more algebraic point of view of lattices, viewing a lattice as a set with two binary operations (meet and join). Seen in this light, the categorification of W by support τ -tilting theory is the categorification of an algebraic object (a lattice) in terms of another algebraic object (a finite-dimensional algebra). In both of these algebraic settings, there is an important algebraic quotient operation. Quotients of the weak order are governed by lattice congruences, while quotients of the preprojective algebra are governed by ideals. A natural question is whether these two notions of quotient are related. The answer is yes, and the relationship turns out to be very nice.

This paper and a companion paper [DIRRT] concern the relationship between the two notions of quotient. In the other paper, we observe, for a more general algebra A and an ideal I of A , that $\text{s}\tau\text{-tilt}(A/I)$ is a lattice quotient of $\text{s}\tau\text{-tilt}A$ and give necessary conditions for lattice congruences which arise in this way from quotients of Π . We study the combinatorics of such *algebraic quotients* of the weak order in general, and in the special case where Π/I is hereditary. We also work out, in detail, the combinatorics of algebraic quotients in type A.

Whereas [DIRRT] starts with algebra quotients and determines what happens to the corresponding lattices, this paper starts from the other direction. Here, we start with the rich combinatorics of (arbitrary, not necessarily algebraic) lattice congruences of W and find that it appears naturally within the representation theory of Π .

The set of all lattice congruences of L form a lattice $\text{Con}L$, and the join-irreducible elements of $\text{Con}L$ are called the join-irreducible congruences (see Section 2.1 for details). The combinatorial approach to congruences of a finite lattice L begins with the connection between arrows in the Hasse quiver of L , join-irreducible elements of L , and join-irreducible congruences on L . We will overload the symbol W , using it to denote the Hasse quiver of the weak order on W .

Our first main theorem connects join-irreducible elements of W and join-irreducible congruences on W to layers of Π . A Π -module is called a *layer* if it is isomorphic to $I(w)/I(ws_i)$ for an arrow $ws_i \rightarrow w$ in the Hasse quiver of W , see [AIRT].

Theorem 1.1. *There exist bijections between the following sets.*

- The set $\text{j-lrr}W$ of join-irreducible elements of W .
- The set $\text{m-lrr}W$ of meet-irreducible elements of W .
- The set $\text{Con}_{\Pi}(W)$ of join-irreducible congruences of W .

- The set $i\tau$ -rigid Π of indecomposable τ -rigid Π -modules.
- The set $i\tau^-$ -rigid Π of indecomposable τ^- -rigid Π -modules.
- The set $\text{layer } \Pi$ of layers of Π .

We prove Theorem 1.1 and give explicit bijections as part of Theorem 4.1. The fact that join-irreducible elements, meet-irreducible elements, and join-irreducible congruences of W are all in bijection is known, and this property of a lattice is called *congruence uniformity* (in the sense of Day [D]). This was proved in [CLM] (where in fact an equivalent property called *boundedness* was established).

The main content of Theorem 1.1 is the unexpectedly deep link between the representation theory of the preprojective algebra and the lattice theory of weak order on the corresponding Weyl group. As part of establishing this link, we have also proved some new results within the representation theory of preprojective algebras which we believe to be of independent interest. Our second main theorem gives two additional algebraic descriptions of layers of Π . We say that a Π -module L is a *brick* if $\text{End}_\Pi(L)$ is a division algebra, and a *stone* if L is a brick satisfying $\text{Ext}_\Pi^1(L, L) = 0$ [HHKU, KL]. Let $\widehat{\Pi}$ be the preprojective algebra of the extended Dynkin type corresponding to Π . Let $\text{D}^b(\text{fd } \widehat{\Pi})$ be the bounded derived category of finite dimensional $\widehat{\Pi}$ -modules. An object $L \in \text{D}^b(\text{fd } \widehat{\Pi})$ is called *2-spherical* if $\text{Hom}_{\text{D}^b(\text{fd } \widehat{\Pi})}(L, L[i])$ has dimension 1 for $i = 0, 2$ and 0 otherwise. (See [ST].)

Theorem 1.2. *The following classes of Π -modules are the same.*



- Layers of Π .
- Bricks of Π .
- Stones of Π .
- Π -modules which are 2-spherical as $\widehat{\Pi}$ -modules,

An observation related to Theorem 1.2 was given by Bolten [Bol] and Sekiya and Yamaura [SY2].

Our third main result concerns the interplay between arrows in the Hasse quiver of W , join-irreducible elements of W , and join-irreducible congruences on W . We refer to Section 2.1 for details about the following notions. Given any arrow $x \rightarrow y$ in the Hasse quiver of an arbitrary finite lattice L , define $\text{con}(x, y)$ to be the smallest congruence on L such that $x \equiv y$. This is a join-irreducible congruence. If j is a join-irreducible element of L , we write j_* for the unique element covered by j in L . The congruence $\text{con}(j, j_*)$ is thus join-irreducible, and it turns out that every join-irreducible element is $\text{con}(j, j_*)$ for some j . When L is the weak order on W , the map $j \mapsto \text{con}(j, j_*)$ is the bijection from join-irreducible elements of W to join-irreducible congruences from Theorem 1.1. Since each Hasse arrow of W specifies a join-irreducible congruence, and since join-irreducible congruences are in bijection with join-irreducible elements, we obtain a labelling of the Hasse arrows of W by join-irreducible elements. We call this the *join-irreducible labelling* of W . Besides this labelling coming from lattice theory, there is a labelling of the Hasse quiver coming from representation theory, namely the *layer labelling*. This labels a Hasse arrow $ws_i \rightarrow w$ by the layer $I(w)/I(ws_i)$. The layer labellings for type A_2 and A_3 are given in Figures 1 and 2.

Theorem 1.3. *The map $j \mapsto I(j_*)/I(j)$ takes the join-irreducible labelling of W to the layer labelling on W . That is, given a Hasse arrow $ws_i \rightarrow w$ labelled by the join-irreducible element j , which covers the element j_* , the layer label on $ws_i \rightarrow w$ is $I(j_*)/I(j)$.*

This is also proved as part of Theorem 4.1, which gives a commutative diagram shown in Figure 3 between important objects. We include this diagram here, although some elements of it have not yet been explained, as a road map to the major

FIGURE 1. Layer labelling for A_2 FIGURE 2. Layer labelling for A_3

results of the paper. The maps are bijections or surjections as marked with tildes “ \sim ” or double-headed arrows.

Given two Hasse arrows $x \rightarrow y$ and $x' \rightarrow y'$ of a lattice L , we say that $x \rightarrow y$ forces $x' \rightarrow y'$ if $\text{con}(x, y) \geq \text{con}(x', y')$ in $\text{Con}(L)$. In other words, $x \rightarrow y$ forces $x' \rightarrow y'$ if every congruence setting $x \equiv y$ also sets $x' \equiv y'$. In the weak order on W , the forcing order on Hasse arrows restricts to a partial order on Hasse arrows of the form $j \rightarrow j_*$ such that j is join-irreducible. We think of this as a partial order on join-irreducible elements and call it the *forcing order* on join-irreducible elements of W ,




FIGURE 3. The correspondences established in Theorem 4.1

We say that a pair of layer modules X, Y form a *doubleton* if $\text{Ext}_{\Pi}^1(Y, X)$ and $\text{Ext}_{\Pi}^1(X, Y)$ are one-dimensional, and the corresponding extensions are again layer modules. We define the *doubleton extension order* on layer modules to be the transitive closure of the relation with $A > B$ if there exists a doubleton A, C such that B is the extension of A by C or of C by A . Our fourth main result is the following,

Theorem 1.4. *The map $j \mapsto I(j_*)/I(j)$ is an isomorphism from the forcing order on join-irreducible elements of W to the doubleton extension order on layer modules of Π .*

In addition to these general results, we show that in type A_n , the doubleton extension order coincides with the reverse of the subfactor order (see Theorem 5.5). Furthermore, we give an explicit combinatorial description of the indecomposable τ -rigid modules in types A_n and D_n in terms of the Young diagrams associated with the join-irreducible elements in the Weyl group W (see Theorems 6.1, 6.5 and 6.12).

2. PRELIMINARIES

In this section, we review the necessary background on lattices, the weak order, and finite-dimensional algebras.

2.1. Lattice-theoretic preliminaries. Proofs and additional details for the material reviewed here can be found in [R2, Sections 9-5 and 9-6].

For any poset P , we say that x covers y , and we write $x \succ y$, if $x > y$, and there is no $z \in P$ such that $x > z > y$. We represent P by its Hasse quiver $\text{Hasse}(P) = (P, \text{Hasse}_1(P))$, whose vertex set is P , and whose arrow set $\text{Hasse}_1(P)$ consists of all arrows $v \rightarrow w$ where v covers w .

Given a subset S of P , if there is a unique smallest element which is greater than or equal to all elements in S , then this least upper bound is called the *join* of S and denoted $\bigvee S$. Similarly, if there is a unique largest element in P that is less than or equal to all elements in S , then this element is called the *meet* of S and denoted $\bigwedge S$. A *lattice* L is a poset in which every pair a, b of elements in L have both a meet $a \wedge b$ and a join $a \vee b$, and a *complete lattice* L is a poset in which every subset S of L has both a meet and a join. (Every finite subset of a lattice L has both a meet and a join, but an infinite lattice fails to be complete if it has some infinite subset without a meet or without a join.)

We restrict our attention to finite lattices in this paper. Some of the assertions made here for finite lattices hold for infinite lattice as well, but some do not.

An element j of a finite lattice L is called *join-irreducible*, whenever $j = a \vee b$ for some $a, b \in L$, either $a = j$ or $b = j$ or both, and j is not the minimum element of L . Equivalently, j is join-irreducible if and only if it covers exactly one element of L . We write j_* for the unique element covered by a join-irreducible element j . Dually, a *meet-irreducible* element of L is an element m that is covered by a unique element m^* . The set of join-irreducible (respectively, meet-irreducible) elements of L is denoted $\text{j-lrr } L$ (respectively, $\text{m-lrr } L$).

A (*lattice*) *congruence* on a lattice L is an equivalence relation Θ having the property that the Θ -class of $a \vee b$ depends only on the Θ -class of a and the Θ -class of b , and having the same property for meets. Given a congruence Θ on L , the set L/Θ of Θ -classes has a well-defined meet and join operation, making L/Θ a lattice called the *quotient* of L modulo Θ .

The set of all equivalence relations on a given set L forms a lattice, where the meet of two relations is given by the intersection of relations and the join of two relations is given by the transitive closure of union of relations. When L is a lattice, the set $\text{Con}(L)$ consisting of congruences of L is a sublattice of the lattice of equivalence relations. Furthermore, $\text{Con}(L)$ is a distributive lattice. We denote by $\text{Con}_{\text{JI}}(L)$ the set of all join-irreducible congruences. As mentioned in the introduction, we have a surjective map $\text{Hasse}_1(L) \rightarrow \text{Con}_{\text{JI}}(L)$ sending an arrow $x \rightarrow y$ to $\text{con}(x, y)$. Here $\text{con}(x, y)$ is the meet, in $\text{Con}(L)$, of all congruences with $x \equiv y$. A congruence Θ on a finite lattice L is determined completely by the set of cover relations $x \succ y$

in L such that $x \equiv y$ modulo Θ . It is also determined uniquely by the set of join-irreducible elements j in L such that $j \equiv j_*$ modulo Θ , and thus we have an injective map $\text{Con}(L) \rightarrow 2^{\text{j-irr}(L)}$.

The map from cover relations $x \succ y$ to join-irreducible congruences is typically not one-to-one. The restriction of the map to cover relations of the form $j \succ j_*$ is also surjective onto join-irreducible congruences, but may still fail to be one-to-one. A lattice is called *congruence uniform* if the map $j \mapsto \text{con}(j, j_*)$ is injective (and thus a bijection) from join-irreducible elements to join-irreducible congruences and the map $m \rightarrow \text{con}(m^*, m)$ is injective (and thus a bijection) from meet-irreducible elements to join-irreducible congruences. A finite congruence uniform lattice is always *semidistributive*. This means that if $x \vee y = x \vee z$ then $x \vee (y \wedge z) = x \vee y$ and if $x \wedge y = x \wedge z$ then $x \wedge (y \vee z) = x \wedge y$.

Since $\text{Con}(L)$ is a finite distributive lattice, the Fundamental Theorem of Finite Distributive Lattices says that its elements are naturally identified with order ideals in the subposet $\text{Con}_{\text{JI}}(L)$ of $\text{Con}(L)$. When L is congruence uniform, the subposet $\text{Con}_{\text{JI}}(L)$ induces a partial order on the join-irreducible elements of L , which we call the *forcing order*. A congruence $\Theta \in \text{Con}(L)$ corresponds to the order ideal in $\text{Con}_{\text{JI}}(L)$ consisting of those join-irreducible congruences below Θ in $\text{Con}(L)$ (i.e. finer than Θ as equivalence relations). These are the join-irreducible congruences $\text{con}(j, j_*)$ such that $j \equiv j_*$ modulo Θ . The forcing order on join-irreducible elements sets $j \leq j'$ if and only if $j \equiv j_*$ modulo $\text{con}(j', j'_*)$.

As mentioned above, each cover relation $x \succ y$ in a finite lattice defines a join-irreducible congruence of L . In a finite congruence uniform lattice L , each join-irreducible congruence is $\text{con}(j, j_*)$ for a unique join-irreducible element j of L . The map $\text{Hasse}_1(L) \rightarrow \text{j-irr}(L)$ sending the arrow $x \rightarrow y$ to the unique j with $\text{con}(j, j_*) = \text{con}(x, y)$ is called the *join-irreducible labelling* of L . Each join-irreducible congruence is also $\text{con}(m^*, m)$ for a unique meet-irreducible element m , and the map $\text{Hasse}_1(L) \rightarrow \text{m-irr}(L)$ sending $x \rightarrow y$ to the unique m with $\text{con}(m^*, m) = \text{con}(x, y)$ is called the *meet-irreducible labelling* of L . These labellings are described explicitly as follows.

The following proposition is [R2, Proposition 9-5.20]. Since that proposition's proof is left to an exercise, we give a proof here.

Proposition 2.1. *Let L be a finite congruence uniform lattice and let $x \rightarrow y$ be an arrow in $\text{Hasse}(L)$.*

- (a) *The join-irreducible label on $x \rightarrow y$ is $j = \bigwedge \{z \in L : z \leq x, z \not\leq y\}$. Furthermore, $j \leq x$ but $j \not\leq y$.*
- (b) *The meet-irreducible label on $x \rightarrow y$ is $m = \bigvee \{z \in L : z \geq y, z \not\geq x\}$. Furthermore, $m \geq y$ but $m \not\geq x$.*

In particular, if j is a join-irreducible element and m is a meet-irreducible element with $\text{con}(j, j_) = \text{con}(m^*, m)$, then $j = \bigwedge \{z \in L : z \leq m^*, z \not\leq m\}$ and $m = \bigvee \{z \in L : z \geq j_*, z \not\geq j\}$.*

Proof. The last statements are special cases of assertions (a) and (b). Assertions (a) and (b) are dual to each other, so by symmetry it is enough to prove (a). To do so, it is enough to show that j is join-irreducible and that $\text{con}(j, j_*) = \text{con}(x, y)$.

Recall that a congruence uniform finite lattice is also semidistributive. Every element z of $\{z \in L : z \leq x, z \not\leq y\}$ has $z \vee y = x$, so applying semidistributivity several times, we see that $j \vee y = x$, so in particular $j \not\leq y$. It is immediate that $j \leq x$. If j covers elements a and b , then $a \leq y$ and $b \leq y$. But if $a \neq b$, then j is a minimal upper bound for a and b , so it must equal $a \vee b$. Since y is another upper bound for a and b , we reach the contradiction $j \leq y$. We conclude that j covers




FIGURE 4. Join-irreducible elements, meet-irreducible elements, and congruences in a finite congruence uniform lattice

at most one element. If j covers no element, then j is the minimal element of L , contradicting again the fact that $j \not\leq y$. We see that j is join-irreducible.

We saw that $j \vee y = x$ and we also verify easily that $j \wedge y = j_*$. If Θ is a congruence with $j \equiv j_*$, then $j \vee y \equiv j_* \vee y$, or in other words $x \equiv y$. Conversely, if Θ has $x \equiv y$, then $j \wedge x \equiv j \wedge y$, or in other words $j \equiv j_*$. We see that Θ has $j \equiv j_*$ if and only if $x \equiv y$, so that $\text{con}(j, j_*) = \text{con}(x, y)$. \square

We summarize some of what we know about join-irreducible elements, meet-irreducible elements, and congruences in a finite congruence uniform lattice in Figure 4. If L is a finite congruence uniform lattice with Hasse quiver $\text{Hasse}(L)$, join-irreducible elements $\text{j-lrr}(L)$, meet-irreducible elements $\text{m-lrr}(L)$, and join-irreducible congruences $\text{Con}_{\text{JI}}(L)$, then the diagram in Figure 4 commutes and the maps are bijections or surjections as marked with tildes “ \sim ” or double-headed arrows.

Let x be an element of a finite lattice L . The expression $x = \bigvee S$ is the *canonical join representation* of x if no proper subset of S joins to x and if every join-representation $x = \bigvee T$ has the property that for all $s \in S$, there exists $t \in T$ with $s \leq t$. An element x may fail to have a canonical join representation, but the canonical join representation of x is unique if it exists. The *canonical meet representation* is defined dually. The semidistributive property of a finite lattice L , described above, is equivalent to the property that every element of L has a canonical join representation and a canonical meet representation. A finite congruence uniform lattice is in particular semidistributive, and canonical join and meet representations can be described in terms of the join-irreducible labelling and meet-irreducible labelling as follows.

The following proposition is [R2, Proposition 9-5.30]. Since that proposition’s proof is also left to an exercise, we give a proof here.

Proposition 2.2. *If L is a finite congruence uniform lattice, then the canonical join representation of $x \in L$ is $x = \bigvee J$, where J is the set of join-irreducible labels on arrows starting at x in $\text{Hasse}(L)$. The canonical meet representation is $x = \bigwedge M$, where M is the set of meet-irreducible labels on arrows ending at x in $\text{Hasse}(L)$.*

Proof. We prove the statement for canonical join representations, using Proposition 2.1 throughout. The other statement is dual.


First, we check that $x = \bigvee J$. On the one hand, each $j \in J$ is below x by Proposition 2.1, so $\bigvee J \leq x$. If $\bigvee J < x$, then there exists y with $x > y \geq \bigvee J$. But the label of $x \rightarrow y$ is in J and is not below y , which is a contradiction.

Next, we show that no proper subset of J joins to x . If $x \rightarrow y$ and $J' = J \setminus \{\bigwedge \{z \in L : z \leq x, z \not\leq y\}\}$, then for any other y' with $x \rightarrow y'$, we have $y \leq x$ by $y \not\leq y'$, so $y \geq \bigwedge \{z \in L : z \leq x, z \not\leq y'\}$. Therefore every element of J' is $\leq y$, so $\bigvee J' < x$.

Finally, we show that, if $x = \bigvee T$ for some $T \subseteq L$, then for all $j \in J$, there exists $t \in T$ with $j \leq t$. Let y have $x \rightarrow y$ and $j = \bigwedge \{z \in L : z \leq x, z \not\leq y\}$. Every element of T is $\leq x$, and if every element of T is $\leq y$, then $\bigvee T \leq y$, contradicting the supposition that $x = \bigvee T$. Thus there exists some element t of T with $t \leq x$ and $t \not\leq y$, and this element is above j by definition. \square

A *polygon* in a finite lattice L is an interval $[x, y]$ such that $\{z \in L : x < z < y\}$ consists of two disjoint nonempty chains. (Thus the Hasse quiver of $[x, y]$ is a cycle with one source and one sink.) The lattice L is *polygonal* if the following two conditions hold: First, if distinct elements y_1 and y_2 both cover an element x , then $[x, y_1 \vee y_2]$ is a polygon; and second, if an element y covers distinct elements x_1 and x_2 , then $[x_1 \wedge x_2, y]$ is a polygon.

If L is a polygonal lattice, then we define a quiver $\text{FPoly}(L)$ whose set of vertices is $\text{Hasse}_1(L)$, with arrows defined in every polygon P of L as follows. The two arrows into the bottom element of P are called *bottom arrows*, while the two arrows from the top element of P are called *top arrows*, and all other arrows of P are called *side arrows*. Every bottom arrow of P has an arrow (in $\text{FPoly}(L)$) to the opposite top arrow in P (i.e. the top arrow in the opposite chain) and has an arrow to every side arrow in P . Every top arrow of P has an arrow to the opposite bottom arrow and every side arrow in P . For example, a square and hexagon in L would contribute to $\text{FPoly}(L)$ as indicated below.



The following is [R2, Theorem 9-6.5].

Theorem 2.3. *If L is a finite polygonal lattice, and $x \rightarrow y$ and $x' \rightarrow y'$ are arrows in $\text{Hasse}(L)$, then $\text{con}(x', y') \leq \text{con}(x, y)$ if and only if there is a directed path from $x \rightarrow y$ to $x' \rightarrow y'$ in $\text{FPoly}(L)$.*

When L is also congruence uniform, the quiver $\text{FPoly}(L)$ is closely related to the forcing order on join-irreducible elements of L , as described in the following corollary, which is immediate by combining Theorem 2.3 with facts about finite congruence uniform lattices already given in this section. We say that a quiver Q is *strongly connected* if given any two vertices x and y of the quiver, there exists a path from x to y and a path from y to x . A *strongly connected component* of Q is a set of vertices of Q that is maximal with respect to inducing a strongly connected subquiver.

Corollary 2.4. *Suppose L is a finite, polygonal, congruence uniform lattice. Then each strongly connected component of $\text{FPoly}(L)$ contains exactly one arrow in $\text{Hasse}(L)$ of the form $j \rightarrow j_*$ where j is join-irreducible. This bijection between join-irreducible*

elements of L and strongly connected components of $\text{FPoly}(L)$ is an isomorphism from the forcing order on join-irreducible elements of L to the partial order induced by $\text{FPoly}(L)$ on its strongly connected components.

Informally, the corollary says that all forcing in a finite polygonal, congruence uniform lattice comes from forcing in polygons. Indeed, the notation $\text{FPoly}(L)$ suggests the phrase “forcing in polygons.”

We define another quiver $\text{SFPoly}(L)$, again with the set of vertices given by $\text{Hasse}_1(L)$, but with strictly fewer arrows than $\text{FPoly}(L)$. The notation $\text{SFPoly}(L)$ suggests the phrase “strong forcing in polygons.” In $\text{SFPoly}(L)$, every bottom arrow of a polygon P has an arrow only to the opposite top arrow in P and every top arrow of P has an arrow only to the opposite bottom arrow in P . Thus from the square and hexagon as labelled above, $\text{SFPoly}(L)$ gets arrows as indicated below.

$$a \leftrightarrow d \quad b \leftrightarrow c \quad e \leftrightarrow j \quad g \leftrightarrow h$$

Strong forcing in polygons controls whether two Hasse arrows in a finite polygonal, congruence uniform lattice determine the same congruence, as described in the following corollary.

Corollary 2.5. *Suppose L is a finite polygonal, congruence uniform lattice and let $x \rightarrow y$ and $x' \rightarrow y'$ be arrows of $\text{Hasse}(L)$. Then $\text{con}(x, y) = \text{con}(x', y')$ if and only if there is a directed path (or equivalently a path) in $\text{SFPoly}(L)$ from $x \rightarrow y$ to $x' \rightarrow y'$.*

Proof. The “if” direction is immediate by Theorem 2.3, the fact that every arrow is $\text{SFPoly}(L)$ is also an arrow in $\text{FPoly}(L)$, and the fact that arrows in $\text{SFPoly}(L)$ come in opposite pairs. We prove the converse by proving that for every Hasse arrow $x \rightarrow y$, there is a directed path in $\text{SFPoly}(L)$ from $x \rightarrow y$ to $j \rightarrow j_*$, where j is the unique join-irreducible element with $\text{con}(j, j_*) = \text{con}(x, y)$. Proposition 2.1 says in particular that $x \geq j$, so we can argue by induction on the length of a longest maximal chain from j to x . Choose z with $x \succ z \geq j$ and set $y' = y \wedge z$. Since L is polygonal, $[y', x]$ is a polygon P , and $x \rightarrow y$ is a top arrow in P . Choosing x' so that $x' \rightarrow y'$ is the bottom arrow opposite $x \rightarrow y$ (i.e. in the other chain of P), we have $\text{con}(x' \rightarrow y') = \text{con}(x \rightarrow y)$ by Theorem 2.3. Thus $\text{con}(j, j_*) = \text{con}(x', y')$, and thus $x' \geq j$. By induction, there is a directed path in $\text{SFPoly}(L)$ from $x' \rightarrow y'$ to $j \rightarrow j_*$, and using an arrow in P , we obtain a directed path in $\text{SFPoly}(L)$ from $x \rightarrow y$ to $j \rightarrow j_*$. \square

2.2. Weak order preliminaries. Fix a simply-laced Dynkin type (i.e., one of A_n for $n \geq 1$, D_n for $n \geq 4$, E_6 , E_7 , or E_8). Let W be the finite Weyl group of that type. For background on Weyl groups, see [BB].

Let $S = \{s_1, \dots, s_n\}$ be the set of simple reflections of W . By definition, any element w of W can be written as a product of the simple reflections. Such an expression for w of minimal length is called *reduced*. The length of a reduced expression is the *length* of w , denoted $\ell(w)$.

The (*right*) *weak order* on W is the partial order with $u \geq v$ if $\ell(u) = \ell(v) + \ell(v^{-1}u)$. We write $\text{Hasse}(W)$ for the Hasse quiver of the weak order on W . The arrows of $\text{Hasse}(W)$ are all arrows $ws_i \rightarrow w$ such that $w \in W$, $s_i \in S$, and $\ell(ws_i) > \ell(w)$, or equivalently $\ell(ws_i) = \ell(w) + 1$. For $w \in W$, we denote by w^+ (respectively, w^-) the set of arrows in $\text{Hasse}(W)$ ending (respectively, starting) at w .

The weak order on W is a finite lattice. In particular, it has a maximal element, denoted w_0 and often called the *longest element* of W . For our purposes, the most important properties of the weak order are the following.

Theorem 2.6. *W is congruence uniform [CLM] and polygonal [R3, Theorem 10-3.7].*

2.3. Algebraic preliminaries. Fix a base field k , and let A be a finite-dimensional k -algebra. We write $\text{mod } A$ for the finite-dimensional left A -modules.

We say that a full subcategory \mathcal{T} of $\text{mod } A$ is a *torsion class* if it is closed under factors, isomorphisms and extensions. *Torsion-free classes* are defined dually. We write $\text{tors } A$ for the torsion classes of A , and $\text{torf } A$ for its torsion-free classes. We view $\text{tors } A$ and $\text{torf } A$ as posets under the inclusion order. Then we have an anti-isomorphism given by $\mathcal{T} \mapsto \mathcal{T}^\perp = \{X \in \text{mod } A \mid \text{Hom}_A(\mathcal{T}, X) = 0\}$

$$\text{tors } A \rightarrow \text{torf } A,$$

whose inverse is given by $\mathcal{F} \mapsto {}^\perp \mathcal{F} = \{X \in \text{mod } A \mid \text{Hom}_A(X, \mathcal{F}) = 0\}$.

We recall that a torsion class \mathcal{T} of $\text{mod } A$ is *functorially finite* if there exists $M \in \text{mod } A$ such that $\mathcal{T} = \text{Fac } M$, where $\text{Fac } M$ is the full subcategory of $\text{mod } A$ consisting of factor modules of finite direct sums of copies of M [AS]. We denote by $\text{f-tors } A$ (respectively, $\text{f-torf } A$) the set of all functorially finite torsion (respectively, torsionfree) classes in $\text{mod } A$. We view $\text{f-tors } A$ also as a poset under inclusion. The above anti-isomorphism restricts to an anti-isomorphism $\text{f-tors } A \rightarrow \text{f-torf } A$.

There is a bijection between $\text{f-tors } A$ and a certain class of A -modules. Recall that a module $M \in \text{mod } A$ is called τ -*rigid* if $\text{Hom}_A(M, \tau M) = 0$, and τ^- -*rigid* if $\text{Hom}_A(\tau^- M, M) = 0$. A module $M \in \text{mod } A$ is called τ -*tilting* if it is τ -rigid and $|M| = |A|$ holds, where $|M|$ is the number of non-isomorphic indecomposable direct summands of M . A module $M \in \text{mod } A$ is called *support τ -tilting* if there exists an idempotent e of A such that M is a τ -tilting $(A/\langle e \rangle)$ -module. We denote by $\text{s}\tau\text{-tilt } A$ the set of isomorphism classes of basic support τ -tilting A -modules, and by $\text{i}\tau\text{-rigid } A$ (respectively, $\text{i}\tau^- \text{-rigid } A$) the set of isomorphism classes of indecomposable τ -rigid (respectively, τ^- -rigid) A -modules. (See [AIR] for more background on these notions.) By [AIR, 2.7], we have a surjection $\{\tau\text{-rigid } A\text{-modules}\} \rightarrow \text{f-tors } A$ given by $M \mapsto \text{Fac } M$, which induces a bijection

$$\text{s}\tau\text{-tilt } A \xrightarrow{\sim} \text{f-tors } A.$$

Recall that A is τ -*tilting finite* if $\text{s}\tau\text{-tilt } A$ is a finite set, or equivalently, $\text{i}\tau\text{-rigid } A$ is a finite set. It is shown in [DIJ] and [IRTT, 0.2] that the following conditions are equivalent.

- A is τ -tilting finite
- $\text{f-tors } A$ is a finite set.
- $\text{f-tors } A$ (respectively, $\text{f-torf } A$) forms a complete lattice.
- $\text{f-tors } A = \text{tors } A$.

Via the bijection between $\text{f-tors } A$ and $\text{s}\tau\text{-tilt } A$, we obtain a partial order on $\text{s}\tau\text{-tilt } A$. We refer to the poset on $\text{s}\tau\text{-tilt } A$ as *generation order*. The arrows of the Hasse diagram of this poset are *mutations* (see [AIR, Theorem 0.6]).

Using τ -tilting theory, we have the following description of join-irreducible elements in $\text{tors } A$.

Theorem 2.7. *Let A be a finite dimensional k -algebra which is τ -tilting finite. Then we have a bijection given by $M \mapsto \text{Fac } M$*

$$\text{i}\tau\text{-rigid } A \rightarrow \text{j-lrr}(\text{tors } A).$$

Proof. Since A is τ -tilting finite, we have $\text{s}\tau\text{-tilt } A \simeq \text{tors } A$. For $\mathcal{T} \in \text{tors } A$, we take $M = M_1 \oplus \cdots \oplus M_n \in \text{s}\tau\text{-tilt } A$ such that $\mathcal{T} = \text{Fac } M$. It is shown in [AIR] that adjacent vertices to \mathcal{T} in the Hasse quiver of $\text{tors } A$ are given by $\text{Fac } \mu_k(M)$ for $1 \leq k \leq n$, where $\mu_k(M)$ is the mutation of M at M_k . Clearly $\text{Fac } \mu_k(M) \subset \mathcal{T}$ holds if and only if $M_k \in \text{Fac}(M/M_k)$. Therefore \mathcal{T} is join-irreducible if and only if there exists a unique k satisfying $\text{Fac } \mu_k(M) \subset \mathcal{T}$. This is equivalent to $\mathcal{T} = \text{Fac } M_k$. Since M_k is an indecomposable τ -rigid A -module, we have the assertion. \square

For a complete lattice L , we denote by $\text{cj-lrr}L$ the set of *completely join-irreducible elements*, that is, elements $a \in L$ such that $a = \bigvee S$ for a subset S of L implies $a \in S$. If we drop the τ -tilting finiteness assumption on A , then we still have a bijection $\text{i}\tau\text{-rigid } A \rightarrow \text{f-tors } A \cap \text{cj-lrr}(\text{tors } A)$ given by $M \mapsto \text{Fac } M$. The proof is the same, we only need to use [DIJ, Theorem 3.1]. Note that completely join-irreducible elements in $\text{tors } A$ are not necessarily functorially finite. For example, consider $\text{mod } kQ$ for a Kronecker quiver Q . Then all preinjective modules together with one tube form such a torsion class.

We denote by $\text{brick } A$ the set of isomorphism classes of bricks of A . A full subcategory \mathscr{W} of $\text{mod } A$ is called *wide* if it is closed under kernels, cokernels and extensions. In this case, \mathscr{W} forms an abelian category and the inclusion functor $\mathscr{W} \rightarrow \text{mod } A$ is exact. A wide subcategory \mathscr{W} is called *local* if it contains a unique simple object up to isomorphism. We denote by $\text{l-wide } A$ the set of local wide subcategories of $\text{mod } A$. We have the following easy observation.

Proposition 2.8. *Let A be a finite dimensional k -algebra. Then we have a bijection*

$$\text{brick } A \rightarrow \text{l-wide } A \quad \text{given by } S \mapsto \text{Filt } S,$$

where $\text{Filt } S$ consists of A -modules X which have a filtration $X = X_0 \supset X_1 \supset \cdots \supset X_{\ell-1} \supset X_{\ell} = 0$ with $\ell \geq 0$ such that $X_i/X_{i+1} \simeq S$ for any $0 \leq i < \ell$.

Proof. Let S be a brick of A . Then $\text{Filt } S$ is a wide subcategory by [Ri, 1.2]. Clearly $\text{Filt } S$ has a unique simple object in S . Conversely, let \mathscr{W} be a local wide subcategory of $\text{mod } A$ with a simple object S . Clearly S is a brick and we have $\mathscr{W} = \text{Filt } S$. \square

2.4. Preliminaries on preprojective algebras. Let $\Pi = \Pi(W)$ be a preprojective algebra of the same Dynkin type as W . To construct Π , take the Dynkin diagram and replace each edge by a pair of opposite arrows a, a^* to obtain the quiver \bar{Q} . Then $\Pi(W)$ is the path algebra of \bar{Q} modulo the ideal generated by $\sum_a (aa^* - a^*a)$. It is a finite-dimensional self-injective algebra. We write S_i for the simple module corresponding to vertex i .

We let e_i be the idempotent corresponding to the vertex i . Let I_i be the two-sided ideal $\Pi(1 - e_i)\Pi$. It is maximal as a left ideal and as a right ideal. For each $w \in W$, we take a reduced word $w = s_{i_1} \cdots s_{i_k}$, and we define

$$I(w) = I_{i_1} \cdots I_{i_k}, \quad \mathsf{T}(w) := \text{Fac } I(w) \quad \text{and} \quad \mathsf{F}(w) := \text{Sub}(\Pi/I(w)).$$

Here $\text{Sub } X$ refers to the subcategory consisting of subobjects of direct sums of copies of X .

The following result due to Mizuno is fundamental.

Theorem 2.9. [Mi, 2.14, 2.21]


- (a) $I(w)$ does not depend on the choice of the reduced word for w .
- (b) We have bijections given by $w \mapsto I(w) \mapsto \mathsf{T}(w)$:

$$W \xrightarrow{\sim} \text{s}\tau\text{-tilt } \Pi \xrightarrow{\sim} \text{tors } \Pi. \tag{2.1}$$

- (c) The bijection above from W to $\text{tors } \Pi$ is an *anti-isomorphism* from weak order on W to inclusion order on torsion classes.

We remark that Mizuno uses right modules, while we use left modules, and therefore his results need to be suitably translated to account for this difference. (In particular, this has the effect that he uses left weak order while we use right weak order.) We also remark here that [AIR] uses right modules, but writes I_w for the ideal which we would refer to as $I_{w^{-1}}$.

Thus $I(e) = \Pi$ gives the maximum torsion class $\mathsf{T}(e) = \text{mod } \Pi$ and $I(w_0) = 0$ gives the minimum torsion class $\mathsf{T}(w_0) = \{0\}$, where $e \in W$ is the identity and $w_0 \in W$ is the longest element.

FIGURE 5. The weak order and $s\tau$ -tilt Π in type A_2

The following is an easy consequence of (2.1).

Lemma 2.10. *Let $w, v \in W$. Then $\ell(wv) = \ell(w) + \ell(v)$ holds if and only if $I(wv) = I(w)I(v)$ holds. If this holds, then we have $I(wv) = I(w)I(v) = I(w) \otimes_{\Pi} I(v)$.*

We show a few examples. Here, 1_2_3 , for example, refers to the indecomposable $\overline{\Pi}$ -module with composition factors S_1, S_2, S_3 from top to bottom.

Example 2.11. The left picture in Figure 5 shows the weak order on permutations in S_3 (the Weyl group of type A_2). The right picture in Figure 5 shows the set $s\tau$ -tilt Π for Π of type A_2 , arranged in generation order. We show in red the meet-irreducible elements of W and the join-irreducible elements of $s\tau$ -tilt Π . The two pictures are arranged so that the map $w \rightarrow I(w)$ is accomplished by a translation of the page.

Example 2.12. Figure 6 shows the weak order on permutations in S_4 (the Weyl group of type A_3) and $s\tau$ -tilt Π for Π of type A_3 , with the same conventions as described in Example 2.11.

3. HOMOLOGICAL CHARACTERIZATIONS OF LAYERS

In this section we prove Theorem 1.2, the homological characterization of layer modules of the preprojective algebra Π . The assertion that every layer module is a stone was shown in [AIRT, 2.3]. It is also easy to show that the stones of Π are exactly the Π -modules which are 2-spherical as $\widehat{\Pi}$ -modules. Indeed, for any Π -module L , we have $\text{Ext}_{\widehat{\Pi}}^1(L, L) = \text{Ext}_{\Pi}^1(L, L)$ and $\text{Ext}_{\widehat{\Pi}}^2(L, L) = D\text{End}_{\Pi}(L)$, and the assertion follows.

We show that any brick L of Π is a stone. Let $\langle -, - \rangle$ be the Euler form on $K_0(\text{fd } \widehat{\Pi})$. For $x = (x_i)_{i \in Q_0} \in \mathbb{Z}^{Q_0}$, we have

$$\langle x, x \rangle = 2 \left(\sum_{i \in Q_0} x_i^2 - \sum_{(i \rightarrow j) \in Q_1} x_i x_j \right),$$

which is a non-negative even integer. Since the restriction of $\langle -, - \rangle$ to $K_0(\text{mod } \Pi)$ is positive definite, we have

$$2 - \dim_k \text{Ext}_{\Pi}^1(L, L) = \sum_{i=0}^2 (-1)^i \dim_k \text{Ext}_{\Pi}^i(L, L) = \langle \underline{\dim} L, \underline{\dim} L \rangle \geq 2,$$

which implies $\text{Ext}_{\Pi}^1(L, L) = 0$. Thus L is a stone.

To complete the proof, we will show that every stone is a layer. As before, $D^b(\text{fd } \widehat{\Pi})$ is the bounded derived category of finite dimensional $\widehat{\Pi}$ -modules. Let $\widehat{I}_i := \widehat{\Pi}(1 - e_i)\widehat{\Pi}$ be an ideal of $\widehat{\Pi}$. Then we have an autoequivalence

$$F_i := \widehat{I}_i \overset{\mathbf{L}}{\otimes}_{\widehat{\Pi}} - : D^b(\text{fd } \widehat{\Pi}) \rightarrow D^b(\text{fd } \widehat{\Pi})$$




FIGURE 6. The weak order and $s\tau$ -tilt Π in type A_3

with quasi-inverse

$$F_i^{-1} := \mathbf{R}\mathrm{Hom}_{\widehat{\Pi}}(\widehat{I}_i, -) : \mathrm{D}^b(\mathrm{fd} \widehat{\Pi}) \rightarrow \mathrm{D}^b(\mathrm{fd} \widehat{\Pi}).$$

We identify Π -modules with $\widehat{\Pi}$ -modules annihilated by e_0 , where e_0 is an idempotent of $\widehat{\Pi}$ satisfying $\Pi = \widehat{\Pi}/\langle e_0 \rangle$.

The following assertions are easy to check.

Lemma 3.1. *Let L be a finite-dimensional Π -module.*

- (a) *We have $I_i = \widehat{I}_i \otimes_{\widehat{\Pi}} \Pi = \Pi \otimes_{\widehat{\Pi}} \widehat{I}_i$.*
- (b) *We have $I_i \otimes_{\Pi} L = \widehat{I}_i \otimes_{\widehat{\Pi}} L$ and $\text{Hom}_{\Pi}(I_i, L) = \text{Hom}_{\widehat{\Pi}}(\widehat{I}_i, L)$.*

The following observation plays a key role.

Proposition 3.2. *Let L be a stone of Π and i a vertex of \overline{Q} .*

- (a) *If $L \not\cong S_i$, then at least one of $\text{Hom}_{\Pi}(S_i, L) = 0$ or $\text{Hom}_{\Pi}(L, S_i) = 0$ holds.*
- (b) *If $\text{Hom}_{\Pi}(S_i, L) = 0$, then $F_i(L) \simeq I_i \otimes_{\Pi} L$ is a stone of Π .*
- (c) *If $\text{Hom}_{\Pi}(L, S_i) = 0$, then $F_i^{-1}(L) \simeq \text{Hom}_{\Pi}(I_i, L)$ is a stone of Π .*

Proof. We already observed that stones of Π are precisely Π -modules which are 2-spherical as $\widehat{\Pi}$ -modules.

(a) If both $\text{Hom}_{\Pi}(S_i, L)$ and $\text{Hom}_{\Pi}(L, S_i)$ are non-zero, the composition $L \rightarrow S_i \rightarrow L$ is a non-zero non-isomorphic endomorphism of L . This is a contradiction since $\text{End}_{\Pi}(L) = k$.

(b) Since F_i is an autoequivalence of $\text{D}^b(\text{fd } \widehat{\Pi})$, we have that $F_i(L)$ is a 2-spherical object. We show that it is a $\widehat{\Pi}$ -module, that is, $\text{Tor}_j^{\widehat{\Pi}}(\widehat{I}_i, L) = 0$ holds for any $j \neq 0$. Since the $\widehat{\Pi}$ -module \widehat{I}_i has projective dimension at most one [BIRS], we only have to check the case $j = 1$. Since

$$\text{Tor}_1^{\widehat{\Pi}}(\widehat{I}_i, L) \simeq \text{Tor}_2^{\widehat{\Pi}}(S_i, L) \simeq D \text{Ext}_{\widehat{\Pi}}^2(L, S_i) \simeq \text{Hom}_{\widehat{\Pi}}(S_i, L) = 0$$

holds by our assumption, we have the assertion. Thus we have $F_i(L) \simeq \widehat{I}_i \otimes_{\widehat{\Pi}} L$, which is isomorphic to $I_i \otimes_{\Pi} L$ by Lemma 3.1.

(c) Similar to (b). \square

We need the following observation.

Lemma 3.3. *Let $w \in W$ and let L be a stone of Π which belongs to $\mathsf{T}(w)$. For any i satisfying $\text{Hom}_{\Pi}(L, S_i) \neq 0$, the following assertions hold.*

- (a) *$\ell(s_i w) > \ell(w)$ and $I(s_i w) = I_i I(w) = I_i \otimes_{\Pi} I(w)$.*
- (b) *If $L \not\cong S_i$, then $\text{Hom}_{\Pi}(S_i, L) = 0$ and $F_i(L)$ is a stone of Π which belongs to $\mathsf{T}(s_i w)$.*

Proof. (a) Since $L \in \mathsf{T}(w)$ and $\text{Hom}_{\Pi}(L, S_i) \neq 0$, we have $\text{Hom}_{\Pi}(I(w), S_i) \neq 0$. Thus $I_i I(w) \neq I(w)$ holds. Thus we have the assertions by Lemma 2.10.

(b) By Proposition 3.2, $\text{Hom}_{\Pi}(S_i, L) = 0$ holds, and $F_i(L)$ is a stone of Π . Since we have a surjection $I(w)^{\oplus m} \rightarrow L$, we have a surjection $I(s_i w)^{\oplus m} = I_i \otimes_{\Pi} I(w)^{\oplus m} \rightarrow I_i \otimes_{\Pi} L = F_i(L)$. Thus $F_i(L)$ belongs to $\mathsf{T}(s_i w)$. \square

For $w \in W$, we have an ideal $\widehat{I}(w)$ of $\widehat{\Pi}$ satisfying $\widehat{I}(w) \supseteq \langle e_0 \rangle$ and $\widehat{I}(w)/\langle e_0 \rangle = I(w)$. Let

$$F(w) := \widehat{I}(w) \otimes_{\widehat{\Pi}}^{\mathbf{L}} - : \text{D}^b(\text{fd } \widehat{\Pi}) \rightarrow \text{D}^b(\text{fd } \widehat{\Pi}).$$

For any reduced expression $w = s_{i_1} \cdots s_{i_\ell}$, we have

$$\widehat{I}(w) = \widehat{I}_{i_1} \cdots \widehat{I}_{i_\ell} \quad \text{and} \quad F(w) = F_{i_1} \circ \cdots \circ F_{i_\ell}.$$

Lemma 3.4. *Let L be a stone of Π . Then there exists $v \in W$ such that $F(v)(L)$ is a simple Π -module S_i .*

Proof. Assume that L is not simple. Taking i_1 such that $\text{Hom}_{\Pi}(L, S_{i_1}) \neq 0$ and applying Lemma 3.3 to the stone $L \in \text{mod } \Pi = \mathsf{T}(e)$, we have that $F_{i_1}(L)$ is a stone in $\text{Fac } I_{i_1}$.

Assume that $F_{i_1}(L)$ is not simple. Taking i_2 such that $\text{Hom}_{\Pi}(F_{i_1}(L), S_{i_2}) \neq 0$ and applying Lemma 3.3 to the stone $F_{i_1}(L) \in \text{Fac } I_{i_1}$, we have that $F_{i_2}F_{i_1}(L)$ is a stone in $\text{Fac}(I_{i_2}I_{i_1})$ and $\ell(s_{i_2}s_{i_1}) = 2$ holds.

We repeat this process. Since the lengths of elements in W are bounded by $\ell(w_0)$, the process must stop, that is, there exists $v \in W$ such that $F(v)(L)$ is a simple Π -module S_i . \square

We need the following general observation.

Lemma 3.5. *For any $v, w \in W$ satisfying $I(v) \supset I(w)$, we have an isomorphism $D(I(v)/I(w)) \simeq I(w_0w^{-1})/I(w_0v^{-1})$ of Π -modules.*

Proof. Applying D to the exact sequence $0 \rightarrow I(w) \rightarrow I(v) \rightarrow I(v)/I(w) \rightarrow 0$ of Π^{op} -modules, we have an exact sequence

$$0 \rightarrow D(I(v)/I(w)) \rightarrow D(I(v)) \rightarrow D(I(w)) \rightarrow 0 \quad (3.1)$$

of Π -modules. On the other hand, we have an exact sequence

$$0 \rightarrow I(w_0w^{-1})/I(w_0v^{-1}) \rightarrow \Pi/I(w_0v^{-1}) \rightarrow \Pi/I(w_0w^{-1}) \rightarrow 0 \quad (3.2)$$

of Π -modules. Using an isomorphism $D(I(w)) \simeq \Pi/I(w_0w^{-1})$ of Π -modules for any $w \in W$ [ORT] and comparing (3.1) and (3.2), we have the assertion. \square

Now we are ready to finish the proof of Theorem 1.2.

Proof of Theorem 1.2. As shown in the first paragraph of this section, we can complete the proof by showing that every stone is a layer. Let L be a stone of Π . By Lemma 3.4, there exists $v \in W$ such that $F(v)(L) = \widehat{I}(v) \otimes_{\widehat{\Pi}}^{\mathbf{L}} L$ is a simple Π -module S_i . Thus L equals

$$\begin{aligned} \mathbf{R}\text{Hom}_{\widehat{\Pi}}(\widehat{I}(v), S_i) &= \text{Hom}_{\widehat{\Pi}}(\widehat{I}(v), S_i) \stackrel{\text{Lem.3.1}}{=} \text{Hom}_{\Pi}(I(v), S_i) = D(DS_i \otimes_{\Pi} I(v)) \\ &= D((\Pi/I_i) \otimes_{\Pi} I(v)) = D(I(v)/I_i I(v)) \stackrel{\text{Lem.2.10}}{=} D(I(v)/I(s_i v)). \end{aligned}$$

This is a layer of Π by Lemma 3.5. \square

4. BIJECTIONS: THEOREMS 1.1 AND 1.3

Throughout this section, let Π be a preprojective algebra of Dynkin type and W the corresponding Weyl group. The main result of this section is that a certain diagram is commutative. Commutativity of that diagram includes Theorems 1.1 and 1.3.

For an arrow $a : ws_i \rightarrow w$ in H , we have a natural inclusion $I(w) \supset I(ws_i)$ of ideals of Π , and we associate to a the Π -module

$$L(a) := I(w)/I(ws_i) = (I(w)e_i)/(I(w)I_i e_i),$$

where the right equality follows from $I(w) = I(w)e_i \oplus I(w)(1 - e_i)$, $I(ws_i) = I(w)I_i e_i \oplus I(w)I_i(1 - e_i)$ and $I_i(1 - e_i) = \Pi(1 - e_i)$. We also define maps $L : \mathfrak{m}\text{-lrr } W \rightarrow \text{layer } \Pi$ and $M : \mathfrak{m}\text{-lrr } W \rightarrow \text{i}\tau\text{-rigid } \Pi$ as

$$L(m) := L(a) \quad \text{and} \quad M(m) := I(m)e_i$$

for $m \in \mathfrak{m}\text{-lrr } W$ and $a : m^* = ms_i \rightarrow m$ in H .

Theorem 4.1. *Let Π be a preprojective algebra of Dynkin type and W the corresponding Weyl group. Then we have the commutative diagram shown in Figure 3. The maps are bijections or surjections as marked with tildes “ \sim ” or double-headed arrows.*

We start with the following simple observation.

Lemma 4.2. $L(m) = M(m)/\text{rad } M(m)_{\text{End}_{\Pi}(M(m))}$ holds for any $m \in \mathfrak{m}\text{-lrr } W$.

Proof. We have $M(m) = I(m)e_i$ and $L(m) = I(m)e_i/I(m)I_i e_i$. Thus we only have to show $I(m)I_i e_i = (I(m)e_i) \text{rad } \text{End}_{\Pi}(I(m)e_i)$.

Since $\mathbb{T}(m) = \text{Fac } I(m)$ is join-irreducible, we have $I(m) \in \text{Fac}(I(m)e_i)$. Thus

$$\begin{aligned} (I(m)e_i) \text{rad } \text{End}_{\Pi}(I(m)e_i) &= \sum_{f \in \text{rad } \text{End}_{\Pi}(I(m)e_i)} \text{Im}(f : I(m)e_i \rightarrow I(m)e_i) \\ &= \sum_{g \in \text{rad}_{\Pi}(I(m), I(m)e_i)} \text{Im}(g : I(m) \rightarrow I(m)e_i), \end{aligned}$$

where rad_{Π} is the radical of the category $\text{mod } \Pi$ and hence $\text{rad}_{\Pi}(I(m), I(m)e_i)$ consists of morphisms which are not split epimorphisms.

By [Mi, Lemma 2.7], we have a surjection $\Pi \rightarrow \text{End}_{\Pi}(I(m))$ given by $x \mapsto (y \mapsto yx)$. This induces surjections $\Pi e_i \rightarrow \text{Hom}_{\Pi}(I(m), I(m)e_i)$ and $I_i e_i \rightarrow \text{rad}_{\Pi}(I(m), I(m)e_i)$. Therefore we have


$$\sum_{g \in \text{rad}_{\Pi}(I(m), I(m)e_i)} \text{Im}(g : I(m) \rightarrow I(m)e_i) = I(m)I_i e_i,$$

which completes the proof. \square

In Section 2.1 (particularly Theorem 2.3 and Corollary 2.4), we described how the forcing order on join-irreducible elements interacts with the polygons (squares and rectangles) in W . One ingredient in the proof of Theorem 4.1 is a similar description of how the layer labelling interacts with polygons.

Recall from the introduction that a pair of layer modules X, Y form a *doubleton* if $\text{Ext}_{\Pi}^1(Y, X)$ and $\text{Ext}_{\Pi}^1(X, Y)$ are one-dimensional, and the corresponding extensions are again layer modules. Recall also that the *doubleton extension order* on layer modules is the transitive closure of the relation with $A > B$ if there exists a doubleton A, C such that B is the extension of A by C or of C by A .

Proposition 4.3. *For a polygon in W (necessarily a square or hexagon), the layer labelling has the following configuration:*



Moreover, in the hexagon case the layers X and Y form a doubleton and there exist short exact sequences of Π -modules:

$$0 \rightarrow X \rightarrow E \rightarrow Y \rightarrow 0 \quad \text{and} \quad 0 \rightarrow Y \rightarrow F \rightarrow X \rightarrow 0.$$

Thus $X \geq E$, $X \geq F$, $Y \geq E$, and $Y \geq F$ in the doubleton extension order.

Proof. Of the two diagrams in the statement of the theorem, the hexagon occurs if and only if i and j are neighbouring in \overline{Q} . We argue the hexagon case. The square case is similar but simpler. What we need to show is the following:

(i) There are isomorphisms of Π -modules:

$$I(u)/I(u)I_i \simeq I(u)I_j I_i / I(u)I_i I_j I_i \quad \text{and} \quad I(u)/I(u)I_j \simeq I(u)I_i I_j / I(u)I_i I_j I_i.$$

(ii) There are short exact sequences of Π -modules:

$$\begin{aligned} 0 \rightarrow I(u)/I(u)I_i \rightarrow I(u)I_i / I(u)I_i I_j \rightarrow I(u)I_i I_j / I(u)I_i I_j I_i \rightarrow 0, \\ 0 \rightarrow I(u)/I(u)I_j \rightarrow I(u)I_j / I(u)I_j I_i \rightarrow I(u)I_j I_i / I(u)I_i I_j I_i \rightarrow 0. \end{aligned}$$

We first show (i). The partial order tells us, by Lemma 2.10, that we have

$$I(u)/I(u)I_i = I(u) \otimes_{\Pi} (\Pi/I_i) \quad \text{and} \quad I(u)I_jI_i/I(u)I_iI_jI_i = I(u) \otimes_{\Pi} (I_jI_i/I_iI_jI_i).$$

Since $\Pi/I_i \simeq S_i \simeq I_jI_i/I_iI_jI_i$ holds by an easy calculation for the preprojective algebra $\Pi/I_iI_jI_i$ of type A_2 (see Figure 5), we have

$$I(u)/I(u)I_i = I(u) \otimes_{\Pi} (\Pi/I_i) \simeq I(u) \otimes_{\Pi} (I_jI_i/I_iI_jI_i) = I(u)I_jI_i/I(u)I_iI_jI_i.$$

The other isomorphism follows by interchanging i and j .

Now we show (ii). Again by an easy calculation for the preprojective algebra $\Pi/I_iI_jI_i$ of type A_2 (see Figure 5), we have $\Pi/I_i \simeq S_i$, $I_i/I_iI_j \simeq \binom{S_j}{S_i}$ and $I_iI_j/I_iI_jI_i \simeq S_j$. Thus there exists an exact sequence

$$0 \rightarrow \Pi/I_i \rightarrow I_i/I_iI_j \rightarrow I_iI_j/I_iI_jI_i \rightarrow 0$$

of Π -modules. Applying $I(u) \otimes_{\Pi} -$, we obtain the first sequence. The second one follows by interchanging i and j .

We verify that X and Y form a doubleton in the hexagon case. If the labels are simple, then the extension groups are certainly one-dimensional, and the extensions are layers. Any hexagon is obtained by applying $I(u) \otimes_{\Pi} -$ to such a hexagon, as above. This implies the desired result once we note that, by Lemma 3.1, we can instead consider applying $\widehat{I}(u) \otimes_{\widehat{\Pi}}^{\mathbf{L}} -$, which is an auto-equivalence of $\mathbf{D}^b(\text{fd } \widehat{\Pi})$ by [BIRS].

Now $X \geq E$, $X \geq F$, $Y \geq E$, and $Y \geq F$ because there exist short exact sequences $0 \rightarrow X \rightarrow E \rightarrow Y \rightarrow 0$ and $0 \rightarrow Y \rightarrow F \rightarrow X \rightarrow 0$. \square

The ideas in the proof above also lead to the following lemma. For vertices $i \neq j$ in \overline{Q} , let $W_{i,j} := \langle s_i, s_j \rangle \subset W$ be a parabolic subgroup of W . For $w \in W$, the coset $wW_{i,j}$ is an interval in the weak order on W . We write $H|_{wW_{i,j}}$ for the restriction of $\text{Hasse}(W)$ to $wW_{i,j}$. Define w^+ (respectively, w^-) to be the set of arrows in $\text{Hasse}(W)$ starting (respectively, ending) at w . For a set S of Π -modules, we denote by $\mathsf{T}(S)$ (respectively, $\mathsf{F}(S)$) the smallest torsion (respectively, torsionfree) class in $\text{mod } \Pi$ containing S . For convenience and brevity, we will omit set braces inside the operator $\mathsf{T}(\bullet)$, so that, for example $\mathsf{T}(L_0, M_i \mid i \in I)$ would mean $\mathsf{T}(\{L_0\} \cup \{M_i \mid i \in I\})$. Recall that the layer labelling of $\text{Hasse}(W)$ maps each Hasse arrow $x \rightarrow y$ to the isomorphism class of the corresponding concrete layer $L(x \rightarrow y) := I(y)/I(x)$.

Lemma 4.4. *If $a : ws_i \rightarrow w$ is an arrow in $\text{Hasse}(W)$, and $j \neq i$ is a vertex in \overline{Q} , then*

$$\mathsf{T}(L(b) \mid b \in w^- \cap H|_{wW_{i,j}}) = \mathsf{T}(L(a), L(b) \mid b \in (ws_i)^- \cap H|_{wW_{i,j}}).$$

Proof. We have either the square or the hexagon in Proposition 4.3. We argue the hexagon case. The square case is similar but simpler. If w coincides with u in Proposition 4.3, the desired equality reduces to an identity $\mathsf{T}(X, Y) = \mathsf{T}(X, E)$, which follows from the exact sequence $0 \rightarrow X \rightarrow E \rightarrow Y \rightarrow 0$. If w coincides with us_j in Proposition 4.3, the desired equality reduces to an identity $\mathsf{T}(F) = \mathsf{T}(F, X)$, which follows from the exact sequence $0 \rightarrow Y \rightarrow F \rightarrow X \rightarrow 0$. If w coincides with $us_i s_j$ in Proposition 4.3, the desired equality reduces to an identity $\mathsf{T}(Y) = \mathsf{T}(Y)$, which clearly holds. \square

Another ingredient in the proof of Theorem 4.1 is a precise connection between torsion classes and layer modules.

Theorem 4.5. *For any $w \in W$, we have*

$$\mathsf{T}(w) = \mathsf{T}(L(a) \mid a \in w^-) \quad \text{and} \quad \mathsf{F}(w) = \mathsf{F}(L(a) \mid a \in w^+).$$

Proof. We only prove the first equality since the second one is proved similarly. We use decreasing induction on W . The statement is clear for the longest element w_0 since both sides are $\{0\}$ in this case.

Let $a : ws_i \rightarrow w$ be an arrow in $\text{Hasse}(W)$. Assume that the assertion holds for ws_i , that is,

$$\mathsf{T}(ws_i) = \mathsf{T}(L(b) \mid b \in (ws_i)^-). \quad (4.1)$$

Using obvious decompositions

$$w^- = \bigcup_{j \neq i} (w^- \cap H|_{wW_{i,j}}) \quad \text{and} \quad (ws_i)^- = \bigcup_{j \neq i} ((ws_i)^- \cap H|_{wW_{i,j}}), \quad (4.2)$$

we have

$$\begin{aligned} \mathsf{T}(L(b) \mid b \in w^-) &\stackrel{(4.2)}{=} \mathsf{T}(\mathsf{T}(L(b) \mid b \in w^- \cap H|_{wW_{i,j}}) \mid j \neq i) \\ &\stackrel{\text{Lem. 4.4}}{=} \mathsf{T}(\mathsf{T}(L(a), L(b) \mid b \in (ws_i)^- \cap H|_{wW_{i,j}}) \mid j \neq i) \\ &\stackrel{(4.2)}{=} \mathsf{T}(L(a), \mathsf{T}(L(b) \mid b \in (ws_i)^-)) \\ &\stackrel{(4.1)}{=} \mathsf{T}(L(a), \mathsf{T}(ws_i)) \\ &= \mathsf{T}(w), \end{aligned}$$

where the last equality follows from having an exact sequence $0 \rightarrow I(ws_i) \rightarrow I(w) \rightarrow L(a) \rightarrow 0$ and $I(ws_i) \in \mathsf{T}(w)$. \square

Proposition 4.6. *Two arrows $x \rightarrow y$ and $x' \rightarrow y'$ in $\text{Hasse}(W)$ have $\text{con}(x, y) = \text{con}(x', y')$ if and only if they have the same layer label.*

Proof. Proposition 4.3 implies in particular that the map $(x \rightarrow y) \mapsto I(y)/I(x)$ is constant on components of $\text{SFPoly}(W)$. Thus by Corollary 2.5, if $x \rightarrow y$ and $x' \rightarrow y'$ have $\text{con}(x, y) = \text{con}(x', y')$, then they have the same layer labelling.

If $x \rightarrow y$ and $x' \rightarrow y'$ have the same layer labelling, let m be the unique meet-irreducible element with $\text{con}(m^*, m) = \text{con}(x, y)$, and similarly let m' be the meet-irreducible corresponding to $x' \rightarrow y'$. Then $L(m^* \rightarrow m) = L((m')^* \rightarrow m)$, so by Theorem 4.5, $\mathsf{T}(m) = \mathsf{T}(L(m^* \rightarrow m)) = \mathsf{T}(L((m')^* \rightarrow m)) = \mathsf{T}(m')$. Since $\mathsf{T} : W \rightarrow \text{tors } \Pi$ is a bijection, we see that $m = m'$, so that $\text{con}(x, y) = \text{con}(x', y')$. \square

Proof of Theorem 4.1. Commutativity of the three triangles involving $\text{m-lrr}(W)$, $\text{j-lrr}(W)$, $\text{Con}_{\text{II}}(W)$, and $\text{Hasse}_1(W)$, and correctness of the markings of the arrows as bijections or surjections was established in Section 2.1 in the more general context of congruence uniform lattices. (See especially Figure 4.)

The map from $\text{Hasse}_1(W)$ to $\text{layer } \Pi$ is surjective by definition. Proposition 4.6, together with the commutativity of the triangles containing $\text{Con}_{\text{II}}(W)$, then implies the commutativity of the two triangles containing the map from $\text{Hasse}_1(W)$ to $\text{layer } \Pi$ and also implies that the map from $\text{m-lrr}(W)$ to $\text{layer } \Pi$ is a bijection. The triangle containing $\text{m-lrr}(W)$, $i\tau\text{-rigid } \Pi$ and $\text{layer } \Pi$ commutes by Lemma 4.2, and the left bottom square commutes by Theorem 4.5. The remaining part of the diagram commutes dually. The maps $M : \text{m-lrr } W \rightarrow i\tau\text{-rigid } \Pi$, $\text{Fac} : i\tau\text{-rigid } \Pi \rightarrow \text{tors } \Pi$ and $\text{Filt} : \text{brick } \Pi \rightarrow \text{l-wide } \Pi$ are bijections by Theorems 2.9, 2.7 and Proposition 2.8. The bijectivity of the other maps follows from commutativity. \square

The bijections $\text{j-lrr } W \simeq \text{layer } \Pi \simeq \text{m-lrr } W$ given in Theorem 4.1, combined with Proposition 2.2, imply the following corollary:

Corollary 4.7. *$\mathsf{T}(w)$ is the smallest subcategory of $\Pi\text{-mod}$ which is closed under extensions and quotients and contains the layers corresponding to the canonical meet representation of w . $\mathsf{F}(w)$ is the smallest subcategory of $\Pi\text{-mod}$ which is closed under*

extensions and subobjects and contains the layers corresponding to the canonical join representation of w .

Remark 4.8. Let $a : ws_i \rightarrow w$ be an arrow in H . One might wonder if $F(w)$ is a unique stone of Π which belongs to $\mathbb{T}(w) \setminus \mathbb{T}(ws_i)$.

This has an easy counterexample: let Π be of type A_2 , and let $w := e$ and $s_i := s_1$. Then both layer modules S_1 and P_1 belong to $(\text{mod } \Pi) \setminus (\text{Fac } I_1)$.

It therefore seems that $L(a)$ should be characterized by some kind of minimality among the stones of $\mathbb{T}(w) \setminus \mathbb{T}(ws_i)$.

5. DOUBLETON EXTENSION ORDER ON LAYER MODULES

In this section, we prove Theorem 1.4 and a characterization of the doubleton extension order on layer modules. The last ingredient needed is the following proposition.

Proposition 5.1. *Suppose A, B , and C are layers of Π such that A, C is a doubleton and B is the extension of C by A . Then there exists a hexagon in weak order such that the layer ordering of one of its chains is (A, B, C) , read either from bottom to top or from top to bottom.*

Before proving Proposition 5.1, we show how it completes the proof of Theorem 1.4.

Proof of Theorem 1.4. Theorem 4.1 already states that the map $j \mapsto I(j_*)/I(j)$ is a bijection from the set of join-irreducible elements of W to the set of layer modules of Π . Proposition 4.3 implies that every arrow in the quiver $\text{FPoly}(W)$ (defined in Section 2.1) gives rise to an order relation in the doubleton extension order. Proposition 5.1 shows that each doubleton extension comes from some arrow in $\text{FPoly}(W)$. Corollary 2.4 (which applies in light of Theorem 2.6) thus implies that $j \mapsto I(j_*)/I(j)$ is an isomorphism from the forcing order on join-irreducible elements of W to the doubleton extension order on layers. \square

We now prepare to prove Proposition 5.1.

Lemma 5.2. *If X, Y is a doubleton, then $\text{Hom}_\Pi(X, Y) = \text{Hom}_\Pi(Y, X) = 0$.*

Proof. Let $\langle -, - \rangle$ be the Euler form on $K_0(\text{fd } \widehat{\Pi})$. Since the extensions E of X, Y are layers by assumption, we have $\langle [X] + [Y], [X] + [Y] \rangle = \langle [E], [E] \rangle = 2$. Thus

$$2\langle [X], [Y] \rangle = \langle [X] + [Y], [X] + [Y] \rangle - \langle [X], [X] \rangle - \langle [Y], [Y] \rangle = -2.$$

Since $\text{Ext}_\Pi^1(X, Y) = \text{Ext}_\Pi^1(Y, X)$ is one-dimensional by assumption, it follows that $\text{Hom}_\Pi(X, Y) = \text{Hom}_{\widehat{\Pi}}(X, Y) = 0$ and $\text{Hom}_\Pi(Y, X) = D \text{Ext}_\Pi^2(X, Y) = 0$. \square

The following lemma is an analogue for doubletons of Lemma 3.3.

Lemma 5.3. *If X, Y is a doubleton contained in $\mathbb{T}(w)$, and $\text{Hom}_\Pi(X, S_i) \neq 0$ and $X \not\prec S_i$, then $\ell(s_i w) = \ell(w) + 1$ and $F_i(X), F_i(Y)$ form a doubleton contained in $\mathbb{T}(s_i w)$.*

Proof. Since $\text{Hom}_\Pi(X, S_i) \neq 0$, we have (Proposition 3.2(a)) that $\text{Hom}_\Pi(S_i, X) = 0$. Since $\text{Hom}_\Pi(X, Y) = 0$ by Lemma 5.2, then $\text{Hom}_\Pi(S_i, Y) = 0$ also. Thus, by Proposition 3.2(b), $F_i(X)$ and $F_i(Y)$ are layer modules. Since F_i is an auto-equivalence of $\text{D}^b(\text{fd } \widehat{\Pi})$, it follows that $F_i(X), F_i(Y)$ still form a doubleton.

$F_i(X)$ belongs to $\mathbb{T}(s_i w)$ by Lemma 3.3(b). If $\text{Hom}_\Pi(Y, S_i) \neq 0$, then Lemma 3.3 is also directly applicable to Y , and tells us that $F_i(Y)$ also belongs to $\mathbb{T}(s_i w)$. In fact, the proof of Lemma 3.3(b) is actually applicable to Y even if $\text{Hom}_\Pi(Y, S_i) = 0$ — all that is really needed is that $\text{Hom}_\Pi(S_i, Y) = 0$, and we have this. Thus, $F_i(Y)$ also belongs to $\mathbb{T}(s_i w)$. \square

The following lemma is an analogue for doubletons of Lemma 3.4.

Lemma 5.4. *If X, Y is a doubleton, then there exists $v \in W$ such that $F(v)(X)$ and $F(v)(Y)$ are simple modules.*

Proof. Suppose that at least one of X and Y is not simple. Without loss of generality, suppose that X is not simple. Choose S_i so that $\text{Hom}_\Pi(X, S_i) \neq 0$. Applying Lemma 5.3, we see that $F_i(X)$ and $F_i(Y)$ are a doubleton in $\text{Fac } I_{S_i}$. Assume that one of $F_i(X)$ and $F_i(Y)$ is not simple. Repeat the previous procedure. As in the proof of Lemma 3.4, the procedure must terminate, at which point we have obtained a doubleton of simple modules. \square

Proof of Proposition 5.1. Suppose that we have a doubleton A, C , with B the extension of C by A . We will establish that there exists a hexagon in weak order such that one side of it is labelled (A, B, C) . By Lemma 5.4, there exists $v \in W$ such that $F(v)(A)$, $F(v)(C)$ form a doubleton of simple modules, say S_i, S_j . As in the proof of Theorem 1.2, we conclude that A, C are isomorphic to $D(I(v)/I(s_i v))$ and $D(I(v)/I(s_j v))$. By Lemma 3.5, these are the labels of the Hasse arrows $w_0 v^{-1} \rightarrow w_0 v^{-1} s_i$ and $w_0 v^{-1} \rightarrow w_0 v^{-1} s_j$ which form the two top arrows of the desired hexagon. Since the extension groups between A and C are one-dimensional, the side of the hexagon whose arrows are labelled $(A, ?, C)$, has B as the label of its middle side. \square

It is immediate from the definition that if $A \geq B$ in the doubleton extension order, then A is a subfactor of B . The converse does not hold in general. In Section 6, we define conventions regarding modules over the preprojective algebra of type D_n . In the notation of that section, in type D_4 ,

$$\begin{array}{|c|c|} \hline -2 & -1 \\ \hline -3 & 1 \\ \hline \end{array} \not\geq \begin{array}{|c|c|} \hline 2 & 3 \\ \hline -2 & -1 \\ \hline \end{array},$$

even though the first of these modules is a subfactor of the second.

We now show that the converse does hold in type A_n . We denote by \mathcal{S} the set of *non-revisiting walks* on the double quiver \overline{Q} . By definition, these are walks in Q which follow a sequence of arrows either with or against the direction of the arrow and which do not visit any vertex more than once. We identify a walk and its reverse walk.

Let I_{cyc} denote the ideal of Π generated by all 2-cycles and let $\overline{\Pi} := \Pi/I_{\text{cyc}}$. To any $p \in \mathcal{S}$, we can associate an indecomposable $\overline{\Pi}$ -module X_p called a *string module*, and these exhaust the indecomposable $\overline{\Pi}$ -modules, see [WW].


Theorem 5.5. *Suppose Π is the preprojective algebra of type A_n .*

- (a) *The layer modules for Π are exactly the indecomposable $\overline{\Pi}$ -modules (which are exactly the string modules).*
- (b) *The doubleton extension order on layer modules is the opposite of subfactor order.*

Proof. (a) Note that I_{cyc} is generated as an ideal of Π by the sum of all the two-cycles, which is a central element $x \in \Pi$. If X is a Π -module, then multiplication by x is a surjective morphism onto $I_{\text{cyc}}X \subset X$. Therefore, if X is a stone, $I_{\text{cyc}}X$ must be zero. Conversely, it is easy to see that any string module for $\overline{\Pi}$ is a stone.

(b) As mentioned above, we need only show that if A is a subfactor of B , then $A \geq B$. If A is a submodule or a quotient module of B , then B is part of a doubleton having A as one of its two extensions; if not, then A is a submodule of some layer module C which is a quotient of B , and we apply the same argument twice. \square

Example 5.6. Let $n = 3$. Then the Hasse quiver of \mathcal{S} with respect to the opposite of subfactor order is the following:



In light of Theorems 1.4 and 5.5, this quiver is the Hasse quiver of the forcing order on join-irreducible elements in type A_3 . Compare [R1, Figure 4].

6. COMBINATORIAL DESCRIPTION OF INDECOMPOSABLE τ -RIGID MODULES

Let Π be a preprojective algebra of Dynkin type and W the corresponding Weyl group. A combinatorial description of join-irreducible elements in W is well-known for type A and D . We refer to Sections 1.5, 2.1, 2.4, and 2.6 in [BB] for type A , and Section 8.2 in [BB] for type D . (The description of join-irreducible elements is not given explicitly in [BB], but it is easily worked out from the combinatorial models developed there.) On the other hand, recall from Theorem 4.1 that we have a bijection

$$J : \text{j-lrr } W \rightarrow i\tau^- \text{-rigid } \Pi$$

given by $J(w) := (\Pi/I(w))e_i$ for a unique arrow $w \rightarrow ws_i$ in the Hasse quiver of W starting at w .

The main result of this section is to give a combinatorial description of the Π -module $J(w)$ for each join-irreducible element $w \in W$ in type A or D . See Theorem 6.1 for type A and Theorems 6.5 and 6.12 for type D . It will be interesting to compare our results with Bongartz's description of bricks for type A and D [Bon].

6.1. Type A. Let Π be a preprojective algebra of type A_n . It is given by a quiver

$$1 \begin{array}{c} \xleftarrow{x_1} \\ \xrightarrow{y_2} \end{array} 2 \begin{array}{c} \xleftarrow{x_2} \\ \xrightarrow{y_3} \end{array} 3 \begin{array}{c} \xleftarrow{x_3} \\ \xrightarrow{y_4} \end{array} \cdots \begin{array}{c} \xleftarrow{x_{n-2}} \\ \xrightarrow{y_{n-1}} \end{array} n-1 \begin{array}{c} \xleftarrow{x_{n-1}} \\ \xrightarrow{y_n} \end{array} n$$

with relations $x_1y_2 = 0$, $x_iy_{i+1} = y_ix_{i-1}$ for $2 \leq i \leq n-1$ and $y_nx_{n-1} = 0$. We denote by S_ℓ the simple Π -module corresponding to the vertex ℓ , and by P_ℓ the projective cover of S_ℓ .

Let $W = \mathfrak{S}_{n+1}$ be the Weyl group of Π . We use the convention that the product ww' of elements $w, w' \in W$ is given by $(ww')(i) = w(w'(i))$ for $i \in \{1, \dots, n+1\}$. The elements of W are the permutations

$$w = [i_1, \dots, i_{n+1}].$$

This is join-irreducible if and only if there exists $\ell \in \{1, \dots, n\}$ such that

$$i_1 < \cdots < i_\ell > i_{\ell+1} < \cdots < i_{n+1}.$$

In this case, we say that w is of *type* ℓ . There exists a unique arrow $w \rightarrow ws_\ell$ starting at w in the Hasse quiver of W . The number of join-irreducible elements of type ℓ is given by

$$\binom{n+1}{\ell} - 1,$$

and therefore $\#\text{j-lrr } W = 2^{n+1} - n - 2$.

As a quiver representation, P_ℓ is given by

$$\begin{array}{cccccccc}
 \ell & \longrightarrow & \ell-1 & \longrightarrow & \ell-2 & \longrightarrow & \dots & \longrightarrow & 3 & \longrightarrow & 2 & \longrightarrow & 1 \\
 \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\
 \ell+1 & \longrightarrow & \ell & \longrightarrow & \ell-1 & \longrightarrow & \dots & \longrightarrow & 4 & \longrightarrow & 3 & \longrightarrow & 2 \\
 \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\
 \ell+2 & \longrightarrow & \ell+1 & \longrightarrow & \ell & \longrightarrow & \dots & \longrightarrow & 5 & \longrightarrow & 4 & \longrightarrow & 3 \\
 \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\
 \vdots & & \vdots & & \vdots & & \dots & & \vdots & & \vdots & & \vdots \\
 \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\
 n-1 & \longrightarrow & n-2 & \longrightarrow & n-3 & \longrightarrow & \dots & \longrightarrow & n-\ell+2 & \longrightarrow & n-\ell+1 & \longrightarrow & n-\ell \\
 \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\
 n & \longrightarrow & n-1 & \longrightarrow & n-2 & \longrightarrow & \dots & \longrightarrow & n-\ell+3 & \longrightarrow & n-\ell+2 & \longrightarrow & n-\ell+1
 \end{array}$$

where each number i shows a k -vector space k lying on the vertex i , and each arrow is the identity map of k . Submodules (respectively, factor modules) of P_ℓ correspond bijectively to subquivers that are closed under successors (respectively, predecessors).

We represent P_ℓ in abbreviated form as an array of numbers in rows as follows:

$$P_\ell = \begin{array}{|c|c|c|c|c|c|c|}
 \hline
 \ell & \ell-1 & \ell-2 & \dots & 3 & 2 & 1 \\
 \hline
 \ell+1 & \ell & \ell-1 & \dots & 4 & 3 & 2 \\
 \hline
 \ell+2 & \ell+1 & \ell & \dots & 5 & 4 & 3 \\
 \hline
 \vdots & \vdots & \vdots & \dots & \vdots & \vdots & \vdots \\
 \hline
 n-1 & n-2 & n-3 & \dots & n-\ell+2 & n-\ell+1 & n-\ell \\
 \hline
 n & n-1 & n-2 & \dots & n-\ell+3 & n-\ell+2 & n-\ell+1 \\
 \hline
 \end{array}$$

Submodules and factor modules are similarly represented by sub-arrays of P_ℓ , as, for example, in the following theorem.

Theorem 6.1. *Let $w = [i_1, \dots, i_{n+1}] \in W$ be a join-irreducible element of type ℓ . Then $J(w)$ is a factor module of P_ℓ which has the form*

$$J(w) = \begin{array}{|c|c|c|c|c|c|c|}
 \hline
 \ell & \ell-1 & \ell-2 & \dots & \dots & \dots & i_{\ell+1} \\
 \hline
 \ell+1 & \ell & \ell-1 & \dots & \dots & \dots & i_{\ell+2} \\
 \hline
 \ell+2 & \ell+1 & \ell & \dots & \dots & i_{\ell+3} & \\
 \hline
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \\
 \hline
 n-1 & \dots & \dots & \dots & i_n & & \\
 \hline
 n & \dots & i_{n+1} & & & & \\
 \hline
 \end{array}$$

In particular, any factor module of P_ℓ is indecomposable τ^- -rigid.

Note that $i_m \leq m$ holds for any $m \in \{\ell+1, \dots, n+1\}$, and if $i_m = m$ holds, then the row starting at $m-1$ is empty. For example, for $n=5$ and $w = [351246]$,

$$\text{then } \ell = 2 \text{ and } J(w) = \begin{array}{|c|c|}
 \hline
 2 & 1 \\
 \hline
 3 & 2 \\
 \hline
 4 & \\
 \hline
 \end{array} = \begin{array}{c}
 2 \rightarrow 1 \\
 \downarrow \quad \downarrow \\
 3 \rightarrow 2 \\
 \downarrow \\
 4
 \end{array}$$

To prove this, we need the following easy observation.

Lemma 6.2. *Let $w = [i_1, \dots, i_{n+1}] \in W$ be a join-irreducible element of type ℓ . Then we have a reduced expression $w = x_{n+1}x_n \dots x_{\ell+2}x_{\ell+1}$, where $x_m = s_{i_m}s_{i_m+1} \dots s_{m-2}s_{m-1}$ for $m \in \{\ell+1, \dots, n+1\}$.*

Proof. If $\ell = n$, then $w = [1, 2, \dots, i-1, i+1, \dots, n+1, i]$ holds for some $i \in \{1, \dots, n\}$. Thus $w = x_{n+1}$ holds clearly. In the rest, assume $\ell \neq n$, and let $v := x_{n+1}^{-1}w$. Since x_{n+1}^{-1} is a cyclic permutation $(n+1, n, \dots, i_{n+1}+1, i_{n+1})$, it preserves the total order on $\{i_1, \dots, i_n\}$, and therefore v is a join-irreducible element of type ℓ and clearly satisfies $v(n+1) = n+1$. Since x_{n+1}^{-1} fixes any element in $\{i_{\ell+1}, \dots, i_n\}$, we have $v(m) = i_m$ for any $m \in \{\ell+1, \dots, n\}$ and $v(n+1) = n+1$. Inductively on n , we have $v = x_n \dots x_{\ell+1}$, and therefore $w = x_{n+1}x_n \dots x_{\ell+1}$. \square

Now we are ready to prove Theorem 6.1.

Proof of Theorem 6.1. Clearly we have

$$I(x_{\ell+1})e_\ell = I_{i_{\ell+1}} \cdots I_\ell e_\ell =$$

				$i_{\ell+1}+1$	\cdots	1
$\ell+1$	ℓ	\cdots	\cdots	\cdots	\cdots	2
$\ell+2$	$\ell+1$	\cdots	\cdots	\cdots	\cdots	3
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
n	$n-1$	\cdots	\cdots	\cdots	\cdots	$n-\ell+1$

and

$$I(x_{\ell+2}x_{\ell+1})e_\ell =$$

				$i_{\ell+1}+1$	\cdots	1
			$i_{\ell+2}+1$	\cdots	\cdots	2
$\ell+2$	$\ell+1$	\cdots	\cdots	\cdots	\cdots	3
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
n	$n-1$	\cdots	\cdots	\cdots	\cdots	$n-\ell+1$

Repeating similar calculations, we have

$$I(w)e_\ell =$$

				$i_{\ell+1}+1$	\cdots	1
			$i_{\ell+2}+1$	\cdots	\cdots	2
		$i_{\ell+3}+1$	\cdots	\cdots	\cdots	3
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
$i_{n+1}+1$	\cdots	\cdots	\cdots	\cdots	\cdots	$n-\ell+1$

Therefore $J(w) = (\Pi/I(w))e_\ell$ has the desired form. \square

Example 6.3. Let $w = [n - \ell + 2, n - \ell + 3, \dots, n + 1, 1, 2, \dots, n - \ell + 1]$ be a join-irreducible element of type ℓ . Then we have a reduced expression

$$w = x_{n+1}x_n \cdots x_{\ell+2}x_{\ell+1} \quad \text{for } x_m = s_{m-\ell} \cdots s_{m-2}s_{m-1},$$

and the corresponding indecomposable τ^- -rigid Π -module is $J(w) = P_\ell$.

Example 6.4. Consider type A_3 .

We have the following 3 join-irreducible elements of type 1.

$$J(2134) = J(s_1) = \boxed{1}, \quad J(3124) = J(s_2 \cdot s_1) = \boxed{\frac{1}{2}}, \quad J(4123) = J(s_3 \cdot s_2 \cdot s_1) = \boxed{\frac{1}{2} \frac{1}{3}}.$$

We have the following 5 join-irreducible elements of type 2.

$$J(1324) = J(s_2) = \boxed{2}, \quad J(2314) = J(s_1 s_2) = \boxed{2 \ 1}, \quad J(1423) = J(s_3 \cdot s_2) = \boxed{\frac{2}{3}},$$


$$J(2413) = J(s_3 \cdot s_1 s_2) = \boxed{\frac{2 \ 1}{3}}, \quad J(3412) = J(s_2 s_3 \cdot s_1 s_2) = \boxed{\frac{2 \ 1}{3 \ 2}}.$$

We have the following 3 join-irreducible elements of type 3.

$$J(1243) = J(s_3) = \boxed{3}, \quad J(1342) = J(s_2 s_3) = \boxed{3 \ 2},$$

$$J(2341) = J(s_1 s_2 s_3) = \boxed{3 \ 2 \ 1}.$$

6.2. Type D . Let Π be a preprojective algebra of type D_n . Then Π is given by a quiver



with relations $x_1 y_2 = 0$, $x'_1 y'_2 = 0$, $x_2 y_3 = y_2 x_1 + y'_2 x'_1$, $x_i y_{i+1} = y_i x_{i-1}$ for $3 \leq i \leq n-2$ and $y_{n-1} x_{n-2} = 0$. Let S_ℓ be the simple Π -module corresponding to the vertex ℓ , and let P_ℓ be the projective cover of S_ℓ .

The Weyl group W of Π is the group of automorphisms w of the set $\{\pm 1, \dots, \pm n\}$ satisfying $w(-\ell) = -w(\ell)$ for any ℓ and $\#\{\ell \in \{1, \dots, n\} \mid w(\ell) < 0\}$ is even. Setting $i_\ell = w(\ell)$ for $\ell \in \{1, \dots, n\}$, we write $w \in W$ as a sequence

$$w = [i_1, \dots, i_n] \in \{\pm 1, \dots, \pm n\}^n$$

satisfying the following two conditions.

- $|i_1|, \dots, |i_n|$ is a permutation of $1, \dots, n$.
- The number of negative integers is even.

We often denote $-i$ by \bar{i} . The simple reflections in W are given by

$$s_{-1} = [\underline{2}, \underline{1}, 3, 4, \dots, n] \quad \text{and} \quad s_\ell = [1, \dots, \ell - 1, \ell + 1, \ell, \ell + 2, \dots, n]$$

with $\ell \in \{1, \dots, n - 1\}$. The length of $w = [i_1, \dots, i_n]$ is given by

$$\ell(w) = \#\{1 \leq \ell < m \leq n \mid i_\ell > i_m\} + \#\{1 \leq \ell < m \leq n \mid -i_\ell > i_m\}.$$

For $\ell \in \{1, \dots, n - 1\}$, $\ell(ws_\ell) - \ell(w)$ is 1 if $i_\ell < i_{\ell+1}$, and -1 otherwise. Moreover $\ell(ws_{-1}) - \ell(w)$ is 1 if $-i_1 < i_2$, and -1 otherwise. Therefore an element $w = [i_1, \dots, i_n] \in W$ is join-irreducible if and only if one of the following conditions is satisfied.


- $i_1 < \dots < i_n$ and $-i_1 > i_2$.
- There exists $\ell \in \{1, \dots, n - 1\}$ such that $i_1 < \dots < i_\ell > i_{\ell+1} < \dots < i_n$ and $-i_1 < i_2$.

We say that w is of *type* -1 (respectively, *type* ℓ) if the first (respectively, second) condition is satisfied. Then there exists a unique arrow $w \rightarrow ws_\ell$ starting at w in the Hasse quiver of W . The number of join-irreducible elements of type $\ell \neq \pm 1$ (respectively, $1, -1$) is

$$2^{n-\ell} \binom{n}{\ell} - 1 \quad (\text{respectively, } 2^{n-1} - 1, 2^{n-1} - 1),$$

and therefore $\#j\text{-lrr } W = 3^n - n2^{n-1} - n - 1$.

As a quiver representation, P_ℓ for $\ell = \pm 1$ is given by



where each number i shows a k -vector space k lying on the vertex i , and each arrow is the identity map of k . Again, submodules (respectively, factor modules) of P_ℓ correspond bijectively to subquivers that are closed under successors (respectively, predecessors).

We represent P_ℓ in abbreviated form as an array of numbers in rows as follows:

$$P_\ell = \begin{array}{|c|} \hline \pm 1 \\ \hline 2 \quad \mp 1 \\ \hline 3 \quad 2 \quad \pm 1 \\ \hline \vdots \quad \vdots \quad \vdots \quad \ddots \\ \hline n-3 \quad n-4 \quad n-5 \quad \dots \quad \pm(-1)^n \\ \hline n-2 \quad n-3 \quad n-4 \quad \dots \quad 2 \quad \mp(-1)^n \\ \hline n-1 \quad n-2 \quad n-3 \quad \dots \quad 3 \quad 2 \quad \pm(-1)^n \\ \hline \end{array} .$$

Submodules and factor modules of P_ℓ are again represented by subarrays.

Theorem 6.5. *Let $w = [i_1, \dots, i_n] \in W$ be a join-irreducible element of type $\ell = \pm 1$. Then $J(w)$ is a factor module of P_ℓ which has the form*

$$J(w) = \begin{array}{|c|} \hline i'_2 \\ \hline 2 \quad i'_3 \\ \hline 3 \quad \dots \quad i'_4 \\ \hline \vdots \quad \vdots \quad \vdots \quad \ddots \\ \hline n-2 \quad n-3 \quad \dots \quad i'_{n-1} \\ \hline n-1 \quad n-2 \quad \dots \quad i'_n \\ \hline \end{array} .$$

where $i'_m := \max\{i_m, (-1)^m \ell\}$ for $m \in \{2, \dots, n\}$.

Note that $i'_m \leq m$ holds for any $m \in \{2, \dots, n\}$, and if $i'_m = m$ holds, then the row starting at $m - 1$ is empty. For example, for $n = 6$ and $w = [5\bar{3}\underline{1}246]$, then

$$\ell = 1 \text{ and } J(w) = \begin{array}{|c|c|} \hline 1 & \\ \hline 2 & -1 \\ \hline 3 & 2 \\ \hline 4 & \\ \hline \end{array} = \begin{array}{c} \downarrow \\ 2 \rightarrow -1 \\ \downarrow \downarrow \\ 3 \rightarrow 2 \\ \downarrow \\ 4 \end{array}.$$

To prove this, we need the following observation.

Lemma 6.6. *Let $w = [i_1, \dots, i_n] \in W$ be a join-irreducible element of type $\ell = \pm 1$. Then we have a reduced expression $w = x_n x_{n-1} \cdots x_3 x_2$, where*

$$x_m = \begin{cases} s_{i_m} s_{i_m+1} \cdots s_{m-2} s_{m-1} & \text{if } i_m \geq 2 \\ s_{(-1)^m \ell} s_2 \cdots s_{m-2} s_{m-1} & \text{if } i_m < 2 \end{cases} \quad \text{for } m \in \{2, \dots, n\}.$$

Proof. The case $n = 2$ can be checked directly. In the rest, we assume $n \geq 3$. Assume that the assertion holds for $n - 1$. Again, let $v := x_n^{-1} w$.

Assume $i_n > 0$. Then v is obtained from w by replacing i_n with n and then, for each i_j with $i_n < |i_j|$ replacing i_j by an entry with the same sign but absolute value $|i_j| - 1$. In particular, v is a join-irreducible element of type ℓ and satisfies $v(n) = n$. Fix $m \in \{2, \dots, n - 1\}$. If $i_m > 0$, then $v(m) = i_m$ holds, and if $i_m < 0$, then $v(m) < 0$ holds. By our assumption on induction, we have $v = x_{n-1} \cdots x_2$, and therefore $w = x_n x_{n-1} \cdots x_2$.

Assume $i_n < 0$. In this case, we have $w = [(-1)^{n-1} n, \underline{n-1}, \underline{n-2}, \dots, \underline{1}]$ and $\ell = (-1)^{n-1}$. Thus $x_n^{-1} = [\underline{n}, \underline{1}, 2, \dots, n-1]$ and $v = [(-1)^{n-1} (n-1), \underline{n-2}, \dots, \underline{2}, 1, n]$ hold. Therefore v is a join-irreducible element of type ℓ and satisfies $v(n) = n$. Since $v(m) < 2$ holds for any $m \in \{2, \dots, n - 1\}$, we have $v = x_{n-1} \cdots x_2$ by our assumption on induction, and therefore $w = x_n x_{n-1} \cdots x_2$. \square

Now we are ready to prove Theorem 6.5.

Proof of Theorem 6.5. Using Lemma 6.6, one can calculate $I(w)$ as in the proof of Theorem 6.1, and we obtain the desired assertion. \square

Example 6.7. Let $w = [n, \underline{n-1}, \underline{n-2}, \dots, \underline{2}, (-1)^n]$ be a join-irreducible element of type 1. Then we have a reduced expression

$$w = x_n x_{n-1} \cdots x_3 x_2 \quad \text{for } x_m = s_{(-1)^m} s_2 s_3 \cdots s_{m-1},$$

and the corresponding indecomposable τ^- -rigid Π -module is $J(w) = P_1$.

Similarly, let $w = [\underline{n}, \underline{n-1}, \underline{n-2}, \dots, \underline{2}, (-1)^{n+1}]$ be a join-irreducible element of type -1 . Then we have a reduced expression

$$w = x_n x_{n-1} \cdots x_3 x_2 \quad \text{for } x_m = s_{(-1)^{m+1}} s_2 s_3 \cdots s_{m-1},$$

and the corresponding indecomposable τ^- -rigid Π -module is $J(w) = P_{-1}$.

Example 6.8. Consider type D_4 .

We have the following 7 join-irreducible elements of type 1.

$$J(2134) = J(s_1) = \begin{array}{|c|} \hline 1 \\ \hline \end{array}, \quad J(3124) = J(s_2 \cdot s_1) = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \end{array},$$

$$J(3\underline{2}14) = J(s_{-1} s_2 \cdot s_1) = \begin{array}{|c|c|} \hline 1 & \\ \hline 2 & -1 \\ \hline \end{array}, \quad J(4123) = J(s_3 \cdot s_2 \cdot s_1) = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array},$$

$$J(4\underline{2}13) = J(s_3 \cdot s_{-1} s_2 \cdot s_1) = \begin{array}{|c|c|} \hline 1 & \\ \hline 2 & -1 \\ \hline 3 & \\ \hline \end{array}, \quad J(4\underline{3}12) = J(s_2 s_3 \cdot s_{-1} s_2 \cdot s_1) = \begin{array}{|c|c|} \hline 1 & \\ \hline 2 & -1 \\ \hline 3 & 2 \\ \hline \end{array},$$

$$J(4\underline{3}21) = J(s_1 s_2 s_3 \cdot s_{-1} s_2 \cdot s_1) = \begin{array}{|c|c|c|} \hline 1 & & \\ \hline 2 & -1 & \\ \hline 3 & 2 & 1 \\ \hline \end{array}.$$

We have the following 7 join-irreducible elements of type -1 .

$$\begin{aligned}
J(\underline{2134}) &= J(s_{-1}) = \boxed{\begin{array}{c} -1 \\ \hline \end{array}}, & J(\underline{3124}) &= J(s_2 \cdot s_{-1}) = \boxed{\begin{array}{c} -1 \\ \hline 2 \end{array}}, \\
J(\underline{3214}) &= J(s_1 s_2 \cdot s_{-1}) = \boxed{\begin{array}{cc} -1 & \\ \hline 2 & 1 \end{array}}, & J(\underline{4123}) &= J(s_3 \cdot s_2 \cdot s_{-1}) = \boxed{\begin{array}{c} -1 \\ \hline 2 \\ \hline 3 \end{array}}, \\
J(\underline{4213}) &= J(s_3 \cdot s_1 s_2 \cdot s_{-1}) = \boxed{\begin{array}{cc} -1 & \\ \hline 2 & 1 \\ \hline 3 & \end{array}}, & J(\underline{4312}) &= J(s_2 s_3 \cdot s_1 s_2 \cdot s_{-1}) = \boxed{\begin{array}{cc} -1 & \\ \hline 2 & 1 \\ \hline 3 & 2 \end{array}}, \\
J(\underline{4321}) &= J(s_{-1} s_2 s_3 \cdot s_1 s_2 \cdot s_{-1}) = \boxed{\begin{array}{ccc} -1 & & \\ \hline 2 & 1 & \\ \hline 3 & 2 & -1 \end{array}}.
\end{aligned}$$

In the rest of this section, let w be a join-irreducible element $i_1 < \cdots < i_\ell > i_{\ell+1} < \cdots < i_n$ of type $\ell \neq \pm 1$. Note that all integers in i_2, \dots, i_ℓ must be positive, and therefore w can be recovered from the latter part $i_{\ell+1}, \dots, i_n$.

We need the following preparation on the structure of P_ℓ .

Lemma 6.9. *Let α and β be scalars satisfying $\alpha + \beta = 1$. As a quiver representation, P_ℓ with $\ell \neq \pm 1$ is given by*

(6.1)

where each number i shows a k -vector space k lying on the vertex i if $i \geq -1$ and $-i$ if $i \leq -2$. Each unlabelled arrow is the identity map of k , and each arrow labelled by a scalar γ is a linear map multiplying by γ .

Proof. Since all relations of Π are satisfied, this gives a Π -module X . It is easy to check that X is generated by ℓ in the upper left corner. Thus we have a surjective morphism $\pi : P_\ell \rightarrow X$ of Π -modules. On the other hand, it is well-known that the Loewy length of any indecomposable projective Π -module is equal to $h - 1$, where h is the Coxeter number. For type D_n , we have $h - 1 = 2n - 3$. Since the length of the path from ℓ in the left corner to $-\ell$ in the lower right corner is $2n - 4$, the above π must be an isomorphism. \square

Example 6.10. Let $n = 6$. Then P_2 is given by the following quiver representations, where the left one is the case $(\alpha, \beta) = (0, 1)$, and the right one is the case $(\alpha, \beta) = (1, 0)$.

We write the quiver P_ℓ of (6.1) in abbreviated form as the following array of numbers.

$$\begin{array}{cccccccccccc}
 \ell & \ell-1 & \cdots & 2 & \overset{1}{-1} & -2 & \cdots & 1-\ell & -\ell & -\ell-1 & \cdots & 2-n & 1-n \\
 \ell+1 & \ell & \cdots & 3 & 2 & \overset{1}{-1} & \cdots & 2-\ell & 1-\ell & -\ell & \cdots & 3-n & 2-n \\
 \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\
 n-2 & n-3 & \cdots & \ell & \ell-1 & \ell-2 & \cdots & \overset{1}{-1} & -2 & -3 & \cdots & -\ell & -\ell-1 \\
 n-1 & n-2 & \cdots & \ell+1 & \ell & \ell-1 & \cdots & 2 & \overset{1}{-1} & -2 & \cdots & 1-\ell & -\ell
 \end{array} \quad (6.2)$$

The description of factor $P_{\pm 1}$ was no more complicated than the analogous description in type A . However, for $\ell > 1$, the description of factor modules of P_ℓ is much more complicated in type D . For example, consider the direct sum k^2 of k 's corresponding to -2 in the first row and 2 in the second row. Then subspaces of k^2 generate distinct submodules of P_ℓ . Fortunately, to describe $J(w)$ for w join-irreducible, we only need the following special class of factor modules.

Definition 6.11. Let S be a subarray of the array (6.2). We say that S is *predecessor-closed* if it is closed under predecessors in the quiver (6.1).

Now we fix scalars α and β satisfying $\alpha + \beta = 1$. We say that S is (α, β) -*predecessor-closed* if it is closed under predecessors in the subquiver of (6.1) obtained by removing all arrows indexed by the scalar 0 . An (α, β) -predecessor-closed subarray S gives a factor module of P_ℓ in a natural way. Clearly, if $(\alpha, \beta) \neq (1, 0), (0, 1)$, then S is (α, β) -predecessor-closed if and only if it is predecessor-closed.

For $m \in \{2, \dots, n-1\}$, let $C(m, j)$ be the following subset of numbers in the row of (6.2) starting at m .

$$C(m, j) := \begin{cases} \emptyset & j > \ell, \\ \boxed{m \ m-1 \ \cdots \ j+2 \ j+1 \ j} & m \geq j \geq 1, \\ \boxed{m \ m-1 \ \cdots \ 3 \ 2 \ -1} & j = -1, \\ \boxed{m \ m-1 \ \cdots \ 3 \ 2 \ \overset{1}{-1}} & j = -2, \\ \boxed{m \ m-1 \ \cdots \ 3 \ 2 \ \overset{1}{-1} \ -2 \ -3 \ \cdots \ j+2 \ j+1} & j \leq -3. \end{cases}$$

We simply write $C(m, j)$ as $\boxed{m \ m-1 \ m-2 \ \cdots \ j'}$, where $j' := j$ if $j \geq -1$ and $j' := j + 1$ if $j \leq -2$.

Let $S(w)$ be the subarray of the array (6.2) given by

$$S(w) = \begin{array}{c} \boxed{C(\ell, i_{\ell+1})} \\ \boxed{C(\ell+1, i_{\ell+2})} \\ \boxed{C(\ell+2, i_{\ell+3})} \\ \vdots \\ \boxed{C(n-2, i_{n-1})} \\ \boxed{C(n-1, i_n)} \end{array} = \begin{array}{cccccccc} \ell & \ell-1 & \ell-2 & \cdots & \cdots & \cdots & \cdots & i'_{\ell+1} \\ \ell+1 & \ell & \ell-1 & \cdots & \cdots & \cdots & \cdots & i'_{\ell+2} \\ \ell+2 & \ell+1 & \ell & \cdots & \cdots & \cdots & \cdots & i'_{\ell+3} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ n-2 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & i'_{n-1} \\ n-1 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & i'_n \end{array} \quad (6.3)$$

Note that $i'_m \leq m$ holds for any $m \in \{\ell+1, \dots, n\}$, and if $i'_m = m$ holds, then the row starting at $m-1$ is empty.

Theorem 6.12. Let $w = [i_1, \dots, i_n] \in W$ be a join-irreducible element of type $\ell \neq \pm 1$, and $S(w)$ subarray of the array (6.2) described in (6.3).

- (a) If $\{1, 2\} \not\subset \{|i_{\ell+1}|, \dots, |i_n|\}$, then $S(w)$ is predecessor-closed.
(b) If $\{1, 2\} \subset \{|i_{\ell+1}|, \dots, |i_n|\}$, then $S(w)$ is $(1, 0)$ or $(0, 1)$ -predecessor-closed.

In either case, we have a factor module of P_ℓ corresponding to $S(w)$.

- (c) $J(w)$ is the factor module of P_ℓ corresponding to $S(w)$.

For example, let $n = 6$ and $w = [\underline{3}4\underline{5}126]$ and $w' = [34\underline{5}216]$. Then

$$S(w) = \begin{array}{cccc} \boxed{2 \ \overset{1}{-1} \ -2 \ -3 \ -4} \\ \boxed{3 \ 2 \ 1} \\ \boxed{4 \ 3 \ 2} \\ \boxed{} \end{array} \quad \text{and} \quad S(w') = \begin{array}{cccc} \boxed{2 \ \overset{1}{-1} \ -2 \ -3 \ -4} \\ \boxed{3 \ 2 \ \overset{1}{-1}} \\ \boxed{4 \ 3 \ 2 \ 1} \\ \boxed{} \end{array}$$

Thus $S(w)$ is $(0, 1)$ -predecessor-closed and $S(w')$ is $(1, 0)$ -predecessor-closed, and

$$J(w) = \begin{array}{ccccccc} & & 2 & \xrightarrow{1} & -2 & \rightarrow & -3 & \rightarrow & -4 \\ & & \downarrow & \xrightarrow{-1} & \downarrow & & & & \\ & & 3 & \xrightarrow{2} & 2 & \xrightarrow{1} & & & \\ & & \downarrow & \downarrow & \downarrow & & & & \\ & & 4 & \xrightarrow{3} & 3 & \xrightarrow{2} & & & \end{array} \quad \text{and} \quad J(w') = \begin{array}{ccccccc} & & 2 & \xrightarrow{-1} & -2 & \rightarrow & -3 & \rightarrow & -4 \\ & & \downarrow & \xrightarrow{-1} & \downarrow & & & & \\ & & 3 & \xrightarrow{2} & 2 & \xrightarrow{1} & & & \\ & & \downarrow & \downarrow & \downarrow & & & & \\ & & 4 & \xrightarrow{3} & 3 & \xrightarrow{2} & & & \end{array}.$$

To prove Theorem 6.12, we need some preparation. Let $w = [i_1, \dots, i_n] \in W$ be a join-irreducible element of type ℓ . For each $m \in \{\ell + 1, \dots, n\}$, let

$$j_m := i_m + \#\{m' \mid m \leq m' \leq n, |i_{m'}| \leq |i_m|\},$$

Clearly $i_m > 0$ implies $j_m = i_m + 1 > 1$, and $i_m < 0$ implies $j_m \leq 0$. In the latter case, we have $j_m = -\#\{m' \mid 1 \leq m' \leq \ell, |i_{m'}| \leq |i_m|\}$. Therefore $j_{\ell+1} \leq j_{\ell+2} \leq \dots \leq j_n$ holds.

Lemma 6.13. *Let $w = [i_1, \dots, i_n] \in W$ be a join-irreducible element of type $\ell \neq \pm 1$. Then we have a reduced expression $w = x_n x_{n-1} \cdots x_{\ell+2} x_{\ell+1}$, where x_m for $m \in \{\ell + 1, \dots, n\}$ is given by*

$$x_m = \begin{cases} s_{i_m} s_{i_m+1} \cdots s_{m-1} & \text{if } j_m > 0, \\ s_{\epsilon_m} s_2 s_3 \cdots s_{m-1} \text{ for } \epsilon_m := (-1)^{\#\{m' \mid m \leq m' \leq n, i_{m'} < 0\}} & \text{if } j_m = 0, \\ s_{-1} s_1 s_2 s_3 \cdots s_{m-1} & \text{if } j_m = -1, \\ s_{-j_m} s_{-j_m-1} \cdots s_3 s_2 s_{-1} s_1 s_2 s_3 \cdots s_{m-1} & \text{if } j_m < -1. \end{cases} \quad (6.4)$$

Proof. Since $i_1 < i_2$ and $-i_1 < i_2$, we have $0 < i_2 < \dots < i_\ell$. Let $h_1 < \dots < h_{\ell+1}$ be the reordering of

- $i_1, i_2, \dots, i_\ell, i_{\ell+1}$ if $j_{\ell+1} > 0$ or ($j_{\ell+1} = 0$ and $i_1 > 0$),
- $-i_1, i_2, \dots, i_\ell, -i_{\ell+1}$ if $j_{\ell+1} < 0$ or ($j_{\ell+1} = 0$ and $i_1 < 0$),

and let $v := [h_1, \dots, h_{\ell+1}, i_{\ell+2}, \dots, i_n] \in W$. It is easy to check that v is either an identity or join-irreducible of type $\ell + 1$. We will show

$$w x_{\ell+1}^{-1} = v \quad \text{and} \quad \ell(w) - \ell(x_{\ell+1}) = \ell(v). \quad (6.5)$$

Then the assertion follows inductively.

Let t be the unique integer satisfying $h_t = |i_{\ell+1}|$.

Assume $j_{\ell+1} > 0$ and so $i_{\ell+1} > 0$. Since all positive integers smaller than $i_{\ell+1}$ appear in $|i_1|, \dots, |i_\ell|$ we have $t = i_{\ell+1}$. Thus (6.5) follows from $x_{\ell+1} = s_t s_{t+1} \cdots s_\ell$.

Assume $j_{\ell+1} = 0$ and so $i_{\ell+1} < 0$. Since all positive integers smaller than $i_{\ell+1}$ appear in $|i_{\ell+2}|, \dots, |i_n|$, we have $t = 1$. If $i_1 > 0$, then $\epsilon_{\ell+1} = 1$ holds, and (6.5) follows from $x_{\ell+1} = s_1 s_2 s_3 \cdots s_\ell$. If $i_1 < 0$, then $\epsilon_{\ell+1} = -1$ holds, and (6.5) follows from $x_{\ell+1} = s_{-1} s_2 s_3 \cdots s_\ell$.

Assume $j_{\ell+1} < 0$ and so $i_{\ell+1} < 0$. Then $t = 1 - j_{\ell+1}$ holds, and (6.5) follows easily from $x_{\ell+1} = s_{t-1} s_{t-2} \cdots s_3 s_2 s_{-1} s_1 s_2 s_3 \cdots s_\ell$. \square

Now we are ready to prove Theorem 6.12.

Proof of Theorem 6.12. (a) and (b) are easily checked.

(c) Choose scalars α and β such that $S(w)$ is (α, β) -predecessor-closed. Using the quiver representation of P_ℓ given in (6.1) and the reduced expression of w given in Lemma 6.13, we calculate $I(w)e_\ell$. Dividing into three cases, we show that the first row of $J(w)$ coincides with that of $S(w)$.

Assume $j_{\ell+1} > 0$. By (6.4), the first row of $I(x_{\ell+1})e_\ell$ is given by

$$\left\{ \begin{array}{l} \boxed{i_{\ell+1}-1 \quad i_{\ell+1}-2 \quad \cdots \quad 2 \quad \frac{1}{-1} \quad -2 \quad \cdots \quad 2-n \quad 1-n} \quad \text{if } i_{\ell+1} \geq 2, \\ \boxed{-1 \quad -2 \quad -3 \quad \cdots \quad 2-n \quad 1-n} \quad \text{if } i_{\ell+1} = 1. \end{array} \right.$$

Since $0 < i_{\ell+1} < i_{\ell+2} < \cdots < i_n$ holds, both $s_{i_{\ell+1}-1}$ (when $i_{\ell+1} \geq 2$) and s_{-1} (when $i_{\ell+1} = 1$) do not appear in $x_n x_{n-1} \cdots x_{\ell+2}$ by (6.4). Thus the first row of $I(w)e_\ell$ coincides with that of $I(x_{\ell+1})e_\ell$, and the first row of $J(w)$ coincides with that of $S(w)$.

Assume $j_{\ell+1} < 0$. By (6.4), the first row of $I(x_{\ell+1})e_\ell$ is given by

$$\boxed{j_{\ell+1}-1 \quad j_{\ell+1}-2 \quad j_{\ell+1}-3 \quad \cdots \quad 2-n \quad 1-n}.$$

Let $m_0 := \min\{m \mid \ell+1 \leq m \leq n, |i_m| \leq |i_{\ell+1}|\}$, which equals $j_{\ell+1} - i_{\ell+1} + \ell$. We show that, for any $\ell+1 \leq m \leq m_0$, the first row of $I(x_m x_{m-1} \cdots x_{\ell+1})e_\ell$ is

$$\boxed{j_{\ell+1}-m+\ell \quad j_{\ell+1}-m+\ell-1 \quad \cdots \quad 2-n \quad 1-n}.$$

The case $m = \ell+1$ was shown above. Assume that this is the case for $m-1 (< m_0)$. It suffices to show that $s_{-j_{\ell+1}+m-1-\ell}$ appears in x_m , and that $s_{-j_{\ell+1}+m-\ell}$ does not appear in the left side of $s_{-j_{\ell+1}+m-1-\ell}$ in x_m . Since $-j_{\ell+1} \leq \ell$ holds, we have

$$2 \leq -j_{\ell+1} + m - 1 - \ell \leq m - 1.$$

If $j_m \geq 0$, then the assertion follows from (6.4) and

$$i_m \leq i_{m_0} - m_0 + m \leq -i_{\ell+1} - m_0 + m - 1 = -j_{\ell+1} + m - 1 - \ell.$$

If $j_m < 0$, then the assertion follows from (6.4) and $-j_m \leq -j_{\ell+1} < -j_{\ell+1} + m - 1 - \ell$. Inductively, we have shown that the first row of $I(x_{m_0} x_{m_0-1} \cdots x_{\ell+1})e_\ell$ is

$$\boxed{i_{\ell+1} \quad i_{\ell+1}-1 \quad \cdots \quad 2-n \quad 1-n} \quad (6.6)$$

since $j_{\ell+1} - m_0 + \ell = i_{\ell+1}$. Since $0 < -i_{\ell+1} < i_{m_0+1} < \cdots < i_n$ holds, $s_{-i_{\ell+1}}$ does not appear in $x_n x_{n-1} \cdots x_{m_0+1}$. Thus the first row of $I(w)e_\ell$ coincides with (6.6), and the first row of $J(w)$ coincides with that of $S(w)$.

Assume $j_{\ell+1} = 0$. By (6.4), the first row of $I(x_{\ell+1})e_\ell$ is given by

$$\boxed{-\epsilon_{\ell+1} \quad -2 \quad -3 \quad \cdots \quad 2-n \quad 1-n}.$$

By a similar argument as in the case $j_{\ell+1} < 0$, one can check that the first row of $J(w)$ coincides with that of $S(w)$.

We have completed to show that the first row of $J(w)$ coincides with that of $S(w)$. All rows of $I(x_{\ell+1})e_\ell$ except the first one coincide with those of P_ℓ since $s_{\ell+1}$ does not appear in $x_{\ell+1}$. Repeating the same calculation, all rows of $J(w)$ coincide with that of $S(w)$. \square

Example 6.14. Let $w = [(-1)^{n-\ell}, 2, \dots, \ell, \underline{n}, n-1, \dots, \underline{\ell+1}]$ be a join-irreducible element of type ℓ . Then we have a reduced expression

$$w = x_n x_{n-1} \cdots x_{\ell+2} x_{\ell+1} \quad \text{for} \quad x_m = s_\ell \cdots s_3 s_2 s_{-1} s_1 s_2 s_3 \cdots s_{m-1},$$

and the corresponding indecomposable τ^- -rigid Π -module is $J(w) = P_\ell$.

Example 6.15. Consider type D_4 .

We have the following 7 join-irreducible elements of type 3.

$$\begin{aligned} J(1243) &= J(s_3) = \boxed{3}, & J(1342) &= J(s_2 s_3) = \boxed{3 \quad 2}, \\ J(2341) &= J(s_1 s_2 s_3) = \boxed{3 \quad 2 \quad 1}, & J(234\bar{1}) &= J(s_{-1} s_2 s_3) = \boxed{3 \quad 2 \quad -1}, \\ J(\underline{1}34\bar{2}) &= J(s_{-1} s_1 s_2 s_3) = \boxed{3 \quad 2 \quad \bar{1}}, & J(\underline{1}24\bar{3}) &= J(s_2 s_{-1} s_1 s_2 s_3) = \boxed{3 \quad 2 \quad \bar{1} \quad -2}, \\ J(\underline{1}23\bar{4}) &= J(s_3 s_2 s_{-1} s_1 s_2 s_3) = \boxed{3 \quad 2 \quad \bar{1} \quad -2 \quad -3}. \end{aligned}$$

We have the following 23 join-irreducible elements of type 2. They are predecessor-closed unless otherwise specified.

$$\begin{aligned} J(1324) &= J(s_2) = \boxed{2}, & J(\underline{2}3\bar{1}4) &= J(s_{-1} s_2) = \boxed{2 \quad -1}, \\ J(2314) &= J(s_1 s_2) = \boxed{2 \quad 1}, & J(1423) &= J(s_3 \cdot s_2) = \boxed{\frac{2}{3}}, \\ J(2413) &= J(s_3 \cdot s_1 s_2) = \boxed{\frac{2 \quad 1}{3}}, & J(\underline{2}4\bar{1}3) &= J(s_3 \cdot s_{-1} s_2) = \boxed{\frac{2 \quad -1}{3}}, \\ J(\underline{1}3\bar{2}4) &= J(s_{-1} s_1 s_2) = \boxed{2 \quad \bar{1}}, & J(\underline{1}4\bar{2}3) &= J(s_3 \cdot s_{-1} s_1 s_2) = \boxed{\frac{2 \quad \bar{1}}{3}}. \end{aligned}$$

$$\begin{aligned}
J(\underline{1234}) &= J(s_2 s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 \\ \hline & -1 & \\ \hline \end{array}, \\
J(\underline{3412}) &= J(s_2 s_3 \cdot s_{-1} s_2) = \begin{array}{|c|} \hline 2 & -1 \\ \hline 3 & 2 \\ \hline \end{array} \text{ (this is } (0, 1)\text{-predecessor closed),} \\
J(\underline{3412}) &= J(s_2 s_3 \cdot s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 \\ \hline 3 & 2 \\ \hline \end{array} \text{ (this is } (1, 0)\text{-predecessor closed),} \\
J(\underline{3421}) &= J(s_{-1} s_2 s_3 \cdot s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 \\ \hline 3 & 2 & -1 \\ \hline \end{array} \text{ (this is } (1, 0)\text{-predecessor closed),} \\
J(\underline{3421}) &= J(s_1 s_2 s_3 \cdot s_{-1} s_2) = \begin{array}{|c|} \hline 2 & -1 \\ \hline 3 & 2 & 1 \\ \hline \end{array} \text{ (this is } (0, 1)\text{-predecessor closed),} \\
J(\underline{1243}) &= J(s_3 \cdot s_2 s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 & -3 \\ \hline & -1 & & \\ \hline 3 & & & \\ \hline \end{array}, \\
J(\underline{1432}) &= J(s_2 s_3 \cdot s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 \\ \hline & -1 & \\ \hline 3 & 2 & \\ \hline \end{array}, \\
J(\underline{2431}) &= J(s_{-1} s_2 s_3 \cdot s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 \\ \hline 3 & 2 & -1 \\ \hline \end{array}, \\
J(\underline{1342}) &= J(s_2 s_3 \cdot s_2 s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 & -3 \\ \hline & -1 & & \\ \hline 3 & 2 & & \\ \hline \end{array}, \\
J(\underline{2431}) &= J(s_1 s_2 s_3 \cdot s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 \\ \hline 3 & 2 & 1 \\ \hline \end{array}, \\
J(\underline{2341}) &= J(s_{-1} s_2 s_3 \cdot s_2 s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 & -3 \\ \hline & -1 & & \\ \hline 3 & 2 & -1 & \\ \hline \end{array}, \\
J(\underline{1432}) &= J(s_{-1} s_1 s_2 s_3 \cdot s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 \\ \hline & -1 & \\ \hline 3 & 2 & -1 \\ \hline \end{array}, \\
J(\underline{2341}) &= J(s_1 s_2 s_3 \cdot s_2 s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 & -3 \\ \hline & -1 & & \\ \hline 3 & 2 & 1 & \\ \hline \end{array}, \\
J(\underline{1342}) &= J(s_{-1} s_1 s_2 s_3 \cdot s_2 s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 & -3 \\ \hline & -1 & & \\ \hline 3 & 2 & 1 & -1 \\ \hline \end{array}, \\
J(\underline{1243}) &= J(s_2 s_{-1} s_1 s_2 s_3 \cdot s_2 s_{-1} s_1 s_2) = \begin{array}{|c|} \hline 2 & 1 & -2 & -3 \\ \hline & -1 & & \\ \hline 3 & 2 & 1 & -2 \\ \hline \end{array}.
\end{aligned}$$

REFERENCES

- [AIR] T. Adachi, O. Iyama, I. Reiten, τ -tilting theory, *Compos. Math.* 150 (2014), no. 3, 415–452.
- [AM] T. Aihara, Y. Mizuno, *Classifying tilting complexes over preprojective algebras of Dynkin type*, arXiv:1509.07387.
- [A] C. Amiot, *Cluster categories for algebras of global dimension 2 and quivers with potential*, *Ann. Inst. Fourier (Grenoble)* 59 (2009), no. 6, 2525–2590.
- [AIRT] C. Amiot, O. Iyama, I. Reiten, G. Todorov, *Preprojective algebras and c-sortable words*, *Proc. Lond. Math. Soc.* (3) 104 (2012), no. 3, 513–539.
- [AS] M. Auslander, S. O. Smalø, *Almost split sequences in subcategories*, *J. Algebra* 69 (1981), no. 2, 426–454.
- [BKT] P. Baumann, J. Kamnitzer, P. Tingley, *Affine Mirković-Vilonen polytopes*, *Publ. Math. Inst. Hautes Études Sci.* 120 (2014), 113–205.
- [BGL] D. Baer, W. Geigle, H. Lenzing, *The preprojective algebra of a tame hereditary Artin algebra*, *Comm. Algebra* 15 (1987), no. 1-2, 425–457.
- [Bol] B. Bolten, *Spiegelungsfunktoren für präprojektive Algebren*, Diplomarbeit (2010), Bonn.
- [Bon] J. Bongartz, *Endotrivial modules over preprojective algebras*, Master’s Thesis (2015), Bonn.
- [BIRS] A. Buan, O. Iyama, I. Reiten, J. Scott, *Cluster structures for 2-Calabi-Yau categories and unipotent groups*, *Compos. Math.* 145 (2009), no. 4, 1035–1079.
- [BB] A. Björner and F. Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, 231. Springer, New York, 2005.
- [CLM] N. Caspard, C. Le Conte de Poly-Barbut and M. Morvan, *Cayley lattices of finite Coxeter groups are bounded*, *Adv. in Appl. Math.* 33 (2004), no. 1, 71–94.
- [CH] W. Crawley-Boevey, M. P. Holland, *Noncommutative deformations of Kleinian singularities*, *Duke Math. J.* 92 (1998), no. 3, 605–635.
- [D] A. Day, *Congruence normality: the characterization of the doubling class of convex sets*, *Algebra Universalis* 31 (1994), no. 3, 397–406.
- [DIJ] L. Demonet, O. Iyama, G. Jasso, τ -tilting finite algebras, g -vectors and brick- τ -rigid correspondence, arXiv:1503.00285.
- [DIRRT] L. Demonet, O. Iyama, I. Reiten, N. Reading, H. Thomas, *Lattices of torsion classes and their algebraic quotients*, in preparation.
- [DR] V. Dlab, C. M. Ringel, *The preprojective algebra of a modulated graph*, Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), pp. 216–231, Lecture Notes in Math., 832, Springer, Berlin-New York, 1980.
- [GLS] C. Geiss, B. Leclerc, J. Schröer, *Kac-Moody groups and cluster algebras*, *Adv. Math.* 228 (2011), no. 1, 329–433.

- [HHKU] D. Happel, S. Hartlieb, O. Kerner, L. Unger, *On perpendicular categories of stones over quiver algebras*, Comment. Math. Helv. 71 (1996), no. 3, 463–474.
- [IR] O. Iyama, I. Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, Amer. J. Math. 130 (2008), no. 4, 1087–1149.
- [IRTT] O. Iyama, I. Reiten, H. Thomas, G. Todorov, *Lattice structure of torsion classes for path algebras*, Bull. Lond. Math. Soc. 47 (2015), no. 4, 639–650.
- [KS] M. Kashiwara, Y. Saito, *Geometric construction of crystal bases*, Duke Math. J. 89 (1997), no. 1, 9–36.
- [KL] O. Kerner, F. Lukas, *Regular stones of wild hereditary algebras*, J. Pure Appl. Algebra 93 (1994), no. 1, 15–31.
- [K] Y. Kimura, *Tilting theory of preprojective algebras and c -sortable elements*, arXiv:1405.4087.
- [Le] B. Leclerc, *Cluster structures on strata of flag varieties*, to appear in Adv. Math., arXiv:1402.4435.
- [Lu] G. Lusztig, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. 4 (1991), no. 2, 365–421.
- [Ma] F. Marks, *Homological embeddings for preprojective algebras*, arXiv:1512.02442.
- [Mi] Y. Mizuno, *Classifying τ -tilting modules over preprojective algebras of Dynkin type*, Math. Z. 277 (2014) no. 3–4, 665–690.
- [N] H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. J. 76 (1994), no. 2, 365–416.
- [ORT] S. Oppermann, I. Reiten, H. Thomas, *Quotient closed subcategories of quiver representations*, Compos. Math. 151 (2015), no. 3, 568–602.
- [R1] N. Reading, *Lattice congruences of the weak order*. Order 21 (2004) no. 4, 315–344.
- [R2] N. Reading, *Lattice Theory of the Poset of Regions*, in Lattice Theory: Special Topics and Applications, ed. G. Grätzer and F. Wehrung, Birkhäuser, to appear.
- [R3] N. Reading, *Finite Coxeter Groups and the Weak Order*, in Lattice Theory: Special Topics and Applications, ed. G. Grätzer and F. Wehrung, Birkhäuser, to appear.
- [Ri] C. M. Ringel, *Representations of K -species and bimodules*, J. Algebra 41 (1976), no. 2, 269–302.
- [ST] P. Seidel, R. Thomas, *Braid group actions on derived categories of coherent sheaves*, Duke Math. J. 108 (2001), no. 1, 37–108.
- [SY1] Y. Sekiya, K. Yamaura, *Tilting theoretical approach to moduli spaces over preprojective algebras*, Algebr. Represent. Theory 16 (2013), no. 6, 1733–1786.
- [SY2] Y. Sekiya, K. Yamaura, private communications.
- [WW] B. Wald, J. Waschbüsch, *Tame biserial algebras*, J. Algebra 95 (1985), no. 2, 480–500.

O. IYAMA: GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, CHIKUSA-KU, NAGOYA, 464-8602 JAPAN

E-mail address: iyama@math.nagoya-u.ac.jp

URL: <http://www.math.nagoya-u.ac.jp/~iyama/>

N. READING: DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695-8205, USA

E-mail address: reading@math.ncsu.edu

URL: <http://www4.ncsu.edu/~nreadin/>

I. REITEN: DEPARTMENT OF MATHEMATICAL SCIENCES NORGES TEKNISK-NATURVITENSKAPELIGE UNIVERSITET 7491 TRONDHEIM NORWAY

E-mail address: idun.reiten@math.ntnu.no

URL: <http://www.ntnu.edu/employees/idun.reiten>

H. THOMAS: DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, CP 8888, SUCCURSALE CENTRE-VILLE, MONTRÉAL, QC, H3C 3P8, CANADA

E-mail address: hugh.ross.thomas@gmail.com

URL: <http://www.lacim.uqam.ca/~hugh>