## Connecting the UMEB in $\mathbb{C}^d \bigotimes \mathbb{C}^d$ with partial Hadamard matrices

Yan-Ling Wang<sup>1</sup>, Mao-Sheng Li<sup>2</sup>, Shao-Ming Fei<sup>3,4</sup>, Zhu-Jun Zheng<sup>1</sup>

<sup>1</sup>Department of Mathematics, South China University of Technology, Guangzhou 510640, P.R.China

<sup>2</sup>Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China

<sup>3</sup>School of Mathematical Sciences, Capital Normal University, Beijing 100048, China

<sup>4</sup>Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany

We study the unextendible maximally entangled bases (UMEB) in  $\mathbb{C}^d \otimes \mathbb{C}^d$  and connect it with the partial Hadamard matrix. Firstly, we show that for a given special UMEB in  $\mathbb{C}^d \otimes \mathbb{C}^d$ , there is a partial Hadamard matrix can not extend to a complete Hadamard matrix in  $\mathbb{C}^d$ . As a corollary, any  $(d-1) \times d$  partial Hadamard matrix can extend to a complete Hadamard matrix. Then we obtain that for any d there is an UMEB except d = p or 2p, where  $p \equiv 3 \mod 4$  and p is a prime. Finally, we argue that there exist different kinds of constructions of UMEB in  $\mathbb{C}^{nd} \otimes \mathbb{C}^{nd}$  for any  $n \in \mathbb{N}$  and  $d = 3 \times 5 \times 7$ .

PACS numbers: 03.67.Hk,03.65.Ud

## I. INTRODUCTION

It is well known that the quantum states are divided into two parts: separable states and entanglement states. The pure product states are special case of the separable states while the maximally entangled states play important roles for the entangled states [1-4]. One of the significant property of the quantum theory is the quantum nonlocality. An unextendible product bases (UPBs) in bipartite quantum system  $\mathbb{C}^m \bigotimes \mathbb{C}^n$  is a set of orthogonal product states less than mn such that no further product states are orthogonal to every state in that set[5, 6]. It is proven that the UPBs display some nonlocality without entanglement [6, 7]. Similar with the UPBs, in 2009, S. Bravyi and J. A. Smolin first proposed the notion of unextendible maximally entangled basis(UMEB): a set of orthonormal maximally entangled states in  $\mathbb{C}^d \bigotimes \mathbb{C}^d$  consisting of fewer than  $d^2$  vectors which have no additional maximally entangled vectors that are orthogonal to all of them. The authors pointed out that the UMEBs are helpful for constructing some quantum states with special property of the entanglement of assistance(EOF) and can be used to find quantum channels that are unital but not convex mixtures of unitary operations [8].

It is proved that there do not exist UMEBs for d = 2, and a 6-member UMEB for d = 3 and a 12-member UMEB for d = 4 were constructed [8]. After that, the construction of UMEB has attracted the attention some authors. Firstly, there are many UMEBs have been constructed in  $\mathbb{C}^d \otimes \mathbb{C}^{d'} (d \neq d')$ [9, 10]. In Ref.[11], the authors studied the UMEB in  $\mathbb{C}^d \otimes \mathbb{C}^d$ , and gave that if there is an UMEB in  $\mathbb{C}^d \otimes \mathbb{C}^d$  then there is also an UMEB in  $\mathbb{C}^{qd} \otimes \mathbb{C}^{qd}$  for any  $q \in \mathbb{N}$ . Nevertheless, for the UMEB in  $\mathbb{C}^d \otimes \mathbb{C}^d$  we only know the cases d = 3, 4, 3n, 4n. So it is interesting to consider the UMEB in other higher-dimensional system  $\mathbb{C}^d \otimes \mathbb{C}^d$  for general d. In addition, Guo generalized the UMEB problem by replacing the condition of maximality of states with states of given Schmidt number[12, 13].

The construction of Hadamard matrix is also an interesting topic. In this paper, we mainly concern about the complex Hadamard matrix. The authors who are interested in complex Hadamard matrix may look Refs.[14–16] for further reading. A partial Hadamard matrix is a matrix  $H \in M_{m \times n}(\mathbb{T})$  (where  $\mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$ ), whose rows are pairwise orthogonal. Given a partial Hadamard matrix  $H \in M_{m \times n}(\mathbb{T})$  one interesting problem is that of deciding whether this matrix extends or not to an  $n \times n$  complex Hadamard matrix. In the real case, there are many results[17, 18]. But for the general complex case, however, very little seems to be known about this question[19].

In this paper, we show a relation between these two basic concepts, and in particular we show that if there are a special UMEB in  $\mathbb{C}^d \bigotimes \mathbb{C}^d$ , then we can find a corresponding partial Hadamard matrix which can not be extended to a complete Hadamard matrix, and vice versa. Then by using the extendibility of any  $d^2 - 1$  orthogonal maximally entangled states, we give an answer to the conjecture in [19]. The above relation between UMEB and partial Hadamard matrix gives us a method construct UMEB. As a example, we first construct a 23member UMEB in  $\mathbb{C}^5 \bigotimes \mathbb{C}^5$ . Then we generalized the example to higher dimensions: we show that for any  $n \in \mathbb{N}$ , there exists an UMEB in  $\mathbb{C}^{4n+1} \bigotimes \mathbb{C}^{4n+1}$ . At last, we show that there exists an UMEB in  $\mathbb{C}^d \bigotimes \mathbb{C}^d$  except d = p or 2p, where  $p \equiv 3 \mod 4$  and p is a prime. In addition, we also give an UMEB in  $\mathbb{C}^7 \bigotimes \mathbb{C}^7$  for the exceptional unsolved cases. Then by using the UMEBs constructed from d = 3, 5, 7, we show there are different kinds of UMEBs in  $\mathbb{C}^{(3 \times 5 \times 7)n} \bigotimes \mathbb{C}^{(3 \times 5 \times 7)n}$  for any  $n \in \mathbb{N}$ .

## II. THE UMEBS IN $\mathbb{C}^d \otimes \mathbb{C}^d$ AND THE PARTIAL HADAMARD MATRIX

**Definition 1.** A set of states  $\{|\phi_a\rangle \in \mathbb{C}^d \otimes \mathbb{C}^d : a = 1, 2, \dots, n, n < d^2\}$  is called an *n*-number UMEB if and only if (i)  $|\phi_a\rangle$ ,  $a = 1, 2, \dots, n$ , are maximally entangled; (ii)  $\langle \phi_a | \phi_b \rangle = \delta_{ab}$ ; (iii) if  $\langle \phi_a | \psi \rangle = 0$  for all  $a = 1, 2, \dots, n$ , then  $|\psi\rangle$  cannot be maximally entangled.

Here under computational basis a maximally entangled state  $|\phi_a\rangle$  can be expressed as

$$|\phi_a\rangle = (I \otimes U_a) \frac{1}{\sqrt{d}} \sum_{i=1}^d |i\rangle \otimes |i\rangle, \qquad (1)$$

where I is the  $d \times d$  identity matrix,  $U_a$  is any unitary matrix. According to (1), a set of unitary matrices  $\{U_a \in M_d(\mathbb{C}) | a = 1, ..., n\}$  gives an *n*-number UMEB in  $\mathbb{C}^d \otimes \mathbb{C}^d$  if and only if

(i)  $n < d^2;$ 

(ii)  $Tr(U_a^{\dagger}U_b) = d\,\delta_{ab}, \quad \forall a, b = 1, \cdots, n;$ 

(iii) For any  $U \in M_d(\mathbb{C})$ , if  $Tr(U_a^{\dagger}U) = 0, \forall a = 1, \dots, n$ , then U cannot be unitary.

In this paper, we only use the latter equivalent difinition of UMEB.

**Definition 2.**[19] Partial Hadamard matrices: A partial Hadamard matrix in  $\mathbb{C}^n$  is a rectangular matrix H with entries in the circle  $\mathbb{T}$  whose rows are pairwise orthogonal. That is,  $H \in M_{m \times n}(\mathbb{T})(m < n)$ , and  $HH^{\dagger} = nI_m$ .

**Definition 3.[19]** We call a partial Hadamard matrice  $H \in M_{m \times n}(\mathbb{T})$  in  $\mathbb{C}^n$  is completable if there exists a Hadamard matrix  $\widetilde{H}$  whose first m rows equal to the rows of H respectively.

Lemma 1. If there is an N-number UMEB in  $\mathbb{C}^d \otimes \mathbb{C}^d$ , then for any  $q \in \mathbb{N}$ , there is a  $\widetilde{N}$ -number,  $\widetilde{N} = (qd)^2 - (d^2 - N)$ , UMEB in  $\mathbb{C}^{qd} \otimes \mathbb{C}^{qd}[11]$ .

In this paper, we mainly study with the UMEB in  $\mathbb{C}^d \bigotimes \mathbb{C}^d$  containing the following set and we call it a special UMEB if exists.

$$S_0 = \{ X^m Z^n \mid m = 1, 2..., d-1, n = 0, 1, ..., d-1 \}$$

where 
$$X = \sum_{j=0}^{d-1} |j+1\rangle \langle j|, \ Z = \sum_{j=0}^{d-1} \omega_d^j |j\rangle \langle j|, \ \omega_d = e^{\frac{2\pi i}{d}}.$$

Suppose  $A = (a_{st})_{k \times d}$  is a  $k \times d$  partial Hadamard matrix in  $\mathbb{C}^d$ ,  $\alpha_1, \alpha_2, \ldots, \alpha_k$  are rows of A. then we can construct a set of unitary matrices , denoted by  $S(A) = \{diag(\alpha_s) \mid s = 1, 2, \ldots, k\}$ , where  $diag(\alpha_s) = \sum_{t=1}^d a_{st}|t - 1\rangle\langle t - 1|$ .

Then the elements in S(A) are unitary and orthogonal with each other under inner product  $\langle A, B \rangle = Tr(AB^{\dagger})$ .

Proposition 1. Given a partial Hadamard matrix A with  $k \times d$ , then  $S_0 \cup S(A)$  can not be extended to a full maximally entangled base (MEB) if and only if A can not extend to a complete Hadamard matrix.

*Proof.* ⇒: Suppose A can extend to a complete Hadamard matrix. That is there are d - k mutually orthogonal vectors  $\nu_1, \nu_2, \ldots, \nu_{d-k}$  with modules 1 for each entry which are orthogonal to all rows of A. Then  $\{U_j = diag(\nu_j) \mid j = 1, 2, ..., d-k\}$  are unitary matrices which are orthogonal with each other, and lies in the orthogonal complement of  $S_0 \cup S(A)$ . Then  $S_0 \cup S(A) \cup \{U_j \mid j = 1, 2, ..., N_{d-k}\}$ .

j = 1, 2, ..., d - k is a MEB. This is contradicted with  $S_0 \cup S(A)$  can not be extended to MEB.

$$\begin{bmatrix} \nu_1 \\ \nu_1 \\ \nu_2 \\ \vdots \\ \nu_4 \\ \nu_4 \end{bmatrix}$$
 is a Hadamard matrix.

Now we give an answer to the conjecture in [5] which conjecture that any partial Hadamard matrix of  $4 \times 5$  can be complemented to a complete Hadamard matrix.

Corollary 1. If d is an integer,  $d \ge 2$ , and A is a partial Hadamard matrix of  $(d-1) \times d$ . Then A can be complemented to a complete Hadamard matrix.

Proof 1. Since A is a  $(d-1) \times d$  matrix, then we have  $S_0 \cup S(A)$  is a set of maximally entangled states with  $d^2 - 1$  states. By [8], it can be extended to MEB. Hence by Proposition 1, A can be complemented to a Hadamard matrix.

Proof 2. Suppose 
$$A = \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_{d-1} \end{pmatrix}$$
, then we have  

$$\dim_{\mathbb{C}}(\operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2, \dots, \alpha_{d-1}\}) = d - 1,$$

$$\dim_{\mathbb{C}}(\operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2, \dots, \alpha_{d-1}\})^{\perp} = 1.$$

Choosing a nonzero vector

$$\nu_d = (\nu_{d1}, \nu_{d2}, \dots, \nu_{dd}) \in \operatorname{span}_{\mathbb{C}} \{\alpha_1, \alpha_2, \dots, \alpha_{d-1}\}^{\perp}$$
  
such that  $\|\nu_d\| = 1$ . Then  $U = \begin{pmatrix} \frac{\alpha_1}{\sqrt{d}} \\ \frac{\alpha_2}{\sqrt{d}} \\ \vdots \\ \frac{\alpha_{d-1}}{\sqrt{d}} \end{pmatrix}$  is a matrix with

normal rows and orthogonal with each other. That is, U is an unitary matrix. Then all columns of U are also normal and orthogonal with each other. Hence,  $|\nu_{dk}| = \frac{1}{\sqrt{d}}$  for

$$k = 1, 2, \dots, d$$
. Then  $\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_{d-1} \\ \sqrt{d}\nu_d \end{pmatrix}$  is a Hadamard matrix.

**Remark 1:** The Proposition 1 give us a method to construct some sets of UMEB. Suppose there is a partial Hadamard matrix A whose orthogonal complement contains no vector with each entry module 1. Then  $S_0 \cup S(A)$ is an UMEB.

*Example* 1. In  $\mathbb{C}^5 \bigotimes \mathbb{C}^5$ , there exists an UMEB with 23 elements.

Let  $A = \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & \omega & \omega^2 \end{pmatrix}$ , where  $\omega = e^{\frac{2\pi i}{3}}$ . If we denote

$$\nu_1 = \left(\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}}, 0, 0\right), \\
\nu_2 = \left(\frac{1}{\sqrt{10}}, 0, \frac{1}{\sqrt{10}}, \frac{2\omega^2}{\sqrt{10}}, \frac{2\omega}{\sqrt{10}}\right), \\
\nu_3 = \left(0, \sqrt{\frac{3}{5}}, 0, \frac{\omega - 1}{\sqrt{15}}, \frac{\omega^2 - 1}{\sqrt{15}}\right),$$

then span<sub>C</sub>{ $\alpha_1, \alpha_2$ }<sup> $\perp$ </sup> = { $\nu_1, \nu_2, \nu_3$ }. Let  $\alpha = k_1\nu_1 +$  $k_2\nu_2 + k_3\nu_3$  is a vector with each entries module 1, that is

$$\begin{cases} |\frac{k_1}{\sqrt{2}} + \frac{k_2}{\sqrt{10}}| = 1, \\ |-\frac{k_1}{\sqrt{2}} + \frac{k_2}{\sqrt{10}}| = 1, \\ |\sqrt{\frac{3}{5}}k_3| = 1, \\ |k_1|^2 + |k_2|^2 + |k_3|^2 = 1. \end{cases}$$

Then from the above equations, we have  $|k_1| = |k_2| =$  $|k_3| = \sqrt{\frac{5}{3}}$ . Moreover,  $k_2 = \pm ik_1, k_3 = \pm ik_1$ . If we let  $\alpha_3 = \sqrt{\frac{5}{3}}\nu_1 + i\sqrt{\frac{5}{3}}\nu_2 + i\sqrt{\frac{5}{3}}\nu_3$ , then  $B = \begin{pmatrix} A \\ \alpha_3 \end{pmatrix}$ is also a partial Hadamard matrix. However, any vector lies in  $\operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2, \alpha_3\}^{\perp} \subseteq \operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2\}^{\perp}$ . Hence, if  $\nu \in \operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2, \alpha_3\}^{\perp}$  with each entry module 1, then  $\nu$  can be written as the form

$$\nu = k_1 \nu_1 \pm i k_1 \nu_2 \pm i k_1 \nu_3$$

However,  $\nu$  can not be orthogonal with  $\alpha_3$ . Hence, the set  $S_0 \cup S(B)$  is an UMEB with 23 elements in  $\mathbb{C}^5 \otimes \mathbb{C}^5$ . 

Proposition 2. In  $\mathbb{C}^{4n+1}$ , there exists a partial Hadamard matrix which can not complete to a Hadamard matrix. *Proof:* Let

where  $\omega = e^{\frac{2\pi i}{2n}}$ ,  $\sigma = e^{\frac{2\pi i}{2n+1}}$ . Firstly, we compute the orthogonal complement of the subspace V spanned by the rows of A. Obviously,

$$\beta_{1} = (\frac{1}{\sqrt{2}}, \overbrace{0, 0, \cdots, 0}^{2n-1}, -\frac{1}{\sqrt{2}}, \overbrace{0, 0, \cdots, 0}^{2n}),$$
  
$$\beta_{2} = (\frac{1}{\sqrt{8n+2}}, \overbrace{0, 0, \cdots, 0}^{2n-1}, \frac{1}{\sqrt{8n+2}}, \frac{2\sigma^{2n}}{\sqrt{8n+2}}, \frac{2\sigma^{2n-1}}{\sqrt{8n+2}}, \cdots, \frac{2\sigma}{\sqrt{8n+2}})$$

are orthogonal with  $\alpha_1, \alpha_2, \ldots, \alpha_{2n}$  and  $\beta_1 \perp \beta_2$ . Now we set

$$\gamma_{1} = (\overbrace{0, 1, 0, 0, \cdots, 0, 0}^{2n}, \overbrace{0, 0, 0, \cdots, 0}^{2n+1}),$$
  

$$\gamma_{2} = (0, 0, 1, 0, \cdots, 0, 0, 0, 0, 0, \cdots, 0),$$
  

$$\vdots$$
  

$$\gamma_{2n-1} = (0, 0, 0, 0, \cdots, 1, 0, 0, 0, \cdots, 0).$$

By Schmidt orthogonalization we have

$$\beta_{3} = (0, \sqrt{\frac{2n+1}{4n+1}}, 0, 0, \dots, 0, 0, 0, \beta_{31}, \beta_{32}, \dots, \beta_{3,2n}),$$
  

$$\beta_{4} = (0, 0, \sqrt{\frac{2n+1}{4n+1}}, 0, \dots, 0, 0, 0, \beta_{41}, \beta_{42}, \dots, \beta_{4,2n}),$$
  

$$\vdots$$
  

$$\beta_{2n+1} = (0, 0, 0, 0, \dots, 0, \sqrt{\frac{2n+1}{4n+1}}, 0, \beta_{2n+1,1}, \dots, \beta_{2n+1,2n})$$

Then we have  $\operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2, \ldots, \alpha_{2n}\}^{\perp}$ =  $\operatorname{span}_{\mathbb{C}}\{\beta_1, \beta_2, \dots, \beta_{2n+1}\}$ . Suppose  $\nu = k_1\beta_1 + k_2\beta_2 + k_$  $\cdots + k_{2n+1}\beta_{2n+1}$  is a vector with entries module 1. Then we must have

$$\begin{cases} |\frac{1}{\sqrt{2}}k_1 + \frac{1}{\sqrt{8n+2}}k_2| = 1, \\ |\frac{1}{\sqrt{2}}k_1 - \frac{1}{\sqrt{8n+2}}k_2| = 1, \\ |\sqrt{\frac{2n+1}{4n+1}}k_3| = 1, \\ |\sqrt{\frac{2n+1}{4n+1}}k_4| = 1, \\ \vdots \\ |\sqrt{\frac{2n+1}{4n+1}}k_{2n+1}| = 1, \\ |k_1|^2 + |k_2|^2 + \dots + |k_{2n+1}|^2 = |\nu|^2 = 4n + 1 \end{cases}$$

Solving the above equations, we have

$$|k_1| = |k_2| = \dots = |k_{2n+1}| = \sqrt{\frac{4n+1}{2n+1}}$$
, and  $k_1 = \pm ik_2$ .

If A can be extended to a Hadamard matrix, by adding 2n + 1 rows  $\nu_1, \nu_2, \ldots, \nu_{2n+1}$ . Then we have

$$\nu_1 = k_{11}\beta_1 + k_{12}\beta_2 + \dots + k_{1,2n+1}\beta_{2n+1}, 
\nu_2 = k_{21}\beta_1 + k_{22}\beta_2 + \dots + k_{2,2n+1}\beta_{2n+1}, 
\vdots$$

$$\nu_{2n+1} = k_{2n+1,1}\beta_1 + k_{2n+1,2}\beta_2 + \dots + k_{2n+1,2n+1}\beta_{2n+1}$$

The above analysis gives that  $|k_{st}| = \sqrt{\frac{4n+1}{2n+1}}$ . Clearly, the orthogonality of  $\nu_1, \nu_2, \ldots, \nu_{2n+1}$  give that vectors  $(k_{11}, k_{12}, \ldots, k_{1,2n+1}), (k_{21}, k_{22}, \ldots, k_{2,2n+1}), \ldots,$ 

 $(k_{2n+1,1}, k_{2n+1,2}, \ldots, k_{2n+1,2n+1})$  are orthogonal with each other. Hence, if we let  $K = (k_{st})_{(2n+1)\times(2n+1)}$ . Then  $\sqrt{\frac{2n+1}{4n+1}}K$  is a matrix with entries module 1 and each row are mutually orthogonal. Hence,  $H = \sqrt{\frac{2n+1}{4n+1}}K$ is a Hadamard matrix. In the following we show that this can not be true and get a contradiction.

If we replace each row by  $(k_{j1}, k_{j2}, \ldots, k_{j,2n+1})$  by  $\frac{1}{k_{j1}}(k_{j1}, k_{j2}, \ldots, k_{j,2n+1})$ , then the new matrix  $\tilde{H}$  is also a Hadamard matrix with the element of first column all are 1. Noticing that  $k_{j2} = \pm i k_{j1}$  for  $j = 1, 2, \cdots, 2n + 1$  so the elements in second column of  $\tilde{H}$  are i or -i. The Hadamard matrix  $\tilde{H}$  also give that the column of  $\tilde{H}$  are orthogonal with each other. Suppose there are p elements of the sencond column are i and q elements are -i. Then the inner product of the first column and the second column is (p-q)i. Here p+q=2n+1, so  $p \neq q$ . Hence,  $(1,1,\ldots,1)^T$  can not orthogonal with the second column.

Hence we can conclude that A can not be extended to a Hadamard matrix.

Corollary 2. There exists an UMEB in  $\mathbb{C}^{4n+1} \otimes \mathbb{C}^{4n+1}$  for any integer n.

Corollary 3. There exists an UMEB in  $\mathbb{C}^d \otimes \mathbb{C}^d$ , whenever  $d \neq p$  or 2p ( $p \equiv 3 \mod 4$  and p is a prime).

*Proof:* Let  $d = p_1^{r_1} p_2^{r_2} \dots p_k^{r_k}$  where  $p_i$  are primes,  $p_1 < p_2 < \dots < p_k$ , and  $r_i \in \mathbb{N}$  for  $i = 1, 2, \dots, k$ . If  $p_1 =$ 

 $2, r_1 \geq 2$  then we have an UMEB for d is multiple of 4. Else if some  $p_j = 4n + 1(n \in \mathbb{N})$ , from the corollary 2 we have an UMEB. So we can suppose that all the primes are of the for  $p_j = 4n + 3$  except if the first one to be 2. Now suppose there are two primes  $p_j = 4n + 3$  and  $p_s = 4m + 3(m \in \mathbb{N})$ , then we can get  $4t + 1|p_jp_s$  for some integer t, we also can get an UMEB. Then we can get only the situation d = p or 2p, where  $p = 3 \mod 4$ and p is a prime are not solved.

We have solved the most situations, only the cases d = p or 2p, where  $p = 3 \mod 4$  and p is a prime are not solved. Among all the numbers d which are unsolved, 7 is the smallest one. In the following, we solve this case by the same method.

Example 2. In  $\mathbb{C}^7 \otimes \mathbb{C}^7$ , there exists an UMEB with 45 elements.

Let 
$$A = \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \omega & \omega^2 & 1 & i & -1 & -i \\ 1 & \omega^2 & \omega & 1 & -1 & 1 & -1 \end{pmatrix}$$
  
where  $\omega = e^{\frac{2\pi i}{3}}$ , and Obviously,  
 $\beta_1 = (\frac{1}{2}, 0, 0, -\frac{1}{2}, 0, 0, 0).$ 

$$\beta_2 = \left(\frac{1}{\sqrt{14}}, 0, 0, \frac{1}{\sqrt{14}}, \frac{-2i}{\sqrt{14}}, \frac{-2}{\sqrt{14}}, \frac{2i}{\sqrt{14}}\right),$$

are orthogonal with  $\alpha_1, \alpha_2, \alpha_3$  and  $\beta_1 \perp \beta_2$ . Now we set

$$\gamma_1 = (0, 0, 0, 0, 0, 1, 0).$$

By Schmidt orthogonalization we have

$$\beta_3 = (0, \frac{2\omega}{\sqrt{14}}, \frac{2\omega^2}{\sqrt{14}}, 0, \frac{-i}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{i}{\sqrt{14}}).$$

 $\beta_3$  is orthogonal with all the vectors  $\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2$ . Let  $\beta_4$  be a normalized vector and orthogonal with  $\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3$ . Then we can get

$$\beta_4 = (0, \frac{2}{\sqrt{14}}, \frac{-2\omega^2}{\sqrt{14}}, 0, \frac{\omega - 1}{\sqrt{14}}, 0, \frac{\omega - 1}{\sqrt{14}}).$$

Hence we obtain that  $\operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2, \alpha_3\}^{\perp} = \operatorname{span}_{\mathbb{C}}\{\beta_1, \beta_2, \beta_3, \beta_4\}$ . Suppose  $\alpha = k_1\beta_1 + k_2\beta_2 + k_3\beta_3 + k_4\beta_4$  is a vector with each entries module 1, that is

$$\begin{cases} \left|\frac{k_1}{\sqrt{2}} + \frac{k_2}{\sqrt{14}}\right| = 1, \\ \left|-\frac{k_1}{\sqrt{2}} + \frac{k_2}{\sqrt{14}}\right| = 1, \\ \left|\frac{2\omega k_3}{\sqrt{14}} + \frac{2k_4}{\sqrt{14}}\right| = 1, \\ \left|\frac{2\omega^2 k_3}{\sqrt{14}} - \frac{2\omega k_4}{\sqrt{14}}\right| = 1, \\ \left|\frac{-2ik_2}{\sqrt{14}} - \frac{ik_3}{\sqrt{14}} + \frac{(\omega-1)k_4}{\sqrt{14}}\right| = 1, \\ \left|\frac{2ik_2}{\sqrt{14}} + \frac{ik_3}{\sqrt{14}} + \frac{(\omega-1)k_4}{\sqrt{14}}\right| = 1, \\ \left|\frac{-2k_2}{\sqrt{14}} + \frac{2k_3}{\sqrt{14}}\right| = 1, \\ \left|\frac{-2k_2}{\sqrt{14}} + \frac{2k_3}{\sqrt{14}}\right| = 1, \\ \left|k_1\right|^2 + \left|k_2\right|^2 + \left|k_3\right|^2 + \left|k_4\right|^2 = 7. \end{cases}$$

Then from the first two equations above, we obtain the first two equations below. And from the third and fourth equations above, we obtain the third and fourth equations below. So are to the fifth and sixth. Here for two complex number  $z_1 = x_1 + iy_1, z_2 = x_2 + iy_2$ , we write  $z_1 \perp z_2$  by meaning that  $x_1x_2 + y_1y_2 = 0$ .

$$\begin{cases} \frac{|k_1|^2}{2} + \frac{|k_2|^2}{14} = 1, \\ k_1 \perp k_2, \\ \frac{4|k_3|^2}{14} + \frac{4|k_4|^2}{14} = 1, \\ \omega k_3 \perp k_4 \Rightarrow k_3 \perp \omega^2 k_4, \\ \frac{|(\omega-1)k_4|^2}{14} + \frac{|2ik_2 + ik_3|^2}{14} = 1, \\ (\omega - 1)k_4 \perp (2ik_2 + ik_3) \Rightarrow \omega^2 k_4 \perp (2k_2 + k_3), \\ |\frac{-2k_2}{\sqrt{14}} + \frac{2k_3}{\sqrt{14}}| = 1, \\ |k_1|^2 + |k_2|^2 + |k_3|^2 + |k_4|^2 = 7. \end{cases}$$

Then from the above equations, we have  $|k_2|^2 = \frac{7}{4}, \frac{|2k_2+k_3|^2}{14} + \frac{3}{14}|k_4|^2 = 1, |k_3|^2 + |k_4|^2 = \frac{7}{2}, |k_2-k_3|^2 = \frac{7}{2}.$ Since  $\omega^2 k_4 \perp k_3, \omega^2 k_4 \perp (2k_2+k_3)$ , we can get  $\omega^2 k_4 \perp 2k_2$ , then we have  $k_2, k_3$  are  $\mathbb{R}$  linear dependence. So we can suppose  $k_2 = rk_3$ , for some real number r. Substituting this into above four equations, we get the following two equantions:

$$\begin{cases} r^2 - r + \frac{1}{2} = 0, \\ r^2 - 2r - 1 = 0. \end{cases}$$

Then r is unsolvable. So there is no  $k_1, k_2, k_3, k_4$  satisfing the condition. That is, there is no vector in  $\operatorname{span}_{\mathbb{C}}\{\alpha_1, \alpha_2, \alpha_3\}^{\perp}$  with each entry module 1. Hence, by Proposition 1, we have an UMEB in  $\mathbb{C}^7 \otimes \mathbb{C}^7$ .

Actually, when d = 3 the UMEB contains 6 states, so it misses 3 states to form a full base. Similarly, when d = 5 there are 2 states missing, when d = 7 there are 4 states missing. Then there are three ways to obtain the UMEBs for  $d = 3 \times 5 \times 7$  by the method of lemma, respectively from d = 3, 5, 7. The one obtained from d =3 is missing  $3 \times 35 = 105$  states. The one obtained from d = 5 is missing  $2 \times 21 = 42$  states, while the last one obtained from d = 7 is missing  $4 \times 15 = 60$ . So the three UMEBs are different with each other. Hence for the case  $d = 3 \times 5 \times 7$ , there are at least three UMEBs. Moreover, it can be generalized to the case  $d = 3 \times 5 \times 7 \times n$  for any integer n.

## **III. CONCLUSION AND DISCUSSION**

We study the UMEBs in  $\mathbb{C}^d \bigotimes \mathbb{C}^d$  and connect it with the partial Hadamard matrix. We show that the existence of a special UMEB in  $\mathbb{C}^d \bigotimes \mathbb{C}^d$  is equivalent to the existence of an uncompletable partial Hadamard matrix. In particular, as a corollary, we get any  $(d-1) \times d$ partial Hadamard matrices can always extend to a complete Hadamard matrix, which gives an answer to the conjecture in [19]. Actually, the Proposition 1 also give us a method to construct UMEB by using an uncompletable partial Hadamard matrix. Then we prove that there exists an uncompletable partial Hadamard matrix for d = 4n + 1 which implies the existence of an UMEB in  $\mathbb{C}^{4n+1} \bigotimes \mathbb{C}^{4n+1}$ . At last, combining the lemma with the proposition 2, we obtain that for any d there is an UMEB except d = p or 2p, where  $p \equiv 3 \mod 4$  and p is a prime. In addition, we also give an UMEB by the partial Hadamard method when d = 7. We conclude there are at least three different sets of UMEBs in  $\mathbb{C}^d \bigotimes \mathbb{C}^d$  when d is multiple of  $3 \times 5 \times 7$ .

We hope that the paper will be helpful both for the construction of UMEB and the partial Hadamard matrices.

Acknowledgments This work is supported by the NSFC 11475178, NSFC 11571119 and NSFC 11275131.

- M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information(Cambridge University Press, Cambridge, U.K., 2004).
- [2] P. Horodecki, Separability criterion and inseparable mixed states with positive partial transposition, Phys. Lett. A 232, 333(1997).
- [3] M. Horodecki, ENTANGLEMENT MEASURES, Quantum Inf. Comput.1, 3 (2001).
- [4] R. Horodecki, P Horodecki, M Horodecki and K Horodecki, Quantum entanglement, Rev. Mod. Phys 81, 865 (2009).
- [5] C.H. Bennett, D.P. DiVincenzo, T. Mor, P.W. Shor, J.A.Smolin and B.M. Terhal, Unextendible product bases and bound entanglement, Phys. Rev. Lett. 82, 5385 (1999).
- [6] D.P. DiVincenzo, T. Mor, P.W. Shor, J.A.Smolin and

B.M. Terhal, Unextendible product bases, uncompletable product bases and bound entanglement, Commun. Math. Phys. 238,379(2003).

- [7] S.De Rinaldis, Distinguishability of complete and unextendible product bases, Phys.Rev.A, 70, 022309(2004).
- [8] S. Bravyi, and J. A. Smolin, Unextendible maximally entangled bases, Phys. Rev. A 84, 042306 (2011).
- B.Chen and S.M.Fei, Unextendible maximally entangled bases and mutually unbiased bases, Phys. Rev. A 88, 034301 (2013).
- [10] M.-S. Li, Y.-L. Wang, Z.-J. Zheng, Unextendible maximally entangled bases in C<sup>d</sup> ⊗ C<sup>d'</sup>, Phys. Rev. A 89, 062313 (2014).
- [11] Y.-L. Wang, M.-S. Li, S.M.Fei, Unextendible maximally entangled bases in C<sup>d</sup> ⊗ C<sup>d</sup>, Phys. Rev. A 90, 034301 (2014).

- [12] Y. Guo, S. Wu, Unextendible entangled bases with fixed Schmidt number, Phys. Rev. A 90, 054303 (2014).
- [13] Y. Guo, Y. Jia, X. Li, Multipartite unextendible entangled basis, Quantum Inf. Proc. 14, 3553 (2015).
- [14] A. T. Butson, Generalized Hadamard matrices, Proc. Amer. Math. Soc. 13, 894C898(1962).
- [15] G. Zauner, 'Quantendesigns: Grundzauge einer nichtkommutativen Designtheorie' (German) [Quantumdesigns: the foundations of a noncommutative design theory], PhD Thesis, Universitat Wien, 1999, http://www.mat.univie.ac.at/neum/ms/zauner.pdf.
- [16] W. Tadej and K. Zyczkowski, 'A concise guide to complex

Hadamard matrices', Open Syst. Inf. Dyn. **13**, 133C177 (2006).

- [17] E. Verheiden, Integral and rational completions of combinatorial matrices, J. Combin. Theory Ser. A 25, 267 (1978).
- [18] W. de Launey, D.A. Levin, A Fourier-analytic approach to counting partial Hadamard matrices, Cryptogr. Commun. 2 307(2010).
- [19] T. Banica, A. Skalski, The quantum algebra of partial Hadamard matrices, Linear Algebra and its Applications, 469, 364(2015).