Anti-van der Waerden numbers of 3-term arithmetic progressions.

Zhanar Berikkyzy[∗] , Alex Schulte† , and Michael Young‡

September 15, 2018

Abstract

The *anti-van der Waerden number*, denoted by $aw([n], k)$, is the smallest r such that every exact r-coloring of $[n]$ contains a rainbow k-term arithmetic progression. Butler et. al. showed that $\lceil \log_3 n \rceil + 2 \leq \text{aw}([n], 3) \leq \lceil \log_2 n \rceil + 1$, and conjectured that there exists a constant C such that $aw([n], 3) \leq \lceil \log_3 n \rceil + C$. In this paper, we show this conjecture is true by determining aw $([n], 3)$ for all n. We prove that for $7 \cdot 3^{m-2} + 1 \le n \le 21 \cdot 3^{m-2}$,

> $aw([n], 3) = \begin{cases} m+2, & \text{if } n = 3^m \\ m+2, & \text{otherwise} \end{cases}$ $m + 3$, otherwise.

Keywords. arithmetic progression; rainbow coloring; unitary coloring; Behrend construction.

1 Introduction

Let *n* be a positive integer and let $G \in \{[n], \mathbb{Z}_n\}$, where $[n] = \{1, \ldots, n\}$. A k-term arithmetic *progression* $(k-AP)$ of G is a sequence in G of the form

$$
a, a+d, a+2d, \ldots, a+(k-1)d,
$$

where $d \geq 1$. For the purposes of this paper, an arithmetic progression is referred to as a set of the form $\{a, a+d, a+2d, \ldots, a+(k-1)d\}$. An *r*-coloring of G is a function $c: G \to [r]$, and such a coloring is called *exact* if c is surjective. Given $c : G \to [r]$, an arithmetic progression is called *rainbow* (under c) if $c(a + id) \neq c(a + jd)$ for all $0 \leq i < j \leq k - 1$.

The *anti-van der Waerden number*, denoted by $aw(G, k)$, is the smallest r such that every exact r-coloring of G contains a rainbow k-AP. If G contains no k-AP, then $aw(G, k) = |G| + 1$; this is consistent with the property that there is a coloring of G with $aw(G, k) - 1$ colors that has no rainbow k-AP.

An r-coloring of G is *unitary* if there is an element of G that is uniquely colored. The smallest r such that every exact unitary r-coloring of G contains a rainbow k -AP is denoted by $aw_u(G, k)$. Similar to the anti-van der Waerden number, $aw_u(G, k) = |G| + 1$ if G has no k-AP.

Problems involving counting and the existence of rainbow arithmetic progressions have been well-studied. The main results of Axenovich and Fon-Der-Flaass [\[1\]](#page-7-0) and Axenovich and Martin [\[2\]](#page-7-1) deal with the existence of 3-APs in colorings that have uniformly sized color classes. Fox,

[∗]Department of Mathematics, Iowa State University, Ames, IA 50011, USA (zhanarb@iastate.edu)

[†]Department of Mathematics, Iowa State University, Ames, IA 50011, USA (aschulte@iastate.edu)

[‡]Department of Mathematics, Iowa State University, Ames, IA 50011, USA (myoung@iastate.edu)

Jungić, Mahdian, Nešetril, and Radoičić also studied anti-Ramsey results of arithmetic progressions in $[6]$. In particular, they showed that every 3-coloring of $[n]$ for which each color class has density more than $1/6$, contains a rainbow 3-AP. Fox et. al. also determined all values of n for which $aw(\mathbb{Z}_n, 3) = 3$.

The specific problem of determining anti-van der Waerden numbers for $[n]$ and \mathbb{Z}_n was studied by Butler et. al. in [\[4\]](#page-7-3). It is proved in [4] that for $k \geq 4$, $aw([n], k) = n^{1-o(1)}$ and $aw(\mathbb{Z}_n, k) = n^{1-o(1)}$. These results are obtained using results of Behrend [\[3\]](#page-7-4) and Gowers [\[5\]](#page-7-5) on the size of a subset of $[n]$ with no k-AP. Butler et. al. also expand upon the results of $[6]$ by determining aw(\mathbb{Z}_n , 3) for all values of n. These results were generalized to all finite abelian groups in [\[7\]](#page-7-6). Butler et. al. also provides bounds for $aw([n], 3)$, as well as many exact values (see Table [1\)](#page-1-0).

Table 1: Values of $aw([n], k)$ for $3 \leq k \leq \frac{n+3}{2}$.

In this paper, we determine the exact value of $aw([n], 3)$, which answers questions posed in [\[4\]](#page-7-3) and confirms the following conjecture:

Conjecture 1. [\[4\]](#page-7-3) There exists a constant C such that $aw([n], 3) \leq \lceil \log_3 n \rceil + C$, for all $n \geq 3$.

Our main result, Theorem [2,](#page-1-1) also determines $aw_u([n], 3)$ which shows the existence of extremal colorings of [n] that are unitary.

Theorem 2. For all integers $n \geq 2$,

$$
aw_u([n],3) = aw([n],3) = \begin{cases} m+2, & \text{if } n = 3^m \\ m+3, & \text{if } n \neq 3^m \text{ and } 7 \cdot 3^{m-2} + 1 \le n \le 21 \cdot 3^{m-2}. \end{cases}
$$

In section [2,](#page-2-0) we provide lemmas that are useful in proving Theorem [2](#page-1-1) and section [3](#page-5-0) contains the proof of Theorem [2.](#page-1-1)

2 Lemmas

In [\[4,](#page-7-3) Theorem 1.6] it is shown that $3 \leq aw(\mathbb{Z}_p, 3) \leq 4$ for every prime number p and that if $aw(\mathbb{Z}_p, 3) = 4$ then $p \geq 17$. Furthermore, it is shown that the value of $aw(\mathbb{Z}_n, 3)$ is determined by the values of $aw(\mathbb{Z}_p, 3)$ for the prime factors p of n. We have included this theorem below with some notation change.

Theorem 3. [\[4\]](#page-7-3) Let n be a positive integer with prime decomposition $n = 2^{e_0} p_1^{e_1} p_2^{e_2} \cdots p_s^{e_s}$ $for e_i \geq 0, i = 0, \ldots, s, where primes are ordered so that $aw(\mathbb{Z}_{p_i}, 3) = 3$ for $1 \leq i \leq \ell$ and$ $aw(\mathbb{Z}_{p_i}, 3) = 4$ *for* $\ell + 1 \leq i \leq s$ *. Then*

$$
aw(\mathbb{Z}_n,3) = \begin{cases} 2 + \sum_{j=1}^{\ell} e_j + \sum_{j=\ell+1}^{s} 2e_j, & \text{if } n \text{ is odd} \\ 3 + \sum_{j=1}^{\ell} e_j + \sum_{j=\ell+1}^{s} 2e_j, & \text{if } n \text{ is even.} \end{cases}
$$

We use Theorem [3](#page-2-1) to prove the following lemma.

Lemma 4. Let $n \geq 3$, then $\text{aw}(\mathbb{Z}_n, 3) \leq \lceil \log_3 n \rceil + 2$ with equality if and only if $n = 3^j$ or $2 \cdot 3^j$ *for* $j \geq 1$ *.*

Proof. Suppose $n = 2^{e_0} p_1^{e_1} p_2^{e_2} \dots p_s^{e_s}$ with $e_i \ge 0$ for $i = 0, \dots, s$, where primes p_1, p_2, \dots, p_s are ordered so that $aw(\mathbb{Z}_{p_i}, \overline{3}) = 3$ for $1 \leq i \leq \ell$ and $aw(\mathbb{Z}_{p_i}, 3) = 4$ for $\ell + 1 \leq i \leq s$. We consider two cases depending on parity of n.

Case 1. Suppose *n* is odd, that is $e_0 = 0$. Then $aw(\mathbb{Z}_n, 3) = 2 + \sum_{n=1}^{\ell}$ $j=1$ $e_j + \sum_{i=1}^s$ $j=\ell+1$ $2e_j$ by Theorem [3.](#page-2-1) Since $\text{aw}(\mathbb{Z}_p, 3) = 3$ for odd primes $p \leq 13$, we have $p_i \geq 17$ for $i \geq \ell + 1$, and clearly $p_i \geq 3$ for $i \leq \ell$, therefore

$$
3^{\mathrm{aw}(\mathbb{Z}_n,3)} = 3^{2+\sum\limits_{j=1}^{\ell} e_j + \sum\limits_{j=\ell+1}^{s} 2e_j} = 9 \cdot 3^{e_1} \cdots 3^{e_\ell} \cdot 9^{e_{\ell+1}} \cdots 9^{e_s} \leq 9 \cdot p_1^{e_1} \cdots p_s^{e_s} = 9n.
$$

Note that the equality holds if and only if n is a power of 3, that is $e_j = 0$ for $2 \leq j \leq s$. Therefore, $aw(\mathbb{Z}_n, 3) \leq \lceil \log_3 n \rceil + 2$ for odd n, with equality if and only if $n = p_1^{e_1}$.

Case 2. Suppose *n* is even, that is $e_0 \geq 1$. Then $aw(\mathbb{Z}_n, 3) = 3 + \sum_{k=1}^{\ell}$ $j=1$ $e_j + \sum^s$ $j = \ell+1$ $2e_j$ by Theorem [3.](#page-2-1) If $n = 2^{e_0} \cdot 3^j$ for $j \ge 1$, then by direct computation $\text{aw}(\mathbb{Z}_n, 3) = 3 + j \le 2 + \lceil \log_3 n \rceil$, with equality if and only if $e_0 = 1$. So suppose there is i such that $p_i \neq 3$, and let $h = \frac{n}{2e_0}$ $\frac{n}{2^{e_0}p_i^{e_i}}.$

If $i \leq \ell$ then $p_i \geq 5$, and so $3 \cdot 3^{e_i} < 2^{e_0} p_i^{e_i}$ for all $e_0 \geq 1$ and $e_i \geq 1$. Therefore, since h is odd, by the previous case

 $3^{\text{aw}(\mathbb{Z}_n,3)} = 3 \cdot 3^{e_i} \cdot 3^{\text{aw}(\mathbb{Z}_h,3)} \leq 3 \cdot 3^{e_i} \cdot 9h < 2^{e_0} p_i^{e_i} \cdot 9h = 9n.$

If $i \geq \ell + 1$ then $p_i \geq 17$, and so $3 \cdot 9^{e_i} < 2^{e_0} p_i^{e_i}$ for all $e_0 \geq 1$ and $e_i \geq 1$. Then by the previous case

$$
3^{\mathrm{aw}(\mathbb{Z}_n,3)} = 3 \cdot 9^{e_i} \cdot 3^{\mathrm{aw}(\mathbb{Z}_h,3)} \le 3 \cdot 9^{e_i} \cdot 9h < 2^{e_0} p_i^{e_i} \cdot 9h = 9n.
$$

A set of consecutive integer I in $[n]$ is called an *interval* and $\ell(I)$ is the number of integers in I. Given a coloring c of some finite nonempty subset S of [n], a *color class* of a color i under c in S is denoted $c_i(S) := \{x \in S : c(x) = i\}$. A coloring c of [n] is *special* if $n = 7q + 1$ for some positive integer q, $c(1)$ and $c(n)$ are both uniquely colored, and there are two colors α and β such that $c_\alpha([n]) = \{q+1, 2q+1, 4q+1\}$ and $c_\beta([n]) = \{3q+1, 5q+1, 6q+1\}.$

Lemma 5. *Let* N *be an integer and* c *be an exact* r*-coloring of* [N] *with no rainbow* 3*-AP, where* 1 and N are colored uniquely. Then either the coloring c is special or $\left|\left\{c(x): x \equiv i\right\}\right|$ $p(\text{mod } 3)$ *and* $x \in [N]$ } $\geq r - 1$ *for* $i = 1$ *or* $i = N$ *.*

Proof. Observe that N is even, otherwise $\{1,(N+1)/2,N\}$ is a rainbow 3-AP. We partition the interval [N] into four subintervals $I_1 = \{1, ..., [N/4]\}, I_2 = \{[N/4] + 1, ..., N/2\}, I_3 =$ $\{N/2 + 1, \ldots, \lfloor 3N/4 \rfloor\}$, and $I_4 = \{\lfloor 3N/4 \rfloor + 1, \ldots, N\}$. Notice that every color other than $c(1)$ and $c(N)$ must be used in the subinterval I_2 . To see this, assume i is the missing color in I_2 distinct from $c(1)$ and $c(N)$. Let x be the largest integer in $c_i(I_1)$. Since N is even, we have $2x - 1 \leq 2\lceil N/4 \rceil - 1 \leq N/2$, and so $2x - 1 \in I_2$ and $c(2x - 1) \neq i$. Therefore the 3-AP $\{1, x, 2x - 1\}$ is a rainbow. If there is no such integer x in I_1 , then the integers colored with i must be in the second half of the interval $[N]$, so we choose the smallest such integer y in $c_i(I_3 \cup I_4)$. Then $\{2y - N, y, N\}$ is a rainbow 3-AP since $c(2y - N) \neq i$, because $2y - N \in I_1 \cup I_2$. Similarly, every color other than $c(1)$ and $c(N)$ must be used in the subinterval I_3 .

Throughout the proof we mostly drop (mod 3) and just say congruent even though we mean congruent modulo 3. We consider the following three cases.

Case 1: $N \equiv 0 \pmod{3}$. Assume $|\{c(x): x \equiv i \pmod{3} \text{ and } x \in [N]\}| < r-1$ for both $i = 1$ and $i = N$. So there are two colors, say red and blue, such that no integer in [N] colored with red is congruent to 1, and no integer in $[N]$ colored with *blue* is congruent to 0. We further partition the interval I_2 into subintervals $I_{2(i)}$ and $I_{2(ii)}$ so that $\ell(I_{2(i)}) \leq \ell(I_{2(ii)}) \leq \ell(I_{2(i)}) + 1$, and partition the interval I_3 into subintervals $I_{3(i)}$ and $I_{3(ii)}$ so that $\ell(I_{3(ii)}) \leq \ell(I_{3(i)}) \leq \ell(I_{3(ii)}) + 1$. Then we have the following observations:

(i) $x \equiv 0$ for all $x \in c_{red}(I_3 \cup I_4)$ and $y \equiv 1$ for all $y \in c_{blue}(I_1 \cup I_2)$.

If there is an integer r in $I_3 \cup I_4$ colored with red and congruent to 2, then $2r - N \equiv 1$, and so $c(2r-N)$ is not red by our assumption. Therefore the 3-AP $\{2r-N, r, N\}$ is rainbow. Similarly, if there is an integer b in $I_1 \cup I_2$ colored with blue and congruent to 2, then $2b - 1 \equiv 0$, and so $c(2b-1)$ is not *blue*, forming a rainbow 3-AP $\{1, b, 2b-1\}.$

(ii) $x \equiv 2$ for all $x \in c_{red}(I_2)$ and $y \equiv 2$ for all $y \in c_{blue}(I_3)$.

If there is an integer r in $c_{red}(I_2)$ congruent to 0, then $2r - 1 \equiv 2$ and $2r - 1 \in I_3 \cup I_4$ since $2r-1 \geq N/2+1$. Therefore, $2r-1$ is not colored with red by the previous observation, and so the 3-AP $\{1, r, 2r - 1\}$ is a rainbow. Similarly, if there is an integer b in $c_{blue}(I_3)$ congruent to 1, then using N we obtain the rainbow 3-AP $\{2b-N, b, N\}$, because $2b-N \equiv 2$ and $2b-N \le N/2$.

 $(iii) \ c_{red}(I_{3(ii)}) = c_{blue}(I_{2(i)}) = \emptyset.$

If there is an integer r in $I_{3(ii)}$ colored with red, then $2r-N \equiv 0$, by observation *(i)*. Furthermore, $2r - N \le N/2$ and $2r - N \ge 2(N/2 + \ell(I_{3(i)}) + 1) - N \ge (2\ell(I_{3(i)}) + 1) + 1 \ge \lceil N/4 \rceil + 1$. So

 \Box

 $2r-N \in I_2$ and hence it is not colored with red by observation (ii). Therefore, $\{2r-N, r, N\}$ is a rainbow 3-AP. Similarly, if there is an integer b in $I_{2(i)}$ colored with blue, then $2b - 1 \equiv 1$ and $N/2 + 1 \leq 2b - 1 \leq |3N/4|$. So $2b - 1 \in I_3$ and hence it is not colored with *blue* by observation (*ii*). Therefore, $\{1, b, 2b - 1\}$ is a rainbow 3-AP.

$(iv) \ c_{red}(I_{2(ii)}) = c_{blue}(I_{3(i)}) = \emptyset.$

Suppose there is an integer r in $I_{2(ii)}$ colored with red. Since the coloring of I_2 contains both red and blue and there is no integer in $I_{2(i)}$ colored with blue, by (iii), there must be an integer b in $I_{2(ii)}$ colored with blue. By (i) and (ii), $b \equiv 1$ and $r \equiv 2$. Wlog, suppose $b > r$. Then $2r - b \equiv 0$ and $2r - b \in I_2$ since $\ell(I_{2(ii)}) \leq \ell(I_{2(i)}) + 1$. So $2r - b$ is not colored red or blue and hence the 3-AP $\{2r - b, r, b\}$ is rainbow. Therefore, there is no integer in $I_{2(ii)}$ that is colored with red. Similarly, there is no integer in $I_{3(i)}$ that is colored with *blue*.

Recall that every color other than $c(1)$ and $c(N)$ is used in both intervals I_2 and I_3 . Therefore, sets $c_{red}(I_{2(i)})$, $c_{blue}(I_{2(ii)})$, $c_{red}(I_{3(i)})$, and $c_{blue}(I_{3(ii)})$ are nonempty. Using above observations we next show that in fact these integers colored with *blue* and red in each subinterval are unique. Let $B = \{b_1, \ldots, b_2\}$ be the shortest interval in $I_{2(ii)}$ which contains all integers colored with blue and let $R = \{r_1, \ldots, r_2\}$ be the shortest interval in $I_{3(i)}$ which contains all integers colored with red. Choose the largest integer x in $c_{red}(I_{2(i)})$ and consider two 3-APs $\{x, b_1, 2b_1 - x\}$ and ${x, b_2, 2b_2 - x}$. Since x is congruent to 2 and both b_1 and b_2 are congruent to 1, we have that both $2b_1 - x$ and $2b_2 - x$ are congruent to 0 and are contained in I_3 , otherwise the 3-APs are rainbow. Since all integers colored with *blue* in I_3 are congruent to 2 by *(ii)*, we have that $2b_1-x$ and $2b_2 - x$ are both colored with red and so contained in R. Therefore, $2\ell(B) - 1 \leq \ell(R)$. Now using the smallest integer in $c_{blue}(I_{3(ii)})$, we similarly have that $2\ell(R) - 1 \leq \ell(B)$. Since $\ell(B) \geq 1$ and $\ell(R) \geq 1$, we have that $\ell(R) = \ell(B) = 1$, i.e. there are unique integers b in $c_{blue}(I_{2(ii)})$ and r in $c_{red}(I_{3(i)})$.

Now for any integer \tilde{r} from $c_{red}(I_{2(i)})$ the integer $2\tilde{r} - 1$ must be colored with red, otherwise the 3-AP $\{1, \tilde{r}, 2\tilde{r} - 1\}$ is rainbow. Since $2\tilde{r} - 1 \in I_3$, it must be equal to the unique red colored integer r of I_3 . Therefore, there is exactly one such \tilde{r} in $I_{2(i)}$, i.e. $c_{red}(I_{2(i)}) = {\tilde{r}}$. Similarly, using N there is a unique integer b in $I_{3(ii)}$ colored with blue. Since $\{1, \tilde{r}, r\}$, $\{\tilde{r}, b, r\}$, $\{b, r, b\}$, and $\{b, \tilde{b}, N\}$ are all 3-APs, $N = 7(\ell(\{b, \ldots, r\}) - 1) + 1 = 7(r - b) + 1$.

Observe that if \tilde{r} is even, the integer $(\tilde{r} + N)/2$ in 3-AP $\{\tilde{r}, (\tilde{r} + N)/2, N\}$ must be red and congruent to 1 since $\tilde{r} \equiv 2$ by *(ii)*, contradicting our assumption. So \tilde{r} is odd, and hence the integer $r' = (\tilde{r} + 1)/2$ in I_1 must be colored with red. Notice that there cannot be another integer x larger than r' in $c_{red}(I_1)$, otherwise $2x - 1$ will be another integer colored with red in I₂ distinct from \tilde{r} . Now, since $\ell(\lbrace r', \ldots, \tilde{r} \rbrace) = \ell(\lbrace b, \ldots, r \rbrace)$ we have that $\lbrace r', r, N \rbrace$ is a 3-AP, and so r' must be even. Suppose there are integers smaller than r' in $c_{red}(I_1)$, and let z be the largest of them. Then $2z - 1$ is also in $c_{red}(I_1)$ and must be equal to or larger than r' in I_1 . However, that is impossible because r' is even and there is no integer in $c_{red}(I_1)$ larger than r'. So r' is a unique integer in I_1 colored with red. Similarly, there is a unique integer b' in I_4 colored with *blue*. Therefore the 8-AP can be formed using integers $1, r', \tilde{r}, b, r, \tilde{b}, b', N$ since $\ell(\{1,\ldots,r'\}) = \ell(\{r',\ldots,\tilde{r}\}) = \ell(\{\tilde{r},\ldots,b\}) = \ell(\{b,\ldots,r\}) = \ell(\{r,\ldots,\tilde{b}\}) = \ell(\{\tilde{b},\ldots,b'\}) = \ell(\{b',\ldots,b'\}) = \$ $\ell({b',\ldots,N}).$

In order for this coloring to be special, it remains to show that $c_{blue}(I_1) = c_{red}(I_4) = \emptyset$. If $c_{blue}(I_1) \neq \emptyset$, then choose the largest integer y in it and consider the 3-AP $\{1, y, 2y - 1\}$. Since $2y - 1$ must be in $c_{blue}(I_2)$ and the only integer in this set is b, we have $2y - 1 = b$. However, we know that b is even because $b = 2b - N$, a contradiction. Similarly, if $c_{red}(I_4) \neq \emptyset$ choose the

smallest integer x in it and consider the 3-AP $\{2x - N, x, N\}$. Since $2x - N$ must be in $c_{red}(I_3)$ and the only integer in this set is r, we have $2x-N=r$. However, we know that r is odd because $r = 2\tilde{r} - 1$, a contradiction. This implies that $c_{red}([N]) = \{r', \tilde{r}, r\}$ and $c_{blue}([N]) = \{b, \tilde{b}, b'\}$, so the coloring is special.

Case 2: $N \equiv 2 \pmod{3}$. This case is analogous to Case 1.

Case 3: $N \equiv 1 \pmod{3}$. Assume $|\{c(x): x \equiv i \pmod{3} \text{ and } x \in [N]\}| < r-1$ i.e. there are two colors, say red and blue, such that no integer in $[N]$ colored with red or blue is congruent to 1. Recall that every color other than $c(1)$ and $c(N)$ appears in I_2 and I_3 . First, notice that all integers colored with red or blue in I_2 must be congruent modulo 3. Otherwise, choosing a red colored integer and a blue colored integer, we obtain a 3-AP whose third term is colored with red or blue and is congruent to 1 contradicting our assumption. Similarly, this is also the case for I₃. So suppose all integers in $c_{red}(I_2) \cup c_{blue}(I_2)$ and $c_{red}(I_3) \cup c_{blue}(I_3)$ are congruent modulo 3 to integers $p \neq 1$ and $q \neq 1$, respectively. Pick the largest integers from $c_{red}(I_2)$ and $c_{blue}(I_2)$ and form a 3-AP whose third term is in I_3 . Then the third term is colored with red or blue and is congruent to p. Therefore, $p \equiv q \not\equiv 1$.

We further partition the interval I_2 into subintervals $I_{2(i)}$ and $I_{2(ii)}$, so that $\ell(I_{2(i)}) \leq$ $\ell(I_{2(ii)}) \leq \ell(I_{2(i)}) + 1$. If there exists $x \in c_{red}(I_{2(i)}) \cup c_{blue}(I_{2(i)})$, the integer $2x - 1$ must be colored with $c(x)$ and contained in I_3 , so $2x - 1 \equiv p$ while $x \equiv p \not\equiv 1$, a contradiction. So $c_{red}(I_{2(i)}) \cup c_{blue}(I_{2(i)}) = \emptyset$. However, then the smallest integers of $c_{red}(I_{2(ii)})$ and $c_{blue}(I_{2(ii)})$ form a 3-AP whose first term is contained in $I_{2(i)}$ and is colored with red or blue, a contradiction. This completes the proof of the lemma.

3 Proof of Theorem [2](#page-1-1)

Given a positive integer n , define the function f as follows:

$$
f(n) = \begin{cases} m+2, & \text{if } n = 3^m \\ m+3, & \text{if } n \neq 3^m \text{ and } 7 \cdot 3^{m-2} + 1 \le n \le 21 \cdot 3^{m-2}. \end{cases}
$$

In this section, we prove Theorem [2](#page-1-1) by showing that $\text{aw}([n], 3) = f(n)$ for all n.

First, we show that $f(n) \leq aw_u([n], 3)$ by inductively constructing a unitary coloring of $[n]$ with $f(n) - 1$ colors and no rainbow 3-AP. The result is true for $n = 1, 2, 3$, by inspection. Suppose $n > 3$ and that the result holds for all positive integers less than n. Let $n = 3h - s$, where $s \in \{0, 1, 2\}$ and $2 \leq h < n$.

Let $r = aw_u([h], 3)$. So there is an exact unitary $(r-1)$ -coloring c of $[h]$ with no rainbow 3-AP. Let red be a color not used in c. Define the coloring c_1 of $[n]$ such that if $x \equiv 1 \pmod{3}$, then $c_1(x) = c((x+2)/3)$, otherwise color x with red. When $s \neq 0$, define the coloring c_2 of $[n]$ as follows: if $x \not\equiv 0 \pmod{3}$ then color x with red; if $x \equiv 0 \pmod{3}$ then $c_2(x) = c(x/3 + 1)$ when $c(h)$ is the only unique color in c and $c_2(x) = c(x/3)$ otherwise. Notice that c_2 is a unitary $aw_u([h-1], 3)$ -coloring when $s \neq 0$ and c_1 is a unitary r-coloring of [n]. Now consider a 3-AP $\{a, b, 2b-a\}$ in [n]. If $a \equiv b \not\equiv 1$, then a and b are colored with red, and so the 3-AP is not a rainbow. If $a \equiv b \equiv 1$, then $2b - a \equiv 1$, so this set corresponds to a 3-AP in [h] with coloring c, and hence the 3-AP is not rainbow. If $a \neq b$, then $2b-a$ is not congruent to a or b, so two of the terms of the 3-AP are colored with red, and hence the 3-AP is not rainbow under c_1 . Similarly, this 3-AP is not rainbow under c_2 . Therefore, c_1 and c_2 are unitary colorings of $[n]$ with no rainbow

 \Box

3-AP. Also note that $aw_u([n], 3) \ge aw_u([n], 3) + 1$ under c_1 and $aw_u([n], 3) \ge aw_u([h-1], 3) + 1$ under c_2 . We proceed with three cases determined by $\frac{n}{3}$.

Case 1. First suppose $7 \cdot 3^{m-2} + 1 \le n \le 3^m - 3$ or $3^m \le n \le 21 \cdot 3^{m-2}$. By the induction hypothesis and using the coloring c_1 ,

$$
aw_u([n],3) \ge aw_u([h],3) + 1 \ge f(h) + 1 = f(n).
$$

Case 2. Suppose $n = 3^m - t$ where $t \in \{1, 2\}$. Notice that $h = 3^{m-1}$, so by induction and using coloring c_2 ,

$$
aw_u([n],3) \ge aw_u([h-1],3) + 1 \ge f(h-1) + 1 = f(3^{m-1} - 1) + 1 = (m+2) + 1 = f(n).
$$

The upper bound, $\text{aw}([n], 3) \leq f(n)$, is also proved by induction on n. For small n, the result follows from Table [1.](#page-1-0) Assume the statement is true for all value less than n , and let $7 \cdot 3^{m-2} + 1 \le n \le 21 \cdot 3^{m-2}$ for some m. Let $aw([n]) = r + 1$, so there is an exact r-coloring ĉ of [n] with no rainbow 3-AP. We need to show that $r \leq f(n) - 1$. Let $[n_1, n_2, \ldots, n_N]$ be the shortest interval in [n] containing all r colors under \hat{c} . Define c to be an r-coloring of [N] so that $c(j) = \hat{c}(n_j)$ for $j \in \{1, ..., N\}$. By minimality of N the colors of 1 and N are unique. If [N] has at least $r-1$ colors congruent to 1 or N, then [n] has at least $r-1$ colors congruent to n_1 or n_N , respectively, so $r \leq aw(\lfloor n/3 \rfloor)$ and by induction $r \leq f(\lfloor n/3 \rfloor) \leq f(n) - 1$. So suppose that is not the case, then by Lemma [5](#page-3-0) we have that the coloring c is special.

Let $N = 7q+1$ for some $q \ge 1$, and let the 8-AP in this special coloring be $\{1, r_1, r_2, b_1, r_3, b_2, b_3, N\}$, where r_1, r_2, r_3 are the only integers colored red, b_1, b_2, b_3 are the only integers colored blue and $q = r_1 - 1$. If $n \geq 9q$, then the 8-AP can be extended to a 9-AP in n by adding the 9th element to either the beginning or the ending. Wlog, suppose $\{1, r_1, r_2, b_1, r_3, b_2, b_3, N, 2N - b_3\}$ correspond to a 9-AP in [n]. Since the coloring has no rainbow 3-AP, the color of $2N - b_3$ is blue or $c(N)$, so we have a 4-coloring of this 9-AP. However, $aw([9], 3) = 4$ and hence there is a rainbow 3-AP in this 9-AP which is in turn a rainbow 3-AP in [n]. Therefore, $n \leq 9q - 1$.

By uniqueness of red colored integer r_1 in interval $\{1, \ldots, r_2 - 1\}$, the colors of integers in interval $\{r_1 + 1, \ldots, r_2 - 1\}$ is the same as the reversed colors of integers in $\{2, \ldots, r_1 - 1\}$, i.e. $c(r_1+i) = c(r_1-i)$ for $i = 1, \ldots, q-1$. Similarly, coloring of integers in interval $\{r_2+1, \ldots, b_1-1\}$ is the reversed of the coloring of integers in interval $\{r_1 + 1, \ldots, r_2 - 1\}$, and so on. This gives a rainbow 3-AP-free $(r-2)$ -coloring of \mathbb{Z}_{2q} . Therefore, $r-2 \leq aw(\mathbb{Z}_{2q}, 3) - 1$.

If $q = 3^i$ for some i, then n can not be a power of 3 because $7 \cdot 3^i + 1 \le n \le 9 \cdot 3^i - 1$. Suppose $n = 3^m$, then 2q is not twice a power of 3 and clearly 2q is not a power of 3. Therefore, by Lemma [4](#page-2-2) we have

$$
r\leq aw(\mathbb{Z}_{2q},3)+1\leq \lceil \log_3(2q) \rceil+2 \leq \lceil \log_3(2n/7) \rceil+2 = \lceil \log_3(2\cdot 3^m/7) \rceil+2=m+1\leq f(n)-1.
$$

Suppose now that $n \neq 3^m$. If $q = 3^i$ for some i then $i \leq m-2$. Otherwise, if $i \geq m-1$ then $q \geq 3^{m-1} \geq 1/7n$ which contradicts the fact that $q < 1/7n$. Therefore, $2q \leq 2 \cdot 3^{m-2} = 18 \cdot 3^{m-4}$ and so by induction and Lemma [4,](#page-2-2) $r \leq aw(\mathbb{Z}_{2q}, 3) + 1 = aw([2q], 3) + 1 \leq m + 2 \leq f(n) - 1$. If q is not a power of 3, then again using Lemma [4,](#page-2-2) $r \leq aw(\mathbb{Z}_{2q}, 3) + 1 \leq aw([2q], 3)$. Notice that $6 \cdot 3^{m-3} + 2/7 \le 2n/7 \le 18 \cdot 3^{m-3}$, and so $aw([2q], 3) \le m+2$ by induction. Therefore, $r \leq m + 2 \leq f(n) - 1.$

References

- [1] M. Axenovich, D. Fon-Der-Flaass, On rainbow arithmetic progressions. *Electronic Journal of Combinatorics* 11 (2004), no. 1, Research Paper 1, 7pp.
- [2] M. Axenovich and R.R. Martin, Sub-Ramsey numbers for arithmetic progressions. *Graphs Comb.* 22 (2006), no. 1, 297–309.
- [3] F.A. Behrend, On sets of integers which contain no three terms in arithmetical progression. *Proc. Nat. Acad. Sci. USA* 32 (1946), 331–332.
- [4] S. Butler, C. Erickson, L. Hogben, K. Hogenson, L. Kramer, R. Kramer, J. Lin, R. Martin, D. Stolee, N. Warnberg, and M. Young, Rainbow arithmetic progressions. arXiv preprint arXiv:1404.7232 (2014).
- [5] W.T. Gowers, A new proof of Szemerédi's theorem. *Geom. Funct. Anal.* **11** (2001), no. 3, 465–588.
- [6] V. Jungić, J. Licht (Fox), M. Mahdian, J. Nešetril, and R. Radoičić, Rainbow arithmetic progressions and anti-Ramsey results. *Comb. Probab. Comput.* 12 (2003), no 5-6, 599–620.
- [7] M. Young, Rainbow arithmetic progressions in finite abelian groups. Manuscript available at arXiv:1603.08153 [math.CO].