

# All januarials constructed from Hecke groups

Saadia Mehwish<sup>a</sup>, Qaiser Mushtaq<sup>b</sup>

<sup>a</sup>Department of Mathematics, The Islamia University of Bahawalpur, Bahawalpur, Pakistan

<sup>b</sup>The Islamia University of Bahawalpur, Bahawalpur, Pakistan

---

## Abstract

Professor Graham Higman defined januarial as a special instance of map constructed from embedding of a coset diagram for an action of  $\Delta(2, \ell, k)$ , on finite sets yielding exactly two orbits of the product of the two generators, having equal sizes. In this paper we determine a condition for the existence of a januarial from  $\Delta(2, \ell, k)$ , the quotients of Hecke groups  $H_{\Lambda_\ell}$ , when acting on the projective lines over finite fields  $PL(F_q)$ . We develop a method to find all the januarials from Hecke groups  $H_{\Lambda_\ell}$ , when the triangle group  $\Delta(2, \ell, k)$  acts on  $PL(F_q)$ . We develop a formula for calculating genus of coset diagram depending on the fixed points. By using it, we determine genus of the januarials.

---

## 1. Introduction

It is well known that triangle group  $\Delta(m, \ell, k)$  has a presentation  $\langle s, t : s^m = t^\ell = (st)^k = 1 \rangle$  (e.g. see [2], [3] and [7]). For januarials, we need to fix  $m = 2$ .

The widely studied Hecke groups  $H_{\Lambda_\ell}$  admit a presentation

$$\langle a, b : a^2 = (ab)^\ell = 1 \rangle$$

where  $a : z \rightarrow \frac{-1}{z}$ ,  $b : z \rightarrow z + \Lambda_\ell$  and  $\Lambda_\ell = 2 \cos(\pi/\ell)$ . In another interpretation  $H_{\Lambda_\ell} \cong C_2 \otimes C_\ell$ , that is,  $H_{\Lambda_\ell} = \Delta(2, \ell, \infty)$ . Thus  $\Delta(2, \ell, k)$  are homomorphic images of  $H_{\Lambda_\ell}$ . Throughout this paper we consider triangle groups as quotients of  $H_{\Lambda_\ell}$ .

Let  $q$  be a prime-power  $p^r$ . Then the projective line over Galois field  $F_q$ , contains elements of  $F_q$ , together with  $\infty$ , and is denoted by  $PL(F_q)$ .

A coset diagram is a graphical representation of an action of any group with finite presentation on a set. For instance, a coset diagram for the permutation action of Hecke group on projective line has  $\ell - gon$  for each cycle of  $y$  and an edge for each transposition of  $x$ .

Genus of a connected orientable surface is the number of handles on it. Genus of a graph is the minimal integer  $m$  such that the graph can be drawn on the surface of genus  $m$  without crossings [10].

Let  $H_{\Lambda_\ell} \cong G = \langle x, y : x^2 = y^\ell = 1 \rangle$  where  $x, y$  are linear fractional transformations  $z \rightarrow \frac{az+b}{cz+d}$  where  $a, b, c, d \in \mathbb{Z}$  and  $ad - bc = 1$ . Let  $\alpha$  be a non-degenerate homomorphism from  $G$  to  $PGL(2, q)$ . The two non-degenerating homomorphisms  $\alpha$  and  $\beta$  are said to be conjugates if there exist an inner automorphism  $\rho$  such that  $\alpha\rho = \beta$ . Let  $\bar{x} = \alpha x$ ,  $\bar{y} = \alpha y$  and  $X, Y$  be the matrices representing  $\bar{x}$  and  $\bar{y}$  respectively. Then  $\theta = (trXY)^2 / \det XY$  is invariant of the conjugacy classes of  $\alpha$ .

---

Email addresses: saadiamehwish@gmail.com (Saadia Mehwish), pir\_qmushtaq@yahoo.com (Qaiser Mushtaq)

If  $S_1, S_2, S_3, \dots, S_n$  are finite sets then inclusion-exclusion principle is symbolically expressed as

$$\begin{aligned}
|S_1 \cup S_2 \cup S_3 \cup \dots \cup S_n| &= \sum_{i=1}^n |S_i| - \sum_{1 \leq i < j \leq n} |S_i \cap S_j| \\
&+ \sum_{1 \leq i < j < l \leq n} |S_i \cap S_j \cap S_l| - \dots + \\
&(-1)^{n+1} |S_1 \cap S_2 \cap S_3 \cap \dots \cap S_n|.
\end{aligned} \tag{1}$$

As every integer can be uniquely written as the product of primes so let  $n = p_1^{r_1} p_2^{r_2} \dots p_s^{r_s}$ . Using inclusion-exclusion principle the Euler's phi function  $\phi(n)$  is expressible as (see [9])

$$\phi(n) = n - \sum_{i=1}^s \frac{n}{p_i} + \sum_{1 \leq i < j \leq s} \frac{n}{p_i p_j} - \dots + (-1)^s \frac{n}{p_1 p_2 \dots p_s}. \tag{2}$$

Professor Graham Higman FRS introduced januarials in his last lectures in 2001. He conceived the idea of januarials during his work on Hurwitz groups, that are non trivial quotients of  $\Delta(2, 3, 7)$  group, which he never published. After his death a brief account of his lectures concerning januarials was published [1].

In [1] associates are used to find januarials from the modular group. But his method does not tell, how many times one should repeat the process of taking associates to ultimately get a januarial.

In this paper we prove when januarials exist and then use some pre-existing tools to get an appropriate method of finding januarials. Further we show all the distinct januarials constructible from the Hecke group  $H_{\Lambda_\ell}$ .

## 2. A condition for the existence of januarials

In this section, we determine when januarials exist in the action of triangle groups that are the quotients of Hecke group  $H_{\Lambda_\ell}$  on  $PL(F_q)$ .

**Theorem 1** *A januarial exists in the action of quotients of  $H_{\Lambda_\ell}$ ,  $\Delta(2, \ell, k)$ , on  $PL(F_q)$  if and only if  $k = (q + 1)/2$ , for all  $q > 3$  where  $q = p^r$  and  $p \neq 2$ .*

*Proof:* If

$$k = (q + 1)/2 \text{ then } (xy)^{(q+1)/2} = 1$$

and the action of  $xy$  on  $PL(F_q)$  gives the permutation  $\rho_{xy} \in S_{q+1}$  such that

$$(\rho_{xy})^{(q+1)/2} = 1.$$

This means that the lengths  $r_i$  of cycles of  $\rho_{xy}$  are divisors of  $(q + 1)/2$ . Here two cases arise: when  $r_i \leq 2$  for all  $i$  and when  $2 < r_m < k$  for any cycle.

Case I: If  $r_i \leq 2$  then  $(xy)^2$ , which is linear fractional transformation, fixes all the points of  $PL(F_q)$  since the only linear fractional transformation which fixes more than two vertices is the identity transformation [4] so  $(xy)^2$  is trivial, that is  $(xy)^2 = 1$ . But order of  $xy$  which is  $k$  for  $q > 3$  and  $k > 2$ , leads to a contradiction.

Case II If  $2 < r_m < k$  for any cycle then  $(xy)^{r_m}$  fixes  $r_m > 2$  points of the cycle since the only linear fractional transformation which fixes more than two vertices is the identity transformation [4] so  $(xy)^{r_m}$  is trivial that is  $(xy)^{r_m} = 1$ . But order of  $xy$  which is  $k$  and  $k > r_m$ , is a contradiction.

So the only possibility is that  $r_i = k$  for all  $i$ . This implies that  $\rho_{xy}$  has exactly two cycles of length  $k$ . That is, there are exactly two orbits of  $\langle xy \rangle$  of same size  $k = (q + 1)/2$ . Hence the result.

The converse follows from the definition of januarials.  $\square$

Since the two equal sized orbits in januarials are the result of action of the cyclic group  $\langle xy \rangle$  on  $PL(F_p)$  so the question arrises, does  $k = (q + 1)/2$  assure the existence of two equal sized orbits under the action of every cyclic group of order  $k$  on  $PL(F_q)$ ? We answer the question in the following theorem.

**Theorem 2** *The action of the cyclic subgroup  $C_k$ , of  $H_{\Lambda_i}$ , on  $PL(F_q)$  when  $k = (q + 1)/2$  give exactly two orbits of the same size  $k$ , for all  $q > 3$  where  $q = p^r$  and  $p \neq 2$ .*

*Proof:* The proof is the same as that of Theorem 1.  $\square$

### 3. Januarials from $PGL(2, q)$

Next we use the above condition to find januarials through a new method. We use the procedure described by F. Shaheen in [8]. Let  $X, Y$  denote the elements in  $GL(2, Z)$  which corresponds to the elements  $x, y$  in  $G$ . Then  $X, Y$  will satisfy the relations

$$X^2 = Y^\ell = \lambda I$$

for some scalar  $\lambda$ . We choose  $X, Y$  to be the matrices

$$\begin{bmatrix} a & ci \\ c & -a \end{bmatrix} \text{ and } \begin{bmatrix} e & fi \\ f & b - e \end{bmatrix}$$

respectively, where  $i \neq 0$ ,  $a, c, e, f, i$  belong to  $F_q$  and  $b \equiv s \pmod{p}$  for some  $s$  in  $F_q$ .

Let  $\Delta$  be the determinant of  $X$ . We assume that the determinant of  $Y$  is 1, so that we have

$$-(a^2 + ic^2) = \Delta \neq 0$$

and

$$1 + if^2 + e^2 - eb = 0.$$

Let  $r$  be the trace of  $XY$ , so that

$$r = a(2e - b) + 2icf.$$

Also

$$\det(XY) = \det X \det Y = \Delta.$$

We define a parameter  $\theta$  as  $r^2/\Delta$ . For a pair  $\bar{x}, \bar{y}$  in  $PGL(2, q)$ , satisfying the relations

$$\bar{x}^2 = \bar{y}^\ell = 1.$$

We denote by  $D(\theta, q, \ell)$ , where  $\theta$  belongs to  $F_q$ , the coset diagram corresponding to the action of  $G$  on  $PL(F_q)$ .

Now by the condition in Theorem 1, we need to consider only those coset diagrams in which every vertex is fixed by  $(xy)^k$ . We consider the case for  $(XY)^k = \lambda I$ , for some scalar  $\lambda$ .

By the following equation, given by Q. Mushtaq and F. Shaheen in [6];

$$\begin{aligned} (XY)^k &= \left\{ \binom{k-1}{0} r^{k-1} - \binom{k-2}{1} r^{k-3} \Delta + \binom{k-3}{2} r^{k-5} \Delta^2 - \dots \right\} XY \\ &\quad - \Delta \left\{ \binom{k-2}{0} r^{k-2} - \binom{k-3}{1} r^{k-4} \Delta + \binom{k-4}{2} r^{k-6} \Delta^2 - \dots \right\} I \end{aligned}$$

we get  $(XY)^k = \lambda I$ , for some scalar  $\lambda$ , if and only if

$$\binom{k-1}{0} r^{k-1} - \binom{k-2}{1} r^{k-3} \Delta + \binom{k-3}{2} r^{k-5} \Delta^2 - \dots = 0.$$

Substituting  $r^2 = \theta \Delta$  in the above equation and simplifying it we get an equation in  $\theta$  say  $g_k(\theta)$ . Let  $\theta_{i's}$  be the roots of  $g_k(\theta)$ . Now using these  $\theta_{i's}$  and by backward substitution we get  $X$  and  $Y$ , and ultimately  $x, y \in PGL(2, Z)$ .

**Theorem 3** *The action of quotients of  $H_{\Lambda_\ell}$ ,  $\Delta(2, \ell, k)$ , on  $PL(F_p)$  for  $k = (p+1)/2$ , yield januarials for all those values of  $\theta$  that are the roots of  $g_k(\theta)$  excluding those of  $g_{k/d}(\theta)$  for all  $d \mid k$ .*

*Proof:* The roots of  $g_{k/d}(\theta)$  which are also the roots of  $g_k(\theta)$  satisfy  $(xy)^{k/d} = 1$ . So there are  $2d$  orbits of  $xy$  (by the arguments as in the proof of Theorem1) and these values of  $\theta$  therefore do not yield januarials. All other values of  $\theta$  yield januarial because they give exactly two orbits of  $\langle xy \rangle$  of the same size.  $\square$

Now we find how many  $\ell$  - januarials exist and determine all the januarials that can be constructed from Hecke groups  $H_{\Lambda_\ell}$ .

#### 4. All $\ell$ -Januarials from $PGL(2, q)$

The following lemma is taken from [1].

**Lemma 4** *The number of conjugacy classes of elements in  $PGL(2, q)$  of order  $(q+1)/2$  is  $1/2 \phi((q+1)/2)$ . If  $z$  is any element of order  $(q+1)/2$  in  $PGL(2, q)$ , then every one of these conjugacy classes intersects the subgroup generated by  $z$  in  $\{z^i, z^{-i}\}$  for exactly one  $i$  coprime to  $(q+1)/2$ .*

The next result was first proved by Q. Mushtaq in [5] for  $y$  of order 3 and then generalized for  $y$  of order  $\ell$  by F. Shaheen in [8] as:

**Theorem 5** *Any element  $g$  whose order is not equal to 1, 2 or 6, of  $PGL(2, q)$  is the image of  $xy$  under some non-degenerate homomorphism from  $G$  to  $PGL(2, q)$ .*

From the above two results we can easily see that:

**Corollary 6** *For any prime  $p > 3$ , the number of distinct  $\ell$ -januarials constructible from  $PGL(2, p)$  is  $\frac{1}{2} \phi((p+1)/2)$ .*

We also show that the number  $\ell$ -januarials constructible from  $PGL(2, p)$  using the method described in Section 2 is  $\frac{1}{2}(\phi(k))$ .

**Theorem 7** *The number of roots,  $N$ , of  $g_k(\theta)$  excluding those of  $g_{k/d}(\theta)$ , for all  $d \in \mathbb{Z}$  such that  $d \mid k$ , is given by*

$$N = \frac{1}{2}(\phi(k)). \quad (3)$$

*Proof:* Let  $N_k$  denote the set of roots of  $g_k(\theta)$ . Since every positive integer can be uniquely written in the form of product of primes so let

$$k = p_1^{r_1} p_2^{r_2} \cdots p_s^{r_s}$$

where  $p_{i's}$  are primes such that  $p_1^{r_1} < p_2^{r_2} < \cdots < p_s^{r_s}$  and  $r_{i's}$  are integers.

$N_{k/p_1} \cup N_{k/p_2} \cup \cdots \cup N_{k/p_s}$  contain the roots of  $g_{k/d}(\theta)$ , for all  $d \in \mathbb{Z}$  such that  $d \mid k$ , as  $N_{k/d} \subset N_{k/p_i}$  for at least one  $i \in \{1, 2, \dots, s\}$ . So

$$N = |N_k| - |N_{k/p_1} \cup N_{k/p_2} \cup \cdots \cup N_{k/p_s}|.$$

Now by equation (1)

$$\begin{aligned} |N_{k/p_1} \cup N_{k/p_2} \cup \dots \cup N_{k/p_s}| &= \sum_{i=1}^s |N_{k/p_i}| - \sum_{1 \leq i < j \leq s} |N_{k/p_i} \cap N_{k/p_j}| \\ &\quad + \sum_{1 \leq i < j < l \leq s} |N_{k/p_i} \cap N_{k/p_j} \cap N_{k/p_l}| \\ &\quad - \dots + (-1)^{s+1} |N_{k/p_1} \cap N_{k/p_2} \cap \dots \cap N_{k/p_s}| \end{aligned}$$

so

$$\begin{aligned} N &= |N_k| - \sum_{i=1}^s |N_{k/p_i}| + \sum_{1 \leq i < j \leq s} |N_{k/p_i} \cap N_{k/p_j}| \\ &\quad - \sum_{1 \leq i < j < l \leq s} |N_{k/p_i} \cap N_{k/p_j} \cap N_{k/p_l}| + \dots + \\ &\quad (-1)^s |N_{k/p_1} \cap N_{k/p_2} \cap \dots \cap N_{k/p_s}|. \end{aligned} \quad (4)$$

Now

$$N_{k/p_i} \cap N_{k/p_j} = N_{k/p_i p_j}.$$

Similarly

$$N_{k/p_1} \cap N_{k/p_2} \cap \dots \cap N_{k/p_s} = N_{k/p_1 p_2 \dots p_s}$$

and equation (4) becomes

$$\begin{aligned} N &= |N_k| - \sum_{i=1}^s |N_{k/p_i}| + \sum_{1 \leq i < j \leq s} |N_{k/p_i p_j}| - \\ &\quad \sum_{1 \leq i < j < l \leq s} |N_{k/p_i p_j p_l}| + \dots + (-1)^s |N_{k/p_1 p_2 \dots p_s}|. \end{aligned} \quad (5)$$

The degree of  $g_k(\theta)$  and hence  $|N_k|$  is  $(k-1)/2$  when  $k$  is odd and  $k/2 - 1$  when  $k$  is even. So there are two cases.

Case I: When  $k$  is odd.

$$|N_k| = (k-1)/2,$$

$$|N_{k/p_i}| = (k/p_i - 1)/2,$$

$$|N_{k/p_i p_j}| = (k/p_i p_j - 1)/2$$

and

$$|N_{k/p_1} \cap N_{k/p_2} \cap \dots \cap N_{k/p_s}| = (k/p_1 p_2 \dots p_s - 1)/2.$$

Then equation (5) becomes

$$\begin{aligned} N &= (k-1)/2 - \sum_{i=1}^s (k/p_i - 1)/2 + \sum_{1 \leq i < j \leq s} (k/p_i p_j - 1)/2 - \sum_{1 \leq i < j < l \leq s} (k/p_i p_j p_l - 1)/2 \\ &\quad + \dots + (-1)^s (k/p_1 p_2 \dots p_s - 1)/2 \end{aligned}$$

implying that

$$N = \frac{1}{2} \left[ (k-1) - \sum_{i=1}^s (k/p_i - 1) + \sum_{1 \leq i < j \leq s} (k/p_i p_j - 1) - \sum_{1 \leq i < j < l \leq s} (k/p_i p_j p_l - 1) + \dots + (-1)^s (k/p_1 p_2 \dots p_s - 1) \right]. \quad (6)$$

Combinatorially it can be easily seen that

$$|\{k/p_i : 1 \leq i \leq s\}| = \binom{s}{1},$$

$$\left| \{k/p_i p_j : 1 \leq i < j \leq s\} \right| = \binom{s}{2}$$

and

$$\left| \{k/p_i p_j p_l : 1 \leq i < j < l \leq s\} \right| = \binom{s}{3}.$$

Putting these values in equation (6) we get

$$N = \frac{1}{2} \left[ \begin{array}{c} k - \binom{s}{0} - \sum_{i=1}^s (k/p_i) + \binom{s}{1} + \sum_{1 \leq i < j \leq s} (k/p_i p_j) - \binom{s}{2} \\ - \sum_{1 \leq i < j < l \leq s} (k/p_i p_j p_l) + \binom{s}{3} + \cdots + (-1)^s (k/p_1 p_2 \cdots p_s - \binom{s}{s}) \end{array} \right]$$

or

$$N = \frac{1}{2} \left[ \begin{array}{c} k - \sum_{i=1}^s k/p_i + \sum_{1 \leq i < j \leq s} (k/p_i p_j) - \sum_{1 \leq i < j < l \leq s} (k/p_i p_j p_l) \\ + \cdots + (-1)^s (k/p_1 p_2 \cdots p_s) \\ - \binom{s}{0} + \binom{s}{1} - \binom{s}{2} + \binom{s}{3} - \cdots + (-1)^{s+1} \binom{s}{s} \end{array} \right]. \quad (7)$$

Since

$$\binom{s}{0} - \binom{s}{1} + \binom{s}{2} - \binom{s}{3} + \cdots + (-1)^{s+1} \binom{s}{s} = 0 \quad (8)$$

therefore using (8) in equation (7) we get

$$N = \frac{1}{2} \left[ \begin{array}{c} k - \sum_{i=1}^s k/p_i + \sum_{1 \leq i < j \leq s} (k/p_i p_j) - \sum_{1 \leq i < j < l \leq s} (k/p_i p_j p_l) \\ + \cdots + (-1)^s (k/p_1 p_2 \cdots p_s) \end{array} \right].$$

But by equation (2)

$$\phi(k) = k - \sum_{i=1}^s \frac{k}{p_i} + \sum_{1 \leq i < j \leq s} \frac{k}{p_i p_j} - \cdots + (-1)^s \frac{k}{p_1 p_2 \cdots p_s} \quad (9)$$

imply that

$$N = \frac{1}{2} (\phi(k)).$$

Case II: When  $k$  is even, we have  $p_1 = 2$ . We have further two cases;

(a) When  $r_1 > 1$  then all the  $k/p_i, k/p_i p_j, \cdots, k/p_1 p_2 \cdots p_s$  are even so we have

$$|N_k| = (k/2 - 1),$$

$$|N_{k/p_i}| = (k/2 p_i - 1),$$

$$|N_{k/p_i p_j}| = (k/2 p_i p_j - 1)$$

and

$$|N_{k/p_1} \cap N_{k/p_2} \cap \cdots \cap N_{k/p_s}| = (k/2 p_1 p_2 \cdots p_s - 1)$$

(b) When  $r_1 = 1$  we have  $k/p_1, k/p_1 p_j, \cdots, k/p_1 p_2 \cdots p_s$  odd whereas all other are even. Simplifying equation (4) for both cases II (a) and (b) we get

$$N = \frac{1}{2} (\phi(k)).$$

So the result is true for all the cases.  $\square$

In Corollary 6 and Theorem 7 it can be seen that all the distinct  $\ell$ -januarials can be constructed from  $PGL(2, p)$  using the method described in Section 2. Hence this method of obtaining januarial is precise as compared to the method of obtaining januarial using associates which is described in [1].

## 5. Genus of the coset diagram for the action of $H_{\Lambda_\ell}$

It is pertinent to find a formula for calculating genus (of the coset diagram for the action of  $H_{\Lambda_\ell}$  on  $PL(F_p)$ ) using the fixed points in a coset diagram instead of edges and vertices, counting which is time consuming for higher primes and increase the possibility of errors.

Graham Higman's formula for the genus  $g$  of a coset diagram is

$$g = \frac{1}{2}\{2 - (v - e + f)\}$$

where  $g$  denotes the genus of the coset diagram,  $v$  is the number of all the orbits and cycles of  $y$ ,  $e$  is the number of nontrivial orbits or cycles of  $x$  and  $f$  is the number of all the orbits or cycles of  $xy$ .

**Theorem 8** Genus  $g$  of the coset diagram for the action of  $H_{\Lambda_\ell} = \langle x, y : x^2 = y^\ell = 1 \rangle$  on  $PL(F_p)$  is

$$g = 1 - \frac{1}{4k\ell}[\{2(k - \ell) - k\ell\}(p + 1) + k\ell\{2(\eta_y + \eta_{xy}) + \eta_x\} - 2(k\eta_y + \ell\eta_{xy})]$$

where  $k$  is the order of  $xy$  in the coset diagram,  $\eta_x, \eta_y$  and  $\eta_{xy}$  are the number of fixed points of  $x, y$  and  $xy$  respectively.

*Proof:* The genus  $g$  of a coset diagram can be calculated by using Graham Higman's formula, that is

$$g = \frac{1}{2}[2 - (v - e + f)]$$

where  $v$  is the number of orbits of  $y$ ,  $e$  is the number of non-trivial orbits of  $x$ ;  $f$  is the number of orbits of  $xy$ .

Let

$$v = \mu_y + \eta_y$$

where  $\mu_y$  is number of non trivial cycles of  $y$  and  $\eta_y$  be the number of trivial cycles or fixed points of  $y$ . As  $y^\ell = 1$  and  $y$  makes a cycle of length  $\ell$  so that values of  $\mu_y$  and  $\eta_y$  would depend on the quotient and the remainder respectively after dividing  $p + 1$  by  $\ell$ , the remainder would be  $\eta_y$ . Then

$$\ell\mu_y = p + 1 - \eta_y$$

that is

$$v = \frac{p + 1 - \eta_y}{\ell} + \eta_y$$

or

$$v = \frac{p + 1 + (\ell - 1)\eta_y}{\ell}.$$

Similarly,  $f$  is the number of all cycles of  $xy$ . We split  $f$  as

$$f = \mu_{xy} + \eta_{xy}$$

where  $\mu_{xy}$  is the number of non-trivial cycles of  $xy$  and  $\eta_{xy}$  is the number of trivial cycles. As  $(xy)^k = 1$  and  $xy$  makes a cycle of length  $k$ .

$$k\mu_{xy} = p + 1 - \eta_{xy}$$

or

$$f = \frac{p + 1 - \eta_{xy}}{k} + \eta_{xy}$$

or

$$f = \frac{p + 1 + (k - 1)\eta_{xy}}{k}.$$

Let  $\eta_x$  denote the number of fixed points of  $x$ . Then

$$p + 1 = 2e + \eta_x$$

or

$$e = \frac{p + 1 - \eta_x}{2}$$

putting values of  $v, f$ , and  $e$  in Graham Higman's formula to obtain the required result. That is,

$$g = \frac{1}{2} \left\{ 2 - \left( \frac{p + 1 + (\ell - 1)\eta_y}{\ell} - \frac{p + 1 - \eta_x}{2} + \frac{p + 1 + (k - 1)\eta_{xy}}{k} \right) \right\}$$

or

$$g = 1 - \frac{1}{4k\ell} [2(k + \ell) - k\ell](p + 1) + k\ell\{2(\eta_y + \eta_{xy}) + \eta_x\} - 2(k\eta_y + \ell\eta_{xy}).$$

□

For januarials we have  $k = \frac{p+1}{2}$  and  $\eta_{xy} = 0$ . Hence the following corollary is in order.

**Corollary 9** *Genus of a januarial is*

$$g = -\frac{p + 1 - \eta_y}{2\ell} + \frac{1}{4}(p + 1 - 2\eta_y - \eta_x)$$

where  $\eta_x$  and  $\eta_y$  are the fixed points of  $x$  and  $y$  respectively.

## 6. Illustration




Figure 1: Coset diagram  $D(7, 31, 4)$ .






Figure 2: The two orbits of  $\bar{x}\bar{y}$  highlighted with bold lines.

In the following we illustrate the above mentioned method for its better understanding. If we take  $p = 31$ , then by Theorem 1 we get  $k = 16$ , and

$$g_{16}(\theta) = 1x^7 - 14x^6 + 78x^5 - 220x^4 + 330x^3 - 252x^2 + 84x - 8.$$

The zeros of  $g_{16}(\theta)$  excluding those of  $g_8(\theta)$ ,  $g_4(\theta)$  and  $g_2(\theta)$  are 7, 16, 19 and 28. Now for a 4 – *januarial* if we consider  $\theta = 7$ , we get

$$\begin{aligned} x &: z \rightarrow \frac{3z + 30}{10z - 3} \\ y &: z \rightarrow \frac{42}{14z + 8}. \end{aligned}$$

The action of  $x$  and  $y$  on  $PL(F_{31})$  give permutations:

$$\begin{aligned} \bar{x} &= (0, 21)(1, 18)(2, 24)(3, 29)(4, 7)(5, 28)(6, 9)(8, 16)(10, 15) \\ &\quad (11, 20)(12, 26)(13, 23)(14, 27)(17, 30)(19, 25)(22, \infty) \\ \bar{y} &= (0, 13, 26, \infty)(1, 16, 9, 29)(2, 27, 3, 12)(4, 21, 18, 19) \\ &\quad (5, 22, 7, 8)(6, 20, 15, 11)(10, 25, 28, 17)(14, 23, 30, 24) \\ \bar{x}\bar{y} &= (0, 18, 16, 5, 17, 24, 27, 23, 26, 2, 14, 3, 1, 19, 28, 22) \\ &\quad (\infty, 7, 21, 13, 30, 10, 11, 15, 25, 4, 8, 9, 20, 6, 29, 12). \end{aligned}$$

Hence the coset diagram  $D(7, 31, 4)$  and the two orbits of  $\bar{x}\bar{y}$  highlighted with bold lines are depicted in the Figures 1 and 2.

According to Corollary 2, genus of this *januarial* is 4.

## References

- [1] M. Conder and T. Riley, Graham Higman's lectures on *januarials*, *Quart. J. Math.*, 65(1), 2014, 113-131.
- [2] M. R. Darafsheh and G. R. Razaeezadeh, Factorization of groups involving symmetric and alternating groups, *Int. J. Math. and Math. Science*, 27(3), 2000, 161-167.
- [3] L. Levai, G. Rosenberger and B. Sounignier, All finite generalized triangle groups, *Trans. AMS*, 347(9), 1995, 3625-3627.

- [4] Q. Mushtaq, Coset diagrams for the modular group, D. Phil. Thesis, Oxford University, 1983.
- [5] Q. Mushtaq, Parametrization of all Homomorphisms from  $PGL(2, \mathbb{Z})$  into  $PGL(2, q)$ , *Comm. in Algebra*, 20(4), 1992, 1023-1040.
- [6] Q. Mushtaq and F. Shaheen, On the subgroups of polyhedral groups, *Math. Japon.*, 35, 1990, 849-853.
- [7] Q. Mushtaq and F. Shaheen, Finite presentation of alternating groups, *Acta Math. Sinica*, 11(2), 1995, 221-224.
- [8] F. Shaheen, Coset diagrams for a two generator group, Ph.D. Thesis, Quaid-i-Azam University, Islamabad, 1991.
- [9] J. H van Lint and R. M Wilson, *A course in combinatorics*, Cambridge University Press, 1992.
- [10] A. T. White, *Graphs, groups and surfaces*, North Holland Pub. Co., Amster., 1973.