

Minimax L_2 -Separation Rate in Testing the Sobolev-type Regularity of a Function

Maurilio Gutzeit, OvGU Magdeburg

Abstract

In this paper we study the problem of testing if an L_2 -function f belonging to a certain l_2 -Sobolev-ball B_t of radius $R > 0$ with smoothness level $t > 0$ indeed exhibits a higher smoothness level $s > t$, that is, belongs to B_s . We assume that only a perturbed version of f is available, where the noise is governed by a standard Brownian motion scaled by $\frac{1}{\sqrt{n}}$. More precisely, considering a testing problem of the form

$$H_0 : f \in B_s \text{ vs. } H_1 : \inf_{h \in B_s} \|f - h\|_{L_2} > \rho$$

for some $\rho > 0$, we approach the task of identifying the smallest value for ρ , denoted ρ^* , enabling the existence of a test φ with small error probability in a minimax sense. By deriving lower and upper bounds on ρ^* , we expose its precise dependence on n :

$$\rho^* \sim n^{-\frac{t}{2t+1/2}}.$$

As a remarkable aspect of this composite-composite testing problem, it turns out that the rate does not depend on s and is equal to the rate in signal detection, i.e. the case of a simple null hypothesis.

1 Introduction

Let $n \in \mathbb{N}^* = \mathbb{N} \setminus \{0\}$, f a fixed unknown element of

$$L_2 := L_2([0, 1]) = \left\{ g : [0, 1] \rightarrow \mathbb{R} ; \int_0^1 g(x)^2 d\lambda(x) < \infty \right\}$$

and $(B(x))_{x \in [0, 1]}$ a standard Brownian motion. Suppose we observe the Gaussian process $(Y(x))_{x \in [0, 1]}$ determined by the stochastic differential equation

$$dY(x) = f(x)dx + \frac{1}{\sqrt{n}}dB(x), \quad x \in [0, 1]. \quad (1.1)$$

The resulting probability measure given f will be written \mathbb{P}_f .

Testing problem

We now fix $s > t > 0$ and $R, \rho > 0$. For any $r > 0$, we denote by $B_r(R)$ the l_2 -Sobolev-ball of radius R of functions on $[0, 1]$ with regularity at least r – see section 2 for a precise definition. Based on that, let

$$\tilde{B}_{s,t}(R, \rho) := \left\{ g \in B_t(R) ; \inf_{h \in B_s(R)} \|g - h\|_{L_2} > \rho \right\}.$$

Hence, if we interpret s and t as degrees of smoothness, $\tilde{B}_{s,t}(R, \rho)$ is the set of functions with smoothness level at least t which are separated from the class $B_s(R)$ with stronger smoothness s by ρ in L_2 -sense. Now, the testing problem of interest is

$$H_0 : f \in B_s(R) \text{ vs. } H_1 : f \in \tilde{B}_{s,t}(R, \rho). \quad (1.2)$$

More specifically, given $\eta \in (0, 1)$, we aim at finding the magnitude in terms of n of the smallest separation distance $\rho^*(\eta) = \rho^*(n, s, t, \eta)$ which enables the existence of a test φ of level η in a minimax sense, i.e. of

$$\rho^*(\eta) = \inf \left\{ \rho > 0 ; \exists \text{ test } \varphi : \sup_{f \in B_s(R)} \mathbb{P}_f(\varphi = 1) + \sup_{f \in \tilde{B}_{s,t}(R, \rho)} \mathbb{P}_f(\varphi = 0) \leq \eta \right\}. \quad (1.3)$$

In the sequel we will omit the index f in \mathbb{P}_f or write a different index depending on the context.

Related questions and literature

There are in essence two lines of work with questions or ideas closely related to the present paper.

Firstly, considering the simpler null hypothesis $H_0 : f \equiv 0$ puts us in the so-called signal-detection setting which has already been studied, see for instance the series of seminal papers [10] as well as [11] or [8] for a more recent treatment. In that context, the order of ρ^* with respect to n is shown to be

$$n^{-\frac{t}{2t+1/2}}.$$

Secondly, another closely related task is the construction of (adaptive and honest) confidence regions for f . In [3], the authors study such sets in terms of L_2 -separation, but rather than the observation $(Y(x))_{x \in [0,1]}$ they use a Gaussian sequence model. However, due to the equivalence of these models in the sense of Le Cam (see [14]), it is possible to derive from their arguments that for our problem (1),

$$n^{-\frac{t}{2t+1/2}} \lesssim \rho^*(\eta) \lesssim \max \left(n^{-\frac{s}{2s+1}}, n^{-\frac{t}{2t+1/2}} \right). \quad (1.4)$$

While the resulting gap in the case $s < 2t$ is not essential in the confidence region setting (see also [4] and [12]), it is quite important from a testing perspective as it raises the question how the complexity of the null hypothesis influences the separation rate.

Now, the article [6] is by far the closest previous work to the present paper. Indeed,

the author studies the same problem with another choice of Sobolev-ball, namely the (r, ∞) -Sobolev-balls $B_{r, \infty}(R)$. In this context, $\rho^*(\eta)$ is proved to be of magnitude

$$n^{-\frac{t}{2t+1/2}}.$$

Note that this quantity is equal to the rate in the signal-detection case and hence in particular does not depend on s . This makes the issue of the gap in (1) even more interesting and, from a technical perspective, it is rather striking given that moving from a simple to the composite null hypothesis is a significant step. On top of that, there are settings where the separation rate strongly depends on the shape of the null hypothesis, see e.g. [2] and [13] or also [5].

To the best of our knowledge, the case of [6] is the only one for which the minimax L_2 -separation rate is known and our main contribution is to extend that result to the $(r, 2)$ -Sobolev-space. While our lower bound (Theorem 2 in section 3) is essentially a corollary of the corresponding result [6, Theorem 3.2], the upper bound (Theorem 1 in section 3) cannot be established through a simple application of [6, Theorem 3.1]. As $B_r(R) \subseteq B_{r, \infty}(R)$, this might be surprising at first sight: Indeed, the test from [6] would perform well in the present setting in terms of type-I-error. However, ensuring sufficient power is significantly more difficult when considering l_2 -Sobolev-balls.

2 Setting

In this section, we describe how the relevant Sobolev balls and the observed Gaussian process will be represented throughout the paper.

Wavelet transform and associated Sobolev ball

Throughout the paper, we make heavy use of a wavelet decomposition of f . As is well-known, we can define a scalar product and associated norm on L_2 by

$$\langle g, h \rangle := \int_0^1 g(x)h(x) \, d\lambda(x) \quad \text{with} \quad \|g\|_{L_2} := \sqrt{\langle g, g \rangle}, \quad g, h \in L_2.$$

There are many orthogonal wavelet bases of L_2 with respect to $\langle \cdot, \cdot \rangle$. A suitable choice for our purposes is a basis developed in [7] that can be written as

$$\mathcal{W} = \bigcup_{j=2}^{\infty} \{\psi_{j,k} : k \in \{1, 2, \dots, 2^j\}\},$$

i.e. it is tailored such that there are exactly 2^j basis functions at resolution $j \geq 2$. Clearly, the coefficients of $g \in L_2$ with respect to \mathcal{W} are given by

$$\langle g, \psi_{j,k} \rangle = \int_0^1 g(x)\psi_{j,k}(x) \, dx, \quad j \geq 2, k \in \{1, 2, \dots, 2^j\}.$$

and yield the representation

$$g = \sum_{j=2}^{\infty} \sum_{k=1}^{2^j} \langle g, \psi_{j,k} \rangle \psi_{j,k}.$$

Let $r > 0$. By virtue of isometry properties discussed for instance in [15] and [9], we may now define a functional $(r, 2)$ -Sobolev-ball of radius R solely through the wavelet coefficients of its elements:

$$B_r(R) := \left\{ g \in L_2 ; \sum_{j=2}^{\infty} 4^{jr} \sum_{k=1}^{2^j} \langle g, \psi_{j,k} \rangle^2 \leq R^2 \right\} \quad (2.1)$$

with associated $(r, 2)$ -Sobolev-norm

$$\|g\|_{\mathcal{B}_r} := \sqrt{\sum_{j=2}^{\infty} 4^{jr} \sum_{k=1}^{2^j} \langle g, \psi_{j,k} \rangle^2}, \quad g \in L_2$$

or also, as mentioned at the end of the previous section,

$$B_{r,\infty}(R) := \left\{ g \in L_2 ; \sup_{j \geq 2} 4^{jr} \sum_{k=1}^{2^j} \langle g, \psi_{j,k} \rangle^2 \leq R^2 \right\}.$$

Discrete observation scheme based on the wavelet basis

Let

$$\mathcal{I} = \{(j, k) \in \mathbb{N}^2 \mid j \geq 2, k \leq 2^j\}.$$

Motivated by (2), for each $(j, k) \in \mathcal{I}$ we consider

$$a_{j,k} := \langle f, \psi_{j,k} \rangle$$

so that

$$f = \sum_{j=2}^{\infty} \sum_{k=1}^{2^j} a_{j,k} \psi_{j,k}.$$

The natural corresponding estimators read

$$\hat{a}_{j,k} := \langle dY, \psi_{j,k} \rangle, \quad \hat{f} = \sum_{j=2}^{\infty} \sum_{k=1}^{2^j} \hat{a}_{j,k} \psi_{j,k}. \quad (2.2)$$

By construction and due to the orthonormality of \mathcal{W} , we know that the family $(\hat{a}_{i,j})_{(j,k) \in \mathcal{I}}$ is independent with

$$\hat{a}_{j,k} \sim \mathcal{N}\left(a_{j,k}, \frac{1}{n}\right).$$

Clearly, observing this family is equivalent to observing the original process $(Y(x))_{x \in [0,1]}$.

3 Main results

In this section, we state and discuss our main results, that is upper and lower bounds on $\rho^*(\eta)$. We also provide a high-level description of the strategy and ideas included in the upper bound proof, which is our main contribution.

3.1 Upper Bound

The test

Note that \widehat{f} from (2) is not a useful estimator as it exhibits infinite variance. Therefore, we need to carefully impose a restriction of the form $j \leq J$ for some fixed $J \in \mathbb{N}$, $J \geq 2$. More specifically, section 5 is primarily concerned with obtaining an upper bound on $\rho_J^*(\eta)$ for the reduced, finite-dimensional problem

$$H'_0 : \underbrace{\left\| \sum_{j=2}^J \sum_{k=1}^{2^j} a_{j,k} \psi_{j,k} \right\|_{\mathcal{B}_s}}_{:=S_J} \leq R \quad \text{vs.} \quad H'_1 : \inf_{h \in B_s(R)} \left\| \sum_{j=2}^J \sum_{k=1}^{2^j} a_{j,k} \psi_{j,k} - h \right\|_{L_2} > \rho_J,$$

where ρ_J and $\rho_J^*(\eta)$ are analogous in definition and relation to their counterparts in (1) and (1).

Finding a sufficient separation distance $\rho_J \geq \rho_J^*(\eta)$ here is the central and most involved part of the paper. Indeed, it turns out that a test based on estimating S_J^2 only cannot perform well enough under the targeted separation distance of order $n^{-t/(2t+1/2)}$. Rather than that, our test estimates $S_2^2, S_3^2, \dots, S_J^2$ through test statistics T_2, T_3, \dots, T_J and, under the alternative H'_1 , relies on the smallest level j^* such that $S_{j^*}^2$ significantly exceeds R^2 (Lemmas 3 and 5 below). It essentially takes the form

$$\varphi = 1 - \prod_{j=2}^J \mathbf{1}_{\{T_j \leq \tau_j\}}$$

of a multi-level test with suitable thresholds τ_j (equation (5.4) below).

Finally, J must be chosen such that an appropriate trade-off between $\rho_J^*(\eta)$ and the error incurred by ignoring the resolutions beyond J is reached.

In terms of technical ingredients, these considerations are remarkable in that they solely rely on elementary computations based on the Sobolev-balls' geometry and classical properties of the χ^2 -distribution. The explicit result reads as follows:

Theorem 1. Let $\eta \in (0, 1)$. Whenever

$$\rho \geq \left(\frac{1346}{\sqrt{\eta}} + \frac{R}{1-2^{-t}} \right) n^{-\frac{t}{2t+1/2}},$$

there is a test φ such that

$$\sup_{f \in B_s(R)} \mathbb{P}_f(\varphi = 1) + \sup_{f \in \widetilde{B}_{s,t}(R,\rho)} \mathbb{P}_f(\varphi = 0) \leq \eta.$$

Hence,

$$\rho^*(\eta) \leq \left(\frac{1346}{\sqrt{\eta}} + \frac{R}{1-2^{-t}} \right) n^{-\frac{t}{2t+1/2}}, \quad \text{i.e.} \quad \rho^*(\eta) \lesssim n^{-\frac{t}{2t+1/2}}.$$

◁

3.2 Lower Bound

Using the same choice for J as indicated above, a lower bound on $\rho^*(\eta)$ of the same order can be derived through studying the statistical distance between specific distributions agreeing with H_0 and H_1 respectively.

Theorem 2. Let $\eta \in (0, 1)$. There are $C_\eta > 0$ and $N_\eta \in \mathbb{N}$ such that whenever $n \geq N_\eta$ and

$$\rho < C_\eta n^{-\frac{t}{2t+1/2}},$$

for any test φ it holds that

$$\sup_{f \in B_s(R)} \mathbb{P}_f(\varphi = 1) + \sup_{f \in \tilde{B}_{s,t}(R,\rho)} \mathbb{P}_f(\varphi = 0) > \eta.$$

Hence,

$$\rho^*(\eta) \geq C_\eta n^{-\frac{t}{2t+1/2}}, \quad \text{i.e. } \rho^*(\eta) \gtrsim n^{-\frac{t}{2t+1/2}}.$$

In particular, one may choose

$$C_\eta := \frac{R}{2} \min \left\{ 1, \frac{1}{R \cdot 2^{t+1/2}}, \frac{\sqrt{\ln(1 + 4(1 - \eta)^2)}}{2^t 16R} \right\}, \quad N_\eta := \left\lceil \left(R \frac{2^{s-t}}{C_\eta} \right)^{\frac{2t+1/2}{s-t}} \right\rceil.$$

◁

Note that, as mentioned in the introduction, Theorems 1 and 2 in conjunction reveal the minimax separation rate to be of order

$$\rho^*(\eta) \sim n^{-\frac{t}{2t+1/2}},$$

which does not depend on the size of the null hypothesis and is equal to the signal detection rate. Indeed, in order to obtain the lower bound of Theorem 2, the fact that H_0 is a composite hypothesis need not be used.

4 Alternative settings

Before presenting the proofs of our main results, we briefly discuss their possible application in two alternative settings which might also be of interest, see also [6, Section 3.3] and references therein.

Heteroscedastic noise

As a generalisation of (1), consider the model

$$dY(x) = f(x)dx + \frac{\sigma(x)}{\sqrt{n}}dB(x), \quad x \in [0, 1], \quad (4.1)$$

where $\sigma \in L_2$ is unknown. The proof of Theorem 1 relies heavily on unbiased estimators of $a_{j,k}^2$, $(j, k) \in \mathcal{I}$, and hence on knowledge of the noise coefficient, so that in this generalised version we cannot directly apply our result. However, there is a relatively

simple solution under certain conditions: Suppose we have access to two independent realisations $(Y^{(1)}(x))_{x \in [0,1]}$ and $(Y^{(2)}(x))_{x \in [0,1]}$ with noise coefficient, say, $\frac{\sigma(x)}{\sqrt{n/2}}$. Then we can still consider the estimates

$$\widehat{a}_{j,k}^{(i)} = \langle dY^{(i)}, \psi_{j,k} \rangle \sim \mathcal{N} \left(a_{j,k}, 2 \frac{\|\sigma \cdot \psi_{j,k}\|_{L_2}^2}{n} \right), \quad i \in \{1, 2\}$$

and define a new unbiased estimator for $a_{j,k}^2$ based on the simple observation

$$\mathbb{E}[\widehat{a}_{j,k}^{(1)} \cdot \widehat{a}_{j,k}^{(2)}] = a_{j,k}^2.$$

If in addition we know an upper bound on $\|\sigma\|_{L_2}$, it turns out that we can state an analogous concentration result as the one for the homoscedastic model (see Lemma 4 below) and obtain essentially the same result.

Regression

Another possible observation scheme for testing the smoothness of f would be collecting n iid samples $(X_i, Y_i)_{i \in \{1, 2, \dots, n\}}$ according to the model

$$Y = f(X) + \frac{\sigma(X)}{\sqrt{n}} \epsilon$$

for $\epsilon \sim \mathcal{N}(0, 1)$ and X uniformly distributed on $[0, 1]$. This situation is particularly interesting since, as mentioned above, it is equivalent to (4) in the sense of Le Cam ([14]) We could then arrive at the same situation as in the previous setting by considering

$$\widehat{a}_{j,k}^{(1)} = \frac{2}{n} \sum_{i=1}^{n/2} Y_i \psi_{j,k}(X_i), \quad \widehat{a}_{j,k}^{(2)} = \frac{2}{n} \sum_{i=n/2+1}^n Y_i \psi_{j,k}(X_i).$$

Note that if X is not uniformly distributed, $\mathbb{E}[\widehat{a}_{j,k}^{(i)}] = a_{j,k}$ is generally not true and it becomes crucial to guarantee a certain spread of the design points $(X_i)_{i \in \{1, 2, \dots, n\}}$ over $[0, 1]$.

5 Proof of Theorem 1

5.1 General preparations

Reduction of the range of resolutions

Let us make this more clear at this point already: For $j_1, j_2 \in \mathbb{N} \cup \{\infty\}$ with $2 \leq j_1 \leq j_2$ and $g \in L_2$, define the projections

$$P_{j_1}^{j_2} g = \sum_{j=j_1}^{j_2} \sum_{k=1}^{2^j} \langle g, \psi_{j,k} \rangle \psi_{j,k}, \quad P_{j_1} := P_{j_1}^{j_1}.$$

Now observe that since $f \in B_t(R)$, for each $j \in \mathbb{N}$, $j \geq 2$, we have

$$\|P_j f\|_{L_2} = \frac{\|P_j f\|_{\mathcal{B}_t}}{2^{jt}} \leq \frac{R}{2^{jt}}$$

and hence

$$\sum_{j=J+1}^{\infty} \|P_j f\|_{L_2} \leq R \sum_{j=J+1}^{\infty} (2^{-t})^j = 2^{-tJ} \frac{2^{-t}R}{1-2^{-t}}.$$

Using the triangle inequality, this tells us that under the alternative hypothesis

$$\begin{aligned} \rho &< \inf_{h \in B_s(R)} \|f - h\|_{L_2} \\ &\leq \inf_{h \in B_s(R)} \|P_2^J f - h\|_{L_2} + \|P_{J+1}^{\infty} f\|_{L_2} \\ &\leq \inf_{h \in B_s(R)} \|P_2^J f - h\|_{L_2} + \sum_{j=J+1}^{\infty} \|P_j f\|_{L_2}, \\ &\leq \inf_{h \in B_s(R)} \|P_2^J f - h\|_{L_2} + 2^{-tJ} \frac{2^{-t}R}{1-2^{-t}}. \end{aligned}$$

Accordingly, under H_1 we consider the assumption

$$\rho - 2^{-tJ} \frac{2^{-t}R}{1-2^{-t}} =: \rho_J < \inf_{h \in B_s(R)} \|P_2^J f - h\|_{L_2}$$

and firstly solve (1) for ρ_J in terms of the reduced range $j \in \{2, 3, \dots, J\}$, that is, subsequently, we will primarily study the testing problem

$$H'_0 : \|P_2^J f\|_{\mathcal{B}_s} \leq R \quad \text{vs.} \quad H'_1 : \inf_{h \in B_s(R)} \|P_2^J f - h\|_{L_2} > \rho_J.$$

Finally, ρ will be determined by choosing J such that a reasonable trade-off between the two summands,

$$\rho = \rho_J + 2^{-tJ} \frac{2^{-t}R}{1-2^{-t}}, \tag{5.1}$$

is realised.

Now, more specifically, with $a = 1346$, for $j^* \in \{2, 3, \dots, J\} =: \mathcal{J}$, let

$$\rho_1 := 0; \quad \rho_{j^*} = a \frac{2^{(3j^*+2J)/20}}{\sqrt{n}}.$$

Under the assumption H'_1 it will be technically useful to detect the level $j^* \in \mathcal{J}$ at which $\inf_{h \in B_s(R)} \|P_2^{j^*} f - h\|_{\mathcal{B}_s}$ firstly exceeds ρ_{j^*} in the sense of Lemma 3 below. That leads to a multiple test across the set \mathcal{J} finally given in (5.4).

Decomposition of H'_1

Lemma 3. Under the assumption H'_1 , we have

$$\exists j^* \in \mathcal{J} : \begin{cases} \inf_{h \in B_s(R)} \|P_2^{j^*-1} f - h\|_{L_2} \leq \rho_{j^*-1}, \\ \inf_{h \in B_s(R)} \|P_2^{j^*} f - h\|_{L_2} > \rho_{j^*}. \end{cases} \tag{5.2}$$

◁

Proof. By contradiction: Assume that (3) is false, i.e.

$$\forall j^* \in \mathcal{J} : \underbrace{\inf_{h \in B_s(R)} \|P_2^{j^*-1} f - h\|_{L_2} > \rho_{j^*-1}}_{E_{j^*}} \vee \underbrace{\inf_{h \in B_s(R)} \|P_2^{j^*} f - h\|_{L_2} \leq \rho_{j^*}}_{F_{j^*}}.$$

Then clearly F_J is false, so that E_J is true. Equivalently, F_{J-1} is false and in turn E_{J-1} must be true. Continued application of this argument leads to the contradiction

$$\inf_{h \in B_s(R)} \|P_2^1 f - h\|_{L_2} = 0 > \rho_1.$$

□

Concentration of $\|P_2^{j^*} \widehat{f}\|_{\mathcal{B}_s}^2$

Lemma 4. Let $j^* \in \mathcal{J}$. Then, with

$$A_{j^*} := \frac{1}{n} \sum_{j=2}^{j^*} (2 \cdot 4^s)^j, \quad B_{j^*} := \frac{2}{n^2} \sum_{j=2}^{j^*} (2 \cdot 4^{2s})^j, \quad V_{j^*} = \frac{4}{n} \sum_{j=2}^{j^*} 4^{2js} \|P_j f\|_{L_2}^2,$$

it holds that

$$\forall \delta \in (0, 1) : \mathbb{P} \left(\left| \|P_2^{j^*} \widehat{f}\|_{\mathcal{B}_s}^2 - A_{j^*} - \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \right| \geq \sqrt{\frac{1}{\delta} (B_{j^*} + V_{j^*})} \right) \leq \delta. \quad (5.3)$$

◁

Proof. For $j \in \mathcal{J}$, let

$$\lambda_j := n \sum_{k=1}^{2^j} a_{j,k}^2 = n \|P_j f\|_{L_2}^2.$$

Then, by construction, we know that

$$n \|P_j \widehat{f}\|_{L_2}^2 = \sum_{k=1}^{2^j} (\sqrt{n} \widehat{a}_{j,k})^2 \sim \chi^2(2^j, \lambda_j),$$

i.e. a χ^2 -distribution with 2^j degrees of freedom and non-centrality parameter λ_j . Classical properties of this distribution now tell us

$$\mathbb{E} \left[\|P_j \widehat{f}\|_{L_2}^2 \right] = \frac{2^j}{n} + \|P_j f\|_{L_2}^2; \quad \text{Var} \left[\|P_j \widehat{f}\|_{L_2}^2 \right] = 2 \left(\frac{2^j}{n^2} + \frac{2}{n} \|P_j f\|_{L_2}^2 \right). \quad (5.4)$$

Since

$$\|P_2^{j^*} f\|_{\mathcal{B}_s}^2 = \sum_{j=2}^{j^*} 4^{js} \|P_j \widehat{f}\|_{L_2}^2,$$

independence in conjunction with (5.1) yields

$$\begin{aligned}
\mathbb{E}[\|P_2^{j^*} \widehat{f}\|_{\mathcal{B}_s}^2] &= \sum_{j=2}^{j^*} 4^{js} \left(\frac{2^j}{n} + \|P_j f\|_{L_2}^2 \right) \\
&= \frac{1}{n} \sum_{j=2}^{j^*} (2 \cdot 4^s)^j + \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \\
&= A_{j^*} + \|P_2^{j^*} f\|_{\mathcal{B}_s}^2; \\
\text{Var}[\|P_2^{j^*} \widehat{f}\|_{\mathcal{B}_s}^2] &= \sum_{j=2}^{j^*} 4^{2js} \left(2 \left(\frac{2^j}{n^2} + \frac{2}{n} \|P_j f\|_{L_2}^2 \right) \right) \\
&= \frac{2}{n^2} \sum_{j=2}^{j^*} (2 \cdot 4^{2s})^j + \frac{4}{n} \sum_{j=2}^{j^*} 4^{2js} \|P_j f\|_{L_2}^2 \\
&= B_{j^*} + V_{j^*}.
\end{aligned}$$

We obtain the desired result directly through Chebyshev's inequality: For $\epsilon > 0$,

$$\mathbb{P} \left(\left| \|P_2^{j^*} \widehat{f}\|_{\mathcal{B}_s}^2 - A_{j^*} - \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \right| \geq \epsilon \right) \leq \frac{B_{j^*} + V_{j^*}}{\epsilon^2}$$

and hence the claim. □

More specifically, observe that

$$\begin{aligned}
B_{j^*} &= \frac{2}{n^2} \sum_{j=2}^{j^*} (2 \cdot 4^{2s})^j \\
&= \frac{2}{n^2} (2 \cdot 4^{2s})^2 \frac{(2 \cdot 4^{2s})^{j^*-1} - 1}{2 \cdot 4^{2s} - 1} \\
&\leq \frac{2}{n^2} \frac{2 \cdot 4^{2s}}{2 \cdot 4^{2s} - 1} (2 \cdot 4^{2s})^{j^*} \\
&\leq \frac{4}{n^2} (2 \cdot 4^{2s})^{j^*},
\end{aligned}$$

(where we use that for $x \geq 2$, $\frac{x}{x-1} \leq 2$) and hence for $\delta \in (0, 1)$

$$\sqrt{\frac{B_{j^*}}{\delta}} \leq \frac{2}{\sqrt{\delta}} 4^{j^* s} \frac{2^{j^*/2}}{n}.$$

Furthermore,

$$\sqrt{\frac{V_{j^*}}{\delta}} = \frac{2}{\sqrt{\delta}} \cdot \frac{1}{\sqrt{n}} \sqrt{\sum_{j=2}^{j^*} 4^{2js} \|P_j f\|_{L_2}^2} \leq \frac{2}{\sqrt{\delta}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} \max_{2 \leq j \leq j^*} 2^{js} \|P_j f\|_{\mathcal{B}_s}.$$

The maximum in the latter computation will play an important role in the sequel. From now on we use the abbreviation

$$M_{j^*} := \max_{2 \leq j \leq j^*} 2^{js} \|P_j f\|_{\mathcal{B}_s}. \tag{5.5}$$

Plugging these bounds in (4) leads to

$$\forall \delta \in (0, 1) : \mathbb{P} \left(\left| \|P_2^{j^*} \widehat{f}\|_{\mathcal{B}_s}^2 - A_{j^*} - \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \right| \geq \frac{2}{\sqrt{\delta}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} M_{j^*} + \frac{2}{\sqrt{\delta}} 4^{j^* s} \frac{2^{j^*/2}}{n} \right) \leq \delta. \quad (5.6)$$

5.2 Preliminary Bounds on $\|P_2^{j^*} f\|_{\mathcal{B}_s}$

As a next step towards controlling the type-I and type-II errors of our test, we study $\|P_2^{j^*} f\|_{\mathcal{B}_s}$ more closely.

On the one hand, under H'_0 , for any $j^* \in \mathcal{J}$ we clearly have $\|P_2^{j^*} f\|_{\mathcal{B}_s} \leq R$.

On the other hand, under H'_1 , we require a lower bound on $\|P_2^{j^*} f\|_{\mathcal{B}_s}$. The following bound is preliminary in the sense that it requires the knowledge of an index $j^* \in \mathcal{J}$ with the property from (3) and the corresponding M_{j^*} . The generalisation will be considered in sections 5.3 and 5.4.

Lemma 5. Let $j^* \in \mathcal{J}$ be an index with the property

$$\inf_{h \in B_s(R)} \|P_2^{j^* - 1} f - h\|_{L_2} \leq \rho_{j^* - 1}, \quad \inf_{h \in B_s(R)} \|P_2^{j^*} f - h\|_{L_2} > \rho_{j^*}. \quad (5.7)$$

Then the following assertion holds for $A = 11$:

$$\|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \geq R^2 + \frac{1}{2 \cdot A^2} \rho_{j^*} M_{j^*} + \frac{1}{2 \cdot A^2} 4^{j^* s} \rho_{j^*}^2. \quad (5.8)$$

◁

Proof. Before giving the main arguments, we need a technical preparation and a general (i.e. only depending on j^*) lower bound on $\|P_2^{j^*} f\|_{\mathcal{B}_s}$:

1. Proxy minimisation of $\inf_{h \in B_s(R)} \|P_2^{j^*} f - h\|_{L_2}$

For $\tilde{j} \in \mathcal{J}$, write $P_{j \neq \tilde{j}} := P_2^{j^*} - P_{\tilde{j}}$. In the case that $\|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s} \leq R$, we can introduce the function \tilde{h} through the wavelet coefficients

$$\begin{aligned} b_{j,k} &:= a_{j,k} && \text{for } (j,k) \in \mathcal{I}, j \neq \tilde{j}, \\ b_{\tilde{j},k} &:= a_{\tilde{j},k} \cdot \frac{\sqrt{R^2 - \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2}}{\|P_{\tilde{j}} f\|_{\mathcal{B}_s}}, && \text{for } k \in \{1, 2, \dots, 2^{\tilde{j}}\}. \end{aligned}$$

Then $\tilde{h} \in B_s(R)$ holds since

$$\begin{aligned} \|\tilde{h}\|_{\mathcal{B}_s}^2 &= \sum_{j=2}^{j^*} 4^{j s} \sum_{k=1}^{2^j} b_{j,k}^2 \\ &= \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2 + \left(\frac{\sqrt{R^2 - \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2}}{\|P_{\tilde{j}} f\|_{\mathcal{B}_s}} \right)^2 \|P_{\tilde{j}} f\|_{\mathcal{B}_s}^2 \\ &= R^2. \end{aligned}$$

Hence, by assumption

$$\rho_{j^*}^2 < \inf_{h \in B_s(R)} \|P_2^{j^*} f - h\|_{L_2}^2 \leq \|P_2^{j^*} f - \tilde{h}\|_{L_2}^2 =: d^2,$$

where

$$\begin{aligned} d^2 &= \|P_2^{j^*} f - \tilde{h}\|_{L_2}^2 \\ &= \sum_{k=1}^{2^{\tilde{j}}} \left(1 - \frac{\sqrt{R^2 - \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2}}{\|P_{\tilde{j}} f\|_{\mathcal{B}_s}} \right)^2 a_{j,k}^2 \\ &= \left(1 - \frac{\sqrt{R^2 - \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2}}{\|P_{\tilde{j}} f\|_{\mathcal{B}_s}} \right)^2 \frac{\|P_{\tilde{j}} f\|_{\mathcal{B}_s}^2}{4^{\tilde{j}s}} \\ &= \left(\|P_{\tilde{j}} f\|_{\mathcal{B}_s} - \sqrt{R^2 - \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2} \right)^2 4^{-\tilde{j}s}. \end{aligned}$$

This tells us that if $\|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s} \leq R$,

$$\|P_{\tilde{j}} f\|_{\mathcal{B}_s} = 2^{\tilde{j}s} d + \sqrt{R^2 - \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2} \geq 2^{\tilde{j}s} d, \quad (5.9)$$

$$\|P_2^{j^*} f\|_{\mathcal{B}_s}^2 = R^2 + 2 \cdot 2^{\tilde{j}s} d \sqrt{R^2 - \|P_{j \neq \tilde{j}} f\|_{\mathcal{B}_s}^2} + 4^{\tilde{j}s} d^2. \quad (5.10)$$

2. Bound in terms of $4^{j^*s} \rho_{j^*}^2$

If $\|P_2^{j^*-1} f\|_{\mathcal{B}_s} \leq R$, we can use (1) with $\tilde{j} = j^*$ and $d \geq \rho_{j^*} \geq 0$ and obtain

$$\|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \geq R^2 + 4^{j^*s} \rho_{j^*}^2.$$

If $\|P_2^{j^*-1} f\|_{\mathcal{B}_s} > R$, observe that by the triangle inequality

$$\inf_{h \in B_s(R)} \|P_2^{j^*} f - h\|_{L_2} \leq \inf_{h \in B_s(R)} \|P_2^{j^*-1} f - h\|_{L_2} + \|P_{j^*} f\|_{L_2} \leq \rho_{j^*-1} + \|P_{j^*} f\|_{L_2}$$

and since

$$\rho_{j^*} - \rho_{j^*-1} \geq a \frac{2^{(3j^*+2J)/20}}{\sqrt{n}} (1 - 2^{-3/20}) \geq \frac{1}{11} \rho_{j^*} = \frac{1}{A} \rho_{j^*},$$

we obtain

$$\begin{aligned} \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 &= \|P_2^{j^*-1} f\|_{\mathcal{B}_s}^2 + \|P_{j^*} f\|_{\mathcal{B}_s}^2 \\ &\geq R^2 + 4^{j^*s} (\rho_{j^*} - \rho_{j^*-1})^2 \\ &\geq R^2 + \frac{1}{A^2} 4^{j^*s} \rho_{j^*}^2. \end{aligned}$$

So, in any case,

$$\|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \geq R^2 + \frac{1}{A^2} 4^{j^*s} \rho_{j^*}^2. \quad (5.11)$$

3. Main arguments

We are now ready to prove (5) effectively. To that end, fix an index

$$\bar{j} \in \operatorname{argmax}_{j \in \mathcal{J}} 2^{j^s} \|P_j f\|_{\mathcal{B}_s}.$$

Case 1: $\|P_{j \neq \bar{j}} f\|_{\mathcal{B}_s} \leq R$

In that case, we can use (1) and (1) with $\tilde{j} = \bar{j}$ in combination with (2) and obtain

$$\begin{aligned} \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 &\geq R^2 + 2^{\bar{j}^s} d \sqrt{R^2 - \|P_{j \neq \bar{j}} f\|_{\mathcal{B}_s}^2} + \frac{1}{2} 4^{\bar{j}^s} d^2 + \frac{1}{2 \cdot A^2} 4^{j^* s} d^2 \\ &= R^2 + 2^{\bar{j}^s} d \left(\|P_{\bar{j}} f\|_{\mathcal{B}_s} - 2^{\bar{j}^s} d \right) + \frac{1}{2} 4^{\bar{j}^s} d^2 + \frac{1}{2 \cdot A^2} 4^{j^* s} d^2 \\ &= R^2 + d \cdot 2^{\bar{j}^s} \left(\|P_{\bar{j}} f\|_{\mathcal{B}_s} - \frac{1}{2} 2^{\bar{j}^s} d \right) + \frac{1}{2 \cdot A^2} 4^{j^* s} d^2 \\ &\geq R^2 + \rho_{j^*} 2^{\bar{j}^s} \left(\|P_{\bar{j}} f\|_{\mathcal{B}_s} - \frac{1}{2} 2^{\bar{j}^s} d \right) + \frac{1}{2 \cdot A^2} 4^{j^* s} \rho_{j^*}^2 \\ &\geq R^2 + \frac{1}{2} \rho_{j^*} M_{j^*} + \frac{1}{2 \cdot A^2} 4^{j^* s} \rho_{j^*}^2, \end{aligned}$$

remembering (5.1).

Case 2: $\|P_{j \neq \bar{j}} f\|_{\mathcal{B}_s} > R$

That case can be handled quickly by considering two subcases:

Subcase 1: $4^{j^* s} \rho_{j^*}^2 \geq \rho_{j^*} M_{j^*}$

Observe that with (2)

$$\|P_2^{j^*} f\|_{\mathcal{B}_s}^2 \geq R^2 + \frac{1}{2 \cdot A^2} 4^{j^* s} \rho_{j^*}^2 + \frac{1}{2 \cdot A^2} 4^{j^* s} \rho_{j^*}^2 \geq R^2 + \frac{1}{2 \cdot A^2} \rho_{j^*} M_{j^*} + \frac{1}{2 \cdot A^2} 4^{j^* s} \rho_{j^*}^2.$$

Subcase 2: $4^{j^* s} \rho_{j^*}^2 < \rho_{j^*} M_{j^*}$

In that case we have

$$\|P_{\bar{j}} f\|_{\mathcal{B}_s} > \frac{4^{j^* s}}{2^{\bar{j}^s}} \rho_{j^*} \geq 2^{\bar{j}^s} \rho_{j^*}$$

and thus

$$\begin{aligned} \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 &= \|P_{j \neq \bar{j}} f\|_{\mathcal{B}_s}^2 + \|P_{\bar{j}} f\|_{\mathcal{B}_s}^2 \\ &> R^2 + 2^{\bar{j}^s} \rho_{j^*} \|P_{\bar{j}} f\|_{\mathcal{B}_s} \\ &= R^2 + \rho_{j^*} M_{j^*} \\ &\geq R^2 + \frac{1}{2} \rho_{j^*} M_{j^*} + \frac{1}{2} 4^{j^* s} \rho_{j^*}^2. \end{aligned}$$

This concludes the proof since in any case (5) holds.

□

5.3 Estimation of M_{j^*}

As a last major step before directly controlling the type-I and type-II error probabilities, we need to find an appropriate estimator for M_{j^*} .

Lemma 6. For $\delta \in (0, 1)$ and $j^* \in \mathcal{J}$, let

$$C_\delta := \sqrt{\frac{2}{\delta}},$$

$$D_{j^*, \delta} := \frac{4^{j^*s}}{\sqrt{n}} \left(\sqrt{2}C_\delta + 2^{j^*/4} \sqrt{C_\delta} \right)$$

and define the events

$$\xi_{j^*, \delta}^0 := \left\{ M_{j^*} \leq \sqrt{\max_{2 \leq j \leq j^*} |Y_j|} + D_{\delta, j^*} \right\}, \quad (5.12)$$

$$\xi_{j^*, \delta}^1 := \left\{ M_{j^*} \geq \sqrt{\max_{2 \leq j \leq j^*} |Y_j|} - D_{\delta, j^*} \right\}. \quad (5.13)$$

Then, for any monotone decreasing sequence $(\beta_j)_{j \in \mathcal{J}}$ in $(0, 1)$, the following holds:

$$\mathbb{P}(\xi_{j^*, \beta_{j^*}}^1) \geq 1 - \sum_{j=2}^{j^*} \beta_j, \quad \mathbb{P}(\xi_{j^*, \beta_{j^*}}^0) \geq 1 - \beta_{j^*}. \quad (5.14)$$

◁

Proof. Remembering (5.1), we know that for $j \in \{2, 3, \dots, j^*\}$

$$Z_j := 4^{js} \|P_j \hat{f}\|_{\mathcal{B}_s}^2 = 16^{js} \|P_j \hat{f}\|_{L_2}^2$$

has the properties

$$\begin{aligned} \mathbb{E}[Z_j] &= 16^{js} \frac{2^j}{n} + 4^{js} \|P_j f\|_{\mathcal{B}_s}^2, \\ \text{Var}[Z_j] &= 2 \cdot 16^{2js} \left(\frac{2^j}{n^2} + \frac{2}{n} \|P_j f\|_{L_2}^2 \right) \\ &= 16^{js} \left(2 \cdot 16^{js} \frac{2^j}{n^2} + \frac{4}{n} 4^{js} \|P_j f\|_{\mathcal{B}_s}^2 \right). \end{aligned}$$

Now observe that for $\delta \in (0, 1)$

$$\begin{aligned} \sqrt{\frac{1}{\delta} \text{Var}[Z_j]} &\leq \sqrt{\frac{2}{\delta}} 2^{j/2} \frac{16^{js}}{n} + \frac{2}{\sqrt{\delta n}} 4^{js} 2^{js} \|P_j f\|_{\mathcal{B}_s} \\ &\leq C_\delta 2^{j^*/2} \frac{16^{j^*s}}{n} + \sqrt{2} C_\delta \frac{4^{j^*s}}{\sqrt{n}} M_{j^*} \\ &=: v_{\delta, j^*}. \end{aligned}$$

With $Y_j = Z_j - 16^{js} \frac{2^j}{n}$, Chebyshev's inequality now tells us that

$$\mathbb{P} \left(|Y_j - m_j^2| \geq v_{\delta, j^*} \right) \leq \delta. \quad (5.15)$$

We derive two bounds from this statement by lower bounding the the left hand side in two different ways:

On the one hand, observe

$$|Y_j - m_j^2| \geq ||Y_j| - m_j^2| \geq |Y_j| - m_j^2 \geq |Y_j| - M_{j^*}^2.$$

Now, since $(\beta_j)_{j \in \mathcal{J}}$ is monotone decreasing, the sequence $(v_{\beta_j, j^*})_{j \in \mathcal{J}}$ is increasing, so that via a union bound we obtain

$$\begin{aligned} \sum_{j=2}^{j^*} \beta_j &\geq \mathbb{P} \left(\exists j \in \{2, 3, \dots, j^*\} : |Y_j| \geq M_{j^*}^2 + v_{\beta_j, j^*} \right) \\ &\geq \mathbb{P} \left(\exists j \in \{2, 3, \dots, j^*\} : |Y_j| \geq M_{j^*}^2 + v_{\beta_{j^*}, j^*} \right) \\ &= \mathbb{P} \left(\sqrt{\max_{2 \leq j \leq j^*} |Y_j|} \geq \sqrt{M_{j^*}^2 + v_{\beta_{j^*}, j^*}} \right). \end{aligned}$$

With

$$\begin{aligned} \sqrt{M_{j^*}^2 + v_{\beta_{j^*}, j^*}} &= \sqrt{\left(M_{j^*} + \frac{C_{\beta_{j^*}}}{\sqrt{2}} \frac{4^{j^*s}}{\sqrt{n}} \right)^2 - \frac{C_{\beta_{j^*}}^2}{2} \frac{16^{j^*s}}{n} + C_{\beta_{j^*}} 2^{j^*/2} \frac{16^{j^*s}}{n}} \\ &\leq \sqrt{\left(M_{j^*} + \frac{C_{\beta_{j^*}}}{\sqrt{2}} \frac{4^{j^*s}}{\sqrt{n}} \right)^2 + \frac{C_{\beta_{j^*}}^2}{2} \frac{16^{j^*s}}{n} + C_{\beta_{j^*}} 2^{j^*/2} \frac{16^{j^*s}}{n}} \\ &\leq M_{j^*} + \frac{4^{j^*s}}{\sqrt{n}} \left(\sqrt{2} C_{\beta_{j^*}} + 2^{j^*/4} \sqrt{C_{\beta_{j^*}}} \right), \end{aligned}$$

we have

$$\mathbb{P} \left(\sqrt{\max_{2 \leq j \leq j^*} |Y_j|} \geq M_{j^*} + \frac{4^{j^*s}}{\sqrt{n}} \left(\sqrt{2} C_{\beta_{j^*}} + 2^{j^*/4} \sqrt{C_{\beta_{j^*}}} \right) \right) \leq \sum_{j=2}^{j^*} \beta_j$$

and hence the first claim from (6).

On the other hand, observe

$$|Y_j - m_j^2| \geq m_j^2 - |Y_j|$$

and consider the specific case $j = \bar{j}$ in (5.3):

$$\begin{aligned} \beta_{j^*} &\geq \mathbb{P} \left(|Y_{\bar{j}}| \leq M_{j^*}^2 - v_{\beta_{j^*}, j^*} \right) \\ &\geq \mathbb{P} \left(\max_{2 \leq j \leq j^*} |Y_j| \leq M_{j^*}^2 - v_{\beta_{j^*}, j^*} \right) \\ &= \mathbb{P} \left(\max_{2 \leq j \leq j^*} |Y_j| + \frac{16^{j^*s}}{n} \left(\frac{C_{\beta_{j^*}}^2}{2} + 2^{j^*/2} C_{\beta_{j^*}} \right) \leq \left(M_{j^*} - \frac{C_{\beta_{j^*}}}{\sqrt{2}} \frac{4^{j^*s}}{\sqrt{n}} \right)^2 \right) \\ &\geq \mathbb{P} \left(\sqrt{\max_{2 \leq j \leq j^*} |Y_j|} + \frac{4^{j^*s}}{\sqrt{n}} \left(\sqrt{2} C_{\beta_{j^*}} + 2^{j^*/4} \sqrt{C_{\beta_{j^*}}} \right) \leq M_{j^*} \right), \end{aligned}$$

which asserts the second claim from (6). □

5.4 Conclusion

We will now assemble the individual results of the previous sections to obtain the claim of Theorem 1. For $j \in \mathcal{J}$ we introduce

$$\rho_j = \frac{1346}{\sqrt{\eta}} \cdot \frac{2^{(3j+2J)/20}}{\sqrt{n}}, \quad \alpha_j = \eta \frac{1 - 2^{-1/5}}{4} 2^{(j-J)/5}, \quad \beta_j = \eta \frac{1 - 2^{-1/2}}{2} 2^{-j/2}, \quad (5.16)$$

so that in particular

$$\sum_{j=2}^J \alpha_j \leq \frac{\eta}{4}, \quad \sum_{j=2}^J \beta_j \leq \frac{\eta}{4}$$

and $(\beta_j)_{j \in \mathcal{J}}$ is monotone decreasing.

Result for fixed index

For $j^* \in \mathcal{J}$ define

$$T_{j^*, \alpha_{j^*}} = \|P_2^{j^*} \widehat{f}\|_{\mathcal{B}_s}^2 - A_{j^*} - \frac{2}{\sqrt{\alpha_{j^*}}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} \sqrt{\max_{2 \leq j \leq j^*} |Y_j|}.$$

Then under $H'_0 \cap \xi_{j^*, \beta_{j^*}}^0$, (6) and (5.1) yield that with probability at least $1 - \alpha_{j^*}$

$$\begin{aligned} T_{j^*, \alpha_{j^*}} &\leq \|P_2^{j^*} f\|_{\mathcal{B}_s}^2 + \frac{2}{\sqrt{\alpha_{j^*}}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} M_{j^*} + \frac{2}{\sqrt{\alpha_{j^*}}} 4^{j^* s} \frac{2^{j^*/2}}{n} - \frac{2}{\sqrt{\alpha_{j^*}}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} \sqrt{\max_{2 \leq j \leq j^*} |Y_j|} \\ &\leq R^2 + \frac{2}{\sqrt{\alpha_{j^*}}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} D_{j^*, \beta_{j^*}} + \frac{2}{\sqrt{\alpha_{j^*}}} 4^{j^* s} \frac{2^{j^*/2}}{n} \end{aligned}$$

so that with

$$\tau_{j^*, \alpha_{j^*}} = R^2 + \frac{2}{\sqrt{\alpha_{j^*}}} \left(\frac{\sqrt{j^* - 1}}{\sqrt{n}} D_{j^*, \beta_{j^*}} + 4^{j^* s} \frac{2^{j^*/2}}{n} \right),$$

we obtain

$$\mathbb{P}_{H'_0}(T_{j^*, \alpha_{j^*}} > \tau_{j^*, \alpha_{j^*}} \mid \xi_{j^*, \beta_{j^*}}^0) \leq \alpha_{j^*}.$$

On the other hand, let \mathbf{j}^* be a transition index with property (5). Then under $H'_1 \cap \xi_{\mathbf{j}^*, \beta_{\mathbf{j}^*}}^1$, (5.1) and (5) tell us that with probability at least $1 - \alpha_{\mathbf{j}^*}$

$$\begin{aligned} T_{\mathbf{j}^*, \alpha_{\mathbf{j}^*}} &\geq \|P_2^{\mathbf{j}^*} f\|_{\mathcal{B}_s}^2 - \frac{2}{\sqrt{\alpha_{\mathbf{j}^*}}} \cdot \frac{\sqrt{\mathbf{j}^* - 1}}{\sqrt{n}} M_{\mathbf{j}^*} - \frac{2}{\sqrt{\alpha_{\mathbf{j}^*}}} 4^{\mathbf{j}^* s} \frac{2^{\mathbf{j}^*/2}}{n} - \frac{2}{\sqrt{\alpha_{\mathbf{j}^*}}} \cdot \frac{\sqrt{\mathbf{j}^* - 1}}{\sqrt{n}} \sqrt{\max_{2 \leq j \leq \mathbf{j}^*} |Y_j|} \\ &\geq R^2 + \left(\frac{1}{2 \cdot A^2} \rho_{\mathbf{j}^*} - \frac{2}{\sqrt{\alpha_{\mathbf{j}^*}}} \cdot \frac{\sqrt{\mathbf{j}^* - 1}}{\sqrt{n}} \right) M_{\mathbf{j}^*} + \frac{1}{2 \cdot A^2} 4^{\mathbf{j}^* s} \rho_{\mathbf{j}^*}^2 \\ &\quad - \frac{2}{\sqrt{\alpha_{\mathbf{j}^*}}} 4^{\mathbf{j}^* s} \frac{2^{\mathbf{j}^*/2}}{n} - \frac{2}{\sqrt{\alpha_{\mathbf{j}^*}}} \cdot \frac{\sqrt{\mathbf{j}^* - 1}}{\sqrt{n}} \sqrt{\max_{2 \leq j \leq \mathbf{j}^*} |Y_j|}. \end{aligned}$$

Provided that

$$\frac{1}{2 \cdot A^2} \rho_{j^*} \geq \frac{4}{\sqrt{\alpha_{j^*}}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}}, \quad (5.17)$$

using (6) this yields

$$T_{j^*, \alpha_{j^*}} \geq R^2 + \frac{1}{2 \cdot A^2} 4^{j^*s} \rho_{j^*}^2 - \frac{2}{\sqrt{\alpha_{j^*}}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} D_{j^*, \beta_{j^*}} - \frac{2}{\sqrt{\alpha_{j^*}}} 4^{j^*s} \frac{2^{j^*/2}}{n} \quad (5.18)$$

Now by explicit computation we see that the choices in (5.4) ensure (5.4) as well as

$$\frac{1}{4 \cdot A^2} 4^{j^*s} \rho_{j^*}^2 \geq \frac{4}{\sqrt{\alpha_{j^*}}} \cdot \frac{\sqrt{j^* - 1}}{\sqrt{n}} D_{j^*, \beta_{j^*}} \quad \text{and} \quad \frac{1}{4 \cdot A^2} 4^{j^*s} \rho_{j^*}^2 \geq \frac{4}{\sqrt{\alpha_{j^*}}} 4^{j^*s} \frac{2^{j^*/2}}{n},$$

so that (5.4) can be continued as

$$T_{j^*, \alpha_{j^*}} \geq R^2 + \frac{2}{\sqrt{\alpha_{j^*}}} \left(\frac{\sqrt{j^* - 1}}{\sqrt{n}} D_{j^*, \beta_{j^*}} + 4^{j^*s} \frac{2^{j^*/2}}{n} \right) = \tau_{j^*, \alpha_{j^*}}$$

and hence, finally,

$$\mathbb{P}_{H'_1}(T_{j^*, \alpha_{j^*}} \leq \tau_{j^*, \alpha_{j^*}} \mid \xi_{j^*, \beta_{j^*}}^1) \leq \alpha_{j^*}.$$

Generalisation to unknown j^*

For our test

$$\varphi(P_2^J \hat{f}) = 1 - \prod_{j^*=2}^J \mathbb{1}_{\{T_{j^*, \alpha_{j^*}} \leq \tau_{j^*, \alpha_{j^*}}\}}, \quad (5.19)$$

we can conclude with (6) and (5.4) that on the one hand

$$\begin{aligned} \mathbb{P}_{H'_0}(\varphi = 1) &\leq \sum_{j^*=2}^J \left(\mathbb{P}_{H'_0}(T_{j^*, \alpha_{j^*}} > \tau_{j^*, \alpha_{j^*}} \mid \xi_{j^*, \beta_{j^*}}^0) + (1 - \mathbb{P}(\xi_{j^*, \beta_{j^*}}^0)) \right) \\ &\leq \frac{\eta}{4} + \frac{\eta}{4} = \frac{\eta}{2} \end{aligned} \quad (5.20)$$

and on the other hand

$$\begin{aligned} \mathbb{P}_{H'_1}(\varphi = 0) &\leq \mathbb{P}_{H'_1}(\forall j^* \in \mathcal{J} : T_{j^*, \alpha} \leq \tau_{j^*, \alpha}) \\ &\leq \mathbb{P}_{H'_1}(T_{j^*, \alpha} \leq \tau_{j^*, \alpha_{j^*}} \mid \xi_{j^*, \beta_{j^*}}^1) + (1 - \mathbb{P}(\xi_{j^*, \beta_{j^*}}^1)) \\ &\leq \alpha_{j^*} + \sum_{j=2}^{j^*} \beta_j \\ &\leq \frac{\eta}{4} + \frac{\eta}{4} = \frac{\eta}{2} \end{aligned} \quad (5.21)$$

Specification of J and conclusion

We are now ready to return to (5.1). Choose

$$J := \left\lfloor \frac{1}{2t + 1/2} \frac{\ln(n)}{\ln(2)} \right\rfloor,$$

so that

$$\frac{1}{2} n^{\frac{1}{2t+1/2}} \leq 2^J \leq n^{\frac{1}{2t+1/2}}. \quad (5.22)$$

That yields

$$2^{-Jt} \leq 2^t n^{-\frac{t}{2t+1/2}}. \quad (5.23)$$

and, on the other hand,

$$\rho_J = \frac{1346}{\sqrt{\eta}} \frac{2^{J/4}}{\sqrt{n}} \leq \frac{1346}{\sqrt{\eta}} \cdot n^{-\frac{t}{2t+1/2}}.$$

Therefore, whenever we choose

$$\rho \geq \left(\frac{1346}{\sqrt{\eta}} + \frac{R}{1 - 2^{-t}} \right) n^{-\frac{t}{2t+1/2}},$$

indeed by (5.4) and (5.4)

$$\sup_{f \in B_s(R)} \mathbb{P}_f(\varphi = 1) + \sup_{f \in \tilde{B}_{s,t}(R,\rho)} \mathbb{P}_f(\varphi = 0) \leq \frac{\eta}{2} + \frac{\eta}{2} = \eta.$$

6 Proof of Theorem 2

6.1 Description of the Strategy

Motivated by (1), given $\eta \in (0, 1)$, we aim at finding $\rho > 0$ such that for any test φ ,

$$\sup_{f \in B_s(R)} \mathbb{P}(\varphi = 1) + \sup_{f \in \tilde{B}_{s,t}(R,\rho)} \mathbb{P}(\varphi = 0) > \eta.$$

This can be achieved through a Bayesian-type approach, see e.g. [1]: Let ν_0, ν_ρ be probability distributions (priors) such that $\text{supp}(\nu_0) \subseteq B_s(R)$ and $\text{supp}(\nu_\rho) \subseteq \tilde{B}_{s,t}(R, \rho)$. Then we have

$$\begin{aligned} \sup_{f \in B_s(R)} \mathbb{P}_f(\varphi = 1) + \sup_{f \in \tilde{B}_{s,t}(R,\rho)} \mathbb{P}_f(\varphi = 0) &\geq \mathbb{P}_{f \sim \nu_0}(\varphi = 1) + \mathbb{P}_{f \sim \nu_\rho}(\varphi = 0) \\ &\geq 1 - \frac{1}{2} \|\mathbb{P}_{f \sim \nu_\rho} - \mathbb{P}_{f \sim \nu_0}\|_{\text{TV}} \\ &\geq 1 - \frac{1}{2} \left(\int \left(\frac{d\mathbb{P}_{f \sim \nu_\rho}}{d\mathbb{P}_{f \sim \nu_0}}(x) \right)^2 d\mathbb{P}_{f \sim \nu_0}(x) - 1 \right)^{\frac{1}{2}}. \end{aligned}$$

This tells us that if we find $\tilde{\rho} > 0$ such that

$$\int \left(\frac{d\mathbb{P}_{f \sim \nu_\rho}}{d\mathbb{P}_{f \sim \nu_0}} \right)^2 d\mathbb{P}_{f \sim \nu_0} < 1 + 4(1 - \eta)^2, \quad (6.1)$$

for any test φ it holds that

$$\sup_{f \in B_s(R)} \mathbb{P}(\varphi = 1) + \sup_{f \in \tilde{B}_{s,t}(R,\rho)} \mathbb{P}(\varphi = 0) > \eta$$

and hence

$$\rho^* \geq \tilde{\rho}.$$

6.2 Application to our Problem

Priors

Since the upper bound does not depend on s and we found the index J from (??) to be critical, we choose the following structurally simple priors: Let ν_0 be the Dirac- δ distribution on $\{0\}$ (i.e. $f \equiv 0$) and ν_ρ be the uniform distribution on

$$\mathcal{A}_{\rho,v} := \left\{ \sum_{k=1}^{2^J} a_{J,k} \psi_{j,k} \mid a_{J,1}, a_{J,2}, \dots, a_{J,2^J} \in \{v, -v\} \right\},$$

where $v > 0$ needs further specification: On the one hand, it is necessary to ensure that each $f \in \mathcal{A}_{\rho,v}$ fulfils $\|f\|_{\mathcal{B}_t} \leq R$ - note that for any such f , $\|f\|_{L_2} = 2^{J/2}v$, so that by construction that condition reads

$$2^{J(t+1/2)}v \leq R.$$

This motivates the choice $v := a_\eta \cdot R \cdot 2^{-J(t+1/2)}$ for some $a_\eta \in (0, 1]$ specified later based on further restrictions. On the other hand, we require

$$\rho \leq \inf_{h \in B_s(R)} \|f - h\|_{L_2}. \quad (6.2)$$

Since only the level J is involved, this is in fact merely the minimum over the Euclidean ball with radius $R \cdot 2^{-Js}$ so that

$$\inf_{h \in B_s(R)} \|f - h\|_{L_2} = \max(0, 2^{J/2}v - R \cdot 2^{-Js}).$$

Now, by explicit computation we see that if

$$n \geq \left(\frac{2^{1+s-t}}{a_\eta} \right)^{\frac{2t+1/2}{s-t}},$$

with our choice of v we have

$$\max(0, 2^{J/2}v - R \cdot 2^{-Js}) \geq \frac{1}{2} 2^{J/2}v = a_\eta \frac{R}{2} 2^{-Jt},$$

so that (6.2) holds if

$$\rho \leq a_\eta \frac{R}{2} 2^{-Jt}$$

Statistical distance

The central task in this proof is to compute the quantities on the left hand side of (6.1): By construction, $\mathbb{P}_{f \sim \nu_0}$ corresponds to the 2^J -fold product of Gaussian distributions with mean 0 and variance $\frac{1}{n}$, so that for $x \in \mathbb{R}^{2^J}$

$$d\mathbb{P}_{f \sim \nu_0}(x) = \sqrt{\frac{n}{2\pi}}^{2^J} \prod_{k=1}^{2^J} \exp\left(-\frac{n}{2}x_k^2\right).$$

On the other hand, $\mathbb{P}_{f \sim \nu_\rho}$ corresponds to a uniform mixture of 2^{2^J} products of 2^J independent Gaussians with means of the form $\pm v$ and variance $\frac{1}{n}$. More precisely, let $\mathcal{S} := \{1, -1\}^{2^J}$ and R be uniformly distributed on \mathcal{S} (i.e. the product of 2^J Rademacher variables). Then

$$\begin{aligned} d\mathbb{P}_{f \sim \nu_\rho}(x) &= \frac{1}{2^{2^J}} \sum_{\alpha \in \mathcal{S}} \sqrt{\frac{n}{2\pi}}^{2^J} \prod_{k=1}^{2^J} \exp\left(-\frac{n}{2} \sum_{k=1}^{2^J} (x_k - \alpha_k v)^2\right) \\ &= \sqrt{\frac{n}{2\pi}}^{2^J} \mathbb{E}_R \left[\prod_{k=1}^{2^J} \exp\left(-\frac{n}{2} (x_k - R_k v)^2\right) \right] \end{aligned}$$

and furthermore, with an independent copy R' of R ,

$$\begin{aligned} (d\mathbb{P}_{f \sim \nu_\rho}(x))^2 &= \left(\frac{n}{2\pi}\right)^{2^J} \mathbb{E}_{R, R'} \left[\prod_{k=1}^{2^J} \exp\left(-\frac{n}{2} [(x_k - R_k v)^2 + (x_k - R'_k v)^2]\right) \right] \\ &= \left(\frac{n}{2\pi}\right)^{2^J} \exp(-2^J n v^2) \mathbb{E}_{R, R'} \left[\prod_{k=1}^{2^J} \exp(-n x_k^2 + n v x_k (R_k + R'_k)) \right]. \end{aligned}$$

The quotient we need to integrate in (6.1) therefore reads

$$\begin{aligned} \frac{(d\mathbb{P}_{f \sim \nu_\rho})^2}{d\mathbb{P}_{f \sim \nu_0}}(x) &= \sqrt{\frac{n}{2\pi}}^{2^J} \exp(-2^J n v^2) \mathbb{E}_{R, R'} \left[\prod_{k=1}^{2^J} \exp\left(-\frac{n}{2} x_k^2 + n v x_k (R_k + R'_k)\right) \right] \\ &= \sqrt{\frac{n}{2\pi}}^{2^J} \exp(-2^J n v^2) \mathbb{E}_{R, R'} \left[\prod_{k=1}^{2^J} \exp\left(-\frac{n}{2} (x_k - v(R_k + R'_k))^2\right) \exp(n v^2 (1 + R_k R'_k)) \right] \\ &= \sqrt{\frac{n}{2\pi}}^{2^J} \mathbb{E}_{R, R'} \left[\prod_{k=1}^{2^J} \exp(n v^2 R_k R'_k) \prod_{k=1}^{2^J} \exp\left(-\frac{n}{2} (x_k - v(R_k + R'_k))^2\right) \right]. \end{aligned}$$

Since the product of independent Rademacher variables is itself a Rademacher variable, we obtain

$$\begin{aligned}
\int_{\mathbb{R}^{2^J}} \frac{(\mathrm{d}\mathbb{P}_{f \sim \nu_\rho})^2}{\mathrm{d}\mathbb{P}_{f \sim \nu_0}}(x) \, dx &= \mathbb{E}_{R, R'} \left[\prod_{k=1}^{2^J} \exp(nv^2 R_k R'_k) \right] \\
&= \mathbb{E}_R \left[\prod_{k=1}^{2^J} \exp(nv^2 R_k) \right] \\
&= \prod_{k=1}^{2^J} \mathbb{E}_{R_k} [\exp(nv^2 R_k)] \\
&= (\cosh(nv^2))^{2^J} \\
&\stackrel{(*)}{\leq} (1 + n^2 v^4)^{2^J} \\
&\leq \exp(2^J n^2 v^4)
\end{aligned}$$

where we require $nv^2 \leq 1$ for (*). Since by (5.4)

$$v = a_\eta \cdot R \cdot 2^{-J(t+1/2)} \leq a_\eta \cdot R (2n^{-1/(2t+1/2)})^{t+1/2} \leq a_\eta \cdot R \cdot 2^{t+1/2} n^{-1/2},$$

this is fulfilled if

$$a_\eta \leq \frac{1}{R \cdot 2^{t+1/2}}. \quad (6.3)$$

Conclusion

We are now ready to combine the preparations from above: Let $\eta \in (0, 1)$. Then, provided that (6.2) is ensured, (6.1) holds if

$$\exp(2^J n^2 v^4) \leq 1 + 4(1 - \eta)^2$$

which, by explicit computation, can be written as

$$a_\eta \leq \frac{2^{J(t+1/4)}}{\sqrt{n}R} \sqrt{\ln(1 + 4(1 - \eta)^2)}.$$

Through (5.4) and (5.4) we find that

$$\frac{2^{J(t+1/4)}}{\sqrt{n}} \geq \frac{2^{-t}}{16}$$

and obtain the stronger condition

$$a_\eta \leq \frac{\sqrt{\ln(1 + 4(1 - \eta)^2)}}{2^t 16R}.$$

In summary: Given $\eta \in (0, 1)$, let

$$a_\eta = \min \left\{ 1, \frac{1}{R \cdot 2^{t+1/2}}, \frac{\sqrt{\ln(1 + 4(1 - \eta)^2)}}{2^t 16R} \right\}.$$

If

$$n \geq \left\lceil \left(\frac{2^{1+s-t}}{a_\eta} \right)^{\frac{2t+1/2}{s-t}} \right\rceil,$$

the priors ν_0 and ν_ρ meet all requirements from section 6.1 and the lower bound

$$\rho^* \geq a_\eta \frac{R}{2} 2^{-Jt} \geq a_\eta \frac{R}{2} n^{-\frac{t}{2t+1/2}}$$

is established, where we write $C_\eta := \frac{R}{2} a_\eta$.

References

- [1] BARAUD, Y. NON-ASYMPTOTIC MINIMAX RATES OF TESTING IN SIGNAL DETECTION. *Bernoulli* 8, 5 (2002), 577–606.
- [2] BLANCHARD, G., CARPENTIER, A., AND GUTZEIT, M. MINIMAX EUCLIDEAN SEPARATION RATES FOR TESTING CONVEX HYPOTHESES IN \mathbb{R}^d . *Electronic Journal of Statistics* 12, 2 (2018), 3713–3735.
- [3] BULL, A. D., AND NICKL, R. ADAPTIVE CONFIDENCE SETS IN L_2 . *Probability Theory and Related Fields* 156, 3 (2013), 889–919.
- [4] CAI, T. T., AND LOW, M. G. ADAPTIVE CONFIDENCE BALLS. *The Annals of Statistics* 34, 1 (2006), 202–228.
- [5] CAI, T. T., AND LOW, M. G. TESTING COMPOSITE HYPOTHESES, HERMITE POLYNOMIALS AND OPTIMAL ESTIMATION OF A NONSMOOTH FUNCTIONAL. *The Annals of Statistics* 39, 2 (2011), 1012–1041.
- [6] CARPENTIER, A. TESTING THE REGULARITY OF A SMOOTH SIGNAL. *Bernoulli* 21, 1 (2015), 465–488.
- [7] COHEN, A., DAUBECHIES, I., AND VIAL, P. WAVELETS ON THE INTERVAL AND FAST WAVELET TRANSFORMS. *Applied and computational harmonic analysis* 1, 1 (1993), 54–81.
- [8] COMMINGES, L., AND DALALYAN, A. MINIMAX TESTING OF A COMPOSITE NULL HYPOTHESIS DEFINED VIA A QUADRATIC FUNCTIONAL IN THE MODEL OF REGRESSION. *Electronic Journal of Statistics* 7 (2013), 146–190.
- [9] GINÉ, E., AND NICKL, R. MATHEMATICAL FOUNDATIONS OF INFINITE-DIMENSIONAL STATISTICAL MODELS.
- [10] INGSTER, Y. ASYMPTOTICALLY MINIMAX HYPOTHESIS TESTING FOR NONPARAMETRIC ALTERNATIVES. I, II, III. *Math. Methods Statist* 2, 2 (1993), 85–114.
- [11] INGSTER, Y., AND SUSLINA, I. A. NON-PARAMETRIC GOODNESS-OF-FIT TESTING UNDER GAUSSIAN MODELS. Springer, 2002.

- [12] JUDITSKY, A., AND LAMBERT-LACROIX, S. NONPARAMETRIC CONFIDENCE SET ESTIMATION. *Math. Methods Statist* 12, 4 (2003), 410–428.
- [13] JUDITSKY, A., AND NEMIROVSKI, A. ON NONPARAMETRIC TESTS OF POSITIVITY/MONOTONICITY/CONVEXITY. *The Annals of Statistics* (2002), 498–527.
- [14] LE CAM, L. ASYMPTOTIC METHODS IN STATISTICAL DECISION THEORY. Springer Science & Business Media, 2012.
- [15] TRIEBEL, H. THEORY OF FUNCTION SPACES II. Birkhäuser Basel, 1992.