# Unique continuation properties for Schredinger operators in Hilbert spaces

Veli Shakhmurov

Department of Mechanical Engineering, Okan University, Akfirat, Tuzla 34959 Istanbul, Turkey,

E-mail: veli.sahmurov@okan.edu.tr

Institute of Mathematics and Mechanics, Azerbaijan National Academy of Sciences, Azerbaijan, AZ1141, Baku, F. Agaev, 9,

E-mail: veli.sahmurov@gmail.com

#### Abstract

Here, the Morgan type uncertainty principle and unique continuation properties of abstract Schredinger equations with time dependent potentials are obtained in Hilbert space valued function classes. The equations include linear operator in abstract Hilbert spaces  $H$  dependent on space variables. So, by selecting appropriate spaces  $H$  and operators, we derive unique continuation properties for numerous classes of Schrödinger type equations and its systems, which occur in a wide variety of physical systems.

Key Word: Schredinger equations, Positive operators, Semigroups of operators, Unique continuation, Morgan type uncertainty principle

## AMS 2010: 35Q41, 35K15, 47B25, 47Dxx, 46E40 1. Introduction, definitions

In this paper, the unique continuation properties of the abstract Schrödinger equations

$$
i\partial_{t}u + \Delta u + A(x)u + V(x,t)u = 0, \ x \in R^{n}, \ t \in [0,1], \tag{1.1}
$$

are studied, where  $A = A(x)$  is a linear operator,  $V(x, t)$  is a given potential operator function in a Hilbert space  $H$ , the subscript t indicates the partial derivative with respect to t,  $\Delta$  denotes the Laplace operator in  $R^n$  and  $u =$  $u(x, t)$  is the H-valued unknown function. The goal is to obtain sufficient conditions on  $A$ , the potential  $V$  and the behavior of the solution  $u$  at two different times,  $t_0 = 0$  and  $t_1 = 1$ , which guarantee that  $u \equiv 0$  in  $R^n \times [0, 1]$ . This linear result is then applied to show that two regular solutions  $u_1$  and  $u_2$  of non-linear Schredinger equation

$$
i\partial_{t}u + \Delta u + Au = F(u, \bar{u}), \ x \in R^{n}, \ t \in [0, 1], \tag{1.2}
$$

for general non-linearities F, must agree in  $R^n \times [0,1]$ , when  $u_1 - u_2$  and its gradient decay faster than any quadratic exponential at times 0 and 1.

Unique continuation properties for Schredinger equations have been studied by [5-8] , [21-23] and the references therein. In contrast to the mentioned above results we will study the unique continuation properties of abstract Schredinger equations with operator potentials. Abstract differential equations were studied e.g. in [1], [10], [13 – 19], [25, 26]. Since the Hilbert space H is arbitrary and A, V are possible linear operators, by choosing  $H$ ,  $\Lambda$  and  $V$  we can obtain numerous classes of Schrödinger type equations and its systems which occur in different applications. If we choose the abstract space  $H$  a concrete Hilbert space, for example  $H = L^2(G)$ ,  $A = L$ ,  $V = q(x, t)$ , where G is a domain in  $R^m$  with smooth boundary, L is a elliptic operator with respect to the variable  $y \in G$ and  $q$  is a complex valued function, then we obtain the unique continuation properties of the following Schrödinger equation

$$
\partial_t u = i \left[ \Delta u + Lu + q(x, t) u \right], \ x \in R^n, \ y \in G, \ t \in [0, 1], \tag{1.3}
$$

where  $u = u(x, y, t)$ . Moreover, let  $H = L<sup>2</sup>(0, 1)$  and let A to be a differential operator with generalized Wentzell-Robin boundary condition defined by

$$
D(A) = \{u \in W^{2,2}(0,1), B_j u = Au(j) = 0, j = 0,1\},\tag{1.4}
$$

$$
Au = au^{(2)} + bu^{(1)},
$$

where a and b are real valued functions on  $R^n \times [0,1]$ . Then, we obtain the unique continuation properties of the Wentzell-Robin type boundary value problem (BVP) for the nonlinear Schredinger type equation

$$
i\partial_t u + \Delta u + a \frac{\partial^2 u}{\partial y^2} + b \frac{\partial u}{\partial y} = F(u, \bar{u}),
$$
  
\n
$$
u = u(x, y, t), x \in R^n, y \in G, t \in [0, 1],
$$
\n
$$
(1.5)
$$

$$
a(j) u_{yy}(x, j, t) + b(j) u_{y}(x, j, t) = 0, j = 0, 1, \text{ for a.e. } x \in R^{n}, t \in (0, 1).
$$
\n(1.6)

Note that, the regularity properties of Wentzell-Robin type BVP for elliptic equations have been studied e.g. in [11, 12 ] and the references therein. Moreover, if put  $H = l_2$  and choose A and  $V(x, t)$  as infinite matrices  $[a_{m i}]$ ,  $[b_{mi}(x,t)]$ ,  $m, j = 1, 2, ..., \infty$  respectively, then we obtain the unique continuation properties of the following system of Schredinger equation

$$
\partial_t u_m = i \left[ \Delta u_m + \sum_{j=1}^N \left( a_{mj} + b_{mj}(x, t) \right) u_j \right], \ x \in R^n, \ t \in (0, 1). \tag{1.7}
$$

Let E be a Banach space and  $\gamma = \gamma(x)$  be a positive measurable function on a domain  $\Omega \subset \mathbb{R}^n$ . Here,  $L_{p,\gamma}(\Omega;E)$  denotes the space of strongly measurable E-valued functions that are defined on  $\Omega$  with the norm

$$
\|f\|_{p,\gamma} = \|f\|_{L_{p,\gamma}(\Omega;E)} = \left(\int \|f(x)\|_{E}^{p} \gamma(x) \, dx\right)^{\frac{1}{p}}, \ 1 \le p < \infty,
$$

$$
||f||_{L_{\infty,\gamma}(\Omega;E)} = \operatorname{ess} \sup_{x \in \Omega} ||f(x)||_{E} \gamma(x), \quad p = \infty.
$$

For  $\gamma(x) \equiv 1$  the space  $L_{p,\gamma}(\Omega;E)$  will be denoted by  $L^p = L^p(\Omega;E)$  for  $p \in [1,\infty]$ .

Let  $\mathbf{p} = (p_1, p_2)$  and  $\Omega = \Omega_1 \times \Omega_2$ , where  $\Omega_k \in R^{n_k}$ .  $L_x^{p_1} L_t^{p_2} (\Omega; E)$  will denote the space of all E-valed, **p**-summable functions with mixed norm, i.e., the space of all measurable functions  $f$  defined on  $G$  equipped with norm

$$
||f||_{L_x^{p_1}L_y^{p_2}(\Omega;E)} = \left( \left( \int_{\Omega_1} ||f(x,t)||_E^{p_2} dt \right)^{\frac{p_1}{p_2}} dx \right)^{\frac{1}{p_1}}.
$$

Let  $H$  be a Hilbert space and

$$
||u|| = ||u||_H = (u, u)_H^{\frac{1}{2}} = (u, u)^{\frac{1}{2}}
$$
 for  $u \in H$ .

For  $p = 2$  and  $E = H$  is a Hilbert space,  $L^p(\Omega; E)$  is to be a Hilbert space with inner product

$$
(f,g)_{L^2(\Omega;H)} = \int_{\Omega} (f(x),g(x))_H dx
$$

for  $f, g \in L^2(\Omega; H)$ .

Let  $C(\Omega; E)$  denote the space of E-valued, bounded uniformly continuous functions on  $\Omega$  with norm

$$
||u||_{C(\Omega;E)} = \sup_{x \in \Omega} ||u(x)||_{E}.
$$

 $C^m(\Omega; E)$  will denote the space of E-valued bounded uniformly strongly continuous and m-times continuously differentiable functions on  $\Omega$  with norm

$$
||u||_{C^m(\Omega;E)} = \max_{0 \leq |\alpha| \leq m} \sup_{x \in \Omega} ||D^{\alpha}u(x)||_{E}.
$$

Moreover,  $C_0^{\infty}(\Omega; E)$  –denotes the space of E-valued infinity many differentiable finite functions.

Let

$$
O_r = \{ x \in R^n, \ |x| < r \}, \ r > 0.
$$

Let  $\mathbb N$  denote the set of all natural numbers,  $\mathbb C$  denote the set of all complex numbers.

Let  $\Omega$  be a domain in  $\mathbb{R}^n$  and let m be a positive integer.  $W^{m,p}(\Omega;E)$ denotes the space of all functions  $u \in L^p(\Omega; E)$  that have generalized derivatives  $\frac{\partial^m u}{\partial x_k^m} \in L^p(\Omega; E), 1 \leq p \leq \infty$  with the norm

$$
||u||_{W^{m,p}(\Omega;E)} = ||u||_{L^p(\Omega;E)} + \sum_{k=1}^n \left\|\frac{\partial^m u}{\partial x_k^m}\right\|_{L^p(\Omega;E)} < \infty.
$$

Let  $E_0$  and E be two Banach spaces and suppose  $E_0$  is continuously and densely embedded into E. Here,  $W^{m,p}(\Omega; E_0, E)$  denotes the space  $W^{m,p}(\Omega; E) \cap$  $L^p(\Omega; E)$  equipped with norm

$$
||u||_{W^{m,p}(\Omega;E_0,E)} = ||u||_{L^p(\Omega;E_0)} + \sum_{k=1}^n \left\| \frac{\partial^m u}{\partial x_k^m} \right\|_{L^p(\Omega;E)} < \infty.
$$

Let  $E_1$  and  $E_2$  be two Banach spaces. Let  $L(E_1, E_2)$  will denote the space of all bounded linear operators from  $E_1$  to  $E_2$ . For  $E_1 = E_2 = E$  it will be denoted by  $L(E)$ .

Let A is a symmetric operator in a Hilbert sapace H with domain  $D(A)$ . Here,  $H(A)$  denotes the domain of A equipped with graphical norm, i.e.

$$
||u||_{H(A)} = ||Au|| + ||u||.
$$

The symmetric operator A is positive defined if  $D(A)$  is dense in H and there exists a positive constant  $C_A$  depend only on  $A$  such that

$$
(Au, u) \ge C_A \|u\|^2
$$

.

It is known that see e.g. [24, §1.15.1]) there exist fractional powers  $A^{\theta}$  of the positive defined operator A.

Let  $[A, B]$  be a commutator operator, i.e.

$$
[A, B] = AB - BA
$$

for linear operators A and B.

Sometimes we use one and the same symbol  $C$  without distinction in order to denote positive constants which may differ from each other even in a single context. When we want to specify the dependence of such a constant on a parameter, say  $\alpha$ , we write  $C_{\alpha}$ .

### 2.1. Main results for absract Scrödinger equation

Consider the problem (1.1). Here,

$$
X = L^{2}(R^{n}; H), X (A) = L^{2}(R^{n}; H (A)), Y^{k} = W^{2,k}(R^{n}; H), k \in \mathbb{N}.
$$

**Definition 2.1.** A function  $u \in L^{\infty}(0,T;H(A))$  is called a local weak solution to (1.1) on  $(0, T)$  if u belongs to  $L^{\infty}(0, T; H(A)) \cap W^{1,2}(0, T; H)$  and satisfies (1.1) in the sense of  $L^{\infty}(0,T;H(A))$ . In particular, if  $(0,T)$  coincides with  $\mathbb{R}$ , then u is called a global weak solution to  $(1.1)$ .

If the solution of (1.1) belongs to  $C([0,T]; H(A)) \cap W^{2,2}(0,T;H)$ , then its called a stronge solution.

**Condition 1.** Assume: (1)  $A = A(x)$  is a symmetric operator in Hilbert space H with independent on x domain  $D(A)$  that is dense on H;

(2)  $\frac{\partial A}{\partial x_k}$  are symmetric operators in Hilbert space H with independent on x domain  $D\left(\frac{\partial A}{\partial x_k}\right) = D(A)$  and

$$
\sum_{k=1}^{n} \left( x_k \left[ A \frac{\partial f}{\partial x_k} - \frac{\partial A}{\partial x_k} f \right], f \right)_X \ge 0,
$$

for each  $f \in L^{\infty}(0,T;Y^1(A));$ (3)

$$
\sum_{k=1}^{n} \left( x_k \left[ A \frac{\partial f}{\partial x_k} + \frac{\partial A}{\partial x_k} f \right], f \right)_X \ge 0
$$

for each  $f \in L^{\infty}(0,T;Y^1(A));$ 

(4) iA generates a Schrödinger grop and  $V(x,t) \in L(H)$  for  $(x,t) \in R^n$  ×  $[0, 1]$ ;

(5) there is a constant  $C_0 > 0$  so that

Im 
$$
((A + V) v, v)_H \geq C_0 \mu (x, t) ||v (x, t)||^2
$$
,

for  $x \in R^n$ ,  $t \in [0, T]$ ,  $T \in (0, 1]$  and  $v \in D(A)$ , where  $\mu$  is a positive function in  $L^1(0,T;L^{\infty}(R^n))$ ;

(6)

$$
||V||_{L^{\infty}(R^n \times (0,1);L(H))} \leq C, \ \ \lim_{R \to \infty} ||V||_{L^1_t L^{\infty}_x(L(H))} = 0,
$$

where

$$
L_{t}^{1} L_{x}^{\infty} (L (H)) = L^{1} (0, 1; L^{\infty} (R^{n}/O_{r}) ; L (H)).
$$

Let  $A = A(x)$  is a symmetric operator in Hilbert space H with independent on  $x$  domain  $D(A)$  and

$$
X(A) = L^2(R^n; H(A)), Y^k(A) = W^{2,k}(R^n; H(A)), k \in \mathbb{N}
$$

Our main result in this paper is the following:

**Theorem 1.** Assume the Condition 1 holds and there exist constants  $a_0$ ,  $a_1, a_2 > 0$  such that for any  $k \in \mathbb{Z}^+$  a solution  $u \in C([0,1]; X(A))$  of  $(1.1)$ satisfy

$$
\int_{R^{n}} \|u(x,0)\|^{2} e^{2a_{0}|x|^{p}} dx < \infty, \text{ for } p \in (1,2), \qquad (2.1)
$$

$$
\int_{R^{n}} \|u(x,1)\|^{2} e^{2k|x|^{p}} dx < a_{2} e^{2a_{1}k^{\frac{q}{q-p}}}, \frac{1}{p} + \frac{1}{q} = 1.
$$
 (2.2)

Moreover, there exists  $M_p > 0$  such that

$$
a_0 a_1^{p-2} > M_p. \tag{2.3}
$$

Then  $u(x,t) \equiv 0$ .

Corollary 1. Assume the Condition1 holds and

$$
\lim_{|R| \to \infty} \int_{0}^{1} \sup_{|x| > R} \|V(x, t)\|_{L(H)} dt = 0.
$$

There exist positive constants  $\alpha$ ,  $\beta$  such that a solution  $u \in C([0,1]; X(A))$  of  $(1.1)$  satisfy

$$
\int_{R^n} \|u(x,0)\|^2 e^{2|\alpha x|^p/p} dx + \int_{R^n} \|u(x,1)\|^2 e^{\frac{2|\beta x|^q}{q}} dx < \infty,
$$
\n(2.4)

with

$$
p\in (1,2)\,,\,\frac{1}{p}+\frac{1}{q}=1
$$

and there exists  $N_p > 0$  such that

$$
\alpha \beta > N_p. \tag{2.5}
$$

Then  $u(x,t) \equiv 0$ .

As a direct consequence of Corollary1 we have the following result regarding the uniqueness of solutions for nonlinear equation (1.2):

**Theorem 2.** Assume the Condition 1 holds and  $u_1, u_2 \in C([0,1]; Y^{2,k}(A))$ strong solutions of (1.2) with  $k \in \mathbb{Z}^+, k > \frac{n}{2}$ . Suppose  $F : H \times H \to H, F \in C^k$ ,  $F(0) = \partial_u F(0) = \partial_{\bar{u}} F(0) = 0$  and there exist positive constants  $\alpha$ ,  $\beta$  such that

$$
e^{\frac{|\alpha x|^p}{p}}(u_1(.,0) - u_2(.,0)) \in X, \ e^{\frac{|\beta x|^q}{q}}(u_1(.,0) - u_2(.,0)) \in X,
$$
 (2.6)

with

$$
p \in (1,2)\, ,\, \frac{1}{p} + \frac{1}{q} = 1
$$

and there exists  $N_p > 0$  such that

$$
\alpha \beta > N_p. \tag{2.7}
$$

Then  $u_1(x,t) \equiv u_2(x,t)$ .

Corollary 2. Assume the Condition 1 hold and there exist positive constants  $\alpha$  and  $\beta$  such that a solution  $u \in C([0,1]; X(A))$  of  $(1.1)$  satisfy

$$
\int_{R^n} \|u(x,0)\|^2 e^{\frac{2|\alpha x_j|^p}{p}} dx + \int_{R^n} \|u(x,1)\|^2 e^{\frac{2|\beta x_j|^q}{q}} dx < \infty,
$$
\n(2.8)

for  $j = 1, 2, ..., n$  and  $p \in (1, 2), \frac{1}{p} + \frac{1}{q} = 1$ . Moreover, there exists  $N_p > 0$  such that

$$
\alpha \beta > N_p. \tag{2.9}
$$

Then  $u(x,t) \equiv 0$ .

**Remark 2.1.** The Theorem 2 still holds, with different constant  $N_p > 0$ , if one replaces the hypothesis (2.6) by

$$
e^{|\alpha x_j|^p/p} (u_1(.,0) - u_2(.,0)) \in X(A), e^{|\beta x_j|^q/q} (u_1(.,0) - u_2(.,0)) \in X(A),
$$

for  $j = 1, 2, ..., n$ .

Next, we shall extend the method used in the proof Theorem 3 to study the blow the nonlinear Schrödinger equations

$$
i\partial_{t}u + \Delta u + Au + F(u, \bar{u})u = 0, \ x \in \mathbb{R}^{n}, \ t \in [0, 1], \tag{2.10}
$$

where A is a linear operator in a Hilbert space H.

Let  $u(x, t)$  be a solution of the equation (2.10). Then it can be shown that the function

$$
u(x,t) = U(x, 1-t) u\left(\frac{x}{1-t}, \frac{t}{1-t}\right),
$$
\n(2.11)

is a solution of the focussing  $L^2$ -critical solution of abstract Schredinger equation

$$
i\partial_t u + \Delta u + Au + ||u||^{\frac{4}{n}} u = 0, \ x \in R^n, \ t \in [0, 1]
$$
 (2.12)

which blows up at time  $t = 1$ , where  $U(x, t)$  is a fundamental solution of the Schrödinger equation

$$
i\partial_t u + \Delta u + Au = 0, \ x \in R^n, \ t \in [0,1],
$$

i.e.

$$
U(x,t) = t^{-n/2} \exp\left\{i\left(A+|x|^2\right)/4t\right\}.
$$

By using the above result we wil prove the following main result

Theorem 3. Assume the Condition 1 holds and there exist positive constants  $b_0$  and  $\theta$  such that a solution  $u \in C([-1,1]; X(A))$  of  $(2.10)$  satisfied  $||F(u, \bar{u})|| \le b_0 ||u||^{\theta}$  for  $||u|| > 1$ . Suppose

$$
||u(.,t)||_X = ||u(.,0)||_X = ||u_0||_X = a, \ t \in (-1,1)
$$

and

$$
||u(x,t)|| \le (1-t)^{-\frac{n}{2}} Q\left(\frac{|x|}{1-t}\right), \ t \in (-1,1) \tag{2.14}
$$

where,

$$
Q\left(x\right) = b_1^{-\frac{n}{2}} e^{-b_2 |x|^p}, \, b_1, \, b_2 > 0, \, p > 1; \tag{2.15}
$$

If  $p > p(\theta) = \frac{2(\theta n - 2)}{(\theta n - 1)}$ , then  $a \equiv 0$ .

## 2.2. Some auxiliary results

First of all, we generalize the result G. W. Morgan (see e.g [7]) about Morgan type uncertainty principle for Fourier transform.

**Lemma 2.0.** Let  $f(x) \in L^1(R^n; H) \cap X$  and

$$
\int_{R^n} \int_{R^n} ||f(x)|| \left\| \hat{f}(\xi) \right\| e^{|x \cdot \zeta|} dx d\xi < \infty.
$$

Then  $f(x) \equiv 0$ .

In particular, using Young's inequality this implies: Result 2.1. Let

$$
f(x) \in L^1(R^n; H) \cap X, \ p \in (1, 2), \ \frac{1}{p} + \frac{1}{q} = 1, \alpha, \beta > 0
$$

and

$$
\int_{R^n} \|f(x)\| \, e^{\frac{\alpha^p |x|^p}{p}} dx + \int_{R^n} \left\| \hat{f}(\xi) \right\| \, e^{\frac{\beta^q |\xi|^q}{q}} d\xi < \infty, \, \alpha\beta > 1.
$$

Then  $f(x) \equiv 0$ .

The Morgan type uncertainty principle, in terms of the solution of the free Schredinger equation will be as:

Let

$$
u_{0}\left(.\right)\in L^{1}\left(R^{n};H\right)\cap X\left(A\right)
$$

and

$$
\int_{R^{n}}\left\Vert u_{0}\left(x\right)\right\Vert e^{\frac{\alpha^{p}\left|x\right|^{p}}{p}}dx+\int_{R^{n}}\left\Vert e^{it\left(\Delta+A\right)}u_{0}\left(x\right)\right\Vert e^{\frac{\beta^{q}\left|\xi\right|}{q\left(2t\right)^{q}}}dx<\infty,\ \alpha\beta>1
$$

for some  $t \neq 0$ . Then  $u_0(x) \equiv 0$ .

Let  $L^2(H) = L^2((0, 1) \times R^n; H)$ .

Consider the abstract Schredinger equation

$$
\partial_{t}u = (a+ib) \left[\Delta u + A\left(x\right)u + V\left(x,t\right)u + F\left(x,t\right)\right], \ x \in R^{n}, \ t \in [0,1], \ (2.16)
$$

where a, b are real numbers, A is a linear operator,  $V(x, t)$  is a given potential operator function in H and  $F(x, t)$  is a given H-valued function.

Let  $S = S(t)$  be a symmetric,  $K = K(t)$  be a skew-symmetric operators in *H*,  $f = f(. , t) \in X$  for  $t \in [0, 1]$  and

$$
Q(t) = (f, f)_X, D(t) = (Sf, f)_X, N(t) = D(t) Q^{-1}(t),
$$
  

$$
\partial_t S = S_t \text{ and } |\nabla v|_H^2 = \sum_{k=1}^n \left\| \frac{\partial v}{\partial x_k} \right\|_H^2 \text{ for } v \in W^{1,2}(R^n; H).
$$

Let

$$
\eta(x) = \frac{\gamma a \, |x|^2}{a + 4\gamma (a^2 + b^2) T}, \ L^{1,\infty}(0,T;H) = L^1(0,T;L^{\infty}(R^n;H)).
$$

**Lemma 2.1.** Let the Condition 1 holds. Assume that  $u \in L^{\infty}([0,1]; X(A)) \cap$  $L^2(0,1;Y^1)$  is a solution of  $(2.16)$  for  $a \in \mathbb{R}_+$  and  $b \in \mathbb{R}$ . Then,

$$
e^{-M_T}\left\|e^{\eta(x)}u\left(.,T\right)\right\|_X\leq
$$

$$
\left\|e^{\gamma|x|^2}u(.,0)\right\|_{X} + (a^2+b^2)\left\|e^{\eta(x)}F(.,t)\right\|_{L^1(0,T;X)} + \left\|A^{\frac{1}{2}}u(.,t)\right\|_{X}^2,
$$

uniformly in  $t \in [0, 1]$ , when  $\gamma \geq 0$ ,  $0 \leq T \leq 1$  and

$$
M_T = \left\| a\left(\text{Re}\,V\right)^+ - b\,\text{Im}\,V \right\|_{L^{1,\infty}(0,T;H)}
$$

**Proof.** Set  $v = e^{\varphi}u$ , where  $\varphi$  is a real-valued function to be chosen later. The function  $v = v(x, t)$  verifies

$$
\partial_t v = Sv + Kv + (a + ib) (Vf + e^{\gamma \varphi} F) \text{ in } R^n \times [0, 1], \quad (2.17)
$$

.

where  $S$ ,  $K$  are symmetric and skew-symmetric operator, respectively given by

$$
S = aA1 - ib\gamma B1 + \varphit + a \operatorname{Re} V - b \operatorname{Im} V, \ K = ibA1 - a\gamma B1 +
$$

$$
i (b \operatorname{Re} v + a \operatorname{Im} v), \tag{2.18}
$$

here

$$
A_1 = \Delta + A(x) + \gamma^2 |\nabla \varphi|^2, B_1 = 2\nabla \varphi \cdot \nabla + \Delta \varphi.
$$

Formally,

$$
\partial_t ||v||_X^2 = 2 \operatorname{Re} (Sv, v)_X + 2 \operatorname{Re} ((a + ib) e^{\gamma \varphi} F, v)_X
$$

for  $t \geq 0$ . Again a formal integration by parts gives that

$$
\operatorname{Re}(Sv, v)_X = -a \int_{R^n} |\nabla v|_H^2 dx + a \int_{R^n} (Av, v) dx + a\gamma^2 \int_{R^n} |\nabla \varphi|^2 ||v(x, t)||^2 dx +
$$
  

$$
\int_{R^n} \varphi_t ||v(x, t)||^2 dx + b \operatorname{Im} \int_{R^n} (\bar{v} \nabla \varphi, \nabla v) dx + \int_{R^n} ((a \operatorname{Re} V - b \operatorname{Im} V) v, v) dx.
$$

By Cauchy-Schwarz's inequality and in view of assumption on A, V we get

$$
\partial_{t} ||v||_{X}^{2} \leq 2a \left||A^{\frac{1}{2}}v\right||_{X}^{2} + 2 \left||\left(a\operatorname{Re}V - b\operatorname{Im}V\right)v\right||_{L^{\infty}(L(H))} \left||v\left(.,t\right)\right||_{X}^{2} +
$$
  

$$
2\sqrt{a^{2} + b^{2}} \left||e^{\gamma\varphi}F\left(.,t\right)\right||_{X} \left||v\left(.,t\right)\right||_{X}, \tag{2.19}
$$

when

$$
\left(a + \frac{b^2}{a}\right) |\nabla v|^2 + \varphi_t \le 0, \text{ for } x \in R^n, t \ge 0.
$$
 (2.20)

Let

$$
\phi_r(x) = \begin{cases} |x|^2, & |x| \le r \\ r^2, & |x| > r \end{cases}, \varphi_{\rho,r} = d(t) \theta_\rho * \phi_r(x), \upsilon_{\rho,r} = e^{\varphi_{\rho,r}} u, \quad (2.21)
$$

where  $\theta_{\rho}$  is a radial mollifier and

$$
d(t) = \frac{\gamma a}{a + 4\gamma (a^2 + b^2)t}.
$$

Then by reasoning as in the end of proof  $[8, \text{Lemma } 1]$ , from  $(2.19)$  we obtain that the estimate holds

$$
||v(., T)||_X \le e^{M_T} \left[ \left\| e^{\gamma |x|^2} u(., T) \right\|_X + 2a \left\| A^{\frac{1}{2}} u(., t) \right\|_X^2 + 2\sqrt{a^2 + b^2} \left\| e^{\varphi_{\rho, r}} F \right\|_{L^1(0, T; X)} \right]
$$

holds uniformly in  $\rho$ , r and  $t \in [0,1]$ . Lemma 2.1 follows after letting  $\rho$  tend to zero and  $r$  to infinity.

**Lemma 2.2.** Let  $S = S(x,t)$  be a symmetric operator,  $K = K(x,t)$  be skew-symmetric in Hilbert space  $H$  with independent on  $x$  domain  $D(S)$  and  $D(K)$ , respectively. Assume  $G(. , t) \in L<sup>2</sup>(R<sup>n</sup>)$  is a positive funtion and  $f(x, t)$ is a H−valued reasonable function. Then,

$$
\frac{d^2}{dt^2}Q(t) = 2\partial_t \operatorname{Re}(\partial_t f - Sf - Kf, f)_X + 2\left(S_t f + [S, K] f, f\right)_X +
$$

$$
\|\partial_t f - Sf + Kf\|_X^2 - \|\partial_t f - Sf - Kf\|_X^2
$$

and

$$
\partial_t N(t) \ge Q^{-1}(t) \left[ (S_t f + [S, K] f, f)_X - \frac{1}{2} ||\partial_t f - Sf - Kf||_X^2 \right].
$$

Moreover, if

$$
\|\partial_t f - Sf - Kf\|_H \le M_1 \|f\|_H + G(x, t), \ S_t + [S, K] \ge -M_0
$$

for  $x \in R^N$ ,  $t \in [0,1]$  and

$$
M_2 = \sup_{t \in [0,1]} \|G(. , t)\|_{L^2(R^n)} \|f(. , t)\|_{X}^{-1} < \infty.
$$

Then  $Q(t)$  is logarithmically convex in [0, 1] and there is a constant M such that

$$
Q(t) \le e^{M(M_0 + M_1 + M_2 + M_1^2 + M_2^2)} Q^{1-t} (0) Q^t (1), 0 \le t \le 1.
$$

Proof. It is clear that

$$
\dot{Q}(t) = 2 \operatorname{Re} \left( \frac{d}{dt} f, f \right)_X = 2 \operatorname{Re} \left( \frac{d}{dt} f - Sf - Kf, f \right)_X + 2 \left( Sf, f \right)_X.
$$

Also,

$$
\dot{Q}(t) = \text{Re}\left(\frac{d}{dt}f + Sf, f\right)_X + \text{Re}\left(\frac{d}{dt}f - Sf, f\right)_X,
$$

$$
\frac{d}{dt}D(t) = \frac{1}{2}\text{Re}\left(\frac{d}{dt}f + Sf, f\right)_X - \frac{1}{2}\text{Re}\left(\frac{d}{dt}f - Sf, f\right)_X.
$$

Then by reasoning as in [8, Lemma 2] we obtain the assertion. Consider the following abstract Schredinger equation

$$
\partial_t u = (a + ib) \left[ \Delta u + A \left( x \right) u + V \left( x, t \right) u \right] + F \left( x, t \right), \ x \in R^n, \ t \in [0, 1], \ (2.22)
$$

Let

$$
\Phi(A, V) v = a \operatorname{Re} ((A + V) v, v) - b \operatorname{Im} ((A + V) v, v),
$$
  
for  $v = v(., t) \in D(A)$ .

**Lemma 2.3.** Let the Condition 1 holds. Assume that  $a, \gamma > 0, b \in \mathbb{R}$ . Let  $u \in L^{\infty}(0, 1; X(A)) \cap L^{2}(0, 1; Y^{1})$  be solution of the equation (2.22) and

$$
|\Phi(A, V) u(x, t)| \leq C_0 \eta(x, t) ||u(x, t)||^2
$$

for  $x \in R^n$ ,  $t \in [0,1]$  and  $u \in D(A)$ , where  $\eta$  is a positive function in  $L^1(0,T;L^{\infty}(R^n))$ . Moreover, suppose

$$
\sup_{t\in[0,1]}\left\|V\left(.,t\right)\right\|_{B}\leq M_{1},\ \left\|e^{\gamma|x|^{2}}u\left(.,0\right)\right\|_{X}<\infty,\ \left\|e^{\gamma|x|^{2}}u\left(.,1\right)\right\|_{X}<\infty
$$

and

$$
M_{2}=\sup_{t\in[0,1]}\frac{\left\|e^{\gamma\left|x\right|^{2}}F\left(.,t\right)\right\|_{X}}{\left\|u\right\|_{X}}<\infty.
$$

Then,  $e^{\gamma |x|^2}u(.,t)$  is logarithmically convex in [0, 1] and there is a constant  $N$  such that  $% \left( N_{1},N_{2}\right)$ 

$$
\left\| e^{\gamma |x|^2} u (., t) \right\|_X \le e^{NM(a, b)} \left\| e^{\gamma |x|^2} u (., 0) \right\|_X^{1-t} \left\| e^{\gamma |x|^2} u (., 1) \right\|_X^t \tag{2.23}
$$

where

$$
M(a,b) = (a2 + b2) (\gamma M_12 + M_22) + \sqrt{a2 + b2} (M_1 + M_2)
$$

when  $0 \le t \le 1$ .

**Proof.** Let  $f = e^{\gamma \varphi} u$ , where  $\varphi$  is a real-valued function to be chosen. The function  $f(x, t)$  verifies

$$
\partial_t f = Sf + Kf + (a + ib) (Vf + e^{\gamma \varphi} F) \text{ in } R^n \times [0, 1], \quad (2.24)
$$

where  $S, K$  are symmetric and skew-symmetric operator, respectively given by

$$
S = aA_1 - ib\gamma B_1 + \gamma \varphi_t, \ K = ibA_1 - a\gamma B_1,
$$

where

$$
A_1 = \Delta + A + \gamma^2 |\nabla \varphi|^2, B_1 = 2\nabla \varphi \cdot \nabla + \Delta \varphi.
$$

A calculation shows that

$$
S_t + [S, K] = \gamma \partial_t^2 \varphi + 2\gamma^2 a \nabla \varphi \cdot \nabla \varphi_t - 2ib \gamma (2 \nabla \varphi_t \cdot \nabla + \Delta \varphi_t) - \qquad (2.25)
$$

$$
\gamma (a^2 + b^2) [4 \nabla \cdot (D^2 \varphi \nabla) - 4\gamma^2 D^2 \varphi \nabla \varphi + \Delta^2 \varphi] + 2 [A(x) \nabla \varphi \cdot \nabla - \nabla \varphi \cdot \nabla A].
$$

If we put  $\varphi = |x|^2$ , then (2.25) reduce the following

$$
S_t + [S, K] = -\gamma \left( a^2 + b^2 \right) \left[ 8\Delta - 32\gamma^2 |x|^2 \right] + 2 \left[ A(x) \nabla \varphi \cdot \nabla - \nabla \varphi \cdot \nabla A \right].
$$

Moreover by assumtion (2) of Condition 1,

$$
(S_t f + [S, K] f, f) = \gamma (a^2 + b^2) \int\limits_{R^n} \left( 8 \left\| |\nabla f| \right\|^2 + 32\gamma^2 |x|^2 \left\| f \right\|^2 \right) dx +
$$
  

$$
4 \sum_{k=1}^n \int\limits_{R^n} x_k \left( \left[ A \frac{\partial f}{\partial x_k} - \frac{\partial A}{\partial x_k} f \right], f \right) dx \ge 0 \text{ for } f \in L^\infty \left( 0, T; Y^1 \left( A \right) \right).
$$

This identity, the condition on  $A = A(x)$ ,  $V = V(x, t)$  and (2.24) imply that

$$
S_t + [S, K] \ge 0,
$$
\n
$$
\|\partial_t f - Sf - Kf\|_X \le \sqrt{a^2 + b^2} \left(M_1 \|f\|_X + e^{\gamma \varphi} \|F\|_X\right).
$$
\n(2.26)

If we knew that the quantities and calculations involved in the proof of Lemma 2.2. ( this fact are derived ın a similar way as in [8, Lemma 2]) were finite and correct, when  $f = e^{\gamma |x|^2} u$ , we would have the logarithmic convexity of  $Q(t) = \left\| e^{\gamma |x|^2} u (., t) \right\|_X$  and get (2.23) from Lemma 2.2. To justify the validity of the previous arguments now we consider the function  $\varphi_a(x)$  constructed in [8, Lemma 3], i.e.,

$$
\varphi_a(x) = \begin{cases} |x|^2, & |x| \le 1 \\ (2-a)^{-1} \left( 2|x|^{2-a} - a \right), & |x| \ge 1 \end{cases}
$$

and replace  $\varphi = |x|^2$  by  $\varphi_{a,\rho} = \theta_\rho * \varphi_a$ , where  $a, \rho \in (0,1)$  and  $\theta \in C_0^\infty(\mathbb{R}^n)$ is a radial function. From the proof [8, Lemma 3] we get that  $\varphi_{a,\rho}$  is convex function and

$$
\left\|\Delta\varphi_{a,\rho}\right\|_{\infty} \le C\left(n,\rho\right)a. \tag{2.27}
$$

Put then,  $f_{a,\rho} = e^{\gamma \varphi_{a,\rho}} u$  for  $u \in X$  and  $Q_{a,\rho}(t) = ||f_{a,\rho}(\cdot,t)||_X^2$  $\chi^2$  in Lemma 2.2. The decay bound in Lemma 2.1 and the interior regularity for solutions of (2.23) can now be used qualitatively to make sure that the quantities or calculations

involved in the proof of  $[8, \text{Lemma 2}]$  (with replicing  $L^2$  by X) are finite and correct for  $f_{a,\rho}$ . In this case,  $f_{a,\rho}$  verifies

$$
\partial_{t} f_{a,\rho} = S^{a,\rho} f_{a,\rho} + K^{a,\rho} f_{a,\rho} + (a+ib) \left[ A(x) f_{a,\rho} + e^{\gamma \varphi_{a,\rho}} F(x,t) \right]
$$

with symmetric and skew-symmetric operators  $S^{a,\rho}$  and  $K^{a,\rho}$  given by (2.24) with  $\varphi$  replaced by  $\varphi_{a,\rho}$ . The formula (2.25), the convexity of  $\varphi_{a,\rho}$ , the bounds  $(2.26)$  and  $(2.27)$  imply that the inequalities

$$
S_t^{a,\rho} + [S^{a,\rho}, K^{a,\rho}] \ge 0,
$$
  

$$
\|\partial_t f^{a,\rho} - S f^{a,\rho} - K^{a,\rho} f\|_X \le \sqrt{a^2 + b^2} \left(M_1 \|f^{a,\rho}\|_X + e^{\gamma \varphi_{a,\rho}} \|F\|_X\right).
$$

hold and  $M_2(a, \rho) \leq e^{C(n)\rho^2} M_2$ , when  $a, \rho \in (0, 1)$ . Particularly,  $Q_{a,\rho}$  is logarithmically convex in [0, 1] and

$$
Q_{a,\rho}(t) \le e^{N\left[\left(a^2+b^2\right)\left(M_1^2+M_2^2\right)+\left(M_1+M_2\right)\sqrt{a^2+b^2}\right]} Q_{a,\rho}^{1-t}(0) Q_{a,\rho}^t(1). \tag{2.28}
$$

Then, (2.23) follows after taking first the limit, when a tends to zero in (2.28) and then, when  $\rho$  tends to zero.

# 3. Some properties of solutions of the abstract Schredinger equations

Let

$$
\sigma(t) = [\alpha (1-t) + \beta t]^{-1}, \ \eta(x,t) = (\alpha - \beta)) |x|^2 [4i(\alpha(1-t) + \beta t)]^{-1},
$$

$$
\nu(s) = \left[\gamma\alpha\beta\sigma^2\left(s\right) + \frac{(\alpha-\beta)a}{4\left(a^2+b^2\right)}\sigma\left(s\right)\right], \ \phi\left(x,t\right) = \frac{\gamma a\left|x\right|^2}{a+4\gamma\left(a^2+b^2\right)t}.
$$

Let  $u = u(x, s)$  be a solution of the equation

$$
\partial_s u = i \left[ \Delta u + Au + V(y, s) u + F(y, s) \right], \ y \in R^n, \ s \in [0, 1].
$$

and  $a + ib \neq 0, \gamma \in \mathbb{R}, \alpha, \beta \in \mathbb{R}_+$ . Set

$$
\tilde{u}(x,t) = \left(\sqrt{\alpha\beta}\sigma(t)\right)^{\frac{n}{2}} u\left(\sqrt{\alpha\beta}x\sigma(t), \beta t\sigma(t)\right) e^{\eta}.
$$
 (3.1)

Then,  $\tilde{u}(x, t)$  verifies the equation

$$
\partial_t \tilde{u} = i \left[ \Delta \tilde{u} + A \left( x \right) \tilde{u} + \tilde{V} \left( x, t \right) \tilde{u} + \tilde{F} \left( x, t \right) \right], \ x \in R^n, \ t \in [0, 1] \tag{3.2}
$$

with

$$
\tilde{V}(x,t) = \alpha \beta \sigma^2(t) V\left(\sqrt{\alpha \beta} x \sigma(t), \beta t \sigma(t)\right), \qquad (3.3)
$$

$$
\tilde{F}(x,t) = \left(\sqrt{\alpha\beta}\sigma(t)\right)^{\frac{n}{2}+2} \left(\sqrt{\alpha\beta}x\sigma(t), \beta t\sigma(t)\right). \tag{3.4}
$$

Moreover,

$$
\left\|e^{\gamma|x|^{2}}\tilde{F}(.,t)\right\|_{X} = \alpha\beta\sigma^{2}(t) e^{\nu|y|^{2}} \left\|F(s)\right\|_{X}, \left\|e^{\gamma|x|^{2}}\tilde{u}(.,t)\right\|_{X} = e^{\nu|y|^{2}} \left\|u(s)\right\|_{X}
$$
\n(3.5)

when  $s = \beta t \sigma(t)$ .

**Remark 3.1.** Let  $\beta = \beta (k)$ . By assumption we have

$$
\left\|e^{a_0|x|^p}u(x,0)\right\|_X = a_0,
$$
  

$$
\left\|e^{k|x|^p}u(x,0)\right\|_X = a_k \le a_2e^{2a_1k^{\frac{q}{q-p}}} = a_2e^{2a_1k^{\frac{1}{2-p}}}.
$$
 (3.6)

Thus, for  $\gamma = \gamma(k) \in [0, \infty)$  to be chosen later, one has

$$
\left\|e^{\gamma|x|^{p}}\tilde{u}_{k}\left(x,0\right)\right\|_{X}=\left\|e^{\gamma\left(\frac{\alpha}{\beta}\right)^{p/2}|x|^{p}}u_{k}\left(x,0\right)\right\|_{X}=b_{0},\tag{3.7}
$$

$$
\left\|e^{\gamma|x|^{p}}\tilde{u}_{k}\left(x,1\right)\right\|_{X}=\left\|e^{\gamma\left(\frac{\beta}{\alpha}\right)^{p/2}|x|^{p}}u_{k}\left(x,1\right)\right\|_{X}=a_{k}.
$$

Let we take

$$
\gamma \left(\frac{\alpha}{\beta}\right)^{p/2} = a_0 \text{ and } \gamma \left(\frac{\beta}{\alpha}\right)^{p/2} = k,
$$

i.e.

$$
\gamma = (ka_0)^{\frac{1}{2}}, \ \beta = k^{\frac{1}{p}}, \ \alpha = a_0^{\frac{1}{p}}.
$$
\n(3.8)

Let

$$
M = \int_{0}^{1} \|V(.,t)\|_{L^{\infty}(R^{n};H)} dt = \int_{0}^{1} \|V(.,s)\|_{L^{\infty}(R^{n};H)} ds.
$$

From (3.2), using energy estimates it follows

$$
e^{-M} \|u(.,0)\|_{X} \le \|u(.,t)\|_{X} = \|\tilde{u}(.,s)\|_{X} \le e^{M} \|u(.,0)\|_{X}, \ t,s \in [0,1], \ (3.9)
$$

where

$$
s=\beta t\sigma\left( t\right) .
$$

Consider the following problem

$$
i\partial_t u + \Delta u + A(x) u + V(x, t) u + F(x, t) = 0, \ x \in R^n, \ t \in [0, 1], \qquad (3.10)
$$

$$
u\left( x,0\right) =u_{0}\left( x\right) ,
$$

where  $A = A(x)$  is a linear operator,  $V(x, t)$  is a given potential operator function in a Hilbert space  $H$  and  $F$  is a  $H\operatorname{\!-valued}$  function.

Let as define operator valued integral operators in  $L^p(\Omega; H)$ . Let  $k: R^n \setminus \{0\} \to$  $L(H)$ . We say  $k(x)$  is a  $L(H)$ -valued Calderon-Zygmund kernel  $(C - Z \text{ kernel})$ if  $k \in C^{\infty}(R^n \setminus \{0\}, L(H)), k$  is homogenous of degree  $-n$ , B  $k(x) d\sigma = 0,$ 

where

$$
B = \{ x \in R^n : \ |x| = 1 \}.
$$

For  $f \in L^p(\Omega; H)$ ,  $p \in (1, \infty)$ ,  $a \in L^{\infty}(R^n)$ , and  $x \in \Omega$  we set the Calderon-Zygmund operator

$$
K_{\varepsilon}f = \int\limits_{|x-y|>\varepsilon, y\in\Omega} k(x, y) f(y) dy, \ Kf = \lim_{\varepsilon \to 0} K_{\varepsilon}f
$$

and commutator operator

$$
[K; a] f = a(x) K f(x) - K (af) (x) =
$$
  

$$
\lim_{\varepsilon \to 0} \int_{|x-y| > \varepsilon, y \in \Omega} k(x, y) [a(x) - a(y)] f(y) dy.
$$

By using Calderón's first commutator estimates  $[4]$ , convolution operators on abstract functions [2] and abstract commutator theorem in [20] we obtain the following result:

**Theorem A**<sub>2</sub>. Assume  $k(.)$  is  $L(H)$  –valued  $C-Z$  kernel that have locally integrable first-order derivatives in  $|x| > 0$ , and

$$
||k(x,y) - k(x',y)||_{L(E)} \le M|x-x'||x-y|^{-(n+1)} \text{ for } |x-y| > 2|x-x'|.
$$

Let  $a(.)$  have first-order derivatives in  $L^r(R^n), 1 \lt r \leq \infty$ . Then for  $p, q \in$  $(1, \infty)$ ,  $q^{-1} = p^{-1} + r^{-1}$  the following estimates hold

$$
\left\| [K; a] \, \partial_{x_j} f \right\|_{L^q(R^n;H)} \leq C \, \|f\|_{L^p(R^n;H)},
$$
  

$$
\left\| \partial_{x_j} [K; a] \, f \right\|_{L^q(R^n;H)} \leq C \, \|f\|_{L^p(R^n;H)},
$$

for  $f \in C_0^{\infty} (R^n; H)$ , where the constant  $C > 0$  is independent of f.

Let  $X_{\gamma}$  denote the weithed Lebesque space  $L^2_{\gamma}(R^n:H)$  with  $\gamma(x) = e^{2\lambda x}$ . By following [5, Lemma 2.1] let us show:

**Lemma 3.1.** Assume that the Condition 1 holds and there exists  $\varepsilon_0 > 0$ such that

$$
||V||_{L_t^1 L_x^{\infty}(R^n \times [0,1]; L(H))} < \varepsilon_0.
$$

Moreover, suppose  $u \in C([0,1]; X(A))$  is a stronge solution of  $(3.10)$  with

$$
u_0, u_1 = u(x, 1) \in X_{\gamma}, F \in L^1(0, 1; X_{\gamma})
$$

for  $\gamma(x) = e^{2\lambda x}$  and for some  $\lambda \in R^n$ . Then there exists a positive constant  $M_0 = M_0 (n, A, H)$  independent of  $\lambda$  such that

$$
\sup_{t \in [0,1]} \|u\left(.,t\right)\|_{X_{\gamma}} \le M_0 \left[ \|u_0\|_{X_{\gamma}} + \|u_1\|_{X_{\gamma}} + \int_0^1 \|F\left(.,t\right)\|_{X_{\gamma}} dt \right]. \tag{3.11}
$$

**Proof.** First, we consider the case, when  $\gamma(x) = \beta(x) = e^{2\beta x_1}$ . Without loss of generality we shall assume  $\beta > 0$ . Let  $\varphi_n \in C^{\infty}(\mathbb{R})$  such that  $\varphi_n(\tau) = 1$ ,  $\tau \leq n$  and  $\varphi_n(\tau) = 0$  for  $\tau \geq 10n$  with  $0 \leq \varphi_n \leq 1$ ,  $\left|\varphi_n^{(j)}(\tau)\right| \leq C_j \tau n^{-j}$ . Let

$$
\theta_n(\tau) = \beta \int_0^{\tau} \varphi_n^2(s) \, ds
$$

so that  $\theta_n \in C^{\infty}(\mathbb{R})$  nondecreasing with  $\theta_n(\tau) = \beta\tau$  for  $\tau < n$ ,  $\theta_n(\tau) = C_n\beta$ for  $\tau > 10n$  and

$$
\theta'_{n}(\tau) = \beta \varphi_{n}^{2}(\tau) \leq \beta, \ \theta_{n}^{(j)}(\tau) = \beta C_{j} n^{1-j}, \ j = 1, 2, .... \tag{3.12}
$$

Let  $\phi_n(\tau) = \exp(2\theta_n(\tau))$  so that  $\phi_n(\tau) \leq \exp(2\beta\tau)$  and  $\phi_n(\tau) \to \exp(2\beta\tau)$ for  $n \to \infty$ . Let  $u(x, t)$  be a solution of the equation (3.10), then one gets the equation  $v_n(x,t) = \phi_n(x_1) u(x,t)$  satisfies the following

$$
i\partial_t v_n + \Delta v_n + Av_n = V_n(x, t) v_n + \phi_n(x_1) F(x, t), \qquad (3.13)
$$

where

$$
V_n(x,t) v_n = V(x,t) v_n + 4\beta \varphi_n(x_1) \partial_{x_1} v_n + \left[4\beta \varphi_n(x_1) \varphi_n'(x_1) - 4\beta^2 v_n^4\right] v_n.
$$

Now, we consider a new function

$$
w_n(x,t) = e^{\mu} v_n(x,t), \ \mu = -i4\beta^2 \varphi_n^4(x_1) t.
$$

Then from (3.13) we get

$$
i\partial_t w_n + \Delta w_n + Aw_n = \tilde{V}_n(x,t) w_n + \tilde{F}_n(x,t),
$$

where

$$
\tilde{V}_n(x,t) w_n = V(x,t) w_n + h(x_1,t) + a^2(x_1) \partial_{x_1} w_n + itb(x_1) \partial_{x_1} w_n
$$
  

$$
\tilde{F}_n(x,t) = e^{\mu} \phi_n(x_1) F(x,t),
$$

when

$$
h(x_1, t) = (i16\beta^2 \varphi_n^3 \varphi_n' t)^2 + i48\beta^2 \varphi_n^2 (\varphi_n')^2 t + i16\beta^2 \varphi_n^3 \varphi_n^{(2)} t +
$$

$$
4\beta\varphi_n\varphi_n' + i64\beta^2\varphi_n^3\varphi_n' t, \ a^2 = 4\beta\varphi_n^2(x_1), \ b = -32\beta^2\varphi_n^3\varphi_n'.
$$

It is clear to see that

$$
\|\partial_{x_1}^j h(x_1, t)\|_{L^{\infty}(\mathbb{R} \times [0, 1])} \le C_j n^{-(j+1)}, \ j = 1, 2, ...,
$$
  

$$
a^2(x_1) \ge 0, \ \|\partial_{x_1}^j a(x_1)\|_{L^{\infty}(\mathbb{R})} \le C_j n^{-j}, \ j = 1, 2, ...,
$$
  

$$
\|\partial_{x_1}^j b(x_1)\|_{L^{\infty}(\mathbb{R})} \le C_j n^{-j}, \ j = 1, 2, ....
$$
  
(3.15)

Then by reasoning as in [5, Lemma 2.1] and by using the properties of symmetric operators  $A$  and  $V$ , we obtain

$$
\partial_t \left| \left( P_{\varepsilon} P_+ w_n, v \right) \right|^2 + 2 \operatorname{Im} \left( \Delta P_{\varepsilon} P_+ w_n, v \right) \left( \overline{K} \left( P_{\varepsilon} P_+ w_n, v \right) \right) +
$$

$$
2 \operatorname{Im} (A P_{\varepsilon} P_{+} w_{n}, v) \left( \overline{K} \left( P_{\varepsilon} P_{+} w_{n}, v \right) \right) = 2 \operatorname{Im} (P_{\varepsilon} P_{+} \left( V w_{n} \right), v) \left( \overline{K} \left( P_{\varepsilon} P_{+} w_{n}, v \right) \right) +
$$

$$
2 \operatorname{Im} \left( \left( P_{\varepsilon} P_{+} h w_{n} \right), v \right) \left( \bar{K} \left( P_{\varepsilon} P_{+} w_{n}, v \right) \right) + 2 \operatorname{Im} \left( \left( P_{\varepsilon} P_{+} w_{n}, v \right) \right) \bar{K} \left( P_{\varepsilon} P_{+} \left( \tilde{F}_{n} \right), v \right) +
$$
  

$$
2 \operatorname{Im} \left( \left( P_{\varepsilon} P_{+} a^{2} \left( x_{1} \right) \partial_{x_{1}} w_{n} \right), v \right) \left( \bar{K} \left( P_{\varepsilon} P_{+} w_{n}, v \right) \right) +
$$
  

$$
2 \operatorname{Im} \left( \left( P_{\varepsilon} P_{+} i b \left( x_{1} \right) \partial_{x_{1}} w_{n} \right), v \right) \left( \bar{K} \left( P_{\varepsilon} P_{+} w_{n}, v \right) \right).
$$
 (3.16)

Since for all  $n \in \mathbb{Z}^+$ ,  $w_n(.) \in X$ ,  $\tilde{F}_n(.,t) \in X$  and for a.e.  $t \in [0,1]$  by integrating both sides of  $(3.16)$  on  $\mathbb{R}^n$  we get

$$
\int_{R^n} \text{Im} \left( \Delta P_{\varepsilon} P_+ w_n, v \right) \left( \bar{K} \left( P_{\varepsilon} P_+ w_n, v \right) \right) dx = 0.
$$

It is clear to see that

$$
(u, v)_X = (u(.), v(.)_H)_{L^2(R^n)}
$$
, for  $u, v \in X$ .

Then applying the Cauchy-Schwartz and Holder inequalites for a.e.  $t \in [0, 1]$ we obtain

$$
\int_{R^n} \text{Im} \left( P_{\varepsilon} P_+ V w_n, v \right) \left( \bar{K} \left( P_{\varepsilon} P_+ w_n, v \right) \right) dx \le C ||V||_B ||w_n||_X^2 ||v||_X^2, \qquad (3.17)
$$

$$
\int_{R^n} \text{Im} \left( P_{\varepsilon} P_+ h w_n, v \right) \bar{K} \left( P_{\varepsilon} P_+ w_n, v \right) dx \le C \left\| h \right\|_{L^{\infty}} \left\| w_n \right\|_{X}^{2} \left\| v \right\|_{X}^{2}, \tag{3.18}
$$

$$
\int_{R^n} \text{Im} \left( P_{\varepsilon} P_+ \tilde{F}_n w_n, v \right) \left( \bar{K} \left( P_{\varepsilon} P_+ w_n, v \right) \right) dx \le C \left\| \tilde{F}_n \right\|_X \| w_n \|_X^2 \| v \|_X^2. \tag{3.19}
$$

Moreover, again applying the Cauchy-Schwartz and Holder inequalities due to symmetricity of the operator A, for a.e.  $t \in [0, 1]$  we get

$$
\int_{R^n} \text{Im}\left(A P_{\varepsilon} P_+ w_n, v\right) \left(\bar{K}\left(P_{\varepsilon} P_+ w_n, v\right)\right) dx \le C \left\|Aw_n\right\|_X \left\|w_n\right\|_X \left\|v\right\|_X^2 \tag{3.20}
$$

where, the constant C in  $(3.18) - (3.20)$  is independent of  $v \in C_0^{\infty} (R^n; H)$ ,  $\varepsilon \in (0, 1]$  and  $n \in \mathbb{Z}^+$ . Since  $C_0^{\infty}(R^n; H)$  is dense in X, from  $(3.17)-(3.20)$  in view of operator theory in Hilbert spaces, we obtain the following

$$
\left| \int_{R^n} \text{Im} \left( P_{\varepsilon} P_+ V w_n, \bar{K} P_{\varepsilon} P_+ w_n \right) dx \right| \leq C \| V \|_{B} \| w_n \|_{X}^2,
$$
\n
$$
\left| \int_{R^n} \text{Im} \left( P_{\varepsilon} P_+ h w_n, \bar{K} P_{\varepsilon} P_+ w_n \right) dx \right| \leq C \| h \|_{L^\infty} \| w_n \|_{X}^2 \leq C \frac{1}{n} \| w_n \|_{X}^2,
$$
\n
$$
\left| \int_{R^n} \text{Im} \left( P_{\varepsilon} P_+ \tilde{F}_n w_n, \bar{K} P_{\varepsilon} P_+ w_n \right) dx \right| \leq C \left\| \tilde{F}_n \right\|_{X} \| w_n \|_{X}^2,
$$
\n
$$
\left| \int_{R^n} \text{Im} \left( A P_{\varepsilon} P_+ w_n, \bar{K} P_{\varepsilon} P_+ w_n \right) dx \right| \leq C \| A w_n \|_{X} \| w_n \|_{X}^2
$$
\n
$$
(3.21)
$$

For bounding the last two terms in (3.16) we will use the abstract version of Calderón's first commutator estimates [4]. Really, by Cauchy-Schvartz inequality and in view of Theorem A<sup>2</sup> we get

$$
\left\| ([P_{\pm}; a] \, \partial_{x_1} f, v) \right\|_X \le C \left\| \partial_{x_1} a \right\|_{L^\infty} \left\| f \right\|_X \left\| v \right\|_X, \tag{3.22}
$$

$$
\|\partial_{x_1} ([P_{\pm}; a] f, v)\|_X \le C \|\partial_{x_1} a\|_{L^\infty} \|f\|_X \|v\|_X, \qquad (3.23)
$$

Also, from the calculus of pseudodifferential operators with operator coefficients (see e.g.  $[5]$ ) and the inequality  $(3.15)$ , we have

$$
\left\| ([P_{\varepsilon}; a] \, \partial_{x_1} f, v) \right\|_X \le \frac{C}{n} \left\| f \right\|_X \left\| v \right\|_X, \tag{3.24}
$$

$$
\|\partial_{x_1} ([P_\varepsilon; a] f, v)\|_X \le \frac{C}{n} \|f\|_X \|v\|_X, \qquad (3.25)
$$

where the constant C in  $(3.22) - (3.25)$  is independent of  $\varepsilon \in [0,1]$  and n. We remark that estimates  $(3.22) - (3.25)$  also hold with  $b(x_1)$  replacing  $a(x_1)$ . Since  $C_0^{\infty}(R^n;H)$  is dense in X, from (3.22)-(3.25) in view of operator theory in Hilbert spaces, we obtain

$$
\left\|[P_{\pm};a]\,\partial_{x_1}f\right\|_X\leq C\left\|\partial_{x_1}a\right\|_{L^{\infty}}\left\|f\right\|_X,\ \ \left\|\partial_{x_1}\left[P_{\pm};a\right]f\right\|_X\leq C\left\|\partial_{x_1}a\right\|_{L^{\infty}}\left\|f\right\|_X,
$$

$$
\|([P_{\varepsilon};a]\,\partial_{x_1}f,v)\|_X \leq \frac{C}{n} \|f\|_X \,,\, \|\partial_{x_1}\left([P_{\varepsilon};a]\,f,v\right)\|_X \leq \frac{C}{n} \|f\|_X \,,\qquad(3.26)
$$

and the same estimates (3.26) with  $b(x_1)$  replacing  $a(x_1)$ . By reasoning as in  $[6, Lemma 2.1]$  (claim 1 and 2) from  $(3.26)$  we obtain

$$
\left|\operatorname{Im}\left(\left(P_{\varepsilon}P_{+}a^{2}\left(x_{1}\right)\partial_{x_{1}}w_{n},\bar{K}P_{\varepsilon}P_{+}w_{n}\right)\right)\right|\leq O\left(n^{-1}\left\|w_{n}\right\|_{X}\right),\tag{3.27}
$$

$$
\left|\operatorname{Im}\left(\left(P_{\varepsilon}P_{+}b\left(x_{1}\right)\partial_{x_{1}}w_{n},\bar{K}P_{\varepsilon}P_{+}w_{n}\right)\right)\right|\leq O\left(n^{-1}\left\|w_{n}\right\|_{X}\right).
$$

Now, the estimates (3.21) and (3.27) implay the assertion.

### 4. Proof of Theorem 1

We will apply Lemma 3.1 to a solution of the equation (3.2). Since  $0 < \alpha <$  $\beta = \beta(k)$  for  $k > k_0$  it follows that  $\alpha \leq \sigma(t) \leq \beta$  for any  $t \in [0, 1]$ . Therefore if  $y = \sqrt{\alpha \beta} x \sigma(t)$ , then from (3.8) we get

$$
\sqrt{\alpha \beta^{-1}} |x| \le |y| \sqrt{\alpha^{-1} \beta} |x| = (ka_0^{-1})^{\frac{1}{2p}} |x|
$$
\n(4.1)

Thus,

$$
\left\|\alpha\beta\sigma^{2}\left(t\right)V\left(\sqrt{\alpha\beta}x\sigma\left(t\right),\beta t\sigma\left(t\right)\right)\right\|_{L(H)} \leq \alpha^{-1}\beta\left\|V\right\|_{B} = \left(ka_{0}^{-1}\right)^{\frac{1}{p}}\left\|V\right\|_{B} \tag{4.2}
$$

and so,

$$
\left\| \tilde{V} \left( ., t \right) \right\|_{L^{\infty}(R^n; H)} \leq \left( k a_0^{-1} \right)^{\frac{1}{p}} \left\| V \left( ., t \right) \right\|_{L^{\infty}(R^n; H)}.
$$
\n(4.3)

Also, for  $s = \beta t \sigma(t)$  it is clear that

$$
\frac{ds}{dt} = \alpha \beta \sigma^2(t), dt = (\alpha \beta)^{-1} \sigma^{-2}(t) ds.
$$
 (4.4)

Therefore,

$$
\int_{0}^{1} \left\| \tilde{V} \left( ., t \right) \right\|_{L^{\infty}(R^{n};H)} dt = \int_{0}^{1} \left\| V \left( ., s \right) \right\|_{L^{\infty}(R^{n};H)} ds,
$$

and from (4.1) we get

$$
\int_{0}^{1} \left\| \tilde{V}(.,t) \right\|_{L^{\infty}(|x|>r;H)} dt = \int_{0}^{1} \left\| V(.,s) \right\|_{L^{\infty}(|y|>\varkappa;H)} ds, \tag{4.5}
$$

where

$$
\varkappa = (a_0 k^{-1})^{\frac{1}{2p}} r.
$$

So, if 
$$
\int_{0}^{1} ||V(.,s)||_{L^{\infty}(|y|>\varkappa;H)} ds < \varepsilon_0
$$
 then,  

$$
\int_{0}^{1} ||\tilde{V}(.,t)||_{L^{\infty}(|y|>r;H)} ds < \varepsilon_0, \text{ for } r = \varkappa \left(ka_0^{-1}\right)^{\frac{1}{2p}}
$$

and we can applay Lemma 3.1 to the equation (3.2) with

$$
\tilde{V} = \tilde{V}_{\chi(|x|>r)}(x,t), \ \tilde{F} = \tilde{V}_{\chi(|x|
$$

to get the following estimate

$$
\sup_{t \in [0,1]} \|e^{\nu} \tilde{u}(.,t)\|_{X} \le M_{0} \left( \|e^{\nu} \tilde{u}(.,0)\|_{X} + \|e^{\nu} \tilde{u}(.,1)\|_{X} \right) + M_{0} e^{M} e^{\nu_{0}} \left\| \tilde{V} \right\|_{B} \|u(.,0)\|_{X}, \tag{4.6}
$$

where  $M$  a positive constant defined in Remark 2.1 and

$$
B=L^\infty\left(R^n\times[0,1]\,;\,L\left(H\right)\right),\,\nu=\left(2p\right)^{\frac{1}{p}}\gamma^{\frac{1}{p}}\lambda.\frac{x}{2},\,\,\nu_0=\left|\lambda\right|\left(2p\right)^{\frac{1}{p}}\gamma^{\frac{1}{p}}\frac{r}{2}.
$$

From (4.6) we have

$$
\sup_{t \in [0,1]} \int_{R^n} \|e^{\nu} \tilde{u}(.,t)\|_{H}^2 dx \le M_0 \int_{R^n} e^{\nu} \left( \|\tilde{u}(.,0)\|_{H}^2 + \|\tilde{u}(.,1)\|_{H}^2 \right) dx +
$$
  

$$
M_0 e^M e^{|\lambda|(2p)^{\frac{1}{p}} \gamma^{\frac{1}{p}r}} \left\| \tilde{V} \right\|_{B} \|u(.,0)\|_{X}^2,
$$

and multiply the above inequality by  $e^{|\lambda|/q} |\lambda|^{n(q-2)/2}$ , integrate in  $\lambda$  and in x, use Fubini theorem and the following formula

$$
e^{\gamma |x|^p/p} \approx \int_{R^n} e^{\gamma^{\frac{1}{p}} \lambda x - |\lambda|^{q/q}} |\lambda|^{n(q-2)/2} d\lambda,
$$
 (4.7)

proven in [7, Appendx] to obtain

$$
\int\limits_{|x|>1}e^{2\gamma|x|^{p}}\left\|\tilde{u}\left(.,t\right)\right\|_{H}^{2}dx\leq M_{0}\int\limits_{R^{n}}e^{2\gamma|x|^{p}}\left(\left\|\tilde{u}\left(.,0\right)\right\|_{H}^{2}+\left\|\tilde{u}\left(.,1\right)\right\|_{H}^{2}\right)dx+
$$

$$
M_0 e^{2M} e^{2\gamma r^p} r^{C_p} \left\| \tilde{V} \right\|_B \| u \left( ., 0 \right) \|^2_X. \tag{4.8}
$$

Hence, the esimates (3.6), (3.8), (3.9), (4.3) and (4.8) imply

$$
\sup_{t \in [0,1]} \left\| e^{\gamma |x|^p} \tilde{u} \left( ., t \right) \right\|_X \leq M_0 \left( \left\| e^{\gamma |x|^p} \tilde{u} \left( ., 0 \right) \right\|_X + \left\| e^{\gamma |x|^p} \tilde{u} \left( ., 1 \right) \right\|_X \right) +
$$
  

$$
M_0 e^M e^\gamma \left\| u \left( ., 0 \right) \right\|_X + M_0 \left( k a_0^{-1} \right)^{C_p} e^M e^{k^p \gamma^{\frac{1}{2}} \left( k a_0^{-1} \right)} \left\| u \left( ., 0 \right) \right\|_X \left\| V \right\|_B \leq (4.9)
$$
  

$$
M_0 \left( a_0 + a_k \right) + M_0 e^M \left\| u \left( ., 0 \right) \right\|_X \left( e^\gamma + \left( k a_0^{-1} \right)^{C_p} \left\| V \right\|_B \right) e^{k^p} \leq
$$
  

$$
M_0 a_k = M_0 e^{a_1 k^{1/2 - p}} \text{ for } k > k_0 \left( M_0 \right) \text{ sufficiently large.}
$$

Next, we shall obtain bounds for the  $\nabla \tilde{u}$ . Let  $\tilde{\gamma} = \frac{\gamma}{2}$  and  $\varphi$  be a strictly convex complex valued function on compact sets of  $R<sup>n</sup>$ , radial such that (see [7])  $(n-2)$ 

$$
D^2 \varphi \ge p(p-1)|x|^{(p-2)}, \text{ for } |x| \ge 1,
$$
  

$$
\varphi \ge 0, \quad \|\partial^{\alpha}\varphi\|_{L^{\infty}} \le C, \quad 2 \le |\alpha| \le 4, \quad \|\partial^{\alpha}\varphi\|_{L^{\infty}(|x|<2)} \le C \text{ for } |\alpha| \le 4,
$$
  

$$
\varphi(x) = |x|^p + O(|x|), \text{ for } |x| > 1.
$$

Let us consider the equation

$$
\partial_t v = i \left( \Delta v + Av + F(x, t) \right), \ x \in R^n, \ t \in [0, 1], \tag{4.10}
$$

where  $F(x,t) = \tilde{V}v$ , A is a symmetric operator in H and  $\tilde{V}$  is a operator in H defined by  $(3.3)$ .

Let

$$
f(x,t) = e^{\tilde{\gamma}\varphi} v(x,t), \ Q(t) = (f(x,t), f(x,t))_H,
$$

where  $v$  is a solution of (4.10). Then, by reasoning as in Lemma 2.3 we have

$$
\partial_t f = Sf + Kf + i \left[ A + e^{\tilde{\gamma}\varphi} F \right], \ (x, t) \in R^n \times [0, 1], \tag{4.11}
$$

here S, K are symmetric and skew-symmetric operator, respectively given by

$$
S = -i\tilde{\gamma} (2\nabla\varphi \cdot \nabla + \Delta\varphi) , K = i\left(\Delta + A + \tilde{\gamma}^2 |\nabla\varphi|^2\right). \tag{4.12}
$$

Let

$$
[S, K] = SK - KS.
$$

A calculation shows that,

$$
SK = \tilde{\gamma} (2\nabla\varphi \cdot \nabla + \Delta\varphi) \left( \Delta + A + \tilde{\gamma}^2 |\nabla\varphi|^2 \right) = \tilde{\gamma} (2\nabla\varphi \cdot \nabla + \Delta\varphi) \Delta +
$$

$$
\tilde{\gamma} (2\nabla\varphi \cdot \nabla + \Delta\varphi) A + \tilde{\gamma}^3 |\nabla\varphi|^2 (2\nabla\varphi \cdot \nabla + \Delta\varphi),
$$

$$
KS = \tilde{\gamma} [\Delta (2\nabla\varphi \cdot \nabla + \Delta\varphi) + A (2\nabla\varphi \cdot \nabla + \Delta\varphi)] + \tilde{\gamma}^3 |\nabla\varphi|^2 (2\nabla\varphi \cdot \nabla + \Delta\varphi),
$$

$$
[S,K] = \tilde{\gamma} \left[ (2\nabla \varphi \cdot \nabla + \Delta \varphi) \Delta - \Delta (2\nabla \varphi \cdot \nabla + \Delta \varphi) \right] + 2\tilde{\gamma} \left( \nabla \varphi \cdot \nabla A - A \nabla \varphi \cdot \nabla \right),
$$

$$
S_t + [S, K] = -2i\tilde{\gamma} (\nabla \varphi \cdot \partial_t \nabla + \Delta \varphi \partial_t) + \tag{4.13}
$$

 $\tilde{\gamma}\left[\left(2\nabla\varphi.\nabla+\Delta\varphi\right)\Delta-\Delta\left(2\nabla\varphi.\nabla+\Delta\varphi\right)\right]+2\tilde{\gamma}\left(\nabla\varphi.\nabla A-A\nabla\varphi.\nabla\right).$ 

By Lemma 2.2

$$
Q^{''}(t) = 2\partial_t \operatorname{Re} (\partial_t f - Sf - Kf, f)_X + 2(S_t f + [S, K] f, f)_X +
$$

$$
\|\partial_t f - Sf + Kf\|_X^2 - \|\partial_t f - Sf - Kf\|_X^2, \tag{4.14}
$$

so,

$$
Q^{''}(t) \ge 2\partial_t \operatorname{Re} (\partial_t f - Sf - Kf, f)_X + 2(S_t f + [S, K] f, f)_X. \tag{4.15}
$$

Multiplying (4.15) by  $t(1-t)$  and integrating in t we obtain

$$
\int_{0}^{1} t(1-t) (S_{t}f + [S, K] f, f)_{X} dt \le
$$
  

$$
M_{0} \left[ \sup_{t \in [0,1]} \left\| e^{\tilde{\gamma}\varphi} \nu(.)t \right\|_{X} + \sup_{t \in [0,1]} \left\| e^{\tilde{\gamma}\varphi} F(. , t) \right\|_{X} \right].
$$

This computation can be justified by parabolic regularization using the fact that we already know the decay estimate for the solution of (4.10). Hence, combining  $(3.8)$ ,  $(4.3)$  and  $(4.9)$  it follows that

$$
\tilde{\gamma} \int_{0}^{1} \int_{R^{n}} t(1-t)D^{2}\varphi(x,t) \left(\nabla f, \nabla f\right)_{H} dxdt + \tilde{\gamma}^{3} \int_{0}^{1} \int_{R^{n}} t(1-t)D^{2}\varphi(x,t) \left(\nabla f, \nabla f\right)_{H} dxdt \leq
$$
  

$$
M_{0} \left[ \sup_{t \in [0,1]} \left\| e^{\tilde{\gamma}\varphi} \nu\left(.,t\right) \right\|_{X} \left(1 + \left\| \tilde{V}\left(.,t\right) \right\|_{L^{\infty}(R^{n};L(H))}\right) + \tilde{\gamma} \sup_{t \in [0,1]} \left\| e^{\tilde{\gamma}\varphi} \nu\left(.,t\right) \right\|_{X} \right] \leq
$$
  

$$
M_{0}k^{C_{p}} a_{k}.
$$
 (4.16)

It is clear to see that

$$
\nabla f = \tilde{\gamma} v e^{\tilde{\gamma}\varphi} \nabla \varphi + e^{\tilde{\gamma}\varphi} \nabla v.
$$

So, by using the properties of  $\varphi$  we get

$$
\left|e^{2\tilde{\gamma}\varphi}D^2\varphi\left|\nabla\varphi\right|^2\right|\leq C_p e^{3\gamma\varphi|2}.
$$

From here, we can conclude that

$$
\gamma \int_{0}^{1} \int_{R^{n}} t(1-t) (1+|x|)^{p-2} \|\nabla v(x,t)\|_{H} e^{\gamma |x|^{p}} dxdt + \sup_{t \in [0,1]} \left\| e^{\frac{\gamma |x|^{p}}{2}} v(.,t) \right\|_{X} \leq C_{0} k^{C_{p}} a_{k}^{2} = C_{0} k^{C_{p}} e^{a(k,p)} \tag{4.17}
$$

for  $k \geq k_0(M_0)$  sufficiently large, where

$$
a(k,p) = C_{\mu} M_0 k^{C_p} e^{2a_1 k^{1/(2-p)}}.
$$

For proving Theorem 1 first, we deduce the following estimate

$$
\int_{|x| < \frac{R}{2}} \int_{\nu_1}^{\nu_2} \left\| \tilde{u}(x, t) \right\|_{H} dt dx \ge C_{\nu} e^{-M} \left\| u(., 0) \right\|_{X},\tag{4.18}
$$

for r sufficiently large,  $\nu_1, \nu_2 \in (0,1)$ ,  $\nu_1 < \nu_2$  and  $\nu = (\nu_1, \nu_2)$ . From (3.1) by using the change of variables  $s = \beta t \sigma(t)$  and  $y = \sqrt{\alpha \beta x \sigma(t)}$  we get

$$
\int_{|x| < \frac{r}{2}} \int_{\nu_1}^{\nu_2} \left\| \tilde{u}(x, t) \right\|_{H}^{2} dt dx = \tag{4.19}
$$

$$
(\alpha\beta)^{\frac{n}{2}}\int\limits_{|x|<\frac{R}{2}}\int\limits_{\nu_1}^{\nu_2}|\sigma\left(t\right)|^n\left\|u\left(\sqrt{\alpha\beta}x\sigma\left(t\right),\beta t\sigma\left(t\right)\right)\right\|_{H}^2dtdx \geq
$$
  

$$
M_0\frac{\beta}{\alpha}\int\limits_{|y|
$$

for  $k > M_0$ ,  $s\nu_1 > \frac{1}{2}$  and  $r_0 = r \left( k a_0^{-1} \right)^{\frac{1}{2p}}$ . Thus, taking

$$
r > \omega \left(ka_0^{-1}\right)^{\frac{1}{2p}} \tag{4.20}
$$

with  $\omega = \omega(u)$  a constant to be determined, it follows that

$$
\Phi \ge M_0 \frac{\beta}{\alpha} \int\limits_{|y|<\omega} \int\limits_{s\nu_1}^{s\nu_2} \|u(y,s)\|_H^2 ds dy,
$$

where the interval  $I = I_k = [s\nu_1, s\nu_2]$  satisfies  $I \subset [1/2, 1]$  for k sufficiently large. Moreover, given  $\varepsilon > 0$  there exists  $k_0(\varepsilon) > 0$  such that for any  $k \geq k_0$ one has that  $I_k \subset [1-\varepsilon,1]$ . By hypothesis on  $u(x,t)$ , i.e. the continuity of  $||u(\cdot, s)||_X$  at  $s = 1$ , it follows that there exists  $\omega > 1$  and  $K_0 = K_0(u)$  such that for any  $k \geq K_0$  and for any  $s \in I_k$ 

$$
\int_{|y|<\omega} \|u(y,s)\|_H^2 dy \ge C_{\nu} e^{-M} \|u(.,0)\|_X,
$$

which yields the desired result. Next, we deduce the following estimate

$$
\int_{|x| < R} \int_{\mu_1}^{\mu_2} \left( \left\| \tilde{u}(x,t) \right\|_{H}^2 + \left\| \nabla \tilde{u}(x,t) \right\|_{H}^2 + \left\| A\tilde{u}(x,t) \right\|_{H}^2 \right) dt dx \leq \tag{4.21}
$$

$$
C_{n\mu}C_0k^{C_p}e^{a(k,p)},
$$

for r sufficiently large,  $\nu_1, \nu_2 \in (0, 1), \mu_1 = \frac{(\nu_2 - \nu_1)}{8}, \mu_2 = 1 - \mu_1, \mu_1 < \mu_2$  and  $\mu = (\mu_1, \mu_2)$ . Indeed, from (3.9) and (4.17) we obtain

$$
\int_{|x|
$$

$$
\int_{\mu_1}^{\mu_2} \int_{|x| < R} \|\nabla \tilde{u}(x, t)\|_{H} \, dt dx \le C_{n\mu} \int_{\mu_1}^{\mu_2} \int_{|x| < R} t(1 - t) \, \|\nabla v(x, t)\|_{H} \, e^{\gamma |x|^{p}} dt dx \le (4.22)
$$

$$
C_{\mu} \gamma^{-1} r^{2-p} C_0 k^{C_p} A_k^2 \le C_{\mu} C_0 k^{C_p} e^{2a_1 k^{(2-p)^{-1}}}.
$$

Hence, from (4.22) we get (4.21) for  $k \geq k_0(C_0)$  sufficiently large.

Let  $Y = L^2(R^n \times [0,1]; H)$ . By reasoning as in [6, Lemma 3.1] we obtain Lemma 4.1. Assume the assumptıions (1) and (3) of Condition 1 are satisfied. Suppose that  $r > 0$  and  $\varphi : [0, 1] \to \mathbb{R}$  is a smooth function. Then, there exists  $C = C(n, \varphi, H, A) > 0$  such that, the inequality

$$
\sum_{k=1}^{n} x_k \left[ \left( \frac{\partial A}{\partial x_k} g, g \right) + \left( \frac{\partial g}{\partial x_k}, Ag \right)_X \right] + \frac{\varkappa^{\frac{3}{2}}}{r^2} \left\| e^{\varkappa |\psi|} g \right\|_Y \le C \left\| e^{\varkappa |\psi|} i \left( \partial_t g + \Delta g + Ag \right) \right\|_Y
$$

holds, for  $\varkappa \geq CR^2$  and  $g \in C_0^{\infty} (R^{n+1}; H(A))$  with support contained in the set

$$
\left\{x, t: \ |\psi(x, t)| = \left|\frac{x}{r} + \varphi(t) e_1\right| \ge 1\right\}.
$$

**Proof.** Let  $f = e^{\alpha |\psi(x,t)|^2} g$ . Then, by acts of Schredinger operator  $(i\partial_t + \Delta + A)$ to  $f \in X(A)$  we get

$$
e^{\alpha |\psi(x,t)|^2} \left( (i\partial_t g + \Delta g + Ag) \right) = S_\alpha f - 4\alpha K_\alpha f,
$$

where

$$
S_{\alpha} = (i\partial_t + \Delta + A) + \frac{4\alpha^2}{r^2} \left| \psi(x, t) \right|^2,
$$

$$
K_{\alpha} = \frac{1}{r} \left( \frac{x}{r} + \varphi(t) e_1 \right) . \nabla + \frac{n}{r^2} + \frac{i \varphi'}{2} \left( \frac{x_1}{r} + \varphi(t) \right) .
$$

Hence,

$$
(S_{\alpha})^* = S_{\alpha}, (K_{\alpha})^* = K_{\alpha}
$$

and

$$
\left\|e^{\alpha |\psi(x,t)|^2} \left(i\partial_t g + \Delta g + Ag\right)\right\|_X^2 = \left(S_\alpha f - 4\alpha K_\alpha f, S_\alpha f - 4\alpha K_\alpha f\right)_X \ge
$$

$$
-4\alpha \left(\left(S_\alpha K_\alpha - K_\alpha S_\alpha\right) f, f\right)_X = -4\alpha \left(\left[S_\alpha, K_\alpha\right] f, f\right)_X.
$$

A calculation shows that

$$
[S_{\alpha}, K_{\alpha}] = \frac{2}{r^2} \Delta - \frac{4\alpha^2}{r^4} \left| \frac{x}{r} + \varphi e_1 \right|^2 - \frac{1}{2} + \frac{2i\varphi'}{r} \partial_{x_1} + [A, K_{\alpha}], \tag{4.23}
$$

where

$$
[A, K_{\alpha}] f = \left[ \left( \frac{x_1}{r} + \varphi \right) A \frac{\partial f}{\partial x_1} + \frac{1}{r} \sum_{k=2}^n x_k A \frac{\partial f}{\partial x_k} \right] -
$$

$$
\left( \frac{x_1}{r} + \varphi \right) \left( \frac{\partial A}{\partial x_1} f + A \frac{\partial f}{\partial x_1} \right) + \frac{1}{r} \sum_{k=2}^n x_k \left( \frac{\partial A}{\partial x_k} f + A \frac{\partial f}{\partial x_k} \right).
$$

Since  $A$  is a symmetric operator in  $H$ , from  $(4.23)$  we have

$$
\left\|e^{\alpha|\psi(x,t)|^2} (i\partial_t g + \Delta g + Ag)\right\|_X^2 \geq \qquad (4.24)
$$

$$
\frac{16\alpha^3}{r^4} \int \left|\frac{x}{r} + \varphi e_1\right|^2 \left\|f(x,t)\right\|^2 dxdt + \frac{8\alpha}{r^2} \int \left\|\nabla f(x,t)\right\|^2 dxdt +
$$

$$
2\alpha \int \left[\left(\frac{x_1}{r} + \varphi\right)\varphi'' + (\varphi')^2\right] \left\|f(x,t)\right\|^2 dxdt - \frac{8\alpha}{r} \int \varphi'\left(\partial_{x_{11}} f, \bar{f}\right) dxdt +
$$

$$
C \sum_{k=1}^n x_k \left[\left(\frac{\partial A}{\partial x_k} f, f\right) + \left(\frac{\partial f}{\partial x_k}, Af\right)_X\right].
$$

In view of the hypothesis on  $g$  and the Cauchy–Schwarz inequality, the absolute value of the third fourth terms in (4.23) can be bounded by a fraction of the first two terms on the right-hand side of (4.24), when  $\alpha > Cr^2$  for some large C depending on  $\|\varphi'\|_{\infty} + \|\varphi''\|_{\infty}$ . Moreover, by using the assumption on  $A = A(x)$ , we get that the last two terms are nonnegative. This yields the assertion. Now, from (3.3) we have

$$
\left\|\tilde{V}(x,t)\right\|_{H} \leq \frac{\alpha}{\beta}\mu_1^{-2} \left\|V\right\|_{B} \leq \mu_1^{-2} a_0^{\frac{1}{p}} k^{-\frac{1}{p}} \left\|V\right\|_{B}.
$$

Then from (4.20) we get

$$
\left\| \tilde{V} \right\|_{L^{\infty}(R^n \times [\mu_1, \mu_2]; L(H))} < R. \tag{4.25}
$$

Define

$$
\delta^2\left(r\right) =
$$

$$
\int_{\mu_{1}}^{\mu_{2}} \int_{R-1 \leq |x| \leq r} \left( \left\| \tilde{u}\left(x,t\right) \right\|_{H}^{2} + \left\| \nabla \tilde{u}\left(x,t\right) \right\|_{H}^{2} + \left\| A\tilde{u}\left(x,t\right) \right\|_{H}^{2} \right) dt dx.
$$

Let  $\nu_1, \nu_2 \in (0,1), \nu_1 < \nu_2$  and  $\nu_2 < 2\nu_1$ . We choose  $\varphi \in C^{\infty}([0,1])$  and  $\theta$ ,  $\theta_r \in C_0^{\infty} (R^n)$  satisfying

$$
0 \le \varphi(t) \le 3, \, \varphi(t) = 3 \text{ for } t \in [\nu_1, \nu_2], \, \varphi(t) = 0 \text{ for}
$$
\n
$$
t \in [0, \nu_2 - \nu_1] \cup \left[ \nu_2 + \frac{\nu_2 - \nu_1}{2}, 1 \right],
$$
\n
$$
\theta_r(x) = 1 \text{ for } |x| < r - 1, \, \theta_R(x) = 0, \text{ for } |x| > r,
$$

and

$$
\theta(x) = 1
$$
 for  $|x| < 1$ ,  $\theta(x) = 0$ , for  $|x| \ge 2$ .

Let

$$
g(x,t) = \theta_r(x) \theta(\psi(x,t)) \tilde{u}(x,t), \qquad (4.26)
$$

where  $\tilde{u}(x, t)$  is a solution of  $(3.2)$  when  $\tilde{V} = \tilde{F} \equiv 0$ . It is clear to see that

$$
|\psi(x,t)| \ge \frac{5}{2}
$$
 for  $|x| < \frac{r}{2}$  and  $t \in [\nu_1, \nu_2]$ .

Hence,

$$
g(x,t) = \tilde{u}(x,t)
$$
 and  $e^{\varkappa |\psi(x,t)|^2} \ge e^{\frac{25}{4}\varkappa}$  for  $|x| < \frac{r}{2}$ ,  $t \in [\nu_1, \nu_2]$ .

Moreover, from (4.26) also we get that

$$
g(x,t) = 0
$$
 for  $|x| \ge r$  or  $t \in [0, \nu_2 - \nu_1] \cup \left[ \nu_2 + \frac{\nu_2 - \nu_1}{2}, 1 \right]$ ,

so

$$
\text{supp } g \subset \{|x| \le R\} \times \left[\nu_2 - \nu_1, \nu_2 + \frac{\nu_2 - \nu_1}{2}\right] \cap \{|b(x, t)| \ge 1\}.
$$

Then, for  $\xi = \psi(x, t)$  we have

$$
\left(i\partial_t + \Delta + A + \tilde{V}\right)g = \left[\theta\left(\xi\right)\left(2\nabla\theta_r\left(x\right).\tilde{u} + \tilde{u}\Delta\theta_r\left(x\right)\right) + 2\nabla\theta\left(\xi\right).\nabla\theta_r\tilde{u}\right] +
$$
  

$$
\theta_r\left(x\right)\left[2r^{-1}\nabla\theta\left(\xi\right).\nabla\tilde{u} + r^{-2}\tilde{u}\Delta\theta\left(\xi\right) + i\varphi'\partial x_1\theta\left(\xi\right)u\right] = B_1 + B_2.
$$

Note that,

$$
supp B_1 \subset \{(x, t): \ r - 1 \le |x| \le r, \ \mu_1 \le t \le \mu_2\}
$$

and

$$
supp B_2 \subset \{(x, t) \in R^n \times [0, 1], 1 \leq |\psi(x, t)| \leq 2\}.
$$

Now applying Lemma 4.1 choosing  $\varkappa = d_n^2 R^2$ ,  $d_n^2 \ge ||\varphi'||_{\infty} + ||\varphi''||_{\infty}$  it follows that

$$
R \left\| e^{\varkappa |\psi|^2} g \right\|_{Y} \leq C \left\| e^{\varkappa |\psi|^2} i \left( \partial_{t} g + \Delta g + A g \right) \right\|_{Y} \leq
$$
\n
$$
C \left[ \left\| e^{\varkappa |\psi|^2} \tilde{V} g \right\|_{Y} + \left\| e^{\varkappa |\psi|^2} B_1 \right\|_{Y} + \left\| e^{\varkappa |\psi|^2} B_2 \right\|_{Y} \right] =
$$
\n
$$
D_1 + D_2 + D_3.
$$
\n(4.27)

.

Since

$$
\Big\|\tilde V\Big\|_{L^\infty(R^n\times [\mu_1,\mu_2];L(H))}
$$

 $D_1$  can be absorbed in the left hand side of (4.27). Moreover,  $|\psi(x, t)| \leq 4$  on the support of  $B_1$ , thus

$$
D_2 \leq C\delta(R) e^{16\varkappa}
$$

Let

$$
R_\mu^n = \{(x, t) : |x| \le r, \ \mu_1 \le t \le \mu_2\}
$$

Then  $R_{\mu}^{n} \subset \text{supp } B_2$ , and  $1 \leq |\psi(x, t)| \leq 2$ , so

$$
D_3 \leq Ce^{4\varkappa} \|\tilde{u} + \nabla \tilde{u}\|_{L^2(R^n \times [\mu_1, \mu_2]; H)}.
$$

By using  $(4.18)$  and  $(4.22)$  we have

$$
C_{\mu}e^{-M}e^{\frac{25}{4}\varkappa}\|u\left(.,0\right)\|_{X}\leq re^{\frac{25}{4}\varkappa}\left[\int_{|x|<\frac{R}{2}}^{\nu_{2}}\|\tilde{u}\left(x,t\right)\|_{H}dtdx\right]^{\frac{1}{2}}\leq\qquad(4.28)
$$

 $C_\mu \delta(r) \, e^{\frac{25}{4}\varkappa} + C_\mu e^{4\varkappa} \, \|\tilde u + \nabla \tilde u\|_{L^2\left(R_\mu^n;H\right)} \leq C_\mu \delta\left(r\right) e^{16\varkappa} + C_\mu C_0 k^{C_p} e^{4\varkappa} e^{2a_1 k^{(2-p)^{-1}}}.$ 

Puting  $\varkappa = d_n r^2 = 2a_1 k^{\frac{1}{2-p}}$  it follows from (4.28) that, if  $||u(.,0)||_X \neq 0$  then

$$
\delta(r) \ge C_{\mu} \|u(.,0)\|_{X} e^{-(M+10\varkappa)} = C_{\mu} \|u(.,0)\|_{X} e^{-(M+20)a_{1}k^{\frac{1}{2-p}}}
$$
(4.29)

for  $k \geq k_0(C_\mu)$  sufficiently large. Now, by (4.22) we get

$$
\delta^{2}(r) = \int_{\mu_{1}}^{\mu_{2}} \int_{r-1 \leq |x| \leq r} \left( \left\| \tilde{u}(x,t) \right\|_{H}^{2} + \left\| \nabla \tilde{u}(x,t) \right\|_{H}^{2} + \left\| A\tilde{u}(x,t) \right\|_{H}^{2} \right) dt dx \leq
$$

$$
\int_{\mu_{1}}^{\mu_{2}} \int_{r-1 \leq |x| \leq r} \left( \left\| \tilde{u}(x,t) \right\|_{H}^{2} + \left\| A\tilde{u}(x,t) \right\|_{H}^{2} \right) dt dx +
$$

$$
C_{\mu} \int_{\mu_1}^{\mu_2} \int_{r-1 \le |x| \le r} t (1-t) \|\nabla \tilde{u}(x,t)\|_{H}^{2} dt dx \le
$$
  
\n
$$
C_{\mu} e^{-\gamma (r-1)^{p}} \sup_{t \in [0,1]} \left\| e^{\gamma |x|^{p/2}} \tilde{u}(x,t) \right\|_{X}^{2} + C_{\mu} \gamma^{-1} r^{2-p} e^{-\gamma (r-1)^{p}} \times
$$
  
\n
$$
\int_{\mu_1}^{\mu_2} \int_{r-1 \le |x| \le r} \left[ \frac{t (1-t)}{(1+|x|)^{2/p}} \|\nabla \tilde{u}(x,t)\|_{H}^{2} + \|A\tilde{u}(x,t)\|_{H}^{2} \right] dt dx \le (4.30)
$$
  
\n
$$
C_{\mu} \gamma^{-1} k^{C_{p}} e^{\eta(p)}, \ \eta(p) = 2a_{1} k^{(2-p)^{-1}} - \gamma (r-1)^{p} .
$$

The estimates  $(4.28) - (4.30)$  imply

$$
C_{\mu}e^{-2M}e^{\frac{25}{4}\varkappa} \|u(.,0)\|_{X} \leq C_{0}k^{C_{p}}e^{\omega(p)} + O\left(k^{1/2(2-p)}\right), \qquad (4.31)
$$

where

$$
\omega(p) = 42a_1k^{1/(2-p)} - a_0^{-\frac{1}{2}} \left(\frac{2a_1}{d_n}\right)^{\frac{p}{2}} k^{1/(2-p)}.
$$

Hence, if  $42a_1 < \sqrt{a_1^p a_0} \left(\frac{2}{d_n}\right)^{\frac{p}{2}}$  by letting k tends to infinity it follows from (4.31) that  $||u (., 0)||_X = 0$ , which gives  $u (x, t) \equiv 0$ .

Proof of Corollary 1. Since

$$
\int\limits_{R^{n}}\left\Vert u\left( x,1\right) \right\Vert _{H}^{2}e^{2b\left\vert x\right\vert ^{q}}dx<\infty\text{ for }b=\frac{\beta^{q}}{q}
$$

one has that

$$
\int_{R^{n}}\left\|u\left(x,1\right)\right\|_{H}^{2}e^{2k|x|^{q}}dx \leq \left\|e^{2k|x|^{q}-2b|x|^{q}}\right\|_{\infty}\int_{R^{n}}\left\|u\left(x,1\right)\right\|_{H}^{2}e^{2b|x|^{q}}dx.
$$

Then, by reasoning as in [7, Corollary 1] we obtain the assertion. Proof of Theorem 2. Indeed, just applying Corollary 1 with

$$
u(x,t) = u_1(x,t) - u_2(x,t)
$$

and

$$
V(x,t) = \frac{F(u_1, \bar{u}_1) - F(u_2, \bar{u}_2)}{u_1 - u_2}
$$

we obtain the assertion of Theorem 2.

# 5. Proof of Theorem 3.

First, we deduce the corresponding upper bounds. Assume

$$
||u\left( .,t\right) ||_{X}=a\neq 0.
$$

Fix  $\bar{t} \in (0,1)$  near 1, and let

$$
\upsilon\left(x,t\right)=u\left(x,t-1+\bar{t}\right),\,t\in\left[0,1\right]
$$

which satisfies the equation (2.10) with

$$
|v(x,0)| \le \frac{b_1}{(2-\bar{t})^{n/2}} e^{-\frac{b_2|x|^p}{(2-\bar{t})^p}}, \ \ |v(x,1)| \le \frac{b_1}{(1-\bar{t})^{n/2}} e^{-\frac{b_2|x|^p}{(1-\bar{t})^p}} \tag{5.1}
$$

where  $A$  is a linear operator,  $V(x, t)$  is a given potential operator function in a Hilbert space H.

From  $(5.1)$  we get

$$
\int_{R^n} \|v(x,0)\|_H^2 e^{A_0|x|^q} dx = a_0^2, \quad \int_{R^n} \|v(x,1)\|_H^2 e^{A_1|x|^q} dx = A_1^2,
$$

where

$$
A_0 = \frac{b_2}{(2 - \bar{t})^p}, A_1 = \frac{b_2}{(1 - \bar{t})^p}.
$$
\n(5.2)

For  $V(x,t) = F(u, \bar{u})$ , by hypothesis

$$
||V(x,t)||_{H} \leq C ||u(x,t-1+\bar{t})||_{H}^{\theta} \leq \frac{C}{(2-t-\bar{t})^{\theta n/2}} e^{\frac{C|x|^{p}}{(2-t-\bar{t})^{p}}}.
$$

By using Appell transformation if we suppose that  $v(y, s)$  is a solution of

$$
\partial_{s} \upsilon = i \left( \Delta \upsilon + A \upsilon + V \left( y, s \right) \upsilon \right), \, y \in R^{n}, \, t \in [0,1],
$$

 $\alpha$  and  $\beta$  are positive, then

$$
\tilde{u}(x,t) = \left(\sqrt{\alpha\beta}\sigma(t)\right)^{\frac{n}{2}} u\left(\sqrt{\alpha\beta}x\sigma(t), \beta t\sigma(t)\right) e^{\eta}.
$$
 (5.3)

verifies the equation

$$
\partial_t \tilde{u} = i \left[ \Delta \tilde{u} + A \tilde{u} + \tilde{V}(x, t) \tilde{u} + \tilde{F}(x, t) \right], \ x \in R^n, \ t \in [0, 1] \tag{5.4}
$$

with  $\tilde{V}\left(x,t\right),$   $\tilde{F}\left(x,t\right)$  defined by (3.3), (3.4) and

$$
\left\|e^{\gamma|x|^{p}}\tilde{u}_{k}\left(x,0\right)\right\|_{X}=\left\|e^{\gamma\left(\frac{\alpha}{\beta}\right)^{p/2}|x|^{p}}\upsilon\left(x,0\right)\right\|_{X}=a_{0},\tag{5.5}
$$

$$
\left\|e^{\gamma|x|^{p}}\tilde{u}_{k}\left(x,1\right)\right\|_{X}=\left\|e^{\gamma\left(\frac{\beta}{\alpha}\right)^{p/2}|x|^{p}}v\left(x,1\right)\right\|_{X}=a_{1}.
$$

It follows from expressions (3.6) and (3.8) that

$$
\gamma \sim \frac{1}{(1-\bar{t})^{p/2}}, \ \beta \sim \frac{1}{(1-\bar{t})^{p/2}}, \ \alpha \sim 1.
$$
 (5.6)

Next, we shall estimate

$$
\left\|\tilde{V}\left(x,t\right)\right\|_{L_{t}^{1}L_{x}^{\infty}\left(L(H),R\right)},
$$

for a  $r > 0$ , where

$$
L_{t}^{1}L_{x}^{\infty}(L(H), r) = L^{1}(0, 1; L^{\infty}(R^{n}/O_{r}); L(H)).
$$

Thus,

$$
\left\|\tilde{V}\left(x,t\right)\right\|_{L(H)} \leq \frac{\beta}{\alpha} \left\|V\left(y,s\right)\right\|_{L(H)} \leq \frac{\beta}{\alpha} \frac{C}{\left(1-\bar{t}\right)^{\theta n/2}} e^{C|y|^p},
$$

with

$$
|y| = \sqrt{\alpha \beta} |x| \sigma(t) \ge R \sqrt{\frac{\alpha}{\beta} \sim \frac{R}{\sqrt{\beta}}} = Cr (1 - \overline{t})^{1/2}.
$$

Hence,

$$
\left\|\tilde{V}\left(.,t\right)\right\|_{L^{\infty}(R^{n};L(H))} \leq \frac{\beta}{\alpha} \left\|V\left(.,s\right)\right\|_{L^{\infty}(R^{n};L(H))} \leq \frac{C}{\left(1-\overline{t}\right)^{1+\theta n/2}}
$$

and

$$
\left\| \tilde{V}(x,t) \right\|_{L_{t}^{1} L_{x}^{\infty}(L(H),r)} \leq \frac{\beta}{\alpha} \left\| V(y,s) \right\|_{L_{t}^{1} L_{x}^{\infty}(r;\beta)} \leq
$$
\n
$$
\frac{C}{(1-t)^{1+\theta n/2}} e^{-Cr^{p}(1-\bar{t})^{1/2}},
$$
\n(5.7)

where

$$
L_t^1 L_x^{\infty}(r, \beta) = L^1 \left( 0, 1; L^{\infty} \left( R^n / O_{Cr/\sqrt{\beta}}; L(H) \right) \right).
$$

To apply Lemma 3.1 we need

$$
\left\|\tilde{V}\left(x,t\right)\right\|_{L_{t}^{1}L_{x}^{\infty}\left(L(H),R\right)} \leq \frac{C}{\left(1-\bar{t}\right)^{1+\theta n/2}}e^{-Cr^{p}\left(1-\bar{t}\right)^{1/2}} \leq \delta_{0}.\tag{5.8}
$$

for some  $R,$  i.e.,

$$
r \sim \frac{C}{\left(1-\bar{t}\right)^{p/2}} \delta\left(t\right),\tag{5.9}
$$

where

$$
\delta(t) = \log^{\frac{1}{p}} \phi(t), \ \phi(t) = \frac{C}{\delta_0 (1 - \bar{t})^{\theta n/2}}.
$$

Let

$$
\mathbb{V} = \tilde{V}_{\chi(|x>R|)}(x,t), \ \mathbb{F} = \tilde{V}_{\chi(|x
$$

By using  $(5.5) - (5.10)$ , by virtue of Lemma 3.1 and  $(4.7)$  we deduced

$$
\sup_{t \in [0,1]} \left\| e^{\gamma |x|^p} \tilde{u}(.,t) \right\|_X^2 \le C \left( \left\| e^{\gamma |x|^p} \tilde{u}(.,0) \right\|_X^2 + \left\| e^{\gamma |x|^p} \tilde{u}(.,1) \right\|_X^2 \right) +
$$
  

$$
Ca^2 e^{C \gamma r^p} \left\| \tilde{V}(x,t) \right\| \le \frac{Ca^2}{(1-\overline{t})^p} e^{\delta(t)}, \ a = \|u_0\|_X.
$$

Next, using the same argument given in section 4,  $(4.10) - (4.22)$ , one finds that

$$
\gamma \int_{0}^{1} \int_{R^{n}} t(1-t) (1+|x|)^{p-2} \|\nabla \tilde{u}(x,t)\|_{H} e^{\gamma |x|^{p}} dxdt \leq \frac{Ca^{2}}{(1-\bar{t})^{p}} e^{\delta(t)}.
$$

Now we turn to the lower bounds estimates. Since they are similar to those given in detail in Section 3, we obtain that the estimate (4.24) for potential operator function  $V(x, t)$  when

$$
\frac{\theta n}{2} - 1 < \frac{p}{2(2-p)}, \text{ i.e., } p > \frac{2(\theta n - 2)}{\theta n - 1}.
$$

Finally, we get

$$
e^{\frac{C}{(1-t)^p} \log \phi(t)} \le e^{C\gamma r^p} \le e^{\frac{C}{(1-t)^{p/2}} \vartheta(t)}, \ \vartheta(t) = C (1-t)^{\frac{-p^2}{2(2-p)}}
$$

for  $p < \frac{p}{2} + \frac{p^2}{2(2-p)}$  $\frac{p}{2(2-p)}$ , i.e.  $p > 1$  that assumed, i.e. we obtain the assertion of Theorem 3.

**Remark 5.1.** Let us consider the case  $\theta = 4/n$  in Theorem 3, i.e.  $p > 4/3$ . Then from Theorem 3 we obtain

Result 5.1. Assume that the conditions of Theorem 3 are satisfied for  $p > 4/3$ . Then  $u(x, t) \equiv 0$ .

## 6. Unique continuation properties for the system of Schredinger equation

Consider the Cauchy problem for the finite or infinite system of Schrödinger equation

$$
\frac{\partial u_m}{\partial t} = i \left[ \Delta u_m + \sum_{j=1}^N a_{mj}(x) u_j + \sum_{j=1}^N b_{mj}(x) u_j \right], \ x \in R^n, \ t \in (0, T), \ (6.1)
$$

where  $u = (u_1, u_2, ..., u_N), u_j = u_j(x, t), a_{mj} = a_{mj}(x)$  are and  $b_{mj} =$  $b_{mj}(x,t)$  are complex valued functions. Let  $l_2 = l_2(N)$  and  $l_2^s = l_2^s(N)$  (see [23, § 1.18]). Let  $A$  be the operator in  $l_2(N)$  defined by

$$
D(A) = \left\{ u = \{u_j\} \in l_2, \ \|u\|_{l_2(N)} = \left(\sum_{j=1}^N 2^{sj} |u_j|^2\right)^{\frac{1}{2}} < \infty\right\},\
$$

$$
A = [a_{mj}], a_{mj} = G_m(x) 2^{sj}, s > 0, m, j = 1, 2, ..., N, N \in \mathbb{N}
$$

and

$$
D\left(V\left(x,t\right)\right) = l_2,
$$

$$
V(x,t) = [b_{mj}(x,t)], b_{mj}(x,t) = g_m(x,t) 2^{sj}, m, j = 1,2,...,N.
$$

Let

$$
X_2 = L^2(R^n; l_2), \; X_2(A) = L^2(R^n; l_2^s), \; Y^{k,2} = W^{k,2}(R^n; l_2).
$$

From Theorem 1 we obtain the following result

**Theorem 6.1.** Suppose  $a_{mj}$  are bounded cotinious on  $R^n$  and  $b_{mj}$  are bounded functions on  $\overline{R}^n \times [0, T]$ . Assume there exist the constants  $a_0, a_1$ ,  $a_2 > 0$  such that for any  $k \in \mathbb{Z}^+$  a solution  $u \in C([0,1]; X_2(A))$  of  $(6.1)$  satisfy

$$
\int_{R^{n}} \|u(x,0)\|_{l_{2}}^{2} e^{2a_{0}|x|^{p}} dx < \infty, \text{ for } p \in (1,2),
$$
  

$$
\int_{R^{n}} \|u(x,1)\|_{l_{2}}^{2} e^{2k|x|^{p}} dx < a_{2} e^{2a_{1}k^{\frac{q}{q-p}}}, \frac{1}{p} + \frac{1}{q} = 1.
$$

Moreover, there exists  $M_p > 0$  such that

$$
a_0 a_1^{p-2} > M_p.
$$

Then  $u(x,t) \equiv 0$ .

**Proof.** It is easy to see that A is a symmetric operator in  $l_2$  and other conditions of Theorem 1 are satisfied. Hence, from Teorem 1 we obtain the conculision.

## 7. Unique continuation properties for nonlinear anisotropic Schredinger equation

The regularity property of BVPs for elliptic equations were studied e.g. in [1, 2]. Let  $\Omega = R^n \times G$ ,  $G \subset R^d$ ,  $d \geq 2$  is a bounded domain with  $(d-1)$ dimensional boundary ∂G. Let us consider the following problem

$$
i\partial_t u + \Delta_x u + \sum_{|\alpha| \le 2m} a_\alpha(x, y) D_y^\alpha u(x, y, t) + F(u, \bar{u}) u = 0,
$$
 (7.1)  

$$
x \in R^n, y \in G, t \in [0, 1],
$$

$$
B_j u = \sum_{|\beta| \le m_j} b_{j\beta}(y) D_y^{\beta} u(x, y, t) = 0, x \in \mathbb{R}^n, y \in \partial G, j = 1, 2, ..., m, (7.2)
$$

where  $a_{\alpha}$ ,  $b_{j\beta}$  are the complex valued functions,  $\alpha = (\alpha_1, \alpha_2, ..., \alpha_d)$ ,  $\beta =$  $(\beta_1, \beta_2, ..., \beta_d), \mu_i < 2m, K = K(x, y, t)$  and

$$
D_x^k = \frac{\partial^k}{\partial x^k}
$$
,  $D_j = -i \frac{\partial}{\partial y_j}$ ,  $D_y = (D_{1, ..., D_d})$ ,  $y = (y_1, ..., y_d)$ .

Let

$$
\xi' = (\xi_1, \xi_2, ..., \xi_{d-1}) \in R^{d-1}, \ \alpha' = (\alpha_1, \alpha_2, ..., \alpha_{d-1}) \in Z^{d-1},
$$

$$
A(y_0, \xi', D_y) = \sum_{|\alpha'| + j \le 2m} a_{\alpha'}(y_0) \xi_1^{\alpha_1} \xi_2^{\alpha_2} ... \xi_{d-1}^{\alpha_{d-1}} D_y^j \text{ for } y_0 \in \bar{G}
$$

$$
B_j(y_0, \xi', D_y) = \sum_{|\beta'| + j \le m_j} b_{j\beta'}(y_0) \xi_1^{\beta_1} \xi_2^{\beta_2} ... \xi_{d-1}^{\beta_{d-1}} D_y^j \text{ for } y_0 \in \partial G.
$$

Let

$$
X_2 = L^2\left(R^n; L^2\left(G\right)\right) = L^2\left(\Omega\right), \ X_2\left(A\right) = L^2\left(R^n; W^{2m,2}\left(G\right)\right), \ Y^{k,2} = W^{k,2}\left(R^n; L^2\left(G\right)\right).
$$

Theorem 7.1. Let the following conditions be satisfied:

(1)  $G \in C^2$ ,  $a_{\alpha} \in C(\overline{\Omega})$  for each  $|\alpha| = 2m$  and  $a_{\alpha} \in L_{\infty}(G)$  for each  $|\alpha| < 2m;$ 

(2) 
$$
b_{j\beta} \in C^{2m-m_j}(\partial G)
$$
 for each j,  $\beta$  and  $m_j < 2m$ ,  $\sum_{j=1}^m b_{j\beta}(y') \sigma_j \neq 0$ , for

 $|\beta| = m_j, y \in \partial G$ , where  $\sigma = (\sigma_1, \sigma_2, ..., \sigma_d) \in R^d$  is a normal to  $\partial G$ ; (3) for  $y \in \bar{G}$ ,  $\xi \in R^d$ ,  $\lambda$  with  $|\arg \lambda| \leq \varphi$  for  $0 \leq \varphi < \pi$ ,  $|\xi| + |\lambda| \neq 0$  let  $\lambda + \sum a_{\alpha}(y) \xi^{\alpha} \neq 0$ ;  $|\alpha| = 2m$  $a_{\alpha}(y)\,\xi^{\alpha}\neq 0;$ 

(4) for each  $y_0 \in \partial G$  local BVP in local coordinates corresponding to  $y_0$ :

$$
\lambda + A(y_0, \xi', D_y) \vartheta(y) = 0,
$$

$$
B_j(y_0, \xi', D_y) \vartheta(0) = h_j, j = 1, 2, ..., m
$$

has a unique solution  $\vartheta \in C_0(\mathbb{R}_+)$  for all  $h = (h_1, h_2, ..., h_d) \in \mathbb{C}^d$  and for  $\xi' \in R^{d-1};$ 

(5) there exist positive constants  $b_0$  and  $\theta$  such that a solution  $u \in C([-1, 1]; X_2(A))$ of  $(7.1) - (7.2)$  satisfied

$$
||F(u,\bar{u})||_{L^{2}(G)} \leq b_0 ||u||_{L^{2}(G)}^{\theta} \text{ for } ||u||_{X_2(A)} > 1;
$$

(6) Suppose

$$
||u (., t)||_{L^{2}(\Omega)} = ||u (., 0)||_{L^{2}(\Omega)} = ||u_{0}||_{L^{2}(\Omega)} = a
$$

for  $t \in [-1,1]$  and that  $(2.15)$  holds with  $Q(.)$  satisfies  $(2.16)$  for  $H = L^2(G)$ . If  $p > p(\theta) = \frac{2(\theta n - 2)}{(\theta n - 1)}$ , then  $a \equiv 0$ .

**Proof.** Let us consider operators A and  $V(x,t)$  in  $H = L^2(G)$  that are defined by the equalities

$$
D(A) = \{ u \in W^{2m,2}(G), B_j u = 0, j = 1, 2, ..., m \}, Au = \sum_{|\alpha| \le 2m} a_{\alpha}(y) D_y^{\alpha} u(y).
$$

Then the problem  $(7.1)$  –  $(7.2)$  can be rewritten as the problem  $(2.12)$ , where  $u(x) = u(x,.)$ ,  $f(x) = f(x,.)$ ,  $x \in \mathbb{R}^n$  are the functions with values in  $H =$  $L^2(G)$ . By virtue of [1] operator  $A + \mu$  is positive in  $L^2(G)$  for sufficiently large  $\mu > 0$ . Moreover, in view of (1)-(6) all conditions of Theorem 3 are hold. Then Theorem 3 implies the assertion.

## 8. The Wentzell-Robin type mixed problem for Schredinger equations

Consider the problem  $(1.5) - (1.6)$ . Let

$$
\sigma = R^{n} \times (0,1), X_2 = L^2(\sigma), Y^{2,k} = W^{2,k}(\sigma).
$$

Suppose  $\nu = (\nu_1, \nu_2, ..., \nu_n)$  are nonnegative real numbers. In this section, from Theorem 1 we obtain the following result:

Theorem 8.1. Suppose the following conditions are satisfied:

(1) a and b are a real-valued functions on  $(0, 1)$  and  $a(t) > 0$ . Moreover, a (.) is bounded continious function on  $R^n \times [0,1]$  and

$$
\exp\left(-\int_{\frac{1}{2}}^{x} b(t) a^{-1}(t) dt\right) \in L_1(0,1);
$$

(2)  $u_1, u_2 \in C([0,1]; Y^{2,k})$  are strong solutions of  $(1.2)$  with  $k > \frac{n}{2}$ ;

(3)  $F: \mathbb{C} \times \mathbb{C} \to \mathbb{C}, F \in C^k, F(0) = \partial_u F(0) = \partial_{\bar{u}} F(0) = 0;$ 

(4) there exist positive constants  $\alpha$  and  $\beta$  such that

$$
e^{\frac{|\alpha x|^p}{p}}(u_1(.,0)-u_2(.,0)) \in X_2, e^{\frac{|\beta x|^q}{q}}(u_1(.,0)-u_2(.,0)) \in X_2,
$$

with

$$
p \in (1, 2), \frac{1}{p} + \frac{1}{q} = 1;
$$

(5) there exists  $N_p > 0$  such that

 $\alpha\beta > N_p$ .

Then  $u_1(x,t) \equiv u_2(x,t)$ .

**Proof.** Let  $H = L^2(0, 1)$  and A is a operator defined by  $(1.4)$ . Then the problem  $(1.5) - (1.6)$  can be rewritten as the problem  $(1.2)$ . By virtue of [10, 11] the operator A generates analytic semigroup in  $L^2(0,1)$ . Hence, by virtue of (1)-(5) all conditons of Theorem 2 are satisfied. Then Theorem 2 implies the assertion.

### Acknowledgements

The author would like to express a gratitude to Dr. Neil. Course for his useful advice in English in preparing of this paper

### References

- 1. H. Amann, Linear and quasi-linear equations,1, Birkhauser, Basel (1995).
- 2. A. Benedek, A. Calderòn, R. Panzone, Convolution operators on Banach space valued functions, Proc. Nat. Acad. Sci. USA, 48(1962), 356–365
- 3. A. Bonami, B. Demange, A survey on uncertainty principles related to quadratic forms, Collect. Math., V. Extra, (2006), 1–36.
- 4. A.-P. Calder´on, Commutators of singular integral operators, Proc. Nat. Acad. Sci. U.S.A. 53(1965), 1092–1099.
- 5. C. E. Kenig, G. Ponce, L. Vega, On unique continuation for nonlinear Schr¨odinger equations, Comm. Pure Appl. Math. 60 (2002), 1247–1262.
- 6. L. Escauriaza, C. E. Kenig, G. Ponce, L. Vega, On uniqueness properties of solutions of Schrödinger Equations, Comm. PDE. 31, 12 (2006) 1811– 1823.
- 7. L. Escauriaza, C. E. Kenig, G. Ponce, and L. Vega, Uncertainty principle of Morgan type and Schrödinger evolution, J. London Math. Soc. 81, (2011) 187–207.
- 8. L. Escauriaza, C. E. Kenig, G. Ponce, and L. Vega, Hardy's uncertainty principle, convexity and Schrödinger Evolutions, J. European Math. Soc. 10, 4 (2008) 883–907.
- 9. L. Hörmander, A uniqueness theorem of Beurling for Fourier transform pairs, Ark. Mat. 29, 2 (1991) 237–240.
- 10. J. A. Goldstain, Semigroups of Linear Operators and Applications, Oxford University Press, Oxfard (1985).
- 11. A. Favini, G. R. Goldstein, J. A. Goldstein and S. Romanelli, Degenerate Second Order Differential Operators Generating Analytic Semigroups in  $L_p$  and  $W^{1,p}$ , Math. Nachr. 238 (2002), 78 –102.
- 12. V. Keyantuo, M. Warma, The wave equation with Wentzell–Robin boundary conditions on Lp-spaces, J. Differential Equations 229 (2006) 680–697.
- 13. S. G. Krein, Linear Differential Equations in Banach space, American Mathematical Society, Providence, (1971).
- 14. A. Lunardi, Analytic Semigroups and Optimal Regularity in Parabolic Problems, Birkhauser (2003).
- 15. J-L. Lions, E. Magenes, Nonhomogenous Boundary Value Broblems, Mir, Moscow (1971).
- 16. V. B. Shakhmurov, Imbedding theorems and their applications to degenerate equations, Differential equations, 24 (4) (1988), 475-482.
- 17. V. B. Shakhmurov, Linear and nonlinear abstract equations with parameters, Nonlinear Anal-Theor., 73 (2010), 2383-2397.
- 18. R. Shahmurov, On strong solutions of a Robin problem modeling heat conduction in materials with corroded boundary, Nonlinear Anal., Real World Appl., 13, (1)( 2011), 441-451.
- 19. R. Shahmurov, Solution of the Dirichlet and Neumann problems for a modified Helmholtz equation in Besov spaces on an annuals, J. Differential equations, 249(3)(2010), 526-550.
- 20. C. Segovia, J. L.Torrea, Vector-valued commutators and applications, Indiana Univ. Math.J. 38(4) (1989), 959–971.
- 21. E. M. Stein, R. Shakarchi, Princeton, Lecture in Analysis II. Complex Analysis, Princeton University Press (2003).
- 22. A. Stefanov, Strichartz estimates for the Schrodinger equation with radial data, Proc. Amer. Math. Soc. 129 (2001), 1395-1401.
- 23. B. Simon, M. Schechter, Unique Continuation for Schrodinger Operators with unbounded potentials, J. Math. Anal. Appl., 77 (1980), 482-492.
- 24. H. Triebel, Interpolation theory, Function spaces, Differential operators, North-Holland, Amsterdam (1978).
- 25. S. Yakubov and Ya. Yakubov, Differential-operator Equations. Ordinary and Partial Differential Equations, Chapman and Hall /CRC, Boca Raton  $(2000).$
- 26. Y. Xiao and Z. Xin, On the vanishing viscosity limit for the 3D Navier-Stokes equations with a slip boundary condition. Comm. Pure Appl. Math. 60 (7) (2007), 1027–1055.