

Non-explicit formula of boundary crossing probabilities by the Girsanov theorem

Yoann Potiron*¹,

¹*Faculty of Business and Commerce, Keio University. 2-15-45 Mita, Minato-ku, Tokyo, 108-8345, Japan, e-mail: potiron@fbc.keio.ac.jp*

Abstract: This paper derives several formulae for the probability that a Wiener process, which has a stochastic drift and random variance, crosses a one-sided stochastic boundary within a finite time interval. A non-explicit formula is first obtained by the Girsanov theorem when considering an equivalent probability measure in which the boundary is constant and equal to its starting value. A more explicit formula is then achieved by decomposing the Radon-Nikodym derivative inverse. This decomposition expresses it as the product of a random variable, which is measurable with respect to the Wiener process's final value, and an independent random variable. We also provide an explicit formula based on a strong theoretical assumption. To apply the Girsanov theorem, we assume that the difference between the drift increment and the boundary increment, divided by the standard deviation, is absolutely continuous. Additionally, we assume that its derivative satisfies Novikov's condition.

Keywords and phrases: Mathematical statistics, sequential analysis, first-passage time problem, boundary crossing probabilities, stochastic boundary process, Wiener process, Girsanov theorem.

1. Introduction

This paper concerns boundary crossing probabilities, i.e., the probability that a stochastic process crosses a boundary. The application of boundary crossing probabilities in statistics dates back to the one-sample Kolmogorov-Smirnov statistic, where the process represents the difference between the true and empirical cumulative distribution functions (cdfs). The primary application of boundary crossing probabilities is in sequential analysis. Initially, the focus was on boundary crossing probabilities for random walks. Due to the complexity of solving this problem, the literature often relies on continuous approximations and develops theoretical tools for cases where the process is a Wiener process

*The online version of this article contains supplementary material.

(see Gut (1974), Woodroffe (1976), Woodroffe (1977), Lai et al. (1977), Lai et al. (1979) and Siegmund (1986)).

Another field of application is in survival analysis. Matthews et al. (1985) show that tests for constant hazard involve the first-passage time (FPT) of an Ornstein-Uhlenbeck process. Butler et al. (1997) present a Bayesian approach when the process is semi-Markovian. Eaton et al. (1977) discuss the application of FPT for hospital stay. Aalen et al. (2001) study the case when the process is Markovian. Detailed reviews on FPT are available in Lee et al. (2006) and Lawless (2011) (Section 11.5, pp. 518-523).

Another application is in pricing barrier options in mathematical finance (see Roberts et al. (1997)). There are also some applications in econometrics. Abbring (2012) studies mixed FPT of a spectrally negative Levy process. Renault et al. (2014) considers mixed FPT of the sum of a Wiener process and a positive linear drift. Potiron et al. (2017) estimate the quadratic covariation between two price processes based on endogenous observations generated by FPT of an Itô-semimartingale to a stochastic boundary process.

Despite their importance for applications, explicit formulae of these boundary crossing probabilities only exist when the boundaries and the drift are linear. More specifically, Doob (1949) gives explicit formulae (Equations (4.2)-(4.3), pp. 397-398) based on elementary geometrical and analytical arguments. They are obtained when the final time is not finite, the variance is nonrandom, the drift is null and the boundaries are nonrandom linear with nonnegative upper trend and nonpositive lower trend. Malmquist (1954) obtains an explicit formula conditioned on the starting and final values of the Wiener process for a finite final time (Theorem 1, p. 526). This is obtained with Doob's transformation (Section 5, pp. 401-402) in the one-sided boundary case. Anderson (1960) derives an explicit formula conditioned on the final value of the Wiener process (Theorem 4.2, pp. 178-179) in the two-sided boundary case with linear drift. Then, he integrates it with respect to the final value of the Wiener process to get an explicit formula (Theorem 4.3, p. 180).

For square root boundaries, Breiman (1967) expresses the problem as boundary crossing probabilities of an Ornstein-Uhlenbeck process to a constant boundary. They are obtained with Doob's transformation. However, the boundary crossing probabilities of an Ornstein-Uhlenbeck process to a constant boundary are only known in the form of Laplace transform. Daniels (1969) uses the same technique and obtains an explicit formula. Finally, the boundary crossing probabilities of a jump diffusion process with linear drift to a constant boundary are obtained in the form of Laplace transform in Kou et al. (2003). Alili et al.

(2005), Doney et al. (2006) and Kypriano et al. (2010) consider a link between the first and last passage time and overshoot above/below a fixed level of a Lévy process. Potiron (2023) obtains an explicit formula when the boundary is constant and the stochastic process is a continuous local martingale.

Since there is no available explicit formula when the drift and the boundaries are not linear, there is a large literature on approximating and computing numerically these boundary crossing probabilities. Strassen (1967) (Lemma 3.3, p. 323) shows that P_g^Z is continuously differentiable when g is continuously differentiable. Durbin (1971, Wang et al. (1997) and Novikov et al. (1999) use piecewise-linear boundaries to approximate the general boundaries. Durbin (1985) gives a formula for a general boundary, which depends on asymptotic conditional expectations whose approximations are studied in Salminen (1988).

In this paper, we derive several formulae for the one-sided and two-sided boundary crossing probability when the boundaries and drift are stochastic processes and the variance is random. Unfortunately, these formulae are either non-explicit or explicit, but based on a strong theoretical assumption. We derive the results in two cases, i.e., (i) a simpler case when the one-sided boundary and the drift are nonrandom time-varying and the variance is nonrandom and (ii) a more complicated case when the one-sided boundary and the drift are stochastic processes and the variance is random.

More specifically, consider a stochastic process $(Z_t)_{t \in \mathbb{R}^+}$ defined as $Z_t = \mu_t + \sigma W_t$. Here, $(\mu_t)_{t \in \mathbb{R}^+}$ is a stochastic drift process, $(W_t)_{t \in \mathbb{R}^+}$ is a standard Wiener process with random time-invariant variance σ^2 , and $(g_t)_{t \in \mathbb{R}^+}$ and $(h_t)_{t \in \mathbb{R}^+}$ are two stochastic boundary processes. We focus on the probabilities of a process crossing one-sided and two-sided boundaries, defined as follows

$$P_g^Z(T) = \mathbb{P}\left(\sup_{0 \leq t \leq T} Z_t - g_t \geq 0\right), \quad (1)$$

$$P_{g,h}^Z(T) = \mathbb{P}\left(\sup_{0 \leq t \leq T} Z_t - g_t \geq 0 \text{ or } \sup_{0 \leq t \leq T} h_t - Z_t \geq 0\right), \quad (2)$$

i.e., the probability that the process Z crosses the boundary or one of both boundaries between 0 and the final time T .

We describe first the main results in the one-sided (i) case. A non-explicit formula is first obtained by the Girsanov theorem when considering an equivalent probability measure in which the boundary is constant equal to its starting value. To apply the Girsanov theorem, the main idea consists in rewriting the boundary crossing probability of a time-varying boundary as an equivalent boundary crossing probability of a constant boundary. More specifically, we define the new nonrandom drift as $u_t = \frac{\mu_t - g_t + g_0}{\sigma}$, the new $\sigma(W_t)$ -measurable

process as $Y_t = u_t + W_t$ and the new constant boundary as $b = \frac{g_0 - \mu_0}{\sigma}$. We then observe that the boundary crossing probability (1) may be rewritten as $P_g^Z(T) = P_b^Y(T)$. We thus obtain (see Proposition 2)

$$\mathbb{P}(T_b^Y \leq T | W_T) = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} | W_T]. \quad (3)$$

Here, $M_T = \exp(\overline{W}_T - \frac{1}{2} \int_0^T \theta_s^2 ds)$ is the Radon-Nikodym derivative with $\overline{W}_T = \int_0^T \theta_s dW_s$, $u_T = \int_0^T \theta_s ds$, and θ is a nonrandom function $\theta : [0, T] \rightarrow \mathbb{R}$.

A more explicit formula is then obtained by using an elementary decomposition of the Radon-Nikodym derivative inverse. This decomposition expresses it as the product of a random variable, which is $\sigma(W_T)$ -measurable, and an independent random variable. More specifically, the decomposition is based on the fact that (W_T, \overline{W}_T) is a centered normal random vector under \mathbb{P} . It consists of rewriting \overline{W}_T as $\overline{W}_T = \alpha W_T + \tilde{\alpha} \tilde{W}$, where \tilde{W} is a standard normal random variable under \mathbb{P} , independent of W_T . Then, we obtain (see Theorem 4)

$$\begin{aligned} \mathbb{P}(T_b^Y \leq T | W_T) = \\ \exp(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds) \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T]. \end{aligned} \quad (4)$$

We also give a formula based on a strong theoretical assumption. More specifically, we assume that

$$\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T] \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha} \tilde{W}) | W_T].$$

Then, the non-explicit factor in Equation (4) is equal to (see Theorem 6)

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T] &= \exp\left(-\frac{2b(b - Y_T)}{T}\right) \\ &\times \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned}$$

Here, \mathcal{L}_N is the Laplace transform of a standard normal variable. This is based on the preliminary result (see Lemma 5)

$$\mathbb{Q}(T_b^Y \leq T | W_T) = \exp\left(-\frac{2b(b - Y_T)}{T}\right),$$

by using the explicit formula from Malmquist (1954) (Theorem 1, p. 526) and since Y is a standard Wiener process under \mathbb{Q} .

Finally, we give formulae of $P_b^Y(T)$ by integrating $\mathbb{P}(T_b^Y \leq T | W_T)$ with respect to the value of W_T (see Corollary 7). We also derive similar formulae in the case (ii). If we define v as $v = (g, \mu, \sigma)$ and we assume that v is independent

of W , the elementary idea in the case (ii) is to condition by both W_T and v , i.e., to derive results of the form $\mathbb{P}(T_b^Y \leq T | W_T, v)$. We also derive similar formulae in the two-sided boundary case when the difference between the deviation of each boundary from their starting value is linear, i.e., there exists $\beta \in \mathbb{R}$ such that $h_t - h_0 = g_t - g_0 + \beta t$. To apply the Girsanov theorem to the two-sided boundary case, we cannot use two different drifts since the process Z is unique in the boundary crossing probability (2). Thus, the elementary idea consists in rewriting the FPT to a two-sided time-varying boundary as an equivalent FPT to a two-sided boundary, with one constant boundary and one linear boundary. More specifically, we define the new drift as $u_t = \frac{\mu_t - \mu_0 - g_t + g_0}{\sigma}$, the new process as $Y_t = u_t + W_t$, the new constant boundary as $b = \frac{\mu_0 - g_0}{\sigma}$ and the new linear boundary as $c_t = \frac{h_0 - \mu_0 + \beta t}{\sigma}$. We then observe that the boundary crossing probability (2) may be rewritten as $P_{g,h}^Z = P_{b,c}^Y$.

When the boundaries and the drift are linear, the one-sided and two-sided boundary crossing probability (1)-(2) can be obtained easily. For instance, one can use a combination of the Girsanov theorem and the reflection principle, or by calculating the Laplace transforms via some appropriate martingales and the optional sampling theorem. Details of both methods can be found in many classical textbooks on stochastic analysis, e.g., Karatzas et al. (2012) or Revuz et al. (2013). More specifically, one can obtain by the Girsanov theorem Equation (3), and since the Radon-Nikodym derivative inverse M_T^{-1} is $\sigma(W_T)$ -measurable in that simpler case, one can conclude by the joint distribution of the maximum and terminal value of a Wiener process based on the reflection principle. When the boundaries and the drift are not linear, however, M_T^{-1} is no longer $\sigma(W_T)$ -measurable. This renders a direct calculation not possible since that would require to extend the arguments based on the reflection principle. As discussed above, we circumvent that difficulty by an elementary decomposition of the Radon-Nikodym derivative inverse. This decomposition expresses it as the product of a $\sigma(W_T)$ -measurable random variable and an independent random variable. The price to pay is that we do not obtain explicit formulae without strong theoretical assumptions.

As it stands, our obtained formulae are based on a strong theoretical assumption. Unfortunately, such assumption is hard to show when the final time is fixed. The main reason is that the boundary is not linear enough. However, we conjecture that this assumption is asymptotically met if we divide the final time interval $[0, T]$ into smaller time intervals with length that converges to 0 asymptotically. This is due to the fact that the boundary gets more and more linear locally. Thus, we can relatively safely apply our obtained formulae locally.

By taking the sum of the local approximations, we can then approximate the boundary crossing probabilities (1)-(2).

2. Main results

2.1. One-sided time-varying boundary case

In this section, we consider the case when the one-sided boundary and the drift are nonrandom time-varying and the variance is nonrandom.

We consider the complete stochastic basis $\mathcal{B} = (\Omega, \mathbb{P}, \mathcal{F}, \mathbf{F})$, where \mathcal{F} is a σ -field and $\mathbf{F} = (\mathcal{F}_t)_{t \in \mathbb{R}^+}$ is a filtration. We define the set of continuous functions from \mathbb{R}^+ to \mathbb{R} as $\mathcal{C}(\mathbb{R}^+, \mathbb{R})$. We first give the definition of the set of boundary functions. We assume that the boundary is continuous since it is required in the assumptions for the Girsanov theorem.

Definition 2.1. We define the set of boundary functions as $\mathcal{G} = \mathcal{C}(\mathbb{R}^+, \mathbb{R})$.

We now give the definition of the FPT. We assume that the stochastic process is continuous since we consider a Wiener process with a continuous drift which is required in the assumptions for the Girsanov theorem.

Definition 2.2. We define the FPT of an \mathbf{F} -adapted continuous process Z to a boundary $g \in \mathcal{G}$ satisfying $Z_0 \leq g_0$ as

$$\mathbb{T}_g^Z = \inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \geq g_t\}. \quad (5)$$

We have that Z is a continuous and \mathbf{F} -adapted stochastic process and $\inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \geq g_t\} = \inf\{t \in \mathbb{R}^+ \text{ s.t. } (t, Z_t) \in G\}$, where $G = \{(t, u) \in \mathbb{R}^+ \times \mathbb{R} \text{ s.t. } u \geq g_t\}$ is a closed subset of \mathbb{R}^2 . Thus, the FPT \mathbb{T}_g^Z is an \mathbf{F} -stopping time by Theorem I.1.27 (p. 7) in Jacod et al. (2003). We can rewrite the boundary crossing probability P_g^Z as the cdf of \mathbb{T}_g^Z , i.e.,

$$P_g^Z(t) = \mathbb{P}(\mathbb{T}_g^Z \leq t) \text{ for any } t \geq 0. \quad (6)$$

We assume that W is an \mathbf{F} -standard Wiener process. We assume that $Z_t = \mu_t + \sigma W_t$. Here, μ is time-varying, nonrandom, and satisfies $\mu_0 < g_0$. The variance σ^2 is time-invariant, nonrandom, and positive, i.e., $\sigma^2 > 0$. To apply the Girsanov theorem, the main elementary idea is to rewrite the FPT to a time-varying boundary as an equivalent FPT to a constant boundary. More specifically, we define the new nonrandom drift as $u_t = \frac{\mu_t - \mu_0 - g_t + g_0}{\sigma}$, the new process as $Y_t = u_t + W_t$ and the new constant boundary as $b = \frac{g_0 - \mu_0}{\sigma}$. We then observe that the FPT (5) may be rewritten as

$$\mathbb{T}_g^Z = \mathbb{T}_b^Y. \quad (7)$$

Then, we will consider an equivalent probability measure under which the new process Y will be a standard Wiener process. Accordingly, we provide the assumption which corresponds to Novikov's condition (see Novikov (1972)) which is required to apply the Girsanov theorem (see Girsanov (1960)). The proofs of this paper would hold with no change with the more general conditions obtained by Kazamaki (1977).

Assumption A. We assume that $u \neq 0$, i.e., that there exists a $t \in [0, T]$ such that $u_t \neq 0$. We also assume that u is absolutely continuous on $[0, T]$, i.e., there exists a nonrandom function $\theta : [0, T] \rightarrow \mathbb{R}$ with $u_t = \int_0^t \theta_s ds$. Finally, we assume that $\int_0^T \theta_s^2 ds < \infty$.

Definition 2.3. We define M as

$$M_t = \exp\left(\int_0^t \theta_s dW_s - \frac{1}{2} \int_0^t \theta_s^2 ds\right) \text{ for any } 0 \leq t \leq T. \quad (8)$$

By **Assumption A**, M satisfies Novikov's condition and thus is a positive martingale. We embed this result and its implications on an equivalent probability measure \mathbb{Q} by the Girsanov theorem in the following lemma.

Lemma 1. *Under **Assumption A**, we have that M is a positive martingale. Thus, we can consider an equivalent probability measure \mathbb{Q} such that the Radon-Nikodym derivative is defined as $\frac{d\mathbb{Q}}{d\mathbb{P}} = M_T$. Finally, Y is a standard Wiener process under \mathbb{Q} .*

Consequently, we obtain that $\mathbb{E}_{\mathbb{P}}[X] = \mathbb{E}_{\mathbb{Q}}[XM_T^{-1}]$ for any \mathcal{F}_T -measurable random variable X by a change of probability in the expectation. The next proposition reexpresses $\mathbb{P}(T_b^Y \leq T|W_T)$ under \mathbb{Q} and is a result of interest in its own right although the obtained formulae are not explicit. The proof is based on Lemma 1 and its consequence in the particular case $X = \mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}$ where E_T is a $\sigma(W_T)$ -measurable event. We define \overline{W}_t as

$$\overline{W}_t = \int_0^t \theta_s dW_s. \quad (9)$$

Proposition 2. *Under **Assumption A**, we have*

$$\mathbb{P}(T_b^Y \leq T|W_T) = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1}|W_T]. \quad (10)$$

This can be reexpressed as

$$\mathbb{P}(T_b^Y \leq T|W_T) = \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}}|W_T, \overline{W}_T]|W_T]. \quad (11)$$

It remains to calculate $\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_b^Y \leq T\}} M_T^{-1} | W_T]$ or

$$\mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_b^Y \leq T\}} | W_T, \overline{W}_T] | W_T]$$

if we want to derive a completely explicit formula. Although Y is a standard Wiener process under \mathbb{Q} by Lemma 1, the presence of M_T^{-1} or \overline{W}_T in the conditional expectation renders a direct calculation not possible. Indeed, we would need to extend the arguments based on the reflection principle. We circumvent that difficulty by an elementary decomposition of the Radon-Nikodym derivative inverse M_T^{-1} . More specifically, the decomposition is based on the fact that (W_T, \overline{W}_T) is a centered normal random vector under \mathbb{P} . It consists of rewriting \overline{W}_T as $\overline{W}_T = \alpha W_T + \tilde{\alpha} \tilde{W}$, where \tilde{W} is a standard normal random variable under \mathbb{P} , independent of W_T . We define the correlation under \mathbb{P} between W_T and \overline{W}_T as ρ , i.e., $\rho = \text{Cor}_{\mathbb{P}}(W_T, \overline{W}_T)$.

Lemma 3. *Under Assumption A, we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . We can also show that $\rho = \frac{\int_0^T \theta_s ds}{T \int_0^T \theta_s^2 ds}$. Moreover, there exists a standard normal random variable \tilde{W} under \mathbb{P} which is independent of W_T , and such that \overline{W}_T when normalized can be reexpressed as*

$$\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}} = \rho \frac{W_T}{\sqrt{T}} + \sqrt{1 - \rho^2} \tilde{W}, \quad (12)$$

This can be reexpressed as

$$\overline{W}_T = \alpha W_T + \tilde{\alpha} \tilde{W}, \quad (13)$$

where $\alpha = \rho \sqrt{T^{-1} \int_0^T \theta_s^2 ds}$ and $\tilde{\alpha} = \sqrt{(1 - \rho^2) \int_0^T \theta_s^2 ds}$. If we define $\tilde{\theta}_t = \frac{\theta_s - \alpha}{\tilde{\alpha}}$, we can reexpress \tilde{W} as

$$\tilde{W} = \int_0^T \tilde{\theta}_s dW_s. \quad (14)$$

Finally, $\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} .

Our main result is the next theorem, which gives a more explicit formula to $\mathbb{P}(\mathbb{T}_b^Y \leq T | W_T)$. The proof is based on Lemma 3.

Theorem 4. *Under Assumption A, we have*

$$\begin{aligned} & \mathbb{P}(\mathbb{T}_b^Y \leq T | W_T) \\ &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_b^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T]. \end{aligned} \quad (15)$$

We first calculate $\mathbb{Q}(T_b^Y \leq T|W_T)$, whose explicit formula is given in the following lemma. This reexpresses Malmquist (1954) (Theorem 1, p. 526) under \mathbb{Q} , who considers the linear case $\mu_t = 0$, $g_t = at + b$ and $\sigma = 1$ under \mathbb{P} . He obtains that

$$\begin{aligned} & \mathbb{P}(T_g^Z \leq T|W_T = x) \\ &= \exp\left(-\frac{2b(aT + b - x)}{T}\right) \mathbf{1}_{\{x \leq aT + b\}} + \mathbf{1}_{\{x > aT + b\}} \end{aligned} \quad (16)$$

for any $x \in \mathbb{R}$.

Lemma 5. *Under **Assumption A**, we have*

$$\mathbb{Q}(T_b^Y \leq T|W_T) = \exp\left(-\frac{2b(b - Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} + \mathbf{1}_{\{Y_T > b\}}. \quad (17)$$

The next theorem gives an explicit formula based on a strong theoretical assumption (19). The proof is based on Lemma 5. Let N be a standard normal random variable under \mathbb{P} . We define the Laplace transform of N as

$$\mathcal{L}_N(u) = \mathbb{E}_{\mathbb{P}}[\exp(-uN)]. \quad (18)$$

Theorem 6. *We assume that **Assumption A** and the following assumption*

$$\begin{aligned} & \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha}\tilde{W})|W_T] \\ &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}}|W_T] \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W})|W_T] \end{aligned} \quad (19)$$

holds. Then, we have

$$\begin{aligned} \mathbb{P}(T_b^Y \leq T|W_T) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \left(\exp\left(-\frac{2b(b - Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} + \mathbf{1}_{\{Y_T > b\}} \right) \\ &\quad \times \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned} \quad (20)$$

Finally, we get $P_b^Y(T)$ in the next corollary by integrating $\mathbb{P}(T_b^Y \leq T|W_T)$ with respect to the value of W_T . The proof follows the steps of Equations (3) in Wang et al (1997) (p. 55). We define the standard Gaussian cdf as $\phi(t) = \int_0^t \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right) du$ for any $t \in \mathbb{R}^+$.

Corollary 7. *Under **Assumption A**, we have*

$$P_b^Y(T) = 1 - \phi\left(\frac{b - u_T}{\sqrt{T}}\right) + \int_{-\infty}^{b - u_T} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right)$$

$$\begin{aligned}
& \times \exp\left(-\alpha x + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\
& \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T = x] dx. \tag{21}
\end{aligned}$$

If we further assume (19), we have

$$\begin{aligned}
P_b^Y(T) &= 1 - \phi\left(\frac{b - u_T}{\sqrt{T}}\right) \\
&+ \int_{-\infty}^{b - u_T} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-\alpha x + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\
&\times \exp\left(-\frac{2b(b - u_T - x)}{T}\right) \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}) dx. \tag{22}
\end{aligned}$$

We consider the particular case when the boundary is linear, there is no drift and the standard deviation is equal to unity $\mu_t = 0$, $g_t = at + b$ and $\sigma = 1$. Then, Corollary 7 reduces to Wang et al. (1997) (Equation (2), p. 55), i.e.,

$$P_b^Y(T) = 1 - \phi\left(\frac{b + aT}{\sqrt{T}}\right) + \exp(-2ba) \phi\left(\frac{b - aT}{\sqrt{T}}\right).$$

2.2. One-sided stochastic boundary process case

In this section, we consider the case when the one-sided boundary and the drift are stochastic processes and the variance is random.

We first give the definition of the set of stochastic boundary processes.

Definition 2.4. We define the set of stochastic boundary processes as $\mathcal{H} = \mathbb{R}^+ \times \Omega \rightarrow \mathbb{R}$ such that for any $g \in \mathcal{H}$ and $\omega \in \Omega$ we have $g(\omega) \in \mathcal{G}$ and g is \mathbf{F} -adapted.

We now give the definition of the FPT.

Definition 2.5. We define the FPT of an \mathbf{F} -adapted continuous process Z to a boundary $g \in \mathcal{H}$ satisfying $Z_0 \leq g_0 \forall \omega \in \Omega$ as

$$T_g^Z = \inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \geq g_t\}. \tag{23}$$

We have that $Z - g$ is an \mathbf{F} -adapted continuous stochastic process and $\inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \geq g_t\} = \inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t - g_t \geq 0\} = \inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t - g_t \in \mathbb{R}^+\}$. Thus, the FPT T_g^Z is an \mathbf{F} -stopping time by Theorem I.1.27 (p. 7) in Jacod et al. (2003). We can rewrite the boundary crossing probability P_g^Z as the cdf of T_g^Z , i.e.,

$$P_g^Z(t) = \mathbb{P}(T_g^Z \leq t) \text{ for any } t \geq 0. \tag{24}$$

We assume that μ is an \mathbf{F} -adapted stochastic process which satisfies $\mathbb{P}(\mu_0 < g_0) = 1$. We also assume that the variance σ^2 is time-invariant, random, and such that $\mathbb{P}(\sigma^2 = 0) = 0$. Finally, we assume that v is independent of W , where v is defined as $v = (g, \mu, \sigma)$.

Assumption B. We assume that $\mathbb{P}(\exists t \in [0, T] \text{ s.t. } u_t \neq 0) = 1$. We also assume that u is absolutely continuous on $[0, T]$, i.e., there exists a stochastic process $\theta : [0, T] \times \Omega \rightarrow \mathbb{R}$ with $u_t = \int_0^t \theta_s ds$, a.s.. Finally, we assume that $\mathbb{E}[\exp(\frac{1}{2} \int_0^T \theta_s^2 ds)] < \infty$.

Definition 2.6. We define M as

$$M_t = \exp\left(\int_0^t \theta_s dW_s - \frac{1}{2} \int_0^t \theta_s^2 ds\right) \text{ for any } 0 \leq t \leq T. \quad (25)$$

By **Assumption B**, M satisfies Novikov's condition and thus is a positive martingale.

Lemma 8. *Under **Assumption B**, we have that M is a positive martingale. Thus, we can consider an equivalent probability measure \mathbb{Q} such that the Radon-Nikodym derivative is defined as $\frac{d\mathbb{Q}}{d\mathbb{P}} = M_T$. Finally, Y is a standard Wiener process under \mathbb{Q} .*

The elementary idea in this section is to condition by both W_T and v , i.e., to derive results of the form $\mathbb{P}(T_b^Y \leq T | W_T, v)$. The next proposition reexpresses $\mathbb{P}(T_b^Y \leq T | W_T, v)$ under \mathbb{Q} . We define \overline{W}_t as

$$\overline{W}_t = \int_0^t \theta_s dW_s. \quad (26)$$

Proposition 9. *Under **Assumption B**, we have*

$$\mathbb{P}(T_b^Y \leq T | W_T, v) = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} | W_T, v]. \quad (27)$$

This can be reexpressed as

$$\mathbb{P}(T_b^Y \leq T | W_T, v) = \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, \overline{W}_T, v] | W_T, v]. \quad (28)$$

To obtain a more explicit formula, we use an elementary decomposition of the Radon-Nikodym derivative inverse M_T^{-1} . This decomposition expresses it as the product of a random variable which is $\sigma(W_T, v)$ -measurable and a random variable conditionally independent from W_T given v . Since θ is a stochastic process, we do not have that \overline{W}_T is a normal random variable, but rather that it is a mixed normal random variable. The elementary idea is to normalize

\overline{W}_T by $\sqrt{\int_0^T \theta_s^2 ds}$, so that $(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}})$ is a centered normal random vector under \mathbb{P} . We define the correlation under \mathbb{P} between W_T and $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ as ρ , i.e., $\rho = \text{Cor}_{\mathbb{P}}(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}})$.

Lemma 10. *Under **Assumption B**, we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . We can also show that $\rho = \frac{1}{T} \mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s ds}{\sqrt{\int_0^T \theta_s^2 ds}} \right]$. Moreover, there exists a standard normal random variable \widetilde{W} under \mathbb{P} , which is independent of W_T , and such that \overline{W}_T , when normalized, can be reexpressed a.s. as*

$$\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}} = \rho \frac{W_T}{\sqrt{T}} + \sqrt{1 - \rho^2} \widetilde{W}. \quad (29)$$

This can be reexpressed a.s. as

$$\overline{W}_T = \alpha W_T + \widetilde{\alpha} \widetilde{W}, \quad (30)$$

where $\alpha = \rho \sqrt{T^{-1} \int_0^T \theta_s^2 ds}$ a.s. and $\widetilde{\alpha} = \sqrt{(1 - \rho^2) \int_0^T \theta_s^2 ds}$ a.s.. If we define $\widetilde{\theta}_t = \frac{\theta_t - \alpha}{\widetilde{\alpha}}$, we can reexpress \widetilde{W} a.s. as

$$\widetilde{W} = \int_0^T \widetilde{\theta}_s dW_s. \quad (31)$$

Moreover, $\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} . Finally, the conditional distribution of $\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds$ given v , i.e., $\mathcal{D}(\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds | v)$, is standard normal under \mathbb{Q} .

Our main result is the next theorem, which gives a more explicit formula to $\mathbb{P}(T_b^Y \leq T | W_T, v)$.

Theorem 11. *Under **Assumption B**, we have*

$$\begin{aligned} \mathbb{P}(T_b^Y \leq T | W_T, v) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\widetilde{\alpha} \widetilde{W}) | W_T, v]. \end{aligned} \quad (32)$$

We first calculate $\mathbb{Q}(T_b^Y \leq T | W_T, v)$.

Lemma 12. *Under **Assumption B**, we have*

$$\mathbb{Q}(T_b^Y \leq T | W_T, v) = \exp\left(-\frac{2b(b - Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} + \mathbf{1}_{\{Y_T > b\}}. \quad (33)$$

The next theorem gives a formula based on a theoretical assumption (34).

Theorem 13. *We assume that **Assumption B** and the following assumption*

$$\begin{aligned} & \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha}\tilde{W})|W_T, v] \\ &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}}|W_T, v] \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W})|W_T, v] \end{aligned} \quad (34)$$

holds. Then, we have

$$\begin{aligned} \mathbb{P}(T_b^Y \leq T|W_T, v) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \left(\exp\left(-\frac{2b(b-Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} + \mathbf{1}_{\{Y_T > b\}} \right) \\ &\quad \times \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned} \quad (35)$$

Finally, we get $P_b^Y(T)$ in the next corollary by integrating $\mathbb{P}(T_b^Y \leq T|W_T, v)$ with respect to the value of (W_T, v) . We define the arrival space and cdf of v as respectively Π_v and P_v . Moreover, we define y_u, y_b, y_θ , etc. following the above definitions when integrating with respect to $y \in \Pi_v$.

Corollary 14. *Under **Assumption B**, we have*

$$\begin{aligned} P_b^Y(T) &= 1 - \phi\left(\frac{b-u_T}{\sqrt{T}}\right) \\ &\quad + \int_{-\infty}^{b-u_T} \int_{\Pi_v} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-y_\alpha x + \frac{1}{2} \int_0^T y_{\theta,s}^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha}\tilde{W})|W_T = x, v = y] dx dP_v(y). \end{aligned} \quad (36)$$

If we further assume (34), we have

$$\begin{aligned} P_b^Y(T) &= 1 - \phi\left(\frac{b-u_T}{\sqrt{T}}\right) \\ &\quad + \int_{-\infty}^{b-u_T} \int_{\Pi_v} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-y_\alpha x + \frac{1}{2} \int_0^T y_{\theta,s}^2 ds\right) \\ &\quad \times p_v(y) \exp\left(-\frac{2y_b(y_b - y_{u,T} - x)}{T}\right) \\ &\quad \times \exp\left(y_{\tilde{\alpha}} \int_0^T y_{\tilde{\theta},s} y_{\theta,s} ds\right) \mathcal{L}_N(y_{\tilde{\alpha}}) dx dP_v(y). \end{aligned} \quad (37)$$

2.3. Two-sided time-varying boundary case

In this section, we consider the case when the two-sided boundary and the drift are nonrandom time-varying and the variance is nonrandom.

We first give the definition of the set of two-sided boundary functions.

Definition 2.7. We define the set of two-sided boundary functions as $\mathcal{I} = \mathcal{G} \times \mathcal{G}$.

We now give the definition of the FPT to a two-sided boundary.

Definition 2.8. We define the FPT of an \mathbf{F} -adapted continuous process Z to a two-sided boundary $(g, h) \in \mathcal{I}$ satisfying $g_0 \leq Z_0 \leq h_0$ as

$$T_{g,h}^Z = \inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \geq g_t \text{ or } Z_t \leq h_t\}. \quad (38)$$

We have that Z is a continuous and \mathbf{F} -adapted stochastic process and $\inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \geq g_t \text{ or } Z_t \leq h_t\} = \inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \in G\}$ where $G = \{(t, u) \in \mathbb{R}^+ \times \mathbb{R} \text{ s.t. } u \geq g_t \text{ or } u \leq h_t\}$ is an open subset of \mathbb{R}^2 . Thus, the FPT $T_{g,h}^Z$ is an \mathbf{F} -stopping time by Theorem I.1.27 (p. 7) in Jacod et al. (2003). We can rewrite the boundary crossing probability $P_{g,h}^Z$ as the cdf of $T_{g,h}^Z$, i.e.,

$$P_{g,h}^Z(t) = \mathbb{P}(T_{g,h}^Z \leq t) \text{ for any } t \geq 0. \quad (39)$$

We assume that $g_0 < \mu_0 < h_0$. To apply the Girsanov theorem to the two-sided boundary case, we cannot use two different drifts since the process Z_t is unique in Definition 2.8. Thus, we have to restrict the class of boundary functions as we will assume that the deviation of g from its starting value is equal to the sum of the deviation of h from its starting value and a linear term, i.e., there exists $\beta \in \mathbb{R}$ such that $h_t - h_0 = g_t - g_0 + \beta t$. Thus, we can rewrite the FPT to a two-sided time-varying boundary as an equivalent FPT to a two-sided boundary with one constant boundary and one linear boundary. More specifically, if we define the new drift as $u_t = \frac{\mu_t - \mu_0 - g_t + g_0}{\sigma}$, the new process as $Y_t = u_t + W_t$, the new constant boundary as $b = \frac{\mu_0 - g_0}{\sigma}$ and the new linear boundary as $c_t = \frac{h_0 - \mu_0 + \beta t}{\sigma}$, we observe that the FPT (38) may be rewritten as $T_{g,h}^Z = T_{b,c}^Y$.

Assumption C. We assume that $u \neq 0$, i.e., that there exists a $t \in [0, T]$ such that $u_t \neq 0$. We also assume that u is absolutely continuous on $[0, T]$, i.e., there exists a nonrandom function $\theta : [0, T] \rightarrow \mathbb{R}$ with $u_t = \int_0^t \theta_s ds$. Finally, we assume that $\int_0^T \theta_s^2 ds < \infty$.

By **Assumption C**, M satisfies Novikov's condition and thus is a positive martingale. We embed this result and its implications on an equivalent probability measure \mathbb{Q} by the Girsanov theorem in the following lemma.

Lemma 15. *Under **Assumption C**, we have that M is a positive martingale. Thus, we can consider an equivalent probability measure \mathbb{Q} such that the Radon-Nikodym derivative is defined as $\frac{d\mathbb{Q}}{d\mathbb{P}} = M_T$. Finally, Y is a standard Wiener process under \mathbb{Q} .*

The next proposition reexpresses $\mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T)$ under \mathbb{Q} . We define \overline{W}_t as

$$\overline{W}_t = \int_0^t \theta_s dW_s. \quad (40)$$

Proposition 16. *Under Assumption C, we have*

$$\mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T) = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} M_T^{-1} | W_T]. \quad (41)$$

This can be reexpressed as

$$\mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T) = \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} | W_T, \overline{W}_T] | W_T]. \quad (42)$$

To obtain a more explicit formula, we elementary decompose the Radon-Nikodym derivative inverse M_T^{-1} as the product of a $\sigma(W_T)$ -measurable random variable and a random variable independent from W_T . We define the correlation under \mathbb{P} between W_T and \overline{W}_T as ρ , i.e., $\rho = \text{Cor}_{\mathbb{P}}(W_T, \overline{W}_T)$.

Lemma 17. *Under Assumption C, we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . We can also show that $\rho = \frac{\int_0^T \theta_s ds}{T \int_0^T \theta_s^2 ds}$. Moreover, there exists a standard normal random variable \widetilde{W} under \mathbb{P} which is independent of W_T and such that \overline{W}_T when normalized can be reexpressed as*

$$\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}} = \rho \frac{W_T}{\sqrt{T}} + \sqrt{1 - \rho^2} \widetilde{W}. \quad (43)$$

This can be reexpressed as

$$\overline{W}_T = \alpha W_T + \widetilde{\alpha} \widetilde{W}, \quad (44)$$

where $\alpha = \rho \sqrt{T^{-1} \int_0^T \theta_s^2 ds}$ and $\widetilde{\alpha} = \sqrt{(1 - \rho^2) \int_0^T \theta_s^2 ds}$. If we define $\widetilde{\theta}_t = \frac{\theta_s - \alpha}{\widetilde{\alpha}}$, we can reexpress \widetilde{W} as

$$\widetilde{W} = \int_0^T \widetilde{\theta}_s dW_s. \quad (45)$$

Finally, $\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} .

Our main result is the next theorem.

Theorem 18. *Under Assumption C, we have*

$$\begin{aligned} \mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} \exp(-\widetilde{\alpha} \widetilde{W}) | W_T]. \end{aligned} \quad (46)$$

We first calculate $\mathbb{Q}(\mathbb{T}_{b,c}^Y \leq T|W_T)$ whose explicit formula is given in the following lemma. This reexpresses Anderson (1960) (Theorem 4.2, pp. 178-179) under \mathbb{Q} , which considers the linear case $\mu_t = \gamma t$, $g_t = at + b$, $h_t = ct + d$ and $\sigma = 1$ under \mathbb{P} . He obtains that

$$\mathbb{P}(\mathbb{T}_{g,h}^Z \leq T|W_T = x) = \sum_{j=1}^{\infty} p_{g,h}^Z(j|x) \mathbf{1}_{\{x \in [h_T - \mu_T, g_T - \mu_T]\}} + \mathbf{1}_{\{x \notin [h_T - \mu_T, g_T - \mu_T]\}}.$$

Here, $p_{g,h}^Z(j|x)$ is defined as

$$\begin{aligned} p_{g,h}^Z(j|x) &= \exp\left(-\frac{2}{T}(j\delta_0 + h_0)(j\delta_T + (h_T - \mu_T) - x)\right) \\ &\quad + \exp\left(-\frac{2j}{T}(j\delta_0\delta_T + \delta_0((h_T - \mu_T) - x) - \delta_T h_0)\right) \\ &\quad + \exp\left(-\frac{2}{T}(j\delta_0 - g_0)(j\delta_T - ((g_T - \mu_T) - x))\right) \\ &\quad + \exp\left(-\frac{2j}{T}(j\delta_0\delta_T - \delta_0((g_T - \mu_T) - x) + \delta_T g_0)\right) \end{aligned}$$

for any $j \in \mathbb{N}_*$, $x \in [h_T - \mu_T, g_T - \mu_T]$ and the difference between g and h is defined as $\delta_t = \delta_t(g, h) = g_t - h_t$ for any $t \in [0, T]$.

Lemma 19. *Under **Assumption C**, we have*

$$\mathbb{Q}(\mathbb{T}_{b,c}^Y \leq T|W_T) = \sum_{j=1}^{\infty} q_{b,c}^Y(j|Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} + \mathbf{1}_{\{Y_T \notin [c_T, b_T]\}}. \quad (47)$$

Here, $q_{b,c}^Y(j|x)$ is defined as

$$\begin{aligned} q_{b,c}^Y(j|x) &= \exp\left(-\frac{2}{T}(j\delta_0 + c_0)(j\delta_T + (c_T - x))\right) \\ &\quad + \exp\left(-\frac{2j}{T}(j\delta_0\delta_T + \delta_0(c_T - x) - \delta_T c_0)\right) \\ &\quad + \exp\left(-\frac{2}{T}(j\delta_0 - b_0)(j\delta_T - (b_T - x))\right) \\ &\quad + \exp\left(-\frac{2j}{T}(j\delta_0\delta_T - \delta_0((b_T - x) + \delta_T b_0)\right), \end{aligned}$$

for any $j \in \mathbb{N}_*$, $x \in [c_T, b_T]$ and $\delta_t = \delta_t(b, c)$.

The next theorem gives a formula based on a strong theoretical assumption (48).

Theorem 20. *We assume that **Assumption C** and the following assumption*

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} \exp(-\widetilde{\alpha W})|W_T] &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}}|W_T] \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\exp(-\widetilde{\alpha W})|W_T] \end{aligned} \quad (48)$$

holds. Then, we have

$$\begin{aligned} \mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \left(\sum_{j=1}^{\infty} q_{b,c}^Y(j | Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} \right. \\ &\quad \left. + \mathbf{1}_{\{Y_T \notin [c_T, b_T]\}} \right) \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned} \quad (49)$$

Finally, we get $P_{b,c}^Y(T)$ in the next corollary by integrating $\mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T)$ with respect to the value of W_T .

Corollary 21. *Under Assumption C, we have*

$$\begin{aligned} P_{b,c}^Y(T) &= 1 - \phi\left(\frac{b_T - u_T}{\sqrt{T}}\right) + \phi\left(\frac{c_T - u_T}{\sqrt{T}}\right) \\ &\quad + \int_{c_T - u_T}^{b_T - u_T} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-\alpha x + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T = x] dx. \end{aligned} \quad (50)$$

If we further assume (48), we have

$$\begin{aligned} P_{b,c}^Y(T) &= 1 - \phi\left(\frac{b_T - u_T}{\sqrt{T}}\right) + \phi\left(\frac{c_T - u_T}{\sqrt{T}}\right) \\ &\quad + \int_{c_T - u_T}^{b_T - u_T} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-\alpha x + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \left(\sum_{j=1}^{\infty} q_{b,c}^{x+u_T}(j | x + u_T) \mathbf{1}_{\{x \in [c_T - u_T, b_T - u_T]\}} \right. \\ &\quad \left. + \mathbf{1}_{\{x \notin [c_T - u_T, b_T - u_T]\}} \right) \\ &\quad \times \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}) dx. \end{aligned} \quad (51)$$

In the particular case when the boundaries and the drift are linear, and the standard deviation is equal to unity, Corollary 21 reduces to Anderson (1960) (Theorem 4.3, p. 180).

3. Proofs

Our proofs rely on an elementary application of the Girsanov theorem.

3.1. One-sided time-varying boundary case

We start with the proofs in the case when the one-sided boundary and the drift are nonrandom time-varying and the variance is nonrandom.

The proof of Proposition 2 is based on Lemma 1 and its consequence in the particular case $X = \mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}$ in which E_T is a $\sigma(W_T)$ -measurable event.

Proof of Proposition 2. By definition of the conditional probability, Equation (10) can be rewritten formally as

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} | W_T]. \quad (52)$$

For any $\sigma(W_T)$ -measurable event E_T , we can use a change of probability in the expectation by Lemma 1 along with **Assumption A** and we obtain that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} \mathbf{1}_{E_T}]. \quad (53)$$

We can deduce Equation (52) from Equation (53) by definition of the conditional expectation. By definition of the conditional probability, Equation (11) can be rewritten formally as

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T] = \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, \overline{W}_T] | W_T]. \quad (54)$$

By definition of the conditional expectation, we can deduce what follows. If we can show that for any E_T which is $\sigma(W_T)$ -measurable that

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}] \\ &= \mathbb{E}_{\mathbb{P}}[\mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, \overline{W}_T] | W_T] \mathbf{1}_{E_T}], \end{aligned} \quad (55)$$

then Equation (54) holds. Let E_T be a $\sigma(W_T)$ -measurable event. By Lemma 1 along with **Assumption A**, we can use a change of probability in the expectation and we obtain that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1}]. \quad (56)$$

Then, we have by the law of total expectation that

$$\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1}] = \mathbb{E}_{\mathbb{Q}}[\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} | W_T, \overline{W}_T]]. \quad (57)$$

Since $\mathbf{1}_{E_T}$ and M_T^{-1} are $\sigma(W_T, \overline{W}_T)$ -measurable random variables, we can pull them out of the conditional expectation and deduce that

$$\begin{aligned} & \mathbb{E}_{\mathbb{Q}}[\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} | W_T, \overline{W}_T]] \\ &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{E_T} M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, \overline{W}_T]]. \end{aligned} \quad (58)$$

If we use Equations (56)-(57)-(58), we can deduce that Equation (55) holds. \square

In what follows, we give the proof of Lemma 3. It is based on the fact that (W_T, \overline{W}_T) is a centered normal random vector under \mathbb{P} so that we can rewrite \overline{W}_T as $\overline{W}_T = \alpha W_T + \tilde{\alpha} \widetilde{W}$.

Proof of Lemma 3. By **Assumption A**, we have that \overline{W}_T is well-defined and $\int_0^T \theta_s^2 ds < \infty$ thus we can deduce that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . Since (W_T, \overline{W}_T) is a centered normal random vector under \mathbb{P} , there exists a standard normal random variable \widetilde{W} under \mathbb{P} which is independent of W_T and such that Equation (12) holds. Then, we can calculate that

$$\begin{aligned} \rho &= \text{Cor}_{\mathbb{P}}(W_T, \int_0^T \theta_s dW_s) \\ &= \frac{\text{Cov}_{\mathbb{P}}(W_T, \int_0^T \theta_s dW_s)}{\text{Var}_{\mathbb{P}}(W_T) \text{Var}_{\mathbb{P}}(\int_0^T \theta_s dW_s)} \\ &= \frac{\int_0^T \theta_s ds}{T \int_0^T \theta_s^2 ds}. \end{aligned}$$

Here, we use the definition of ρ and Equation (9) in the first equality, the fact that $\theta \neq 0$ by **Assumption A** in the second equality, and the Itô isometry in the last equality. Equation (13) can be deduced directly from Equation (12). Moreover, we can reexpress \widetilde{W} as

$$\begin{aligned} \widetilde{W} &= \frac{1}{\tilde{\alpha}} (\overline{W}_T - \alpha W_T) \\ &= \int_0^T \frac{\theta_s - \alpha}{\tilde{\alpha}} dW_s \\ &= \int_0^T \tilde{\theta}_s dW_s. \end{aligned}$$

Here, we use Equation (13) in the first equality, Equation (9) in the second equality and the definition of $\tilde{\theta}_t$ in the last equality. Finally, we can deduce that $\widetilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} by its expression (14) and since by Lemma 1 along with **Assumption A**, Y is a Wiener process under \mathbb{Q} . \square

We provide now the proof of Theorem 4 which is based on Lemma 3.

Proof of Theorem 4. We can reexpress M_T as

$$\begin{aligned} M_T &= \exp\left(\int_0^T \theta_s dW_s - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &= \exp\left(\overline{W}_T - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \end{aligned}$$

$$\begin{aligned}
&= \exp\left(\alpha W_T + \widetilde{\alpha}\widetilde{W} - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\
&= \exp\left(\alpha W_T - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \exp(\widetilde{\alpha}\widetilde{W}). \tag{59}
\end{aligned}$$

Here, we use Equation (8) in the first equality, Equation (9) in the second equality, Equation (13) from Lemma 3 in the third equality and algebraic manipulation in the last equality. Then, we have

$$\begin{aligned}
\mathbb{P}(T_b^Y \leq T | W_T) &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} | W_T] \\
&= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\
&\quad \times \exp(-\widetilde{\alpha}\widetilde{W}) | W_T] \\
&= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\
&\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\widetilde{\alpha}\widetilde{W}) | W_T].
\end{aligned}$$

Here, we use Equation (10) from Proposition 2 along with **Assumption A** in the first equality, Equation (73) in the second equality, and the fact that W_T is a $\sigma(W_T)$ -measurable random variable in the third equality. Thus, we have shown Equation (15). \square

We now give the proof which reexpresses Malmquist (1954) (Theorem 1, p. 526) under \mathbb{Q} .

Proof of Lemma 5. By definition of the conditional probability, Equation (17) can be rewritten formally as

$$\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T] = \exp\left(-\frac{2b(b - Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} + \mathbf{1}_{\{Y_T > b\}}. \tag{60}$$

By Lemma 1 along with **Assumption A**, Y is a Wiener process under \mathbb{Q} . Then, we have by Malmquist (1954) (Theorem 1, p. 526) that Equation (60) holds. \square

We provide now the proof of Theorem 6, which is based on Lemma 5.

Proof of Theorem 6. We have

$$\begin{aligned}
\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\widetilde{\alpha}\widetilde{W}) | W_T] &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T] \mathbb{E}_{\mathbb{Q}}[\exp(-\widetilde{\alpha}\widetilde{W}) | W_T] \\
&= \left(\exp\left(-\frac{2b(b - Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} + \mathbf{1}_{\{Y_T > b\}}\right) \\
&\quad \times \mathbb{E}_{\mathbb{Q}}[\exp(-\widetilde{\alpha}\widetilde{W}) | W_T], \tag{61}
\end{aligned}$$

where we use Assumption (19) in the first equality, Equation (17) from Lemma 5 along with **Assumption A** in the second equality. Finally, we have

$$\begin{aligned}
\mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W})|W_T] &= \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W})] \\
&= \exp\left(\tilde{\alpha}\int_0^T \tilde{\theta}_s\theta_s ds\right) \\
&\quad \times \mathbb{E}_{\mathbb{Q}}\left[\exp\left(-\tilde{\alpha}\left(\tilde{W} + \int_0^T \tilde{\theta}_s\theta_s ds\right)\right)\right] \\
&= \exp\left(\tilde{\alpha}\int_0^T \tilde{\theta}_s\theta_s ds\right)\mathbb{E}_{\mathbb{P}}[\exp(-\tilde{\alpha}N)] \\
&= \exp\left(\tilde{\alpha}\int_0^T \tilde{\theta}_s\theta_s ds\right)\mathcal{L}_N(\tilde{\alpha}). \tag{62}
\end{aligned}$$

Here, we use the fact that \tilde{W} is independent from W_T in the first equality, algebraic manipulation in the second equality, the fact that $\tilde{W} + \int_0^T \tilde{\theta}_s\theta_s ds$ is a standard normal variable under \mathbb{Q} by Lemma 3 along with **Assumption A** in the third equality, and Equation (18) in the last equality. We can deduce Equation (20) from Equations (15), (61) and (62). \square

Finally, the proof of Corollary 7 follows the steps of Equations (3) in Wang et al. (1997) (p. 55).

Proof of Corollary 7. We can calculate that

$$\begin{aligned}
P_b^Y(T) &= \int_{-\infty}^{\infty} \mathbb{P}(T_b^Y \leq T | W_T = x) \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx \\
&= 1 - \phi\left(\frac{b-u_T}{\sqrt{T}}\right) \\
&\quad + \int_{-\infty}^{b-u_T} \mathbb{P}(T_b^Y \leq T | W_T = x) \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx \\
&= 1 - \phi\left(\frac{b-u_T}{\sqrt{T}}\right) \\
&\quad + \int_{-\infty}^{b-u_T} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-\alpha x + \frac{1}{2}\int_0^T \theta_s^2 ds\right) \\
&\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\tilde{\alpha}\tilde{W}) | W_T = x] dx.
\end{aligned}$$

Here, we use Equation (6) and regular conditional probability in the first equality, the fact that $\mathbb{P}(T_b^Y \leq T | W_T = x) = 1$ for any $x \geq b - u_T$ in the second equality, and Equation (15) in the third equality. We have thus shown Equation (21). Equation (22) can be shown following the same first two equalities and using Assumption (20) in the third equality. \square

3.2. One-sided stochastic boundary process case

We continue with the proofs in the case when the one-sided boundary and the drift are stochastic processes and the variance is random.

The elementary idea in the proofs of this section is to condition by both W_T and v , i.e., to derive results of the form $\mathbb{P}(T_b^Y \leq T | W_T, v)$. The proof of Proposition 9 is based on Lemma 8 and its consequence in the particular case $X = \mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}$, in which E_T is a $\sigma(W_T, v)$ -measurable event.

Proof of Proposition 9. By definition of the conditional probability, Equation (27) can be rewritten formally as

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, v] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} | W_T, v]. \quad (63)$$

For any $\sigma(W_T, v)$ -measurable event E_T , we can use a change of probability in the expectation by Lemma 8 along with **Assumption B** and we obtain that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} \mathbf{1}_{E_T}]. \quad (64)$$

We can deduce Equation (63) from Equation (64) by definition of the conditional expectation. By definition of the conditional probability, Equation (28) can be rewritten formally as

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, v] \\ &= \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, \overline{W}_T, v] | W_T, v]. \end{aligned} \quad (65)$$

By definition of the conditional expectation, we can deduce what follows. If we can show that for any E_T which is $\sigma(W_T, v)$ -measurable that

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}] \\ &= \mathbb{E}_{\mathbb{P}}\left[\mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, \overline{W}_T, v] | W_T, v] \mathbf{1}_{E_T}\right], \end{aligned} \quad (66)$$

then Equation (65) holds. Let E_T be a $\sigma(W_T, v)$ -measurable event. By Lemma 8 along with **Assumption B**, we can use a change of probability in the expectation and we obtain that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T}] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1}]. \quad (67)$$

Then, we have by the law of total expectation that

$$\begin{aligned} & \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1}] = \\ & \mathbb{E}_{\mathbb{Q}}\left[\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} | W_T, \overline{W}_T, v]\right]. \end{aligned} \quad (68)$$

Since $\mathbf{1}_{E_T}$ and M_T^{-1} are $\sigma(W_T, \overline{W}_T, v)$ -measurable random variables, we can pull them out of the conditional expectation and deduce that

$$\begin{aligned} & \mathbb{E}_{\mathbb{Q}} \left[\mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{\tau_b^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} \mid W_T, \overline{W}_T, v \right] \right] \\ &= \mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{E_T} M_T^{-1} \mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{\tau_b^Y \leq T\}} \mid W_T, \overline{W}_T, v \right] \right]. \end{aligned} \quad (69)$$

If we use Equations (67)-(68)-(69), we can deduce that Equation (66) holds. \square

In what follows, we give the proof of Lemma 10. It is based on the fact that $(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}})$ is a centered normal random vector under \mathbb{P} .

Proof of Lemma 10. By **Assumption B**, we can deduce that $0 < \int_0^T \theta_s^2 ds < \infty$ a.s.. Thus, we can normalize \overline{W}_T by $\sqrt{\int_0^T \theta_s^2 ds}$ a.s. and we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a mixed normal random variable a.s. by definition. We have that its conditional mean under \mathbb{P} is a.s. equal to

$$\begin{aligned} \mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \mid v \right] &= \frac{1}{\sqrt{\int_0^T \theta_s^2 ds}} \mathbb{E}_{\mathbb{P}} \left[\int_0^T \theta_s dW_s \mid v \right] \\ &= 0. \end{aligned} \quad (70)$$

Here, we use the fact that $\frac{1}{\sqrt{\int_0^T \theta_s^2 ds}}$ is $\sigma(v)$ -measurable in the first equality, and the fact that $\int_0^T \theta_s dW_s$ is a.s. a martingale since $\int_0^T \theta_s^2 ds < \infty$ a.s. in the second equality. We also have that its conditional variance under \mathbb{P} is a.s. equal to

$$\begin{aligned} \text{Var}_{\mathbb{P}} \left(\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \mid v \right) &= \mathbb{E}_{\mathbb{P}} \left[\left(\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \right)^2 \mid v \right] \\ &= \frac{1}{\int_0^T \theta_s^2 ds} \mathbb{E}_{\mathbb{P}} \left[\left(\int_0^T \theta_s dW_s \right)^2 \mid v \right] \\ &= 1. \end{aligned} \quad (71)$$

Here, we use Equation (70) in the first equality, the fact that $\frac{1}{\sqrt{\int_0^T \theta_s^2 ds}}$ is $\sigma(v)$ -measurable in the second equality, and the Itô isometry in the third equality. Since its conditional mean and conditional variance are nonrandom, we obtain that its mean under \mathbb{P} is equal to $\mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \right] = \mathbb{E}_{\mathbb{P}} \left[\mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \mid v \right] \right] = 0$ by the law of total expectation and Equation (70), and similarly that its variance is equal to 1 by the law of total expectation and Equation (71). Thus, we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . Since $(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}})$ is a centered normal random vector under \mathbb{P} , there exists a standard normal random variable \widetilde{W} under \mathbb{P} which is independent of W_T and such that Equation

(29) holds. Then, we can calculate that the covariance between W_T and $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ under \mathbb{P} is equal to

$$\begin{aligned}
\text{Cov}_{\mathbb{P}}\left(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}\right) &= \text{Cov}_{\mathbb{P}}\left(W_T, \frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}}\right) \\
&= \mathbb{E}_{\mathbb{P}}\left[W_T \frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}}\right] \\
&= \mathbb{E}_{\mathbb{P}}\left[\mathbb{E}\left[W_T \frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \middle| v\right]\right] \\
&= \mathbb{E}_{\mathbb{P}}\left[\frac{1}{\sqrt{\int_0^T \theta_s^2 ds}} \mathbb{E}\left[W_T \int_0^T \theta_s dW_s \middle| v\right]\right] \\
&= \mathbb{E}_{\mathbb{P}}\left[\frac{\int_0^T \theta_s ds}{\sqrt{\int_0^T \theta_s^2 ds}}\right]. \tag{72}
\end{aligned}$$

Here, we use Equation (9) in the first equality, Equation (70) in the second equality, the law of total expectation in the third equality, the fact that $\frac{1}{\sqrt{\int_0^T \theta_s^2 ds}}$ is $\sigma(v)$ -measurable in the fourth equality, and the Itô isometry in the last equality. Now, we can calculate that the correlation between W_T and $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ under \mathbb{P} is equal to

$$\begin{aligned}
\rho &= \text{Cor}_{\mathbb{P}}\left(W_T, \frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}}\right) \\
&= \frac{\text{Cov}_{\mathbb{P}}\left(W_T, \frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}}\right)}{\text{Var}_{\mathbb{P}}(W_T) \text{Var}_{\mathbb{P}}\left(\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}}\right)} \\
&= \frac{1}{T} \mathbb{E}_{\mathbb{P}}\left[\frac{\int_0^T \theta_s ds}{\sqrt{\int_0^T \theta_s^2 ds}}\right].
\end{aligned}$$

Here, we use the definition of ρ and Equation (9) in the first equality, and Equations (71) and (72) in the last equality. Equation (30) can be deduced directly from Equation (29). Moreover, we can reexpress \widetilde{W} as

$$\begin{aligned}
\widetilde{W} &= \frac{1}{\widetilde{\alpha}}(\overline{W}_T - \alpha W_T) \\
&= \int_0^T \frac{\theta_s - \alpha}{\widetilde{\alpha}} dW_s
\end{aligned}$$

$$= \int_0^T \tilde{\theta}_s dW_s,$$

where we use Equation (30) in the first equality, Equation (9) in the second equality and the definition of $\tilde{\theta}_t$ in the last equality. Moreover, we can deduce that $\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} . This is due to its expression (31) and since by Lemma 8 along with **Assumption B**, Y is a Wiener process under \mathbb{Q} . Finally, $\mathcal{D}(\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds | v)$ is standard normal under \mathbb{Q} by Equation (31). \square

We provide now the proof of Theorem 11, which is based on Lemma 10.

Proof of Theorem 11. We can reexpress M_T as

$$\begin{aligned} M_T &= \exp\left(\int_0^T \theta_s dW_s - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &= \exp\left(\bar{W}_T - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &= \exp\left(\alpha W_T + \tilde{\alpha} \tilde{W} - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &= \exp\left(\alpha W_T - \frac{1}{2} \int_0^T \theta_s^2 ds\right) \exp(\tilde{\alpha} \tilde{W}). \end{aligned} \quad (73)$$

Here, we use Equation (8) in the first equality, Equation (9) in the second equality, Equation (30) from Lemma 10 in the third equality, and algebraic manipulation in the last equality. Then, we have

$$\begin{aligned} \mathbb{P}(T_b^Y \leq T | W_T, v) &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} M_T^{-1} | W_T, v] \\ &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \exp\left(-\tilde{\alpha} \tilde{W}\right) | W_T, v] \\ &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp\left(-\tilde{\alpha} \tilde{W}\right) | W_T, v]. \end{aligned}$$

Here, we use Equation (27) from Proposition 9 along with **Assumption B** in the first equality, Equation (73) in the second equality, the fact that W_T and θ_t for any $t \in [0, T]$ are $\sigma(W_T, v)$ -measurable random variables in the third equality. Thus, we have shown Equation (32). \square

We now give the proof of Lemma 12.

Proof of Lemma 12. By definition of the conditional probability, Equation (33) can be rewritten formally as

$$\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\tau_b^Y \leq T\}} | W_T, v] = \exp\left(-\frac{2b(b-Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} + \mathbf{1}_{\{Y_T > b\}}. \quad (74)$$

By Lemma 8 along with **Assumption B**, Y is a Wiener process under \mathbb{Q} . Then, we have by Malmquist (1954) (Theorem 1, p. 526) that Equation (74) holds. \square

We provide now the proof of Theorem 13, which is based on Lemma 12.

Proof of Theorem 13. We have

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\tau_b^Y \leq T\}} \exp(-\tilde{\alpha}\tilde{W}) | W_T, v] &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\tau_b^Y \leq T\}} | W_T, v] \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | W_T, v] \\ &= \left(\exp\left(-\frac{2b(b-Y_T)}{T}\right) \mathbf{1}_{\{Y_T \leq b\}} \right. \\ &\quad \left. + \mathbf{1}_{\{Y_T > b\}} \right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | W_T, v], \end{aligned} \quad (75)$$

where we use Assumption (34) in the first equality, Equation (33) from Lemma 12 along with **Assumption B** in the second equality. Finally, we have

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | W_T, v] &= \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | v] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}\left[\exp\left(-\tilde{\alpha}\left(\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds\right)\right) | v\right] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}\left[\exp\left(-\tilde{\alpha}\left(\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds\right)\right)\right] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathbb{E}_{\mathbb{P}}[\exp(-\tilde{\alpha}N)] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned} \quad (76)$$

Here, we use the fact that \tilde{W} is independent from W_T in the first equality, the fact that θ_t and $\tilde{\theta}_t$ for any $t \in [0, T]$ are $\sigma(v)$ -measurable random variables in the second equality, the fact that $\mathcal{D}(\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds | v)$ is standard normal under \mathbb{Q} by Lemma 10 along with **Assumption B** in the third equality, the fact that

$\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} by Lemma 10 along with **Assumption B** in the fourth equality, and Equation (18) in the last equality. We can deduce Equation (35) from Equations (32), (75) and (76). \square

Finally, we get $P_b^Y(T)$ in the next corollary, by integrating $\mathbb{P}(T_b^Y \leq T | W_T, v)$ with respect to the value of (W_T, v) .

Proof of Corollary 14. We can calculate that

$$\begin{aligned}
P_b^Y(T) &= \int_{-\infty}^{\infty} \int_{\Pi_v} \mathbb{P}(T_b^Y \leq T | W_T = x, v = y) \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx dP_v(y) \\
&= 1 - \phi\left(\frac{b - u_T}{\sqrt{T}}\right) \\
&\quad + \int_{-\infty}^{b - u_T} \int_{\Pi_v} \mathbb{P}(T_b^Y \leq T | W_T = x, v = y) \\
&\quad \times \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx dP_v(y) \\
&= 1 - \phi\left(\frac{b - u_T}{\sqrt{T}}\right) + \\
&\quad \int_{-\infty}^{b - u_T} \int_{\Pi_v} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-y_\alpha x + \frac{1}{2} \int_0^T y_{\theta,s}^2 ds\right) \\
&\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_b^Y \leq T\}} \exp(-\widetilde{\alpha} \widetilde{W}) | W_T = x, v = y] dx dP_v(y).
\end{aligned}$$

Here, we use Equation (24), regular conditional probability and the fact that W_T and v are independent in the first equality, the fact that $\mathbb{P}(T_b^Y \leq T | W_T = x) = 1$ for any $x \geq b - u_T$ in the second equality, and Equation (32) in the third equality. We have thus shown Equation (36). Equation (37) can be shown following the same first two equalities and using Equation (35) in the third equality. \square

3.3. Two-sided time-varying boundary case

In this section, we consider the proofs in the case when the two-sided boundary and the drift are nonrandom time-varying and the variance is nonrandom.

The proof of Proposition 16 is based on Lemma 15.

Proof of Proposition 16. By definition of the conditional probability, Equation (41) can be rewritten formally as

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} M_T^{-1} | W_T]. \quad (77)$$

For any $\sigma(W_T)$ -measurable event E_T , we can use a change of probability in the expectation by Lemma 15 along with **Assumption C** and we obtain that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T}] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} M_T^{-1} \mathbf{1}_{E_T}]. \quad (78)$$

We can deduce Equation (77) from Equation (78) by definition of the conditional expectation. By definition of the conditional probability, Equation (42) can be rewritten formally as

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T] = \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T, \overline{W}_T] | W_T]. \quad (79)$$

By definition of the conditional expectation, we can deduce what follows. If we can show that for any E_T which is $\sigma(W_T)$ -measurable that

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T}] \\ &= \mathbb{E}_{\mathbb{P}}[\mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T, \overline{W}_T] | W_T] \mathbf{1}_{E_T}], \end{aligned} \quad (80)$$

then Equation (79) holds. Let E_T be a $\sigma(W_T)$ -measurable event. By Lemma 15 along with **Assumption C**, we obtain that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T}] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1}]. \quad (81)$$

Then we have by the law of total expectation that

$$\begin{aligned} & \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1}] \\ &= \mathbb{E}_{\mathbb{Q}}[\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} | W_T, \overline{W}_T]]. \end{aligned} \quad (82)$$

Since $\mathbf{1}_{E_T}$ and M_T^{-1} are $\sigma(W_T, \overline{W}_T)$ -measurable random variables, we can pull them out of the conditional expectation and deduce that

$$\begin{aligned} & \mathbb{E}_{\mathbb{Q}}[\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} | W_T, \overline{W}_T]] \\ &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{E_T} M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T, \overline{W}_T]]. \end{aligned} \quad (83)$$

If we use Equations (81)-(82)-(83), we can deduce that Equation (80) holds. \square

In what follows, we give the proof of Lemma 17.

Proof of Lemma 17. By **Assumption C**, we can deduce that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . Since (W_T, \overline{W}_T) is a centered normal random vector under \mathbb{P} , there exists a standard normal random variable \widetilde{W} under \mathbb{P} which is independent of W_T and such that Equation (43) holds. Using the same arguments from the proof of Lemma 3, we can calculate that

$$\rho = \frac{\int_0^T \theta_s ds}{T \int_0^T \theta_s^2 ds}.$$

Moreover, we can reexpress \widetilde{W} as

$$\widetilde{W} = \int_0^T \widetilde{\theta}_s dW_s.$$

Finally, we can deduce that $\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} . This is due to its expression (45) and since by Lemma 15 along with **Assumption C**, Y is a Wiener process under \mathbb{Q} . \square

We provide now the proof of Theorem 18, which is based on Lemma 17.

Proof of Theorem 18. By the same arguments from the proof of Theorem 4, we can reexpress M_T as

$$M_T = \exp(\alpha W_T - \frac{1}{2} \int_0^T \theta_s^2 ds) \exp(\widetilde{\alpha} \widetilde{W}).$$

Then, we have

$$\begin{aligned} \mathbb{P}(\mathbf{T}_{b,c}^Y \leq T | W_T) &= \exp(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbf{T}_{b,c}^Y \leq T\}} \exp(-\widetilde{\alpha} \widetilde{W}) | W_T]. \end{aligned}$$

Thus, we have shown Equation (46). \square

We now give the proof which reexpresses Anderson (1960) (Theorem 4.2, pp. 178-179) under \mathbb{Q} .

Proof of Lemma 19. By definition of the conditional probability, Equation (47) can be rewritten formally as

$$\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbf{T}_b^Y \leq T\}} | W_T] = \sum_{j=1}^{\infty} q_{b,c}^Y(j | Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} + \mathbf{1}_{\{Y_T \notin [c_T, b_T]\}}. \quad (84)$$

By Lemma 15 along with **Assumption C**, Y is a Wiener process under \mathbb{Q} . Then, we have by Anderson (1960) (Theorem 4.2, pp. 178-179) that Equation (84) holds. \square

We provide now the proof of Theorem 20, which is based on Lemma 19.

Proof of Theorem 20. We have

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbf{T}_{b,c}^Y \leq T\}} \exp(-\widetilde{\alpha} \widetilde{W}) | W_T] &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbf{T}_{b,c}^Y \leq T\}} | W_T] \mathbb{E}_{\mathbb{Q}}[\exp(-\widetilde{\alpha} \widetilde{W}) | W_T] \\ &= \left(\sum_{j=1}^{\infty} q_{b,c}^Y(j | Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} \right) \quad (85) \end{aligned}$$

$$+\mathbf{1}_{\{Y_T \notin [c_T, b_T]\}} \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W})|W_T],$$

where we use Assumption (48) in the first equality, and Equation (47) from Lemma 19 along with **Assumption C** in the second equality. Finally, we have

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W})|W_T] &= \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W})] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}(\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds))] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathbb{E}_{\mathbb{P}}[\exp(-\tilde{\alpha}N)] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned} \quad (86)$$

Here, we use the fact that \tilde{W} is independent from W_T in the first equality, algebraic manipulation in the second equality, the fact that $\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} by Lemma 17 along with **Assumption C** in the third equality, and Equation (18) in the last equality. We can deduce Equation (49) from Equations (46), (85) and (86). \square

Finally, we get $P_{b,c}^Y(T)$ in the next proof, by integrating $\mathbb{P}(T_{b,c}^Y \leq T|W_T)$ with respect to the value of W_T .

Proof of Corollary 21. We can calculate that

$$\begin{aligned} P_{b,c}^Y(T) &= \int_{-\infty}^{\infty} \mathbb{P}(T_{b,c}^Y \leq T|W_T = x) \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx \\ &= 1 - \phi\left(\frac{b_T - u_T}{\sqrt{T}}\right) + \phi\left(\frac{c_T - u_T}{\sqrt{T}}\right) \\ &\quad + \int_{c_T - u_T}^{b_T - u_T} \mathbb{P}(T_{b,c}^Y \leq T|W_T = x) \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx \\ &= 1 - \phi\left(\frac{b_T - u_T}{\sqrt{T}}\right) + \phi\left(\frac{c_T - u_T}{\sqrt{T}}\right) \\ &\quad + \int_{c_T - u_T}^{b_T - u_T} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-\alpha x + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \exp(-\tilde{\alpha}\tilde{W})|W_T = x] dx. \end{aligned}$$

Here, we use Equation (39) and regular conditional probability in the first equality, the fact that $\mathbb{P}(T_{b,c}^Y \leq T|W_T = x) = 1$ for any $x \geq b_T - u_T$ and any $x \leq c_T - u_T$ in the second equality, and Equation (46) in the third equality. We

have thus shown Equation (50). Equation (51) can be shown following the same first two equalities and using Equation (49) in the third equality. \square

Funding

The author was supported in part by Japanese Society for the Promotion of Science Grants-in-Aid for Scientific Research (B) 23H00807 and Early-Career Scientists 20K13470.

Supplementary Material

Supplement A: Results in the two-sided stochastic boundary process case

Supplement A gives the results in the two-sided stochastic boundary process case.

Supplement B: Proofs in the two-sided stochastic boundary process case

Supplement B collects the proofs in the two-sided stochastic boundary process case.

Supplement A gives the results in the two-sided stochastic boundary process case. Supplement B collects the proofs in the two-sided stochastic boundary process case.

Appendix A: Results in the two-sided stochastic boundary process case

In this appendix, we consider the case when the two-sided boundary and the drift are stochastic processes and the variance is random.

We first give the definition of the set of stochastic boundary processes.

Definition A.1. We define the set of stochastic two-sided boundary processes as $\mathcal{J} = \mathbb{R}^+ \times \Omega \rightarrow \mathbb{R}^2$ such that for any $(g, h) \in \mathcal{J}$ and $\omega \in \Omega$ we have $(g, h)(\omega) \in \mathcal{I}$ as well as g and h are \mathbf{F} -adapted.

We now give the definition of the FPT.

Definition A.2. We define the FPT of an \mathbf{F} -adapted continuous process Z to the two-sided boundary $(g, h) \in \mathcal{J}$ satisfying $g_0 \leq Z_0 \leq h_0 \forall \omega \in \Omega$ as

$$T_{g,h}^Z = \inf\{t \in \mathbb{R}^+ \text{ s.t. } Z_t \geq g_t \text{ or } Z_t \leq h_t\}. \quad (87)$$

We can rewrite $T_{g,h}^Z$ as the infimum of two \mathbf{F} -stopping times, i.e., $T_{g,h}^Z = \inf(T_h^Z, T_g^-)$. Thus, it is an \mathbf{F} -stopping time. We can rewrite the boundary crossing probability $P_{g,h}^Z$ as the cdf of $T_{g,h}^Z$, i.e.,

$$P_{g,h}^Z(t) = \mathbb{P}(T_{g,h}^Z \leq t) \text{ for any } t \geq 0. \quad (88)$$

We assume that μ is an \mathbf{F} -adapted stochastic process which satisfies $\mathbb{P}(g_0 < \mu_0 < h_0) = 1$. We also assume that the variance σ^2 is time-invariant, random, and such that $\mathbb{P}(\sigma^2 = 0) = 0$. Finally, we assume that v is independent of W where v is defined as $v = (g, h, \mu, \sigma)$.

Assumption D. We assume that $\mathbb{P}(\exists t \in [0, T] \text{ s.t. } u_t \neq 0) = 1$. We also assume that u is absolutely continuous on $[0, T]$, i.e., there exists a stochastic process $\theta : [0, T] \times \Omega \rightarrow \mathbb{R}$ with $u_t = \int_0^t \theta_s ds$, a.s.. Finally, we assume that $\mathbb{E}[\exp(\frac{1}{2} \int_0^T \theta_s^2 ds)] < \infty$.

By **Assumption D**, M satisfies Novikov's condition and thus is a positive martingale.

Lemma 22. *Under Assumption D, we have that M is a positive martingale. Thus, we can consider an equivalent probability measure \mathbb{Q} such that the Radon-Nikodym derivative is defined as $\frac{d\mathbb{Q}}{d\mathbb{P}} = M_T$. Finally, Y is a standard Wiener process under \mathbb{Q} .*

The elementary idea in this appendix is to condition by both W_T and v , i.e., to derive results of the form $\mathbb{P}(T_{b,c}^Y \leq T | W_T, v)$. The next proposition reexpresses $\mathbb{P}(T_{b,c}^Y \leq T | W_T, v)$ under \mathbb{Q} . We define \overline{W}_t as

$$\overline{W}_t = \int_0^t \theta_s dW_s. \quad (89)$$

Proposition 23. *Under Assumption D, we have*

$$\mathbb{P}(T_{b,c}^Y \leq T | W_T, v) = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} M_T^{-1} | W_T, v]. \quad (90)$$

This can be reexpressed as

$$\begin{aligned} \mathbb{P}(T_{b,c}^Y \leq T | W_T, v) &= \\ \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T, \overline{W}_T, v] | W_T, v]. \end{aligned} \quad (91)$$

We define the correlation under \mathbb{P} between W_T and $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ as ρ , i.e., $\rho = \text{Cor}_{\mathbb{P}}(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}})$.

Lemma 24. *Under Assumption D, we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . We can also show that $\rho = \frac{1}{T} \mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s ds}{\sqrt{\int_0^T \theta_s^2 ds}} \right]$ a.s.. Moreover, there exists a standard normal random variable \widetilde{W} under \mathbb{P} , which is independent of W_T , and such that \overline{W}_T when normalized can be reexpressed a.s. as*

$$\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}} = \rho \frac{W_T}{\sqrt{T}} + \sqrt{1 - \rho^2} \widetilde{W}. \quad (92)$$

This can be reexpressed a.s. as

$$\overline{W}_T = \alpha W_T + \tilde{\alpha} \widetilde{W}, \quad (93)$$

where $\alpha = \rho \sqrt{T^{-1} \int_0^T \theta_s^2 ds}$ a.s. and $\tilde{\alpha} = \sqrt{(1 - \rho^2) \int_0^T \theta_s^2 ds}$ a.s.. If we define $\tilde{\theta}_t = \frac{\theta_t - \alpha}{\tilde{\alpha}}$, we can reexpress \widetilde{W} a.s. as

$$\widetilde{W} = \int_0^T \tilde{\theta}_s dW_s. \quad (94)$$

Moreover, $\widetilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} . Finally, the conditional distribution of $\widetilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds$ given v , i.e., $\mathcal{D}(\widetilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds | v)$, is standard normal under \mathbb{Q} .

Our main result is the next theorem.

Theorem 25. *Under **Assumption D**, we have*

$$\begin{aligned} \mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T, v) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T, v]. \end{aligned} \quad (95)$$

We first calculate $\mathbb{Q}(\mathbb{T}_{b,c}^Y \leq T | W_T, v)$.

Lemma 26. *Under **Assumption D**, we have*

$$\mathbb{Q}(\mathbb{T}_{b,c}^Y \leq T | W_T, v) = \sum_{j=1}^{\infty} q_{b,c}^Y(j | Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} + \mathbf{1}_{\{Y_T \notin [c_T, b_T]\}}. \quad (96)$$

The next theorem gives a formula based on the strong theoretical assumption (97).

Theorem 27. *We assume that **Assumption D** and the following assumption*

$$\begin{aligned} &\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T, v] \\ &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} | W_T, v] \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha} \tilde{W}) | W_T, v] \end{aligned} \quad (97)$$

holds. Then, we have

$$\begin{aligned} \mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T, v) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \left(\sum_{j=1}^{\infty} q_{b,c}^Y(j | Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} \right. \\ &\quad \left. + \mathbf{1}_{\{Y_T \notin [c_T, b_T]\}} \right) \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned} \quad (98)$$

Finally, we get $P_b^Y(T)$ in the next corollary, by integrating $\mathbb{P}(\mathbb{T}_b^Y \leq T | W_T, v)$ with respect to the value of (W_T, v) . We define the arrival space and cdf of v as respectively Π_v and P_v . Moreover, we define y_u, y_b, y_θ , etc. following the above definitions when integrating with respect to $y \in \Pi_v$.

Corollary 28. *Under **Assumption D**, we have*

$$\begin{aligned} P_{b,c}^Y(T) &= 1 - \phi\left(\frac{b_T - u_T}{\sqrt{T}}\right) + \phi\left(\frac{c_T - u_T}{\sqrt{T}}\right) \\ &\quad + \int_{c_T - u_T}^{b_T - u_T} \int_{\Pi_v} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-y_\alpha x + \frac{1}{2} \int_0^T y_{\theta,s}^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_{b,c}^Y \leq T\}} \exp(-\tilde{\alpha} \tilde{W}) | W_T = x, v = y] dx dP_v(y). \end{aligned} \quad (100)$$

If we further assume (97), we have

$$\begin{aligned}
P_{b,c}^Y(T) &= 1 - \phi\left(\frac{b_T - u_T}{\sqrt{T}}\right) + \phi\left(\frac{c_T - u_T}{\sqrt{T}}\right) \\
&+ \int_{c_T - u_T}^{b_T - u_T} \int_{\Pi_v} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-y_\alpha x + \frac{1}{2} \int_0^T y_{\theta,s}^2 ds\right) \\
&\times \left(\sum_{j=1}^{\infty} y_{q,y_b,y_c}^{x+y_{u,T}}(j|x+y_{u,T}) \mathbf{1}_{\{x \in [y_{c,T-y_{u,T}}, y_{b,T-y_{u,T}}]\}} \right. \\
&\left. + \mathbf{1}_{\{x \notin [y_{c,T-y_{u,T}}, y_{b,T-y_{u,T}}]\}} \right) \\
&\times \exp\left(y_{\bar{\alpha}} \int_0^T y_{\bar{\theta},s} y_{\theta,s} ds\right) \mathcal{L}_N(y_{\bar{\alpha}}) dx dP_v(y). \tag{101}
\end{aligned}$$

Appendix B: Proofs in the two-sided stochastic boundary process case

In this section, we consider the proofs in the case when the two-sided boundary and the drift are stochastic processes and the variance is random.

The elementary idea in the proofs of this section is to condition by both W_T and v , i.e., to derive results of the form $\mathbb{P}(T_{b,c}^Y \leq T | W_T, v)$. The proof of Proposition 23 is based on Lemma 22.

Proof of Proposition 23. By definition of the conditional probability, Equation (90) can be rewritten formally as

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_b^Y \leq T\}} | W_T, v] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} M_T^{-1} | W_T, v]. \tag{102}$$

For any $\sigma(W_T, v)$ -measurable event E_T , we can use a change of probability in the expectation by Lemma 22, along with **Assumption D**, and we obtain that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T}] = \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} M_T^{-1} \mathbf{1}_{E_T}]. \tag{103}$$

We can deduce Equation (102) from Equation (103) by definition of the conditional expectation. By definition of the conditional probability, Equation (91) can be rewritten formally as

$$\begin{aligned}
&\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T, v] \\
&= \mathbb{E}_{\mathbb{Q}}[M_T^{-1} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} | W_T, \bar{W}_T, v] | W_T, v].
\end{aligned} \tag{104}$$

By definition of the conditional expectation, we can deduce what follows. If we can show that for any E_T , which is $\sigma(W_T, v)$ -measurable, that

$$\mathbb{E}_{\mathbb{P}}[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T}] \tag{105}$$

$$= \mathbb{E}_{\mathbb{P}} \left[\mathbb{E}_{\mathbb{Q}} \left[M_T^{-1} \mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mid W_T, \overline{W}_T, v \right] \mid W_T, v \right] \mathbf{1}_{E_T} \right],$$

then Equation (104) holds. Let E_T be a $\sigma(W_T, v)$ -measurable event. By Lemma 22 along with **Assumption D**, we obtain that

$$\mathbb{E}_{\mathbb{P}} \left[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} \right] = \mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} \right]. \quad (106)$$

Then, we have by the law of total expectation that

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} \right] &= \\ \mathbb{E}_{\mathbb{Q}} \left[\mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} \mid W_T, \overline{W}_T, v \right] \right]. \end{aligned} \quad (107)$$

Since $\mathbf{1}_{E_T}$ and M_T^{-1} are $\sigma(W_T, \overline{W}_T, v)$ -measurable random variables, we can pull them out of the conditional expectation and deduce that

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}} \left[\mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mathbf{1}_{E_T} M_T^{-1} \mid W_T, \overline{W}_T, v \right] \right] &= \\ = \mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{E_T} M_T^{-1} \mathbb{E}_{\mathbb{Q}} \left[\mathbf{1}_{\{T_{b,c}^Y \leq T\}} \mid W_T, \overline{W}_T, v \right] \right]. \end{aligned} \quad (108)$$

If we use Equations (106)-(107)-(108), we can deduce that Equation (105) holds. \square

In what follows, we give the proof of Lemma 24.

Proof of Lemma 24. By **Assumption D**, we can deduce that $0 < \int_0^T \theta_s^2 ds < \infty$ a.s.. Thus, we can normalize \overline{W}_T by $\sqrt{\int_0^T \theta_s^2 ds}$ a.s. and we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a mixed normal random variable a.s. by definition. Using the same arguments from the proof of Lemma 2.10, we have that its conditional mean under \mathbb{P} is a.s. equal to

$$\mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \mid v \right] = 0.$$

We also have that its conditional variance under \mathbb{P} is a.s. equal to

$$\text{Var}_{\mathbb{P}} \left(\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \mid v \right) = 1. \quad (109)$$

Since its conditional mean and conditional variance are nonrandom, we obtain that its mean under \mathbb{P} is equal to $\mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \right] = \mathbb{E}_{\mathbb{P}} \left[\mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s dW_s}{\sqrt{\int_0^T \theta_s^2 ds}} \mid v \right] \right] = 0$ by the the law of total expectation and Equation (109). Similarly, we obtain that its variance is equal to 1 by the law of total expectation and Equation (109).

Thus, we have that $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ is a standard normal random variable under \mathbb{P} . Since $(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}})$ is a centered normal random vector under \mathbb{P} , there exists a standard normal random variable \widetilde{W} under \mathbb{P} which is independent of W_T and such that Equation (92) holds. Then, we can calculate that the covariance between W_T and $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ under \mathbb{P} is equal to

$$\text{Cov}_{\mathbb{P}} \left(W_T, \frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}} \right) = \mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s ds}{\sqrt{\int_0^T \theta_s^2 ds}} \right]. \quad (110)$$

Now, we can calculate that the correlation between W_T and $\frac{\overline{W}_T}{\sqrt{\int_0^T \theta_s^2 ds}}$ under \mathbb{P} is equal to

$$\rho = \frac{1}{T} \mathbb{E}_{\mathbb{P}} \left[\frac{\int_0^T \theta_s ds}{\sqrt{\int_0^T \theta_s^2 ds}} \right].$$

Equation (93) can be deduced directly from Equation (92). Moreover, we can reexpress \widetilde{W} as

$$\widetilde{W} = \int_0^T \widetilde{\theta}_s dW_s.$$

Moreover, we can deduce that $\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} . This is due to its expression (94) and since by Lemma 22 along with **Assumption D**, Y is a Wiener process under \mathbb{Q} . Finally, $\mathcal{D}(\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds | v)$ is standard normal under \mathbb{Q} by Equation (94). \square

We provide now the proof of Theorem 25, which is based on Lemma 24.

Proof of Theorem 25. Using the same arguments from the proof of Theorem 2.11, we can reexpress M_T as

$$M_T = \exp(\alpha W_T - \frac{1}{2} \int_0^T \theta_s^2 ds) \exp(\widetilde{\alpha} \widetilde{W}).$$

Then, we have

$$\begin{aligned} \mathbb{P}(\mathbf{T}_{b,c}^Y \leq T | W_T, v) &= \exp\left(-\alpha W_T + \frac{1}{2} \int_0^T \theta_s^2 ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbf{T}_{b,c}^Y \leq T\}} \exp(-\widetilde{\alpha} \widetilde{W}) | W_T, v]. \end{aligned}$$

Thus, we have shown Equation (95). \square

We now give the proof of Lemma 26.

Proof of Lemma 26. By definition of the conditional probability, Equation (96) can be rewritten formally as

$$\mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\tau_{b,c}^Y \leq T\}} | W_T] = \sum_{j=1}^{\infty} q_{b,c}^Y(j|Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} + \mathbf{1}_{\{Y_T \notin [c_T, b_T]\}}. \quad (111)$$

By Lemma 22 along with **Assumption D**, Y is a Wiener process under \mathbb{Q} . Then, we have by Anderson (1960) (Theorem 4.2, pp. 178-179) that Equation (111) holds. \square

We provide now the proof of Theorem 27, which is based on Lemma 26.

Proof of Theorem 27. We have

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\tau_{b,c}^Y \leq T\}} \exp(-\tilde{\alpha}\tilde{W}) | W_T, v] &= \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\tau_{b,c}^Y \leq T\}} | W_T, v] \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | W_T, v] \\ &= \left(\sum_{j=1}^{\infty} q_{b,c}^Y(j|Y_T) \mathbf{1}_{\{Y_T \in [c_T, b_T]\}} \right. \\ &\quad \left. + \mathbf{1}_{\{Y_T \notin [c_T, b_T]\}} \right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | W_T, v], \end{aligned} \quad (112)$$

where we use Assumption (97) in the first equality, and Equation (96) from Lemma 26 along with **Assumption D** in the second equality. Finally, we have

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | W_T, v] &= \mathbb{E}_{\mathbb{Q}}[\exp(-\tilde{\alpha}\tilde{W}) | v] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}\left[\exp\left(-\tilde{\alpha}\left(\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds\right)\right) | v\right] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \\ &\quad \times \mathbb{E}_{\mathbb{Q}}\left[\exp\left(-\tilde{\alpha}\left(\tilde{W} + \int_0^T \tilde{\theta}_s \theta_s ds\right)\right)\right] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathbb{E}_{\mathbb{P}}[\exp(-\tilde{\alpha}N)] \\ &= \exp\left(\tilde{\alpha} \int_0^T \tilde{\theta}_s \theta_s ds\right) \mathcal{L}_N(\tilde{\alpha}). \end{aligned} \quad (113)$$

Here, we use the fact that \widetilde{W} is independent from W_T in the first equality, the fact that θ_t and $\widetilde{\theta}_t$ for any $t \in [0, T]$ are $\sigma(v)$ -measurable random variables in the second equality, the fact that $\mathcal{D}(\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds | v)$ is standard normal under \mathbb{Q} by Lemma 24 along with **Assumption D** in the third equality, the fact that $\widetilde{W} + \int_0^T \widetilde{\theta}_s \theta_s ds$ is a standard normal variable under \mathbb{Q} by Lemma 24 along with **Assumption D** in the fourth equality, and Equation (2.14) in the last equality. We can deduce Equation (98) from Equations (95), (112) and (113). \square

Finally, we get $P_{b,c}^Y(T)$ in the next theorem, by integrating $\mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T, v)$ with respect to the value of (W_T, v) .

Proof of Corollary 28. We can calculate that

$$\begin{aligned}
P_{b,c}^Y(T) &= \int_{-\infty}^{\infty} \int_{\Pi_v} \mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T = x, v = y) \\
&\quad \times \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx dP_v(y) \\
&= 1 - \phi\left(\frac{b_T - u_T}{\sqrt{T}}\right) + \phi\left(\frac{c_T - u_T}{\sqrt{T}}\right) \\
&\quad + \int_{c_T - u_T}^{b_T - u_T} \int_{\Pi_v} \mathbb{P}(\mathbb{T}_b^Y \leq T | W_T = x, v = y) \\
&\quad \times \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) dx dP_v(y) \\
&= 1 - \phi\left(\frac{b - u_T}{\sqrt{T}}\right) \\
&\quad + \int_{-\infty}^{b - u_T} \int_{\Pi_v} \frac{1}{\sqrt{2\pi T}} \exp\left(-\frac{x^2}{2T}\right) \exp\left(-y_\alpha x + \frac{1}{2} \int_0^T y_{\theta,s}^2 ds\right) \\
&\quad \times \mathbb{E}_{\mathbb{Q}}[\mathbf{1}_{\{\mathbb{T}_b^Y \leq T\}} \exp(-\widetilde{\alpha}\widetilde{W}) | W_T = x, v = y] dx dP_v(y).
\end{aligned}$$

Here, we use Equation (88), regular conditional probability and the fact that W_T and v are independent in the first equality, the fact that $\mathbb{P}(\mathbb{T}_{b,c}^Y \leq T | W_T = x) = 1$ for any $x \geq b_T - u_T$ and any $x \leq c_T - u_T$ in the second equality, and Equation (95) in the third equality. We have thus shown Equation (100). Equation (101) can be shown following the same first two equalities and using Equation (98) in the third equality. \square

References

AALLEN, O. O. and GJESSING, H. K. (2001). Understanding the shape of the hazard rate: A process point of view (with comments and a rejoinder by the authors). *Statistical Science* **16** 1–22.

- ABBRING, J. H. (2012). Mixed Hitting-Time Models. *Econometrica* **80** 783–819.
- ALILI, L. and KYPRIANOU, A. E. (2005). Some remarks on first passage of Lévy processes, the American put and pasting principles. *Annals of Applied Probability* **15** 2062–2080.
- ANDERSON, T. W. (1960). A modification of the sequential probability ratio test to reduce the sample size. *The Annals of Mathematical Statistics* **31** 165–197.
- BREIMAN, L. (1967). First exit times from a square root boundary. In *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability* **5** 9–17. University of California Press.
- BUTLER, R. W. and HUZURBAZAR, A. V. (1997). Stochastic network models for survival analysis. *Journal of the American Statistical Association* **92** 246–257.
- DANIELS, H. E. (1969). The minimum of a stationary Markov process superimposed on a U-shaped trend. *Journal of Applied Probability* **6** 399–408.
- DONEY, R. A. and KYPRIANOU, A. E. (2006). Overshoots and undershoots of Lévy processes. *Annals of Applied Probability* **16** 91–106.
- DOOB, J. L. (1949). Heuristic approach to the Kolmogorov-Smirnov theorems. *The Annals of Mathematical Statistics* **20** 393–403.
- DURBIN, J. (1971). Boundary-crossing probabilities for the Brownian motion and Poisson processes and techniques for computing the power of the Kolmogorov-Smirnov test. *Journal of Applied Probability* **8** 431–453.
- DURBIN, J. (1985). The first-passage density of a continuous Gaussian process to a general boundary. *Journal of Applied Probability* **22** 99–122.
- EATON, W. W. and WHITMORE, G. (1977). Length of stay as a stochastic process: A general approach and application to hospitalization for schizophrenia. *Journal of Mathematical Sociology* **5** 273–292.
- GIRSANOV, I. V. (1960). On transforming a certain class of stochastic processes by absolutely continuous substitution of measures. *Theory of Probability & Its Applications* **5** 285–301.
- GUT, A. (1974). On the moments and limit distributions of some first passage times. *The Annals of Probability* **2** 277–308.
- JACOD, J. and SHIRYAEV, A. (2003). *Limit theorems for stochastic processes*, 2nd ed. Berlin: Springer-Verlag.
- KARATZAS, I. and SHREVE, S. (2012). *Brownian motion and stochastic calculus* **113**. Springer Science & Business Media.
- KAZAMAKI, N. (1977). On a problem of Girsanov. *Tohoku Mathematical Journal*

- 29** 597–600.
- KOU, S. G. and WANG, H. (2003). First passage times of a jump diffusion process. *Advances in Applied probability* **35** 504–531.
- KYPRIANOU, A. E., PARDO, J. and RIVERO, V. (2010). Exact and asymptotic n-tuple laws at first and last passage. *Annals of Applied Probability* **20** 522–564.
- LAI, T. L. and SIEGMUND, D. (1977). A nonlinear renewal theory with applications to sequential analysis I. *The Annals of Statistics* **5** 946–954.
- LAI, T. L. and SIEGMUND, D. (1979). A nonlinear renewal theory with applications to sequential analysis II. *The Annals of Statistics* **7** 60–76.
- LAWLESS, J. F. (2011). *Statistical models and methods for lifetime data*. John Wiley & Sons.
- LEE, M.-L. T. and WHITMORE, G. A. (2006). Threshold regression for survival analysis: modeling event times by a stochastic process reaching a boundary. *Statistical Science* **21** 501–513.
- MALMQUIST, S. (1954). On certain confidence contours for distribution functions. *The Annals of Mathematical Statistics* **25** 523–533.
- MATTHEWS, D., FAREWELL, V. and PYKE, R. (1985). Asymptotic score-statistic processes and tests for constant hazard against a change-point alternative. *The Annals of Statistics* **13** 583–591.
- NOVIKOV, A. A. (1972). On an identity for stochastic integrals. *Teoriya Veroyatnostei i ee Primeneniya* **17** 761–765.
- NOVIKOV, A., FRISHLING, V. and KORDZAKHIA, N. (1999). Approximations of boundary crossing probabilities for a Brownian motion. *Journal of Applied Probability* **39** 1019–1030.
- POTIRON, Y. (2023). Explicit formula of boundary crossing probabilities for continuous local martingales to constant boundary. *arXiv preprint arXiv:2312.00287*.
- POTIRON, Y. and MYKLAND, P. A. (2017). Estimation of integrated quadratic covariation with endogenous sampling times. *Journal of Econometrics* **197** 20–41.
- RENAULT, E., VAN DER HEIJDEN, T. and WERKER, B. J. (2014). The dynamic mixed hitting-time model for multiple transaction prices and times. *Journal of Econometrics* **180** 233–250.
- REVUZ, D. and YOR, M. (2013). *Continuous martingales and Brownian motion* **293**. Springer Science & Business Media.
- ROBERTS, G. and SHORTLAND, C. (1997). Pricing barrier options with time-dependent coefficients. *Mathematical Finance* **7** 83–93.

- SALMINEN, P. (1988). On the first hitting time and the last exit time for a Brownian motion to/from a moving boundary. *Advances in Applied Probability* **20** 411–426.
- SIEGMUND, D. (1986). Boundary crossing probabilities and statistical applications. *The Annals of Statistics* **14** 361–404.
- STRASSEN, V. (1967). Almost sure behavior of sums of independent random variables and martingales. In *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability* **2** 315–343.
- WANG, L. and PÖTZELBERGER, K. (1997). Boundary crossing probability for Brownian motion and general boundaries. *Journal of Applied Probability* **34** 54–65.
- WOODROOFE, M. (1976). A renewal theorem for curved boundaries and moments of first passage times. *The Annals of Probability* **4** 67–80.
- WOODROOFE, M. (1977). Second order approximations for sequential point and interval estimation. *The Annals of Statistics* **7** 984–995.