Construction of Pseudo-hermitian matrices describing systems with balanced loss-gain

Pijush K. Ghosh*

Department of Physics, Siksha-Bhavana, Visva-Bharati University, Santiniketan, PIN 731 235, India. (Dated: January 3, 2024)

We present a general construction of pseudo-hermitian matrices in an arbitrary large, but finite dimensional vector space. The positive-definite metric which ensures reality of the entire spectra of a pseudo-hermitian operator, and is used for defining a modified inner-product in the associated vector space is also presented. The construction for an N dimensional vector space is based on the generators of SU(N) in the fundamental representation and the identity operator. We apply the results to construct a generic pseudo-hermitian lattice model of size N with balanced loss-gain. The system is amenable to periodic as well as open boundary conditions and by construction, admits entirely real spectra along with unitary time-evolution. The tight binding and Su-Schrieffer-Heeger(SSH) models with nearest neighbour(NN) and next-nearest neighbour(NNN) interaction with balanced loss-gain appear as limiting cases.

Keywords: Pseudo-hermiticity, SU(N) Group, Tight-Binding Model

I. INTRODUCTION

Pseudo-hermitian operators play an important role in the description of consistent quantum theory for a class of non-hermitian Hamiltonians which admit entirely real spectra[1]. The advent of \mathcal{PT} -symmetric quantum systems[2] is followed by the developments in pseudo-hermitian operators, and in a sense, these two concepts are complementary to each other. The basic idea of a pseudo-hermitian operator is to endow the vector space associated with a non-hermitian operator with a positive-definite metric such that it becomes hermitian with respect to a modified inner product. The unitarity and completeness of states are ensured with respect to the modified inner product. All non-hermitian operators are not amenable to such a construction —the positive-definite metric can be found only for the operators satisfying the condition of pseudo-hermiticity and admitting entirely real spectra. A prescription for constructing the metric in the Hilbert space for a given non-hermitian Hamiltonian is well known which requires a priory the complete knowledge of eigenspectra[1]. This requirement makes it difficult to find the exact metric analytically for systems with infinite dimensional Hilbert space, and various approximate methods have been used. The scenario for systems with finite-dimensional Hilbert space is better, and exact metric has been constructed for a few specific physical models[3–5].

The main focus of this article is the construction of generic pseudo-hermitian matrices and the associated metric for arbitrary large, but finite dimensional vector space. The motivation stems from the fact that finite size lattice models have representations in terms of matrices. The various types of tight binding models and spin chains with non-hermitian interaction may be brought under the purview of such a construction. Pseudo-hermitian matrices are also relevant for constructing a class of solvable models of vector non-linear Schrödinger equation, non-linear Dirac equation and

^{*} pijushkanti.ghosh@visva-bharati.ac.in

Oligomers with balanced loss-gain[6, 7]. Further, systems with infinite dimensional Hilbert space may be projected to specific sectors to obtain finite dimensional matrices —multi-level quantum systems with non-hermitian interaction may be amenable for such studies. A construction of pseudo-hermitian operators is also important in its own right from the viewpoint of mathematics.

The product of two hermitian matrices, with one of them being identified as the metric, satisfies the condition of pseudo-hermiticity [8–12]. A pseudo-anti-symmetric matrix D = MR was used in the Hamiltonian formulation of classical systems with balanced loss-gain, where the matrices M, R, D correspond to a generalized mass-term/metric, gauge-potential, loss-gain terms, respectively [8, 9]. The product-form of the pseudo-hermitian matrix has been used in the study of pseudo-hermitian random matrix theory [10, 11]. This product-form of pseudo-hermitian matrix is suitable for describing balanced loss-gain systems, since the trace of the non-hermitian part of the matrix is identically zero.

In this article, we use the generators of SU(N) and the $N \times N$ identity matrix as a basis to construct a hermitian matrix with N^2 independent real parameters. The product of two such hermitian matrices gives a non-hermitian matrix depending on $2N^2$ independent real parameters. The resultant matrix is pseudo-hermitian by construction, and we use the algebra of SU(N) generators to express it in terms of the basis formed by the SU(N) generators and the identity matrix. A sufficient condition for the metric to be positive definite is derived ensuring that the pseudo-hermitian matrix admits entirely real spectra. The coefficients of the SU(N) generators may be chosen judiciously to identify the pseudo-hermitian matrix with some physical models. We construct a generic pseudo-hermitian lattice model of size N with balanced loss-gain. The system is amenable to both periodic as well open boundary conditions, and admits entirely real spectra along with unitary time-evolution. We present a tight-binding model of size N with NN and NNN interaction. The lattice contains loss-gain terms at all sites and the strength of the hopping term is non-uniform. As a special case, we obtain a tight binding Hamiltonian of size N with defects at the two endpoints and uniform coupling. The positive-definite metric has been constructed. We also present a SSH-type tight-binding model with explicit construction of the positive definite metric.

The plan of this article is the following. We present some basic results pertaining to pseudo-hermitian operators [1] in the next section. The construction of the pseudo-hermitian operator is described in Sec. II. A. A sufficient condition for the generic pseudo-hermitian operator to admit entirely spectra is also presented in this section. Various known results related to the SU(N) group are presented in order to facilitate model building from the generic pseudo-hermitian operator. In Sec. III, the generic pseudo-hermitian matrix with NN and NNN interaction is presented. The balanced loss-gain terms are present at all sites and hopping strengths are non-uniform. A uniform tight-binding model and a SSH-type model are presented in Sec. III.A and III.B, respectively. Finally, results are summarized along with discussions in Sec. IV.

II. PSEUDO-HERMITIAN MATRIX

We will be using pseudo-hermitian matrices in order to analyze a class of systems with balanced loss and gain. We present the known basic results involving pseudo-hermitian matrices in order to facilitate further discussions. An operator \mathcal{O} is said to be pseudo-hermitian if it is related to its adjoint through a similarity transformation,

$$\mathcal{O}^{\dagger} = \eta \mathcal{O} \eta^{-1}. \tag{1}$$

The hermitian operator η is not unique for a given operator \mathcal{O} and may or may not be positive-definite. The operator \mathcal{O} admits entirely real spectra and norm of the associated eigen-vectors

are non-negative if a positive-definite $\eta := \eta_+$ exists. It should be noted that the operator \mathcal{O} is non-hermitian in a vector-space with the standard inner product. However, due to the pseudo-hermiticity property, it is hermitian in a vector space endowed with a metric η_+ such that the scalar-product is defined as $\langle \langle \cdot, \cdot \rangle \rangle_{\eta_+} \equiv \langle \cdot, \eta_+ \cdot \rangle$. The metric η_+ is hermitian with respect to the standard inner-product and its positivity excludes negative-norm states. The standard results for a hermitian operator \mathcal{O} is reproduced for $\eta_+ = I$, where I is the identity operator.

A non-hermitian operator that is related to a hermitian operator through a similarity transformation is known as quasi-hermitian operator[13]. The pseudo-hermiticity implies that \mathcal{O} is also crypto-hermitian, since the operator $\hat{\mathcal{O}} = \rho \mathcal{O} \rho^{-1}$ is hermitian with respect to the standard innerproduct in the associated vector-space. It should be noted that the operator $\rho := \sqrt{\eta_+}$ is well defined, since η_+ is positive-definite. The defining relation for a pseudo-unitary operator[14] V is $V^{\dagger}\eta_+V = \eta_+$ which can be realized in terms of a pseudo-hermitian operator \mathcal{O} as $V = e^{-i\mathcal{O}}$. The standard results for a unitary operator is reproduced for $\eta_+ = I$ for which \mathcal{O} is hermitian. The identity $\rho^{-1}e^{-i\hat{\mathcal{O}}} = e^{-i\mathcal{O}}\rho^{-1}$ is useful in various contexts which implies that the action of the operator ρ^{-1} on the unitary operator $\rho^{-i\hat{\mathcal{O}}}$ is to change it to a pseudo-unitary operator $\rho^{-i\hat{\mathcal{O}}}$. Similarly, ρ changes $\rho^{-i\hat{\mathcal{O}}}$ to $\rho^{-i\hat{\mathcal{O}}}$, i.e. $\rho e^{-i\hat{\mathcal{O}}} = e^{-i\hat{\mathcal{O}}}\rho$.

A. Construction of Pseudo-hermitian Operator

We consider three hermitian operators S, η and M in an N-dimensional vector space with the condition $[\eta, M] = 0$ and η is invertible. The pseudo-hermitian operator O is constructed in terms of S, η and M as follows,

$$\mathcal{O} = a_0 \mathcal{S} \eta + a_1 \mathcal{M}$$

$$= \underbrace{\frac{a_0}{2} \{\mathcal{S}, \eta\} + a_1 \mathcal{M}}_{\mathcal{O}_1} + \underbrace{\frac{a_0}{2} [\mathcal{S}, \eta]}_{\mathcal{O}_2}, \ a_0, a_1 \in \mathbb{R}$$

where \mathcal{O}_h is hermitian, while \mathcal{O}_{nh} is non-hermitian. The matrix \mathcal{O} is η -pseudo-hermitian, i.e. $\mathcal{O}^{\dagger} = \eta \mathcal{O} \eta^{-1}$, and admits entirely real spectra for a positive-definite η . The operator \mathcal{O}_{nh} is trace less by construction, i.e. $Tr(\mathcal{O}_{nh}) = 0$ and is suitable for describing systems with balanced loss-gain terms. The operator \mathcal{O} is defined in terms of $3N^2 + 2$ real parameters $-2N^2$ parameters corresponding to hermitian matrices \mathcal{S} and η , while a_0, a_1, \mathcal{M} account for the extra $N^2 + 2$ parameters. There are $N^2 + 2$ redundant parameters in this formalism and these parameters may be redefined appropriately so that there are $2N^2$ independent real parameters as required for a general pseudo-hermitian operator in N-dimensional vector space. The operator \mathcal{O} is pseudo-hermitian even for $a_1 = 0$. The sole purpose of adding the term is to have more flexibility in constructing physical systems by using \mathcal{O} with $a_1 \neq 0$. For example, it may be convenient to define the metric η in terms of total $p < N^2$ parameters to keep it simple. The operator \mathcal{O} with $a_1 = 0, a_0 = 1$ for such a choice is described in terms of $N^2 + p < 2N^2$ number of real parameters. The deficiency of $N^2 - p$ parameters can be compensated partly or fully by considering $a_1 \neq 0$. The operator \mathcal{M} modifies the hermitian part \mathcal{O}_h of \mathcal{O} and it may be chosen in various ways for a given η . A particular choice that is suitable for describing a host of physical systems in terms of pseudo-hermitian operators is as follows:

$$\mathcal{M} = \sum_{k=1}^{N-1} b_k \eta^k, \quad b_k \in \mathbb{R} \ \forall \ k. \tag{2}$$

The highest power of η in \mathcal{M} is N-1, since $\eta^k \ \forall \ k \geq N$ can be expressed in terms of a polynomial in η with the highest order N-1. This can be understood simply in terms of matrix representation of the operator η and Cayley-Hamilton theorem which states that every matrix satisfies its own characteristic polynomial. We choose $a_0 = 1, a_1 = 0$ in rest of this article such that \mathcal{O} is described in terms of $2N^2$ real parameters. One may add an anti-hermitian operator ξ to \mathcal{O} with the condition $\{\xi,\eta\}=0$ for which $\tilde{\mathcal{O}}=\mathcal{O}+\xi$ is still η -pseudo-hermitian. However, the anti-commutation relation between η and ξ implies that η is not positive-definite. Consequently, $\tilde{\mathcal{O}}$ can not admit entirely spectra and this possibility is abandoned.

The operators \mathcal{M} and η may be expressed as linear combinations of SU(N) generators T^a , $a=1,\ldots,N^2-1$ and the identity operator T^0 . We use the normalization convention $Tr(T^aT^b)=2\delta^{ab}$, where Tr(X) denotes the trace of X. The commutation and the ant-commutation relations among T^a are given by [15],

$$[T^a, T^b] = 2if^{abc}T^c, \quad \{T^a, T^b\} = \frac{4}{N}\delta^{ab} + 2d^{abc}T^c,$$
 (3)

where the completely anti-symmetric tensor $f^{abc} = \frac{1}{4i}Tr([T^a, T^b]T^c)$ denote the structure constants and the completely symmetric tensor $d^{abc} = \frac{1}{4}Tr(\{T^a, T^b\}T^c)$. The operators η and \mathcal{S} are expressed in terms of $2N^2$ real parameters $\alpha_a, \beta_a, a = 0, 1, \ldots, N^2 - 1$ as follows:

$$\eta = \sum_{a=0}^{N^{2}-1} \alpha_{a} T^{a} = \underbrace{\frac{1}{N} Tr(\eta)}_{\alpha_{0}} T^{0} + \sum_{a=1}^{n^{2}-1} \underbrace{\frac{1}{2} Tr(\eta T^{a})}_{\alpha_{a}; a \ge 1} T^{a},$$

$$S = \sum_{a=0}^{N^{2}-1} \beta_{a} T^{a} = \underbrace{\frac{1}{N} Tr(S)}_{\beta_{0}} T^{0} + \sum_{a=1}^{n^{2}-1} \underbrace{\frac{1}{2} Tr(ST^{a})}_{\beta_{a}; a \ge 1} T^{a}.$$
(4)

The operator $\eta - \alpha_0 T^0$ is trace less and admits positive as well as negative eigenvalues. The lowest negative eigenvalue of $\eta - \alpha_0 T^0$ is denoted as $-\lambda_{min}$, $\lambda_{min} > 0$ so that the lowest eigenvalue of η for $\alpha_0 > 0$ is $\alpha_0 - \lambda_{min}$. The condition $\alpha_0 > \lambda_{min}$ ensures that η is positive-definite. In absence of any closed form expression for λ_{min} , a lower bound on α_0 which ensures a positive-definite η may be specified as,

$$\alpha_0 > \alpha_0^{min} \equiv \sqrt{\frac{1}{2} Tr((\eta - \alpha_0 T^0)^2)} = \sqrt{\sum_{a=1}^{N^2 - 1} (\alpha_a)^2}$$
 (5)

The condition $\alpha_0 > \alpha_0^{min}$ is sufficient to ensure a positive-definite η and this is not a necessary condition. For all practical purposes, λ_{min} may have to be found analytically or numerically for a given η . The important message here is that it is always possible to choose α_0 such that η is positive-definite. It is obvious from the above discussions that η can not be positive definite for $\alpha_0 \leq 0$. The expressions for the operator \mathcal{O} may be evaluated by using Eq. (3) as,

$$\mathcal{O} = \sum_{a=0}^{N^2 - 1} \delta_a T^a$$

$$= \underbrace{\left(\alpha_{0}\beta_{0} + \frac{2}{N}\sum_{a=1}^{N^{2}-1}\alpha_{a}\beta_{a}\right)}_{\delta_{0}}T^{0} + \sum_{a=1}^{N^{2}-1}\underbrace{\left(\alpha_{0}\beta_{a} + \beta_{0}\alpha_{a} + \sum_{b,c=1}^{N^{2}-1}\left(d^{abc} - if^{abc}\right)\alpha_{b}\beta_{c}\right)}_{\delta_{a},a \geq 1}T^{a}. (6)$$

The term $d^{abc} - if^{abc}$ may be substituted as $\frac{1}{2}(Tr(T^aT^bT^c))^*$ by using the identity $Tr(T^aT^bT^c) = 2\left(d^{abc} + if^{abc}\right)$. The trace may be computed analytically for a given representation of SU(N) generators T^a by using any computer system algebra. The operator \mathcal{O} is non-hermitian with respect to the standard norm in the vector space due to the term containing the structure constants f^{abc} . However, with modified inner-product in a vector space endowed with the metric η , \mathcal{O} is a hermitian operator in terms $2N^2$ real parameters in an N dimensional vector space. The associated positive-definite metric operator η with N^2 real parameters is also constructed. It may be noted that previous attempts [3–5] in this regard in the context of \mathcal{PT} -symmetric and/or pseudo-hermitian systems in N dimensional Hilbert space produced non-hermitian Hamiltonian with total number of real parameters less than $2N^2$. To the best of our knowledge, this is the first time that a construction of the most general pseudo-hermitian operators with the maximum number of allowed real parameters is presented for a given N. The large number of parameters allow more flexibility in constructing physical systems in addition to providing a complete formalism. Further, the known properties of SU(N) generators may be useful for analyzing the system.

The explicit construction of pseudo-hermitian operator in N dimensional vector space involves choosing a representation of the SU(N) generators T^a . We introduce a complete set of N orthonormal basis vectors $|j\rangle$ satisfying the relations,

$$\langle j|k\rangle = \delta_{jk}, \quad \sum_{j=1}^{N} |j\rangle\langle j| = I.$$
 (7)

The N(N-1)/2 symmetric generators Λ_{ij}^S , N(N-1)/2 anti-symmetric generators Λ_{ij}^A and N-1 diagonal generators $\Lambda_{n^2-1}^D$ of SU(N) in the fundamental representation can be expressed in terms of these basis vectors as follows[16, 17]:

$$\Lambda_{k,j}^{S} = |k\rangle\langle j| + |j\rangle\langle k|, \quad \Lambda_{k,j}^{A} = -i\left(|k\rangle\langle j| - |j\rangle\langle k|\right), \quad 1 \le k < j \le N,
\Lambda_{n^{2}-1}^{D} = \sqrt{\frac{2}{n^{2}-n}} \left(\sum_{j=1}^{n-1} |j\rangle\langle j| - (n-1)|n\rangle\langle n|\right), \quad n = 2, 3, \dots, N$$
(8)

The SU(N) generators are chosen to be orthogonal $Tr(\Lambda_{ij}^a \Lambda_{kl}^b) = 2\delta_{ab}\delta_{ik}\delta_{jl}$. The following identifications may be made between T^a and Λ operators by expressing the index 'a' of the generator T^a in terms of j, k[18]:

$$\Lambda_{kj}^S = T^{j^2 + 2(k-j) - 1}, \ \Lambda_{kj}^A = T^{j^2 + 2(k-j)}, \ \Lambda_{n^2 - 1}^D = T^{n^2 - 1} \text{ for } n = 2, \dots N$$
(9)

The matrix representation of Λ operators reproduce the generalized Gell-Mann matrices [15]:

$$\left[\Lambda_{k,j}^{S} \right]_{lm} = \delta_{kl} \delta_{jm} + \delta_{km} \delta_{jl}, \quad \left[\Lambda_{k,j}^{A} \right]_{lm} = -i \left(\delta_{kl} \delta_{jm} - \delta_{km} \delta_{jl} \right), \quad 1 \le k < j \le N$$

$$\left[\Lambda_{n^{2}-1}^{D} \right]_{kl} = \sqrt{\frac{2}{n^{2}-n}} \begin{pmatrix} I_{n-1,n-1} & \mathbf{O}_{n-1,N-n+1} \\ \mathbf{O}_{N-n+1,n-1} & P_{N-n+1,N-n+1} \end{pmatrix}, \quad n = 2, \dots, N;$$

$$[P_{N-n+1,N-n+1}]_{k'l'} = -(n-1)\delta_{k'1}\delta_{l'1}, \ k',l'=1,2,\ldots,N-n+1, \ k,l=1,2,\ldots,N$$
 (10)

where I_m and \mathbf{O}_{mn} denote $m \times m$ identity and $m \times n$ null matrices, respectively. Recently, closed form expressions for f^{abc} , d^{abc} for any N have been given for the above representation of the SU(N) generators[18], which may be used to evaluate the term $d^{abc} - if^{abc}$ appearing in \mathcal{O} . We present our results in terms of Λ^s , Λ^a and $\Lambda^d_{n^2-1}$ and specific representations either in terms of ket-vectors or matrices may be chosen depending on the physical scenario.

III. SYSTEM WITH NN AND NNN INTERACTION

The general construction of a pseudo-hermitian operator is given in Sec. II. An analytical and/or numerical analysis of a system with $2N^2$ parameters becomes non-trivial for large N. There are plenty of physical systems which are described in terms a few independent parameters. For example, the matrix representation of \mathcal{O} may correspond to a tight-binding model of size N in which each atom interacts with rest of the atoms on the lattice and defects are present at all sites. The strength of the inter-atomic coupling decreases with the increasing separation between two atoms. If the interaction among the atoms is restricted to uniform NN and/or NNN coupling, there is a significant reduction in the total number of independent parameters, yet it corresponds to a physical model. Similar considerations for various types of lattice models of size N or coupled discrete and continuum nonlinear Schröinger equations are fruitful in analyzing a large class of pseudo-hermitian systems in terms of a few real parameters. In this section, we describe explicit construction of pseudo-hermitian operators which may correspond to physical systems with NN and NNN interactions.

A general approach for constructing a physical model from Eq. (6) would be to choose the $2N^2$ parameters appropriately such that it reduces to the desired form. A lattice model which can be expressed in terms of a finite-dimensional matrix is amenable for this approach. For example, a real symmetric tridiagonal matrix $\mathcal{O} - \delta_0 T^0$ with non-hermitian diagonal elements is suitable for describing a tight binding Hamiltonian with balanced loss-gain and NN interaction. This may be achieved by imposing the following conditions:

$$\alpha_0 \beta_a + \beta_0 \alpha_a + \sum_{b,c=1}^{N^2 - 1} \left(d^{abc} - i f^{abc} \right) \alpha_b \beta_c = \delta_a \neq 0, \ a = k^2 + 2k - 2 \text{ and } a = n^2 - 1$$

$$\alpha_0 \beta_a + \beta_0 \alpha_a + \sum_{b,c=1}^{N^2 - 1} \left(d^{abc} - i f^{abc} \right) \alpha_b \beta_c = 0, \ a \neq k^2 + 2k - 2 \text{ and } a \neq n^2 - 1, \tag{11}$$

where $k=1,2,\ldots,N-1$ and $n=2,3,\ldots,N$. It is apparent from Eqs. (7,8,9,10) that the first condition ensures non-vanishing elements in the superdiagonal, subdiagonal and main diagonal, while the second condition implies that the remaining elements are zero. All non-vanishing δ^a 's may be considered to be equal for describing a tight binding model with uniform coupling. Eq. (11) describes an underdetermined system —there are N^2-1 equations in terms of $2N^2$ variables. One may choose α_0,β_0,β_a as parameters and α_a 's as independent variables. This choice reduces Eq. (11) to a set of N^2-1 coupled linear equations in terms of N^2-1 variables. Analytical solutions may be pursued for a few smaller values of N. However, for large N, computer system algebra may be useful to find solutions. We present below a tight binding model by following a different approach.

We consider metric η and S as follows,

$$\eta = \gamma_0 I_N + \sum_{j=1}^N \gamma_j \Lambda_{j,j+1}^S, \quad \mathcal{S} = \xi_0 I_N + \sum_{j=1}^N \xi_j \Lambda_{j,j+1}^A, \quad \gamma_0, \xi_0, \gamma_i, \xi_i \in \mathbb{R} \ \forall i,$$
 (12)

where the identifications $\Lambda_{N,N+j}^S = \Lambda_{j,N}^S$, $\Lambda_{N,N+j}^a = -\Lambda_{j,N}^a$ have been made. We introduce the complex parameters Γ_j and the operators $\Lambda_{k,j}^+, \Lambda_{k,j}^-$ as follows:

$$\Gamma_{j} = \xi_{0}\gamma_{j} + i\xi_{j}\gamma_{0}, \ \Lambda_{k,j}^{+} = \frac{1}{2} \left(\Lambda_{k,j}^{S} + i\Lambda_{k,j}^{A} \right) = |k\rangle\langle j|, \ \Lambda_{k,j}^{-} = (\Lambda_{k,j}^{+})^{\dagger} = |j\rangle\langle k|, \ j > k.$$
 (13)

The pseudo-hermitian operator \mathcal{O} may be expressed as,

$$\mathcal{O} = \gamma_0 \xi_0 I_N + \sum_{j=1}^{N} \left(\Gamma_j^* \Lambda_{j,j+1}^+ + \Gamma_j \Lambda_{j,j+1}^- \right) - i \sum_{j=1}^{N} \left(\gamma_j \xi_{j+1} \Lambda_{j,j+2}^+ - \gamma_{j+1} \xi_j \Lambda_{j,j+2}^- \right) \\
- i \sum_{j=1}^{N} \left(\gamma_j \xi_j - \gamma_{j+1} \xi_{j+1} \right) \left[\sqrt{\frac{j}{2(j+1)}} \Lambda_{(j+1)^2 - 1}^D - \sum_{l=0}^{N-j-2} \frac{\Lambda_{(N-l)^2 - 1}^D}{[2(j+l+1)(j+l+2)]^{\frac{1}{2}}} \right] (14)$$

where the identifications $\Lambda^D_{(N+j)^2-1} = \Lambda^D_{(j+1)^2-1}$, $\gamma_{N+j} = \gamma_j$, $\xi_{N+j} = \xi_j$ have been made. The term with diagonal generators in $\mathcal O$ are non-hermitian and correspond to balanced loss-gain. The second term of $\mathcal O$ is hermitian and corresponds to NN interaction, while the third term describes non-hermitian NNN interaction for generic values of the parameters. A particular physical scenario is that only balanced loss-gain terms are non-hermitian, while the remaining terms of $\mathcal O$ are hermitian. This may be achieved by fixing the parameters as $\xi_i = -C\gamma_i, C \in \mathbb R, i=1,\ldots,N$ which make the terms describing NNN interaction hermitian. The third term is still non-hermitian for this choice of parameters and describe loss-gain terms. The complex parameters $\Gamma_i = (\xi_0 - iC\gamma_0)\gamma_i$ for this particular choice of ξ_i . The pseudo-hermitian operator $\mathcal O$ with balanced loss-gain is described in terms N+3 real parameters as follows:

$$\mathcal{O} = \gamma_{0}\xi_{0}I_{N} + \sum_{j=1}^{N} \left(\Gamma_{j}^{*}\Lambda_{j,j+1}^{+} + \Gamma_{j}\Lambda_{j,j+1}^{-}\right) + iC\sum_{j=1}^{N} \gamma_{j}\gamma_{j+1} \left(\Lambda_{j,j+2}^{+} - \Lambda_{j,j+2}^{-}\right)$$

$$+ iC\sum_{j=1}^{N} \left(\gamma_{j}^{2} - \gamma_{j+1}^{2}\right) \left[\sqrt{\frac{j}{2(j+1)}}\Lambda_{(j+1)^{2}-1}^{D} - \sum_{l=0}^{N-j-2} \frac{\Lambda_{(j+l+2)^{2}-1}^{D}}{\left[2(j+l+1)(j+l+2)\right]^{\frac{1}{2}}}\right]$$

$$= \gamma_{0}\xi_{0} + \sum_{j=1}^{N} \left[\left(\Gamma_{j}^{*}|j\rangle\langle j+1| + iC\gamma_{j}\gamma_{j+1}|j\rangle\langle j+2| + h.c.\right) - iC\left(\gamma_{j}^{2} - \gamma_{j+1}^{2}\right)|j+1\rangle\langle j+1|\right]$$
(15)

where $|N+j\rangle = |j\rangle$ and h.c. denotes hermitian conjugate. The following identity[17] relating the matrix representations of the diagonal generators $\Lambda_{n^2-1}^D$ and the projection operator $|j\rangle\langle j|$ has been used in the last step of Eq. (15):

$$|j+1\rangle\langle j+1| = \frac{1}{N}I_N - \sqrt{\frac{j}{2(j+1)}}\Lambda_{(j+1)^2-1}^D + \sum_{l=0}^{N-j-2} \frac{\Lambda_{(j+l+2)^2-1}^D}{[2(j+l+1)(j+l+2)]^{\frac{1}{2}}}, j=0,\dots,N-1$$

The matrix representation of \mathcal{O} is pentagonal sans the three corner elements $[\mathcal{O}_h]_{1,N-1}$, $[\mathcal{O}_h]_{1,N}$ and $[\mathcal{O}_h]_{2,N}$. The matrix is suitable for describing systems with nearest neighbour and next-nearest-neighbour interaction with periodic boundary conditions. It may be noted that system with open boundary condition may also be described by the same matrix by taking all three corner elements to be identically zero. In particular, the choice $\gamma_N = \delta_N = 0$ eliminates the corner elements and the system is amenable for open boundary condition.

A. Uniform coupling

We present an example of a Pseudo-hermitian Tight-Binding Model with uniform couplings for NN and NNN terms and two defects at the ends. We choose $\gamma_1 = \gamma_2 = \cdots = \gamma_{N-1} \equiv \gamma, \gamma_N = 0, \xi_0 = 0$ for which $\mathcal{H} := \frac{1}{C\gamma_0} (\mathcal{O} - \gamma_0 \delta_0 I_N)$ with \mathcal{O} given in Eq. (15) takes the following form:

$$\mathcal{H} = i\gamma_0 \sum_{j=1}^{N-1} \left[|j\rangle\langle j+1| - |j+1\rangle\langle j| \right] + i\gamma \sum_{j=1}^{N-2} \left[|j\rangle\langle j+2| - |j+2\rangle\langle j| \right] + i\gamma \left[|1\rangle\langle 1| - |N\rangle\langle N| \right]$$
(16)

The first two terms of \mathcal{H} are hermitian, while the last term is non-hermitian with respect to the standard norm in the Hilbert space. The Hamiltonian is pseudo-hermitian with respect to the metric η ,

$$\eta = \gamma_0 \sum_{i=1}^{N} |i\rangle\langle i| + \gamma \sum_{i=1}^{N-1} \left(|i\rangle\langle i+1| + |i+1\rangle\langle i| \right)$$
(17)

which has the form of a tridiagonal matrix. In particular, $\eta = \gamma_0 I_N + \gamma \sum_{i=1}^{N-1} \Lambda_{i,i+1}^S$ and its eigenvalues are,

$$e_k = \gamma_0 + 2\gamma \cos\left(\frac{k\pi}{N+1}\right), \ k = 1, \dots, N.$$
(18)

Note that $e_i - \gamma_0 = -(e_{N+1-i} - \gamma_0)$ implying that all the roots are positive definite for

$$\gamma_0 > 2|\gamma|\cos(\frac{\pi}{N+1})\tag{19}$$

which is the condition for a η -pseudo-hermitian \mathcal{H} with entirely real spectra. Denoting $\eta_D = \operatorname{diagonal}(e_1, \ldots, e_N)$ and X^T as the transpose of X, the orthogonal matrix O that diagonalizes η , i.e. $\eta_D = O\eta O^T$, has the expression $[O]_{ij} = \sqrt{\frac{2}{N+1}} \sin(\frac{ij\pi}{N+1})$. We define a matrix $\rho := \sqrt{\eta} = O^T EO$, where $E = \operatorname{diagonal}(\sqrt{e_1}, \ldots, \sqrt{e_N})$. The matrix ρ can be used to construct a hermitian Hamiltonian $h = \rho \mathcal{H} \rho^{-1}$ that is isospectral with \mathcal{H} . We do not investigate the eigenspectra and physical properties of \mathcal{H} in this article which will be taken elsewhere[19].

B. SSH Model

We present a pseudo-hermitian modified SSH model with defects at all sites. We choose N=2m, $\gamma_{2i-1,2i}=\delta_1, \gamma_{2i,2i+1}=\delta_2, \xi_0=0$ for which the Hamiltonian $\mathcal{H}:=(\mathcal{O}-\gamma_0\xi_0I_N)/C$ takes the form,

$$\mathcal{H} = i\gamma_0 \delta_1 \sum_{i=1}^m \left(|2i - 1\rangle\langle 2i| - |2i\rangle\langle 2i + 1| \right) + i\gamma_0 \delta_2 \sum_{i=1}^m \left(|2i\rangle\langle 2i + 1| - |2i + 1\rangle\langle 2i| \right)$$

+
$$i\delta_1\delta_2 \sum_{i=1}^{N} \left(|i\rangle\langle i+2| - |i\rangle\langle i+2| \right) + i\left(\delta_2^2 - \delta_1^2\right) \sum_{i=1}^{N} (-1)^{i+1} |i\rangle\langle i|$$
 (20)

The Hamiltonian contains NNN interaction in addition to the balanced loss-gain. The Hamiltonian is pseudo-hermitian with respect to the metric η ,

$$\eta = \gamma_0 \sum_{i=1}^{N} |i\rangle\langle i| + \delta_1 \sum_{i=1}^{m} \left(|2i - 1\rangle\langle 2i| + |2i\rangle\langle 2i - 1| \right) + \delta_2 \sum_{i=1}^{m} \left(|2i\rangle\langle 2i + 1| + |2i + 1\rangle\langle 2i| \right)$$
 (21)

The condition $\gamma_0 > \sqrt{m(\delta_1^2 + \delta_2^2)}$ is sufficient to ensure that the eigenvalues of η are positive-definite.

IV. CONCLUSIONS & DISCUSSIONS

We have presented a construction of a generic $N \times N$ pseudo-hermitian matrix in terms of $2N^2$ real parameters. This is the most general form of an $N \times N$ pseudo-hermitian matrix. The explicit realization of such a matrix utilizes the generators of SU(N) group, the identity matrix I_N and their properties. We have also presented the explicit form of the metric in the associated vector space which is required to define a modified norm. A sufficient condition has been stated ensuring a positive-definite metric, and hence entirely real spectra for the pseudo-hermitian matrix.

The generic pseudo-hermitian matrix is suitable for describing various types of lattice models appearing in physics. We have presented a generic pseudo-hermitian tight-binding model of size N with NN and NNN interaction, and defects at all sites leading to balanced loss-gain. The strength of the hopping term is non-uniform for this model. Further, we have specialized to uniform coupling and defects at the two extreme end-points. The metric for the tight-binding model and the condition for its positivity have also been presented. We have also presented a SSH-type model as a special case of the generic model.

There are a few new directions related to the present construction. For example, the eigen-spectra, transport properties, topological properties, ..., etc. of the tight binding models presented in this article have not been investigated in this article. It will be reported in a separate publication[19]. Further, it may be interesting to derive other physically interesting lattice models from the generic pseudo-hermitian matrix. It can be applied to the construction of exactly solvable vector non-linear Schrödinger equation, non-linear Dirac equation, Oligomer etc. by following the methods outlined in Ref. [6]. Some of these issues will be addressed in future publications.

A. Mostafazadeh, Pseudo-Hermitian Representation of Quantum Mechanics, Int. J. Geom. Meth. Mod. Phys. 7, 1191 (2010).

^[2] C. M. Bender, PT Symmetry in Quantum and Classical Physics (World Scientific, 2018).

^[3] Carl M. Bender, Peter N. Meisinger and Qinghai Wang, Finite-Dimensional PT-Symmetric Hamiltonians, J. Phys. A: Math. Gen. 36, 6791(2003).

 ^[4] Qing-hai Wang, Song-zhi Chia and Jie-hong Zhang, PT-symmetry as a generalization of Hermiticity, J. Phys. A: Math. Theor. 43, 295301(2010).

^[5] Qing-hai Wang, 2×2 PT-symmetric matrices and their applications, Phil. Trans. Royal Soc. A: Math. Phys. Eng. Sci. 371, 20120045 (2013).

^[6] P. K. Ghosh, Classical Hamiltonian Systems with balanced loss and gain, J. Phys.: Conf. Ser. 2038, 012012(2021).

- [7] P K Ghosh, Constructing Solvable Models of Vector Non-linear Schrödinger Equation with Balanced Loss and Gain via Non-unitary transformation, Phys. Lett. A402, 127361(2021).
- [8] P K Ghosh and D Sinha, Hamiltonian formulation of systems with balanced loss–gain and exactly solvable models, Annals of Physics 388, 276 (2018).
- [9] Pijush K. Ghosh, Taming Hamiltonian systems with balanced loss and gain via Lorentz interaction: General results and a case study with Landau Hamiltonian, J. Phys. A: Math. Theor. **52**, 415202 (2019).
- [10] Y. N. Joglekar, W. A. Karr, Level density and level-spacing distribution of random, self-adjoint, non-hermitian matrices, Phys. Rev. E 83, 031122 (2011).
- [11] J. Feinberg and R. Riser, Pseudo-Hermitian Random Matrix Models: General Formalism, Nucl. Phys. B 975, 115678 (2022).
- [12] Joshua Feinberg and Miloslav Znojil, Which metrics are consistent with a given pseudo-hermitian matrix?, J. Math. Phys. 63, 013505 (2022)
- [13] F G Scholtz, H B Geyer and F J W Hahne, Quasi-Hermitian operators in quantum mechanics and the variational principle, Annals of Physics 213, 74(1992).
- [14] A. Mostafazadeh, Pseudounitary operators and pseudounitary quantum dynamics, J. Math. Phys. 45, 932 (2004).
- [15] W. Greiner and B. Müller, Quantum Mechanics: Symmetries, Springer, Berlin 1989.
- [16] G. Kimura, The bloch vector for n-level systems, Phys. Lett. A 314, 339 (2003).
- [17] R. A. Bertlmann and P. Krammer, Bloch vectors for qudits, J. Phys. A: Math.Theor. 41, 235303 (2008).
- [18] D. Bossion and and P. Huo, General Formulas of the Structure Constants in the su(N) Lie Algebra, arXiv:2108.07219.
- [19] S. Ghosh and P. K. Ghosh, Under preparation