# Circulant graphs with valency up to 4 that admit perfect state transfer in Grover walks

Sho Kubota<sup>∗</sup> Kiyoto Yoshino†

#### Abstract

We completely characterize circulant graphs with valency up to 4 that admit perfect state transfer. Those of valency 3 do not admit it. On the other hand, circulant graphs with valency 4 admit perfect state transfer only in two infinite families: one discovered by Zhan and another new family, while no others do. The main tools for deriving these results are symmetry of graphs and eigenvalues. We describe necessary conditions for perfect state transfer to occur based on symmetry of graphs, which mathematically refers to automorphisms of graphs. As for eigenvalues, if perfect state transfer occurs, then certain eigenvalues of the corresponding isotropic random walks must be the halves of algebraic integers. Taking this into account, we utilize known results on the rings of integers of cyclotomic fields.

Keywords: perfect state transfer, Grover walk, circulant graph, automorphism MSC 2020 subject classifications: 05C50; 81Q99

## 1 Introduction

Quantum walks are a powerful tool in quantum computing and quantum information processing, with potential applications in quantum algorithms [\[1\]](#page-22-0), quantum simulations [\[21\]](#page-23-0), and quantum cryptography [\[22\]](#page-23-1). They have also been studied from a purely mathematical point of view as they show interesting properties not found in classical random walks, such as localization [\[14\]](#page-22-1) and periodicity [\[13\]](#page-22-2). One of the most attractive properties of quantum walks is perfect state transfer, which roughly speaking refers to transferring between specific quantum states with probability 1. Perfect state transfer has important applications in quantum communication and quantum information processing [\[8\]](#page-22-3).

The subject of this paper is perfect state transfer in Grover walks, which are typical discrete time quantum walks. Although perfect state transfer in continuous time quantum walks has been well studied, those in discrete time quantum walks have not been as well investigated as in continuous time quantum walks. For literature on perfect state transfer in continuous time quantum walks, we refer to Godsil's survey [\[9\]](#page-22-4). Representative studies of perfect state transfer in discrete time quantum walks are summarized in Section 1 of Zhan's paper [\[23\]](#page-23-2). Among these, the studies dealt with Grover walks are summarized in Table [1.](#page-1-0)

We focus on Zhan's work [\[23\]](#page-23-2). Zhan discovered an infinite family of circulant graphs with valency 4 that admits perfect state transfer. We extend her result and completely characterize circulant graphs up to valency 4 that admit perfect state transfer. In order to describe our results in detail, we will introduce the symbols and terms used in this paper

Let  $\Gamma = (V, E)$  be a graph with the vertex set V and the edge set E. Throughout this paper, we assume that graphs are simple and finite, i.e.,  $|V| < \infty$  and  $E \subset \{ \{x, y\} \subset V \mid x \neq y \}$ . Define  $\mathcal{A} = \mathcal{A}(\Gamma) = \{(x, y), (y, x) \mid \{x, y\} \in E\},\$  which is the set of *symmetric arcs* of  $\Gamma$ . The origin x and

<sup>∗</sup>Department of Mathematics Education, Aichi University of Education, 1 Hirosawa, Igaya-cho, Kariya, Aichi 448- 8542, Japan. skubota@auecc.aichi-edu.ac.jp

<sup>†</sup>Department of Computer Science, Faculty of Applied Information Science, Hiroshima Institute of Technology, Saeki Ward, Hiroshima, 731-5143, Japan. k.yoshino.n9@cc.it-hiroshima.ac.jp

Graphs	Ref.
Diamond chains	
$K_2 + K_n, K_2 + C_n$	'2
Circulant graphs with valency 4	23
$K_{m,m,\dots,m}$	

<span id="page-1-0"></span>Table 1: Previous works on perfect state transfer in Grover walks

terminus y of  $a = (x, y) \in A$  are denoted by  $o(a)$  and  $t(a)$ , respectively. We write the inverse arc of a as  $a^{-1}$ . We define several matrices on Grover walks. Note that Grover walks are also referred to as arc-reversal walks or arc-reversal Grover walks, and they are known as a special case of bipartite walks [\[7\]](#page-22-8). Let  $\Gamma = (V, E)$  be a graph, and set  $\mathcal{A} = \mathcal{A}(\Gamma)$ . Denote by  $I_V$  and  $I_A$  the identity matrices indexed by V and A, respectively. The *boundary matrix*  $d = d(\Gamma) \in \mathbb{C}^{V \times A}$  is defined by

$$
d_{x,a} = \frac{1}{\sqrt{\deg x}} \delta_{x,t(a)},
$$

where  $\delta_{a,b}$  is the Kronecker delta. The *shift matrix*  $R = R(\Gamma) \in \mathbb{C}^{\mathcal{A} \times \mathcal{A}}$  is defined by  $R_{a,b} = \delta_{a,b-1}$ . Note that  $R^2 = I_{\mathcal{A}}$ . Define the *time evolution matrix*  $U = U(\Gamma) \in \mathbb{C}^{\mathcal{A} \times \mathcal{A}}$  by  $U = R(2d^*d - I_{\mathcal{A}})$ . The entries of U are computed as

$$
U_{a,b} = \frac{2}{\deg_{\Gamma} t(b)} \delta_{o(a),t(b)} - \delta_{a,b^{-1}},
$$
\n(1.1)

whose substantial proof is in Lemma 5.1 of [\[19\]](#page-23-3). The quantum walk defined by U is called the *Grover walk* on Γ.

The matrix U can be visually captured by regarding it as a linear mapping. Let  $\Gamma = (V, E)$  be a graph. We write the entries of a vector  $\Psi \in \mathbb{C}^{\mathcal{A}}$  on the arcs of the graph as in Figure [1.](#page-1-1) If an entry of



<span id="page-1-1"></span>Figure 1: The entries of a vector written on the arcs of the graph

 $\Psi$  is 0, we omit the arc itself corresponding to the entry. If an entry of  $\Psi$  is 1, we may omit the value on the arc corresponding to the entry. See Figure [2.](#page-1-2) An arrow transmits with weight  $\frac{2}{\deg_{\Gamma} x}$  except

 $\overline{x}$ 2 de ∞ 1  $\stackrel{U}{\mapsto}$ 2  $\frac{2}{\deg x} - 1$ 2  $\deg x$ 2  $\deg x$ 2  $\overline{\deg x}$  $\overline{x}$ 

<span id="page-1-2"></span>Figure 2: The action of  $U$ 

for the opposite arrow, and reflects to the opposite arrow with weight  $\frac{2}{\deg_{\Gamma} x} - 1$ . See Section 6 in [\[19\]](#page-23-3)



<span id="page-2-0"></span>Figure 3: The action of  $U$  in the case where the degree of the terminus of an arc is  $2$ 

for details. In particular, an arrow is perfectly transmitted if the degree of the terminus of the arc is 2, as shown in Figure [3.](#page-2-0)

Let  $\Gamma$  be a graph, and let  $U = U(\Gamma)$  be the time evolution matrix. A vector  $\Phi \in \mathbb{C}^{\mathcal{A}}$  is a *state* if  $\|\Phi\|=1$ . For distinct states  $\Phi$  and  $\Psi$ , we say that *perfect state transfer* occurs from  $\Phi$  to  $\Psi$  at time  $\tau \in \mathbb{Z}_{\geq 1}$  if there exists  $\gamma \in \mathbb{C}$  with norm one such that  $U^{\tau} \Phi = \gamma \Psi$ . A state  $\Phi$  is said to be *vertex type* if there exists a vertex  $x \in V$  such that  $\Phi = d^*e_x$ , where  $e_x \in \mathbb{C}^V$  is the unit vector defined by  $(e_x)_z = \delta_{x,z}$ . A vertex type state represents a state that has values only on arcs reaching toward some vertex. Indeed, letting  $\epsilon_a \in \mathbb{C}^{\mathcal{A}}$  be the unit vector defined by  $(\epsilon_a)_z = \delta_{a,z}$ , we have

$$
d^*e_x = \frac{1}{\sqrt{\deg x}} \sum_{\substack{a \in \mathcal{A} \\ t(a) = x}} \epsilon_a.
$$

This can be visually represented as shown in the left of Figure [4.](#page-2-1) A vertex type state corresponds to the situation that a quantum walker is only on some vertex in the model where a walker moves on vertices. See also the right of Figure [4.](#page-2-1) We study perfect state transfer between vertex type states. See [\[17\]](#page-22-7) for motivation to restrict states under consideration to vertex type states.



<span id="page-2-1"></span>Figure 4: A vertex type state  $d^*e_x$  with  $\deg x = 3$ 

Definition 1.1. For a graph Γ, we say that Γ *admits perfect state transfer* if perfect state transfer occurs for  $U(\Gamma)$  between two distinct vertex type states.

In this paper, we completely characterize circulant graphs up to valency 4 that admit perfect state transfer. Our main tools are symmetry of graphs and eigenvalues. We will show a useful necessary condition for perfect state transfer to occur, which asserts that if it occurs between vertex type states  $d^*e_x$  and  $d^*e_y$ , then automorphisms that fix the vertex x must also fix the vertex y. See Corollary [3.4](#page-5-0) for details. This fact is well-known in the context of continuous-time quantum walks. As for eigenvalues, if perfect state transfer occurs, then certain eigenvalues of the corresponding isotropic random walks must be the halves of algebraic integers. Taking this into account, we can use certain facts on the rings of integers of cyclotomic fields to completely characterize circulant graphs up to valency 4 that admit perfect state transfer. The main theorems are the following two. For the symbol X representing a circulant graph, refer to Section [4.](#page-6-0)

<span id="page-3-0"></span>**Theorem 1.2.** *Let*  $X = X(\mathbb{Z}_{2l}, \{\pm a, l\})$  *be a connected* 3*-regular circulant graph, where*  $a \in \{1, \ldots, l-\}$ 1}*. Then, perfect state transfer does not occur on* X *between vertex type states.*

**Theorem 1.3.** Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  be a connected 4*-regular circulant graph.* Let  $x, y \in \mathbb{Z}_{2l}$ . *Then, the graph* X *admits perfect state transfer from*  $d^*e_x$  *to*  $d^*e_y$  *at minimum time*  $\tau \in \mathbb{Z}_{\geq 1}$  *if and only if*  $a + b = l$ ,  $y = x + l$  *and one of the following hold.* 

- (i) l *is odd and*  $\tau = 2l$ .
- (ii)  $l \equiv 2 \pmod{4}$  *and*  $\tau = l$ .

We supplement Theorem [1.3.](#page-3-0) In Theorem 6.1 of [\[23\]](#page-23-2), Zhan showed that perfect state transfer occurs in the case (i) of our Theorem [1.3.](#page-3-0) We newly show that perfect state transfer also occurs in the case (ii) and reveal that it does not occur in any other 4-regular circulant graphs.

## 2 Preliminaries

This section provides supplementary details that were not covered in the introduction. First, with respect to boundary matrices, we have the following by direct calculation.

<span id="page-3-1"></span>
$$
dd^* = I_V. \tag{2.1}
$$

The *discriminant*  $P = P(\Gamma) \in \mathbb{C}^{V \times V}$  is defined by  $P = dRd^*$ . The matrix P is isomorphic to the transition matrix of the isotropic random walk. Indeed, the matrix  $P$  is in some sense a normalized adjacency matrix of a graph. Note that the *adjacency matrix*  $A = A(\Gamma) \in \mathbb{C}^{V \times V}$  of a graph  $\Gamma = (V, E)$ is defined by

$$
A_{x,y} = \begin{cases} 1 & \text{if } \{x,y\} \in E, \\ 0 & \text{otherwise.} \end{cases}
$$

If a graph is regular, then the two matrices are closely related:

Lemma 2.1. *Let* Γ *be a* k*-regular graph, and let* A *and* P *be the adjacency matrix and the discriminant of*  $\Gamma$ *, respectively. Then we have*  $P = \frac{1}{k}A$ *.* 

This can be verified straightforwardly. A more general claim and its proof can be found in Theorem 3.1 of  $[18]$  and Proposition 3.3 of  $[18]$ . Since the discriminant P is a normal matrix, it has a spectral decomposition. Let  $\lambda_1, \ldots, \lambda_s$  be the distinct eigenvalues of P, and let  $W_i$  be the eigenspace associated to  $\lambda_i$ . Let  $d_i = \dim W_i$ , and let  $u^{(i,1)}, \ldots, u^{(i,d_i)}$  be an orthonormal basis of  $W_i$ . Then, the matrix

<span id="page-3-2"></span>
$$
E_i = \sum_{j=1}^{d_i} u^{(i,j)} (u^{(i,j)})^*
$$
\n(2.2)

is called the *projection matrix* associated to the eigenvalue  $\lambda_i$ , and it satisfies

$$
P = \sum_{i=1}^{s} \lambda_i E_i,
$$

$$
E_i E_j = \delta_{i,j} E_i,\tag{2.3}
$$

$$
E_i^* = E_i,\tag{2.4}
$$

$$
\sum_{i=1}^{s} E_i = I.
$$
\n(2.5)

See [\[20\]](#page-23-5) for the spectral decomposition of matrices. Using the above properties, the following can be obtained.

<span id="page-3-3"></span>**Lemma 2.2.** Let M be a normal matrix with the spectral decomposition  $M = \sum_{i=1}^{s} \lambda_i E_i$ , and let  $p(x)$ *be a polynomial. Then we have*  $p(M) = \sum_{i=1}^{s} p(\lambda_i) E_i$ .

## 3 Automorphisms of graphs

In this section, we discuss relationships between symmetry of graphs and perfect state transfer. For a wide readership, we begin by introducing automorphisms of graphs and their associated permutation matrices. Let  $\Gamma = (V, E)$  be a graph. A mapping  $g: V \to V$  is an *automorphism* of  $\Gamma$  if g is bijective, and  $\{x, y\} \in E$  if and only if  $\{g(x), g(y)\} \in E$ . We denote the set of all automorphisms of Γ by Aut(Γ), and let  $Aut(\Gamma)_x = \{g \in Aut(\Gamma) \mid g(x) = x\}$  for  $x \in V$ . We note that automorphisms preserve the degrees of vertices, i.e.,

<span id="page-4-4"></span>
$$
\deg g(x) = \deg x \tag{3.1}
$$

for any  $x \in V$  and  $g \in Aut(\Gamma)$ . See Lemma 1.3.1 in [\[11\]](#page-22-9) for its proof.

Let g be an automorphism of a graph Γ. Since g is bijective, it defines the permutation matrix  $M_g \in \mathbb{C}^{V \times V}$  given by  $(M_g)_{x,y} = \delta_{x,g(y)}$ . Basic properties of permutation matrices are

<span id="page-4-0"></span>
$$
M_g^{\top} = M_g^{-1} = M_{g^{-1}},\tag{3.2}
$$

and

<span id="page-4-2"></span>
$$
M_g e_x = e_{g(x)} \tag{3.3}
$$

for any  $x \in V$  and  $q \in Aut(\Gamma)$ .

The action of automorphisms of a graph is naturally extended to the set of symmetric arcs. Let g be an automorphism of a graph Γ, and let  $\mathcal{A} = \mathcal{A}(\Gamma)$  be the symmetric arc set. Define  $\tilde{g} : \mathcal{A} \to \mathcal{A}$  by  $\tilde{g}((x,y)) = (g(x), g(y))$ . Cleary,  $\tilde{g}$  is a bijection. Thus, the permutation matrix  $N_{\tilde{g}} \in \mathbb{C}^{\mathcal{A}\times\mathcal{A}}$  is defined by  $(N_{\tilde{g}})_{a,b} = \delta_{a,\tilde{g}(b)}$ , and it holds that

<span id="page-4-1"></span>
$$
N_{\tilde{g}}^{\top} = N_{\tilde{g}}^{-1} = N_{\tilde{g}^{-1}} \tag{3.4}
$$

as in Equality [\(3.2\)](#page-4-0). Moreover, we have

<span id="page-4-3"></span>
$$
N_{\tilde{g}}\epsilon_a = \epsilon_{\tilde{g}(a)}\tag{3.5}
$$

for  $a \in \mathcal{A}$  and  $q \in Aut(\Gamma)$ . The following claims are relationships between permutation matrices defined by automorphisms and matrices associated with Grover walks, which plays an important role in this study.

<span id="page-4-5"></span>**Lemma 3.1.** Let g be an automorphism of a graph  $\Gamma$ . Then we have  $d^*M_g = N_{\tilde{g}}d^*$ .

**Proof.** We want to show that  $N_{\tilde{g}}^{-1}d^*M_g = d^*$ . For  $a \in \mathcal{A}(\Gamma)$  and  $x \in V(\Gamma)$ , we have

$$
(N_{\tilde{g}}^{-1}d^*M_g)_{a,x} = \epsilon_a^{-\top} (N_{\tilde{g}}^{-1}d^*M_g)e_x
$$
  
\n
$$
= (N_{\tilde{g}}\epsilon_a)^{\top} d^*(M_ge_x)
$$
 (by (3.4))  
\n
$$
= \epsilon_{\tilde{g}(a)}^{-\top} d^*e_{g(x)}
$$
 (by (3.3) and (3.5))  
\n
$$
= (d^*)_{\tilde{g}(a),g(x)}
$$
  
\n
$$
= \frac{1}{\sqrt{\deg g(x)}} \delta_{g(x),t(\tilde{g}(a))}
$$
  
\n
$$
= \frac{1}{\sqrt{\deg x}} \delta_{g(x),g(t(a))}
$$
 (by (3.1))  
\n
$$
= \frac{1}{\sqrt{\deg x}} \delta_{x,t(a)}
$$
  
\n
$$
= d_{x,a}
$$
  
\n
$$
= (d^*)_{a,x}.
$$

Thus, we have  $N_{\tilde{g}}^{-1}d^*M_g = d^*$ , i.e.,  $d^*M_g = N_{\tilde{g}}d^*$ .

口

<span id="page-5-1"></span>**Lemma 3.2.** *Let* g *be an automorphism of a graph* Γ*. Then we have*  $UN_{\tilde{g}} = N_{\tilde{g}}U$ *.* 

**Proof.** Let  $a, b \in \mathcal{A}(\Gamma)$ . In a similar way as in the proof of Lemma [3.1,](#page-4-5) we have

$$
(N_{\tilde{g}}^{-1}RN_{\tilde{g}})_{a,b} = R_{\tilde{g}(a),\tilde{g}(b)} = \delta_{\tilde{g}(a),\tilde{g}(b)^{-1}} = \delta_{a,b^{-1}} = R_{a,b}.
$$

This means  $N_{\tilde{g}}R = RN_{\tilde{g}}$ . Since  $U = R(2d^*d - I_A)$ , we have

$$
N_{\tilde{g}}^* UN_{\tilde{g}} = RN_{\tilde{g}}^* (2d^*d - I_A) N_{\tilde{g}}
$$
  
=  $R (2d^* M_g^* M_g d - I_A)$  (by Lemma 3.1)  
= U.

Thus, we have  $UN_{\tilde{g}} = N_{\tilde{g}}U$ .

These lemmas imply that automorphisms preserve the occurrence of perfect state transfer between vertex type states.

<span id="page-5-3"></span>**Theorem 3.3.** Let g be an automorphism of a graph  $\Gamma = (V, E)$ , and let  $x, y \in V$ . If

<span id="page-5-2"></span>
$$
U^{\tau}d^*e_x = \gamma d^*e_y \tag{3.6}
$$

*for a positive integer*  $\tau$  *and a complex number*  $\gamma$ *, then we have*  $U^{\tau}d^*e_{g(x)} = \gamma d^*e_{g(y)}$ *. In particular, if perfect state transfer occurs from*  $d^*e_x$  *to*  $d^*e_y$  *on*  $\Gamma$  *at time*  $\tau$ *, then so does from*  $d^*e_{g(x)}$  *to*  $d^*e_{g(y)}$ *.* 

Proof. Indeed,

$$
U^{\tau}d^*e_{g(x)} = U^{\tau}d^*M_ge_x \qquad \qquad \text{(by (3.3))}
$$
  
\n
$$
= U^{\tau}N_{\tilde{g}}d^*e_x \qquad \qquad \text{(by Lemma 3.1)}
$$
  
\n
$$
= N_{\tilde{g}}U^{\tau}d^*e_x \qquad \qquad \text{(by Lemma 3.2)}
$$
  
\n
$$
= N_{\tilde{g}}\gamma d^*e_y \qquad \qquad \text{(by (3.6))}
$$
  
\n
$$
= \gamma d^*M_ge_y \qquad \qquad \text{(by Lemma 3.1)}
$$
  
\n
$$
= \gamma d^*e_{g(y)} \qquad \qquad \text{(by Lemma 3.1)}
$$

 $\Box$ 

 $\Box$ 

As a consequence of the above theorem, a necessary condition for perfect state transfer to occur between vertex type states  $d^*e_x$  and  $d^*e_y$  can be formulated in terms of automorphism groups. In short, an automorphism that fixes the vertex  $x$  must also fix the vertex  $y$ .

<span id="page-5-0"></span>Corollary 3.4. Let g be an automorphism of a graph  $\Gamma = (V, E)$ , and let  $x, y \in V$ . If perfect state *transfer occurs from*  $d^*e_x$  *to*  $d^*e_y$  *on*  $\Gamma$ *, then we have*  $Aut(\Gamma)_x = Aut(\Gamma)_y$ *.* 

**Proof.** Let  $g \in Aut(\Gamma)_x$ , and we want to show that  $g(y) = y$ . Since  $U^{\tau}d^*e_x = \gamma d^*e_y$  for a positive integer  $\tau$  and a complex number  $\gamma$ , we have

$$
d^*e_y = \gamma^{-1}U^{\tau}d^*e_x
$$
  
=  $\gamma^{-1}U^{\tau}d^*e_{g(x)}$  (by  $g \in \text{Aut}(\Gamma)_x$ )  
=  $\gamma^{-1}\gamma d^*e_{g(y)}$  (by Theorem 3.3)  
=  $d^*e_{g(y)}$ .

Multiply both sides by the matrix d. Then by Equality [\(2.1\)](#page-3-1), we have  $g(y) = y$ , and hence  $g \in Aut(\Gamma)_y$ .<br>The reverse inclusion can be verified in a similar way The reverse inclusion can be verified in a similar way.

In fact, the same statement holds true for continuous-time quantum walks as well. See Lemma 4.1 in [\[10\]](#page-22-10) for details.

Using Corollary [3.4,](#page-5-0) we demonstrate in Example [3.5](#page-6-1) how to verify that perfect state transfer does not occur between a specific pair of vertices. As this example shows, the corollary is useful if we know part of the automorphism group.

<span id="page-6-1"></span>**Example 3.5.** Let  $K_3$  be the complete graph of order 3 with vertex set  $[3] = \{0, 1, 2\}$  in Figure [5.](#page-6-2) Let g be the automorphism of  $K_3$  defined by  $g(0) = 0$ ,  $g(1) = 2$  and  $g(2) = 1$ . Then,  $g \in Aut(K_3)_0$ 



<span id="page-6-2"></span>Figure 5: The complete graph of order 3

and  $g \notin \text{Aut}(K_3)_1$ . Hence, by Corollary [3.4,](#page-5-0) we see that perfect state transfer does not occur from 0 to 1.

We will use Corollary [3.4](#page-5-0) to provide a necessary condition for perfect state transfer to occur. This assertion can also be interpreted as saying that the Grover walk captures symmetry of graphs through perfect state transfer.

## <span id="page-6-0"></span>4 Circulant graphs

One of the main objects in this paper is circulant graphs, which are Cayley graphs over the additive group  $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$  of integers modulo n. Here, we may assume that n is a positive integer. If no confusion arises, we identify an integer  $x \in \mathbb{Z}$  with the equivalent class  $x + n\mathbb{Z} \in \mathbb{Z}_n$ .

In this section, we present two useful lemmas for circulant graphs. Combining Corollary [3.4](#page-5-0) with some automorphisms of circulant graphs, we present Lemma [4.2](#page-7-0) to identify candidates for pairs of vertices in circulant graphs between which perfect state transfer can occur. After that, we introduce the concept of eigenvalue support to investigate whether perfect state transfer occurs. We then prove Lemma [4.5,](#page-7-1) which asserts that the eigenvalue support coincides with the set of eigenvalues for circulant graphs.

We recall the definition and basic properties of circulant graphs. Let S be a subset of  $\mathbb{Z}_n \setminus \{0\}$  such that

<span id="page-6-3"></span>
$$
S = -S,\tag{4.1}
$$

i.e.,  $s \in S$  if and only if  $-s \in S$ . The *circulant graph*  $X(\mathbb{Z}_n, S)$  is the graph with the vertex set  $\mathbb{Z}_n$  and the edge set  $\{x, y\} | y - x \in S\}$ . In circulant graphs, "rotations" and "inversions" are automorphisms as claimed in the following lemma.

<span id="page-6-4"></span>**Lemma 4.1** (pp. 8–9 [\[11\]](#page-22-9)). Let  $X = X(\mathbb{Z}_n, S)$ . Then we have the following.

- (i) *For*  $z \in \mathbb{Z}_n$ *, the mapping*  $\rho_z : \mathbb{Z}_n \to \mathbb{Z}_n$  *defined by*  $\rho_z(x) = x + z$  *is an automorphism of* X.
- (ii) *The mapping*  $r : \mathbb{Z}_n \to \mathbb{Z}_n$  *defined by*  $r(x) = -x$  *is an automorphism of* X.

In considering perfect state transfer on circulant graphs between vertex type states, from Theo-rem [3.3](#page-5-3) we may assume that the initial state is  $d^*e_0$  without loss of generality. Moreover, we obtain simple necessary conditions for perfect state transfer to occur on  $X(\mathbb{Z}_n, S)$ .

<span id="page-7-0"></span>**Lemma 4.2.** Let  $X = X(\mathbb{Z}_n, S)$ , and let x, y be distinct vertices of X. If perfect state transfer occurs *on X* from  $d^*e_x$  to  $d^*e_y$ , then *n* is even and  $y = x + \frac{n}{2}$ .

**Proof.** Suppose that perfect state transfer occurs from  $d^*e_x$  to  $d^*e_y$ . From  $\rho_{-x} \in \text{Aut}(X)$  and Theorem [3.3,](#page-5-3) perfect state transfer occurs from  $d^*e_{\rho_{-x}(x)} = d^*e_0$  to  $d^*e_{\rho_{-x}(y)} = d^*e_{y-x}$ . Also, the automorphism r fixes the vertex 0, so Corollary [3.4](#page-5-0) implies that r also fixes the vertex  $y - x$ , i.e.,  $y - x = r(y - x) = -(y - x)$ . We have

$$
2(y - x) = 0.\t\t(4.2)
$$

We note that the operations and equal signs here are on  $\mathbb{Z}_n$ . Since  $x \neq y$  in  $\mathbb{Z}_n$ , we see that n is even.<br>Furthermore, we have either  $y - x = 0$  or  $y - x = \frac{n}{L}$ . As  $x \neq y$ , we conclude  $y = x + \frac{n}{L}$ . Furthermore, we have either  $y - x = 0$  or  $y - x = \frac{n}{2}$ . As  $x \neq y$ , we conclude  $y = x + \frac{n}{2}$ .

Next, we describe the eigenvalues and eigenvectors of circulant graphs. A general discussion on eigenvalues of Cayley graphs on abelian groups is given in Section 1.4.9 in [\[5\]](#page-22-11). In this paper, we avoid representation theoretic discussion and explicitly provide eigenvectors of circulant graphs. Let  $[n] = \{0, 1, \ldots, n-1\}$  for a positive integer n. Let  $\zeta_n = e^{\frac{2\pi}{n}i}$ . For  $j \in [n]$ , define the vector  $u_j \in \mathbb{C}^{\mathbb{Z}_n}$ by

<span id="page-7-2"></span>
$$
(u_j)_k = \zeta_n^{jk} \tag{4.3}
$$

<span id="page-7-3"></span>for  $k \in [n]$ .

**Lemma 4.3** (Section 1.4.9 in [\[5\]](#page-22-11)). Let  $X = X(\mathbb{Z}_n, S)$ . Then  $u_j$  is an eigenvector of  $A(X)$  of the *eigenvalue*  $\sum_{s \in S} \zeta_n^{js}$ . Furthermore, the eigenvectors  $u_0, u_1, \ldots, u_{n-1}$  are pairwise orthogonal.

**Definition 4.4.** Let  $M \in \mathbb{C}^{n \times n}$  be a normal matrix with the spectral decomposition  $M = \sum_{i=1}^{s} \lambda_i E_i$ , where  $\lambda_1, \ldots, \lambda_s$  are the distinct eigenvalues of M, and  $E_1, \ldots, E_s$  are the orthogonal projections associated to these eigenvalues, respectively. We denote by  $\sigma(M)$  the set of distinct eigenvalues of M. For a vector  $x \in \mathbb{C}^n$ , we define  $\Theta_M(x) = \{\lambda_i \in \sigma(M) \mid E_i x \neq 0\}$ . This is called the *eigenvalue support* of the vector  $x$  with respect to  $M$ .

Note that  $\sigma(M)$  is simply a set, so the multiplicities of the eigenvalues are ignored.

<span id="page-7-1"></span>**Lemma 4.5.** Let  $X = X(\mathbb{Z}_n, S)$ , and let A be the adjacency matrix of X. Then we have  $\Theta_A(e_0) =$  $\sigma(A)$ .

**Proof.** It is enough to show that  $\Theta_A(e_0) \supset \sigma(A)$ . Let  $\lambda \in \sigma(A)$ . Define

$$
I_{\lambda} = \left\{ j \in [n] \, \middle| \, \sum_{s \in S} \zeta_n^{js} = \lambda \right\}.
$$

Let  $E_{\lambda}$  be the projection matrix associated to the eigenvalue  $\lambda$ . Then by [\(4.3\)](#page-7-2) and [\(2.2\)](#page-3-2), we have

$$
E_{\lambda} = \frac{1}{n} \sum_{j \in I_{\lambda}} u_j u_j^*.
$$

Hence

$$
E_{\lambda}e_0 = \frac{1}{n} \left( \sum_{j \in I_{\lambda}} u_j u_j^* \right) e_0 = \frac{1}{n} \sum_{j \in I_{\lambda}} u_j.
$$

By Lemma [4.3,](#page-7-3)  $u_i$ 's are pairwise orthogonal, so they are linearly independent. This implies that  $E_{\lambda}e_0 \neq 0$ . Therefore, we have  $\lambda \in \Theta_A(e_0)$ , i.e.,  $\sigma(A) = \Theta_A(e_0)$ .  $\Box$ 

## 5 Cycle graphs

As a simple application of the previous sections, we characterize circulant graphs with valency 2 that admit perfect state transfer. Connected 2-regular graphs are cycle graphs. Thus, it is sufficient to consider only  $C_n = X(\mathbb{Z}_n, \{\pm 1\})$  without loss of generality. By Lemma [4.2,](#page-7-0) if the cycle graph  $C_n$ admits perfect state transfer, then n is even. Moreover, let  $n = 2l$ , and if perfect state transfer occurs from  $d^*e_x$ , then it must occur from  $d^*e_x$  to  $d^*e_{x+l}$ . Conversely, it can be easily verified that  $C_{2l}$  indeed admits perfect state transfer by visually capturing the action of the time evolution matrix  $U$ . Readers should recall the action of U described in Figure [3.](#page-2-0) For simplicity, we confirm this with  $C_6$  as shown in Figure [6,](#page-8-0) but readers will be convinced that the same is true for general  $C_{2l}$ .



<span id="page-8-0"></span>Figure 6: Visual proof that  $C_6$  admits perfect state transfer at time 3

In summarizing the above discussion, we have the following.

**Theorem 5.1.** Let  $C_n$  be the cycle graph on n vertices.

- (i) If n is odd, then  $C_n$  does not admit perfect state transfer between distinct vertex type states.
- (ii) If n is even, say  $n = 2l$ , then  $C_n$  admits perfect state transfer from  $d^*e_x$  to  $d^*e_{x+l}$  at minimum *time* l*.*

### 6 Characterization of perfect state transfer

For circulant graphs with valency 3 or greater, the superposition of quantum walkers, which corresponds to the arrows in our model, is too complex to consider in the same way as in the previous section. Therefore, it is necessary to switch strategies to use of eigenvalues. The main goal of this section is to characterize perfect state transfer by eigenvalues of graphs. We slightly generalize Zhan's result and connect it to Chebyshev polynomials.

The *Chebyshev polynomial of the first kind*, denoted by  $T_n(x)$ , is the polynomial defined by  $T_0(x)$  = 1,  $T_1(x) = x$  and  $T_n(x) = 2xT_{n-1}(x) - T_{n-2}(x)$  for  $n \ge 2$ . It is well-known that  $T_n(\cos \theta) = \cos(n\theta)$ , and hence we have the following.

<span id="page-8-1"></span>**Lemma 6.1.** *Let*  $\lambda$  ∈ [−1, 1]*.* 

- (i)  $|T_\tau(\lambda)| \leq 1$ .
- (ii)  $T_{\tau}(\lambda) = 1$  *if and only if*  $\lambda = \cos \frac{j}{\tau} \pi$  for some even j.
- <span id="page-8-2"></span>(iii)  $T_{\tau}(\lambda) = -1$  *if and only if*  $\lambda = \cos \frac{j}{\tau} \pi$  *for some odd j.*

**Lemma 6.2.** Let P be the discriminant of a graph  $\Gamma$ , and  $\tau$  be a non-negative integer. Then, the *spectral radius of* P *is at most* 1*, and that of*  $T_{\tau}(P)$  *is at most* 1*. In particular,*  $|T_{\tau}(\lambda)| \leq 1$  *for every*  $\lambda \in \sigma(P)$ .

**Proof.** It follows from [\[18,](#page-23-4) Proposition 3.3] with  $\eta = 0$  that the spectral radius of P is at most 1. This together with Lemma [6.1](#page-8-1) implies that the spectral radius of  $T_{\tau}(P)$  is at most 1.  $\Box$ 

Furthermore, Chebyshev polynomials connect Grover walks and random walks. Kubota and Segawa presented the following equality and obtained a necessary condition on eigenvalues for perfect state transfer to occur [\[17\]](#page-22-7).

<span id="page-9-0"></span>Lemma 6.3 (Lemma 3.1 in [\[17\]](#page-22-7)). *Let* Γ *be a graph with the time evolution matrix* U *and the discriminant* P. Then we have  $dU^{\tau}d^* = T_{\tau}(P)$  for any non-negative integer  $\tau$ .

Interestingly, an equality analogous to Lemma [6.3](#page-9-0) has recently been presented also in another quantum walk model, which is called vertex-face walks. For more details see Section 4 of the paper written by Guo and Schmeits [\[12\]](#page-22-12).

Zhan has characterized in Theorem 5.3 of [\[23\]](#page-23-2) the occurrence of perfect state transfer in terms of eigenvalues and eigenprojections. We slightly extend Zhan's theorem by using Lemma [6.3.](#page-9-0) Our claim does not require that graphs are regular.

<span id="page-9-6"></span>**Lemma 6.4** (Section 4 in [\[6\]](#page-22-13)). Let  $\Gamma$  be a graph, and let  $U = U(\Gamma)$  be the time evolution matrix. *Perfect state transfer occurs from a state*  $\Phi$  *to a state*  $\Psi$  *at time*  $\tau$  *if and only if*  $|\langle U^{\tau} \Phi, \Psi \rangle| = 1$ *.* 

<span id="page-9-7"></span>**Theorem 6.5.** Let  $\Gamma = (V, E)$  be a graph. Let  $x, y \in V$  and  $\tau \in \mathbb{Z}_{\geq 1}$ . For each eigenvalue  $\lambda$  of  $P = P(\Gamma)$ , let  $E_{\lambda}$  be the eigenprojection of P associated to  $\lambda$ . The following are equivalent.

- <span id="page-9-3"></span><span id="page-9-1"></span>(*A*) *Perfect state transfer occurs on*  $\Gamma$  *from*  $d^*e_x$  *to*  $d^*e_y$  *at time*  $\tau$ *.*
- <span id="page-9-2"></span>(*B*) *There exists*  $\gamma \in \{\pm 1\}$  *such that*  $T_{\tau}(P)e_x = \gamma e_y$ .
- <span id="page-9-8"></span><span id="page-9-4"></span>(*C*) *There exists*  $\gamma \in \{\pm 1\}$  *such that all of the following three conditions are satisfied:* 
	- (*a*) *For any*  $\lambda \in \sigma(P)$ ,  $E_{\lambda}e_x = \pm E_{\lambda}e_y$ .
	- (*b*) *For any*  $\lambda \in \Theta_P(e_x)$ , if  $E_{\lambda}e_x = \gamma E_{\lambda}e_y$ , then  $\lambda = \cos \frac{j}{\tau}\pi$  for some even j.
	- (*c*) *For any*  $\lambda \in \Theta_P(e_x)$ , if  $E_{\lambda}e_x = -\gamma E_{\lambda}e_y$ , then  $\lambda = \cos \frac{j}{\tau} \pi$  *for some odd j.*

<span id="page-9-5"></span>*Moreover, if*  $\Gamma$  *is regular, then*  $\gamma = 1$  *in both*  $(B)$  $(B)$  $(B)$  *and*  $(C)$  $(C)$  $(C)$ *.* 

**Proof.** Let U be the time evolution matrix of Γ, and  $P = \sum_{\lambda \in \sigma(P)} \lambda E_{\lambda}$  the spectral decomposition. We assume [\(A\),](#page-9-3) and prove [\(B\).](#page-9-1) By (A), we have  $U^{\tau}d^*e_x = \gamma d^*e_y$  for some  $\gamma \in \mathbb{C}$  with  $|\gamma| = 1$ . Since  $U = R(2d^*d - I_{\mathcal{A}})$  and  $d^*$  are real matrices,  $\gamma \in \{\pm 1\}$  follows. By [\(A\)](#page-9-3) again and Lemma [6.3,](#page-9-0) we have

$$
T_{\tau}(P)e_x = (dU^{\tau}d^*)e_x = d(\gamma d^*e_y) = (dd^*)\gamma e_y = \gamma e_y.
$$

Next, we assume [\(B\),](#page-9-1) and prove [\(C\).](#page-9-2) Take  $\gamma \in \{\pm 1\}$  such that  $T_{\tau}(P)e_x = \gamma e_y$ . Since

$$
1=\|\gamma e_y\|^2=\|T_\tau(P)e_x\|^2=\|\sum_{\lambda\in\sigma(P)}T_\tau(\lambda)(E_\lambda e_x)\|^2=\sum_{\lambda\in\sigma(P)}|T_\tau(\lambda)|^2\|E_\lambda e_x\|^2\leq\sum_{\lambda\in\sigma(P)}\|E_\lambda e_x\|^2=1
$$

by Lemmas [2.2](#page-3-3) and [6.2,](#page-8-2) we have  $T_{\tau}(\lambda) \in \{\pm 1\}$  for every  $\lambda \in \Theta_P(e_x)$ . Also, from  $T_{\tau}(P)e_x = \gamma e_y$ , we have by Lemma [2.2,](#page-3-3)  $T_{\tau}(\lambda)E_{\lambda}e_{x} = \gamma E_{\lambda}e_{y}$  for every  $\lambda \in \sigma(P)$ . These together with Lemma [6.1](#page-8-1) implies [\(C\).](#page-9-2)

We assume  $(C)$ , and prove  $(A)$ . By Lemmas [6.3](#page-9-0) and [2.2,](#page-3-3) we have

$$
(d^*e_y)^*U^\tau(d^*e_x) = e_y^*T_\tau(P)e_x = \sum_{\lambda \in \sigma(P)} T_\tau(\lambda)(E_\lambda e_y)^*(E_\lambda e_x).
$$
\n(6.1)

Let  $\Theta_{\pm} := {\lambda \in \Theta_P(e_x) | E_{\lambda}e_x = \pm \gamma E_{\lambda}e_y}.$  Then, we have

$$
\sum_{\lambda \in \sigma(P)} T_{\tau}(\lambda)(E_{\lambda}e_{y})^{*}(E_{\lambda}e_{x}) = \sum_{\lambda \in \Theta_{P}(e_{x})} T_{\tau}(\lambda)(E_{\lambda}e_{y})^{*}(E_{\lambda}e_{x})
$$
\n
$$
= \sum_{\lambda \in \Theta_{+}} T_{\tau}(\lambda)(E_{\lambda}e_{y})^{*}(E_{\lambda}e_{x}) + \sum_{\lambda \in \Theta_{-}} T_{\tau}(\lambda)(E_{\lambda}e_{y})^{*}(E_{\lambda}e_{x})
$$
\n
$$
= \sum_{\lambda \in \Theta_{+}} (E_{\lambda}e_{y})^{*}(E_{\lambda}e_{x}) + \sum_{\lambda \in \Theta_{-}} -(E_{\lambda}e_{y})^{*}(E_{\lambda}e_{x}) \qquad \text{(by (b) and (c))}
$$
\n
$$
= \sum_{\lambda \in \Theta_{+}} \gamma(E_{\lambda}e_{y})^{*}(E_{\lambda}e_{y}) + \sum_{\lambda \in \Theta_{-}} \gamma(E_{\lambda}e_{y})^{*}(E_{\lambda}e_{y})
$$
\n
$$
= \gamma \sum_{\lambda \in \sigma(P)} (E_{\lambda}e_{y})^{*}(E_{\lambda}e_{y})
$$
\n
$$
= \gamma.
$$

By Lemma [6.4,](#page-9-6) we derive [\(A\)](#page-9-3) as desired.

Finally, we assume that  $\Gamma$  is regular of valency k and [\(A\)](#page-9-3) holds, and prove  $\gamma = 1$  in [\(B\)](#page-9-1) and [\(C\).](#page-9-2) By [\(A\),](#page-9-3) we have  $U^{\tau}d^*e_x = \gamma d^*e_y$  for some  $\gamma \in {\pm 1}$ . As discussed above, [\(B\)](#page-9-1) and [\(C\)](#page-9-2) hold for this γ. By the assumption that Γ is regular, we see that  $P = A/k$ , where A is the adjacency matrix of Γ. Hence, 1 is an eigenvector of  $P$ , and the all-ones vector j is an eigenvector of  $P$  belonging to 1. This implies  $1 \in \Theta_P(e_x)$ . By [\(b\)](#page-9-4) and [\(c\)](#page-9-5) in [\(C\),](#page-9-2)  $E_1e_x = \gamma E_1e_y$  holds. Thus,

$$
1 = \langle \mathbf{j}, e_x \rangle = \langle \mathbf{j}, E_1 e_x \rangle = \langle \mathbf{j}, \gamma E_1 e_y \rangle = \gamma \langle \mathbf{j}, e_y \rangle = \gamma.
$$

Therefore,  $\gamma = 1$  holds.

Currently, we have not found examples of non-regular graphs that admit perfect state transfer with  $\gamma = -1.$ 

## 7 Circulant graphs with valency 3

In this section, we show that 3-regular circulant graphs do not admit perfect state transfer. We note from [\(4.1\)](#page-6-3) that any 3-regular circulant graph  $X(\mathbb{Z}_n, S)$  requires that n is even,  $a \neq 0, \frac{n}{2}$  and  $S = {\pm a, \frac{n}{2}}.$ 

Let  $a', b' \in \mathbb{Z}$ , and set  $a := a' + n\mathbb{Z}$  and  $b := b' + n\mathbb{Z}$ . In the case of  $n \in 2\mathbb{Z}$ , a is said to be *odd* if a ′ is an odd integer, and *even* otherwise. The elements a and b are said to be *coprime* if the greatest common divisor of a', b' and n equals 1. In other words, they are coprime if  $n_1a + n_2b = 1$  in  $\mathbb{Z}_n$  for some integers  $n_1$  and  $n_2$ . Also, for a function f, we will write  $f(a')$  as  $f(a)$  if  $f(a') = f(a'')$  for any  $a'' \in a$ . For example,  $e^{\frac{2\pi}{n}a} = e^{\frac{2\pi}{n}a'}$  and  $\cos(\frac{2\pi a}{n}) = \cos(\frac{2\pi a'}{n})$ .

Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, l\})$  be a 3-regular circulant graph, and let  $P = P(X)$ . Then by Lemma [4.3,](#page-7-3) the eigenvalues of P are

$$
\lambda_j = \frac{\zeta_{2l}^{aj} + \zeta_{2l}^{-aj} + \zeta_{2l}^{jl}}{3} = \frac{2\cos\frac{aj}{l}\pi + (-1)^j}{3}
$$

for  $j \in [2l]$ . Note that for X to be connected, a and l are coprime since there exists a path from 0 to 1, i.e., there exist non-negative integers  $n_1$  and  $n_2$  such that  $n_1a + n_2l = 1$  in  $\mathbb{Z}_{2l}$ . Here, the length of the path is  $n_1 + n_2$ .

**Definition 7.1.** Let r be a rational number  $p/q$  with  $p \in \mathbb{Z}$  and  $q \in \mathbb{Z}_{\geq 1}$  where p and q are coprime. Then define  $N(r) := q$ .

<span id="page-10-0"></span>**Lemma 7.2** (Fact 1.1 in [\[3\]](#page-22-14)). Let  $r \in \mathbb{Q}$ . The numbers 1 and  $\cos(r\pi)$  are  $\mathbb{Q}$ -linearly independent if *and only if*  $N(r) \geq 4$ *. In particular,*  $\mathbb{Q} \cap \{ \cos(r\pi) \mid r \in \mathbb{Q} \} = \{0, \pm 1, \pm \frac{1}{2}\}.$ 

 $\Box$ 

<span id="page-11-0"></span>**Lemma 7.3** (Theorem 1.2 in [\[3\]](#page-22-14)). *Let*  $r_1, r_2$  *be rational numbers such that neither*  $r_1 - r_2$  *nor*  $r_1 + r_2$ *is an integer. Then the following are equivalent.*

- (i) *The numbers*  $1, \cos(r_1\pi)$  *and*  $\cos(r_2\pi)$  *are*  $\mathbb{Q}$ *-linearly independent.*
- (ii)  $N(r_i) \geq 4$  *for*  $j \in \{1, 2\}$ *, and*  $(N(r_1), N(r_2)) \neq (5, 5)$ *.*

**Theorem 7.4.** Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, l\})$  be a connected 3-regular graph, where  $a \in \{1, ..., l-1\}$ . *Then, perfect state transfer does not occur on* X *between vertex type states.*

**Proof.** Let  $x, y \in \mathbb{Z}_{2l}$  be distinct vertices in X. We suppose that perfect state transfer occurs from  $d^*e_x$  to  $d^*e_y$  at time  $\tau$ , and want to derive a contradiction. Let  $P = P(X)$ . By Lemma [4.5,](#page-7-1) the eigenvalue  $\lambda_1 = \frac{1}{3}(2 \cos \frac{a}{l}\pi - 1)$  of P is in the eigenvalue support  $\Theta_P(e_x)$ . Thus, Theorem [6.5](#page-9-7) implies that there exists an integer s such that

<span id="page-11-1"></span>
$$
\frac{1}{3}(2\cos\frac{a}{l}\pi - 1) = \cos\frac{s}{\tau}\pi.
$$
 (7.1)

In particular, this means that the elements 1,  $\cos \frac{a}{l}\pi$  and  $\cos \frac{a}{l}\pi$  are not Q-linearly independent.

To apply Lemma [7.3,](#page-11-0) we want to check that neither  $\frac{a}{l} + \frac{s}{\tau}$  nor  $\frac{a}{l} - \frac{s}{\tau}$  is an integer. We suppose that either  $\frac{a}{l} + \frac{s}{\tau}$  or  $\frac{a}{l} - \frac{s}{\tau}$  is an integer. Then we have  $\cos \frac{a}{l}\pi = \pm \cos \frac{s}{\tau}\pi$ . If  $\cos \frac{a}{l}\pi = -\cos \frac{s}{\tau}\pi$ , then Equality [\(7.1\)](#page-11-1) if and only if  $\cos \frac{s}{\tau} \tau = -\frac{1}{5}$ . This contradicts  $\cos \mathbb{Q} \tau \cap \mathbb{Q} = {\pm 1, \pm \frac{1}{2}, 0}$  which is asserted in Lemma [7.2.](#page-10-0) Thus, we have  $\cos \frac{a}{l}\pi \neq -\cos \frac{s}{l}\pi$ . Then Equality [\(7.1\)](#page-11-1) if and only if  $\cos \frac{a}{l}\pi = -1$ . In particular, we have  $a \in l\mathbb{Z}$ , but this is inconsistent with the way a is taken. We are therefore in a situation to apply Lemma [7.3,](#page-11-0) and hence either  $N(\frac{a}{l}) < 4$ ,  $N(\frac{s}{l}) < 4$ , or  $N(\frac{a}{l}) = N(\frac{s}{l}) = 5$  happens.

First, we want to reject  $N(\frac{a}{l}) < 4$ . We assume  $N(\frac{a}{l}) < 4$ . Since a and l are coprime, we have  $l = N(\frac{a}{l}) < 4$ , i.e.,  $l \leq 3$ . If  $l = 2$ , then  $X = X(\mathbb{Z}_4, {\{\pm 1, 2\}})$  is isomorphic to the complete graph  $K_4$ with 4 vertices. However,  $K_4$  does not admit perfect state transfer. See Section 4.3 in [\[17\]](#page-22-7) for details. Let  $l = 3$ . If  $a = 1$ , then  $X = X(\mathbb{Z}_6, \{\pm 1, 3\})$  is isomorphic to the complete bipartite graph  $K_{3,3}$ . It has also been shown in Lemma 4.2 of [\[17\]](#page-22-7) that this graph does not admit perfect state transfer. If  $a=2$ , then we have by  $(7.1) \cos \frac{s}{\tau} \pi = \frac{-1+2 \cos \frac{2}{3}\pi}{3} = -\frac{2}{3}$ . However, this contradicts Lemma [7.2](#page-10-0)

Next, we want to reject  $N(\frac{s}{\tau}) < 4$ . We assume  $N(\frac{s}{\tau}) < 4$ . Then we have  $\cos \frac{s}{\tau} \pi \in \mathbb{Q}$  by Lemma [7.2,](#page-10-0) so from Equality [\(7.1\)](#page-11-1) we obtain  $\cos \frac{a}{l}\pi \in \mathbb{Q}$ . Thus we have  $N(\frac{a}{l}) < 4$  by Lemma [7.2.](#page-10-0) The situation has been returned to the case  $N(\frac{a}{l}) < 4$ , which has already been rejected.

Finally, we want to reject the case  $N(\frac{a}{l}) = N(\frac{s}{\tau}) = 5$ . Then we have

$$
\cos\frac{a}{l}\pi, \cos\frac{s}{\tau}\pi \in \left\{\frac{\pm 1 \pm \sqrt{5}}{4}\right\}.
$$

However, it can be seen by inspecting the coefficient of  $\sqrt{5}$  that none of the values satisfy Equality [\(7.1\)](#page-11-1).  $\Box$ 

### 8 Circulant graphs with valency 4 and  $a + b = l$

Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  be a 4-regular circulant graph. Zhan [\[23\]](#page-23-2) proved that X admits perfect state transfer between vertex type states when l is odd and  $a + b = l$ . We completely characterize X which admits perfect state transfer between vertex type states for every l. First, we consider the case of  $l \not\equiv 0 \pmod{4}$  and  $a + b = l$ .

<span id="page-11-4"></span>**Lemma 8.1.** Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  be a 4-regular circulant graph. Assume that  $a + b = l$  with  $l \not\equiv 0 \pmod{4}$ .

- <span id="page-11-3"></span><span id="page-11-2"></span>(i) If  $l \equiv 2 \pmod{4}$ , then X admits perfect state transfer from  $d^*e_x$  to  $d^*e_{x+l}$  at time l.
- (ii) If l is odd, then X admits perfect state transfer from  $d^*e_x$  to  $d^*e_{x+l}$  at time 2l.

#### *Moreover, if* X *is connected, then these times at which perfect state transfer occurs are minimum.*

**Proof.** First, we prove the case [\(i\).](#page-11-2) Let  $l \equiv 2 \pmod{4}$ . By Lemma [4.1](#page-6-4) [\(ii\)](#page-11-3) and Theorem [3.3,](#page-5-3) we may assume that  $x = 0$  without loss of generality. Let  $P = P(X)$ . We will check the three conditions [\(a\)–](#page-9-8)[\(c\)](#page-9-5) of [\(C\)](#page-9-2) in Theorem [6.5.](#page-9-7) Before that, we calculate eigenvalues and some projections. The eigenvalues of P are

<span id="page-12-0"></span>
$$
\lambda_j = \begin{cases} 0 & \text{if } j \text{ is odd,} \\ \cos \frac{aj}{l} \pi & \text{if } j \text{ is even} \end{cases}
$$
 (8.1)

for  $j \in [2l]$ , because

$$
\lambda_{j} = \frac{\zeta_{2l}^{aj} + \zeta_{2l}^{-aj} + \zeta_{2l}^{bj} + \zeta_{2l}^{-bj}}{4}
$$
 (by Lemma 4.3)  
\n
$$
= \frac{2 \cos \frac{aj}{l} \pi + 2 \cos \frac{bj}{l} \pi}{4}
$$
  
\n
$$
= \frac{2 \cos \frac{aj}{l} \pi + 2 \cos \frac{(l-a)j}{l} \pi}{4}
$$
 (by  $a + b = l$ )  
\n
$$
= \frac{1 + (-1)^{j}}{2} \cos \frac{aj}{l} \pi.
$$

Next we verify

$$
E_{\lambda}e_0 = \begin{cases} -E_{\lambda}e_l & \text{if } \lambda = 0\\ E_{\lambda}e_l & \text{otherwise} \end{cases}
$$
 (8.2)

Recall the vectors  $u_j$  defined by [\(4.3\)](#page-7-2). We have

<span id="page-12-4"></span><span id="page-12-1"></span>
$$
\langle u_j, e_0 \rangle = \zeta_{2l}^0 = 1 \tag{8.3}
$$

and

<span id="page-12-2"></span>
$$
\langle u_j, e_l \rangle = \zeta_{2l}^{lj} = \zeta_2^j = (-1)^j. \tag{8.4}
$$

Let  $\mathcal{I}_{\lambda} = \{j \in [2l] | \lambda_j = \lambda\}$  for a real number  $\lambda$ . We check

<span id="page-12-3"></span>
$$
\mathcal{I}_0 = \{1, 3, \dots, 2l - 1\}.\tag{8.5}
$$

By [\(8.1\)](#page-12-0), we immediately have  $\mathcal{I}_0 \supset \{1, 3, \ldots, 2l-1\}$ . We suppose that an even integer j satisfies  $\lambda_j = 0$ . Then by [\(8.1\)](#page-12-0) we have  $0 = \lambda_j = \cos \frac{aj\pi}{l}$ , so there exists an odd integer s such that  $\frac{aj\pi}{l} = \frac{s}{2}\pi$ . Thus, we obtain  $j = \frac{ls}{2a} = \frac{\frac{l}{2}s}{a}$  $\frac{5s}{a}$ . Since  $l \equiv 2 \pmod{4}$ , both  $\frac{l}{2}$  and s are odd, which contradicts that j is even. Therefore, we have  $\mathcal{I}_0 = \{1, 3, \ldots, 2l - 1\}$ . Let  $\lambda \in \sigma(P)$ , and calculate  $E_{\lambda}e_0$ . If  $\lambda = 0$ , then

$$
2l \cdot E_0 e_0 = \sum_{j \in \mathcal{I}_0} u_j u_j^* e_0 \qquad \qquad \text{(by (2.2))}
$$

$$
=\sum_{j\in\mathcal{I}_0}u_j\cdot1\tag{b y (8.3)}
$$

$$
= -\sum_{j \in \mathcal{I}_0} u_j \cdot (-1)
$$
  
=  $-\sum_{j \in \mathcal{I}_0} u_j u_j^* e_l$  (by (8.4) and (8.5))  
=  $-2l \cdot E_0 e_l$ . (by (2.2))

On the other hand, if  $\lambda \neq 0$ , then from Equality [\(8.5\)](#page-12-3) it is guaranteed that the indices in the set  $[2l] \setminus \mathcal{I}_0$  are even. Thus we have

$$
2l \cdot E_{\lambda} e_0 = \sum_{j \in \mathcal{I}_{\lambda}} u_j u_j^* e_0 \qquad \qquad \text{(by (2.2))}
$$

$$
=\sum_{j\in\mathcal{I}_{\lambda}}u_j\cdot 1\tag{b y (8.3)}
$$

$$
=\sum_{j\in\mathcal{I}_{\lambda}}u_{j}u_{j}^{*}e_{l}\tag{b y (8.4)}
$$

$$
= 2l \cdot E_{\lambda} e_l.
$$

Next, we check the three conditions [\(a\)–](#page-9-8)[\(c\)](#page-9-5) of [\(C\)](#page-9-2) in Theorem [6.5](#page-9-7) with  $\gamma = 1$  and  $\tau = l$ . First, [\(a\)](#page-9-8) follows from [\(8.2\)](#page-12-4). Next we check [\(b\).](#page-9-4) Note that  $\Theta_P(e_0) = \sigma(P)$  holds by Lemma [4.5.](#page-7-1) Let  $\lambda \in \sigma(P)$ , and assume  $E_{\lambda}e_0 = E_{\lambda}e_l$ . Then,  $\lambda \neq 0$  follows from [\(8.2\)](#page-12-4), and this together with [\(8.5\)](#page-12-3) implies that  $\lambda = \lambda_j$  holds for some even j. From [\(8.1\)](#page-12-0), we see that [\(b\)](#page-9-4) is satisfied. We check [\(c\).](#page-9-5) Assume  $E_{\lambda}e_0 = -E_{\lambda}e_l$ . Then, by [\(8.2\)](#page-12-4), we have  $\lambda = 0 = \cos \frac{\pi}{2} = \cos \frac{\frac{1}{2}}{l}\pi$ . Since  $l/2$  is odd, we see [\(c\)](#page-9-5) is satisfied. Therefore, by Theorem [6.5,](#page-9-7) we see that X admits perfect state transfer from  $d^*e_0$  to  $d^*e_l$ at time l.

Finally, we show minimality of the time at which perfect state transfer occurs. By Theorem [6.5](#page-9-7) [\(C\),](#page-9-2) if perfect state transfer occurs at time  $\tau$ , then  $T_{\tau}(\lambda) = \pm 1$  for any  $\lambda \in \Theta_P(e_0)$ . By [\(8.1\)](#page-12-0), we have  $0 \in \sigma(P) = \Theta_P(e_0)$ , so  $\pm 1 = T_\tau(0) = T_\tau(\cos \frac{\pi}{2}) = \cos \frac{\tau}{2}\pi$ , and hence  $\frac{\tau}{2} \in \mathbb{Z}$ , i.e.,

<span id="page-13-0"></span>
$$
\tau \in 2\mathbb{Z}.\tag{8.6}
$$

In addition, since  $l \equiv 2 \pmod{4}$ ,  $\frac{l}{2} - 1$  is even integer. Thus, [\(8.1\)](#page-12-0) implies that  $\cos \frac{a(\frac{l}{2}-1)\pi}{l} \in \sigma(P)$ , and hence

$$
\pm 1 = T_{\tau} \left( \cos \frac{a(\frac{l}{2} - 1)}{l} \pi \right) = \cos \frac{a \tau(\frac{l}{2} - 1)}{l} \pi.
$$

We have  $\frac{a\tau(\frac{1}{2}-1)}{l} \in \mathbb{Z}$ , so  $a\tau(\frac{l}{2}-1) \in l\mathbb{Z} \subset \frac{l}{2}\mathbb{Z}$ . Since a and  $\frac{l}{2}$  are coprime by the connectivity of X, and  $\frac{l}{2} - 1$  and  $\frac{l}{2}$  are coprime, we have  $\tau \in \frac{l}{2}\mathbb{Z}$ . Hence, by [\(8.6\)](#page-13-0), we obtain  $\tau \in l\mathbb{Z}$ , i.e.,  $\tau \geq l$ . The minimum time at which perfect state transfer occurs is  $l$ .

The case [\(ii\)](#page-11-3) has already been shown in Theorem 6.1 of [\[23\]](#page-23-2). We can show that the time  $l$  is minimum in the same manner in [\(i\).](#page-11-2) □

In contrast to the above lemma, perfect state transfer does not occur if  $l \equiv 0 \pmod{4}$ .

**Lemma 8.2.** Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  be a connected 4-regular graph. Assume that  $a + b = l$ . If l ≡ 0 (mod 4)*, then* X *does not admit perfect state transfer between vertex type states.*

**Proof.** As confirmed in Lemma [4.2,](#page-7-0) if perfect state transfer occurs from  $d^*e_x$  to  $d^*e_y$ , then  $y = x + l$ . By Lemma [4.1](#page-6-4) (i) and Theorem [3.3,](#page-5-3) we may assume that  $x = 0$  without loss of generality. We check that the first condition of Theorem [6.5](#page-9-7) (C) does not hold. We use the same notation as in the proof of Lemma [8.1.](#page-11-4) Since X is connected, either a or b is odd. Furthermore, by the assumptions  $a + b = l$ and  $l \equiv 0 \pmod{4}$ , both a and b are odd. Noting that  $\frac{l}{2}$  is even, we have by  $(8.1)$ ,

$$
\lambda_1 = 0
$$
 and  $\lambda_{\frac{l}{2}} = \cos\left(\frac{a \cdot \frac{l}{2}}{l}\pi\right) = \cos\frac{a}{2}\pi = 0,$ 

and obtain

<span id="page-13-1"></span>
$$
1, \frac{l}{2} \in \mathcal{I}_0 = \{ j \in [2l] \mid \lambda_j = 0 \}.
$$
\n(8.7)

By Equalities [\(8.3\)](#page-12-1) and [\(8.4\)](#page-12-2), we have  $2l \cdot E_0 e_0 = \sum_{j \in \mathcal{I}_0} u_j$  and  $2l \cdot E_0 e_l = \sum_{j \in \mathcal{I}_0} (-1)^j u_j$ . Supposing that the first condition of Theorem [6.5](#page-9-7) (C) holds, from linearly independence of eigenvectors we obtain either  $1 = (-1)^j$  for any  $j \in \mathcal{I}_0$ , or  $1 = -(-1)^j$  for any  $j \in \mathcal{I}_0$ . However, by  $(8.7)$  and  $\frac{l}{2} \in 2\mathbb{Z}$ , none of the above is valid. We see that perfect state transfer does not occur.

## 9 Circulant graphs with valency 4 and  $a + b \neq l$

Following the previous section, we study 4-regular circulant graphs  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  with  $a+b \neq 0$ l. To conclude first, such circulant graphs do not admit perfect state transfer. However, its proof requires a lot of effort. To begin, we check obvious conditions required by the circulant graph  $X$  with valency 4. For the graph  $X$  to have valency 4, the following must be satisfied:

<span id="page-14-4"></span><span id="page-14-3"></span>
$$
a, b \not\equiv 0 \pmod{l},\tag{9.1}
$$

$$
a \pm b \not\equiv 0 \pmod{2l}.\tag{9.2}
$$

Furthermore, we may assume that  $a, b \equiv 1, 2, \ldots, l - 1 \pmod{2l}$  without loss of generality, and since  $a - b \equiv \pm (l - 2), \ldots, \pm 1 \pmod{2l}$ , we must have

<span id="page-14-5"></span>
$$
a - b \not\equiv l \pmod{2l}.\tag{9.3}
$$

#### 9.1 Necessary conditions for a graph to admit perfect state transfer

A complex number  $\alpha$  is said to be an *algebraic integer* if there exists a monic polynomial  $p(x) \in \mathbb{Z}[x]$ such that  $p(\alpha) = 0$ . Let  $\Omega$  denote the set of algebraic integers. The set of algebraic integers forms a ring, in particular,  $\Omega$  is closed under addition and multiplication. A specific example is that  $\zeta_m^k$  and  $\zeta_m^{-k}$  are algebraic integers because  $(\zeta_m^k)^m = (\zeta_m^{-k})^m = 1$ . Hence, for a rational number  $2k/m$ ,  $2 \cos \frac{2k}{m} \pi = \zeta_m^k + \zeta_m^{-k}$  is also an algebraic integer.

Let K be a finite extension field of the field Q of rational numbers. The set  $\{\omega_1, \ldots, \omega_t\}$  is an *integral basis* for K when every element of  $K \cap \Omega$  is uniquely expressible as a Z-linear combination of elements of the set. For example, an integral basis for  $\mathbb Q$  is  $\{1\}$ , i.e.,  $\mathbb Q \cap \Omega = \mathbb Z$ , and an integral basis for  $\mathbb{Q}(\sqrt{2})$  is  $\{1,\sqrt{2}\}$ , i.e.,  $\mathbb{Q}(\sqrt{2}) \cap \Omega = \{p+q\sqrt{2} \mid p,q \in \mathbb{Z}\}\.$  Readers who want to know more details on integral bases should refer to [\[15\]](#page-22-15).

<span id="page-14-0"></span>**Lemma 9.1.** Let  $\Gamma$  be a graph with the discriminant P, and let x, y be vertices of  $\Gamma$ . If perfect state *transfer occurs from*  $d^*e_x$  *to*  $d^*e_y$  *at time*  $\tau$ *, then*  $2\lambda$  *is an algebraic integer for any*  $\lambda \in \Theta_P(e_x)$ *.* 

**Proof.** Assume perfect state transfer occurs from  $d^*e_x$  to  $d^*e_y$  at time  $\tau$ . Let  $\lambda \in \Theta_P(e_x)$ . By Theorem [6.5,](#page-9-7) we have  $\lambda = \cos \frac{j}{\tau} \pi$  for some integer j. Hence  $2\lambda$  is an algebraic integer.

<span id="page-14-6"></span>**Lemma 9.2.** Let  $l \in \mathbb{Z}_{\geq 1}$ . If a 4*-regular circulant graph*  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  *admits perfect state transfer, then*

<span id="page-14-1"></span>
$$
\frac{\zeta_{2l}^a + \zeta_{2l}^{-a} + \zeta_{2l}^b + \zeta_{2l}^{-b}}{2}
$$
\n(9.4)

*is an algebraic integer.*

**Proof.** From Lemmas [4.3](#page-7-3) and [4.5,](#page-7-1) we have  $(\zeta_{2l}^a + \zeta_{2l}^b + \zeta_{2l}^b) / 4 \in \sigma(P(X)) = \Theta_{P(X)}(e_0)$ . Therefore, Lemma [9.1](#page-14-0) implies that if  $X$  admits perfect state transfer, then  $(9.4)$  is an algebraic integer.

<span id="page-14-2"></span>**Lemma 9.3.** Let l be an odd integer, and let c be an integer. Then we have  $\zeta_{2l}^c = (-1)^c \zeta_l^{\varepsilon(c)}$  $\int_l^{\varepsilon(c)}$ , where

$$
\varepsilon(c) = \begin{cases} \frac{c}{2} & \text{if } c \text{ is even,} \\ \frac{c+l}{2} & \text{if } c \text{ is odd.} \end{cases}
$$

*In particular, if a* 4*-regular circulant graph*  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  *admits perfect state transfer, then* 

$$
\frac{\zeta_{2l}^a + \zeta_{2l}^{-a} + \zeta_{2l}^b + \zeta_{2l}^{-b}}{2} = \frac{(-1)^a \zeta_l^{\varepsilon(a)} + (-1)^a \zeta_l^{-\varepsilon(a)} + (-1)^b \zeta_l^{\varepsilon(b)} + (-1)^b \zeta_l^{-\varepsilon(b)}}
$$

*is an algebraic integer.*

**Proof.** If c is even, then we have  $\zeta_{2l}^c = \zeta_l^{\frac{c}{2}}$ . If c is odd, then we have  $\zeta_{2l}^c = -\zeta_{2l}^l \zeta_{2l}^c = -\zeta_{2l}^{c+l} = -\zeta_l^{\frac{c+l}{2}}$ . Hence these together with Lemma [9.1](#page-14-0) implies the desired result.

By this lemma, what we need to discuss is whether

<span id="page-15-3"></span>
$$
\Delta = \frac{\pm \zeta_m^p \pm \zeta_m^{-p} \pm \zeta_m^q \pm \zeta_m^{-q}}{2} \tag{9.5}
$$

is an algebraic integer for some positive integers  $m$ ,  $p$  and  $q$ . To do so, we once forget the condition  $a + b \neq l$  for the exponents with respect to  $\zeta_{2l}$  and study a sufficient condition for  $\Delta$  not to be an algebraic integer. Readers who want to first know its sufficient condition should refer to Lemma [9.16.](#page-19-0) Due to circumstances of this sufficient condition, we first need to study Q-linearly independence of  $\zeta_m^p, \zeta_m^{-p}, \zeta_m^q, \zeta_m^{-q}.$ 

#### 9.2 Q-linearly independence

<span id="page-15-1"></span>The following is easily verified.

**Lemma 9.4.** Let s, t be integers such that  $s \pm t \neq 0 \pmod{5}$ . Then  $\cos \frac{s}{5} \pi$  and  $\cos \frac{t}{5} \pi$  are Q-linearly *independent.*

<span id="page-15-0"></span>**Lemma 9.5.** Let m be a positive integer, and let p be an integer. Then  $4p \equiv 0 \pmod{m}$  if and only *if*  $N(\frac{2p}{m}) \in \{1, 2\}$ .

**Proof.** First, we suppose that  $4p \equiv 0 \pmod{m}$ . There exists an integer k such that  $4p = km$ , i.e.,  $\frac{2p}{m} = \frac{k}{2}$ . Thus, we have  $N(\frac{2p}{m}) = N(\frac{k}{2}) \leq 2$ . Conversely, we suppose that  $N(\frac{2p}{m}) \in \{1,2\}$ . If  $N(\frac{2p}{m}) \in \{1,2\}$ , then there exists an integer s such that  $\frac{2p}{m} = \frac{s}{2}$ , and hence  $4p = ms \equiv 0 \pmod{m}$ .

<span id="page-15-2"></span>**Lemma 9.6.** Let m be a positive integer, and let p, q be distinct integers. If  $4p, 4q, 2(p \pm q) \neq 0$ (mod *m*), then  $\cos \frac{2p}{m}\pi$  and  $\cos \frac{2q}{m}\pi$  are Q-linearly independent.

**Proof.** By Lemma [9.5,](#page-15-0) we have  $N(\frac{2p}{m})$ ,  $N(\frac{2q}{m}) \geq 3$ . First, we suppose that  $N(\frac{2p}{m}) = N(\frac{2q}{m}) = 3$ . Then, there exist  $s, t \in \mathbb{Z} \setminus 3\mathbb{Z}$  such that  $\frac{2p}{m} = \frac{s}{3}$  and  $\frac{2q}{m} = \frac{t}{3}$ . Since  $\frac{m}{3}(s+t) = 2(p+q) \not\equiv 0 \pmod{m}$ , either  $s \equiv t \equiv 1 \pmod{3}$  or  $s \equiv t \equiv 2 \pmod{3}$  has to be held. However, in both cases we have  $2(p-q) = \frac{m}{3}(s-t) \equiv 0 \pmod{m}$ , which contradicts our assumption. Therefore, either  $N(\frac{2p}{m})$  or  $N(\frac{2q}{m})$  is greater than 3. Without loss of generality, we may assume that  $N(\frac{2q}{m}) \geq 4$ .

First, we consider the case  $N(\frac{2p}{m}) = 3$ . Lemma [7.2](#page-10-0) implies that  $\cos \frac{2p}{m}\pi$  is a rational number while cos  $\frac{2q}{m}\pi$  is an irrational number. Moreover, since  $N(\frac{2p}{m})=3$ , we have  $\cos\frac{2p}{m}\pi\neq 0$ . Thus, we see that  $\cos \frac{2p}{m}\pi$  and  $\cos \frac{2q}{m}\pi$  are Q-linearly independent.

Next, we consider the case  $N(\frac{2p}{m}) \ge 4$ . If  $(N(\frac{2p}{m}), N(\frac{2q}{m})) \ne (5,5)$ , then Lemma [7.3](#page-11-0) implies the conclusion. If  $(N(\frac{2p}{m}), N(\frac{2q}{m})) = (5, 5)$ , then there exist  $s, t \in \mathbb{Z} \setminus 5\mathbb{Z}$  such that  $\frac{2p}{m} = \frac{s}{5}$  and  $\frac{2q}{m} = \frac{t}{5}$ . By our assumption, we have  $s \pm t \neq 0 \pmod{5}$  since

$$
2(p \pm q) \not\equiv 0 \pmod{m} \iff 10(p \pm q) \not\equiv 0 \pmod{5m}
$$
  

$$
\iff m(s \pm t) \not\equiv 0 \pmod{5m}
$$
  

$$
\iff s \pm t \not\equiv 0 \pmod{5}.
$$

Therefore, Lemma [9.4](#page-15-1) leads to the conclusion.

口

<span id="page-16-1"></span>**Lemma 9.7.** Let m be a positive integer, and let p, q be distinct integers. If  $4p, 4q, 2(p \pm q) \neq 0$ (mod *m*), then  $\sin \frac{2p}{m}\pi$  and  $\sin \frac{2q}{m}\pi$  are Q-linearly independent.

**Proof.** Since  $\sin \frac{2p}{m}\pi = \cos \frac{2(m-4p)}{4m}\pi$  and  $\sin \frac{2q}{m}\pi = \cos \frac{2(m-4q)}{4m}\pi$ , it is sufficient to confirm that Lemma [9.6](#page-15-2) is applicable, i.e.,  $4(m - 4p)$ ,  $4(m - 4q)$ ,  $4(m - 2p - 2q)$ ,  $8(p - q) \not\equiv 0 \pmod{4m}$ . This can be verified straightforwardly.

<span id="page-16-4"></span>**Lemma 9.8.** Let m be a positive integer, and let p, q be distinct integers. If  $4p, 4q, 2(p \pm q) \neq 0$ (mod *m*), then the numbers  $\zeta_m^p, \zeta_m^{-p}, \zeta_m^q, \zeta_m^{-q}$  are Q-linearly independent.

**Proof.** We assume that a Q-linear combination

<span id="page-16-0"></span>
$$
c_1\zeta_m^p + c_2\zeta_m^{-p} + c_3\zeta_m^q + c_4\zeta_m^{-q} = 0.
$$
\n(9.6)

Take the conjugate of both sides and add it to Equality [\(9.6\)](#page-16-0). We have

$$
0 = (c_1 + c_2)(\zeta_m^p + \zeta_m^{-p}) + (c_3 + c_4)(\zeta_m^q + \zeta_m^{-q})
$$
  
= 2(c\_1 + c\_2) cos  $\frac{2p}{m}\pi$  + 2(c\_3 + c\_4) cos  $\frac{2q}{m}\pi$ .

By Lemma [9.6,](#page-15-2) we have  $c_1 + c_2 = 0$  and  $c_3 + c_4 = 0$ . Substituting them into Equality [\(9.6\)](#page-16-0), we have

$$
0 = c_1 \zeta_m^p - c_1 \zeta_m^{-p} + c_3 \zeta_m^q - c_3 \zeta_m^{-q}
$$
  
=  $c_1 (\zeta_m^p - \zeta_m^{-p}) + c_3 (\zeta_m^q - \zeta_m^{-q})$   
=  $2ic_1 \sin \frac{2p}{m} \pi + 2ic_3 \sin \frac{2q}{m} \pi$ .

Lemma [9.7](#page-16-1) implies that  $c_1 = c_3 = 0$ . Therefore, we have  $c_1 = c_2 = c_3 = c_4 = 0$ , and hence we see that  $\zeta_m^p, \zeta_m^{-p}, \zeta_m^q, \zeta_m^{-q}$  are Q-linearly independent. □

#### 9.3 Canonical bases of cyclotomic fields

For a set A and a positive integer k, we denote by  $\begin{pmatrix} A \\ b \end{pmatrix}$ k the set of all  $k$ -subsets of  $A$ , i.e.,

$$
\binom{A}{k} = \{ S \mid S \subset A, |S| = k \}.
$$

For a prime number  $p$ , we define

<span id="page-16-3"></span>
$$
p^* = \begin{cases} 4 & \text{if } p = 2, \\ p & \text{otherwise.} \end{cases}
$$

Let  $\varphi$  denote Euler's totient function, i.e.,  $\varphi(n) = |Z_n^{\times}|$  for a positive integer n. The following is on integral bases of the cyclotomic fields, which plays an important role in this section. We cite a result by Bosma, but slightly change notation.

<span id="page-16-2"></span>**Theorem 9.9** (Theorem 5.1 in [\[4\]](#page-22-16)). *Let* n *be a positive positive integer with*  $n \not\equiv 2 \pmod{4}$ *, and let*  $n = p_1^{f_1} p_2^{f_2} \cdots p_t^{f_t}$  be the prime factorization of n. For each prime factor  $p_j$ , let  $A_{p_j} \in \binom{[p_j]}{\varphi(p_j)}$  $\varphi(p_j^*)$  *, and*  $B_{p_j} = \left[\frac{p_j^{f_j}}{p_j^*}\right]$  *. Then the set*  $I =$  $\sqrt{ }$ J  $\mathcal{L}$  $\frac{t}{\prod}$  $j=1$  $\left(\zeta_{p_j^*}^{a_j} \zeta_{p_j^*}^{b_j}\right)$  $p_j^{f_j}$  $\Bigg)$  $a_1 \in A_{p_1}, \ldots, a_t \in A_{p_t}, b_1 \in B_{p_1}, \ldots, b_t \in B_{p_t}$  $\mathcal{L}$  $\mathcal{L}$  $\left| \right|$ (9.7)

*forms a basis for*  $\mathbb{Q}(\zeta_n)$  *over*  $\mathbb{Q}$ *. Moreover, for every divisor d of n, the set*  $I_d = I \cap \mathbb{Q}(\zeta_d)$  *forms an integral basis for*  $\mathbb{Q}(\zeta_d)$ *.* 

**Example 9.10.** We take  $n = 36 = 2^2 \cdot 3^2$  as an example to help understand Theorem [9.9.](#page-16-2) Since the Qminimal polynomial of  $\zeta_{36}$  is  $x^{12} - x^6 + 1$ , we can take  $1, \zeta_{36}, \zeta_{36}^2, \ldots, \zeta_{36}^{11}$  as a Q-basis of Q( $\zeta_{36}$ ). On the other hand, we can take another basis of  $\mathbb{Q}(\zeta_{36})$  using Theorem [9.9.](#page-16-2) The sets  $A_2 = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$  $\varphi(2^*)$  $= \{0, 1\},\$  $B_2 = \left[\frac{2^2}{2^*}\right]$  $\left[\frac{2^2}{2^*}\right] = \{0\}, \text{and } B_3 = \left[\frac{3^2}{3^*}\right]$  $\left[\frac{3^2}{3^*}\right] = \{0,1,2\}$  are automatically determined, while the set  $A_3 \in {\binom{3}{\varnothing}}$  $\varphi(3^*)$  $\setminus$ 

can be chosen arbitrarily, for example  $A_3 = \{0, 2\}$ . Then,

$$
I = \left\{ \zeta_{2^*}^{a_2} \zeta_{2^2}^{b_3} \zeta_{3^2}^{b_3} \middle| a_2 \in A_2, a_3 \in A_3, b_2 \in B_2, b_3 \in B_3 \right\}
$$
  
=  $\left\{ \zeta_4^{a_2} \zeta_4^{b_2} \zeta_3^{a_3} \zeta_9^{b_3} \middle| a_2 \in \{0, 1\}, a_3 \in \{0, 2\}, b_2 \in \{0\}, b_3 \in \{0, 1, 2\} \right\}$   
=  $\left\{ \zeta_{36}^{0}, \zeta_{36}^{1}, \zeta_{36}^{4}, \zeta_{36}^{5}, \zeta_{36}^{8}, \zeta_{36}^{9}, \zeta_{36}^{13}, \zeta_{36}^{17}, \zeta_{36}^{24}, \zeta_{36}^{28}, \zeta_{36}^{32}, \zeta_{36}^{23} \right\}$ 

forms a Q-basis for  $\mathbb{Q}(\zeta_{36})$  by Theorem [9.9.](#page-16-2) Furthermore,  $I_n = I \cap \mathbb{Q}(\zeta_{36}) = I$  forms an integral basis for  $\mathbb{Q}(\zeta_{36})$ .

## 9.4 Decompositions of  $\zeta_n^x$

Display the prime factorization of an integer  $n \geq 2$  as  $n = p_1^{f_1} p_2^{f_2} \cdots p_t^{f_t}$ , where  $p_1, \ldots, p_t$  are pairwise distinct prime numbers, and  $f_1, \ldots, f_t$  are positive integers. Then, we define the mapping  $\Phi_n: \prod_{j=1}^t [p_j^{f_j}] \to \{\zeta_n^j \mid j \in [n]\}$  by

$$
\Phi_n(x_1, x_2, \dots, x_t) = \prod_{j=1}^t \zeta_{p_j^{s_j}}^{x_j}.
$$

<span id="page-17-0"></span>**Lemma 9.11.** *The mapping*  $\Phi_n$  *is bijective for any integer*  $n \geq 2$ *.* 

**Proof.** Since the number of elements in the domain of  $\Phi_n$  and that in the codomain are equal, it is sufficient to show that  $\Phi_n$  is surjective. Take  $\zeta_n^m \in \{\zeta_{n_\epsilon}^j | j \in [n]\}\)$ . Let  $n = p_1^{f_1} p_2^{f_2} \cdots p_t^{f_t}$  be the prime factorization of *n*. The greatest common divisor of  $np_1^{-f_1}, np_2^{-f_2}, \ldots, np_t^{-f_t}$  is 1, so there exist integers  $x_1, x_2, \ldots, x_t$  such that

$$
m = \sum_{j=1}^{t} x_j (np_j^{-f_j}) = \sum_{j=1}^{t} nx_j p_j^{-f_j}.
$$

Let  $x'_j$  be the remainder of  $x_j$  divided by  $p_j^{f_j}$  for each j. Then, we have  $x'_j \in [p_j^{f_j}]$ , and

$$
\Phi(x'_1, x'_2, \dots, x'_t) = \prod_{j=1}^t \zeta_{p_j^{f_j}}^{x'_j} = \prod_{j=1}^t \zeta_{p_j^{f_j}}^{x_j} = \prod_{j=1}^t \zeta_n^{nx_j p_j^{-f_j}} = \zeta_n^{\sum_{j=1}^t nx_j p_j^{-f_j}} = \zeta_n^m
$$

<span id="page-17-1"></span>Therefore, we see that  $\Phi_n$  is surjective.

**Definition 9.12.** Let  $n \geq 2$  be an integer, and let  $n = p_1^{f_1} p_2^{f_2} \cdots p_t^{f_t}$  be the prime factorization of n, where  $f_1, \ldots, f_t$  are positive integers. Since  $\Phi_n$  is bijective by Lemma [9.11,](#page-17-0) let  $\Psi_n$  be the inverse mapping of  $\Phi_n$ . For  $\zeta_n^x \in {\{\zeta_n^j \mid j \in [n]\}}$ , we write

$$
\Psi_n(\zeta_n^x) = (x_1, x_2, \dots, x_t) \in [p_1^{f_1}] \times [p_2^{f_2}] \times \dots \times [p_t^{f_t}].
$$

For each prime factor  $p_j$ , we define  $\Psi_n^{(p_j)}(\zeta_n^x) := x_j$ . In the case of  $p_j \neq 2$ , we define

$$
\pi_n^{(p_j)}(\zeta_n^x) := x_j' \quad \text{ and } \quad \theta_n^{(p_j)}(\zeta_n^x) := x_j''
$$

 $\Box$ 

.

where  $x'_j \in [p_j]$  and  $x''_j \in [p_j^{f_j-1}]$  such that  $x_j = x'_j p_j^{f_j-1} + x''_j$ . In the case of  $p_j = 2$  and  $f_j \ge 2$ , define

$$
\pi_n^{(2)}(\zeta_n^x) := x'_j
$$
 and  $\theta_n^{(2)}(\zeta_n^x) := x''_j$ 

where  $x'_{j} \in [4]$  and  $x''_{j} \in [2^{f_j - 2}]$  satisfy  $x_{j} = x'_{j} 2^{f_j - 2} + x''_{j}$ .

Note that we do not define  $\Phi_n$  in the case of  $n \equiv 2 \pmod{4}$ , i.e.,  $p_j = 2$  and  $f_j = 1$  since this case is reduced to the case of  $n \equiv 0 \pmod{4}$  in proofs of our results.

**Example 9.13.** We take  $n := 36 = 2^2 \cdot 3^2$  and  $x := 5$  as an example to help understand the mappings in Definition [9.12.](#page-17-1) Then, since  $\zeta_{36}^5 = \zeta_4^1 \zeta_9^8$ , we have  $\Psi_n(\zeta_n^x) = (1,8)$ ,  $\Psi_n^{(2)}(\zeta_n^x) = 1$  and  $\Psi_n^{(3)}(\zeta_n^x) = 8$ . Also, since  $1 = 1 \cdot 2^{2-2} + 0$  and  $8 = 2 \cdot 3^{2-1} + 2$ , we have  $\pi_n^{(2)}(\zeta_n^x) = 1$ ,  $\theta_n^{(2)} = 0$   $\pi_n^{(3)}(\zeta_n^x) = 2$ , and  $\theta_n^{(3)}(\zeta_n^x) = 2.$ 

The reason for defining the functions  $\pi_m^{(p)}$  and  $\theta_m^{(p)}$  is that they are useful tools for checking whether the element

$$
\Delta = \frac{\pm \zeta_m^p \pm \zeta_m^{-p} \pm \zeta_m^q \pm \zeta_m^{-q}}{2}
$$

in [\(9.5\)](#page-15-3) of the cyclotomic field  $\mathbb{Q}(\zeta_m)$  is an algebraic integer. Indeed, the element  $\Delta$  is an algebraic integer if and only if ∆ can be expressed as a Z-linear combination of an integral basis of the cyclotomic field  $\mathbb{Q}(\zeta_m)$ . Therefore, we will construct an integral basis I containing as many of  $\zeta_m^p$ ,  $\zeta_m^{-p}$ ,  $\zeta_m^{-q}$ ,  $\zeta_m^{-q}$ as possible, and then check whether  $\Delta$  is a  $\mathbb{Z}$ -linear combination of I. In fact, we can construct such an integral basis by Theorem [9.9,](#page-16-2) which asserts that  $\zeta_m^x$  is contained in I if  $A_p$  and  $B_p$  are chosen such that  $\pi_m^{(p)}(\zeta_m^x) \in A_p$  and  $\theta_m^{(p)}(\zeta_m^x) \in B_p$  for every prime factor p of m. This is the reason why we introduced  $\pi_m^{(p)}$  and  $\theta_m^{(p)}$ .

In the following subsection, we will determine whether  $\Delta$  is an algebraic integer for some specific values of m, p, q. Before proceeding, we prepare a lemma concerning  $\pi_m^{(p)}$  and  $\theta_m^{(p)}$ .

<span id="page-18-2"></span><span id="page-18-0"></span>**Lemma 9.14.** Let  $n \geq 2$  be an integer, and let  $x \in [n]$ . Let p be a prime factor of n.

(i) *If*  $n \equiv 0 \pmod{4}$ *, then* 

$$
\pi_n^{(p)}(-\zeta_n^x) = \begin{cases} \left(\pi_n^{(2)}(\zeta_n^x) + 2\right) \bmod 4 & \text{if } p = 2, \\ \pi_n^{(p)}(\zeta_n^x) & \text{otherwise,} \end{cases}
$$

and  $\theta_n^{(p)}(-\zeta_n^x) = \theta_n^{(p)}(\zeta_n^x)$  for every prime factor p of n.

<span id="page-18-1"></span>(ii) If 
$$
n \equiv 0 \pmod{3}
$$
, then  $\pi_n^{(3)}(\zeta_n^{-x}) = \begin{cases} \left(3 - \pi_n^{(3)}(\zeta_n^{x})\right) \mod 3 & \text{if } \theta_m^{(3)}(\zeta_m^{x}) = 0, \\ 2 - \pi_n^{(3)}(\zeta_n^{x}) & \text{otherwise.} \end{cases}$ 

**Proof.** Let  $n = p_1^{f_1} p_2^{f_2} \cdots p_t^{f_t}$  be the prime factorization of n, and set  $q_i := p_i^{f_i}$ . Write  $z := \zeta_n^x$  and  $x_i := \Psi_n^{(p_i)}(z)$  for short. First we show [\(i\).](#page-18-0) Assume  $n \equiv 0 \pmod{4}$ . Then we may assume  $p_1 = 2$  and  $f_1 \geq 2$ . We have

$$
-\zeta_n^x = -\zeta_{q_1}^{x_1} \cdot \zeta_{q_2}^{x_2} \cdots \zeta_{q_t}^{x_3} = -\zeta_4^{\pi_4^{(2)}(z)} \zeta_{q_1}^{\theta_4^{(2)}(z)} \zeta_{q_2}^{x_2} \cdots \zeta_{q_t}^{x_3} = \zeta_4^{\pi_4^{(2)}(z) + 2} \zeta_{q_1}^{\theta_4^{(2)}(z)} \zeta_{q_2}^{x_2} \cdots \zeta_{q_t}^{x_3}.
$$

Hence desired conclusion holds.

Next we show [\(ii\).](#page-18-1) Assume  $n \equiv 0 \pmod{3}$ . Then we may assume  $p_1 = 3$  and  $f_1 \geq 1$ . We have

$$
\zeta_{3f_1}^{\Psi_n^{(3)}(z^{-1})} = \zeta_3^{\pi_n^{(3)}(z^{-1})} \zeta_{3f_1}^{\theta_n^{(3)}(z^{-1})} = \zeta_3^{-\pi_n^{(3)}(z)} \zeta_{3f_1}^{-\theta_n^{(3)}(z)} = \begin{cases} \zeta_3^{3-\pi_n^{(3)}(z)} & \text{if } \theta_m^{(3)}(z) = 0, \\ \zeta_3^{2-\pi_n^{(3)}(z)} \zeta_{3f_1}^{3f_1-1} - \theta_n^{(3)}(z) & \text{otherwise.} \end{cases}
$$

Hence the desired result holds.

#### 9.5 Proof of Theorem [1.3](#page-3-0)

By using Theorem [9.9,](#page-16-2) we can change integral bases of cyclotomic fields. First, we provide one lemma as a situation where Theorem [9.9](#page-16-2) can be easily applied.

<span id="page-19-2"></span>**Lemma 9.15.** Let  $m \geq 2$  be an integer with  $m \not\equiv 2 \pmod{4}$ , and let  $S \subset \{1, 2, \ldots, m-1\}$ . If  $|S| \leq \varphi(p^*)$  *for every prime factor*  $p \neq 2$  *of* m, then there exists an integral basis for  $\mathbb{Q}(\zeta_m)$  *containing*  $\epsilon$ *either*  $\zeta_m^r$  *or*  $-\zeta_m^r$  *for each*  $r \in S$ *. In particular, if all*  $\pm \zeta_m^r$  ( $r \in S$ ) *are pairwise-distinct, then* 

<span id="page-19-3"></span><span id="page-19-1"></span>
$$
\frac{1}{2} \sum_{r \in S} \pm \zeta_m^r \tag{9.8}
$$

*is not an algebraic integer.*

**Proof.** Let  $m = 2^{f_0} p_1^{f_1} p_2^{f_2} \cdots p_t^{f_t}$  be the prime factorization of m where  $f_0$  is a non-negative integer not equal to 1 and  $f_1, \ldots, f_t$  are positive integers.

Assume  $|S| \leq \varphi(p^*)$  for every prime factor  $p \neq 2$ . We give an integral basis of  $\mathbb{Q}(\zeta_m)$  containing either  $\zeta_m^r$  or  $-\zeta_m^r$ . For every  $j \geq 1$ , we let  $B_{p_j} = [p_j^{f_j-1}]$ , and can take  $A_{p_j}$  by the assumption such that

$$
\{\pi_m^{(p_j)}(\zeta_m^r) \mid r \in S\} \subset A_{p_j} \subset [p_j] \quad \text{and} \quad |A_{p_j}| = \varphi(p_j^*). \tag{9.9}
$$

In addition, if  $m \equiv 0 \pmod{4}$ , then we let

$$
B_2 = [2^{f_0}/2^*] = [2^{f_0 - 2}] \text{ and } A_2 = \{0, 1\}. \tag{9.10}
$$

By Theorem [9.9,](#page-16-2) we derive an integral basis I in [\(9.7\)](#page-16-3) of  $\mathbb{Q}(\zeta_m)$ .

Next, we show that for each  $r \in S$ , either  $\zeta_m^r$  or  $-\zeta_m^r$  is contained in *I*. Since

<span id="page-19-4"></span>
$$
\zeta_m^r = \prod_{j=0}^t \zeta_{p_j^*}^{\pi_m^{(p_j)}(\zeta_m^r)} \zeta_{q_j}^{\theta_m^{(p_j)}(\zeta_m^r)},
$$

where  $p_0 = 2$ , it suffices to verify that there exists  $\varepsilon \in {\pm 1}$  such that for every  $j \in [t]$ ,  $\pi_m^{(p_j)}(\varepsilon \zeta_m^r) \in A_{p_j}$ and  $\theta_m^{(p_j)}(\varepsilon \zeta_m^r) \in B_{p_j}$ .

In the case of  $f_0 = 0$ , we see that all  $\zeta_m^r$  is contained in *I*. Hence we assume  $f_0 \geq 2$ . If  $\pi_m^{(2)}(\zeta_m^r) \in$  $\{0,1\}$  for some  $r \in S$ , then  $\zeta_m^r \in I$ . Otherwise, we have by Lemma [9.14,](#page-18-2)  $\pi_m^{(2)}(-\zeta_m^r) \equiv \pi_m^{(2)}(\zeta_m^r) + 2 \equiv 0,1$ (mod 4) and  $\pi_m^{(p_j)}(-\zeta_m^r) \in A_{p_j}$  for every  $j \geq 1$ . These imply  $-\zeta_m^r \in I$ . Thus, I is a desired integral basis.

Lastly, if all  $\pm \zeta_m^r$   $(r \in S)$  are pairwise-distinct, then  $\zeta_m^r$   $(r \in S)$  are Q-linearly independent. Thus, [\(9.8\)](#page-19-1) is not an algebraic integer.

More precise discussion shows that  $\Delta$  in [\(9.5\)](#page-15-3), which is our interest, is not an algebraic integer.

<span id="page-19-0"></span>**Lemma 9.16.** *Let*  $m \geq 2$  *be an integer with*  $m \not\equiv 2 \pmod{4}$ *, and let* p, q *be positive integers.* If  $\zeta_m^p, \zeta_m^{-p}, \zeta_m^q, \zeta_m^{-q}$  are Q-linearly independent, then

$$
\Delta = \frac{\pm \zeta_m^p \pm \zeta_m^{-p} \pm \zeta_m^q \pm \zeta_m^{-q}}{2}
$$

*is not an algebraic integer.*

**Proof.** Let  $S = {\pm p, \pm q}$ . Let  $m = 2^{f_0} 3^{f_1} p_2^{f_2} p_3^{f_3} \cdots p_t^{f_t}$  be the prime factorization of m where  $f_0$  and  $f_1$  are non-negative integers and  $f_2, \ldots, f_t$  are positive integers.

First, since  $2 = \varphi(3) < 4 \leq \varphi(p)$  for any prime number  $p > 3$ , we consider the case where m is not divisible by 3 or where  $|\{\pi_m^{(3)}(\zeta_m^r) \mid r \in S\}| \leq \varphi(3)$ . In this case, Lemma [9.15](#page-19-2) is applicable, and hence we see that  $\Delta$  is not an algebraic integer. Below, we consider the case where m is divisible by 3, i.e.,  $f_2 \geq 1$ , and  $|\{\pi_m^{(3)}(\zeta_m^r) \mid r \in S\}| = 3$ . Let  $p_1 := 3$ . Let  $B_3 := [3^{f_1-1}]$  and  $A_3 := \{1, 2\}$ . For every  $j \geq 2$ , we define  $B_{p_j} = [p_j^{f_j-1}]$ , and  $A_{p_j}$  as in [\(9.9\)](#page-19-3). In addition, if  $m \equiv 0 \pmod{4}$ , then we define  $A_2$  and  $B_2$ by [\(9.10\)](#page-19-4). By Theorem [9.9,](#page-16-2) we derive an integral basis I in [\(9.7\)](#page-16-3) of  $\mathbb{Q}(\zeta_m)$ .

Since  $\{\pi_m^{(3)}(\zeta_m^n)\mid r \in S\} = \{0,1,2\}$ , there exists  $r' \in S$  such that  $\pi_m^{(3)}(\zeta_m^{r'}) = 0$ . We may assume that  $r' = p$ , i.e.,  $\pi_m^{(3)}(\zeta_m^p) = 0$  without loss of generality. We have  $\zeta_m^p = -\zeta_3^1 \zeta_m^p - \zeta_3^2 \zeta_m^p$ ,  $\pi_m^{(3)}(\zeta_3^1 \zeta_m^p) = 1$ and  $\pi_m^{(3)}(\zeta_3^2 \zeta_m^p) = 2$ . Also, we have by Lemma [9.14,](#page-18-2)

<span id="page-20-0"></span>
$$
\pi_m^{(3)}(\zeta_m^{-p}) \in \{0, 2\}.
$$
\n(9.11)

<span id="page-20-1"></span> $\Box$ 

Hence, either  $\pi_m^{(3)}(\zeta_m^q)$  or  $\pi_m^{(3)}(\zeta_m^{-q})$  must be 1. By Lemma [9.14,](#page-18-2) we may assume that

$$
\pi_m^{(3)}(\zeta_m^q) = 1
$$
 and  $\pi_m^{(3)}(\zeta_m^{-q}) \in \{1,2\}$ 

without loss of generality.

Assume  $\pi^{(3)}(\zeta_m^{-p}) = 2$ . Take an arbitrary element  $z \in {\zeta_3^1\zeta_m^p, \zeta_3^2\zeta_m^p, \zeta_m^{-p}, \zeta_m^{\pm q}}$ . Then, we have by Lemma [9.14,](#page-18-2)

$$
\pi_m^{(p_i)}(z) = \pi_m^{(p_i)}(-z) \in A_{p_i} \quad \text{and} \quad \theta_m^{(p_i)}(z) = \theta_m^{(p_i)}(-z) \in B_{p_i}
$$
\n(9.12)

for every  $1 \leq i \leq t$ . Furthermore, we have by Lemma [9.14](#page-18-2) again,

$$
\pi_m^{(2)}(z) \text{ or } \pi_m^{(2)}(-z) \in A_2 \quad \text{ and } \quad \theta_m^{(2)}(z), \theta_m^{(2)}(-z) \in B_2 \tag{9.13}
$$

if  $f_0 \geq 2$ . Hence,  $\zeta_3^1 \zeta_m^p, \zeta_3^2 \zeta_m^p, \zeta_m^{-p}, \zeta_m^{\pm q}$  are contained in I, up to multiplication by  $\pm 1$ . Since Q-linear independence of  $\zeta_m^{-p}, \zeta_m^{\pm q}$  implies that  $\zeta_m^{-p}, \zeta_m^{\pm q}$  are pairwise distinct up to multiplication by  $\pm 1$ , we see

$$
\Delta = \frac{\pm \zeta_m^p \pm \zeta_m^{-p} \pm \zeta_m^q \pm \zeta_m^{-q}}{2} = \frac{\mp \zeta_3^1 \zeta_m^p \mp \zeta_3^2 \zeta_m^p \pm \zeta_m^{-p} \pm \zeta_m^q \pm \zeta_m^{-q}}{2}
$$

is not an algebraic integer.

Assume  $\pi^{(3)}(\zeta_m^{-p}) = 0$ . Then,

$$
\zeta_m^{-p} = -\zeta_3^1 \zeta_m^{-p} - \zeta_3^2 \zeta_m^{-p}, \quad \pi_m^{(3)}(\zeta_3^1 \zeta_m^{-p}) = 1 \quad \text{and} \quad \pi_m^{(3)}(\zeta_3^2 \zeta_m^{-p}) = 2.
$$

We take an arbitrary element  $z \in \{\zeta_3^1\zeta_m^{\pm p}, \zeta_3^2\zeta_m^{\pm p}, \zeta_m^{\pm q}\}\.$  Then, by Lemma [9.14,](#page-18-2) we have [\(9.12\)](#page-20-0) for every  $1 \leq i \leq t$ , and obtain  $(9.13)$  if  $f_0 \geq 2$ . Thus,  $\zeta_3^1 \zeta_2^{\pm p}$ ,  $\zeta_3^2 \zeta_2^{\pm p}$ ,  $\zeta_m^{\pm q}$  are contained in I, up to multiplying by  $\pm 1$ . Also,  $\pm \zeta_3^1 \zeta_m^{\pm p}$ ,  $\pm \zeta_3^2 \zeta_m^{\pm p}$  are pairwise distinct by the assumption that  $\zeta_m^{\pm p}$  are Q-linearly independent. Therefore,

$$
\Delta = \frac{\pm \zeta_m^p \pm \zeta_m^{-p} \pm \zeta_m^q \pm \zeta_m^{-q}}{2} = \frac{\mp \zeta_3^1 \zeta_m^p \mp \zeta_3^2 \zeta_m^p \mp \zeta_3^1 \zeta_m^{-p} \mp \zeta_3^2 \zeta_m^{-p} \pm \zeta_m^q \pm \zeta_m^{-q}}{2}
$$

is not an algebraic integer.

We are now almost ready to prove our second main result (Theorem [1.3\)](#page-3-0). We show the case where  $l$  is odd first, and later we deal with the case where  $l$  is even.

**Theorem 9.17.** Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  be a connected 4-regular circulant graph, and let  $a + b \neq l$ . *If* l *is odd, then* X *does not admit perfect state transfer between vertex type states.*

**Proof.** Let l be an odd integer. By Lemma [9.3,](#page-14-2) it suffices to prove

$$
\Delta := \frac{(-1)^{a} \zeta_l^{\varepsilon(a)} + (-1)^{a} \zeta_l^{-\varepsilon(a)} + (-1)^{b} \zeta_l^{\varepsilon(b)} + (-1)^{b} \zeta_l^{-\varepsilon(b)}}{2}
$$

is not an algebraic integer. To confirm that the assumptions of Lemma [9.16](#page-19-0) are satisfied, we want to show that  $4\varepsilon(a)$ ,  $4\varepsilon(b)$ ,  $2(\varepsilon(a)\pm\varepsilon(b))\neq 0 \pmod{l}$  as these will be needed in order to apply Theorem [9.8.](#page-16-4) This can be proved by dividing the case by the parity of  $a$  and  $b$ . Since tedious calculations follow, we show only the case  $(\varepsilon(a), \varepsilon(b)) = (\frac{a}{2}, \frac{b+l}{2})$  as a representative. Suppose  $4\varepsilon(a) \equiv 0 \pmod{l}$ . We have  $2a \equiv 0 \pmod{l}$ . Since l is odd,  $a \equiv 0 \pmod{l}$ , which contradicts [\(9.1\)](#page-14-3). Suppose  $4\varepsilon(b) \equiv 0 \pmod{l}$ . We have  $0 \equiv 2(b + l) \equiv 2b \pmod{l}$ . Since l is odd,  $b \equiv 0 \pmod{l}$ , which contradicts [\(9.1\)](#page-14-3). Suppose  $2(\varepsilon(a) + \varepsilon(b)) \equiv 0 \pmod{l}$ . We have  $0 \equiv a + b + l \equiv a + b \pmod{l}$ , so  $a + b \equiv 0, l \pmod{2l}$ . Both contradict [\(9.2\)](#page-14-4) or our assumption. Suppose  $2(\varepsilon(a)-\varepsilon(b))\equiv 0\pmod{l}$ . We have  $0\equiv a-b-l\equiv a-b$ (mod l), so  $a - b \equiv 0, l \pmod{2l}$ . Both contradict [\(9.2\)](#page-14-4) or [\(9.3\)](#page-14-5). Therefore by Lemma [9.8](#page-16-4) and Lemma 9.16, we see that  $\Delta$  is not an algebraic integer. Lemma [9.16,](#page-19-0) we see that  $\Delta$  is not an algebraic integer.

The reason why the above proof worked is that the assumption that  $l$  is odd guaranteed  $\mathbb Q$ -linear independence of  $\zeta_l^{\varepsilon(r)}$  $\ell_l^{(r)}$   $(r \in S = {\pm a, \pm b})$ . However, when l is even, for example  $l = 6$  and  $(a, b) =$ (1, 3), then it holds that  $\zeta_{2l}^{-b} = \zeta_4^3 = -\zeta_4^1 = -\zeta_{2l}^b$ . This means that the case where l is even does not necessarily guarantee  $\overline{Q}$ -linear independence of  $\zeta_{2l}^r$  ( $r \in S$ ). Therefore, more devised discussion is required when  $l$  is even.

<span id="page-21-1"></span>**Lemma 9.18.** Let  $X = X(\mathbb{Z}_2, \{\pm a, \pm b\})$  be a connected 4-regular graph. If  $4a \equiv 0 \pmod{2l}$ , then  $4b \not\equiv 0 \pmod{2l}$ . Moreover,  $\zeta_{2l}^b$  and  $\zeta_{2l}^{-b}$  are Q-linearly independent.

**Proof.** Our assumption derives that  $2a \equiv 0 \pmod{l}$ , so  $2a \equiv 0, l \pmod{2l}$ . By [\(9.1\)](#page-14-3), we have  $2a \equiv l$ (mod 2l). Next, we suppose that  $4b \equiv 0 \pmod{2l}$ . Similarly, we obtain  $2b \equiv l \pmod{2l}$ . This implies that  $a - b \equiv 0, l \pmod{2l}$ . However, this contradicts [\(9.2\)](#page-14-4) or [\(9.3\)](#page-14-5).

Next, we assume that a Q-linear combination

<span id="page-21-0"></span>
$$
c_1 \zeta_{2l}^b + c_2 \zeta_{2l}^{-b} = 0. \tag{9.14}
$$

Take the conjugate of both sides and add it to Equality [\(9.14\)](#page-21-0). We have  $2(c_1 + c_2) \cos \frac{2b}{2l} \pi = 0$ , and hence  $c_1 + c_2 = 0$  since  $4b \neq 0 \pmod{2l}$ . Substituting it into Equality [\(9.14\)](#page-21-0). Then we have  $2ic_1 \sin \frac{2b}{2l}\pi = 0$ , and hence  $c_1 = 0$  since  $4b \not\equiv 0 \pmod{2l}$ . Therefore,  $\zeta_{2l}^b$  and  $\zeta_{2l}^{-b}$  are Q-linearly independent. П

**Theorem 9.19.** Let  $X = X(\mathbb{Z}_{2l}, \{\pm a, \pm b\})$  be a connected 4-regular with  $a + b \neq l$ . If l is even, then X *does not admit perfect state transfer between vertex type states.*

**Proof.** Let  $\Delta$  be the element of  $\mathbb{Q}(\zeta_{2l})$  in [\(9.4\)](#page-14-1). By Lemma [9.2,](#page-14-6) it suffices to check that  $\Delta$  is not an algebraic integer. First, we consider the case where  $4a, 4b \neq 0 \pmod{2l}$ . Then we claim that  $2(a \pm b) \not\equiv 0 \pmod{2l}$ . Indeed, we suppose that  $2(a \pm b) \equiv 0 \pmod{2l}$ . Then we have  $a \pm b \equiv 0, l$  $\pmod{2l}$ , which contradicts  $(9.2)$ ,  $(9.3)$ , or our assumption. Therefore by Lemma [9.8](#page-16-4) and Lemma [9.16,](#page-19-0) we see that  $\Delta$  is not an algebraic integer. Hence Lemma [9.1](#page-14-0) implies that the graph X does not admit perfect state transfer.

Next, we consider the case where either  $4a \equiv 0 \pmod{2l}$  or  $4b \equiv 0 \pmod{2l}$ . We may assume  $4a \equiv 0 \pmod{2l}$  without loss of generality. By Lemma [9.18,](#page-21-1) we have  $4b \not\equiv 0 \pmod{2l}$ . Since  $4a \equiv 0$ (mod 2l) and [\(9.1\)](#page-14-3), we have  $2a \equiv l \pmod{2l}$ . Thus, there exists an integer s such that  $2a - l = 2sl$ , i.e.,  $2a = (2s+1)l$ . We have  $\zeta_{2l}^{\pm a} = \zeta_{4l}^{\pm 2a} = \zeta_{4l}^{\pm (2s+1)l} = \zeta_4^{\pm (2s+1)}$ . In particular,  $\zeta_{2l}^a + \zeta_{2l}^{-a} = 0$ . Hence,

$$
\Delta = \frac{\zeta_{2l}^a + \zeta_{2l}^{-a} + \zeta_{2l}^b + \zeta_{2l}^{-b}}{2} = \frac{\zeta_{2l}^b + \zeta_{2l}^{-b}}{2}.
$$

By Lemma [9.18,](#page-21-1)  $\zeta_{2l}^b$  and  $\zeta_{2l}^{-b}$  are Q-linearly independent, so Lemma [9.15](#page-19-2) derives that  $\Delta$  is not an algebraic integer. Therefore, Lemma [9.1](#page-14-0) implies that the graph  $X$  does not admit perfect state transfer.  $\Box$ 

## Acknowledgements

We are deeply grateful to the anonymous reviewers for their insightful feedback and constructive suggestions, which have significantly contributed to improving the manuscript. We would also like to thank Akihiro Higashitani for his helpful advice and insightful discussions. S.K. is supported by JSPS KAKENHI (Grant Number JP20J01175). K.Y. is supported by JSPS KAKENHI (Grant Number JP21J14427).

## <span id="page-22-0"></span>References

- <span id="page-22-6"></span>[1] A. Ambainis. Quantum walk algorithm for element distinctness. *SIAM Journal on Computing*, 37(1):210–239, 2007.
- [2] K. Barr, T. Proctor, D. Allen, and V. Kendon. Periodicity and perfect state transfer in quantum walks on variants of cycles. *Quantum Information & Computation*, 14(5&6):417–438, 2014.
- <span id="page-22-16"></span><span id="page-22-14"></span>[3] A. Berger. On linear independence of trigonometric numbers. *Carpathian Journal of Mathematics*, 34(2):157–166, 2018.
- [4] W. Bosma. Canonical bases for cyclotomic fields. *Applicable Algebra in Engineering, Communication and Computing*, 1(2):125–134, 1990.
- <span id="page-22-13"></span><span id="page-22-11"></span>[5] A.E. Brouwer and W.H. Haemers. *Spectra of graphs*. Universitext. Springer, New York, 2012.
- [6] A. Chan and H. Zhan. Pretty good state transfer in discrete-time quantum walks. *Journal of Physics A: Mathematical and Theoretical*, 56(16):165305, mar 2023.
- <span id="page-22-8"></span>[7] Q. Chen, C. Godsil, M. Sobchuk, and H. Zhan. Hamiltonians of Bipartite Walks. *Electron. J. Combin.*, 31(4):Paper No. 4.10, 2024.
- <span id="page-22-3"></span>[8] M. Christandl, N. Datta, A. Ekert, and A. J. Landahl. Perfect state transfer in quantum spin networks. *Physical review letters*, 92(18):187902, 2004.
- <span id="page-22-10"></span><span id="page-22-4"></span>[9] C. Godsil. State transfer on graphs. *Discrete Mathematics*, 312(1):129–147, 2012.
- <span id="page-22-9"></span>[10] C. Godsil. When can perfect state transfer occur? *Electron. J. Linear Algebra*, 23:877–890, 2012.
- [11] C. Godsil and G. Royle. *Algebraic graph theory*, volume 207 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001.
- <span id="page-22-12"></span>[12] K. Guo and V. Schmeits. Perfect state transfer in quantum walks on orientable maps. *Algebr. Comb.*, 7(3):713–747, 2024.
- <span id="page-22-2"></span>[13] Y. Higuchi, N. Konno, I. Sato, and E. Segawa. Periodicity of the discrete-time quantum walk on a finite graph. *Interdisciplinary Information Sciences*, 23(1):75–86, 2017.
- <span id="page-22-1"></span>[14] N. Inui, Y. Konishi, and N. Konno. Localization of two-dimensional quantum walks. *Physical Review A*, 69(5):052323, 2004.
- <span id="page-22-15"></span><span id="page-22-5"></span>[15] F. Jarvis. *Algebraic number theory*, volume 1303. Springer, 2014.
- [16] V. Kendon and C. Tamon. Perfect state transfer in quantum walks on graphs. *Journal of Computational and Theoretical Nanoscience*, 8(3):422–433, 2011.
- <span id="page-22-7"></span>[17] S. Kubota and E. Segawa. Perfect state transfer in grover walks between states associated to vertices of a graph. *Linear Algebra and its Applications*, 646:238–251, 2022.
- <span id="page-23-4"></span>[18] S. Kubota, E. Segawa, and T. Taniguchi. Quantum walks defined by digraphs and generalized hermitian adjacency matrices. *Quantum Information Processing*, 20(3):1–30, 2021.
- <span id="page-23-5"></span><span id="page-23-3"></span>[19] S. Kubota, H. Sekido, and H. Yata. Periodicity of quantum walks defined by mixed paths and mixed cycles. *Linear Algebra and its Applications*, 630:15–38, 2021.
- <span id="page-23-0"></span>[20] C. D. Meyer. *Matrix analysis and applied linear algebra*, volume 71. Siam, 2000.
- [21] F. D. Nicola, L. Sansoni, A. Crespi, R. Ramponi, R. Osellame, V. Giovannetti, R. Fazio, P. Mataloni, and F. Sciarrino. Quantum simulation of bosonic-fermionic noninteracting particles in disordered systems via a quantum walk. *Physical Review A*, 89(3):032322, 2014.
- <span id="page-23-1"></span>[22] C. Vlachou, J. Rodrigues, P. Mateus, N. Paunković, and A. Souto. Quantum walk public-key cryptographic system. *International Journal of Quantum Information*, 13(07):1550050, 2015.
- <span id="page-23-2"></span>[23] H. Zhan. An infinite family of circulant graphs with perfect state transfer in discrete quantum walks. *Quantum Information Processing*, 18(12):1–26, 2019.