

FUNCTIONAL DIFFERENTIAL EQUATIONS DRIVEN BY CÀDLÀG ROUGH PATHS

ANNA P. KWOSSEK, ANDREAS NEUENKIRCH, AND DAVID J. PRÖMEL

ABSTRACT. The existence of unique solutions is established for rough differential equations (RDEs) with path-dependent coefficients and driven by càdlàg rough paths. Moreover, it is shown that the associated solution map, also known as Itô–Lyons map, is locally Lipschitz continuous. These results are then applied to various classes of rough differential equations, such as controlled RDEs and RDEs with delay, as well as to stochastic differential equations with delay. To that end, a joint rough path is constructed for a càdlàg martingale and its delayed version, that corresponds to stochastic Itô integration.

Key words: Itô–Lyons map, martingale, rough differential equation, delayed rough path, stochastic functional differential equation, stochastic differential equation with delay.

MSC 2020 Classification: 60L20, 60H10.

1. INTRODUCTION

Stochastic functional differential equations, also known as stochastic delay differential equations, are a natural generalization of stochastic ordinary differential equations, allowing for path-dependent coefficients which may depend on past values of the generated random dynamics. Since numerous real-world phenomena show evidence of a dependence on the past as well as a stochastic behaviour, stochastic functional differential equations serve as mathematical models in many areas ranging from biology to finance. For classical introductions to stochastic functional differential equations we refer, e.g., to [Moh84, Moh98].

A deterministic approach to stochastic differential equations is provided by rough path theory, initiated by Lyons [Lyo98]. Originally designed to treat stochastic ordinary differential equations, it has been extended in various directions, for instance, allowing to deal with stochastic Volterra equations [DT09], reflected stochastic differential equations [Aid15], stochastic inclusion equations [BBC21], and different classes of stochastic partial differential equations [Hai11, CFO11]. These rough path approaches contributed many novel insights to the study of the aforementioned equations, such as, but not limited to, new well-posedness and stability results. Comprehensive introductions to rough path theory can be found, e.g., in [LCL07, FH20].

In the present paper, we study *rough functional differential equations* (RFDEs)

$$(1.1) \quad Y_t = y_t + \int_0^t b_s(Y) ds + \int_0^t \sigma_s(Y) d\mathbf{X}_s, \quad t \in [0, T],$$

where the driving signal \mathbf{X} is a càdlàg p -rough path for $p \in (2, 3)$, the initial condition y is a given controlled path, and the coefficients b, σ are non-anticipative functionals, mapping a controlled path to a controlled path. Assuming a quadratic growth and a Lipschitz type condition on the path-dependent coefficients b, σ , which both are formulated on the space

of controlled paths, we establish the existence of a unique solution to the RFDE (2.2). To that end, we rely on the theory of càdlàg rough paths, as introduced by Friz, Shekhar and Zhang [FS17, FZ18], as well as Banach’s fixed point theorem. Moreover, we show that the solution map, also known as Itô–Lyons map, mapping the input (initial condition, coefficients, driving signal) of an RFDE to its solution, is locally Lipschitz continuous with respect to suitable distances on the associated spaces of coefficients, controlled paths and rough paths. Let us remark that the continuity of the Itô–Lyons map is one of the most fundamental insights of rough path theory, with many applications to stochastic differential equations, cf. e.g. [FH20].

The presented results on rough functional differential equations provide a unifying theory, recovering and extending various previous results on different classes of rough differential equations with path-dependent coefficients. Indeed, we deduce the existence of unique solutions as well as the local Lipschitz continuity of the Itô–Lyons map for classical rough differential equations (RDEs), controlled RDEs, RDEs with discrete time dependence and RDEs with constant/variable delay, that are all driven by càdlàg p -rough paths for $p \in (2, 3)$.

In the existing literature, there are several different approaches to deal with rough functional differential equations driven by continuous rough paths. Since the theory of (continuous) rough paths works nicely for infinite dimensional Banach spaces, RDEs with path-dependent coefficients can be treated as Banach space-valued RDEs, see e.g. [Bai14], which requires the coefficients to be Fréchet differentiable and, thus, excludes some interesting examples. Existence, uniqueness and stability results are established by Neuenkirch, Nourdin and Tindel [NNT08] for RDEs with constant delay. The existence of a solution is proven by Ananova [Ana23] for RDEs with path-dependent coefficients, which are assumed to be Dupire differentiable [Dup19], and by Aida [Aid24] for RDEs with coefficients containing path-dependent bounded variation. The latter two approaches rely on Schauder’s fixed point theorem. Another exemplary class of RFDEs are reflected rough differential equations, see e.g. [Aid15, DGHT19], which, in general, do not possess a unique solution, see [Gas21].

Most applications of rough path theory to stochastic differential equations (SDEs) crucially rely on the construction of suitable (random) rough paths. To apply the developed theory on RFDEs to Itô SDEs with constant delays, we show that a càdlàg martingale together with its delayed version can be lifted to a random rough path in the spirit of stochastic Itô integration. The key challenge to obtain the ‘delayed’ rough path is that a martingale together with its delayed version is, in general, not a martingale itself, thus preventing the direct use of stochastic Itô integration. For related constructions of random rough paths above fractional Brownian motions we refer to [NNT08, TT12, BMCR14, CE20]. Consequently, one can apply the continuity of the Itô–Lyons map to derive pathwise stability results for stochastic differential equations with constant delay, which plays an important role in many applications, see e.g. [BDNHP18]. In particular, the map $y \mapsto Y$, mapping the initial condition y to the associated solution Y of an SDE with constant delay, is continuous on the space of controlled paths. This resolves an old observation, pointed out by Mohammed [Moh86], about the non-continuity of the flow of stochastic differential equations with delay, for which the initial condition is in fact an initial path.

The organization of the paper is as follows. In Section 2 we provide existence, uniqueness and continuity results for rough functional differential equations. In Section 3 we prove that various classes of rough differential equations are covered by the presented results on rough functional differential equations. In Section 4 we establish the existence of the Itô rough path

lift of delayed martingales and discuss applications to stochastic differential equations with delay. Appendix A contains some auxiliary estimates for rough integrals.

2. EXISTENCE, UNIQUENESS AND CONTINUITY

Before treating rough functional differential equations (RFDEs), we recall the necessary definitions and some essentials from the theory of càdlàg rough paths, as introduced by Friz and Shekhar [FS17] and Friz and Zhang [FZ18]. The theory of càdlàg rough paths extends the classical rough path theory, allowing to deal with many stochastic processes with jumps [Che18], and has numerous applications, e.g., in probability theory [FZK23], numerical analysis [FZ18] and mathematical finance [ALP24].

2.1. Essentials on rough path theory. Throughout, let $T > 0$ be a fixed finite time horizon. Let $\Delta_T := \{(s, t) \in [0, T]^2 : s \leq t\}$ be the standard 2-simplex. A function $w: \Delta_T \rightarrow [0, \infty)$ is called a *control function* if it is superadditive, in the sense that $w(s, u) + w(u, t) \leq w(s, t)$ for all $0 \leq s \leq u \leq t \leq T$. We write $w(s, t-) := \lim_{u \uparrow t} w(s, u)$ if $s < t$, and $w(s, t-) := 0$ if $s = t$.

Whenever $(B, \|\cdot\|)$ is a normed space and $f, g: B \rightarrow \mathbb{R}$ are two functions on B , we shall write $f \lesssim g$ or $f \leq Cg$ to mean that there exists a constant $C > 0$ such that $f(x) \leq Cg(x)$ for all $x \in B$. The constant C may depend on the normed space, e.g. through its dimension or regularity parameters, and, if we want to emphasize the dependence of the constant C on some particular variables, say $\alpha_1, \dots, \alpha_n$, then we will write $\lesssim_{\alpha_1, \dots, \alpha_n}$ or $C = C_{\alpha_1, \dots, \alpha_n}$. Unless otherwise stated, the dependence of the implicit constants on the variables is locally bounded; that is, if $\alpha_1 \in A_1, \dots, \alpha_n \in A_n$, where A_1, \dots, A_n are compact subsets of the range of $\alpha_1, \dots, \alpha_n$ respectively, then we have that $\sup_{\alpha_1 \in A_1, \dots, \alpha_n \in A_n} C_{\alpha_1, \dots, \alpha_n} < \infty$.

The space of linear maps from $\mathbb{R}^d \rightarrow \mathbb{R}^n$ is denoted by $\mathcal{L}(\mathbb{R}^d; \mathbb{R}^n)$ and we write $C_b^l = C_b^l(\mathbb{R}^n; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^n))$ for the space of l -times differentiable (in the Fréchet sense) functions $f: \mathbb{R}^n \rightarrow \mathcal{L}(\mathbb{R}^d; \mathbb{R}^n)$ such that f and all its derivatives up to order l are continuous and bounded. We equip this space with the norm

$$\|f\|_{C_b^l} := \|f\|_\infty + \|Df\|_\infty + \dots + \|D^l f\|_\infty,$$

where $\|\cdot\|_\infty$ denotes the supremum norm on the corresponding spaces of operators.

For a normed space $(E, |\cdot|)$, let $D([0, T]; E)$ be the set of càdlàg (right-continuous with left-limits) paths from $[0, T]$ to E . For $p \geq 1$, the p -variation of a path $X \in D([0, T]; E)$ is given by

$$\|X\|_p := \|X\|_{p, [0, T]} \quad \text{with} \quad \|X\|_{p, [s, t]} := \left(\sup_{\mathcal{P} \subset [s, t]} \sum_{[u, v] \in \mathcal{P}} |X_v - X_u|^p \right)^{\frac{1}{p}}, \quad (s, t) \in \Delta_T,$$

where the supremum is taken over all possible partitions \mathcal{P} of the interval $[s, t]$. We recall that, given a path X , we have that $\|X\|_p < \infty$ if and only if there exists a control function w such that¹

$$\sup_{(u, v) \in \Delta_T} \frac{|X_v - X_u|^p}{w(u, v)} < \infty.$$

¹Here and throughout, we adopt the convention that $\frac{0}{0} := 0$.

We write $D^p = D^p([0, T]; E)$ for the space of paths $X \in D([0, T]; E)$ which satisfy $\|X\|_p < \infty$. Moreover, for a path $X \in D([0, T]; \mathbb{R}^d)$, we will often use the shorthand notation:

$$X_{s,t} := X_t - X_s, \quad \text{for } (s, t) \in \Delta_T.$$

For $p > 2$ and a two-parameter function $\mathbb{X}: \Delta_T \rightarrow E$, we similarly define

$$\|\mathbb{X}\|_{\frac{p}{2}} := \|\mathbb{X}\|_{\frac{p}{2}, [0, T]} \quad \text{with} \quad \|\mathbb{X}\|_{\frac{p}{2}, [s, t]} := \left(\sup_{\mathcal{P} \subset [s, t]} \sum_{[u, v] \in \mathcal{P}} |\mathbb{X}_{u, v}|^{\frac{p}{2}} \right)^{\frac{2}{p}}, \quad (s, t) \in \Delta_T.$$

We write $D_2^{\frac{p}{2}} = D_2^{\frac{p}{2}}(\Delta_T; E)$ for the space of all functions $\mathbb{X}: \Delta_T \rightarrow E$ which satisfy $\|\mathbb{X}\|_{\frac{p}{2}} < \infty$, and are such that the maps $s \mapsto \mathbb{X}_{s, t}$ for fixed t , and $t \mapsto \mathbb{X}_{s, t}$ for fixed s , are both càdlàg.

For $p \in (2, 3)$, a pair $\mathbf{X} = (X, \mathbb{X})$ is called a *càdlàg p -rough path* over \mathbb{R}^d if

- (i) $X \in D^p([0, T]; \mathbb{R}^d)$ and $\mathbb{X} \in D_2^{\frac{p}{2}}(\Delta_T; \mathbb{R}^{d \times d})$, and
- (ii) Chen's relation: $\mathbb{X}_{s, t} = \mathbb{X}_{s, u} + \mathbb{X}_{u, t} + X_{s, u} \otimes X_{u, t}$ holds for all $0 \leq s \leq u \leq t \leq T$,

where \otimes denotes the usual tensor product. In component form then, condition (ii) states that $\mathbb{X}_{s, t}^{ij} = \mathbb{X}_{s, u}^{ij} + \mathbb{X}_{u, t}^{ij} + X_{s, u}^i X_{u, t}^j$ for every i and j . We will denote the space of càdlàg p -rough paths by $\mathcal{D}^p = D^p([0, T]; \mathbb{R}^d)$. On the space $\mathcal{D}^p([0, T]; \mathbb{R}^d)$, we use the natural seminorm

$$\|\mathbf{X}\|_p := \|X\|_{p, [0, T]} \quad \text{with} \quad \|\mathbf{X}\|_{p, [s, t]} := \|X\|_{p, [s, t]} + \|\mathbb{X}\|_{\frac{p}{2}, [s, t]}$$

for $(s, t) \in \Delta_T$, and the induced distance

$$\|\mathbf{X}; \tilde{\mathbf{X}}\|_p := \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [0, T]} \quad \text{with} \quad \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [s, t]} := \|X - \tilde{X}\|_{p, [s, t]} + \|\mathbb{X} - \tilde{\mathbb{X}}\|_{\frac{p}{2}, [s, t]},$$

for $(s, t) \in \Delta_T$.

Let $p \in (2, 3)$, and $X \in D^p([0, T]; \mathbb{R}^d)$. We say that a pair (Y, Y') is a *controlled path* (with respect to X), if

$$Y \in D^p([0, T]; E), \quad Y' \in D^p([0, T]; \mathcal{L}(\mathbb{R}^d; E)), \quad \text{and} \quad R^Y \in D_2^{\frac{p}{2}}(\Delta_T; E),$$

where R^Y is defined by

$$Y_{s, t} = Y'_s X_{s, t} + R_{s, t}^Y \quad \text{for all } (s, t) \in \Delta_T.$$

We write $\mathcal{V}_X^p = \mathcal{V}_X^p([0, T]; E)$ for the space of E -valued controlled paths, which becomes a Banach space when equipped with the norm $(Y, Y') \mapsto |Y_0| + \|Y, Y'\|_{X, p}$, where

$$\|Y, Y'\|_{X, p} := \|Y, Y'\|_{X, p, [0, T]} \quad \text{with} \quad \|Y, Y'\|_{X, p, [s, t]} := |Y'_s| + \|Y'\|_{p, [s, t]} + \|R^Y\|_{\frac{p}{2}, [s, t]}$$

for $(s, t) \in \Delta_T$. We point out that, by definition, for $(s, t) \in \Delta_T$,

$$|Y_{s, t}| \leq |Y'_s| |X_{s, t}| + |R_{s, t}^Y| \quad \text{and} \quad |Y'_t| \leq |Y'_0| + |Y'_{0, t}|,$$

which implies that

$$\|Y\|_p \leq C_p (\|Y'\|_\infty \|X\|_p + \|R^Y\|_{\frac{p}{2}}) \quad \text{and} \quad \|Y'\|_\infty \leq |Y'_0| + \|Y'\|_p.$$

Given $X, \tilde{X} \in D^p$, we further introduce the standard ‘distance’

$$\|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p} := \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p, [0, T]}$$

with

$$\|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p, [s, t]} := |Y'_s - \tilde{Y}'_s| + \|Y' - \tilde{Y}'\|_{p, [s, t]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [s, t]}$$

for $(s, t) \in \Delta_T$, whenever $(Y, Y') \in \mathcal{V}_X^p, (\tilde{Y}, \tilde{Y}') \in \mathcal{V}_{\tilde{X}}^p$. Note that \mathcal{V}_X^p and $\mathcal{V}_{\tilde{X}}^p$ are, in general, different Banach spaces; if $X = \tilde{X}$, we write $\|\cdot; \cdot\|_{X,p,[s,t]}$.

Given $p \in (2, 3)$, $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ and $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, the (forward) rough integral

$$(2.1) \quad \int_s^t Y_r \, d\mathbf{X}_r := \lim_{|\mathcal{P}| \rightarrow 0} \sum_{[u,v] \in \mathcal{P}} (Y_u X_{u,v} + Y'_u \mathbb{X}_{u,v}), \quad (s, t) \in \Delta_T,$$

exists (in the classical mesh Riemann–Stieltjes sense), where the limit is taken along any sequence of partitions $(\mathcal{P}^n)_{n \in \mathbb{N}}$ of the interval $[s, t]$ such that $|\mathcal{P}^n| \rightarrow 0$ as $n \rightarrow \infty$. More precisely, in writing the product $Y_u X_{u,v}$, we apply the operator $Y_u \in \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)$ onto $X_{u,v} \in \mathbb{R}^d$; and in writing the product $Y'_u \mathbb{X}_{u,v}$, we use the natural identification of $\mathcal{L}(\mathbb{R}^d; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ with $\mathcal{L}(\mathbb{R}^d \otimes \mathbb{R}^d; \mathbb{R}^k)$. The rough integral comes with the estimate

$$\left| \int_s^t Y_r \, d\mathbf{X}_r - Y_s X_{s,t} - Y'_s \mathbb{X}_{s,t} \right| \leq C \left(\|R^Y\|_{\frac{p}{2}, [s,t]} \|X\|_{p, [s,t]} + \|Y'\|_{p, [s,t]} \|\mathbb{X}\|_{\frac{p}{2}, [s,t]} \right)$$

for some constant C depending only on p ; see [FZ18, Proposition 2.6], where

$$\|Y'\|_{p, [s,t]} := \sup_{u < t} \|Y'\|_{p, [s,u]} \quad \text{and} \quad \|R^Y\|_{\frac{p}{2}, [s,t]} := \sup_{u < t} \|R^Y\|_{\frac{p}{2}, [s,u]}.$$

The estimate implies that $(\int_0^\cdot Y_r \, d\mathbf{X}_r, Y) \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ is a controlled path with respect to X , see also [FZ18, Remark 2.8].

For details on the construction of the rough integral with respect to càdlàg p -rough paths and its properties, we refer to [FS17, FZ18], and we provide some auxiliary estimates for the rough integral in Appendix A.

Let us now consider the rough functional differential equation (RFDE)

$$(2.2) \quad Y_t = y_t + \int_0^t F_s(Y) \, d\mathbf{X}_s, \quad t \in [0, T],$$

where $\mathbf{X} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ is a càdlàg p -rough path for $p \in (2, 3)$, $(y, y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ is a given controlled path with respect to X and further, where $(F, F') : \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ is a non-anticipative functional, i.e.

- (i) $(F(Y), F'(Y, Y')) \in \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$,
- (ii) $(F_t(Y), F'_t(Y, Y')) = (F_t(Y_{\cdot \wedge t}), F'_t(Y_{\cdot \wedge t}, Y'_{\cdot \wedge t}))$ for all $t \in [0, T]$,

for every $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$. The integral in (2.2) is defined as a (forward) rough integral, see (2.1) for its definition. Note that the RFDE (1.1) can be re-written in the form of (2.2), using a standard time-extension of the driving rough path.

2.2. Existence and uniqueness. To prove the existence of a unique solution to the rough functional differential equation (2.2), we postulate a quadratic growth and a Lipschitz type condition on the path-dependent coefficient (F, F') , formulated on the associated path spaces. While a Lipschitz type condition is expected, the quadratic growth condition appears to be natural in the presented context of (second order) controlled paths, which corresponds to a Taylor expansion with quadratic remainder term.

Assumption 2.1. For any $X, \tilde{X} \in D^p([0, T]; \mathbb{R}^d)$ and for every $K > 0$, there exist constants $C_F > 0$, which depends on p and the functional F , and $C_{F, K, X, \tilde{X}} > 0$, which additionally depends on K, X, \tilde{X} such that the maps

$$\begin{aligned} (F, F') &: \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \\ (F, F') &: \mathcal{V}_{\tilde{X}}^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_{\tilde{X}}^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)) \end{aligned}$$

satisfy, for all $(Y, Y') \in \mathcal{V}_X^p$, $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_{\tilde{X}}^p$, and every $0 \leq s < t \leq T$,

(i) the growth conditions:

$$\begin{aligned} |F_t(Y)| &\leq C_F, \\ |F_{t-, t}(Y)| &\leq C_F(1 + \|Y\|_{p, [s, t]} + |Y_{t-, t}|), \\ \|F(Y)\|_{p, [s, t]} &\leq C_F(1 + (|Y'_s| + \|Y'\|_{p, [s, t]})\|X\|_{p, [s, t]} + \|R^Y\|_{\frac{p}{2}, [s, t]}), \quad \text{and} \\ \|F(Y), F'(Y, Y')\|_{X, p, [s, t]} &\leq C_F(1 + \|Y, Y'\|_{X, p, [s, t]})^2(1 + \|X\|_{p, [s, t]})^2; \end{aligned}$$

(ii) the Lipschitz conditions:

$$\begin{aligned} \|F(Y) - F(\tilde{Y})\|_{p, [s, t]} &\leq C_{F, K, X, \tilde{X}}(|Y_s - \tilde{Y}_s| + \|Y - \tilde{Y}\|_{p, [s, t]}), \quad \text{and} \\ \|F(Y), F'(Y, Y'); F(\tilde{Y}), F'(\tilde{Y}, \tilde{Y}')\|_{X, \tilde{X}, p, [s, t]} & \\ &\leq C_{F, K, X, \tilde{X}}(|Y_s - \tilde{Y}_s| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p, [s, t]} + \|X - \tilde{X}\|_{p, [s, t]}), \end{aligned}$$

if $\|Y, Y'\|_{X, p, [s, t]}, \|\tilde{Y}, \tilde{Y}'\|_{\tilde{X}, p, [s, t]} \leq K$.

Remark 2.2. When referring to a functional (F, F') which satisfies Assumption 2.1, we consider a functional which maps a controlled path with respect to some path $X \in D^p$ to a controlled path with respect to X , where X is not being fixed a priori. That is, $(F(Y), F'(Y, Y')) \in \mathcal{V}_X^p$ for $(Y, Y') \in \mathcal{V}_X^p$, for any $X \in D^p$. Particularly, Assumption 2.1 is to hold for any $X, \tilde{X} \in D^p$, which are not assumed fixed.

Remark 2.3. The growth and Lipschitz conditions in Assumption 2.1 are formulated in terms of both the p -variation of Y and the controlled path norm $(Y, Y') \mapsto |Y_0| + \|Y, Y'\|_{X, p}$ on the space \mathcal{V}_X^p of controlled paths (Y, Y') . To deduce the existence of a unique solution to the RFDE (2.2) under a growth and Lipschitz conditions formulated only in terms of the controlled path norm seems to be far from being obvious. Moreover, notice that the common examples of RDEs with path-dependent coefficients do satisfy Assumption 2.1, see Section 3 below, demonstrating that Assumption 2.1 is, indeed, a natural generalization of the conditions on the coefficients postulated in the existing literature.

Based on Assumption 2.1, we obtain the following global existence and uniqueness result for rough functional differential equations.

Theorem 2.4. Let $\mathbf{X} \in D^p([0, T]; \mathbb{R}^d)$ be a càdlàg p -rough path for $p \in (2, 3)$, and $(y, y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ be a given controlled path with respect to X . Suppose that the non-anticipative functional $(F, F') : \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ satisfies Assumption 2.1. Then, there exists a unique solution to the rough functional differential equation (2.2), i.e. there exists a unique controlled path $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, with $Y' = y' + F(Y)$, such that

$$Y_t = y_t + \int_0^t F_s(Y) d\mathbf{X}_s, \quad t \in [0, T].$$

Moreover, there exists a componentwise non-decreasing function $K_p: [0, \infty)^3 \rightarrow [0, \infty)$ such that

$$\|Y, Y'\|_{X,p} \leq K_p(\|y, y'\|_{X,p}, C_F, \|\mathbf{X}\|_p).$$

The proof relies on a fixed point approach using Banach's fixed point theorem.

Proof. Step 1: Local solution. We may assume that

$$\|\mathbf{X}\|_p \leq 1 \quad \text{and} \quad \|y'\|_p + \|R^y\|_{\frac{p}{2}} \leq 1.$$

For $t \in (0, T]$, we define the map $\mathcal{M}_t: \mathcal{V}_X^p([0, t]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, t]; \mathbb{R}^k)$ by

$$(Y, Y') \mapsto (Z, Z') := \mathcal{M}_t(Y, Y') := \left(y + \int_0^\cdot F_s(Y) d\mathbf{X}_s, y' + F(Y) \right),$$

noting that $\mathcal{M}_t(Y, Y')$ is a controlled path with respect to X as \mathcal{V}_X^p is a Banach space, and introduce the subset of controlled paths

$$\mathcal{B}_t := \left\{ (Y, Y') \in \mathcal{V}_X^p([0, t]; \mathbb{R}^k) : \begin{array}{l} (Y_0, Y'_0) = (y_0, y'_0 + F_0(y)), \\ \|(Y - y)'\|_{p,[0,t]} \leq 4C_F, \|R^Y - R^y\|_{\frac{p}{2},[0,t]} \leq 1 \end{array} \right\}.$$

Invariance. For any $(Y, Y') \in \mathcal{B}_t$, we have that

$$\begin{aligned} \|(Z - y)'\|_{p,[0,t]} &= \|F(Y)\|_{p,[0,t]} \\ &\leq C_F(1 + (|Y'_0| + \|Y'\|_{p,[0,t]})\|X\|_{p,[0,t]} + \|R^Y\|_{\frac{p}{2},[0,t]}) \\ &\leq C_F + C_F(|Y'_0| + \|Y'\|_{p,[0,t]})\|X\|_{p,[0,t]} + C_F\|R^{Y-y}\|_{\frac{p}{2},[0,t]} + C_F\|R^y\|_{\frac{p}{2},[0,t]} \\ &\leq C_F(1 + |Y'_0| + \|Y'\|_{p,[0,t]})\|X\|_{p,[0,t]} + 3C_F, \end{aligned}$$

since (F, F') satisfies Assumption 2.1 (i), and by the local estimate for rough integration, see Lemma A.1,

$$\begin{aligned} \|R^Z - R^y\|_{\frac{p}{2},[0,t]} &= \|R^{\int_0^\cdot F(Y) d\mathbf{X}}\|_{\frac{p}{2},[0,t]} \\ &\lesssim C_F(1 + \|Y, Y'\|_{X,p,[0,t]})^2(1 + \|X\|_{p,[0,t]})\|\mathbf{X}\|_{p,[0,t]}, \end{aligned}$$

where the implicit multiplicative constant depends only on p . Hence, for $t = t_1$ sufficiently small we obtain that \mathcal{B}_{t_1} is invariant under \mathcal{M}_{t_1} . Note that t_1 depends on p , $|y'_0|$, C_F and $\|\mathbf{X}\|_p$.

Contraction. Let $(Y, Y'), (\tilde{Y}, \tilde{Y}') \in \mathcal{B}_t$ for some $t \in (0, t_1]$, that is, setting $K := 5(1 + \|y, y'\|_{X,p} + C_F)$, it holds that $\|Y, Y'\|_{X,p,[0,t]}, \|\tilde{Y}, \tilde{Y}'\|_{p,[0,t]} \leq K$. We have that

$$\begin{aligned} \|Z' - \tilde{Z}'\|_{p,[0,t]} &= \|F(Y) - F(\tilde{Y})\|_{p,[0,t]} \\ &\leq C_{F,K,X,X}\|Y - \tilde{Y}\|_{p,[0,t]} \\ &\lesssim_p C_{F,K,X,X}(\|Y' - \tilde{Y}'\|_{p,[0,t]}\|X\|_{p,[0,t]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2},[0,t]}). \end{aligned}$$

Further, due to Assumption 2.1 (ii) and Lemma A.2, it holds that

$$\begin{aligned} \|R^Z - R^{\tilde{Z}}\|_{\frac{p}{2},[0,t]} &= \|R^{\int_0^\cdot F(Y) d\mathbf{X} - \int_0^\cdot F(\tilde{Y}) d\mathbf{X}}\|_{\frac{p}{2},[0,t]} \\ &\lesssim C_{F,K,X,X}\|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X,p,[0,t]}\|\mathbf{X}\|_{p,[0,t]}, \end{aligned}$$

where the implicit multiplicative constant depends on p and $\|\mathbf{X}\|_p$. Defining an equivalent norm on \mathcal{V}_X^p by

$$\|Y, Y'\|_{X,p,[0,t]}^{(\delta)} := |Y_0'| + \|Y'\|_{p,[0,t]} + \delta \|R^Y\|_{\frac{p}{2},[0,t]}, \quad \text{for } \delta \geq 1,$$

we then deduce that

$$\begin{aligned} & \|Z - \tilde{Z}, Z' - \tilde{Z}'\|_{X,p,[0,t]}^{(\delta)} \\ & \lesssim C_{F,K,X,X} (\|Y' - \tilde{Y}'\|_{p,[0,t]} \|X\|_{p,[0,t]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2},[0,t]}) \\ & \quad + \delta C_{F,K,X,X} (\|Y' - \tilde{Y}'\|_{p,[0,t]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2},[0,t]}) \|\mathbf{X}\|_{p,[0,t]} \\ & \lesssim C_{F,K,X,X} (1 + \delta) \|\mathbf{X}\|_{p,[0,t]} \|Y' - \tilde{Y}'\|_{p,[0,t]} + C_{F,K,X,X} (1 + \delta) \|\mathbf{X}\|_{p,[0,t]} \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2},[0,t]} \\ & \lesssim C_{F,K,X,X} \left((1 + \delta) \|\mathbf{X}\|_{p,[0,t]} \vee \frac{1 + \delta \|\mathbf{X}\|_{p,[0,t]}}{\delta} \right) \|Y - \tilde{Y}, Y' - \tilde{Y}'\|_{X,p,[0,t]}^{(\delta)}, \end{aligned}$$

where the implicit multiplicative constant depends on p and $\|\mathbf{X}\|_p$. Hence, we can choose δ sufficiently large and $t = t_2 \leq t_1$ sufficiently small such that

$$C_{F,K,X,X} \left((1 + \delta) \|\mathbf{X}\|_{p,[0,t_2]} \vee \frac{1 + \delta \|\mathbf{X}\|_{p,[0,t_2]}}{\delta} \right) \leq 1,$$

where the left-hand side is up to a multiplicative constant which depends on p and $\|\mathbf{X}\|_p$. It follows that \mathcal{M}_{t_2} is a contraction on the subset of controlled paths $(\mathcal{B}_{t_2}, \|\cdot\|_{X,p,[0,t_2]}^{(\delta)})$. Hence, by Banach's fixed point theorem, there exists a unique fixed point of the map \mathcal{M}_{t_2} , which is the unique solution of the RFDE (2.2) over the time interval $[0, t_2]$.

Step 2: Global solution. Let $w: \Delta_T \rightarrow [0, \infty)$ be the right-continuous control function given by

$$w(s, t) := \|X\|_{p,[s,t]}^p + \|\mathbb{X}\|_{\frac{p}{2},[s,t]}^{\frac{p}{2}}, \quad (s, t) \in \Delta_T.$$

We infer from Step 1 that there exists a constant $\gamma > 0$, which depends on p , $\|y, y'\|_{X,p}$, C_F , $C_{F,K,X,X}$ and $\|\mathbf{X}\|_p$, such that the local solution (Y, Y') established above exists on any interval $[s, t]$ such that $w(s, t) \leq \gamma$, given any initial condition $\xi \in \mathcal{V}_X^p$ with $|\xi'_s| \leq \|y, y'\|_{X,p}$.

By [FZ18, Lemma 1.5], there exists a partition $\mathcal{P} = \{0 = \tau_0 < \tau_1 < \dots < \tau_N = T\}$ of $[0, T]$, such that $w(\tau_i, \tau_{i+1}-) < \gamma$ for every $i = 0, 1, \dots, N-1$. We can then define the solution (Y, Y') on each of the half-intervals $[\tau_i, \tau_{i+1})$. Given the solutions on $[\tau_i, \tau_{i+1})$, the values $Y_{\tau_{i+1}}$ at the right end-point of the interval are uniquely determined by the jumps of \mathbf{X} at time τ_{i+1} . More precisely, let $y_{0,\cdot} = y$, and define y_i , $i = 1, \dots, N-1$, by

$$y_{i;t} = y_t + Y_{\tau_i-} - y_{\tau_i-} + F_{\tau_i-}(Y)X_{\tau_i-, \tau_i} + F'_{\tau_i-}(Y, Y')\mathbb{X}_{\tau_i-, \tau_i}, \quad t \in [\tau_i, \tau_{i+1}).$$

We note that $|y'_{i,\tau_i}| = |y'_i| \leq \|y, y'\|_{X,p}$. Given the initial condition $(y_i, y'_i) \in \mathcal{V}_X^p$, we obtain the solution (Y, Y') on $[\tau_i, \tau_{i+1})$, $i = 0, 1, \dots, N-1$. By pasting the solutions on each of these subintervals together, with $Y_T = y_{N;T}$, we obtain a unique global solution (Y, Y') of the RFDE (2.2) on the interval $[0, T]$.

Step 3: A priori estimate. It remains to show the existence of a componentwise non-decreasing function $K_p: [0, \infty)^3 \rightarrow [0, \infty)$ such that

$$\|Y, Y'\|_{X,p} \leq K_p(\|y, y'\|_{X,p}, C_F, \|\mathbf{X}\|_p).$$

As stated in Step 2, we obtain the existence of a partition $\mathcal{P} = \{0 = \tau_0 < \tau_1 < \dots < \tau_N = T\}$ and define the solution on (Y, Y') on each of the half-intervals $[t_i, t_{i+1})$, which then satisfies

$$(2.3) \quad \|Y'\|_{p, [t_i, t_{i+1})} \leq 4C_F + \|y'\|_{p, [t_i, t_{i+1})}$$

as well as

$$(2.4) \quad \|R^Y\|_{\frac{p}{2}, [\tau_i, \tau_{i+1})} \leq 1 + \|R^y\|_{\frac{p}{2}, [\tau_i, \tau_{i+1})}$$

for all $i = 0, \dots, N-1$. Here, N depends on $p, \|y, y'\|_{X,p}, C_F, C_{F,K,X,X}, \|\mathbf{X}\|_p$ and is, for p fixed, non-decreasing in the other variables. Observe that

$$Y_{t-,t} = y_{t-,t} + \left(\int_0^{\cdot} F_s(Y) d\mathbf{X}_s \right)_{t-,t} = y_{t-,t} + F_{t-}(Y)X_{t-,t} + F'_{t-}(Y, Y')\mathbb{X}_{t-,t},$$

for any $t \in (0, T]$, so we have

$$R_{t-,t}^Y = R_{t-,t}^y + F'_{t-}(Y, Y')\mathbb{X}_{t-,t}.$$

This yields

$$|R_{\tau_{i+1}^-, \tau_{i+1}}^Y| \leq \|R^y\|_{p, [\tau_i, \tau_{i+1})} + (|F'_{\tau_i}(Y, Y')| + \|F'(Y, Y')\|_{p, [\tau_i, \tau_{i+1})}) |\mathbb{X}_{\tau_{i+1}^-, \tau_{i+1}}|.$$

Now, we use Assumption 2.1 (i), i.e.

$$|F'_{\tau_i}(Y, Y')| + \|F'(Y, Y')\|_{p, [\tau_i, \tau_{i+1})} \leq C_F(1 + \|Y, Y'\|_{X,p, [\tau_i, \tau_{i+1})})^2(1 + \|X\|_{p, [\tau_i, \tau_{i+1})})^2.$$

Since

$$\|Y, Y'\|_{X,p, [\tau_i, \tau_{i+1})} \leq |y'_{\tau_i}| + |F_{\tau_i}(Y)| + \|Y'\|_{p, [\tau_i, \tau_{i+1})} + \|R^Y\|_{\frac{p}{2}, [\tau_i, \tau_{i+1})},$$

It follows from Assumption 2.1 (i), (2.3) and (2.4) that

$$\|Y, Y'\|_{X,p, [\tau_i, \tau_{i+1})} \leq |y'_{\tau_i}| + 5(1 + C_F) + \|y'\|_{p, [\tau_i, \tau_{i+1})} + \|R^y\|_{\frac{p}{2}, [\tau_i, \tau_{i+1})}.$$

Consequently, there exists a componentwise non-decreasing polynomial $Q_p^{(R)}: [0, \infty)^3 \rightarrow [0, \infty)$ such that

$$|R_{\tau_{i+1}^-, \tau_{i+1}}^Y| \leq Q_p^{(R)}(\|y, y'\|_{X,p, [\tau_i, \tau_{i+1})}, C_F, \|\mathbf{X}\|_{p, [\tau_i, \tau_{i+1})})$$

as well as

$$(2.5) \quad \|R^Y\|_{\frac{p}{2}, [\tau_i, \tau_{i+1})} \leq 1 + \|R^y\|_{\frac{p}{2}, [\tau_i, \tau_{i+1})} + Q_p^{(R)}(\|y, y'\|_{X,p, [\tau_i, \tau_{i+1})}, C_F, \|\mathbf{X}\|_{p, [\tau_i, \tau_{i+1})})$$

for all $i = 0, \dots, N-1$. Moreover, since

$$Y'_{t-,t} = y'_{t-,t} + F_{t-,t}(Y),$$

for any $t \in (0, T]$, we have

$$|Y'_{\tau_{i+1}^-, \tau_{i+1}}| \leq \|y'\|_{p, [\tau_i, \tau_{i+1})} + |F_{\tau_{i+1}^-, \tau_{i+1}}(Y)|.$$

By Assumption 2.1 (i), it holds that

$$|F_{\tau_{i+1}^-, \tau_{i+1}}(Y)| \leq C_F(1 + \|Y\|_{p, [\tau_i, \tau_{i+1})} + |Y_{\tau_{i+1}^-, \tau_{i+1}}|),$$

thus, we need to control the jump of Y at τ_{i+1} . For this, note that

$$\begin{aligned}
& |Y_{\tau_{i+1}-, \tau_{i+1}}| \\
& \leq |y_{\tau_{i+1}-, \tau_{i+1}}| + |F_{\tau_{i+1}-}(Y)| |X_{\tau_{i+1}-, \tau_{i+1}}| + |F'_{\tau_{i+1}-}(Y)| |\mathbb{X}_{\tau_{i+1}-, \tau_{i+1}}| \\
& \leq |y_{\tau_{i+1}-, \tau_{i+1}}| + (|F_{\tau_i}(Y)| + \|F(Y)\|_{p, [\tau_i, \tau_{i+1}]}) |X_{\tau_{i+1}-, \tau_{i+1}}| \\
& \quad + (|F'_{\tau_i}(Y, Y')| + \|F'(Y, Y')\|_{p, [\tau_i, \tau_{i+1}]}) |\mathbb{X}_{\tau_{i+1}-, \tau_{i+1}}| \\
& \leq \|y, y'\|_{p, [\tau_i, \tau_{i+1}]} + (C_F + C_F(1 + (\|Y'_{\tau_i}\| + \|Y'\|_{\frac{p}{2}, [\tau_i, \tau_{i+1}]})) \|X\|_{p, [\tau_i, \tau_{i+1}]}) \\
& \quad + \|R^Y\|_{\frac{p}{2}, [\tau_i, \tau_{i+1}]} \|X\|_{p, [\tau_i, \tau_{i+1}]} \\
& \quad + C_F(1 + \|Y, Y'\|_{p, [\tau_i, \tau_{i+1}]})^2 (1 + \|X\|_{p, [\tau_i, \tau_{i+1}]})^2 \|\mathbb{X}\|_{\frac{p}{2}, [\tau_i, \tau_{i+1}]}.
\end{aligned}$$

Using (2.3) and (2.4), we can now conclude that there exist componentwise non-decreasing polynomials $Q_p^{(Y, J)}, Q_p^{(Y')}: [0, \infty)^3 \rightarrow [0, \infty)$ such that

$$|F_{\tau_{i+1}-, \tau_{i+1}}(Y)| \leq Q_p^{(Y, J)}(\|y, y'\|_{X, p, [\tau_i, \tau_{i+1}]}, C_F, \|\mathbf{X}\|_{p, [\tau_i, \tau_{i+1}]})$$

as well as

$$(2.6) \quad \|Y'\|_{p, [\tau_i, \tau_{i+1}]} \leq Q_p^{(Y')}(\|y, y'\|_{X, p, [\tau_i, \tau_{i+1}]}, C_F, \|\mathbf{X}\|_{p, [\tau_i, \tau_{i+1}]}).$$

Combining these estimates, we obtain that

$$\|Y'_0\| + \|Y'\|_p + \|R^Y\|_{\frac{p}{2}} \leq K_p(\|y, y'\|_{X, p}, C_F, \|\mathbf{X}\|_p),$$

which is the assertion. \square

2.3. Continuity of the Itô–Lyons map. A fundamental contribution of the theory of rough paths is the insight that the solution map, mapping the input (initial condition, coefficients, driving rough path, ...) of a rough differential equation to its solution, is continuous with respect to suitable distances on the spaces of controlled paths as well as of rough paths, see e.g. [FH20]. In the context of rough differential equations, this solution map is also known as Itô–Lyons map. In the next theorem we present the local Lipschitz continuity of the Itô–Lyons map for rough functional differential equations.

Theorem 2.5. *Let $\mathbf{X}, \tilde{\mathbf{X}} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ be càdlàg p -rough paths for $p \in (2, 3)$, $(y, y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $(\tilde{y}, \tilde{y}') \in \mathcal{V}_{\tilde{X}}^p([0, T]; \mathbb{R}^k)$ be given controlled paths with respect to X and \tilde{X} , respectively. Suppose that the non-anticipative functionals (F, F') , (\tilde{F}, \tilde{F}') satisfy Assumption 2.1 and $(F - \tilde{F}, F' - \tilde{F}')$ satisfies Assumption 2.1 (i) with some constant $C_{F-\tilde{F}}$.*

Let $(Y, Y) \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ be the solution given by Theorem 2.4 to the rough functional differential equation (2.2), and $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_{\tilde{X}}^p([0, T]; \mathbb{R}^k)$ be the solution to the rough functional differential equation (2.2) driven by $\tilde{\mathbf{X}}$ with initial condition (\tilde{y}, \tilde{y}') and functional (\tilde{F}, \tilde{F}') , and suppose that $\|Y, Y'\|_{X, p}, \|\tilde{Y}, \tilde{Y}'\|_{\tilde{X}, p} \leq K$, for some $K > 0$. Then, we have the estimate

$$\begin{aligned}
& |Y_0 - \tilde{Y}_0| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p} \\
& \lesssim |y_0 - \tilde{y}_0| + |F_0(y) - \tilde{F}_0(\tilde{y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_p,
\end{aligned}$$

where the implicit multiplicative constant depends on $p, C_F \vee C_{\tilde{F}}, C_{F, K, X, \tilde{X}} \vee C_{\tilde{F}, K, X, \tilde{X}}, K, \|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$.

Proof. Step 1: Local estimate. Let $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_{\tilde{X}}^p([0, T]; \mathbb{R}^k)$ be the global solutions to the RFDE (2.2), with data $((y, y'), (F, F'), \mathbf{X})$, $((\tilde{y}, \tilde{y}'), (\tilde{F}, \tilde{F}'), \tilde{\mathbf{X}})$, respectively, see Theorem 2.4. Let $t \in (0, T]$. Without loss of generality assume that $C_{\tilde{F}} \leq C_F$, $C_{\tilde{F}, K, X, \tilde{X}} \leq C_{F, K, X, \tilde{X}}$. Assumption 2.1 gives that

$$\begin{aligned}
& \|F(Y) - \tilde{F}(\tilde{Y})\|_{p, [0, t]} \\
& \leq \|F(Y) - F(\tilde{Y})\|_{p, [0, t]} + \|(F - \tilde{F})(\tilde{Y})\|_{p, [0, t]} \\
& \leq C_{F, K, X, \tilde{X}}(|Y_0 - \tilde{Y}_0| + \|Y - \tilde{Y}\|_{p, [0, t]}) \\
& \quad + C_{F - \tilde{F}}(1 + (\|\tilde{Y}'_0\| + \|\tilde{Y}'\|_{p, [0, t]}))\|X\|_{p, [0, t]} + \|R^{\tilde{Y}}\|_{\frac{p}{2}, [0, t]} \\
& \lesssim_p C_{F, K, X, \tilde{X}}(|Y_0 - \tilde{Y}_0| + (\|Y'_0 - \tilde{Y}'_0\| + \|Y' - \tilde{Y}'\|_{p, [0, t]}))(\|X\|_{p, [0, t]} \vee \|\tilde{X}\|_{p, [0, t]}) \\
& \quad + C_{F, K, X, \tilde{X}}\|R^y - R^{\tilde{y}}\|_{\frac{p}{2}, [0, t]} + C_{F, K, X, \tilde{X}}\|R^{\int_0^\cdot F(Y)d\mathbf{X}} - R^{\int_0^\cdot \tilde{F}(\tilde{Y})d\tilde{\mathbf{X}}}\|_{\frac{p}{2}, [0, t]} \\
& \quad + C_{F, K, X, \tilde{X}}\|X - \tilde{X}\|_{p, [s, t]} + C_{F - \tilde{F}}(1 + K)(1 + \|X\|_{p, [0, t]} \vee \|\tilde{X}\|_{p, [0, t]}).
\end{aligned}$$

Further, by Lemma A.1 and Lemma A.2 we have that

$$\begin{aligned}
& \|R^{\int_0^\cdot F(Y)d\mathbf{X}} - R^{\int_0^\cdot \tilde{F}(\tilde{Y})d\tilde{\mathbf{X}}}\|_{\frac{p}{2}, [0, t]} \\
& \leq \|R^{\int_0^\cdot F(Y)d\mathbf{X}} - R^{\int_0^\cdot F(\tilde{Y})d\tilde{\mathbf{X}}}\|_{\frac{p}{2}, [0, t]} + \|R^{\int_0^\cdot (F - \tilde{F})(\tilde{Y})d\tilde{\mathbf{X}}}\|_{\frac{p}{2}, [0, t]} \\
& \lesssim C_{F, K, X, \tilde{X}}(|Y_0 - \tilde{Y}_0| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p, [0, t]} + \|X - \tilde{X}\|_{p, [0, t]})(\|\mathbf{X}\|_{p, [0, t]} \vee \|\tilde{\mathbf{X}}\|_{p, [0, t]}) \\
& \quad + C_F(1 + K)^2(1 + \|X\|_{p, [0, t]} \vee \|\tilde{X}\|_{p, [0, t]})^2\|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [0, t]} \\
& \quad + C_{F - \tilde{F}}(1 + K)^2(1 + \|X\|_{p, [0, t]} \vee \|\tilde{X}\|_{p, [0, t]})^2(\|\mathbf{X}\|_{p, [0, t]} \vee \|\tilde{\mathbf{X}}\|_{p, [0, t]}),
\end{aligned}$$

where the implicit multiplicative constant depends on p , $\|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$. Combining the results, we get that

$$\begin{aligned}
& \|Y' - \tilde{Y}'\|_{p, [0, t]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [0, t]} \\
& \leq \|y' - \tilde{y}'\|_{p, [0, t]} + \|F(Y) - \tilde{F}(\tilde{Y})\|_{p, [0, t]} \\
& \quad + \|R^y - R^{\tilde{y}}\|_{\frac{p}{2}, [0, t]} + \|R^{\int_0^\cdot F(Y)d\mathbf{X}} - R^{\int_0^\cdot \tilde{F}(\tilde{Y})d\tilde{\mathbf{X}}}\|_{\frac{p}{2}, [0, t]} \\
& \leq C_1(\|\mathbf{X}\|_{p, [0, t]} \vee \|\tilde{\mathbf{X}}\|_{p, [0, t]})(\|Y' - \tilde{Y}'\|_{p, [0, t]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [0, t]}) \\
& \quad + C_2(|Y_0 - \tilde{Y}_0| + |F_0(Y) - \tilde{F}_0(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p, [0, t]} + C_{F - \tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [0, t]})
\end{aligned}$$

for some constants $C_1 > 0$, which depends on p , $C_{F, K, X, \tilde{X}}$, $\|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$, and $C_2 > 1$, which depends additionally on C_F and K . Hence, we can choose t sufficiently small such that $C_1(\|\mathbf{X}\|_{p, [0, t]} \vee \|\tilde{\mathbf{X}}\|_{p, [0, t]}) \leq \frac{1}{2}$, which implies that

$$\begin{aligned}
& \|Y' - \tilde{Y}'\|_{p, [0, t]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [0, t]} \\
(2.7) \quad & \lesssim |Y_0 - \tilde{Y}_0| + |F_0(Y) - \tilde{F}_0(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p, [0, t]} + C_{F - \tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [0, t]}.
\end{aligned}$$

Step 2: Global estimate. Recall the right-continuous control function $w: \Delta_T \rightarrow [0, \infty)$ given by

$$w(s, t) := \|X\|_{p, [s, t]}^p + \|\mathbb{X}\|_{\frac{p}{2}, [s, t]}^{\frac{p}{2}}, \quad (s, t) \in \Delta_T,$$

as introduced in the proof of Theorem 2.4, and let $\tilde{w}(s, t) := \|\tilde{X}\|_{p, [s, t]}^p + \|\tilde{X}\|_{\frac{p}{2}, [s, t]}^{\frac{p}{2}}$, $(s, t) \in \Delta_T$. We infer from Step 1 that there exists a constant $\gamma > 0$, which depends on p , $C_{F, K, X, \tilde{X}}$, $\|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$, such that on any interval $[s, t]$ with $w(s, t) \vee \tilde{w}(s, t) \leq \gamma$ the local solutions satisfy an estimate of the form (2.7).

Let $c(s, t) := w(s, t) + \tilde{w}(s, t)$, $(s, t) \in \Delta_T$. Since c is right-continuous, there exists a partition $\mathcal{P} = \{0 = t_0 < \dots < t_N = T\}$ of $[0, T]$, such that

$$c(t_i, t_{i+1}-) = \gamma, \quad \text{or} \quad c(t_i, t_{i+1}-) < \gamma \quad \text{and} \quad c(t_i, t_{i+1}-) + c(t_{i+1}-, t_{i+1}) \geq \gamma,$$

for every $i = 0, 1, \dots, N-1$. Since w and \tilde{w} and, thus, c is superadditive, we have that

$$N\gamma \leq \sum_{i=0}^{N-1} c(t_i, t_{i+1}-) + c(t_{i+1}-, t_{i+1}) \leq c(0, T).$$

Therefore, the number of partition points N may be bounded by a constant depending only on γ , $w(0, T)$ and $\tilde{w}(0, T)$. Thus, in this step, we may combine the local estimates on each of the subintervals $[t_i, t_{i+1})$, together with simple estimates on the jumps at the end-points of these subintervals, which we aim to derive, to obtain the global estimate. More precisely, by Step 1, we have the local estimate

$$\begin{aligned} (2.8) \quad & \|Y' - \tilde{Y}'\|_{p, [t_i, t_{i+1})} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [t_i, t_{i+1})} \\ & \lesssim |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + |y'_{t_i} - \tilde{y}'_{t_i}| \\ & \quad + \|y' - \tilde{y}'\|_{p, [t_i, t_{i+1})} + \|R^y - R^{\tilde{y}}\|_{\frac{p}{2}, [t_i, t_{i+1})} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_i, t_{i+1})}, \end{aligned}$$

for $i = 0, \dots, N-1$, where the implicit multiplicative constant depends on p and $C_{F, K, X, \tilde{X}}$, C_F , K , $\|\mathbf{X}\|_p$, $\|\tilde{\mathbf{X}}\|_p$, but not on the index i . So, it remains to bound

$$|Y'_{t_{i+1}-, t_{i+1}} - \tilde{Y}'_{t_{i+1}-, t_{i+1}}| + |R^Y_{t_{i+1}-, t_{i+1}} - R^{\tilde{Y}}_{t_{i+1}-, t_{i+1}}|$$

to extend the previous estimate to $[t_i, t_{i+1}]$.

We note that $(\int_0^\cdot F_s(Y) d\mathbf{X}_s)_{t-, t} = F_{t-}(Y)X_{t-, t} + F'_{t-}(Y, Y')\mathbb{X}_{t-, t}$, that is, with $Y_{t-, t} = y_{t-, t} + (\int_0^\cdot F_s(Y) d\mathbf{X}_s)_{t-, t}$, it follows that $Y'_{t-, t} = y'_{t-, t} + F_{t-, t}(Y)$ and $R^Y_{t-, t} = R^y_{t-, t} + F'_{t-}(Y, Y')\mathbb{X}_{t-, t}$, for $t \in (0, T]$. Given the assumptions, we then have

$$\begin{aligned} (2.9) \quad & \|Y' - \tilde{Y}'\|_{p, [t_i, t_{i+1}]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [t_i, t_{i+1}]} \\ & \leq \|Y' - \tilde{Y}'\|_{p, [t_i, t_{i+1})} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [t_i, t_{i+1})} \\ & \quad + |Y'_{t_{i+1}-, t_{i+1}} - \tilde{Y}'_{t_{i+1}-, t_{i+1}}| + |R^Y_{t_{i+1}-, t_{i+1}} - R^{\tilde{Y}}_{t_{i+1}-, t_{i+1}}| \\ & \lesssim |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p, [t_i, t_{i+1}]} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_i, t_{i+1}]}, \end{aligned}$$

where the implicit multiplicative constant depends on p , C_F , $C_{F,K,X,\tilde{X}}$, K , $\|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$, and not on the index i . Here, we used Assumption (2.1) and the estimate (2.8) to derive that

$$\begin{aligned}
& |(R_{t_{i+1}^-, t_{i+1}}^Y - R_{t_{i+1}^-, t_{i+1}}^{\tilde{Y}}) - (R_{t_{i+1}^-, t_{i+1}}^y - R_{t_{i+1}^-, t_{i+1}}^{\tilde{y}})| \\
& \leq |F'_{t_{i+1}^-}(Y, Y') \mathbb{X}_{t_{i+1}^-, t_{i+1}} - \tilde{F}'_{t_{i+1}^-}(\tilde{Y}, \tilde{Y}') \tilde{\mathbb{X}}_{t_{i+1}^-, t_{i+1}}| \\
& \leq |F'_{t_{i+1}^-}(Y, Y') - \tilde{F}'_{t_{i+1}^-}(\tilde{Y}, \tilde{Y}')| |\mathbb{X}_{t_{i+1}^-, t_{i+1}}| + |\tilde{F}'_{t_{i+1}^-}(\tilde{Y}, \tilde{Y}')| |\mathbb{X}_{t_{i+1}^-, t_{i+1}} - \tilde{\mathbb{X}}_{t_{i+1}^-, t_{i+1}}| \\
& \leq (|F'_{t_i}(Y, Y') - \tilde{F}'_{t_i}(\tilde{Y}, \tilde{Y}')| + \|F'(Y, Y') - \tilde{F}'(\tilde{Y}, \tilde{Y}')\|_{p, [t_i, t_{i+1}]}) \|\mathbb{X}\|_{\frac{p}{2}, [t_i, t_{i+1}]} \\
& \quad + (|\tilde{F}'_{t_i}(\tilde{Y}, \tilde{Y}')| + \|\tilde{F}'(\tilde{Y}, \tilde{Y}')\|_{p, [t_i, t_{i+1}]}) \|\mathbb{X} - \tilde{\mathbb{X}}\|_{\frac{p}{2}, [t_i, t_{i+1}]} \\
& \lesssim |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, p, [t_i, t_{i+1}]} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_i, t_{i+1}]},
\end{aligned}$$

and

$$\begin{aligned}
& |(Y'_{t_{i+1}^-, t_{i+1}} - \tilde{Y}'_{t_{i+1}^-, t_{i+1}}) - (y'_{t_{i+1}^-, t_{i+1}} - \tilde{y}'_{t_{i+1}^-, t_{i+1}})| \\
& = |F_{t_{i+1}^-, t_{i+1}}(Y) - \tilde{F}_{t_{i+1}^-, t_{i+1}}(\tilde{Y})| \\
& \leq |(F - \tilde{F})_{t_{i+1}^-, t_{i+1}}(Y)| + |\tilde{F}_{t_{i+1}^-, t_{i+1}}(Y) - \tilde{F}_{t_{i+1}^-, t_{i+1}}(\tilde{Y})| \\
& \leq C_{F-\tilde{F}}(1 + (|Y'_{t_i}| + \|Y'\|_{p, [t_i, t_{i+1}]}) \|X\|_{p, [t_i, t_{i+1}]} + \|R^Y\|_{\frac{p}{2}, [t_i, t_{i+1}]}) \\
& \quad + C_{F,K,X,\tilde{X}}(|Y_{t_i} - \tilde{Y}_{t_i}| + \|Y - \tilde{Y}\|_{p, [t_i, t_{i+1}]} + |Y_{t_{i+1}^-, t_{i+1}} - \tilde{Y}_{t_{i+1}^-, t_{i+1}}|) \\
& \lesssim |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, p, [t_i, t_{i+1}]} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_i, t_{i+1}]},
\end{aligned}$$

where the implicit multiplicative constant depends on p , C_F , $C_{F,K,X,\tilde{X}}$, K and $\|\mathbf{X}\|_p \vee \|\tilde{\mathbf{X}}\|_p$, and not on the index i .

Now, we need to control the term $|Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + |y'_{t_i} - \tilde{y}'_{t_i}|$. For this, we note that

$$\begin{aligned}
& |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + |y'_{t_i} - \tilde{y}'_{t_i}| \\
& \leq |Y_{t_{i-1}} - \tilde{Y}_{t_{i-1}}| + |F_{t_{i-1}}(Y) - \tilde{F}_{t_{i-1}}(\tilde{Y})| + |y'_{t_{i-1}} - \tilde{y}'_{t_{i-1}}| \\
& \quad + \|Y - \tilde{Y}\|_{p, [t_{i-1}, t_i]} + \|F(Y) - \tilde{F}(\tilde{Y})\|_{p, [t_{i-1}, t_i]} + \|y' - \tilde{y}'\|_{p, [t_{i-1}, t_i]} \\
& \lesssim |Y_{t_{i-1}} - \tilde{Y}_{t_{i-1}}| + |F_{t_{i-1}}(Y) - \tilde{F}_{t_{i-1}}(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p, [t_{i-1}, t_i]} \\
& \quad + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_{i-1}, t_i]} \\
& \quad + \|Y' - \tilde{Y}'\|_{p, [t_{i-1}, t_i]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [t_{i-1}, t_i]},
\end{aligned}$$

where the implicit multiplicative constant depends on p , C_F , $C_{F,K,X,\tilde{X}}$, K , $\|X\|_p$ and $\|\tilde{X}\|_p$, and not on the index i ; thus applying the estimate (2.9) for $i-1$ gives that

$$\begin{aligned}
& |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + |y'_{t_i} - \tilde{y}'_{t_i}| \\
& \lesssim |Y_{t_{i-1}} - \tilde{Y}_{t_{i-1}}| + |F_{t_{i-1}}(Y) - \tilde{F}_{t_{i-1}}(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p, [t_{i-1}, t_i]} \\
& \quad + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_{i-1}, t_i]}.
\end{aligned}$$

Iteratively, we obtain for any $i = 1, \dots, N$ that

$$\begin{aligned} & |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + |y'_{t_i} - \tilde{y}'_{t_i}| \\ & \lesssim |y_0 - \tilde{y}_0| + |F_0(y) - \tilde{F}_0(\tilde{y})| + |y'_0 - \tilde{y}'_0| \\ & \quad + \sum_{j=0}^{i-1} \left(\|y' - \tilde{y}'\|_{p, [t_j, t_{j+1}]} + \|R^y - R^{\tilde{y}}\|_{\frac{p}{2}, [t_j, t_{j+1}]} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_j, t_{j+1}]} \right), \end{aligned}$$

that is,

$$\begin{aligned} & (|Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + |y'_{t_i} - \tilde{y}'_{t_i}|)^p \\ & \lesssim (|y_0 - \tilde{y}_0| + |F_0(y) - \tilde{F}_0(\tilde{y})| + |y'_0 - \tilde{y}'_0| + NC_{F-\tilde{F}})^p \\ & \quad + \sum_{j=0}^{i-1} \left(\|y' - \tilde{y}'\|_{p, [t_j, t_{j+1}]}^p + \|R^y - R^{\tilde{y}}\|_{\frac{p}{2}, [t_j, t_{j+1}]}^p + \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [t_j, t_{j+1}]}^p \right). \end{aligned}$$

This implies that

$$\begin{aligned} & |Y_{t_i} - \tilde{Y}_{t_i}| + |F_{t_i}(Y) - \tilde{F}_{t_i}(\tilde{Y})| + |y'_{t_i} - \tilde{y}'_{t_i}| \\ & \lesssim |y_0 - \tilde{y}_0| + |F_0(y) - \tilde{F}_0(\tilde{y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_p, \end{aligned}$$

which is the desired control.

If we plug this into (2.9), it follows that

$$\begin{aligned} & \|Y' - \tilde{Y}'\|_{p, [t_i, t_{i+1}]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}, [t_i, t_{i+1}]} \\ & \lesssim |y_0 - \tilde{y}_0| + |F_0(y) - \tilde{F}_0(\tilde{y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, \tilde{X}, p} + C_{F-\tilde{F}} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_p. \end{aligned}$$

Since $\|\cdot\|_{p, [0, T]} \leq N \sum_{i=0}^{N-1} \|\cdot\|_{p, [t_i, t_{i+1}]}$ for any $p \geq 1$, see e.g. [All21, Lemma A.1], the estimate finally follows. \square

3. EXAMPLES OF RFDES

The general framework of rough functional differential equations, presented in Section 2, allows to treat various classes of rough differential equations. In this section, some exemplary rough functional differential equations are discussed, aiming to develop the main conceptual ideas and demonstrating the scope of RFDEs rather than pushing for the most general results.

3.1. Classical RDEs. Let us start with the classical rough differential equation (RDE)

$$(3.1) \quad Y_t = y_0 + \int_0^t f(Y_s) d\mathbf{X}_s, \quad t \in [0, T],$$

where $y_0 \in \mathbb{R}^k$, $f \in C_b^3(\mathbb{R}^k; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ and $\mathbf{X} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ for $p \in (2, 3)$. While the existence and uniqueness of solutions to the RDE (3.1) driven by a continuous rough path and the continuity of the solution map were first proven by Lyons [Lyo98], the analogous results for RDEs driven by càdlàg rough paths were more recently obtained by Friz and Zhang [FZ18]. As an application of Theorem 2.4 and Theorem 2.5, one can recover these results, demonstrating that Assumption 2.1 is, indeed, a natural generalization of the classical assumptions of the coefficients of a rough differential equation. Furthermore, note that Corollary 3.1 presents the continuity of the solution map with respect to the controlled path norm, which slightly generalizes [FZ18, Theorem 3.8].

Corollary 3.1.

- (i) If $f \in C_b^3(\mathbb{R}^k; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, there exists a unique solution to the RDE (3.1). Moreover, there exists a non-decreasing function $K_p: [0, \infty)^2 \rightarrow [0, \infty)$ such that

$$\|Y, Y'\|_{X,p} \leq K_p(\|f\|_{C_b^2}, \|\mathbf{X}\|_p).$$

- (ii) Let $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ be the unique solution to the RDE (3.1). Moreover, let $\tilde{y}_0 \in \mathbb{R}^k$, $\tilde{f} \in C_b^3(\mathbb{R}^k; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, $\tilde{\mathbf{X}} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ with corresponding solution $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_{\tilde{X}}^p([0, T]; \mathbb{R}^d)$, and suppose that $\|Y, Y'\|_{X,p}, \|\tilde{Y}, \tilde{Y}'\|_{\tilde{X},p} \leq K$, for some $K > 0$. Then, we have the estimate

$$|Y_0 - \tilde{Y}_0| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p} \lesssim |y_0 - \tilde{y}_0| + \|f - \tilde{f}\|_{C_b^2} + \|\mathbf{X}; \tilde{\mathbf{X}}\|_p,$$

where the implicit multiplicative constant depends on $p, \|f\|_{C_b^3} \vee \|\tilde{f}\|_{C_b^3}, K, \|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$.

In order to apply the existence and uniqueness result presented in Theorem 2.4, and also the continuity result presented in Theorem 2.5, we need to check that the vector field f in the RDE (3.1) satisfies Assumption 2.1. This is the content of the next lemma, which we formulate slightly more general, with regard to the dimensions of the underlying spaces, for later use.

Lemma 3.2. Let $f \in C_b^3(\mathbb{R}^m; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ and $X, \tilde{X} \in D^p([0, T]; \mathbb{R}^d)$. The non-anticipative functional

$$(F, F'): \mathcal{V}_X^p([0, T]; \mathbb{R}^m) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \quad (F(Y), F'(Y, Y')) := (f(Y), \text{D}f(Y)Y'),$$

satisfies Assumption 2.1.

Proof. Since the proof is fairly standard, we provide only a sketch of a proof, following, e.g., the proofs of [FZ18, Lemma 3.5, Lemma 3.6, Lemma 3.7].

Fix $(s, t) \in \Delta_T$ and let $(Y, Y') \in \mathcal{V}_X^p, (\tilde{Y}, \tilde{Y}') \in \mathcal{V}_{\tilde{X}}^p$.

Growth conditions. It is clear that $|F_t(Y)| \leq \|f\|_{C_b^2}$, and it follows from the Lipschitz continuity of f that

$$|F_{t-,t}(Y)| \leq \|F(Y)\|_{p,[s,t]} \leq \|f\|_{C_b^2} \|Y\|_{p,[s,t]}.$$

We now note that $\|Y\|_{p,[s,t]} \leq \|Y\|_{p,[s,t]} + |Y_{t-,t}|$ as well as $\|Y\|_{p,[s,t]} \leq C_p(1 + (|Y'_s| + \|Y'\|_{p,[s,t]})) \cdot \|X\|_{p,[s,t]} + \|R^Y\|_{\frac{p}{2},[s,t]}$. Further, it holds that

$$\begin{aligned} |F'_s(Y, Y')| + \|F'(Y, Y')\|_{p,[s,t]} &= |\text{D}f(Y_s)Y'_s| + \|\text{D}f(Y)Y'\|_{p,[s,t]} \\ &\leq \|f\|_{C_b^2} (|Y'_s| + \|Y'\|_{p,[s,t]}) (1 + \|Y\|_{p,[s,t]}) \\ &\lesssim_p \|f\|_{C_b^2} (|Y'_s| + \|Y'\|_{p,[s,t]}) (1 + (|Y'_s| + \|Y'\|_{p,[s,t]})) \|X\|_{p,[s,t]} + \|R^Y\|_{\frac{p}{2},[s,t]} \\ &\lesssim_p \|f\|_{C_b^2} (1 + \|Y, Y'\|_{X,p,[s,t]}) (1 + \|X\|_{p,[s,t]}) \end{aligned}$$

and by Taylor's expansion,

$$\begin{aligned} R_{s,t}^{F(Y)} &= R_{s,t}^{f(Y)} = f(Y_t) - f(Y_s) - \text{D}f(Y_s)Y_{s,t} + \text{D}f(Y_s)R_{s,t}^Y \\ &= \frac{1}{2} \text{D}^2 f(Y_s + \lambda Y_{s,t}) Y_{s,t}^2 + \text{D}f(Y_s)R_{s,t}^Y, \end{aligned}$$

with $\lambda \in [0, 1]$, which implies that

$$\begin{aligned} \|R^{F(Y)}\|_{\frac{p}{2}, [s, t]} &= \|R^{f(Y)}\|_{\frac{p}{2}, [s, t]} \leq \|f\|_{C_b^2} (\|Y\|_{p, [s, t]}^2 + \|R^Y\|_{\frac{p}{2}, [s, t]}) \\ &\lesssim_p \|f\|_{C_b^2} ((|Y'_s| + \|Y'\|_{p, [s, t]}) \|X\|_{p, [s, t]} + \|R^Y\|_{\frac{p}{2}, [s, t]})^2 + \|R^Y\|_{\frac{p}{2}, [s, t]} \\ &\lesssim_p \|f\|_{C_b^2} (1 + \|Y, Y'\|_{X, p, [s, t]})^2 (1 + \|X\|_{p, [s, t]})^2. \end{aligned}$$

Assumption 2.1 (i) therefore holds with some constant $C_F = \|f\|_{C_b^2}$ up to a multiplicative constant which depends on p .

Lipschitz conditions. Fix $K > 0$ and assume that $\|Y, Y'\|_{X, p, [s, t]}, \|\tilde{Y}, \tilde{Y}'\|_{\tilde{X}, p, [s, t]} \leq K$. The proofs work verbatim as the proofs of [FZ18, Lemma 3.1 and Lemma 3.7]. The constant $C_{F, K, X, \tilde{X}}$ depends on p and $\|f\|_{C_b^3}$, K , $\|X\|_p$ and $\|\tilde{X}\|_p$. \square

Proof of Corollary 3.1. (i) The existence and uniqueness of the solution follows immediately from Lemma 3.2 and Theorem 2.4. For the a priori estimate, note that $\|y, y'\|_{X, p} = 0$ and $C_F \lesssim_p \|f\|_{C_b^2}$.

(ii) To apply the continuity result presented in Theorem 2.5, we need to ensure that the functionals satisfy the required assumptions. For (F, F') , (\tilde{F}, \tilde{F}') , this is given in Lemma 3.2, and further,

$$\begin{aligned} (F - \tilde{F}, (F - \tilde{F})') &: \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \\ ((F - \tilde{F})(Y), (F - \tilde{F})'(Y, Y')) &:= (f(Y) - \tilde{f}(Y), Df(Y)Y' - D\tilde{f}(Y)Y'), \end{aligned}$$

satisfies Assumption 2.1 (i), since C_b^3 is a vector space. Thus we have, $C_{F - \tilde{F}} \lesssim_p \|f - \tilde{f}\|_{C_b^2}$. \square

3.2. Controlled RDEs. Motivated by pathwise stochastic control, see e.g. [DFG17, AC20], and robust stochastic filtering, see e.g. [All21], as well as analogously to controlled stochastic differential equations, see e.g. [Pha09], we consider the controlled rough differential equation

$$(3.2) \quad Y_t = y_t + \int_0^t f(\alpha_s, Y_s) d\mathbf{X}_s, \quad t \in [0, T],$$

where $\mathbf{X} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ for $p \in (2, 3)$, $(y, y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $f \in C_b^3(\mathbb{R}^{k+e}; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, and $(\alpha, \alpha') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^e)$ is a fixed controlled path, with $e \in \mathbb{N}$. In case of continuous rough paths and controls α of finite $\frac{p}{2}$ -variation, controlled RDEs were treated in [All21, Theorem 2.3]. The following corollary provides an existence, uniqueness and continuity result for controlled RDEs driven by càdlàg p -rough paths and with controls α , which are only required to be controlled paths.

Corollary 3.3.

- (i) *If $f \in C_b^3(\mathbb{R}^{k+e}; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ and $(\alpha, \alpha') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^e)$, then there exists a unique solution to the controlled rough differential equation (3.2). Moreover, there exists a componentwise non-decreasing function $K_p: [0, \infty)^5 \rightarrow [0, \infty)$ such that*

$$\|Y, Y'\|_{X, p} \leq K_p(\|f\|_{C_b^3}, \|y, y'\|_{X, p}, \|\alpha\|_p, \|\alpha, \alpha'\|_{X, p}, \|\mathbf{X}\|_p).$$

- (ii) *Let $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ be the unique solution to the controlled rough differential equation (3.2). Moreover, let $(\tilde{y}, \tilde{y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $(\tilde{\alpha}, \tilde{\alpha}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^e)$,*

$\tilde{f} \in C_b^3(\mathbb{R}^{k+e}; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, with corresponding solution $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, and suppose that $\|Y, Y'\|_{X,p}, \|\tilde{Y}, \tilde{Y}'\|_{X,p} \leq K$, for some $K > 0$. Then, we have the estimate

$$\begin{aligned} & |Y_0 - \tilde{Y}_0| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X,p} \\ & \lesssim |y_0 - \tilde{y}_0| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X,p} + \|f - \tilde{f}\|_{C_b^2} + |\alpha_0 - \tilde{\alpha}_0| + \|\alpha, \alpha'; \tilde{\alpha}, \tilde{\alpha}'\|_{X,p}, \end{aligned}$$

where the implicit multiplicative constant depends on $p, \|f\|_{C_b^3} \vee \|\tilde{f}\|_{C_b^3}, \|\alpha, \alpha'\|_{X,p}, \|\tilde{\alpha}, \tilde{\alpha}'\|_{X,p}, K$, and $\|\mathbf{X}\|_p$.

In order to apply the existence and uniqueness result presented in Theorem 2.4, and also the continuity result in Theorem 2.5, we need to check that the vector field in the RDE (3.2) satisfies Assumption 2.1. This is the content of the next lemma. Note that it will be sufficient to check Assumption 2.1 (ii) for $X = \tilde{X}$, since we do not establish stability results with respect to the driving rough path in this subsection. (This also applies to the following subsections.)

Lemma 3.4. *Let $f \in C_b^3(\mathbb{R}^{k+e}; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ and $X \in D^p([0, T]; \mathbb{R}^d)$ for $p \in (2, 3)$. Further, let $(\alpha, \alpha') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^e)$. The non-anticipative functional*

$$\begin{aligned} & (F, F'): \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \\ & (F(Y), F'(Y, Y')) := (f((\alpha, Y)), Df((\alpha, Y))(\alpha', Y')), \end{aligned}$$

satisfies Assumption 2.1 for $X = \tilde{X}$.

Proof. Fix $(s, t) \in \Delta_T$ and let $(Y, Y') \in \mathcal{V}_X^p$. It is clear that $|F_t(Y)| \leq \|f\|_{C_b^2}$, and we note that

$$\begin{aligned} |F_{t-,t}(Y)| & \leq \|F(Y)\|_{p,[s,t]} = \|f((\alpha, Y))\|_{p,[s,t]} \\ & \leq \|f\|_{C_b^2} \|(\alpha, Y)\|_{p,[s,t]} \\ & \leq \|f\|_{C_b^2} (1 + \|\alpha\|_{p,[s,t]})(1 + \|Y\|_{p,[s,t]}), \end{aligned}$$

and it holds that $\|Y\|_{p,[s,t]} \leq \|Y\|_{p,[s,t]} + |Y_{t-,t}|$ as well as $\|Y\|_{p,[s,t]} \leq C_p(1 + (|Y'_s| + \|Y'\|_{p,[s,t]}) \cdot \|X\|_{p,[s,t]} + \|R^Y\|_{\frac{p}{2},[s,t]})$. Applying Lemma 3.2 to the enlarged controlled path $((\alpha, Y), (\alpha', Y'))$,

$$\begin{aligned} & \|F(Y), F'(Y, Y')\|_{X,p,[s,t]} \\ & \lesssim \|f\|_{C_b^2} (1 + |(\alpha'_s, Y'_s)| + \|(\alpha', Y')\|_{p,[s,t]} + \|R^{(\alpha, Y)}\|_{\frac{p}{2},[s,t]})^2 (1 + \|X\|_{p,[s,t]})^2 \\ & \lesssim \|f\|_{C_b^2} (1 + \|\alpha, \alpha'\|_{X,p})^2 (1 + \|Y, Y'\|_{X,p,[s,t]})^2 (1 + \|X\|_{p,[s,t]})^2. \end{aligned}$$

The growth conditions thus hold with constant $C_F = \|f\|_{C_b^2}$ up to a multiplicative constant, which depends on $p, \|\alpha\|_p$ and $\|\alpha, \alpha'\|_{X,p}$.

Proceeding as in the proof of Lemma 3.2, we can show the Lipschitz conditions, observing that

$$|(\alpha, Y)_s - (\alpha, \tilde{Y})_s| + \|(\alpha, Y) - (\alpha, \tilde{Y})\|_{p,[s,t]} = |Y_s - \tilde{Y}_s| + \|Y - \tilde{Y}\|_{p,[s,t]}$$

and

$$\|(\alpha, Y), (\alpha', Y'); (\alpha, \tilde{Y}), (\alpha', \tilde{Y}')\|_{X,p} = \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X,p},$$

and similarly for each summand of the norm, so the Lipschitz conditions hold with constant $C_{F,K,X,X}$, which depends on $p, \|f\|_{C_b^3}, K$, for $K > 0, \|\alpha, \alpha'\|_{X,p}$, and $\|X\|_p$. \square

Proof of Corollary 3.3. (i) The existence and uniqueness of the solution follows immediately from Lemma 3.4 and Theorem 2.4. For the a priori estimate, note that $C_F \lesssim_p \|f\|_{C_b^2} (1 + \|\alpha\|_p + \|\alpha, \alpha'\|_{X,p})^2$.

(ii) To apply the continuity result presented in Theorem 2.5, we need to ensure that the functionals satisfy the required assumptions. For $(F, F'), (\tilde{F}, \tilde{F}')$, this is given in Lemma 3.4. Analogously, since $f - \tilde{f} \in C_b^3$, we note that for

$$(Y, Y') \mapsto ((f - \tilde{f})(\alpha, Y), D(f - \tilde{f})(\alpha, Y)(\alpha', Y'))$$

the growth conditions hold with constant equal to $\|f - \tilde{f}\|_{C_b^2}$ up to a multiplicative constant which depends on p , $\|\alpha\|_p$ and $\|\alpha, \alpha'\|_{X,p}$. Further, it follows from the proofs of [FZ18, Lemma 3.1 and Lemma 3.5] that the growth conditions hold for

$$(Y, Y') \mapsto (f(\alpha, Y) - f(\tilde{\alpha}, Y), Df(\alpha, Y)(\alpha', Y') - Df(\tilde{\alpha}, Y)(\alpha', Y'))$$

with constant equal to $|\alpha_0 - \tilde{\alpha}_0| + \|\alpha - \tilde{\alpha}\|_p + \|\alpha, \alpha'; \tilde{\alpha}, \tilde{\alpha}'\|_{X,p}$ up to a multiplicative constant which depends on p , $\|f\|_{C_b^3}$, $\|\alpha\|_p$, $\|\tilde{\alpha}\|_p$, $\|\alpha, \alpha'\|_{X,p}$, $\|\tilde{\alpha}, \tilde{\alpha}'\|_{X,p}$.

This implies that

$$\begin{aligned} (F - \tilde{F}, F' - \tilde{F}'): \mathcal{V}_X^p([0, T]; \mathbb{R}^k) &\rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \\ ((F - \tilde{F})(Y), (F - \tilde{F})'(Y, Y')) & \\ := (f(\alpha, Y) - \tilde{f}(\tilde{\alpha}, Y), Df(\alpha, Y)(\alpha', Y') - D\tilde{f}(\tilde{\alpha}, Y)(\alpha', Y')), & \end{aligned}$$

satisfies Assumption 2.1 (i). Here, $C_{F-\tilde{F}} = \|f - \tilde{f}\|_{C_b^2} + |\alpha_0 - \tilde{\alpha}_0| + \|\alpha - \tilde{\alpha}\|_p + \|\alpha, \alpha'; \tilde{\alpha}, \tilde{\alpha}'\|_{X,p}$ up to a multiplicative constant, which depends on p , $\|f\|_{C_b^3} \vee \|\tilde{f}\|_{C_b^3}$, $\|\alpha\|_p$, $\|\tilde{\alpha}\|_p$, $\|\alpha, \alpha'\|_{X,p}$, $\|\tilde{\alpha}, \tilde{\alpha}'\|_{X,p}$. \square

3.3. RDEs with discrete time dependence. Let us consider the rough differential equation with discrete time dependence

$$(3.3) \quad Y_t = y_t + \int_0^t f(Y_s, Y_{s \wedge r_1}, \dots, Y_{s \wedge r_\ell}) d\mathbf{X}_s, \quad t \in [0, T],$$

where $\mathbf{X} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ for $p \in (2, 3)$, $(y, y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $f \in C_b^3(\mathbb{R}^{k(\ell+1)}; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, and $r_1 < \dots < r_\ell$ be given time points in $[0, T]$, with $\ell \in \mathbb{N}$. For continuous rough paths as driving signals, the existence (without uniqueness) of a solution to the RDE (3.3) was proven in [Ana23, Example 4.2 and Theorem 4.4]. The next proposition provides an existence, uniqueness and continuity result for RDEs with discrete time dependence driven by càdlàg p -rough paths.

Proposition 3.5.

- (i) *In the above setting, there exists a unique solution to the RDE with discrete time dependence (3.3). Moreover, there exists a componentwise non-decreasing function $K_p: \mathbb{N} \times [0, \infty)^3 \rightarrow [0, \infty)$ such that*

$$\|Y, Y'\|_{X,p} \leq K_p(\ell, \|f\|_{C_b^2}, \|y, y'\|_{X,p}, \|\mathbf{X}\|_p).$$

- (ii) *Let $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ be the unique solution to the RDE with time discrete dependence (3.3). Moreover, let $(\tilde{y}, \tilde{y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ with corresponding solution $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, and suppose that $\|Y, Y'\|_{X,p}, \|\tilde{Y}, \tilde{Y}'\|_{X,p} \leq K$, for some*

$K > 0$. Then, we have the estimate

$$|Y_0 - \tilde{Y}_0| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X,p} \lesssim |y_0 - \tilde{y}_0| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X,p},$$

where the implicit multiplicative constant depends on $p, \ell, \|f\|_{C_b^3}, K$, and $\|\mathbf{X}\|_p$.

Proof. (i) On the interval $[0, r_1]$, we extend the vector field f to map into the space of controlled paths by setting

$$(F, F'): \mathcal{V}_X^p([0, r_1]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, r_1]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \quad (F(Y), F'(Y, Y')) := (f(\bar{Y}), Df(\bar{Y})\bar{Y}'),$$

with $(\bar{Y}, \bar{Y}') = ((Y, Y, \dots, Y), (Y', Y', \dots, Y')) \in \mathcal{V}_X^p([0, T]; \mathbb{R}^{k(\ell+1)})$. It follows analogously to Lemma 3.2 that the functional satisfies Assumption 2.1 with constants depending additionally on ℓ , that is, $C_F = \|f\|_{C_b^2}$ up to a multiplicative constant, which depends on p and ℓ , and $C_{F,K,X,X}$ depends on $p, \ell, \|f\|_{C_b^3}, K$, for $K > 0$, and $\|X\|_p$. Note that it is sufficient to check Assumption 2.1 (ii) for $X = \tilde{X}$. We can thus apply Theorem 2.4 to show that there exists a unique solution to the RDE (3.3) on the interval $[0, r_1]$. We now aim to solve the RDE (3.3) iteratively on the subintervals $[r_i, r_{i+1})$, $i = 1, \dots, \ell$, with $r_{\ell+1} = T$. Given the solution on $[r_{i-1}, r_i)$, with $r_0 = 0$, the value Y_{r_i} is determined by the jump of \mathbf{X} at time r_i . We therefore consider $(y_i, y'_i) \in \mathcal{V}_X^p([r_i, r_{i+1}]; \mathbb{R}^k)$, where

$$y_{i;t} = y_t + Y_{r_i-} - y_{r_i-} + F_{r_i-}(Y)X_{r_i-,r_i} + F'_{r_i-}(Y, Y')\mathbb{X}_{r_i-,r_i}, \quad t \in [r_i, r_{i+1}],$$

for every $i = 1, \dots, \ell$, and $(\alpha_i, \alpha'_i) \in \mathcal{V}_X^p([r_i, r_{i+1}]; \mathbb{R}^{ik})$ be a fixed controlled path, with $\alpha_{i,t} = (Y_{r_1}, \dots, Y_{r_i})$, $t \in [r_i, r_{i+1})$. We set

$$\begin{aligned} (F, F'): \mathcal{V}_X^p([r_i, r_{i+1}]; \mathbb{R}^k) &\rightarrow \mathcal{V}_X^p([r_i, r_{i+1}]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \\ (F(Y), F'(Y, Y')) &= (f((Y, \alpha_i, Y, \dots, Y)), Df((Y, \alpha_i, Y, \dots, Y))(Y', \alpha'_i, Y', \dots, Y')) \end{aligned}$$

for $i = 1, \dots, \ell - 1$, and

$$(F(Y), F'(Y, Y')) = (f((Y, \alpha_\ell)), Df((Y, \alpha_\ell))(Y', \alpha'_\ell)),$$

for $i = \ell$. Analogously to Lemma 3.4, we can show that the functional satisfies Assumption 2.1 with constants depending additionally on ℓ , that is $C_F = \|f\|_{C_b^2}$ up to a multiplicative constant, which depends on p and ℓ , see the definition of $(\alpha_i, \alpha'_i) \in \mathcal{V}_X^p([r_i, r_{i+1}])$, and $C_{F,K,X,X}$ depends on $p, \ell, \|f\|_{C_b^3}, K$, for $K > 0$, and $\|X\|_p$. Note that it is again sufficient to check Assumption 2.1 (ii) for $X = \tilde{X}$. We can thus again apply Theorem 2.4 to show that there exists a unique solution to the RDE (3.3) on the interval $[r_i, r_{i+1})$, that is

$$Y_t = y_{i;t} + \int_{r_i}^t F_s(Y) d\mathbf{X}_s, \quad t \in [r_i, r_{i+1}),$$

for every $i = 1, \dots, \ell$. Then, by pasting the solutions on each of these subintervals together, we obtain a unique global solution Y , which holds over the entire interval $[0, T]$.

The a priori estimate follows by iteratively combining the a priori estimate of Corollary 3.3, noting that $\alpha_{i,t} = (Y_{r_1}, \dots, Y_{r_i})$, $t \in [r_i, r_{i+1})$, for $i = 1, \dots, \ell$.

(ii) *Local estimate on $[0, r_1]$.* To apply the continuity result presented in Theorem 2.5 on the subinterval $[0, r_1]$, we need to ensure that the functionals satisfy the required assumptions.

For (F, F') , this is shown in the proof of (i), and as we aim to obtain continuity of the solution map as a function of the initial condition (y, y') , not the vector field f , on the interval

$[0, r_1]$ we may consider $(F, F') = (\tilde{F}, \tilde{F}')$, so, $(F - \tilde{F}, F' - \tilde{F}') = 0$. Theorem 2.5 now gives that

$$\|Y' - \tilde{Y}'\|_{p, [0, r_1]} + \|R^Y - R^{\tilde{Y}}\|_{p, [0, r_1]} \lesssim |y_0 - \tilde{y}_0| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, p, [0, r_1]},$$

where the implicit multiplicative constant depends on $p, \ell, \|f\|_{C_b^3}, K$, and $\|\mathbf{X}\|_p$.

Local estimate on $[r_i, r_{i+1}]$, $i = 1, \dots, \ell$. To apply the continuity result presented in Theorem 2.5, we need to ensure that the functionals satisfy the required assumptions. For $(F, F'), (\tilde{F}, \tilde{F}')$, this is shown in the proof of (i), and for $(F - \tilde{F}, F' - \tilde{F}')$, in the proof of Corollary 3.3 (ii), where the constant $C_{F-\tilde{F}}$ depends additionally on ℓ . Theorem 2.5 then implies that

$$\|Y' - \tilde{Y}'\|_{p, [r_i, r_{i+1}]} + \|R^Y - R^{\tilde{Y}}\|_{p, [r_i, r_{i+1}]} \lesssim |Y_{r_i} - \tilde{Y}_{r_i}| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, p, [r_i, r_{i+1}]},$$

where the implicit multiplicative constant depends on $p, \|f\|_{C_b^3}, K$ and $\|\mathbf{X}\|_p$, see the definition of $(\alpha_i, \alpha'_i), (\tilde{\alpha}_i, \tilde{\alpha}'_i) \in \mathcal{V}_X^p([r_i, r_{i+1}]; \mathbb{R}^{ik}), (y_i, y'_i), (\tilde{y}_i, \tilde{y}'_i) \in \mathcal{V}_X^p([r_i, r_{i+1}]; \mathbb{R}^k)$.

Global estimate. Using the methods of the proof of Theorem 2.5, and applying the local estimates on the subintervals $[r_i, r_{i+1}]$, $i = 0, 1, \dots, \ell$, one can then derive that

$$\|Y' - \tilde{Y}'\|_p + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2}} \lesssim |y_0 - \tilde{y}_0| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X, p},$$

which implies the estimate. \square

3.4. RDEs with constant delay. Maybe the most prominent example of rough functional differential equations are RDEs with constant delays, cf. e.g. [FR06, NNT08, BMCR14, CE20, BBR20]. In the present subsection we consider the delayed rough differential equation

$$(3.4) \quad Y_t = y_t + \int_0^t f(Y_s, Y_{s-r_1}, \dots, Y_{s-r_\ell}) d\mathbf{X}_s, \quad t \in [0, T],$$

where $\mathbf{X} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ for $p \in (2, 3)$, $(y, y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $f \in C_b^3(\mathbb{R}^{k(\ell+1)}; \mathcal{L}(\mathbb{R}^d, \mathbb{R}^k))$, and constant delays $0 < r_1 < \dots < r_\ell$ with $\ell \in \mathbb{N}$. To give a rigorous mathematical meaning to the RDE (3.4), we follow the approach of Neuenkirch, Nourdin and Tindel [NNT08]: we assume that the driving rough path \mathbf{X} is of the form

$$X_t = (Z_t, Z_{t-r_1}, \dots, Z_{t-r_\ell}), \quad t \in [0, T],$$

for a path $Z \in D^p([-r_\ell, T]; \mathbb{R}^e)$ with $d = e(\ell + 1)$. We extend the vector field f to map into the space of controlled paths by setting

$$(F(Y), F'(Y, Y')) := (f((Y, \alpha)), Df((Y, \alpha))(Y', \alpha')),$$

for $(\alpha, \alpha') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^{k\ell})$, where

$$(3.5) \quad \alpha = (\alpha_1, \dots, \alpha_\ell) \quad \text{with} \quad \alpha_{j,t} := \begin{cases} Y_{t-r_j}, & t \in [r_j, T] \\ Y_{j;t}, & t \in [0, r_j] \end{cases},$$

for fixed controlled paths $(Y_j, Y'_j) \in \mathcal{V}_{Z_{\cdot-r_j}}^p([0, T]; \mathbb{R}^k)$ and every $j = 1, \dots, \ell$. This includes the natural case $Y_j = \xi_{\cdot-r_j}$ for an initial path $\xi \in \mathcal{V}_Z^p([-r_\ell, T]; \mathbb{R}^k)$.

Note that the postulated form of the rough path \mathbf{X} is essential to ensure the well-posedness of the rough integral appearing in (3.4) and the extension of the solution Y to the interval $[-r_\ell, 0]$ is a standard and necessary way to give a meaning to $f(Y_s, Y_{s-r_1}, \dots, Y_{s-r_\ell})$ on the entire interval $[0, T]$.

For delayed RDEs of the form (3.4) driven by α -Hölder continuous rough paths, existence, uniqueness and continuity of the Itô–Lyons map were first proven in [NNT08] for $\alpha \in (\frac{1}{3}, \frac{1}{2})$. These results were extended in [TT12] to α -Hölder continuous rough paths for $\alpha \in (\frac{1}{4}, \frac{1}{3})$. A paracontrolled distribution approach to RDEs with constant delay can be found in [PT21]. Based on the general results of Section 2, we can derive the follow proposition.

Proposition 3.6.

- (i) *In the above setting, there exists a unique solution to the delayed RDE (3.4). Moreover, there exists a componentwise non-decreasing function $K_p: \mathbb{N} \times [0, \infty)^4 \rightarrow [0, \infty)$ such that*

$$\|Y, Y'\|_{X,p} \leq K_p \left(\ell, \|f\|_{C_b^2}, \|y, y'\|_{X,p}, \sum_{j=1}^{\ell} \|Y_j, Y'_j\|_{Z_{-r_j,p}}, \|\mathbf{X}\|_p \right).$$

- (ii) *Let $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ be the unique solution to the rough differential equation with constant delay (3.4). Moreover, consider $(\tilde{y}, \tilde{y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, and fixed controlled paths $(\tilde{Y}_j, \tilde{Y}'_j) \in \mathcal{V}_{Z_{-r_j}}^p([0, T]; \mathbb{R}^k)$, $j = 1, \dots, \ell$, with corresponding solution $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$.*

Suppose that $\|Y, Y'\|_{X,p}, \|\tilde{Y}, \tilde{Y}'\|_{X,p} \leq K$, for some $K > 0$, and $\|Y_j\|_p, \|\tilde{Y}_j\|_p, \|Y_j, Y'_j\|_{X,p}, \|\tilde{Y}_j, \tilde{Y}'_j\|_{X,p} \leq L$, for some $L > 0$, $j = 1, \dots, \ell$. Then, we have the estimate

$$\begin{aligned} & |Y_0 - \tilde{Y}_0| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X,p} \\ & \lesssim |y_0 - \tilde{y}_0| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X,p} + \sum_{j=1}^{\ell} |Y_{j;0} - \tilde{Y}_{j;0}| + \sum_{j=1}^{\ell} \|Y_j, Y'_j; \tilde{Y}_j, \tilde{Y}'_j\|_{Z_{-r_j,p}}, \end{aligned}$$

where the implicit multiplicative constant depends on $p, \ell, r_1, T, \|f\|_{C_b^3}, K, L$, and $\|\mathbf{X}\|_p$.

Proof. (i) The existence and uniqueness of the solution follows by iteratively applying Corollary 3.3 (i) to intervals of the length r_1 .

More precisely, we consider the functional $(F, F'): \mathcal{V}_X^p([0, r_1]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, r_1]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, for $(\alpha, \alpha') \in \mathcal{V}_X^p([0, r_1]; \mathbb{R}^{k\ell})$ given by (3.5), and apply Corollary 3.3 (i) to show that there exists a unique solution to the RDE (3.4) on the interval $[0, r_1)$.

We now aim to solve the RDE (3.4) iteratively on the subintervals $[ir_1, (i+1)r_1]$, $i = 1, \dots, N-1$, assuming that $T = Nr_1$ for some $N \in \mathbb{N}$. Given the solution on $[(i-1)r_1, ir_1)$, the value Y_{ir_1} is determined by the jump of \mathbf{X} on ir_1 . We therefore consider $(y_i, y'_i) \in \mathcal{V}_X^p([ir_1, (i+1)r_1]; \mathbb{R}^k)$, where

$$y_{i;t} = y_t + Y_{ir_1-} - y_{ir_1-} + F_{ir_1-}(Y)X_{ir_1-,ir_1} + F'_{ir_1-}(Y, Y')\mathbb{X}_{ir_1-,ir_1}, \quad t \in [ir_1, (i+1)r_1],$$

for every $i = 1, \dots, N-1$, and

$$(F, F'): \mathcal{V}_X^p([ir_1, (i+1)r_1]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([ir_1, (i+1)r_1]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)),$$

for $(\alpha, \alpha') \in \mathcal{V}_X^p([ir_1, (i+1)r_1]; \mathbb{R}^{k\ell})$ given by (3.5). We again apply Corollary 3.3 (i) to show that there exists a unique solution to the RDE (3.4) on the interval $[ir_1, (i+1)r_1)$, that is

$$Y_t = y_{i;t} + \int_{ir_1}^t F_s(Y) d\mathbf{X}_s, \quad t \in [ir_1, (i+1)r_1)$$

for every $i = 1, \dots, N-1$. Then, by pasting solutions on each of these subintervals together, we obtain a unique global solution Y to the RDE (3.4), which holds over the interval $[0, T]$.

The a priori estimate follows by iteratively combining the a priori estimate of Corollary 3.3, and by the definition of α in (3.5).

(ii) *Local estimate on $[ir_1, (i+1)r_1]$, $i = 0, \dots, N-1$.* To apply the continuity result presented in Theorem 2.5 on the subintervals $[ir_1, (i+1)r_1]$, we need to ensure that the functionals satisfy the required assumptions. This is given for $(F, F'), (\tilde{F}, \tilde{F}')$ in Lemma 3.4, and for $(F - \tilde{F}, F' - \tilde{F}')$ we refer to the proof of Corollary 3.3 (ii), and write $C_{F-\tilde{F},i}$ for the corresponding constant. By Theorem 2.5, it then holds the estimate

$$\begin{aligned} & \|Y' - \tilde{Y}'\|_{p,[ir_1,(i+1)r_1]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2},[ir_1,(i+1)r_1]} \\ & \lesssim |Y_{ir_1} - \tilde{Y}_{ir_1}| + |F_{ir_1}(Y) - \tilde{F}_{ir_1}(\tilde{Y})| + \|y, y'; \tilde{y}, \tilde{y}'\|_{p,[ir_1,(i+1)r_1]} + C_{F-\tilde{F},i} \\ & \lesssim |y_0 - \tilde{y}_0| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X,p} + \sum_{j=1}^{\ell} |Y_{j;0} - \tilde{Y}_{j;0}| + \|Y_j, Y'_j; \tilde{Y}_j, \tilde{Y}'_j\|_{Z_{-\tau_j},p} \\ & \quad + \|Y' - \tilde{Y}'\|_{p,[0,ir_1]} + \|R^Y - R^{\tilde{Y}}\|_{\frac{p}{2},[0,ir_1]}, \end{aligned}$$

where the implicit multiplicative constant depends on $p, \ell, \|f\|_{C_b^2}, K, L$, and $\|\mathbf{X}\|_p$, see the definition of $(\alpha, \alpha'), (\tilde{\alpha}, \tilde{\alpha}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^{k\ell})$.

Global estimate. Iteratively applying the local estimates and, as before, using that $\|\cdot\|_p \leq N \sum_{i=0}^{N-1} \|\cdot\|_{p,[ir_1,(i+1)r_1]}$, one can then derive the estimate. \square

3.5. RDEs with variable delay. Rough differential equations with variable delays represent a slight generalization of RDEs with constant delays. More precisely, let us consider the rough differential equation with variable delay

$$(3.6) \quad Y_t = y_t + \int_0^t f(Y_s, Y_{s-\eta(s)}) d\mathbf{X}_s, \quad t \in [0, T],$$

where $\mathbf{X} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ for $p \in (2, 3)$, $(y, y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $f \in C_b^3(\mathbb{R}^{2k}; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, and $\eta(\cdot)$ be a bounded continuous function with $\eta(t) \geq \varepsilon$, $t \in [0, T]$, for some $\varepsilon > 0$, and $\bar{\eta} = \sup\{\eta(t) - t : t \in [0, T]\}$. We assume that the driving rough path \mathbf{X} is of the form

$$X_t = (Z_t, Z_{t-\eta(t)}), \quad t \in [0, T],$$

for a path $Z \in D^p([-\bar{\eta}, T]; \mathbb{R}^e)$ with $d = 2e$. We extend the vector field f into the space of controlled paths by setting

$$\begin{aligned} (F, F') &: \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k)), \\ (F(Y), F'(Y, Y')) &= (f((Y, \alpha)), Df((Y, \alpha))(Y', \alpha')), \end{aligned}$$

for $(\alpha, \alpha') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, where

$$\alpha_t := \begin{cases} Y_{t-\eta(t)}, & t \geq \eta(t) \\ Y_{\eta;t}, & t < \eta(t) \end{cases},$$

for some fixed controlled path $(Y_\eta, Y'_\eta) \in \mathcal{V}_{Z_{-\eta(\cdot)}}^p([0, T]; \mathbb{R}^k)$.

Corollary 3.7.

- (i) In the above setting, there exists a unique solution to the delayed RDE (3.6). Moreover, there exists a componentwise non-decreasing function $K_p: (0, \infty) \times [0, \infty)^4 \rightarrow [0, \infty)$ such that

$$\|Y, Y'\|_{X,p} \leq K_p(\varepsilon^{-1}, \|f\|_{C_b^2}, \|y, y'\|_{X,p}, \|Y_\eta, Y'_\eta\|_{X,p}, \|\mathbf{X}\|_p).$$

- (ii) Let $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$ be the unique solution to the RDE with variable delay (3.6). Moreover, consider $(\tilde{y}, \tilde{y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, and a fixed controlled path $(\tilde{Y}_\eta, \tilde{Y}'_\eta) \in \mathcal{V}_{Z_{-\eta(\cdot)}}^p([0, T]; \mathbb{R}^k)$ with corresponding solution $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$.

Suppose that $\|Y, Y'\|_{X,p}, \|\tilde{Y}, \tilde{Y}'\|_{X,p} \leq K$, for some $K > 0$, and $\|Y_\eta, Y'_\eta\|_{X,p}, \|\tilde{Y}_\eta, \tilde{Y}'_\eta\|_{X,p} \leq L$, for some $L > 0$. Then, we have the estimate

$$\begin{aligned} & |Y_0 - \tilde{Y}_0| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X,p} \\ & \lesssim |y_0 - \tilde{y}_0| + \|y, y'; \tilde{y}, \tilde{y}'\|_{X,p} + |Y_{\eta;0} - \tilde{Y}_{\eta;0}| + \|Y_\eta, Y'_\eta; \tilde{Y}_\eta, \tilde{Y}'_\eta\|_{Z_{-\eta(\cdot)}, p}, \end{aligned}$$

where the implicit multiplicative constant depends on $p, \varepsilon, T, \eta, \|f\|_{C_b^3}, K, L$, and $\|\mathbf{X}\|_p$.

Proof. (i) The existence and uniqueness of the solution follows by iteratively applying Corollary 3.3 (i) to intervals of the length ε , see the proof of Proposition 3.6 (i). The a priori estimate follows analogously.

(ii) The continuity of the solution map follows analogously to Proposition 3.6 (ii). \square

4. APPLICATION TO STOCHASTIC DIFFERENTIAL EQUATIONS WITH DELAY

One main application of rough path theory is a pathwise and robust approach to stochastic differential equations, see e.g. [FH20]. In this section we show how a càdlàg martingale and its delayed version can be lifted to a joint random rough path in the spirit of stochastic Itô integration. Consequently, this allows to apply the results on rough functional differential equations, provided in Section 2, to Itô stochastic differential equations (SDEs) with constant delays.

Throughout the entire section, let us consider constant delays $0 < r_1 < \dots < r_\ell$ with $\ell \in \mathbb{N}$, and let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space with a complete and right-continuous filtration $(\mathcal{F}_t)_{t \in [-r_\ell, T]}$. Let $Z = (Z_t)_{t \in [-r_\ell, T]}$ be an e -dimensional square-integrable càdlàg martingale that is defined on $(\Omega, \mathcal{F}, \mathbb{P})$, with $Z_t = 0$ for $t < 0$. The space of all square-integrable random variables on $(\Omega, \mathcal{F}, \mathbb{P})$ is denoted by L^2 and equipped with the standard L^2 -norm.

4.1. Delayed martingales as rough paths. The aim of this subsection is to construct a random rough path above the stochastic process $X = (X_t)_{t \in [0, T]}$, defined as

$$X_t := (Z_t, Z_{t-r_1}, \dots, Z_{t-r_\ell}), \quad t \in [0, T],$$

in the spirit of stochastic Itô integration. Notice, although Z is a martingale, the stochastic process X is, in general, not a martingale anymore and, thus, we cannot directly apply stochastic Itô integration to define a rough path lift of X . Moreover, if Z is a semimartingale, it seems not obvious under which conditions the associated process X is again a semimartingale.

As a first step to construct a random rough path above the stochastic process X , in the next lemma, we derive the existence of an auxiliary process, inspired by the quadratic co-variation of martingales.

Lemma 4.1. *Let $Z = (Z_t)_{t \in [-r_\ell, T]}$ be an e -dimensional square-integrable càdlàg martingale that is defined on $(\Omega, \mathcal{F}, \mathbb{P})$, with $Z_t = 0$ for $t < 0$. Then, for $i_1, i_2 = 1, \dots, e$, $j_1, j_2 = 0, \dots, \ell$, $j_1 \neq j_2$, we have*

$$\mathbb{E} \left[\sup_{t \in [0, T]} \left| \sum_{k=0}^{N_n-1} Z_{t_k^n \wedge t - r_{j_1}, t_{k+1}^n \wedge t - r_{j_1}}^{i_1} Z_{t_k^n \wedge t - r_{j_2}, t_{k+1}^n \wedge t - r_{j_2}}^{i_2} - \sum_{s \leq t} \Delta_s Z_{\cdot - r_{j_1}}^{i_1} \Delta_s Z_{\cdot - r_{j_2}}^{i_2} \right|^2 \right] \rightarrow 0$$

along any sequence of partitions $\mathcal{P}^n = \{0 = t_0^n < t_1^n < \dots < t_{N_n}^n = T\}$, $n \in \mathbb{N}$, of the interval $[0, T]$ with vanishing mesh size, so that $|\mathcal{P}^n| \rightarrow 0$ as $n \rightarrow \infty$. Here, we write $\Delta_t H = H_{t-, t}$, with $H_{t-} = \lim_{s \uparrow t} H_s$, for the jump of a stochastic process H at time t .

We define the stochastic process

$$[Z_{\cdot - r_{j_1}}^{i_1}, Z_{\cdot - r_{j_2}}^{i_2}]_t := \sum_{s \leq t} \Delta_s Z_{\cdot - r_{j_1}}^{i_1} \Delta_s Z_{\cdot - r_{j_2}}^{i_2}, \quad t \in [0, T].$$

This process is càdlàg and has almost surely finite $\frac{p}{2}$ -variation, that is, $[Z_{\cdot - r_{j_1}}^{i_1}, Z_{\cdot - r_{j_2}}^{i_2}] \in D^{\frac{p}{2}}([0, T]; \mathbb{R})$ \mathbb{P} -almost surely.

Proof. We assume w.l.o.g. that $j_1 = 0$, and write $r = r_{j_2}$, and $M = Z^{i_1}$, $\tilde{M} = Z^{i_2}$. For $n \in \mathbb{N}$, we define

$$H_t^n := \sum_{k=0}^{N_n-1} \tilde{M}_{t_k^n - r, t_{k+1}^n - r} \mathbf{1}_{(t_k^n, t_{k+1}^n]}(t), \quad t \in [0, T],$$

and note that for $|\mathcal{P}^n| < r$, H^n is indeed a simple predictable process, see [Pro05, Chapter II]. The Itô integral is then given by

$$\int_0^t H_s^n dM_s = \sum_{k=0}^{N_n-1} \tilde{M}_{t_k^n \wedge t - r, t_{k+1}^n \wedge t - r} M_{t_k^n \wedge t, t_{k+1}^n \wedge t}.$$

We now aim to show that

$$(4.1) \quad \mathbb{E} \left[\int_0^T (H_s^n - H_s)^2 d[M]_s \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

where $H := \Delta \tilde{M}_{\cdot - r}$, and $[\cdot]$ denotes the quadratic variation. Using the localizing sequence $\tau_m = T \wedge \inf\{t : |\tilde{M}_t| \geq m\}$, $m \in \mathbb{N}$, and replacing H^n by $H_{\cdot \wedge \tau_m}^n$ and H by $H_{\cdot \wedge \tau_m}$, we may assume that the integrand is uniformly bounded, so that we can apply the dominated convergence theorem. Since $H^n \rightarrow H$ converges pointwise as $n \rightarrow \infty$, this shows (4.1).

By [JS03, Chapter I.4], it follows that

$$\mathbb{E} \left[\sup_{t \in [0, T]} \left| \int_0^t H_s^n dM_s - \int_0^t H_s dM_s \right|^2 \right] \rightarrow 0$$

as $n \rightarrow \infty$, thus uniformly in L^2 , and as

$$\int_0^t H_s dM_s = \sum_{s \leq t} \Delta_s \tilde{M}_{\cdot - r} \Delta_s M,$$

this implies the convergence result. Further,

$$4[M, \tilde{M}_{\cdot - r}] = \sum_{s \leq \cdot} (\Delta_s M + \Delta_s \tilde{M}_{\cdot - r})^2 - \sum_{s \leq \cdot} (\Delta_s M - \Delta_s \tilde{M}_{\cdot - r})^2$$

has càdlàg sample paths of finite variation, as both terms on the right hand side are monotonically increasing, which implies that $[M, \tilde{M}_{\cdot-r}]$ has almost surely finite $\frac{p}{2}$ -variation, and concludes the proof. \square

Proposition 4.2. *Let $p \in (2, 3)$, and let $Z = (Z_t)_{t \in [-r, T]}$ be an e -dimensional square-integrable càdlàg martingale that is defined on $(\Omega, \mathcal{F}, \mathbb{P})$, with $Z_t = 0$ for $t < 0$. We set $X = (Z, Z_{\cdot-r_1}, \dots, Z_{\cdot-r_\ell})$. Then, X can be lifted to a random rough path, by defining $\mathbf{X} = (X, \mathbb{X}) \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$, \mathbb{P} -almost surely, with $d = e(\ell + 1)$, where*

$$\mathbb{X}_{s,t}^{ij} := \int_s^t X_{u-}^i dX_u^j - X_s^i X_{s,t}^j := \int_0^t X_{u-}^i dX_u^j - \int_0^s X_{u-}^i dX_u^j - X_s^i X_{s,t}^j,$$

for $i, j = 1, \dots, d$ with $i = j$ and $i > j$ such that $X^i = Z_{\cdot-r_{j_1}}^{i_1}$, $X^j = Z_{\cdot-r_{j_2}}^{i_2}$ with $i_1, i_2 = 0, \dots, e$, $j_1, j_2 = 0, \dots, \ell$, $j_1 > j_2$, and else

$$\mathbb{X}_{s,t}^{ji} = -\mathbb{X}_{s,t}^{ij} + X_{s,t}^i X_{s,t}^j - [X^i, X^j]_{s,t}$$

for any $(s, t) \in \Delta_T$, and where the integration is defined as a stochastic Itô integral, and $[X^i, X^j]$ is defined in Lemma 4.1.

Remark 4.3. *The stochastic integral $\int_0^t Z_{u-r_{j_1}}^{i_1} dZ_{u-r_{j_2}}^{i_2}$ can be defined using classical stochastic Itô integration if $r_{j_1} > r_{j_2}$. Indeed, the stochastic process $(Z_{t-r_{j_2}}^{i_2})_{t \in [0, T]}$ is a martingale and the stochastic process $(Z_{t-r_{j_1}}^{i_1})_{t \in [0, T]}$ is predictable, both with respect to the filtration $(\mathcal{F}_{t-r_{j_2}})_{t \in [0, T]}$ with $\mathcal{F}_t := \{\Omega, \emptyset\}$ for $t < 0$.*

Further, for $i \geq j$, we have that $\mathbb{X}_{s,t}^{ij} := \int_0^t X_{u-}^i dX_u^j - \int_0^s X_{u-}^i dX_u^j - X_s^i X_{s,t}^j = \int_s^t X_{s,u-}^i dX_u^j = \int_s^t \int_s^{u-} dX_r^i dX_u^j$, that is, $\mathbb{X}_{s,t}^{ij}$ coincides with the 2-fold iterated integral, for $(s, t) \in \Delta_T$.

Proof of Proposition 4.2. First, by definition Chen's relation does hold: Let $0 \leq s \leq v \leq t \leq T$. Then, we have that

$$\begin{aligned} & \mathbb{X}_{s,v}^{ii} + \mathbb{X}_{v,t}^{ii} + X_{s,v}^i X_{v,t}^i \\ &= \int_s^t X_{u-}^i dX_u^i - X_s^i X_{s,v}^i - X_v^i X_{v,t}^i + X_{s,v}^i X_{v,t}^i \\ &= \int_s^t X_{u-}^i dX_u^i - X_s^i X_{s,t}^i \\ &= \mathbb{X}_{s,t}^{ii}, \end{aligned}$$

similarly for \mathbb{X}^{ij} , and

$$\begin{aligned} & \mathbb{X}_{s,v}^{ji} + \mathbb{X}_{v,t}^{ji} + X_{s,v}^j X_{v,t}^i \\ &= -\mathbb{X}_{s,v}^{ij} + X_{s,v}^i X_{s,v}^j - [X^i, X^j]_{s,v} - \mathbb{X}_{v,t}^{ij} + X_{v,t}^i X_{v,t}^j - [X^i, X^j]_{v,t} + X_{s,v}^j X_{v,t}^i \\ &= -\mathbb{X}_{s,t}^{ij} + X_{s,v}^i X_{v,t}^j + X_{s,v}^i X_{s,v}^j + X_{v,t}^i X_{v,t}^j + X_{v,t}^i X_{s,v}^j - [X^i, X^j]_{s,t} \\ &= -\mathbb{X}_{s,t}^{ij} + X_{s,t}^i X_{s,t}^j - [X^i, X^j]_{s,t} \\ &= \mathbb{X}_{s,t}^{ji}. \end{aligned}$$

Further, Z has almost surely finite p -variation, see e.g. [Lép76], therefore $X \in D^p([0, T]; \mathbb{R}^d)$ \mathbb{P} -almost surely. Since the maps $s \mapsto \mathbb{X}_{s,t}$ for fixed t , and $t \mapsto \mathbb{X}_{s,t}$ for fixed s are both càdlàg, it thus remains to show that $\|\mathbb{X}\|_{\frac{p}{2}} < \infty$ \mathbb{P} -almost surely.

We define the dyadic stopping times $(\tau_k^n)_{n,k \in \mathbb{N}}$ by

$$t_0^n := 0, \quad \tau_{k+1}^n := \inf\{t \geq \tau_k^n : |X_t - X_{\tau_k^n}| \geq 2^{-n}\} \wedge T.$$

For $t \in [0, T]$ and $n \in \mathbb{N}$ we introduce the dyadic approximation

$$X_t^n := \sum_{k=0}^{\infty} X_{\tau_k^n} \mathbf{1}_{(\tau_k^n, \tau_{k+1}^n]}(t) \quad \text{and} \quad \int_0^t X_r^{i,n} dX_r^j := \sum_{k=0}^{\infty} X_{\tau_k^n}^i X_{\tau_k^n \wedge t, \tau_{k+1}^n \wedge t}^j,$$

for $i = j$ or $i \neq j$ such that $X^i = Z_{\cdot - r_{j_1}}^{i_1}$, $X^j = Z_{\cdot - r_{j_2}}^{i_2}$ for $i_1, i_2 = 1, \dots, e$, $j_1, j_2 = 0, \dots, \ell$, $j_1 > j_2$.

We now show that for almost every $\omega \in \Omega$, for every $t \in [0, T]$ and for every $\varepsilon \in (0, 1)$, there exists a constant $C = C(\omega, \varepsilon)$ such that for all $n \in \mathbb{N}$, we have

$$(4.2) \quad \left| \left(\int_0^t X_u^{i,n} dX_u^j - \int_0^t X_u^i dX_u^j \right) (\omega) \right| \leq C 2^{-n(1-\varepsilon)}.$$

Applying the Burkholder–Davis–Gundy inequality, we have that

$$\mathbb{E} \left[\left(\sup_{t \in [0, T]} \int_0^t (X_u^{i,n} - X_u^i) dX_u^j \right)^2 \right] \lesssim \mathbb{E} \left[\int_0^T (X_u^{i,n} - X_u^i)^2 d[X^j]_u \right] \lesssim 2^{-2n}, \quad n \in \mathbb{N},$$

where the implicit multiplicative constant depends on the quadratic variation $[X^j]$ of X^j . Combining this with Chebyshev's inequality, we obtain for any $\varepsilon \in (0, 1)$ that

$$\mathbb{P} \left(\left| \int_0^t (X_u^{i,n} - X_u^i) dX_u^j \right| \geq 2^{-n(1-\varepsilon)} \right) \lesssim 2^{2n(1-\varepsilon)} 2^{-2n} = 2^{-n\varepsilon}.$$

So by the Borel–Cantelli lemma, we have that

$$\sup_{t \in [0, T]} \left(\int_0^t X_u^{i,n} dX_u^j - \int_0^t X_u^i dX_u^j \right) \lesssim 2^{-n(1-\varepsilon)},$$

where the implicit multiplicative constant is a random variable which does not depend on n , which shows (4.2). Proceeding as in the proof of [LP18, Theorem 3.1], we can show that $\|\mathbb{X}^{ij}\|_{\frac{p}{2}} < \infty$ \mathbb{P} -almost surely. Further, let $i \neq j$ as above, then we have for any $(s, t) \in \Delta_T$ that

$$\|\mathbb{X}_{s,t}^{ji}\|_{\frac{p}{2}} \lesssim \|\mathbb{X}^{ij}\|_{\frac{p}{2}, [s,t]} + \|X\|_{p, [s,t]}^p + \|[X^i, X^j]\|_{\frac{p}{2}, [s,t]}.$$

Lemma 4.1 then ensures that $\|\mathbb{X}^{ji}\|_{\frac{p}{2}} < \infty$ \mathbb{P} -almost surely. \square

Remark 4.4. For a fractional Brownian motion with Hurst index H and its delayed version a joint rough path was constructed in [NNT08] based on the Russo–Vallois integral [RV93], assuming that $H > \frac{1}{3}$. This construction was generalized in [TT12] allowing for $H > \frac{1}{4}$. A related construction of a delayed rough path above a fractional Brownian motion can be found in [BMCR14]. For a standard Brownian motion a delayed rough path was also defined in [CE20] based on stochastic Itô integration. While the delayed rough path provided in Proposition 4.2 corresponds to stochastic Itô integration, see Proposition 4.5 below, the aforementioned constructions of delayed rough paths above (fractional) Brownian motion correspond to Stratonovich integration.

4.2. SDEs with delays as random RDEs. Let us consider the SDE with constant delays

$$(4.3) \quad \begin{aligned} Y_t &= y_0 + \int_0^t f(Y_{s-}, Y_{s-r_1-}, \dots, Y_{s-r_\ell-}) dZ_s, \quad t \in [0, T], \\ Y_t &= y_t, \quad t \in [-r_\ell, 0), \end{aligned}$$

where $y \in D^{\frac{k}{2}}([-r_\ell, T]; \mathbb{R}^k)$, $f \in C_b^3(\mathbb{R}^{k(\ell+1)}; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ and the integral is defined as a stochastic Itô integral. For a comprehensive introduction to stochastic Itô integration and SDEs we refer, e.g., to the textbook [Pro05]. It is known that the SDE (4.3) possesses a unique (strong) solution, see e.g. [Pro05, Chapter V, Theorem 7]. It turns out that the solutions to the SDE (4.3) and to the RDE (3.4) driven by the random rough path $\mathbf{X} = (X, \mathbb{X})$, with \mathbb{X} as defined in Proposition 4.2, coincide \mathbb{P} -almost surely.

Proposition 4.5. *Let $p \in (2, 3)$, and let $Z = (Z_t)_{t \in [-r, T]}$ be an e -dimensional square-integrable càdlàg martingale, that is defined on $(\Omega, \mathcal{F}, \mathbb{P})$, with $Z_t = 0$ for $t < 0$. We set $X := (Z, Z_{-r_1}, \dots, Z_{-r_\ell})$, and let $\mathbf{X} = (X, \mathbb{X})$ be the random rough path, with \mathbb{X} defined as in Proposition 4.2.*

- (i) *Let (V, V') be an adapted stochastic process such that $(V(\omega), V'(\omega)) \in \mathcal{V}_{X(\omega)}^p$ for almost every $\omega \in \Omega$, and assume that $\sup_{t \in [0, T]} |V'(\omega)|_t$ is uniformly bounded in ω . Then, the rough integral exists and coincides \mathbb{P} -almost surely with the stochastic Itô integral, that is*

$$\int_0^T V_u d\mathbf{X}_u = \int_0^T V_{u-} dX_u, \quad \mathbb{P}\text{-almost surely.}$$

- (ii) *The solution of the SDE (4.3) driven by X , and the solution of the RDE (3.4) driven by \mathbf{X} , coincide \mathbb{P} -almost surely.*

Proof. (i) *Step 1.* [FS17, Theorem 31] gives that

$$\int_0^T V_u d\mathbf{X}_u = \lim_{|\mathcal{P}| \rightarrow 0} \sum_{(s,t) \in \mathcal{P}} (V_s X_{s,t} + V'_s \mathbb{X}_{s,t}),$$

where the limit is taken over any sequence of partitions \mathcal{P} of the interval $[0, T]$ with mesh size $|\mathcal{P}| \rightarrow 0$, and it is known that

$$\sum_{(s,t) \in \mathcal{P}} V_s X_{s,t} \rightarrow \int_0^T V_{u-} dX_u,$$

in probability as $|\mathcal{P}| \rightarrow 0$, see e.g. [Pro05, Chapter II, Theorem 21], therefore the convergence holds \mathbb{P} -almost surely, possibly along some subsequence.

Step 2. Since V' is assumed to be bounded, we are left to show that for $i, j = 1, \dots, d$,

$$(4.4) \quad \sup_{\tau \in [0, T]} \left| \sum_{(s,t) \in \mathcal{P} \cap [0, \tau]} \mathbb{X}_{s,t}^{ij} \right| \rightarrow 0, \quad \text{as } |\mathcal{P}| \rightarrow 0,$$

in probability, which then implies \mathbb{P} -almost sure convergence, along some subsequence. From here on, we take the sequence of partitions $\mathcal{P}^n = \{0 = t_0^n < t_1^n < \dots < t_{N_n}^n = T\}$, $n \in \mathbb{N}$, of the interval $[0, T]$ with vanishing mesh size, so that $|\mathcal{P}^n| \rightarrow 0$ as $n \rightarrow \infty$. Moreover, let $i > j$.

Recall that

$$\mathbb{X}_{t_k^n, t_{k+1}^n}^{ij} = \int_{t_k^n}^{t_{k+1}^n} X_{t_k^n, u-}^i dX_u^j, \quad k = 0, \dots, N_n - 1.$$

Thus, the Burkholder–Davis–Gundy inequality gives that

$$\mathbb{E} \left[\sup_{\tau \in [0, T]} \left| \sum_{(s, t) \in \mathcal{P}^n \cap [0, \tau]} \mathbb{X}_{s, t}^{ij} \right|^2 \right] \lesssim \mathbb{E} \left[\int_0^T |X_{u|\mathcal{P}^n, u-}^i|^2 d[X^j]_u \right],$$

where we write $t_{|\mathcal{P}^n} := \max\{t_k^n \in \mathcal{P}^n : t_k^n \leq t\}$, $t \in [0, T]$. Proceeding as in the proof of [FZK23, Lemma 6.1], one can then show that

$$\mathbb{E} \left[\int_0^T |X_{u|\mathcal{P}^n, u-}^i|^2 d[X^j]_u \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

which gives (4.4).

Therefore, by definition and Lemma 4.1 it holds that

$$\begin{aligned} \sum_{k=0}^{N_n-1} \mathbb{X}_{t_k^n, t_{k+1}^n}^{ji} &= - \sum_{k=0}^{N_n-1} \mathbb{X}_{t_k^n, t_{k+1}^n}^{ij} + X_{t_k^n, t_{k+1}^n}^i X_{t_k^n, t_{k+1}^n}^j - [X^i, X^j]_{t_k^n, t_{k+1}^n} \\ &= - \sum_{k=0}^{N_n-1} \mathbb{X}_{t_k^n, t_{k+1}^n}^{ij} + \sum_{k=0}^{N_n-1} X_{t_k^n, t_{k+1}^n}^i X_{t_k^n, t_{k+1}^n}^j - \sum_{s \leq T} \Delta_s X^i \Delta_s X^j \\ &\rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$, where the convergence holds uniformly in probability, which then concludes the proof.

(ii) Let Y be the solution of the rough differential equation (3.4) driven by the random rough path $\mathbf{X} = (X, \mathbb{X})$, see Proposition 3.6 (i). We note that the assumption on (V, V') in (i) does fit into this setting, where $(V(\omega), V'(\omega)) = (F(Y(\omega)), F'(Y(\omega), Y'(\omega)))$ for some functional F , see Section 3.4. As the rough and Itô integral do coincide \mathbb{P} -almost surely by (i), we infer that Y is also a solution of the SDE (4.3), which has a unique solution (by e.g. [Pro05, Chapter V, Theorem 7]). \square

Remark 4.6. *As a consequence of Proposition 3.6 and Proposition 4.5 (ii), one can apply the continuity of the Itô–Lyons map (Theorem 2.5) to derive pathwise stability results for stochastic differential equations with delay like (4.3). In particular, the map $y \mapsto Y$, mapping the initial path y to the associated solution Y of the SDE (4.3), is continuous on the space of controlled paths, which resolves an old observation, pointed out by Mohammed [Moh86], about the non-continuity of the flow of stochastic differential equations with delay. The latter is a consequence of the discontinuity of stochastic integration when using an unsuitable topology for the integrands.*

Remark 4.7. *While we considered square-integrable martingales and the associated stochastic differential equations with constant delays in this section, the presented results can be generalized in a fairly straightforward manner to:*

- (i) càdlàg local martingales using standard localization arguments;
- (ii) càdlàg semimartingales using the classical estimates for Young integrals, see e.g. [FZ18, Proposition 2.4] and [LP18, Theorem 3.1], to show that one can suitably lift X to a random rough path with additional Young integrals;
- (iii) Young semimartingales (also known as semimartingales in the sense of Norvaiša [Nor03]), i.e. $Z = M + \varphi$, for some martingale M and some càdlàg adapted process φ with $\varphi(\omega) \in D^q([0, T]; \mathbb{R}^e)$ for almost every $\omega \in \Omega$, for some $q \in [1, 2)$;

- (iv) *SDEs/RDEs with variable delays of the form (3.6), as long as η is assumed to be bounded with $\eta(t) \geq \varepsilon$, $t \in [0, T]$, for some $\varepsilon > 0$.*

APPENDIX A. LOCAL ESTIMATES FOR ROUGH INTEGRATION

The following local estimates are needed to prove the existence and uniqueness result in Theorem 2.4, and the continuity result in Theorem 2.5.

Lemma A.1. *Let $\mathbf{X} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$ for $p \in (2, 3)$ and $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$. Suppose that the non-anticipative functional $(F, F'): \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ satisfies Assumption 2.1 (i) with some constant C_F . Then, we have the local estimate*

$$\|R^{\int_0^\cdot F(Y) d\mathbf{X}}\|_{\frac{p}{2}, [s, t]} \lesssim C_F (1 + \|Y, Y'\|_{X, p, [s, t]})^2 (1 + \|X\|_{p, [s, t]})^2 \|\mathbf{X}\|_{p, [s, t]},$$

for all $(s, t) \in \Delta_T$, where the implicit multiplicative constant depends only on p .

Proof. Let $(V, V') \in \mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$, and set $\Xi_{u,v} := V_u X_{u,v} + V'_u \mathbb{X}_{u,v}$ and $\delta \Xi_{u,r,v} := \Xi_{u,v} - \Xi_{u,r} - \Xi_{r,v}$ for $s \leq u < r < v \leq t$. Here, strictly speaking, in writing $V'_u \mathbb{X}_{u,v}$, we use the canonical identification of $\mathcal{L}(\mathbb{R}^d; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ with $\mathcal{L}(\mathbb{R}^d \otimes \mathbb{R}^d; \mathbb{R}^k)$. We note that

$$\begin{aligned} \left| R^{\int_0^\cdot V d\mathbf{X}} \right| &\leq \left| \int_u^v V_r d\mathbf{X}_r - \Xi_{u,v} \right| + |V'_u| |\mathbb{X}_{u,v}| \\ &\leq \left| \int_u^v V_r d\mathbf{X}_r - \Xi_{u,v} \right| + (|V'_s| + \|V'\|_{p, [s, t]}) |\mathbb{X}_{u,v}|. \end{aligned}$$

Using Chen's relation, one can show that

$$-\delta \Xi_{u,r,v} = R_{u,r}^V X_{r,v} + V'_{u,r} \mathbb{X}_{r,v},$$

which gives that

$$\begin{aligned} |\delta \Xi_{u,r,v}| &\leq \|R^V\|_{\frac{p}{2}, [u, r]} \|X\|_{p, [r, v]} + \|V'\|_{p, [u, r]} \|\mathbb{X}\|_{\frac{p}{2}, [r, v]} \\ &= w_{1,1}(u, r)^{\frac{2}{p}} w_{2,1}(r, v)^{\frac{1}{p}} + w_{1,2}(u, r)^{\frac{1}{p}} w_{2,2}(r, v)^{\frac{2}{p}}, \end{aligned}$$

where $w_{1,1}(s, t) := \|R^V\|_{\frac{p}{2}, [s, t]}^{\frac{2}{p}}$, $w_{2,1}(s, t) := \|X\|_{p, [s, t]}^p$, $w_{1,2}(s, t) := \|V'\|_{p, [s, t]}^p$, $w_{2,2}(s, t) := \|\mathbb{X}\|_{\frac{p}{2}, [s, t]}^{\frac{2}{p}}$, $(s, t) \in \Delta_T$, are control functions and $\frac{1}{p} + \frac{2}{p} > 1$. It then follows from the generalized sewing lemma, see [FZ18, Theorem 2.5], that

$$\begin{aligned} \|R^{\int_0^\cdot V d\mathbf{X}}\|_{\frac{p}{2}, [s, t]} &\lesssim (\|R^V\|_{\frac{p}{2}, [s, t]} \|X\|_{p, [s, t]} + \|V'\|_{p, [s, t]} \|\mathbb{X}\|_{\frac{p}{2}, [s, t]} + (|V'_s| + \|V'\|_{p, [s, t]}) \|\mathbb{X}\|_{\frac{p}{2}, [s, t]}) \\ &\lesssim \|V, V'\|_{X, p, [s, t]} \|\mathbf{X}\|_{p, [s, t]}, \end{aligned}$$

where the implicit multiplicative constant depends only on p . For $(V, V') = (F(Y), F'(Y, Y'))$, using Assumption 2.1 (i), we therefore obtain the estimate. \square

Lemma A.2. *For $p \in (2, 3)$, suppose $\mathbf{X}, \tilde{\mathbf{X}} \in \mathcal{D}^p([0, T]; \mathbb{R}^d)$, $(Y, Y') \in \mathcal{V}_X^p([0, T]; \mathbb{R}^k)$, $(\tilde{Y}, \tilde{Y}') \in \mathcal{V}_{\tilde{X}}^p([0, T]; \mathbb{R}^k)$, and that the non-anticipative functional $(F, F'): \mathcal{V}_X^p([0, T]; \mathbb{R}^k) \rightarrow$*

$\mathcal{V}_X^p([0, T]; \mathcal{L}(\mathbb{R}^d; \mathbb{R}^k))$ satisfies Assumption 2.1. Then, we have the local estimate

$$\begin{aligned} & \|R \int_0^\cdot F(Y) d\mathbf{X} - R \int_0^\cdot F(\tilde{Y}) d\tilde{\mathbf{X}}\|_{\frac{p}{2}, [s, t]} \\ & \lesssim C_{F, K, X, \tilde{X}} (\|Y_s - \tilde{Y}_s\| + \|Y, Y'; \tilde{Y}, \tilde{Y}'\|_{X, \tilde{X}, p, [s, t]} + \|X - \tilde{X}\|_{p, [s, t]}) (\|\mathbf{X}\|_{p, [s, t]} \vee \|\tilde{\mathbf{X}}\|_{p, [s, t]}) \\ & \quad + C_F (1 + K)^2 (1 + \|X\|_{p, [s, t]} \vee \|\tilde{X}\|_{p, [s, t]})^2 \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [s, t]} \end{aligned}$$

for all $(s, t) \in \Delta_T$, if $\|Y, Y'\|_{X, p, [s, t]}$, $\|\tilde{Y}, \tilde{Y}'\|_{\tilde{X}, p, [s, t]} \leq K$, for some $K > 0$, where the implicit multiplicative constant depends on p , $\|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$.

Proof. It follows from [FZ18, Lemma 3.4] that for any $(V, V') \in \mathcal{V}_X^p$, $(\tilde{V}, \tilde{V}') \in \mathcal{V}_{\tilde{X}}^p$,

$$\begin{aligned} & \|R \int_0^\cdot V d\mathbf{X} - R \int_0^\cdot \tilde{V} d\tilde{\mathbf{X}}\|_{\frac{p}{2}, [s, t]} \\ & \lesssim_p (1 + \|\mathbf{X}\|_{p, [s, t]} + \|\tilde{\mathbf{X}}\|_{p, [s, t]}) (\|V, V'; \tilde{V}, \tilde{V}'\|_{X, \tilde{X}, p, [s, t]} \|\mathbf{X}\|_{p, [s, t]} \\ & \quad + \|\tilde{V}, \tilde{V}'\|_{\tilde{X}, p, [s, t]} \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [s, t]}) \\ & \lesssim \|V, V'; \tilde{V}, \tilde{V}'\|_{X, \tilde{X}, p, [s, t]} \|\mathbf{X}\|_{p, [s, t]} + \|\tilde{V}, \tilde{V}'\|_{\tilde{X}, p, [s, t]} \|\mathbf{X}; \tilde{\mathbf{X}}\|_{p, [s, t]}, \end{aligned}$$

where the implicit multiplicative constant depends on p , $\|\mathbf{X}\|_p$ and $\|\tilde{\mathbf{X}}\|_p$. For $(V, V') = (F(Y), F'(Y, Y'))$, $(\tilde{V}, \tilde{V}') = (F(\tilde{Y}), F'(\tilde{Y}, \tilde{Y}'))$, using Assumption 2.1 (ii), we therefore obtain the estimate. \square

REFERENCES

- [AC20] Andrew L. Allan and Samuel N. Cohen, *Pathwise stochastic control with applications to robust filtering*, Ann. Appl. Probab. **30** (2020), no. 5, 2274–2310.
- [Aid15] Shigeki Aida, *Reflected rough differential equations*, Stochastic Process. Appl. **125** (2015), no. 9, 3570–3595.
- [Aid24] ———, *Rough differential equations containing path-dependent bounded variation terms*, Journal of Theoretical Probability (2024).
- [All21] Andrew L. Allan, *Robust filtering and propagation of uncertainty in hidden Markov models*, Electron. J. Probab. **26** (2021), Paper No. 73, 37.
- [ALP24] Andrew L. Allan, Chong Liu, and David J. Prömel, *A càdlàg rough path foundation for robust finance*, Finance Stoch. **28** (2024).
- [Ana23] Anna Ananova, *Rough differential equations with path-dependent coefficients*, Ann. H. Lebesgue **6** (2023), 1–29.
- [Bai14] Ismaël Bailleul, *Flows driven by Banach space-valued rough paths*, Séminaire de Probabilités XLVI, Lecture Notes in Math., vol. 2123, Springer, Cham, 2014, pp. 195–205.
- [BBC21] Ismaël Bailleul, Antoine Brault, and Laure Coutin, *Young and rough differential inclusions*, Rev. Mat. Iberoam. **37** (2021), no. 4, 1489–1512.
- [BBR20] Mireia Besalú, Giulia Binotto, and Carles Rovira, *Convergence of delay equations driven by a Hölder continuous function of order $1/3 < \beta < 1/2$* , Electron. J. Differ. Equ. (2020), Paper No. 65, 27.
- [BDNHP18] D. R. Baños, G. Di Nunno, H. H. Haferkorn, and F. Proske, *Stochastic functional differential equations and sensitivity to their initial path*, Computation and combinatorics in dynamics, stochastics and control, Abel Symp., vol. 13, Springer, Cham, 2018, pp. 37–70.
- [BMCR14] Mireia Besalú, David Márquez-Carreras, and Carles Rovira, *Delay equations with non-negativity constraints driven by a Hölder continuous function of order $\beta \in (\frac{1}{3}, \frac{1}{2})$* , Potential Anal. **41** (2014), no. 1, 117–141.
- [CE20] Reda Chhaibi and Ibrahim Ekren, *A Hörmander condition for delayed stochastic differential equations*, Ann. H. Lebesgue **3** (2020), 1023–1048.

- [CFO11] Michael Caruana, Peter K. Friz, and Harald Oberhauser, *A (rough) pathwise approach to a class of non-linear stochastic partial differential equations*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **28** (2011), no. 1, 27–46.
- [Che18] Ilya Chevyrev, *Random walks and Lévy processes as rough paths*, Probab. Theory Related Fields **170** (2018), no. 3-4.
- [DFG17] Joscha Diehl, Peter K. Friz, and Paul Gassiat, *Stochastic control with rough paths*, Appl. Math. Optim. **75** (2017), no. 2, 285–315.
- [DGHT19] Aurélien Deya, Massimiliano Gubinelli, Martina Hofmanová, and Samy Tindel, *One-dimensional reflected rough differential equations*, Stochastic Process. Appl. **129** (2019), no. 9, 3261–3281.
- [DT09] Aurélien Deya and Samy Tindel, *Rough Volterra equations. I. The algebraic integration setting*, Stoch. Dyn. **9** (2009), no. 3, 437–477.
- [Dup19] Bruno Dupire, *Functional Itô calculus*, Quant. Finance **19** (2019), no. 5, 721–729.
- [FH20] Peter K. Friz and Martin Hairer, *A course on rough paths*, second ed., Universitext, Springer, Cham, 2020, With an introduction to regularity structures.
- [FR06] Marco Ferrante and Carles Rovira, *Stochastic delay differential equations driven by fractional Brownian motion with Hurst parameter $H > \frac{1}{2}$* , Bernoulli **12** (2006), no. 1, 85–100.
- [FS17] Peter K. Friz and Atul Shekhar, *General rough integration, Lévy rough paths and a Lévy-Kintchine-type formula*, Ann. Probab. **45** (2017), no. 4, 2707–2765.
- [FZ18] Peter K. Friz and Huilin Zhang, *Differential equations driven by rough paths with jumps*, J. Differ. Equ. **264** (2018), no. 10, 6226–6301.
- [FZK23] Peter K. Friz and Pavel Zorin-Kranich, *Rough semimartingales and p -variation estimates for martingale transforms*, Ann. Probab. **51** (2023), no. 2, 397–441. MR 4546622
- [Gas21] Paul Gassiat, *Non-uniqueness for reflected rough differential equations*, Ann. Inst. Henri Poincaré Probab. Stat. **57** (2021), no. 3, 1369–1387.
- [Hai11] Martin Hairer, *Rough stochastic PDEs*, Comm. Pure Appl. Math. **64** (2011), no. 11, 1547–1585.
- [JS03] Jean Jacod and Albert N. Shiryaev, *Limit theorems for stochastic processes*, 2nd ed., Springer, 2003.
- [LCL07] Terry J. Lyons, Michael Caruana, and Thierry Lévy, *Differential equations driven by rough paths*, Lecture Notes in Mathematics, vol. 1908, Springer, Berlin, 2007.
- [Lép76] Dominique Lépingle, *La variation d'ordre p des semi-martingales*, Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete **36** (1976), 295–316.
- [LP18] Chong Liu and David J. Prömel, *Examples of Itô càdlàg rough paths*, Proc. Amer. Math. Soc. **146** (2018), no. 11, 4937–4950.
- [Lyo98] Terry J. Lyons, *Differential equations driven by rough signals*, Rev. Mat. Iberoam. **14** (1998), no. 2, 215–310.
- [Moh84] Salah-Eldin A. Mohammed, *Stochastic functional differential equations*, Research Notes in Mathematics, vol. 99, Pitman (Advanced Publishing Program), Boston, MA, 1984.
- [Moh86] ———, *Nonlinear flows of stochastic linear delay equations*, Stochastics **17** (1986), no. 3, 207–213.
- [Moh98] ———, *Stochastic differential systems with memory: theory, examples and applications*, Stochastic analysis and related topics, VI (Geilo, 1996), Progr. Probab., vol. 42, Birkhäuser Boston, Boston, MA, 1998, pp. 1–77.
- [NNT08] Andreas Neuenkirch, Ivan Nourdin, and Samy Tindel, *Delay equations driven by rough paths*, Electron. J. Probab. **13** (2008), no. 67, 2031–2068.
- [Nor03] R. Norvaiša, *The p -variation and an extension of the class of semimartingales*, Acta Appl. Math. **79** (2003), no. 1-2, 47–53.
- [Pha09] Huyên Pham, *Continuous-time stochastic control and optimization with financial applications*, Stochastic Modelling and Applied Probability, vol. 61, Springer-Verlag, Berlin, 2009.
- [Pro05] Philip E. Protter, *Stochastic integration and differential equations*, Stochastic Modelling and Applied Probability, vol. 21, Springer-Verlag, Berlin, 2005, Second edition. Version 2.1, Corrected third printing.
- [PT21] David J. Prömel and Mathias Trabs, *Paracontrolled distribution approach to stochastic Volterra equations*, J. Differ. Equ. **302** (2021), 222–272.
- [RV93] Francesco Russo and Pierre Vallois, *Forward, backward and symmetric stochastic integration*, Probab. Theory Related Fields **97** (1993), no. 3, 403–421.

- [TT12] Samy Tindel and Iván Torrecilla, *Some differential systems driven by a fBm with Hurst parameter greater than 1/4*, Stochastic analysis and related topics, Springer Proc. Math. Stat., vol. 22, Springer, Heidelberg, 2012, pp. 169–202.

ANNA P. KWOSSEK, UNIVERSITY OF MANNHEIM, GERMANY
Email address: anna.kwossek@uni-mannheim.de

ANDREAS NEUENKIRCH, UNIVERSITY OF MANNHEIM, GERMANY
Email address: neuenkirch@uni-mannheim.de

DAVID J. PRÖMEL, UNIVERSITY OF MANNHEIM, GERMANY
Email address: proemel@uni-mannheim.de