Study of transport properties of a hot and dense QCD matter using a novel approximation method

Anowar Shaikh^{∗1}, Shubhalaxmi Rath^{†2}, Sadhana Dash^{‡3}, and Binata Panda^{§1}

¹Department of Physics, Indian Institute of Technology (Indian School of Mines) Dhanbad, Jharkhand 826004, India

²Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica, Chile ³Department of Physics, Indian Institute of Technology Bombay, Mumbai 400076, India

Abstract

We have studied the charge and heat transport properties of a hot and dense QCD matter using a novel approximation method within the quasiparticle model. Utilizing the novel collision integrals for both the relaxation time approximation (RTA) and the Bhatnagar-Gross-Krook (BGK) models, we have solved the relativistic Boltzmann transport equation to estimate the electrical conductivity and the thermal conductivity of the medium. We have investigated the temperature dependence of these transport coefficients. We have also provided a comparison between our findings and those of the standard RTA and standard BGK models. Further, we have explored the temperature dependence of the thermal diffusion constant and the Lorenz number using the novel approaches of the aforesaid models.

[∗]anowar.19dr0016@ap.iitism.ac.in

[†] shubhalaxmirath@gmail.com

[‡] sadhana@phy.iitb.ac.in

[§]binata@iitism.ac.in

1 Introduction

A long time goal of the physics of ultrarelativistic heavy ion collisions is the observation of new states of matter predicted by the quantum chromodynamics (QCD). The experimental evidences of the relativistic heavy ion collider (RHIC) [\[1](#page-33-0)] at BNL and the large hadron collider (LHC) [\[2](#page-33-1)] at CERN indicate a new state of the strongly interacting matter known as quark-gluon plasma (QGP). Following the discovery of QGP, various properties of the strongly coupled QCD matter became a topic of high interest. Different research groups have studied different features of QGP in recent years, such as the thermodynamic properties [\[3](#page-33-2), [4](#page-33-3)], the relativistic hydrodynamics [\[5](#page-34-0)[–7](#page-34-1)], the heavy quark diffusion in QCD matter $[8–10]$ $[8–10]$, the memory effect on heavy quark dynamics $[11, 12]$ $[11, 12]$, the dissipative spin dynamics in relativistic matter [\[13](#page-34-6), [14\]](#page-34-7) etc. Nowadays, there is much interest in the phase diagram region at finite temperature and baryon chemical potential [\[15](#page-34-8), [16\]](#page-35-0). Many observations have also been made recently to investigate the impact of a strong magnetic field on the characteristics of hot QCD matter, such as the chiral magnetic effect [\[17,](#page-35-1) [18\]](#page-35-2), the chiral vortical effect [\[19\]](#page-35-3), the axial Hall current [\[20](#page-35-4)], the refractive indices and decay constant [\[21,](#page-35-5) [22](#page-35-6)], the photon and dilepton productions from QGP [\[23](#page-35-7)[–25\]](#page-35-8), the thermodynamic and magnetic properties [\[26](#page-35-9)[–31](#page-36-0)], the axial magnetic effect [\[32](#page-36-1), [33\]](#page-36-2), the thermoelectric responses of hot QCD medium in a time varying magnetic field [\[34](#page-36-3), [35](#page-36-4)], the magnetic and inverse magnetic catalysis [\[36](#page-36-5), [37\]](#page-36-6), the nonlinear electromagnetic current [\[38](#page-36-7), [39\]](#page-36-8) and the magnetohydrodynamics [\[40,](#page-37-0) [41\]](#page-37-1).

It is crucial to investigate different transport coefficients since they shed light on the genesis of the hot QCD medium. Among these transport coefficients, the thermal conductivity (κ) and the electrical conductivity (σ_{el}) characterize the movement of heat and charges in the medium, respectively. The hydrodynamic development of the strongly interacting matter at finite baryon density also depends on these two conductivities [\[42,](#page-37-2) [43\]](#page-37-3). Therefore, significant effects on the observables at heavy ion collisions might result from changes in the charge and the heat transport coefficients. There are several methods for determining these conductivities, such as the relativistic Boltzmann transport equation [\[44,](#page-37-4) [45\]](#page-37-5), the Chapman-Enskog approximation [\[46,](#page-37-6) [47\]](#page-37-7), the lattice simulation [\[48](#page-37-8)[–50\]](#page-37-9), the correlator technique using Green-Kubo formula [\[51](#page-37-10)[–53](#page-38-0)] etc. Previous research has examined the impact of magnetic field on the electrical conductivity using a variety of methods, including the diluted instanton-liquid model $[51]$, the quenched $SU(2)$ lattice gauge theory [\[54\]](#page-38-1), the effective fugacity technique [\[55](#page-38-2)], the diagrammatic method utilizing the real-time formalism [\[56\]](#page-38-3) etc. Studies on the electrical and the thermal conductivities of hot QCD matter in the presence of magnetic field within the kinetic theory approach have been carried out in references [\[57](#page-38-4)[–60\]](#page-38-5).

The application of theoretical methodologies intended for systems of this type has been

made viable by the strongly coupled fluid behavior of QGP. A simplified version of the kinetic equation with a simplified form of the collision term is frequently used because of various technical limitations. It is referred as the relaxation time approximation (RTA) [\[61\]](#page-38-6). In this model, the mean-free approach reduced the complexity of the collision term. The development in the nonrelativistic case has already been achieved by Bhatnagar-Gross-Krook (BGK) model [\[62](#page-38-7)] and Welander model [\[63](#page-38-8)]. This approximation is utilized in many domains of physics and is recently used to investigate the hydrodynamization of the matter generated by the ultrarelativistic collisions of heavy ions. The relaxation time approximation has become essential for exploring the domain and microscopic basis of the relevance of hydrodynamics with relativistic effects. Although widely used, the approximation presented in ref. [\[61](#page-38-6)] has some fundamental challenges. It conflicts with some macroscopic and microscopic conservation laws. Recently, some works $[64-66]$ provided a novel relaxation time approximation that fixes such fundamental issues and maintains the core features of the linearized Boltzmann collision operator. In light of the momentum-dependent relaxation time, the novel relaxation time approximation is significant as it maintains the fundamental collision features associated with the microscopic conservation laws. They showed how this adjustment impacts the particle diffusion and the bulk viscosity. In a recent study [\[67](#page-39-2)], the transport coefficients related to charge and heat for a rotating QGP medium have been derived within the RTA model utilizing the aforementioned collision integral.

In this study, we estimate the response function of charge flow and heat flow, such as the electrical conductivity (σ_{el}) and the thermal conductivity (κ) , to investigate the charge and the thermal transport properties of the hot and dense QCD matter. These transport coefficients have been computed using a modified collision integral in the relativistic Boltzmann transport equation within the kinetic theory approach. We also explore the thermal diffusion constant and the Lorenz number as the applications of charge and heat transport properties in an analogous environment. The thermal diffusion constant shows the overall rate of heat transport in the medium, while the Lorenz number illustrates the connection between the thermal and electrical conductivities. We use the quasiparticle description of partons, where the rest masses are replaced by the masses generated in the medium. The quark-gluon plasma is viewed as a medium made up of thermally massive quasiparticles in this framework [\[68](#page-39-3), [69](#page-39-4)].

The rest of the paper has been organized as follows: Using the modified collision integral, the charge and heat transport coefficients from the relativistic Boltzmann transport equation are calculated in section 2. In section 3, we have discussed the results of the aforesaid transport coefficients using the quasiparticle description of partons. Section 4 describes various applications and their results obtained from these coefficients, such as the Lorenz number and the thermal diffusion constant. In section 5, the results are summarized.

Notations and conventions: The covariant derivatives ∂_{μ} and $\partial_{\mu}^{(p)}$ denote $\frac{\partial}{\partial x^{\mu}}$ and $\frac{\partial}{\partial p^{\mu}}$ respectively. The fluid four-velocity in the rest frame $(u^{\mu} = (1,0,0,0))$ is normalized to unity. In this work, the subscript f stands for the flavor index, with $f=u$, d, s. The symbols q_f , g_f and δf_f ($\delta \bar{f}_f$) stand for electric charge, degeneracy factor and the infinitesimal change in the distribution function for the quark (antiquark) of the f^{th} flavor, respectively. Here, $g_g = 2(N_c^2 - 1)$ represents the degeneracy factor for the gluonic degree of freedom, and $g_f(g_{\bar{f}})=2N_c$ denotes the degeneracy factor for the quark (antiquark) degree of freedom with flavor f, where $N_c = 3$ is the number of colours.

2 Charge and heat transport properties of a hot and dense QCD medium

For a single particle distribution function, the relativistic Boltzmann transport equation (RBTE) is written as

$$
\frac{\partial f_f}{\partial t} + \frac{\mathbf{p}}{m} \nabla f_f + \mathbf{F} \cdot \frac{\partial f_f}{\partial \mathbf{p}} = \left(\frac{\partial f_f}{\partial t}\right)_{\mathbf{c}},\tag{1}
$$

where \bf{F} is the force field acting on the particles in the medium and m is the mass of the particle. The term on the right-hand side results from particle collisions within the medium. This term's zero value denotes the Vlasov equation, which describes a collisionless system. The intricate character of the collision term presents a number of challenges even for the simplest solution of the above eq. [\(1\)](#page-3-0). The term $\left(\frac{\partial f_f}{\partial t}\right)$ c represents the instantaneous change in the distribution function due to collisions between particles. In order to arrive at a comprehensive solution, we will first study a few mathematical models those allow us to address the collision term in a suitable manner.

When using the relaxation time approximation (RTA), the collision term is substituted with a relaxation term of the following form,

$$
\left(\frac{\partial f_f}{\partial t}\right)_c = -\frac{1}{\tau_{fp}} \Big(f_f(x, p, t) - f_{eq, f}(p) \Big),\tag{2}
$$

where we can write $f_f(x, p, t) = f_{eq,f}(p) + \delta f_f(x, p, t)$ and τ_{fp} is the relaxation time for collisions, *i.e.* the time required to bring the perturbed system back to its equilibrium state. We can assume that the quark distribution function is near the equilibrium with a slight divergence from equilibrium in order to permit linearization of the relativistic Boltzmann equation. Here, the term $\frac{1}{\tau_{fp}}$ acts as a damping frequency. The initial issue with this model is that, there is no immediate conservation of charge. To remove the crunch, the Bhatnagar-Gross-Krook (BGK) model presents a new type of collision kernel,

denoted as

$$
\left(\frac{\partial f_f}{\partial t}\right)_c = -\frac{1}{\tau_{fp}} \left(f_f(x, p, t) - \frac{n_f(x, t)}{n_{eq, f}} f_{eq, f}(p) \right),\tag{3}
$$

where $n_{eq,f}$ represents the equilibrium particle number density and $n_f(x,t)$ is known as the perturbed particle number density or the fluctuating particle number density. The perturbed particle number density is calculated as $\int_p f_f(x, p, t)$, which gives zero value after integrating over momentum, thus indicating instantaneous conservation of the particle number during collisions in the system. Such fundamental problems are resolved and the essential characteristics of the linearized Boltzmann collision operator are preserved via a novel relaxation time approximation that was recently presented in a research $[64]$.

Here, we are considering the general relativistic covariant form of the Boltzmann transport equation for an isotropic medium of quarks, antiquarks and gluons, which is given as

$$
p^{\mu}\partial_{\mu}f_f + q_f F^{\mu\nu} p_{\nu}\partial_{\mu}^{(p)} f_f = \mathbf{C}[f_f],\tag{4}
$$

where $F^{\mu\nu}$ is the electromagnetic field strength tensor. In order to see the effects of electric field, we take only $\mu = i$ and $\nu = 0$ and vice versa components of the electromagnetic field strength tensor, i.e. $F^{i0} = \mathbf{E}$ and $F^{0i} = -\mathbf{E}$ in our calculation, such that the RBTE [\(4\)](#page-4-0) takes the following form,

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = \mathbf{C} [f_f]. \tag{5}
$$

The infinitesimal change in quark distribution function is denoted as δf_f and $f_{eq,f}$ indicates the equilibrium distribution function in the medium for f^{th} flavor.

2.1 Electrical conductivity in the novel RTA and the novel BGK models

The electric current density of the medium must be determined in order to see how the electric field affects the charge transport phenomena. An external electric field can cause an infinitesimal disturbance to a thermal QCD medium that contains quarks, antiquarks, and gluons of distinct flavors. Due to this electric field, there is an induced current density in the medium, which is given by

$$
J^{i} = \sum_{f} g_{f} q_{f} \int \frac{d^{3}p}{(2\pi)^{3}} \frac{p^{i}}{\omega_{f}} \left[\delta f_{f}(x, p) + \delta \bar{f}_{f}(x, p) \right], \tag{6}
$$

where J^i represents the spatial part of the electric current density vector. Ohm's law states that the spatial component of the electric current density is directly proportional to the electric field with a proportionality factor, known as the electrical conductivity (σ_{el}) . Thus, we have

$$
J^i = \sigma_{el} E^i. \tag{7}
$$

By comparing eq. (6) and eq. (7) , one can obtain the expression of the electrical conductivity.

By solving the relativistic Boltzmann transport equation, the infinitesimal changes in the quark and antiquark distribution functions can be obtained. The relativistic Boltzmann transport equation in the novel RTA model can be written as

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = \mathbf{C} [f_f], \tag{8}
$$

where the collision integral is given [\[64\]](#page-39-0) by

$$
\mathbf{C}[f_f] = -\frac{\omega_f}{\tau_{fp}} \left[\delta f_f - \frac{\langle (\omega_f/\tau_{fp}) \delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp}) P_1^{(0)} \delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \delta f_f \rangle_0}{\langle 1/3 \rangle \langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} \right]. \tag{9}
$$

Here,

$$
P_1^{(0)} = 1 - \frac{\langle \omega_f / \tau_{fp} \rangle_0}{\langle \omega_f^2 / \tau_{fp} \rangle_0} \omega_f,
$$
\n(10)

$$
p^{\langle \mu \rangle} = \Delta^{\mu \nu} p_{\nu},\tag{11}
$$

$$
\Delta^{\mu\nu} = g^{\mu\nu} - u^{\mu}u^{\nu},\tag{12}
$$

and we define the momentum integrals relative to the local equilibrium distribution function $f_{eq,f}$ as follows,

$$
\langle \ldots \rangle_0 = \int \frac{d^3 p}{(2\pi)^3 p_0} f_{eq,f}(\ldots)
$$

$$
= \int dP f_{eq,f}(\ldots), \qquad (13)
$$

where τ_{fp} is the relaxation time and $\nu_f = \frac{1}{\tau_f}$ $\frac{1}{\tau_{fp}}$ denotes the collision frequency of the medium. The equilibrium distribution functions for f^{th} flavor of quark and antiquark are given by

$$
f_{eq,f} = \frac{1}{e^{\beta(\omega_f - \mu_f)} + 1},\tag{14}
$$

$$
\bar{f}_{eq,f} = \frac{1}{e^{\beta(\omega_f + \mu_f)} + 1},\tag{15}
$$

respectively, where $\omega_f = \sqrt{\mathbf{p}^2 + m_f^2}$, $\beta = \frac{1}{T}$ $\frac{1}{T}$, and μ_f represents the chemical potential of fth flavor of quark. Getting inspired by references [\[70](#page-39-5)[–72\]](#page-39-6), we have used the relaxation time of the following form,

$$
\tau_{fp}(\omega_f) = (\beta \omega_f)^\gamma \tau_f,\tag{16}
$$

where γ is an arbitrary constant which controls the energy dependence of the relaxation time and the expression for the τ_f is given [\[73\]](#page-39-7) by

$$
\tau_f(T) = \frac{1}{5.1T\alpha_s^2 \log(1/\alpha_s) \left[1 + 0.12(2N_f + 1)\right]}.
$$
\n(17)

Here, α_s is the QCD running coupling constant, which is a function of both temperature and chemical potential and it has the following [\[74\]](#page-39-8) form,

$$
\alpha_s = \frac{g^2}{4\pi} = \frac{12\pi}{(11N_c - 2N_f)\ln\left(\Lambda^2/\Lambda_{\overline{\rm MS}}^2\right)}\,,\tag{18}
$$

where $\Lambda_{\overline{\rm MS}} = 0.176 \text{ GeV}, \Lambda = 2\pi \sqrt{T^2 + \mu_f^2/\pi^2}$ for electrically charged particles (quarks and antiquarks) and $\Lambda = 2\pi T$ for gluons. Using the expression of $\mathbb{C}[f_f]$ in eq. [\(8\)](#page-5-0), we have

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\frac{\omega_f}{\tau_{fp}} \left[\delta f_f - \frac{\langle (\omega_f / \tau_{fp}) \delta f_f \rangle_0}{\langle \omega_f / \tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f / \tau_{fp}) P_1^{(0)} \delta f_f \rangle_0}{\langle (\omega_f / \tau_{fp}) P_1^{(0)} P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f / \tau_{fp}) p_{\langle \mu \rangle} \delta f_f \rangle_0}{\langle (1/3) \langle (\omega_f / \tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} \right]. \tag{19}
$$

Now, solving the above equation one can extract the expression of δf_f (for a detailed calculation see appendix [A\)](#page-19-0) as

$$
\delta f_f = \frac{2q_f \beta \tau_{fp} (\mathbf{E} \cdot \mathbf{p}) f_{eq,f} (1 - f_{eq,f})}{\omega_f A},\tag{20}
$$

where

$$
A = \left[1 - \frac{\omega_f \int p^2(f_{eq,f}/\tau_{fp})dp}{\int p^2 \omega_f(f_{eq,f}/\tau_{fp})dp}\right] \frac{\int p^2(f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2(f_{eq,f}/\tau_{fp})dp}{\int p^2 \omega_f(f_{eq,f}/\tau_{fp})dp}\right) \omega_f\right] dp}{\int p^2(f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2(f_{eq,f}/\tau_{fp})dp}{\int p^2 \omega_f(f_{eq,f}/\tau_{fp})dp}\right) \omega_f\right]^2 dp} + \frac{3p \int p^3(f_{eq,f}/\tau_{fp})dp}{\int p^4(f_{eq,f}/\tau_{fp})dp}.
$$
 (21)

Similarly for antiquarks,

$$
\delta \bar{f}_f = \frac{2q_{\bar{f}} \beta \tau_{\bar{f}p} (\mathbf{E} \cdot \mathbf{p}) \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\omega_f \bar{A}},\tag{22}
$$

where

$$
\bar{A} = \left[1 - \frac{\omega_f \int p^2(\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}{\int p^2 \omega_f (\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}\right] \frac{\int p^2(\bar{f}_{eq,f}/\tau_{\bar{f}p}) \left[1 - \left(\frac{\int p^2(\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}{\int p^2 \omega_f (\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}\right) \omega_f\right] dp}{\int p^2(\bar{f}_{eq,f}/\tau_{\bar{f}p}) \left[1 - \left(\frac{\int p^2(\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}{\int p^2 \omega_f (\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}\right) \omega_f\right]^2 dp} + \frac{3p \int p^3(\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}{\int p^4(\bar{f}_{eq,f}/\tau_{\bar{f}p})dp}.
$$
 (23)

Finally, replacing the values of δf_f and $\delta \bar{f}_f$ in eq. [\(6\)](#page-4-1) and comparing it with eq. [\(7\)](#page-4-2), we get the final expression of the electrical conductivity in the novel RTA model as

$$
\sigma_{el}^{Novel - RTA} = \frac{\beta}{3\pi^2} \sum_f g_f q_f^2 \int dp \frac{p^4}{\omega_f^2} \left[\frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f})}{A} + \frac{\tau_{\bar{f}p} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\bar{A}} \right].
$$
 (24)

In the standard BGK model, the relativistic Boltzmann transport equation is written as

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = \mathbf{C} [f_f] = -\frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left(f_f(x, p, t) - \frac{n_f(x, t)}{n_{eq, f}} f_{eq, f}(p) \right), \quad (25)
$$

where the collision integral is given as

$$
\mathbf{C}[f_f] = -\frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left(f_f(x, p, t) - \frac{n_f(x, t)}{n_{eq, f}} f_{eq, f}(p) \right). \tag{26}
$$

In the above equation, n_f and $n_{eq,f}$ represent the perturbed particle number density and the equilibrium particle number density, respectively for the fth flavor, which has a degeneracy of g_f . Both can be calculated from the equations as given below,

$$
n_f = g_f \int dP \omega_f (f_{eq,f} + \delta f_f), \qquad (27)
$$

$$
n_{eq,f} = g_f \int dP \omega_f f_{eq,f}.
$$
\n(28)

The BGK collisional-kernel's instantaneous conservation of the particle number is its main benefit over the RTA collisional-kernel [\[31](#page-36-0), [62,](#page-38-7) [75\]](#page-39-9), *i.e.*, in case of the BGK model, we have

$$
\int dP \mathbf{C}[f_f] = 0. \tag{29}
$$

Furthermore, the ref. [\[76\]](#page-39-10) had shown that, by implementing a straightforward adjustment, the BGK collisional kernel effectively maintains the conservation of the color current in a covariant manner. The Boltzmann equation has been recently formulated using the BGK collision kernel to represent relativistic dissipative hydrodynamics [\[77](#page-39-11)].

In the present work, we proposed a collision integral in the following manner,

$$
\mathbf{C}[f_f]^{NBGK} = -\frac{p^{\mu}u_{\mu}}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{\langle (1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \bigg] + g_f n_{eq,f}^{-1} (p^{\mu}u_{\mu}) f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{\langle (1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \bigg] dP. \tag{30}
$$

To fulfill the basic conservation equations in microscopic interactions, the zeroth and first moments of the collision kernel must vanish (detailed calculations are given in appendix [B\)](#page-21-0).

In the novel BGK model, the relativistic Boltzmann transport equation can be expressed as

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left[\delta f_f - \frac{\langle (\omega_f/\tau_{fp}) \delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp}) P_1^{(0)} \delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \rangle_0} \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \delta f_f \rangle_0}{\langle (1/3) \langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} + g_f n_{eq,f}^{-1} (p^{\mu} u_{\mu}) f_{eq,f} \int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left[\delta f_f - \frac{\langle (\omega_f/\tau_{fp}) \delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle 0 \rangle} \delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp}) P_1^{(0)} \delta f_f \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \delta f_f \rangle_0}{\langle (1/3) \langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} \right] dP. \quad (31)
$$

Now, neglecting higher order terms $(\mathcal{O}(\delta f_f)^2)$, we get the solution of the above equation as

$$
\delta f_f = \delta f_f^{(0)} + g_f n_{eq,f}^{-1} f_{eq,f} \int_{p'} \delta f_f^{(0)1},\tag{32}
$$

where $\delta f_f^{(0)}$ is given by

$$
\delta f_f^{(0)} = \frac{2q_f \beta \tau_{fp} (\mathbf{E} \cdot \mathbf{p}) f_{eq,f} (1 - f_{eq,f})}{\omega_f A}.
$$
\n(33)

The infinitesimal change in the quark distribution function is obtained (details are given in appendix C) as

$$
\delta f_f = \left[2q_f \beta \tau_{fp} \frac{\mathbf{E} \cdot \mathbf{p}}{\omega_f A} f_{eq,f} (1 - f_{eq,f}) \right] + g_f n_{eq,f}^{-1} f_{eq,f} \int_{p'} \left[2q_f \beta \tau_{fp} \frac{\mathbf{E} \cdot \mathbf{p}}{\omega_f A} f_{eq,f} (1 - f_{eq,f}) \right]. \tag{34}
$$

Similarly, $\delta \bar{f}_f$ is obtained as

$$
\delta \bar{f}_f = \left[2q_{\bar{f}} \beta \tau_{\bar{f}p} \frac{\mathbf{E} \cdot \mathbf{p}}{\omega_f \bar{A}} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f}) \right] + g_f n_{eq,f}^{-1} \bar{f}_{eq,f} \int_{p'} \left[2q_{\bar{f}} \beta \tau_{\bar{f}p} \frac{\mathbf{E} \cdot \mathbf{p}}{\omega_f \bar{A}} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f}) \right]. \tag{35}
$$

Now, replacing the values of δf_f and $\delta \bar{f}_f$ in eq. [\(6\)](#page-4-1) and comparing it with eq. [\(7\)](#page-4-2), we get the final expression of the electrical conductivity in the novel BGK model as

$$
\sigma_{el}^{Novel-BGK} = \frac{\beta}{3\pi^2} \sum_{f} g_f q_f^2 \int dp \frac{p^4}{\omega_f^2} \left[\frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f})}{A} + \frac{\tau_{fp} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\bar{A}} \right]
$$

$$
+ \frac{2\beta}{\sqrt{3}} \sum_{f} g_f^2 q_f^2 n_{eq,f}^{-1} \left[\int_p \frac{p}{\omega_f} f_{eq,f} \int_{p'} \frac{p'}{\omega_f} \frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f})}{A} + \int_p \frac{p}{\omega_f} \bar{f}_{eq,f} \int_{p'} \frac{p'}{\omega_f} \frac{\tau_{fp} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\bar{A}} \right].
$$
(36)

¹Here, we are using the momentum integration symbol, $\int_p = \int d^3p/(2\pi)^3$.

2.2 Thermal conductivity in the novel RTA and the novel BGK models

Heat flow results from the deviation of the system from its equilibrium condition caused by the presence of a temperature gradient. Through a proportionality constant, *i.e.* the thermal conductivity, the heat flow vector is directly proportional to the temperature gradient. The calculation of the thermal conductivity is necessary in examining the heat transmission in a medium. In four-vector notation, the heat flow is written as

$$
Q_{\mu} = \Delta_{\mu\alpha} T^{\alpha\beta} u_{\beta} - h \Delta_{\mu\alpha} N^{\alpha}.
$$
\n(37)

In the above equation, the projection operator $(\Delta_{\mu\alpha})$ and the enthalpy per particle (h) are respectively defined as

$$
\Delta_{\mu\alpha} = g_{\mu\alpha} - u_{\mu}u_{\alpha},\tag{38}
$$

$$
h = (\varepsilon + P)/n,\tag{39}
$$

where n, ε and P represent the particle number density, the energy density and the pressure, respectively, which are respectively given by the following equations,

$$
n = N^{\alpha} u_{\alpha},\tag{40}
$$

$$
\varepsilon = u_{\alpha} T^{\alpha \beta} u_{\beta},\tag{41}
$$

$$
P = -\frac{\Delta_{\alpha\beta} T^{\alpha\beta}}{3}.
$$
\n(42)

Here, the particle flow four-vector N^{α} and the energy-momentum tensor $T^{\alpha\beta}$ are respectively defined as

$$
N^{\alpha} = \sum_{f} g_f \int \frac{d^3 p}{(2\pi)^3} \frac{p^{\alpha}}{\omega_f} \left[f_f(x, p) + \bar{f}_f(x, p) \right],\tag{43}
$$

$$
T^{\alpha\beta} = \sum_{f} g_f \int \frac{d^3 p}{(2\pi)^3} \frac{p^{\alpha} p^{\beta}}{\omega_f} \left[f_f(x, p) + \bar{f}_f(x, p) \right]. \tag{44}
$$

The spatial part of the heat flow four-vector is given by

$$
Q^i = \sum_f g_f \int \frac{d^3 p}{(2\pi)^3} \frac{p^i}{\omega_f} \left[(\omega_f - h_f) \delta f_f(x, p) + (\omega_f - \bar{h}_f) \delta \bar{f}_f(x, p) \right],\tag{45}
$$

where we use the fact that in the rest frame of the fluid, the heat flow four-vector and fluid four-vector are perpendicular to each other, *i.e.* $Q_{\mu}u^{\mu} = 0$ and the enthalpy per particle for the f^{th} flavor is given by

$$
h_f = \frac{(\varepsilon_{eq,f} + P_{eq,f})}{n_{eq,f}},\tag{46}
$$

where

$$
\varepsilon_{eq,f} = g_f \int dP \; \omega_f^2 f_{eq,f},\tag{47}
$$

$$
P_{eq,f} = \frac{g_f}{3} \int dP \ p^2 f_{eq,f}.
$$
\n(48)

In the Navier-Stokes equation, the heat flow is related to the gradients of temperature and pressure [\[78](#page-40-0)] as

$$
Q^i = -\kappa \delta^{ij} \left[\partial_j T - \frac{T}{\varepsilon + P} \partial_j P \right]. \tag{49}
$$

Here, κ represents the thermal conductivity. One can find the expression of the thermal conductivity by comparing eq. [\(45\)](#page-9-0) and eq. [\(49\)](#page-10-0).

In order to obtain the expression of thermal conductivity, we start with the relativistic Boltzmann transport equation, which can be rewritten as

$$
p^{\mu}\partial_{\mu}T\frac{\partial f_{eq,f}}{\partial T} + p^{\mu}\partial_{\mu}(p^{\nu}u_{\nu})\frac{\partial f_{eq,f}}{\partial p^{0}} + q_{f}\Big[F^{0i}p_{i}\frac{\partial f_{eq,f}}{\partial p^{0}} + F^{i0}p_{0}\frac{\partial f_{eq,f}}{\partial p^{i}}\Big] = \mathbf{C}\big[f_{f}\big].\tag{50}
$$

With the help of eq. [\(9\)](#page-5-1), the above equation can be written in the novel RTA model as

$$
p^{\mu}\partial_{\mu}T\frac{\partial f_{eq,f}}{\partial T} + p^{\mu}\partial_{\mu}(p^{\nu}u_{\nu})\frac{\partial f_{eq,f}}{\partial p^{0}} + q_{f}\left[F^{0i}p_{i}\frac{\partial f_{eq,f}}{\partial p^{0}} + F^{i0}p_{0}\frac{\partial f_{eq,f}}{\partial p^{i}}\right]
$$

$$
= -\frac{\omega_{f}}{\tau_{fp}}\left[\delta f_{f} - \frac{\langle(\omega_{f}/\tau_{fp})\delta f_{f}\rangle_{0}}{\langle\omega_{f}/\tau_{fp}\rangle_{0}} + P^{(0)}_{1}\frac{\langle(\omega_{f}/\tau_{fp})P^{(0)}_{1}\delta f_{f}\rangle_{0}}{\langle(\omega_{f}/\tau_{fp})P^{(0)}_{1}P^{(0)}_{1}\rangle_{0}} + p^{\langle\mu\rangle}\frac{\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\delta f_{f}\rangle_{0}}{\langle(1/3)\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p^{\langle\mu\rangle}\rangle_{0}}\right],
$$
(51)

where $p_0 = \omega_f - \mu_f$ and $p_0 \approx \omega_f$ for very small μ_f . After solving the above equation one can extract the expression of δf_f (for a detailed calculation see appendix [D\)](#page-32-0) as

$$
\delta f_f = -\frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f})}{TA} \left[\frac{p_0}{T} \partial_0 T + \left(\frac{p_0 - h_f}{p_0} \right) \frac{p^i}{T} \left(\partial_i T - \frac{T}{\varepsilon_{eq,f} + P_f} \partial_i P_f \right) + T \partial_0 \left(\frac{\mu_f}{T} \right) - \frac{p^i p^\nu}{p_0} \partial_i u_\nu - \frac{2q_f}{p_0} \mathbf{E} \cdot \mathbf{p} \right]. \tag{52}
$$

Similarly for antiquarks, we have

$$
\delta \bar{f}_f = -\frac{\tau_{\bar{f}p} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{T\bar{A}} \Big[\frac{p_0}{T} \partial_0 T + \left(\frac{p_0 - \bar{h}_f}{p_0} \right) \frac{p^i}{T} \Big(\partial_i T - \frac{T}{\varepsilon_{eq,f} + P_f} \partial_i P_f \Big) - T \partial_0 \Big(\frac{\mu_f}{T} \Big) - \frac{p^i p^\nu}{p_0} \partial_i u_\nu - \frac{2q_{\bar{f}}}{p_0} \mathbf{E} \cdot \mathbf{p} \Big]. \tag{53}
$$

Finally, replacing the values of δf_f and $\delta \bar{f}_f$ in eq. [\(45\)](#page-9-0) and comparing it with eq. [\(49\)](#page-10-0), we get the final expression of the thermal conductivity in the novel RTA model as

$$
\kappa^{Novel - RTA} = \frac{\beta^2}{6\pi^2} \sum_f g_f \int dp \frac{p^4}{\omega_f^2} \left[\frac{(\omega_f - h_f)^2 \tau_{fp} f_{eq,f} (1 - f_{eq,f})}{A} + \frac{(\omega_f - \bar{h}_f)^2 \tau_{fp} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\bar{A}} \right]
$$
\n(54)

.

In the novel BGK model, the relativistic Boltzmann transport equation takes the following form,

$$
p^{\mu}\partial_{\mu}T\frac{\partial f_{eq,f}}{\partial T} + p^{\mu}\partial_{\mu}(p^{\nu}u_{\nu})\frac{\partial f_{eq,f}}{\partial p^{0}} + q_{f}\left[F^{0i}p_{i}\frac{\partial f_{eq,f}}{\partial p^{0}} + F^{i0}p_{0}\frac{\partial f_{eq,f}}{\partial p^{i}}\right]
$$

\n
$$
= -\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left[\delta f_{f} - \frac{\left\langle(\omega_{f}/\tau_{fp})\delta f_{f}\right\rangle_{0}}{\left\langle\omega_{f}/\tau_{fp}\right\rangle_{0}} + P_{1}^{(0)}\frac{\left\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f}\right\rangle_{0}}{\left\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\right\rangle_{0}} + p^{\langle\mu\rangle}\frac{\left\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\delta f_{f}\right\rangle_{0}}{\left(1/3\right)\left\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p^{\langle\mu\rangle}\right\rangle_{0}}\right]
$$

\n
$$
+ g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\left\langle(\omega_{f}/\tau_{fp})\delta f_{f}\right\rangle_{0}}{\left\langle\omega_{f}/\tau_{fp}\right\rangle_{0}} + P_{1}^{(0)}\frac{\left\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f}\right\rangle_{0}}{\left\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)}\right\rangle_{0}} + p^{\langle\mu\rangle}\frac{\left\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\delta f_{f}\right\rangle_{0}}{\left(1/3\right)\left\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p^{\langle\mu\rangle}\right\rangle_{0}}\right]dP. (55)
$$

Now, neglecting higher order terms $(\mathcal{O}(\delta f_f)^2)$, we get the solution of this equation as

$$
\delta f_f = \delta f_f^{(0)} + g_f n_{eq,f}^{-1} f_{eq,f} \int_{p'} \delta f_f^{(0)}, \tag{56}
$$

where

$$
\delta f_f^{(0)} = -\frac{\tau_{fp}f_{eq,f}(1 - f_{eq,f})}{TA} \left[\frac{p_0}{T} \partial_0 T + \left(\frac{p_0 - h_f}{p_0} \right) \frac{p^i}{T} \left(\partial_i T - \frac{T}{\varepsilon_{eq,f} + P_f} \partial_i P_f \right) + T \partial_0 \left(\frac{\mu_f}{T} \right) - \frac{p^i p^\nu}{p_0} \partial_i u_\nu - \frac{2q_f}{p_0} \mathbf{E} \cdot \mathbf{p} \right].
$$
 (57)

The infinitesimal change in the quark distribution function is obtained (for a detailed calculation see appendix E) as

$$
\delta f_f = -\frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f})}{TA} \left[\frac{p_0}{T} \partial_0 T + \left(\frac{p_0 - h_f}{p_0} \right) \frac{p^i}{T} \left(\partial_i T - \frac{T}{\varepsilon_{eq,f} + P_f} \partial_i P_f \right) + T \partial_0 \left(\frac{\mu_f}{T} \right) - \frac{p^i p^{\nu}}{p_0} \partial_i u_{\nu} \right]
$$

$$
- \frac{2q_f}{p_0} \mathbf{E} \cdot \mathbf{p} \right] - g_f n_{eq,f}^{-1} f_{eq,f} \int_{p'} \frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f})}{TA} \left[\frac{p_0}{T} \partial_0 T + \left(\frac{p_0 - h_f}{p_0} \right) \frac{p^i}{T} \left(\partial_i T - \frac{T}{\varepsilon_{eq,f} + P_f} \partial_i P_f \right) \right]
$$

$$
+ T \partial_0 \left(\frac{\mu_f}{T} \right) - \frac{p^i p^{\nu}}{p_0} \partial_i u_{\nu} - \frac{2q_f}{p_0} \mathbf{E} \cdot \mathbf{p} \right]. \tag{58}
$$

Similarly, $\delta \bar{f}_f$ for antiquark case is obtained as

$$
\delta \bar{f}_f = -\frac{\tau_{fp}\bar{f}_{eq,f}(1-\bar{f}_{eq,f})}{T\bar{A}} \Big[\frac{p_0}{T}\partial_0 T + \left(\frac{p_0-\bar{h}_f}{p_0}\right) \frac{p^i}{T} \Big(\partial_i T - \frac{T}{\varepsilon_{eq,f}+P_f}\partial_i P_f\Big) - T\partial_0 \Big(\frac{\mu_f}{T}\Big) - \frac{p^i p^{\nu}}{p_0}\partial_i u_{\nu}
$$

$$
-\frac{2q_{\bar{f}}}{p_0} \mathbf{E} \cdot \mathbf{p} \Big] - g_f n_{eq,f}^{-1} \bar{f}_{eq,f} \int_{p'} \frac{\tau_{fp}\bar{f}_{eq,f}(1-\bar{f}_{eq,f})}{T\bar{A}} \Big[\frac{p_0}{T}\partial_0 T + \left(\frac{p_0-\bar{h}_f}{p_0}\right) \frac{p^i}{T} \Big(\partial_i T - \frac{T}{\varepsilon_{eq,f}+P_f}\partial_i P_f\Big)
$$

$$
-T\partial_0 \Big(\frac{\mu_f}{T}\Big) - \frac{p^i p^{\nu}}{p_0}\partial_i u_{\nu} - \frac{2q_{\bar{f}}}{p_0} \mathbf{E} \cdot \mathbf{p} \Big]. \tag{59}
$$

Finally, replacing the values of δf_f and $\delta \bar{f}_f$ in eq. [\(45\)](#page-9-0) and comparing it with eq. [\(49\)](#page-10-0), we get the final expression of the thermal conductivity in the novel BGK model as

$$
\kappa^{Novel-BGK} = \frac{\beta^2}{6\pi^2} \sum_{f} g_f \int dp \frac{p^4}{\omega_f^2} \left[\frac{(\omega_f - h_f)^2 \tau_{fp} f_{eq,f} (1 - f_{eq,f})}{A} + \frac{(\omega_f - \bar{h}_f)^2 \tau_{fp} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\bar{A}} \right]
$$

+
$$
\frac{\beta^2}{\sqrt{3}} \sum_{f} g_f^2 n_{eq,f}^{-1} \left[\int_p \frac{p}{\omega_f} (\omega_f - h_f) f_{eq,f} \int_{p_f} \frac{p'}{\omega_f} \frac{(\omega_f - h_f) \tau_{fp} f_{eq,f} (1 - f_{eq,f})}{A} + \int_p \frac{p}{\omega_f} (\omega_f - \bar{h}_f) \bar{f}_{eq,f} \int_{p'} \frac{p'}{\omega_f} \frac{(\omega_f - \bar{h}_f) \tau_{fp} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\bar{A}} \right].
$$
 (60)

1

3 Results and discussions

The interaction of the partons with the thermal medium gives rise to a certain mass known as the thermal mass or the quasiparticle mass. Within the framework of the quasiparticle model, we may see QGP as a system of massive noninteracting quasiparticles. The quasiparticle quark-gluon plasma (qQGP) model is a commonly employed framework to explain the collective properties of the QGP medium. The quasiparticle masses of particles in a dense thermal medium rely on both temperature and chemical potential. For a dense QCD medium, the thermal masses (squared) of quark and gluon up to oneloop are given [\[68,](#page-39-3) [69](#page-39-4)] by

$$
m_{fT}^2 = \frac{g^2 T^2}{6} \left(1 + \frac{\mu_f^2}{\pi^2 T^2} \right),\tag{61}
$$

$$
m_{gT}^2 = \frac{g^2 T^2}{6} \left(N_c + \frac{N_f}{2} + \frac{3}{2\pi^2 T^2} \sum_f \mu_f^2 \right),\tag{62}
$$

respectively, with $g = (4\pi\alpha_s)^{1/2}$, where the expression of α_s is given in eq. [\(18\)](#page-6-0). It can be seen from eq. [\(18\)](#page-6-0) that, α_s depends on both temperature and chemical potential. In this work, we have used $\mu_f = \mu = 0.04$ GeV. The accessible energy scales for a thermal QCD medium are related to the temperature, quark mass and chemical potential. The QGP phase, which is made up of light quarks, light antiquarks and gluons, has a temperature high enough to reach the maximum energy scale. The hard thermal loop (HTL) perturbation theory is one of the effective theories that is applied to tackle the divergences that arise in the computation of the QCD thermodynamic observables, the transport coefficients and the amplitudes in the high temperature regime.

Figure 1: The variation of A (a) as a function of temperature at a fixed momentum and (b) as a function of momentum at a fixed temperature.

Figure 2: Variation of \tilde{A} (a) as a function of temperature at a fixed momentum and (b) as a function of momentum at a fixed temperature.

In this section, the charge and the heat transport coefficients are investigated with the thermal masses as given in equations [\(61\)](#page-12-0) and [\(62\)](#page-12-0). In the novel RTA and the novel BGK models, quantities A and \overline{A} appear in the denominators of the electrical conductivity and the thermal conductivity expressions. Thus the knowledge on the variations of these quantities with temperature and momentum is important to understand the variations of the conductivities. In figures [1](#page-13-0) and [2,](#page-13-1) we have plotted A and \overline{A} for the momentumdependent relaxation time ($\gamma = 0.5$) and for the momentum-independent relaxation time $(\gamma = 0)$, respectively. We have found decreasing trends of A and A with the increase of temperature (figures [1a](#page-13-0) and [2a\)](#page-13-1) and their increasing trends with the increase of momentum

(figures [1b](#page-13-0) and [2b\)](#page-13-1). In both the cases, it is observed that the momentum dependence of the relaxation time enhances both A and \overline{A} . Thus, through the effects of the momentumdependent relaxation time on A and \overline{A} , it is possible to comprehend how the momentumdependent relaxation time would modulate the transport coefficients.

Figure 3: Variation of the electrical conductivity (σ_{el}) with the temperature, (a) comparison with the result in the RTA model and (b) comparison with the result in the BGK model.

Figures [3a](#page-14-0) and [3b](#page-14-0) show the variations of the electrical conductivity (σ_{el}) with the temperature in the novel RTA and the novel BGK models at finite chemical potential, respectively. More precisely, we have compared our results on the electrical conductivity with those of the standard RTA and the standard BGK models. We have also compared the result for $\gamma = 0.5$ with that for $\gamma = 0$. The relaxation time appears to be momentum-dependent (eq. (16)) in the first case, whereas the relaxation time is momentum-independent in the second case. We have observed that, regardless of the relaxation type ($\gamma = 0$ or $\gamma = 0.5$), there is a noticeable reduction in the magnitudes of σ_{el} in the novel RTA and the novel BGK models as compared to those in the RTA and the BGK models, respectively. This reduction in the electrical conductivity in the novel RTA and the novel BGK models is mainly attributed to the quantities A and A appearing in the denominators of equations (24) and (36) .

Figure 4: Variation of the thermal conductivity (κ) with the temperature, (a) comparison with the result in the RTA model and (b) comparison with the result in the BGK model.

$T=0.16$ GeV	σ_{NR} σ _R	σ_{NB}	κ_{NR}	K_{NB} κ_B	$\frac{D_{NR}}{D_{B}}$	D_{NB}	L_{NR}	L_{NB}
$\gamma = 0$					0.50 0.51 0.64 0.78 0.64 0.78 1.26			1.51
$\gamma = 0.5$					0.40 0.45 0.45 0.57 0.45 0.57 1.11			

Table 1: Effect of collision integral on various transport coefficients in the low temperature regime.

$T=1$ GeV			$\begin{array}{ c c c c c c c c c } \hline \sigma_{NR} & \sigma_{RB} & \sigma_{RB} & \sigma_{RR} & \sigma_{RB} & \sigma_{RB} & D_{NR} & D_{RB} & D_{RB} & L_{RB} \ \hline \sigma_{RB} & \sigma_{RB} & \sigma_{RB} & \sigma_{RB} & D_{RB} & D_{RB} & L_{RB} \ \hline \end{array}$		L_{NB}
$\gamma = 0$			$\vert 0.52 \vert 0.54 \vert 0.54 \vert 0.54 \vert 0.68 \vert 0.83 \vert 1.55 \vert 1.30$		
$\gamma=0.5$			$\vert 0.41 \vert 0.46 \vert 0.46 \vert 0.46 \vert 0.47 \vert 0.60 \vert 1.29 \vert 1.26$		

Table 2: Effect of collision integral on various transport coefficients in the high temperature regime.

In figures [4a](#page-15-0) and [4b,](#page-15-0) the thermal conductivity (κ) has been plotted as a function of the temperature in the novel RTA and the novel BGK models at finite chemical potential, respectively. These figures show the comparison of our results on the thermal conductivity with those of the standard RTA and BGK models. We have found that, irrespective of the relaxation type ($\gamma = 0$ or $\gamma = 0.5$), there is a discernible decrease in the magnitudes of κ in the novel RTA and the novel BGK models as compared to those in the RTA and the BGK models, respectively. The quantities A and \overline{A} appearing in the denominators of equations [\(54\)](#page-10-1) and [\(60\)](#page-12-1) mainly contribute in the decrease of the thermal conductivity in the novel RTA and the novel BGK models.

The electrical and thermal conductivities are found to be significantly impacted by the adjustments we made to the RTA model as well as the BGK model when an energydependent relaxation time ($\gamma = 0.5$) is used. Tables [1](#page-15-1) and [2](#page-15-2) show how the updated collision integral affects the charge and heat transfer in general in the low temperature and

the high temperature regimes, respectively. Based on the electrical conductivity related ratios in tables [1](#page-15-1) and [2,](#page-15-2) we can conclude that, for the novel RTA and the novel BGK models, the drop in the magnitude of σ_{el} becomes larger when considering the momentum dependent relaxation time as compared to the standard RTA and BGK models. We may infer from the thermal conductivity related ratios in tables [1](#page-15-1) and [2](#page-15-2) that, as compared to the standard RTA and BGK models, the reduction in the magnitude of κ for the novel RTA and the novel BGK models increases when the momentum dependent relaxation time is taken into account. Thus the momentum dependence of the relaxation time decelerates the flow of charge as well as the flow of heat in the medium.

4 Applications

The purpose of this section is to study some applications of electrical and thermal conductivities for the hot and dense QCD medium. In subsection [4.1,](#page-16-0) we analyze the heat transfer in the medium through the thermal diffusion constant in the novel RTA and the novel BGK models. The relative behavior of the thermal conductivity and the electrical conductivity is explored in subsection [4.2](#page-17-0) using the Lorenz number in the aforesaid models.

4.1 Thermal diffusion constant

Instead of being measured directly, the thermal conductivity in common materials is usually determined from measurements of the thermal diffusion constant (D_T) . This is related to the rate of heat transfer in the medium. The thermal diffusion constant in the first order relativistic viscous hydrodynamics is defined as

$$
D_T = \frac{\kappa}{C_p}.\tag{63}
$$

Here, C_p represents the specific heat at constant pressure and it can be determined by using the following formula.

$$
C_p = \frac{\partial (\varepsilon + P)}{\partial T},\tag{64}
$$

where ε and P are given in equations [\(41\)](#page-9-1) and [\(42\)](#page-9-2), respectively. Increase of the thermal diffusion constant explains the faster heat transfer in the medium.

Figure 5: Variation of the thermal diffusion constant (D_T) with the temperature, (a) comparison with the result in the RTA model and (b) comparison with the result in the BGK model.

Figure [5](#page-17-1) shows the variation of the thermal diffusion constant with the temperature at a fixed chemical potential in the novel RTA and the novel BGK models. It is evident from figures [5a](#page-17-1) and [5b](#page-17-1) that the magnitude of D_T gets decreased regardless of the relaxation type ($\gamma = 0$ or $\gamma = 0.5$) for both RTA and BGK models with the modified integral term. When we compare it with the standard RTA and BGK models, we have found that the momentum-dependent relaxation time ($\gamma = 0.5$) leads to a considerable shift in the thermal diffusion constant. We can conclude from the thermal diffusion constant related ratios in tables [1](#page-15-1) and [2](#page-15-2) that, for the novel RTA and the novel BGK models, the decline in the magnitude of D_T becomes larger when considering the momentumdependent relaxation time as compared to the standard RTA and BGK models.

4.2 Lorenz number

The Lorenz number (L) is the proportionality factor in the Wiedemann-Franz law, which states that the ratio of the heat transport coefficient (κ) to the charge transport coefficient (σ_{el}) is directly proportional to the temperature, *i.e.*,

$$
\frac{\kappa}{\sigma_{el}} = LT.
$$
\n(65)

This law describes how the phenomena of heat transfer and charge transfer behave in relation to one another in a medium.

Figure 6: Variation of the Lorenz number (L) with the temperature, (a) comparison with the result in the RTA model and (b) comparison with the result in the BGK model.

Figure [6](#page-18-0) shows the variation of the Lorenz number with temperature at a fixed chemical potential. It is apparent from figures [6a](#page-18-0) and [6b](#page-18-0) that the Lorenz number exhibits a notable increase in magnitude in case of the novel RTA and the novel BGK models, regardless of the relaxation type ($\gamma = 0$ or $\gamma = 0.5$). A significant change in the Lorenz number is also observed as a result of the momentum dependence of the relaxation time ($\gamma = 0.5$) for both RTA and BGK models. From the Lorenz number related ratios in tables [1](#page-15-1) and [2,](#page-15-2) it is observed that, the increase in the magnitude of L for the novel RTA and the novel BGK models as compared to the normal RTA and BGK models gets smaller when the momentum-dependent relaxation time is used.

5 Summary

In this work, we studied the transport properties of a hot and dense QCD matter in the novel RTA and the novel BGK models. This study could help to get an understanding of the charge and heat transport phenomena in a strongly coupled thermal QCD medium with a finite chemical potential background through the novel approaches of the aforesaid models. Using the kinetic theory technique, we solved the relativistic Boltzmann transport equation with a modified collision integral for estimating the thermal and electrical conductivities of the aforementioned medium. We examined the effects of the modified collision integral on the charge and heat transport in a thermal QCD medium, in contrast to the widely used collision terms of the RTA and the BGK models. In this research, we used the quasiparticle description of partons, where both the temperature and the chemical potential affect the constituent masses in the medium. Our modifications to the

collision integral for the RTA and the BGK models were shown to result in a considerable drop in the electrical and thermal conductivities. Additionally, we found that these transport coefficients exhibit a discernible shift when we take into account the momentumdependent relaxation time ($\gamma = 0.5$). In the same regime, the thermal diffusion constant and the Lorenz number were also investigated. We observed that the novel collision integral leads to an increase in the Lorenz number and a decrease in the thermal diffusion constant of the medium.

6 Acknowledgments

One of the present authors (S. R.) acknowledges financial support from ANID Fondecyt postdoctoral grant 3240349 and S. D. acknowledges the SERB Power Fellowship, SPF/2022/000014 for the support on this work.

Appendices

A Derivation of equation [\(20\)](#page-6-1)

Using the result of ref. [\[64](#page-39-0)], one can write the relativistic Boltzmann transport equation as

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\frac{\omega_f}{\tau_{fp}} \left[\delta f_f - \frac{\langle (\omega_f/\tau_{fp}) \delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp}) P_1^{(0)} \delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \delta f_f \rangle_0}{\langle (1/3) \langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} \right].
$$
 (A.66)

The second term in the right-hand side can be reduced to

$$
\frac{\langle (\omega_f/\tau_{fp}) \delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} = (1/\langle \omega_f/\tau_{fp} \rangle_0) \int dP \frac{\omega_f}{\tau_{fp}} f_{eq,f} \delta f_f = (1/\langle \omega_f/\tau_{fp} \rangle_0)
$$

$$
\times \left[\delta f_f \int dP \frac{\omega_f}{\tau_{fp}} f_{eq,f} - \int \left[\frac{d(\delta f_f)}{dP} \int dP \frac{\omega_f}{\tau_{fp}} f_{eq,f} \right] dP \right]
$$

$$
= (1/\langle \omega_f/\tau_{fp} \rangle_0) \delta f_f \left(\langle \omega_f/\tau_{fp} \rangle_0 \right) = \delta f_f. \quad (A.67)
$$

Here, we have used the fact that the shift of the distribution function from its equilibrium value is infinitesimal, thus one can use the derivative $\frac{d(\delta f_f)}{dP} \approx 0$. Similarly, the third and the fourth terms can be respectively reduced to

$$
P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp}) P_1^{(0)} \delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \rangle_0} = P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp}) P_1^{(0)} \rangle_0}{\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \rangle_0} \delta f_f,
$$
(A.68)

$$
p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \delta f_f \rangle_0}{(1/3) \langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} = p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \rangle_0}{(1/3) \langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} \delta f_f.
$$
 (A.69)

Now, with the help of equations [\(A.67\)](#page-19-1), [\(A.68\)](#page-20-0) and [\(A.69\)](#page-20-1), eq. [\(A.66\)](#page-19-2) takes the following form,

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\frac{\omega_f}{\tau_{fp}} \bigg[P_1^{(0)} \frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} \delta f_f
$$

$$
+ p^{\langle \mu \rangle} \frac{\left\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \right\rangle_0}{(1/3) \left\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \right\rangle_0} \delta f_f \bigg] = -\frac{\omega_f A}{\tau_{fp}} \delta f_f, \quad (A.70)
$$

where,

$$
A = \left[1 - \frac{\omega_f \int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right] \frac{\int p^2 (f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right) \omega_f\right] dp}{\int p^2 (f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right) \omega_f\right]^2 dp} + \frac{3p \int p^3 (f_{eq,f}/\tau_{fp}) dp}{\int p^4 (f_{eq,f}/\tau_{fp}) dp}.
$$
 (A.71)

Using the following partial derivatives,

$$
\frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\frac{\mathbf{p}}{\omega_f} \beta f_{eq,f} (1 - f_{eq,f}),
$$
\n
$$
\frac{\partial f_{eq,f}}{\partial p^0} = -\beta f_{eq,f} (1 - f_{eq,f}),
$$
\n(A.72)

we solve eq. [\(A.70\)](#page-20-2) to get the expression of δf_f as given in eq. [\(20\)](#page-6-1).

B The particle number and the energy-momentum conservation laws

In this study, we have introduced a novel collision integral in the BGK model of the following form,

$$
\mathbf{C}[f_f]^{NBGK} = -\frac{p^{\mu}u_{\mu}}{\tau_{fp}} \Bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{\langle (1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \Bigg] + g_f n_{eq,f}^{-1}(p^{\mu}u_{\mu}) f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \Bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{\langle (1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \Bigg] dP. \quad (B.73)
$$

We will be subject to the following constraints if we insist that the conservation laws be upheld.

$$
\partial_{\mu}N^{\mu} = \int \mathbf{C} \left[f_f\right]^{NBGK} dP = 0, \tag{B.74}
$$

$$
\partial_{\mu}T^{\mu\nu} = \int p^{\nu}\mathbf{C}[f_f]^{NBGK}dP = 0.
$$
 (B.75)

Validity of the particle number conservation law in the novel BGK model:

$$
\partial_{\mu}N^{\mu} = \int \mathbf{C}[f_{f}]^{NBGK}dP = -\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \Bigg[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\gamma_{0}^{(0)} \rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\gamma_{0}^{(0)} \rangle_{0}} \Bigg] dP + \int \Bigg[g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \Bigg[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\gamma_{0}^{(0)} \rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\gamma_{0}^{(0)} \rangle_{0}} \Bigg] dP \Bigg] dP
$$

\n
$$
= - \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \Bigg[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\gamma_{0}^{(0)} \rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{
$$

where,

$$
T1 = -\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{\langle 1/3 \rangle \langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \right] dP, \quad (B.77)
$$

$$
T2 = \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{(1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}^{(0)}p^{\langle \mu \rangle_{0}} \rangle_{0}} \left] dP \times \int g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f}dP = \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}^{(0)}\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{(1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}^{(0)}p^{\langle \mu \rangle_{0}} \rangle_{0}} \left] dP \times n_{eq,f}^{-1} \left(g_{f} \int (p^{\mu}u_{\mu})f_{eq,f}dP \right) = \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{P_{1}}^{(0)}p^{\langle \mu \rangle_{0}} \rangle_{0}} \right] dP \times n_{eq,f}^{-1} \left(n_{eq,f} \right) \qquad \left[\text{Using eq. (28)} \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p
$$

$$
T3 = -\int \left[\frac{d}{dP} \left[\int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp}) \delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} P_{1}^{(0)} \rangle_{0}} \right] \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} \delta f_{f} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} dP \right] f g_{f} n_{eq, f}^{-1} (p^{\mu} u_{\mu}) f_{eq, f} dP \right] dP.
$$

= 0. [Due to the $\mathcal{O}((\delta f_{f})^{2})$] (B.79)

Now, putting the results of equations [\(B.77\)](#page-22-0), [\(B.78\)](#page-23-0) and [\(B.79\)](#page-23-1) in eq. [\(B.76\)](#page-22-1), we get

$$
\partial_{\mu}N^{\mu} = -\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} \right] \n+ p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{\langle (1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_{0}} \right] dP + \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} \right] \n+ p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{\langle (1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_{0}} \right] dP
$$
\n
$$
= 0.
$$
\n(B.80)

From eq. [\(B.80\)](#page-24-0), it is confirmed that the particle number is instantaneously conserved in the novel BGK model.

Validity of the energy-momentum conservation law in the novel BGK model:

$$
\partial_{\mu}T^{\mu\nu} = \int p^{\nu} \mathbf{C}[f_{f}]^{NBGK}dP = -\int p^{\nu}\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}}\right]dP + \int \left[p^{\nu}g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})p_{\mu\nu}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})p_{\mu\nu}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})p_{\mu\nu}\delta f_{f} \rangle_{0}}\right]dP\right]dP
$$

\n
$$
= -\int p^{\nu}\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})p_{\mu\nu}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})p_{\mu}\delta f_{f} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})p_{\mu}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})p_{\mu}\delta f_{f} \rangle_{0}}\right]dP
$$

where $S1^{\nu}$, $S2^{\nu}$ and $S3^{\nu}$ are respectively expressed as

$$
S1^{\nu} = -\int p^{\nu} \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp}) \delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} P_{1}^{(0)} \rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} \delta f_{f} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} \right] dP
$$

\n
$$
= -\int p^{\nu} \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \delta f_{f} dP + \int p^{\nu} \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \frac{\langle (\omega_{f}/\tau_{fp}) \delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} dP
$$

\n
$$
- \int p^{\nu} \frac{p^{\mu} u_{\mu}}{\tau_{fp}} P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \rangle_{0}} dP
$$

\n
$$
- \int p^{\nu} \frac{p^{\mu} u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} \delta f_{f} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} dP
$$

\n= S11^u + S12^u + S13^u + S14^u, (B.82)

$$
S2^{\nu} = \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{(1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_{0}} \left] dP \times \int p^{\nu}g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f}dP \right.
$$

\n
$$
= \left[\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \delta f_{f}dP - \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} dP + \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} dP \right.
$$

\n
$$
+ \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{(1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_{0}} dP \right] \int p^{\nu}g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f}dP
$$

\n= (S21 + S22 + S23 + S24)S20°,

$$
(B.83)
$$

$$
S3^{\nu} = -\int \left[\frac{d}{dP} \left[\int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp}) \delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} P_{1}^{(0)} \rangle_{0}} \right] \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} \delta f_{f} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} \right] dP \right] \int p^{\nu} g_{f} n_{eq,f}^{-1}(p^{\mu} u_{\mu}) f_{eq,f} dP \right] dP. \quad (B.84)
$$

The energy-momentum conservation requires that,

$$
u_{\nu} \partial_{\mu} T^{\mu \nu} = 0
$$

\n
$$
\implies u_{\nu} (S1^{\nu} + S2^{\nu} + S3^{\nu}) = 0.
$$
 (B.85)

The term $u_{\nu}S1^{\nu}$ can be calculated as follows.

$$
u_{\nu}S1^{\nu} = u_{\nu} \left(S11^{\nu} + S12^{\nu} + S13^{\nu} + S14^{\nu} \right), \tag{B.86}
$$

where,

$$
u_{\nu}S11^{\nu} = -u_{\nu}\int p^{\nu}\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\delta f_f dP,
$$
\n(B.87)

$$
u_{\nu}S12^{\nu} = u_{\nu} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} dP = u_{\nu} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \frac{\langle (\omega_f/\tau_{fp}) \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} \delta f_f dP
$$

= $u_{\nu} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \delta f_f dP$, [Using eq. (A.67)] (B.88)

$$
u_{\nu}S13^{\nu} = -u_{\nu} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} P_{1}^{(0)} \frac{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \delta f_{f} \right\rangle_{0}}{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} P_{1}^{(0)} \right\rangle_{0}} dP
$$
\n
$$
= -u_{\nu} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} P_{1}^{(0)} \frac{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}}{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}} \delta f_{f} dP \qquad \qquad \text{[Using eq. (A.68)]}
$$
\n
$$
= -u_{\nu} \frac{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}}{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} P_{1}^{(0)} \delta f_{f} dP
$$
\n
$$
+ u_{\nu} \int \left[\frac{d}{dP} \left[\frac{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}}{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}} \right] \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} P_{1}^{(0)} \delta f_{f} dP \right] dP
$$
\n
$$
= -u_{\nu} \frac{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}}{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} P_{1}^{(0)} \delta f_{f} dP, \qquad \text{[2nd term vanishes due to } \mathcal{O}((\delta f_{f})^{2})\text{]}
$$
\n(B.89)

$$
u_{\nu}S14^{\nu} = -u_{\nu} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} \delta f_{f} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} dP
$$

\n
$$
= -u_{\nu} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} \delta f_{f} dP \qquad \qquad \text{[Using eq. (A.69)]}
$$

\n
$$
= -u_{\nu} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \delta f_{f} dP
$$

\n
$$
+u_{\nu} \int \left[\frac{d}{dP} \left[\frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} \right] \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \delta f_{f} dP \right] dP
$$

\n
$$
= -u_{\nu} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} \int p^{\nu} \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \delta f_{f} dP. \quad \text{[2nd term vanishes due to } \mathcal{O}((\delta f_{f})^{2})]
$$

\n(B.90)

Now, the term $u_{\nu}S1^{\nu}$ turns out to be

$$
u_{\nu}S1^{\nu} = -\left[u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)}\rangle_{0}}\int p^{\nu}\frac{p^{\mu}u_{\mu}}{\tau_{fp}}P_{1}^{(0)}\delta f_{f}dP\right.+u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\rangle_{0}}{\langle1/3\rangle\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\rangle_{0}}\int p^{\nu}\frac{p^{\mu}u_{\mu}}{\tau_{fp}}p^{\langle\mu\rangle}\delta f_{f}dP\right]= -\left[u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)}\rangle_{0}}\int p^{\nu}\frac{p^{\mu}u_{\mu}}{\tau_{fp}}P_{1}^{(0)}\phi_{f}f_{eq,f}dP\right.+u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\rangle_{0}}{\langle1/3\rangle\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p^{\langle\mu\rangle}\rangle_{0}}\int p^{\nu}\frac{p^{\mu}u_{\mu}}{\tau_{fp}}p^{\langle\mu\rangle}\phi_{f}f_{eq,f}dP\right],= -\left[\frac{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}\langle\omega_{f}P_{1}^{(0)}\phi_{f}\rangle+\frac{\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\rangle_{0}}{\langle1/3\rangle\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p^{\langle\mu\rangle}\rangle_{0}}\langle\omega_{f}p^{\langle\mu\rangle}\phi_{f}\rangle\right].(B.91)
$$

In getting the above equation, we have used the following substitutions (in this case, f_f is expressed in terms of an equilibrium contribution $(f_{eq,f})$ and a nonequilibrium correction (ϕ_f) as $f_f \equiv (1 + \phi_f) f_{eq,f}),$

$$
\delta f_f = \phi_f f_{eq,f},\tag{B.92}
$$

$$
\left\langle \omega_f P_1^{(0)} \phi_f \right\rangle = \int \left(\omega_f / \tau_{fp} \right) \omega_f P_1^{(0)} \phi_f f_{eq,f} dP, \tag{B.93}
$$

$$
\langle \omega_f p^{\langle \mu \rangle} \phi_f \rangle = \int (\omega_f / \tau_{fp}) \, \omega_f p^{\langle \mu \rangle} \phi_f f_{eq,f} dP. \tag{B.94}
$$

The term $u_{\nu}S2^{\nu}$ can be calculated as follows.

$$
u_{\nu}S2^{\nu} = u_{\nu}(S21 + S22 + S23 + S24) S20^{\nu}, \tag{B.95}
$$

where,

$$
u_{\nu}S20^{\nu} = u_{\nu} \int p^{\nu}(p^{\mu}u_{\mu})g_{f}n_{eq,f}^{-1}f_{eq,f}dP
$$

= $n_{eq,f}^{-1}g_{f}u_{\nu} \int p^{\nu}(p^{\mu}u_{\mu})f_{eq,f}dP = \frac{\varepsilon_{eq,f}}{n_{eq,f}},$ [Using eq. (47)] (B.96)

$$
S21 = \int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \delta f_f dP, \tag{B.97}
$$

$$
S22 = -\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} dP = -\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \frac{\langle (\omega_f/\tau_{fp}) \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} \delta f_f dP
$$

$$
= -\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \delta f_f dP, \qquad \qquad \text{[Using eq. (A.67)]} \quad (B.98)
$$

$$
S23 = \int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} P_1^{(0)} \frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \delta f_f \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} dP
$$

\n
$$
= \int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} P_1^{(0)} \frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} \delta f_f dP \qquad \text{[Using eq. (A.68)]}
$$

\n
$$
= \frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} \int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} P_1^{(0)} \delta f_f dP
$$

\n
$$
- \int \left[\frac{d}{dP} \left[\frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} \right] \int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} P_1^{(0)} \delta f_f dP \right] dP
$$

\n
$$
= \frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} \int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} P_1^{(0)} \delta f_f dP, \qquad \text{[2nd term vanishes due to } \mathcal{O}((\delta f_f)^2)] \qquad (B.99)
$$

$$
S24 = \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{(1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\rho^{\langle \mu \rangle} \rangle_0} dP
$$

\n
$$
= \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\rho^{\langle \mu \rangle} \rangle_0}{(1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\rho^{\langle \mu \rangle} \rangle_0} \delta f_f dP
$$
 [Using eq. (A.69)]
\n
$$
= \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\rho^{\langle \mu \rangle} \rangle_0}{(1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\rho^{\langle \mu \rangle} \rangle_0} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \delta f_f dP
$$

\n
$$
- \int \left[\frac{d}{dP} \left[\frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle} \rangle_0}{(1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\rho^{\langle \mu \rangle} \rangle_0} \right] \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \delta f_f dP \right] dP
$$

\n
$$
= \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle} \rangle_0}{(1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\rho^{\langle \mu \rangle} \rangle_0} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} p^{\langle \mu \rangle} \delta f_f dP.
$$
 [2nd term vanishes due to $\mathcal{O}((\delta f_f)^2)$] (B.100)

Thus the term $u_{\nu}S2^{\nu}$ becomes

$$
u_{\nu}S2^{\nu} = \left[u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)}\rangle_{0}}\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}}P_{1}^{(0)}\delta f_{f}dP + u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\rangle_{0}}{\langle(1/3)\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p_{\langle\mu\rangle}\rangle_{0}}\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}}p^{\langle\mu\rangle}\delta f_{f}dP\right]\frac{\epsilon_{eq,f}}{n_{eq,f}}
$$

$$
= \left[u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)}\rangle_{0}}\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}}P_{1}^{(0)}\phi_{f}f_{eq,f}dP + u_{\nu}\frac{\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\rangle_{0}}{\langle(1/3)\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p_{\langle\mu\rangle}\rangle_{0}}\int \frac{p^{\mu}u_{\mu}}{\tau_{fp}}p^{\langle\mu\rangle}\phi_{f}f_{eq,f}dP\right]\frac{\epsilon_{eq,f}}{n_{eq,f}}
$$

$$
= \left[\frac{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}{\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\rangle_{0}}\langle P_{1}^{(0)}\phi_{f}\rangle + \frac{\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\rangle_{0}}{\langle(1/3)\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p_{\langle\mu\rangle}\rangle_{0}}\langle p^{\langle\mu\rangle}\phi_{f}\rangle\right]\frac{\epsilon_{eq,f}}{n_{eq,f}}.
$$
(B.101)

We have obtained the above equation by using the following substitutions,

$$
\delta f_f = \phi_f f_{eq,f},\tag{B.102}
$$

$$
\left\langle P_1^{(0)}\phi_f \right\rangle = \int \left(\omega_f / \tau_{fp} \right) P_1^{(0)} \phi_f f_{eq,f} dP, \tag{B.103}
$$

$$
\langle p^{\langle \mu \rangle} \phi_f \rangle = \int \left(\omega_f / \tau_{fp} \right) p^{\langle \mu \rangle} \phi_f f_{eq,f} dP. \tag{B.104}
$$

The term $u_{\nu}S3^{\nu}$ can be calculated as follows.

$$
u_{\nu} S3^{\nu} = -u_{\nu} \int \left[\frac{d}{dP} \left[\int \frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp}) \delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} P_{1}^{(0)} \rangle_{0}} \right] \right] + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} \delta f_{f} \rangle_{0}}{(1/3) \langle (\omega_{f}/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} \right] dP \left[\int p^{\nu} g_{f} n_{eq,f}^{-1} (p^{\mu} u_{\mu}) f_{eq,f} dP \right] dP
$$

= 0. \qquad \left[\text{Due to } \mathcal{O}((\delta f_{f})^{2}) \right] (B.105)

Now, substituting the results of equations [\(B.91\)](#page-27-0), [\(B.101\)](#page-29-0) and [\(B.105\)](#page-30-0) in eq. [\(B.85\)](#page-26-0), we get

$$
\left[\frac{\left\langle (\omega_f/\tau_{fp})P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \right\rangle_0} \left\langle \omega_f P_1^{(0)} \phi_f \right\rangle + \frac{\left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle} \right\rangle_0}{(1/3) \left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \right\rangle_0} \left\langle \omega_f p^{\langle \mu \rangle} \phi_f \right\rangle \right] -\left[\frac{\left\langle (\omega_f/\tau_{fp})P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp})P_1^{(0)} P_1^{(0)} \right\rangle_0} \left\langle P_1^{(0)} \phi_f \right\rangle + \frac{\left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle} \right\rangle_0}{(1/3) \left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \right\rangle_0} \left\langle p^{\langle \mu \rangle} \phi_f \right\rangle \right] \frac{\varepsilon_{eq,f}}{n_{eq,f}} = 0.
$$
\n(B.106)

The above equation serves as the matching condition for the novel BGK model and can be expressed as

$$
Q2 n_{eq,f} = Q1 \varepsilon_{eq,f}, \tag{B.107}
$$

where,

$$
Q1 = \left[\frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} \left\langle P_1^{(0)} \phi_f \right\rangle + \frac{\left\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \right\rangle_0}{(1/3) \left\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \right\rangle_0} \left\langle p^{\langle \mu \rangle} \phi_f \right\rangle \right], \quad (B.108)
$$

$$
Q2 = \left[\frac{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} \right\rangle_0}{\left\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \right\rangle_0} \left\langle \omega_f P_1^{(0)} \phi_f \right\rangle + \frac{\left\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \right\rangle_0}{(1/3) \left\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \right\rangle_0} \left\langle \omega_f p^{\langle \mu \rangle} \phi_f \right\rangle \right].
$$
\n(B.109)

In the present scenario, Q1 and Q2 represent the constraints for the conservation of the particle number and the energy-momentum, respectively. The consistency of our formulation within the generalised hydrodynamic frame described in ref. [\[64](#page-39-0)] may be readily observed from equations [\(B.108\)](#page-30-1) and [\(B.109\)](#page-30-2).

C Derivation of equation [\(34\)](#page-8-2)

In the BGK model, the relativistic Boltzmann transport equation is written as

$$
q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\frac{p^\mu u_\mu}{\tau_{fp}} \left(f_f(x, p, t) - \frac{n_f(x, t)}{n_{eq,f}} f_{eq,f}(p) \right), \tag{C.110}
$$

where,

$$
-\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left(f_{f}(x,p,t)-\frac{n_{f}(x,t)}{n_{eq,f}}f_{eq,f}(p)\right) = -\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left(f_{f}(x,p,t)-\frac{g_{f}\int_{p}(f_{eq,f}+\delta f_{f})}{n_{eq,f}}f_{eq,f}(p)\right)
$$

$$
=-\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left(f_{f}(x,p,t)-\frac{(g_{f}\int_{p}f_{eq,f}+g_{f}\int_{p}\delta f_{f})}{n_{eq,f}}f_{eq,f}(p)\right)
$$

$$
=-\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left(\delta f_{f}-g_{f}n_{eq,f}^{-1}f_{eq,f}\int_{p}\delta f_{f}\right).
$$
 (C.111)

Now, following the formulation of ref. [\[64](#page-39-0)] in the novel BGK model, we get

$$
\mathbf{C}[f_f]^{NBGK} = -\frac{p^{\mu}u_{\mu}}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{\langle (1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \bigg] + g_f n_{eq,f}^{-1} (p^{\mu}u_{\mu}) f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{\langle (1/3)\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \bigg] dP. \quad \text{(C.112)}
$$

Using eq. $(C.112)$ and the partial derivatives as given in eq. $(A.72)$, eq. $(C.110)$ takes the following form,

$$
\frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{\langle (1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_{0}} \right]
$$

$$
- g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{\langle 1/3\rangle \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_{0}} \right] dP = 2q_{f} \beta(\mathbf{E} \cdot \mathbf{p}) f_{eq,f} (1 - f_{eq,f}). \quad \text{(C.113)}
$$

For a proper treatment of the left-hand side of the above equation, we are considering an infinitesimal perturbation to the equilibrium distribution function, *i.e.* $f_f = f_{eq,f} + \delta f_f$ with $\delta f_f \ll f_{eq,f}$ and after linearizing it, we get

$$
\frac{\omega_f A}{\tau_{fp}} \delta f_f^{(0)} = 2q_f \beta (\mathbf{E} \cdot \mathbf{p}) f_{eq,f} (1 - f_{eq,f}), \tag{C.114}
$$

where $\delta f_f^{(0)}$ is related to the infinitesimal change of the quark distribution function (δf_f) by the relation given in eq. [\(32\)](#page-8-3). After putting the value of $\delta f_f^{(0)}$ in eq. (32), we get δf_f as given in eq. [\(34\)](#page-8-2).

D Derivation of equation (52)

With the help of equations $(A.67)$, $(A.68)$ and $(A.69)$, eq. (51) can be rewritten as

$$
p^{\mu}\partial_{\mu}T\frac{\partial f_{eq,f}}{\partial T} + p^{\mu}\partial_{\mu}(p^{\nu}u_{\nu})\frac{\partial f_{eq,f}}{\partial p^{0}} + q_{f}\left[F^{0i}p_{i}\frac{\partial f_{eq,f}}{\partial p^{0}} + F^{i0}p_{0}\frac{\partial f_{eq,f}}{\partial p^{i}}\right]
$$

$$
= -\frac{\omega_{f}}{\tau_{fp}}\left[P_{1}^{(0)}\frac{\left\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}\right\rangle_{0}}{\left\langle(\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)}\right\rangle_{0}}\delta f_{f}\right]
$$

$$
+ p^{\langle\mu\rangle}\frac{\left\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}\right\rangle_{0}}{(1/3)\left\langle(\omega_{f}/\tau_{fp})p_{\langle\mu\rangle}p^{\langle\mu\rangle}\right\rangle_{0}}\delta f_{f}\right] = -\frac{\omega_{f}A}{\tau_{fp}}\delta f_{f}.
$$
 (D.115)

Using the following partial derivatives,

$$
\frac{\partial f_{eq,f}}{\partial T} = -\beta^2 \omega_f f_{eq,f} (1 - f_{eq,f})
$$
\n
$$
\frac{\partial f_{eq,f}}{\partial p^0} = -\beta f_{eq,f} (1 - f_{eq,f})
$$
\n
$$
\frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\beta \frac{\mathbf{p}}{\omega_f} f_{eq,f} (1 - f_{eq,f})
$$
\n(D.116)

we solve eq. [\(D.115\)](#page-32-2) to get the expression of δf_f as given in eq. [\(52\)](#page-10-2).

E Derivation of equation [\(58\)](#page-11-0)

In the BGK model, the relativistic Boltzmann transport equation can be written as

$$
p^{\mu} \partial_{\mu} T \frac{\partial f_{eq,f}}{\partial T} + p^{\mu} \partial_{\mu} (p^{\nu} u_{\nu}) \frac{\partial f_{eq,f}}{\partial p^{0}} + q_{f} \left[F^{0i} p_{i} \frac{\partial f_{eq,f}}{\partial p^{0}} + F^{i0} p_{0} \frac{\partial f_{eq,f}}{\partial p^{i}} \right]
$$

$$
= -\frac{p^{\mu} u_{\mu}}{\tau_{fp}} \left(f_{f}(x, p, t) - \frac{n_{f}(x, t)}{n_{eq,f}} f_{eq,f}(p) \right). \quad (E.117)
$$

With the help of eq. $(C.111)$, eq. $(C.112)$ and the partial derivatives as given in eq. [\(D.116\)](#page-32-3), we can express the above equation in the novel BGK model as

$$
\frac{p^{\mu}u_{\mu}}{\tau_{fp}}\left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)} \rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{\langle 1/3 \rangle \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} \rho^{\langle \mu \rangle} \rangle_{0}}\right]
$$

$$
- g_{f}n_{eq,f}^{-1}(p^{\mu}u_{\mu})f_{eq,f} \int \frac{p^{\mu}u_{\mu}}{\tau_{fp}} \left[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f} \rangle_{0}}{\langle \omega_{f}/\tau_{fp} \rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f} \rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)} \rangle_{0}}\right]
$$

$$
+ p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f} \rangle_{0}}{\langle 1/3 \rangle \langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_{0}} dP
$$

$$
= -\frac{f_{eq,f}(1 - f_{eq,f})p_{0}}{T} \left[\frac{p_{0}}{T} \partial_{0}T + \left(\frac{p_{0} - h_{f}}{p_{0}}\right) \frac{p^{i}}{T} \left(\partial_{i}T - \frac{T}{\varepsilon_{eq,f} + P_{f}} \partial_{i}P_{f}\right) + T \partial_{0} \left(\frac{\mu_{f}}{T}\right) - \frac{p^{i}p^{\nu}}{p_{0}} \partial_{i}u_{\nu} - \frac{2q_{f}}{p_{0}} \mathbf{E}.\mathbf{p}\right].
$$
(E.118)

After some mathematical calculation, we get

$$
\frac{\omega_f A}{\tau_{fp}} \delta f_f^{(0)} = -\frac{\omega_f f_{eq,f} (1 - f_{eq,f})}{T} \left[\frac{p_0}{T} \partial_0 T + \left(\frac{p_0 - h_f}{p_0} \right) \frac{p^i}{T} \left(\partial_i T - \frac{T}{\varepsilon_{eq,f} + P_f} \partial_i P_f \right) \right. \\
\left. + T \partial_0 \left(\frac{\mu_f}{T} \right) - \frac{p^i p^\nu}{p_0} \partial_i u_\nu - \frac{2q_f}{p_0} \mathbf{E} \cdot \mathbf{p} \right], \quad (E.119)
$$

where $\delta f_f^{(0)}$ is related to the infinitesimal change of the distribution function (δf_f) by the relation given in eq. [\(56\)](#page-11-1). After putting the value of $\delta f_f^{(0)}$ in eq. (56), the value of δf_f , *i.e.* eq. [\(58\)](#page-11-0) is obtained.

References

- [1] V. Skokov, A. Y. Illarionov and V. Toneev, "Estimate of the magnetic field strength in heavy-ion collisions", Int. J. Mod. Phys. A 24 (2009), 5925-5932 [arXiv:0907.1396 [nucl-th]].
- [2] A. Bzdak and V. Skokov, "Event-by-event fluctuations of magnetic and electric fields in heavy ion collisions", Phys. Lett. B 710 (2012), 171-174 [arXiv:1111.1949 [hep-ph]].
- [3] P. Moreau, O. Soloveva, I. Grishmanovskii, V. Voronyuk, L. Oliva, T. Song, V. Kireyeu, G. Coci and E. Bratkovskaya, "Properties of the quark-gluon plasma created in heavy-ion collisions", Astron. Nachr. 342 (2021) no.5, 715-726 [arXiv:2101.05688 [nucl-th]].
- [4] M. He, H. van Hees and R. Rapp, "Heavy-quark diffusion in the quark–gluon plasma", Prog. Part. Nucl. Phys. 130 (2023), 104020 [arXiv:2204.09299 [hep-ph]].
- [5] S. Mitra, "Relativistic hydrodynamics with momentum dependent relaxation time", Phys. Rev. C 103 (2021) no.1, 014905 [arXiv:2009.06320 [nucl-th]].
- [6] S. Mitra, "Correspondence between momentum-dependent relaxation time and field redefinition of relativistic hydrodynamic theory", Phys. Rev. C 105 (2022) no.1, 014902 [arXiv:2104.11380 [nucl-th]].
- [7] S. Mitra, "Causality and stability analysis of first-order field redefinition in relativistic hydrodynamics from kinetic theory", Phys. Rev. C 105 (2022) no.5, 054910 [arXiv:2106.08510 [nucl-th]].
- [8] F. Scardina, S. K. Das, V. Minissale, S. Plumari and V. Greco, "Estimating the charm quark diffusion coefficient and thermalization time from D meson spectra at energies available at the BNL Relativistic Heavy Ion Collider and the CERN Large Hadron Collider", Phys. Rev. C 96 (2017) no.4, 044905 [arXiv:1707.05452 [nucl-th]].
- [9] S. K. Das, J. e. Alam and P. Mohanty, "Dragging Heavy Quarks in Quark Gluon Plasma at the Large Hadron Collider", Phys. Rev. C 82 (2010), 014908 [arXiv:1003.5508 [nucl-th]].
- [10] P. Khowal, S. K. Das, L. Oliva and M. Ruggieri, "Heavy quarks in the early stage of high energy nuclear collisions at RHIC and LHC: Brownian motion versus diffusion in the evolving Glasma", Eur. Phys. J. Plus 137 (2022) no.3, 307 [arXiv:2110.14610 [hep-ph]].
- [11] A. Viñales and M. Desposito, "Anomalous diffusion induced by a Mittag-Leffler correlated noise", *Physical Review E*. 75, 042102 (2007) [arXiv:0903.1050v2 [condmat.stat-mech.]].
- [12] S. K. Das, J. M. Torres-Rincon, L. Tolos, V. Minissale, F. Scardina and V. Greco, "Propagation of heavy baryons in heavy-ion collisions", Phys. Rev. D 94 (2016) no.11, 114039 [arXiv:1604.05666 [nucl-th]].
- [13] S. Bhadury, W. Florkowski, A. Jaiswal, A. Kumar and R. Ryblewski, "Dissipative Spin Dynamics in Relativistic Matter", Phys. Rev. D 103 (2021) no.1, 014030 [arXiv:2008.10976 [nucl-th]].
- [14] S. Bhadury, W. Florkowski, A. Jaiswal, A. Kumar and R. Ryblewski, "Relativistic dissipative spin dynamics in the relaxation time approximation", Phys. Lett. B 814 (2021), 136096 [arXiv:2002.03937 [hep-ph]].
- [15] Y. Aoki, S. Borsanyi, S. Durr, Z. Fodor, S. D. Katz, S. Krieg and K. K. Szabo, "The QCD transition temperature: results with physical masses in the continuum limit II.", JHEP 06 (2009), 088 [arXiv:0903.4155 [hep-lat]].
- [16] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz and K. K. Szabo, "The Order of the quantum chromodynamics transition predicted by the standard model of particle physics", Nature 443 (2006), 675-678 [arXiv:hep-lat/0611014 [hep-lat]].
- [17] K. Fukushima, D. E. Kharzeev and H. J. Warringa, "The Chiral Magnetic Effect", Phys. Rev. D 78 (2008), 074033 [arXiv:0808.3382 [hep-ph]].
- [18] D. E. Kharzeev, L. D. McLerran and H. J. Warringa, "The Effects of topological charge change in heavy ion collisions: 'Event by event P and CP violation'", Nucl. Phys. A 803 (2008), 227-253 [arXiv:0711.0950 [hep-ph]].
- [19] D. E. Kharzeev and D. T. Son, "Testing the chiral magnetic and chiral vortical effects in heavy ion collisions", Phys. Rev. Lett. 106 (2011), 062301 [arXiv:1010.0038 [hep-ph]].
- [20] S. Pu, S. Y. Wu and D. L. Yang, "Chiral Hall Effect and Chiral Electric Waves", Phys. Rev. D 91 (2015) no.2, 025011 [arXiv:1407.3168 [hep-th]].
- [21] S. Fayazbakhsh and N. Sadooghi, "Weak decay constant of neutral pions in a hot and magnetized quark matter", Phys. Rev. D 88 (2013) no.6, 065030 [arXiv:1306.2098 [hep-ph]].
- [22] S. Fayazbakhsh, S. Sadeghian and N. Sadooghi, "Properties of neutral mesons in a hot and magnetized quark matter", Phys. Rev. D 86 (2012), 085042 [arXiv:1206.6051 [hep-ph]].
- [23] H. van Hees, C. Gale and R. Rapp, "Thermal Photons and Collective Flow at the Relativistic Heavy-Ion Collider", Phys. Rev. C 84 (2011), 054906 [arXiv:1108.2131 [hep-ph]].
- [24] C. Shen, U. W. Heinz, J. F. Paquet and C. Gale, "Thermal photons as a quarkgluon plasma thermometer reexamined", Phys. Rev. C 89 (2014) no.4, 044910 [arXiv:1308.2440 [nucl-th]].
- [25] K. Tuchin, "Magnetic contribution to dilepton production in heavy-ion collisions", Phys. Rev. C 88 (2013), 024910 [arXiv:1305.0545 [nucl-th]].
- [26] S. Rath and B. K. Patra, "One-loop QCD thermodynamics in a strong homogeneous and static magnetic field", JHEP 12 (2017), 098 [arXiv:1707.02890 [hep-th]].
- [27] A. Bandyopadhyay, B. Karmakar, N. Haque and M. G. Mustafa, "Pressure of a weakly magnetized hot and dense deconfined QCD matter in one-loop hard-thermalloop perturbation theory", Phys. Rev. D 100 (2019) no.3, 034031 [arXiv:1702.02875 [hep-ph]].
- [28] S. Rath and B. Krishna Patra, "Thermomagnetic properties and Bjorken expansion of hot QCD matter in a strong magnetic field", Eur. Phys. J. A 55 (2019) no.11, 220 [arXiv:1806.03008 [hep-th]].
- [29] B. Karmakar, R. Ghosh, A. Bandyopadhyay, N. Haque and M. G. Mustafa, "Anisotropic pressure of deconfined QCD matter in presence of strong magnetic field within one-loop approximation", Phys. Rev. D 99 (2019) no.9, 094002 [arXiv:1902.02607 [hep-ph]].
- [30] S. Rath and B. K. Patra, "Effect of magnetic field on the charge and thermal transport properties of hot and dense QCD matter", Eur. Phys. J. C 80 (2020) no.8, 747 [arXiv:2005.00997 [hep-ph]].
- [31] S. A. Khan and B. K. Patra, "Cumulative effects of collision integral, strong magnetic field, and quasiparticle description on charge and heat transport in a thermal QCD medium", Phys. Rev. D 104 (2021) no.5, 054024 [arXiv:2011.02682 [hep-ph]].
- [32] V. Braguta, M. N. Chernodub, V. A. Goy, K. Landsteiner, A. V. Molochkov and M. I. Polikarpov, "Temperature dependence of the axial magnetic effect in two-color quenched QCD", Phys. Rev. D 89 (2014) no.7, 074510 [arXiv:1401.8095 [hep-lat]].
- [33] M. N. Chernodub, A. Cortijo, A. G. Grushin, K. Landsteiner and M. A. H. Vozmediano, "Condensed matter realization of the axial magnetic effect", Phys. Rev. B 89 (2014) no.8, 081407 [arXiv:1311.0878 [hep-th]].
- [34] M. Kurian and V. Chandra, "Electromagnetic response of hot QCD medium in the presence of background time-varying fields", Phys. Rev. D 104 (2021) no.9, 094037 [arXiv:2108.06791 [hep-ph]].
- [35] M. Kurian and V. Chandra, "Thermal and thermoelectric responses of hot QCD medium in time-varying magnetic fields", Phys. Rev. D 106 (2022) no.3, 034008 [arXiv:2205.14427 [hep-ph]].
- [36] V. P. Gusynin, V. A. Miransky and I. A. Shovkovy, "Catalysis of dynamical flavor symmetry breaking by a magnetic field in $(2+1)$ -dimensions", Phys. Rev. Lett. **73** (1994), 3499-3502 [erratum: Phys. Rev. Lett. 76 (1996), 1005 [arXiv:hep-ph/9405262 [hep-ph]].
- [37] D. S. Lee, C. N. Leung and Y. J. Ng, "Chiral symmetry breaking in a uniform external magnetic field", Phys. Rev. D 55 (1997), 6504-6513 [arXiv:hep-th/9701172 [hep-th]].
- [38] D. E. Kharzeev, "The Chiral Magnetic Effect and Anomaly-Induced Transport", Prog. Part. Nucl. Phys. 75 (2014), 133-151 [arXiv:1312.3348 [hep-ph]].
- [39] D. Satow, "Nonlinear electromagnetic response in quark-gluon plasma", Phys. Rev. D 90 (2014) no.3, 034018 [arXiv:1406.7032 [hep-ph]].
- [40] V. Roy, S. Pu, L. Rezzolla and D. Rischke, "Analytic Bjorken flow in onedimensional relativistic magnetohydrodynamics", Phys. Lett. B 750 (2015), 45-52 [arXiv:1506.06620 [nucl-th]].
- [41] S. I. Godunov, A. N. Rozanov, M. I. Vysotsky and E. V. Zhemchugov, "Extending the Higgs sector: an extra singlet", Eur. Phys. J. C 76 (2016), 1 [arXiv:1503.01618 [hep-ph]].
- [42] J. I. Kapusta and J. M. Torres-Rincon, "Thermal Conductivity and Chiral Critical Point in Heavy Ion Collisions", Phys. Rev. C 86 (2012), 054911 [arXiv:1209.0675 [nucl-th]].
- [43] G. S. Denicol, H. Niemi, I. Bouras, E. Molnar, Z. Xu, D. H. Rischke and C. Greiner, "Solving the heat-flow problem with transient relativistic fluid dynamics", Phys. Rev. D 89 (2014) no.7, 074005 [arXiv:1207.6811 [nucl-th]].
- [44] A. Muronga, "Relativistic Dynamics of Non-ideal Fluids: Viscous and heatconducting fluids. II. Transport properties and microscopic description of relativistic nuclear matter", Phys. Rev. C 76 (2007), 014910 [arXiv:nucl-th/0611091 [nucl-th]].
- [45] A. Puglisi, S. Plumari and V. Greco, "Electric Conductivity from the solution of the Relativistic Boltzmann Equation", Phys. Rev. D 90 (2014), 114009 [arXiv:1408.7043 [hep-ph]].
- [46] S. Mitra and V. Chandra, "Thermal relaxation, electrical conductivity, and charge diffusion in a hot QCD medium", Phys. Rev. D 94 (2016) no.3, 034025 [arXiv:1606.08556 [nucl-th]].
- [47] S. Mitra and V. Chandra, "Transport coefficients of a hot QCD medium and their relative significance in heavy-ion collisions", Phys. Rev. D 96 (2017) no.9, 094003 [arXiv:1702.05728 [nucl-th]].
- [48] S. Gupta, "The Electrical conductivity and soft photon emissivity of the QCD plasma", Phys. Lett. B 597 (2004), 57-62 [arXiv:hep-lat/0301006 [hep-lat]].
- [49] G. Aarts, C. Allton, A. Amato, P. Giudice, S. Hands and J. I. Skullerud, "Electrical conductivity and charge diffusion in thermal QCD from the lattice", JHEP 02 (2015), 186 [arXiv:1412.6411 [hep-lat]].
- [50] H. T. Ding, O. Kaczmarek and F. Meyer, "Thermal dilepton rates and electrical conductivity of the QGP from the lattice", Phys. Rev. D 94 (2016) no.3, 034504 [arXiv:1604.06712 [hep-lat]].
- [51] S. i. Nam, "Electrical conductivity of quark matter at finite T under external magnetic field", Phys. Rev. D 86 (2012), 033014 [arXiv:1207.3172 [hep-ph]].
- [52] M. Greif, I. Bouras, C. Greiner and Z. Xu, "Electric conductivity of the quark-gluon plasma investigated using a perturbative QCD based parton cascade", Phys. Rev. D 90 (2014) no.9, 094014 [arXiv:1408.7049 [nucl-th]].
- [53] B. Feng, "Electric conductivity and Hall conductivity of the QGP in a magnetic field", Phys. Rev. D 96 (2017) no.3, 036009.
- [54] P. V. Buividovich, M. N. Chernodub, D. E. Kharzeev, T. Kalaydzhyan, E. V. Luschevskaya and M. I. Polikarpov, "Magnetic-Field-Induced insulator-conductor transition in $SU(2)$ quenched lattice gauge theory", Phys. Rev. Lett. **105** (2010), 132001 [arXiv:1003.2180 [hep-lat]].
- [55] M. Kurian and V. Chandra, "Effective description of hot QCD medium in strong magnetic field and longitudinal conductivity", Phys. Rev. D 96 (2017) no.11, 114026 [arXiv:1709.08320 [nucl-th]].
- [56] K. Hattori and D. Satow, "Electrical Conductivity of Quark-Gluon Plasma in Strong Magnetic Fields", Phys. Rev. D 94 (2016) no.11, 114032 [arXiv:1610.06818 [hep-ph]].
- [57] M. Kurian, S. Mitra, S. Ghosh and V. Chandra, "Transport coefficients of hot magnetized QCD matter beyond the lowest Landau level approximation", Eur. Phys. J. C 79 (2019) no.2, 134 [arXiv:1805.07313 [nucl-th]].
- [58] S. Rath and B. K. Patra, "Revisit to electrical and thermal conductivities, Lorenz and Knudsen numbers in thermal QCD in a strong magnetic field", Phys. Rev. D 100 (2019) no.1, 016009 [arXiv:1901.03855 [hep-ph]].
- [59] S. Rath and S. Dash, "Effects of weak magnetic field and finite chemical potential on the transport of charge and heat in hot QCD matter", Eur. Phys. J. A 59 (2023) no.2, 25 [arXiv:2112.11802 [hep-ph]].
- [60] A. Shaikh, S. Rath, S. Dash and B. Panda, "Flow of charge and heat in thermal QCD within the weak magnetic field limit: A Bhatnagar-Gross-Krook model approach", Phys. Rev. D 108 (2023) no.5, 056021 [arXiv:2210.15388 [hep-ph]].
- [61] J. Anderson and H. Witting, "A relativistic relaxation-time model for the Boltzmann equation", *Physica*. 74, 466-488 (1974).
- [62] P. Bhatnagar, E. Gross and M. Krook, "A Model for Collision Processes in Gases 1. Small Amplitude Processes in Charged and Neutral One-Component Systems", *Phys. Rev.*. **94** pp. 511-525 (1954).
- [63] P. Welander, "On the temperature jump in a rarefied gas", *Ark. Fys.* 7 pp. 507-553 (1954).
- [64] G. S. Rocha, G. S. Denicol and J. Noronha, "Novel Relaxation Time Approximation to the Relativistic Boltzmann Equation", Phys. Rev. Lett. 127 (2021) no.4, 042301 [arXiv:2103.07489 [nucl-th]].
- [65] G. S. Rocha and G. S. Denicol, "Transient fluid dynamics with general matching conditions: A first study from the method of moments," Phys. Rev. D 104 (2021) no.9, 096016 [arXiv:2108.02187 [nucl-th]].
- [66] G. S. Rocha, M. N. Ferreira, G. S. Denicol and J. Noronha, "Transport coefficients of quasiparticle models within a new relaxation time approximation of the Boltzmann equation," Phys. Rev. D 106 (2022) no.3, 036022 [arXiv:2203.15571 [nucl-th]].
- [67] S. Rath and S. Dash, "Analyzing the transport coefficients and observables of a rotating QGP medium in kinetic theory framework with a novel approach to the collision integral", [arXiv:2403.01240 [hep-ph]].
- [68] E. Braaten and R. D. Pisarski, "Simple effective Lagrangian for hard thermal loops", Phys. Rev. D 45 (1992) no.6, R1827.
- [69] A. Peshier, B. Kampfer and G. Soff, "From QCD lattice calculations to the equation of state of quark matter", Phys. Rev. D 66 (2002), 094003 [arXiv:hep-ph/0206229 [hep-ph]].
- [70] K. Dusling, G. D. Moore and D. Teaney, "Radiative energy loss and v(2) spectra for viscous hydrodynamics", Phys. Rev. C 81 (2010), 034907 [arXiv:0909.0754 [nucl-th]].
- [71] K. Dusling and T. Schäfer, "Bulk viscosity, particle spectra and flow in heavy-ion collisions", Phys. Rev. C 85 (2012), 044909 [arXiv:1109.5181 [hep-ph]].
- [72] A. Kurkela and U. A. Wiedemann, "Analytic structure of nonhydrodynamic modes in kinetic theory", Eur. Phys. J. C 79 (2019) no.9, 776 [arXiv:1712.04376 [hep-ph]].
- [73] A. Hosoya and K. Kajantie, "Transport Coefficients of QCD Matter", Nucl. Phys. B 250 (1985), 666-688.
- [74] J. Kapusta and P. Landshoff, "Finite-temperature field theory", *Journal Of Physics G: Nuclear And Particle Physics*. 15, 267 (1989).
- [75] B. Schenke, M. Strickland, C. Greiner and M. H. Thoma, "A Model of the effect of collisions on QCD plasma instabilities," Phys. Rev. D 73 (2006), 125004 [arXiv:hepph/0603029 [hep-ph]].
- [76] C. Manuel and S. Mrowczynski, "Whitening of the quark gluon plasma," Phys. Rev. D 70 (2004), 094019 [arXiv:hep-ph/0403024 [hep-ph]].
- [77] P. Singha, S. Bhadury, A. Mukherjee and A. Jaiswal, "Relativistic BGK hydrodynamics," Eur. Phys. J. C 84 (2024) no.4, 417 [arXiv:2301.00544 [nucl-th]].

[78] M. Greif, F. Reining, I. Bouras, G. S. Denicol, Z. Xu and C. Greiner, "Heat conductivity in relativistic systems investigated using a partonic cascade", Phys. Rev. E 87 (2013), 033019 [arXiv:1301.1190 [hep-ph]].