

# Semilinear Klein-Gordon equation in space-time of black hole, which is gaining mass in the universe with accelerating expansion\*

Karen Yagdjian and Anahit Galstian

School of Mathematical and Statistical Sciences,  
University of Texas RGV, 1201 W. University Drive,  
Edinburg, TX 78539, USA  
e-mails: karen.yagdjian@utrgv.edu, anahit.galstyan@utrgv.edu

## Abstract

In this paper, we consider the propagation of waves in the space-time of a single black hole with a static Schwarzschild radius in the expanding universe, namely, the solutions of the linear and semilinear Klein-Gordon equations.

## 1 Main results

In this paper, we consider the propagation of waves in the space-time of a single black hole (BH) with a static Schwarzschild radius in the expanding universe, namely, the solutions of the linear and semilinear Klein-Gordon equations. The line element of this space-time, which we consider, is given by

$$ds^2 = - \left(1 - \frac{R_{Sch}}{r}\right) c^2 dt^2 + \left(1 - \frac{R_{Sch}}{r}\right)^{-1} a^2(t) dr^2 + a^2(t) r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (1.1)$$

where the Schwarzschild radius is

$$R_{Sch} := \frac{2GM_{bh}}{c^2}, \quad M_{bh} = \text{constant}.$$

Here,  $G$  is Newton's gravitational constant, and  $c$  is the vacuum speed of light. In this paper, we consider the case of space-time with the scale factor  $a(t) = \alpha_{acc} t^\ell$ ,  $\ell > 1$ . In what follows, we set  $\alpha_{acc} = 1$  and  $c = 1$ , if no obscurity arises.

Thus, we assume that the universe is expanding with acceleration and that the mass of the BHs is changing in time. (See, e.g., [5], [11].) In [5], it is suggested to assume that the gaining mass of BH is proportional to the scale factor raised to some power  $k$ . In this paper, we set  $k = 1$ , and consequently, the Schwarzschild radius is static.

In [15, 18], the BH with the static Schwarzschild radius embedded in the FLRW space-time with de Sitter scale factor  $a(t) = e^{Ht}$  was studied.

We consider another family of the Klein-Gordon equations for the self-interacting waves, with the mass term ingoing redshift proportional to the Hubble parameter  $\dot{a}(t)/a(t)$ , that is,  $m(t) = t^{-1}m_c$ , where  $m_c := mc^2/h$ . For a justification of this assumption, see, e.g., [9, 17]. Thus, the equation is,

$$\frac{\partial^2 \psi}{\partial t^2} + \frac{3\ell}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}(x, \partial_x) \psi + t^{-2} \frac{m^2 c^4}{h^2} \left(1 - \frac{R_{Sch}}{r}\right) \psi = c^2 \left(1 - \frac{R_{Sch}}{r}\right) \Psi(x, \psi), \quad (1.2)$$

---

\*AMS Subject Classifications: 35A01, 35L15, 35L71, 35Q75; 83C57

where  $\mathcal{A}(x, \partial_x)$ , written in spherical coordinates, is

$$\mathcal{A}(x, \partial_x) := \frac{c^2}{\alpha_{acc}^2} \left\{ \left(1 - \frac{R_{Sch}}{r}\right)^2 \frac{\partial^2}{\partial r^2} + \frac{2}{r} \left(1 - \frac{R_{Sch}}{2r}\right) \left(1 - \frac{R_{Sch}}{r}\right) \frac{\partial}{\partial r} + \left(1 - \frac{R_{Sch}}{r}\right) \frac{1}{r^2} \Delta_{\mathbb{S}^2} \right\}, \quad (1.3)$$

and  $\Delta_{\mathbb{S}^2}$  is the Laplace operator on the unit sphere  $\mathbb{S}^2$ . According to Lemma 2.1[18], the space-time of the BH with a static Schwarzschild radius is the only one that dissipates the waves independently of their location in space. This is related to the cosmological principle.

We can generalize (1.2) as follows

$$\frac{\partial^2 \psi}{\partial t^2} + \frac{3\ell}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}(x, \partial_x) \psi + t^{-2} \frac{m^2 c^4}{h^2} \psi + V(r, t) \psi = c^2 \left(1 - \frac{R_{Sch}}{r}\right) \Psi(x, \psi). \quad (1.4)$$

In (1.4)  $V(r, t)$  is the potential that, in particular, includes the case of the gravitational potential  $V(r, t) = -t^{-2} \frac{m^2 c^4}{h^2} \frac{R_{Sch}}{r}$ . The term  $\Psi(x, \psi)$  represents the self-interaction of the field. Consider the radial null geodesics of the space-time (1.1) with  $\ell > 1$ , when

$$A(t) := \frac{c}{\alpha_{acc}} \left( \frac{t^{1-\ell}}{1-\ell} - \frac{1}{1-\ell} \right) \leq A(\infty) = \frac{c}{\alpha_{acc}} \frac{1}{\ell-1} \quad \text{for all } t \in [1, \infty).$$

We consider waves (solutions of the equation) in the exterior  $B_{Sch}^{ext} := \{x \in \mathbb{R}^3 \mid |x| > R_{Sch}\}$  of the BH. The choice of the initial data with supports sufficiently distant from the horizon of the BH allows us to avoid the manifold, where the equation degenerates. Hence, we set

$$\text{supp } \psi_0, \text{supp } \psi_1 \subseteq \left\{ x \in \mathbb{R}^3 \mid |x| \geq R_{ID} > \frac{c}{\alpha_{acc}} \frac{1}{\ell-1} + R_{Sch} \right\} \subset B_{Sch}^{ext}. \quad (1.5)$$

Let  $H_{(s)} := H_{(s)}(\mathbb{R}^3)$  be the Sobolev space. The set of all smooth functions with uniformly bounded derivatives of any order will be denoted  $\mathcal{B}^\infty(B_{Sch}^{ext} \times [0, \infty))$ . For the projection operator on  $\mathbb{R}^3$  we write  $\pi_x$ . The space of solutions is

$$X(R, H_{(s)}, \gamma) := \left\{ \psi \in C([0, \infty); H_{(s)}) \mid \|\psi\|_X := \sup_{t \in [1, \infty)} t^\gamma \|\psi(x, t)\|_{H_{(s)}} \leq R \right\},$$

where  $\gamma \geq 0$ , with the metric

$$d(\psi_1, \psi_2) := \sup_{t \in [1, \infty)} t^\gamma \|\psi_1(x, t) - \psi_2(x, t)\|_{H_{(s)}}.$$

We apply the integral transform approach developed in [15]. (One can refer to [6, 7, 4] on the approach to the semilinear Klein-Gordon equation in the FLRW spacetimes based on the energy estimate.) Those integral transforms make it possible to bring the problem with unbounded time down to the problem with bounded time. This is possible since the space-time has a permanently restricted domain of influence.

Thus, the setting of the Cauchy Problem is as follows. Let the number  $R_{ID}$  be such that  $R_{ID} > R_{Sch}$ . For every given  $\psi_0 \in H_{(s)}$ ,  $\psi_1 \in H_{(s-1)}$  with  $\text{supp } \psi_0 \cup \text{supp } \psi_1 \subseteq \{x \in \mathbb{R}^3 \mid |x| \geq R_{ID}\} \subset B_{Sch}^{ext}$ , find a *global in time solution*  $\psi \in C^1([1, \infty); H_{(s)}) \cap C([1, \infty); H_{(s-1)})$  of the equation (1.4), such that  $\text{supp } \psi(t) \subseteq B_{Sch}^{ext}$  for all times  $t > 1$  and which takes the initial values  $\psi(x, 1) = \psi_0(x)$ ,  $\psi_t(x, 1) = \psi_1(x)$ , for all  $x \in \mathbb{R}^3$ .

We relate the properties of the potential  $V(x, t) \in C^2(B_{Sch}^{ext} \times [0, \infty))$  to the setting of the Cauchy problem, more precisely, with the support of the initial functions. We assume that the potential  $V(x, t) \in \mathcal{B}^\infty(B_{Sch}^{ext} \times [1, \infty))$  is such that for given  $s$  there are  $\varepsilon_P > 0$  and  $\delta \geq 0$  such that

$$\|V(x, t)\Phi(x)\|_{H_{(s)}} \leq t^{-\delta} \varepsilon_P \|\Phi\|_{H_{(s)}} \quad \text{for all } t \in [1, \infty) \quad \text{and all } \Phi \in H_{(s)}, \quad (1.6)$$

with compact  $\text{supp } \Phi \subseteq B_P \subset B_{Sch}^{ext}$ . The number  $\varepsilon_P$  may depend on the distance between the sphere  $\partial B_P$  and  $B_{Sch}$ .

The non-linear term is supposed to satisfy the following Lipschitz continuity condition ( $\mathcal{L}$ ). The smooth in  $x \in B_{Sch}^{ext}$  function  $\Psi = \Psi(x, \psi)$  is said to be Lipschitz continuous with exponent  $\alpha \geq 0$  in the space  $H_{(s)}$  if  $\text{supp } \Psi(x, \psi) \subseteq \text{supp } \psi$  and there is a constant  $C_s \geq 0$  such that

$$\|\Psi(x, \psi_1(x)) - \Psi(x, \psi_2(x))\|_{H_{(s)}} \leq C_s \|\psi_1 - \psi_2\|_{H_{(s)}} \left( \|\psi_1\|_{H_{(s)}}^\alpha + \|\psi_2\|_{H_{(s)}}^\alpha \right) \quad \text{for all } \psi_1, \psi_2 \in H_{(s)}.$$

The semilinear terms  $\Psi(\psi) = |\psi|^{1+\alpha}$  and  $\Psi(\psi) = \psi|\psi|^\alpha$  are of special interest.

**Theorem 1.1** *Consider the Cauchy problem for the equation (1.4) in  $\mathbb{R}^3 \times [1, \infty)$  with the initial conditions*

$$\psi(x, 1) = \psi_0(x) \in H_{(s)}, \quad \partial_t \psi(x, 1) = \psi_1(x) \in H_{(s)}, \quad (1.7)$$

with (1.5). Assume that the potential  $V(x, t) \in \mathcal{B}^\infty(B_{Sch}^{ext} \times [1, \infty))$  satisfies (1.6) with  $\delta = 2$ . Assume also that the physical mass of the field is large, that is,  $4m_c^2 \geq (3\ell - 1)^2$ , and the nonlinear term  $\Psi(x, \psi)$  satisfies condition ( $\mathcal{L}$ ) with  $\alpha > \frac{4}{3\ell-1}$  and  $\Psi(x, 0) = 0$ .

If  $\varepsilon_P$  and the norms  $\|\psi_0\|_{H_{(s)}}, \|\psi_1\|_{H_{(s)}}$  with  $s > 3/2$  are sufficiently small, then the Cauchy problem (1.4)  $\mathcal{E}$ (1.5)  $\mathcal{E}$ (1.7) has a global solution  $\psi \in C^2([1, \infty); H_{(s)})$ .

Moreover, if we denote  $M := \frac{\sqrt{4m_c^2 - (3\ell-1)^2}}{2(\ell-1)}$ , then the solution  $\psi$  belongs to the space  $X(R, H_{(s)}, \gamma)$ , with  $\gamma \leq \ell$  such that:

- (a) if  $\alpha > \frac{\ell+3}{2\ell}$ , then  $\gamma > \frac{3(\ell+1)}{2(\alpha+1)}$  for  $M > 0$ , while  $\ell > \gamma > \frac{3(\ell+1)}{2(\alpha+1)}$  for  $M = 0$ ;
- (b) if  $\alpha > \frac{4}{3\ell-1}$ , then  $\frac{2}{\alpha} \leq \gamma \leq \frac{3(\ell+1)}{2(\alpha+1)}$  for  $M > 0$ , while  $\frac{2}{\alpha} \leq \gamma \leq \frac{3(\ell+1)}{2(\alpha+1)}$  and  $\gamma < \ell$  for  $M = 0$ .

Thus, the solution  $\psi$  decays according to  $\|\psi(x, t)\|_{H_{(s)}} \leq Rt^{-\gamma}$ ,  $t \in [1, \infty)$ .

Note,  $\lim_{\ell \rightarrow \infty} \frac{\ell+3}{2\ell} = \frac{1}{2}$  and  $2 > \frac{\ell+3}{2\ell} > \frac{4}{3\ell-1}$  for all  $\ell > 1$ , while  $\frac{\ell+3}{2\ell} = \frac{4}{3\ell-1}$  if  $\ell = 1$ .

**Theorem 1.2** *Consider the Cauchy problem for the equation (1.4) in  $\mathbb{R}^3 \times [1, \infty)$  with the initial conditions (1.7) satisfying (1.5). Assume that the potential  $V(x, t) \in \mathcal{B}^\infty(B_{Sch}^{ext} \times [1, \infty))$  satisfies (1.6). Assume also that the physical mass  $m$  of the field is small, that is,  $4m_c^2 < (3\ell - 1)^2$ , and the nonlinear term  $\Psi(x, \psi)$  satisfies condition ( $\mathcal{L}$ ) with  $\alpha > 0$  and  $\Psi(x, 0) = 0$ .*

If  $\varepsilon_P$  and the norms  $\|\psi_0\|_{H_{(s)}}, \|\psi_1\|_{H_{(s)}}$  with  $s > 3/2$  are sufficiently small, then the Cauchy problem (1.4)  $\mathcal{E}$ (1.5)  $\mathcal{E}$ (1.7) has a global solution  $\psi \in C^2([1, \infty); H_{(s)})$ .

Moreover, if  $\delta = 0$ , then the solution  $\psi$  belongs to the space  $X(R, H_{(s)}, \gamma)$ , with  $\gamma$  such that  $\gamma \leq \ell$  and

$$\gamma < \frac{1}{2} \left( 3\ell - 1 - \sqrt{(3\ell - 1)^2 - 4m_c^2} \right),$$

that is, the solution  $\psi$  decays according to  $\|\psi(x, t)\|_{H_{(s)}} \leq Rt^{-\gamma}$ ,  $t \in [1, \infty)$ .

If  $\delta > 0$ , then one can choose  $\gamma \leq \min \left\{ \ell, \frac{1}{2} \left( 3\ell - 1 - \sqrt{(3\ell - 1)^2 - 4m_c^2} \right) \right\}$ .

Note, if  $\delta > 0$  and  $(3\ell - 1)^2/4 > m_c^2 > 1$  in the last theorem, then  $\gamma$  attains its largest value  $\gamma = \ell$ .

The Sobolev embedding theorem implies the decay of  $L_\infty$ -norm of solution. In particular, we recover some results of [9] if the charge of the model is zero. Further, the results of Theorems 1.1, 1.2, if  $M_{bh} = 0$ , agree with those previously known, see [6, 8, 16, 1] and references therein.

For the semilinear Klein-Gordon equation with integrable speed of propagation and time-dependent coefficients, that is, for the equation with a permanently restricted domain of influence, in flat FLRW models, to prove a global existence Aslan, Ebert, and Reissig [1] applied a partial Fourier transform and a splitting of the phase space into three zones.

Nataro and Sasane [9] obtained exact decay rates for solutions to the Klein-Gordon equation in the de Sitter universe in flat FLRW form. For the derivative of the solutions to the massless equation in the cosmological region of the Reissner–Nordström–de Sitter model, under some assumption on the boundedness of the smooth solution in the norms over the two components of the future cosmological horizon, parametrized by the flow of the global Killing vector field  $\partial/\partial t$ , they proved  $L_\infty$ -norm decay at infinity. In the last model, the mass and charge of the black hole are assumed to be static.

**Remark 1.3** *Theorems 1.1,1.2 can be extended to the equation*

$$\frac{\partial^2 \psi}{\partial t^2} + \frac{\mu}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}(x, \partial_x) \psi + t^{-2} \frac{m^2 c^4}{h^2} \psi + V(r, t) \psi = c^2 \left( 1 - \frac{R_{Sch}}{r} \right) \Psi(x, \psi),$$

with the parameter  $\mu \in \mathbb{R}$ , which is not necessarily relevant to the dimension of the spatial variable (see, e.g., [3]); they can be extended to  $n$ -dimensional black holes. Moreover, to derive a better decay at infinite time, one can appeal to the estimate in homogeneous Sobolev spaces and to the Gagliardo-Nirenberg inequality.

The rest of this paper is organized as follows. In Section 2, the Liouville transformation and representation of the solution to the linear equation are given. Moreover, the equation is transformed into the self-adjoint Cauchy-Kowalewski form. The case of large mass is considered in Section 3, while the case of small mass is discussed in Section 4. The proof of existence of the solution of semilinear equation is based on Banach's fixed point theorem. The decay estimates for the solution of the linear equation are derived in subsections 3.2-3.5 and subsections 4.2-4.5, respectively. A significant part of the text (subsection 3.1,3.2,4.1,4.2) is devoted to the estimation of the integrals of the kernel functions of the integral transform containing Gauss's hypergeometric function.

## 2 Preliminaries. Linear equation

We will use the notations  $\vec{x} = (x_1, x_2, x_3) := (x, y, z)$ ,  $x_0 = t$ , and  $\vec{\xi} = (\xi_1, \xi_2, \xi_3)$ . The inner product in  $\mathbb{R}^3$  will be denoted  $\vec{x} \cdot \vec{\xi}$ . In the case of  $\ell = 0$ , the linear Klein-Gordon equation without source in Cartesian coordinates can be written (see, e.g., [18]) in the form (1.2), where

$$\mathcal{A}(\vec{x}; \vec{\xi}) = A_2(\vec{x}; \vec{\xi}) + A_1(\vec{x}; \vec{\xi}), \quad (2.1)$$

while  $A_2(\vec{x}; \vec{\xi})$  and  $A_1(\vec{x}; \vec{\xi})$  are the principal symbol and the low order symbol, respectively, and

$$F(|\vec{x}|) = F(r) := 1 - \frac{R_{Sch}}{|\vec{x}|}, \quad r := |\vec{x}| := \sqrt{x^2 + y^2 + z^2}, \quad r > R_{Sch}.$$

Thus, the symbol  $\mathcal{A}(\vec{x}; \vec{\xi})$  of the operator  $\mathcal{A}(\vec{x}, \partial_x)$  (1.3) is given by

$$A_2(\vec{x}; \vec{\xi}) = -c^2 F(r) \left( |\vec{\xi}|^2 - \frac{R_{Sch} (\vec{x} \cdot \vec{\xi})^2}{|\vec{x}|^3} \right), \quad A_1(\vec{x}; \vec{\xi}) = -c^2 F(r) \frac{i R_{Sch} (\vec{x} \cdot \vec{\xi})}{|\vec{x}|^3}.$$

On every compact set in  $B_{Sch}^{ext} \times \mathbb{R}$ , the equation (1.4) is strictly hyperbolic. It has multiple characteristics on the event horizon  $r = R_{Sch}$ . Consequently, the well-posedness of the Cauchy problem requires some kind of Levi condition. (See, for details, [13].)

**The finite propagation speed property.** Consider the equation

$$\frac{\partial^2 \psi}{\partial t^2} + \frac{3\ell}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}(x, \partial_x) \psi + t^{-2} F(r) m_c^2 \psi = 0, \quad (2.2)$$

in  $B_{Sch}^{ext}$ . We apply the Liouville transform with  $k_{\pm}$  defined in subsection 2.1  $\psi(x, t) = t^{k_{\pm}} \sqrt{F(r)} u(x, t)$  to (2.2), then the covariant Klein-Gordon equation (2.2) reads

$$\frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} - t^{-2\ell} \mathcal{A}_{3/2}(x, \partial_x) u + (t^{-1}(\ell + 2i\mathcal{M})) \frac{\partial}{\partial t} u - \frac{R_{Sch}}{r} t^{-2} \frac{m_c^2}{c^2} u = 0, \quad (2.3)$$

where  $\mathcal{M}$  is written in (2.10) and the operator  $\mathcal{A}_{3/2}$  in the spherical coordinates is defined (see, e.g., [18, Sec.2.3]) by

$$\mathcal{A}_{3/2}(x, \partial_x) v := F(r)^{3/2} \frac{\partial^2}{\partial r^2} \sqrt{F(r)} v + \sqrt{F(r)} \frac{2}{r} \left( 1 - \frac{R_{Sch}}{2r} \right) \frac{\partial}{\partial r} \sqrt{F(r)} v + F(r) \frac{1}{r^2} \Delta_{S^2} v,$$

while the term  $-t^{-2} \frac{R_{Sch}}{r} \frac{m^2}{c^2}$  can be regarded as a potential  $V = V(\vec{x}, t)$ . According to [18], the operator  $\mathcal{A}_{3/2}(x, \partial_x)$  can be written as follows

$$\mathcal{A}_{3/2}(\vec{x}; D_x)v = \sum_{i,j=1}^3 \frac{\partial}{\partial x_i} \left( a_{ij}(\vec{x}) \frac{\partial}{\partial x_j} v \right) + \frac{G^2 M_{bh}^2}{c^4 |\vec{x}|^4} v,$$

with the coefficients  $a_{ij}(\vec{x})$  such that

$$\begin{aligned} a_{ii}(\vec{x}) &:= -F(r) \left( \frac{R_{Sch} x_i^2}{|\vec{x}|^3} - 2 \right), \quad i = 1, 2, 3, \\ a_{ji}(\vec{x}) &= a_{ij}(\vec{x}) := -F(r) \frac{R_{Sch} x_i x_j}{|\vec{x}|^3}, \quad i, j = 1, 2, 3, \quad i \neq j. \end{aligned}$$

The finite propagation speed property for the linear equation (2.3) can be proved by standard arguments (see, e.g., [12]).

**The radial geodesics.** Next, consider the radial geodesic solving

$$\frac{dr}{dt} = -\frac{c}{\alpha_{acc}} t^{-\ell} \left( 1 - \frac{R_{Sch}}{r} \right) \quad (2.4)$$

and starting at  $R_{ID}$ . The existence of global in time geodesic is given by the following statement.

**Lemma 2.1** *The implicit function  $r = r(t)$ , which is defined for all  $t \in [1, \infty)$  by*

$$R_{ID} - r - R_{Sch} \ln \left( \frac{r - R_{Sch}}{R_{ID} - R_{Sch}} \right) = \frac{c}{\alpha_{acc}} \left( \frac{t^{1-\ell}}{1-\ell} - \frac{1}{1-\ell} \right), \quad (2.5)$$

with some positive number  $\varepsilon$  satisfies the inequality

$$r(t) \geq R_{Sch} + \varepsilon \quad \text{for all } t > 1. \quad (2.6)$$

**Proof.** The solution to (2.4) is given by (2.5). For  $\tau \in [0, \infty)$ , consider an implicit function  $z(\tau)$  given by the problem

$$z - R_{Sch} \ln \left( 1 - \frac{z}{R_{ID} - R_{Sch}} \right) = \tau, \quad z(1) = 0.$$

The function  $z = z(\tau)$  is well defined if  $z(\tau) \in [0, R_{ID} - R_{Sch})$ . Moreover, it is positive, and is continuous. Indeed,

$$\frac{dz}{d\tau} \left( 1 + R_{Sch} \frac{1}{R_{ID} - R_{Sch} - z} \right) = 1$$

as long as  $z(\tau) < R_{ID} - R_{Sch}$  implies  $\frac{dz}{d\tau} > 0$ . Denote  $z_\infty = z(A(\infty))$ , where  $A(\infty) = \frac{c}{\alpha(\ell-1)}$ , then  $z(\tau) \leq z_\infty$  for all  $\tau \in [0, cA(\infty)]$ . The existence of the number  $z_\infty \in [0, R_{ID} - R_{Sch})$  follows from

$$\lim_{z \nearrow R_{ID} - R_{Sch}} \left( z - R_{Sch} \ln \left( 1 - \frac{z}{R_{ID} - R_{Sch}} \right) \right) = \infty.$$

Hence, there is a number  $\varepsilon > 0$  such that  $z(\tau) \leq R_{ID} - R_{Sch} - \varepsilon$  for all  $\tau \in [0, cA(\infty)]$ . Next, we define  $\tau$  by

$$\tau(t) := A(t) \quad \text{and} \quad \frac{d\tau}{dt} = \frac{c}{a(t)} > 0 \quad \text{for all } t \in [1, \infty).$$

Thus, the function  $r = r(t)$ , which is defined on  $[0, \infty)$  and  $r(t) = R_{ID} - z(\tau(t))$ , where  $\tau(t) < cA(\infty)$  for all  $t \in [1, \infty)$ . The lemma is proved.  $\square$

Hence, if the initial functions have compact supports in  $K \subset B_{Sch}^{ext} \subset \mathbb{R}^3$ ,  $\text{dist}(\partial K, B_{Sch}) > \frac{c}{\alpha_{acc}(\ell-1)}$ , then there is  $\varepsilon > 0$  such that for all  $t \in [1, \infty)$ , the solution of (1.4) with  $\Psi = 0$  has a compact support in  $\{x \in \mathbb{R}^3 \mid |x| > R_{Sch} + \varepsilon\} \subset B_{Sch}^{ext}$ . This follows from the dependence domain.

Thus, for every given fixed compact  $K \subset B_{Sch}^{ext}$ , the coefficients of the equation can be continued smoothly outside of the small  $\delta$ -neighborhood of  $K$  to be constants. This continuation does change solutions, which takes initial data supported by  $K$ , at least for time duration  $d/c$ . Consequently, we can replace the operator  $\mathcal{A}(x, \partial_x)$  with such continuation (auxiliary operator)  $\mathcal{A}_\varepsilon(\vec{x}; \vec{\xi})$  that has a symbol

$$\mathcal{A}_\varepsilon(\vec{x}; \vec{\xi}) := c^2 \left( 1 - \chi_\varepsilon(\vec{x}) \frac{R_{Sch}}{|\vec{x}|} \right) \left( -|\xi|^2 + \chi_\varepsilon(\vec{x}) \frac{R_{Sch} (\vec{x} \cdot \vec{\xi})^2}{|\vec{x}|^3} - \chi_\varepsilon(\vec{x}) \frac{i R_{Sch} (\vec{x} \cdot \vec{\xi})}{|\vec{x}|^3} \right), \quad (2.7)$$

where  $\chi_\varepsilon$  is a cutoff function such that  $\chi_\varepsilon(\vec{x}) = 0$  if  $|\vec{x}| < R_{Sch} + \varepsilon/2$  while  $\chi_\varepsilon(\vec{x}) = 1$  if  $|\vec{x}| > R_{Sch} + \varepsilon$ . This allows us to apply the  $H_{(s)}$ -estimates.

## 2.1 The partial Liouville transform

To get rid of the mass term in the initial value problem for the equation

$$\frac{\partial^2}{\partial t^2} \psi + \frac{3\ell}{t} \frac{\partial}{\partial t} \psi - t^{-2\ell} \mathcal{A}(x, \partial_x) \psi + t^{-2} m_c^2 \psi = f$$

we use the partial Liouville transform  $\psi(x, t) = \omega(t)u(x, t)$ . Then the equation reads

$$u_{tt}(x, t) - t^{-2\ell} \mathcal{A}(x, \partial_x) u(x, t) + \left( 2 \frac{\omega_t(t)}{\omega(t)} + \frac{3\ell}{t} \right) u_t(x, t) + \left( \frac{\omega_{tt}(t)}{\omega(t)} + \frac{3\ell}{t} \frac{\omega_t(t)}{\omega(t)} + \frac{m_c^2}{t^2} \right) u(x, t) = \frac{1}{\omega(t)} f.$$

We set

$$\frac{\omega_{tt}(t)}{\omega(t)} + \frac{3\ell}{t} \frac{\omega_t(t)}{\omega(t)} + \frac{m_c^2}{t^2} = 0.$$

If we look for  $\omega(t) = t^k$ , then  $k(k-1) + 3\ell k + m_c^2 = 0$ . The solutions to the last equation are

$$k_\pm = \frac{1}{2} \left( 1 - 3\ell \pm \sqrt{(1 - 3\ell)^2 - 4m_c^2} \right). \quad (2.8)$$

We consider the equation with  $\ell > 1$  that implies that we have to consider three cases: (1)  $m_c < \frac{3\ell-1}{2}$ , (2)  $m_c = \frac{3\ell-1}{2} \implies k_\pm = \frac{1}{2}(1 - 3\ell)$ , and (3)  $m_c > \frac{3\ell-1}{2}$ . We say that *mass is small* if  $m_c < \frac{3\ell-1}{2}$ , while in other cases we say mass is *large*.

Thus, we arrived at the generalized Euler-Poisson-Darboux equation

$$u_{tt}(x, t) - t^{-2\ell} \mathcal{A}(x, \partial_x) u(x, t) + \frac{1 \pm \sqrt{(1 - 3\ell)^2 - 4m_c^2}}{t} u_t(x, t) = t^{-k_\pm} f. \quad (2.9)$$

Denote

$$\mathcal{M}_\pm := \frac{1}{2i} (1 - \ell \pm \sqrt{(1 - 3\ell)^2 - 4m_c^2}), \quad \frac{1 \pm \sqrt{(1 - 3\ell)^2 - 4m_c^2}}{t} := t^{-1} (\ell + 2i\mathcal{M}_\pm). \quad (2.10)$$

Then, for the case of small mass, that is,  $4m_c^2 < (1 - 3\ell)^2$ , we have  $2i\frac{\mathcal{M}_-}{1-\ell} > 1$ ,  $2i\frac{\mathcal{M}_+}{1-\ell} < 1$ , while for the large mass, that is,  $4m_c^2 \geq (1 - 3\ell)^2$ , we have  $\Re 2i\frac{\mathcal{M}_-}{1-\ell} = \Re 2i\frac{\mathcal{M}_+}{1-\ell} = 1$ .

## 2.2 Representation of the solution of linear equation

We set the initial time  $\varepsilon = 1$  in the formulas of [17] (for some particular cases, see also [8, 10]), then

$$\phi(1) := \frac{1}{1-\ell} < 0, \quad \phi(t) := \phi(1)t^{1-\ell} \quad \text{for all } t \geq 1, \quad \phi(t) - \phi(b) > 0, \quad \text{for all } 1 \leq b < t.$$

Next, we need the following kernel functions  $E(r, t; t_0; \mathcal{M})$ ,  $K_1(r, t; \mathcal{M}; 1)$ , and  $K_0(r, t; \mathcal{M}; 1)$ , from [17]

$$E(r, t; t_0; \mathcal{M}) = 2^{\frac{2i\mathcal{M}}{1-\ell}} (1-\ell)^{\frac{\ell}{1-\ell}} \phi(t_0)^{\frac{\ell+2i\mathcal{M}}{1-\ell}} \left( (\phi(t) + \phi(t_0))^2 - r^2 \right)^{-\frac{i\mathcal{M}}{1-\ell}} \quad (2.11)$$

$$\times F \left( i \frac{\mathcal{M}}{1-\ell}, i \frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(t_0))^2 - r^2}{(\phi(t) + \phi(t_0))^2 - r^2} \right),$$

$$K_1(r, t; \mathcal{M}; 1) := 2^{2i\frac{\mathcal{M}}{1-\ell}} \phi(1)^{2i\frac{\mathcal{M}}{1-\ell}} \left( (\phi(t) + \phi(1))^2 - r^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \quad (2.12)$$

$$\times F \left( i \frac{\mathcal{M}}{1-\ell}, i \frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(1))^2 - r^2}{(\phi(t) + \phi(1))^2 - r^2} \right),$$

$$K_0(r, t; \mathcal{M}; 1) := -i 2^{1+2i\frac{\mathcal{M}}{1-\ell}} \frac{\mathcal{M}}{1-\ell} \phi^{\frac{2i\mathcal{M}}{1-\ell}}(1) \left( (\phi(t) + \phi(1))^2 - r^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \quad (2.13)$$

$$\times \left\{ \frac{r^2 - \phi(t)(\phi(t) - \phi(1))}{r^2 - (\phi(t) - \phi(1))^2} F \left( i \frac{\mathcal{M}}{1-\ell}, i \frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(1))^2 - r^2}{(\phi(t) + \phi(1))^2 - r^2} \right) \right.$$

$$\left. - \frac{2\phi(t)\phi(1) (\phi^2(t) - \phi^2(1) - r^2)}{\left( (\phi(t) - \phi(1))^2 - r^2 \right) \left( (\phi(t) + \phi(1))^2 - r^2 \right)} \right.$$

$$\left. \times F \left( i \frac{\mathcal{M}}{1-\ell} + 1, i \frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(1))^2 - r^2}{(\phi(t) + \phi(1))^2 - r^2} \right) \right\},$$

respectively.<sup>1</sup> Here  $F(a, b; c; \zeta)$  is the Gauss's hypergeometric function (see, e.g., [2]). We also define

$$K_2(r, t; \mathcal{M}; 1) := K_0(r, t; \mathcal{M}; 1) + 2i \frac{\mathcal{M}}{1-\ell} K_1(r, t; \mathcal{M}; 1). \quad (2.14)$$

We rewrite equation (2.9) as follows:

$$u_{tt} - t^{-2\ell} A(x, \partial_x)u + t^{-1}(\ell + 2i\mathcal{M}_{\pm})u_t = t^{-k_{\pm}} f(x, t), \quad t > 1, \quad x \in \Omega,$$

where (2.10) has been used.

**Theorem 2.2** [17, Theorem 4.1] *Assume that the function  $v = v_g(x, r; b) \in C_{x,r,b}^{d,2,0}$  solves the problem*

$$\begin{cases} v_{rr}(x, r; b) - A(x, \partial_x)v(x, r; b) = 0, & x \in \Omega, \\ v(x, 0; b) = g(x, b), \quad v_r(x, 0; b) = 0, & x \in \Omega, \end{cases}$$

while the function  $v = v_{\varphi}(x, r) \in C_{x,r}^{d,2}$  solves the problem

$$\begin{cases} v_{rr}(x, r) - A(x, \partial_x)v(x, r) = 0, & x \in \Omega, \\ v(x, 0) = \varphi(x), \quad v_r(x, 0) = 0, & x \in \Omega. \end{cases}$$

Then the function  $u = u(x, t)$  defined by

$$u(x, t) = \int_1^t db \int_0^{\phi(t)-\phi(b)} E(r, t; b; \mathcal{M}) v_g(x, r; b) dr$$

$$+ \int_0^{\phi(t)-\phi(1)} K_1(r, t; \mathcal{M}; 1) v_{u_1}(x, r) dr$$

$$+ \left( \frac{\phi(t)}{\phi(1)} \right)^{-i\frac{\mathcal{M}}{1-\ell}} v_{u_0}(x, \phi(t) - \phi(1)) + \frac{1}{\phi(1)} \int_0^{\phi(t)-\phi(1)} K_2(r, t; \mathcal{M}; 1) v_{u_0}(x, r) dr,$$

where  $\phi(t) := \frac{1}{1-\ell} t^{1-\ell}$  and the kernels  $E(r, t; b; \mathcal{M})$ ,  $K_1(r, t; \mathcal{M}; 1)$ ,  $K_0(r, t; \mathcal{M}; 1)$ , and  $K_2(r, t; \mathcal{M}; 1)$  are defined by (2.11), (2.12), (2.13), and (2.14), respectively, is a solution to the Cauchy problem

$$\begin{cases} u_{tt} - t^{-2\ell} A(x, \partial_x)u + t^{-1}(\ell + 2i\mathcal{M})u_t = g(x, t), & t > 1, \quad x \in \Omega, \\ u(x, 1) = u_0(x), \quad u_t(x, 1) = u_1(x), & x \in \Omega. \end{cases}$$

<sup>1</sup>The definition of  $E$  and  $K_1$  in [17] differs from the one used in this paper by factors.

We study the Cauchy problem (1.4)&(1.7) through the integral equation. To define that integral equation we appeal to the operator

$$G := \mathcal{K} \circ \mathcal{E}\mathcal{E}, \quad (2.15)$$

where  $\mathcal{E}\mathcal{E}$  stands for the evolution (wave) equation in the exterior of BH in the universe without expansion as follows. For the function  $f(x, t)$  we define

$$v(x, t; b) := \mathcal{E}\mathcal{E}[f](x, t; b),$$

where the function  $v(x, t; b)$  is a solution to the Cauchy problem

$$\begin{cases} \partial_t^2 v - \mathcal{A}(x, \partial_x)v = 0, & x \in B_{Sch}^{ext} \subset \mathbb{R}^3, \quad t \geq 0, \\ v(x, 0; b) = f(x, b), \quad v_t(x, 0; b) = 0, & x \in B_{Sch}^{ext} \subset \mathbb{R}^3, \quad b \geq 1, \end{cases} \quad (2.16)$$

while  $\mathcal{K}$  is introduced by

$$\mathcal{K}[v](x, t) := 2t^k \int_1^t b^{-k} db \int_0^{\phi(t)-\phi(b)} E(r, t; b; \mathcal{M})v(x, r; b) dr. \quad (2.17)$$

The kernel  $E(r, t; 0, b; M)$  is given by (2.11).

### 2.3 Energy estimates and energy conservation

Let  $\partial_t^2 - A(x, \partial_x) = \partial_t^2 - \sum_{|\alpha| \leq 2} a_\alpha(x) \partial_x^\alpha$  be a second-order strictly hyperbolic operator with coefficients  $a_\alpha \in \mathcal{B}^\infty$ , where  $\mathcal{B}^\infty$  is the space of all  $C^\infty(\mathbb{R}^3)$  functions with uniformly bounded derivatives of all orders. Let  $v = v(x, t)$  be the solution of the problem

$$\begin{cases} \partial_t^2 v - A(x, \partial_x)v = 0, & x \in \mathbb{R}^n, \quad t \geq 0, \\ v(x, 0) = v_0(x), \quad v_t(x, 0) = v_1(x), & x \in \mathbb{R}^n. \end{cases}$$

The following energy estimate is well known. (See, e.g., [12].) For every  $s \in \mathbb{R}$  and given  $T > 0$  there is  $C_s(T)$  such that

$$\|v_t(t)\|_{H_{(s)}} + \|v(t)\|_{H_{(s+1)}} \leq C_s(T)(\|v_1\|_{H_{(s)}} + \|v_0\|_{H_{(s+1)}}), \quad 0 \leq t \leq T. \quad (2.18)$$

We note that although in the last estimate the time interval is bounded, however, due to the integral transform approach, it is possible to reduce the problem with infinite time to the problem with finite time and to apply (2.18). Here, the permanently restricted domain of influence of the space-time with  $\ell > 1$  comes into play.

We are going to apply the estimate (2.18) to the problem

$$\begin{cases} \partial_t^2 v - \mathcal{A}_\varepsilon(x, \partial_x)v = 0, & x \in \mathbb{R}^3, \quad t \geq 0, \\ v(x, 0) = v_0(x), \quad v_t(x, 0) = v_1(x), & x \in \mathbb{R}^3, \end{cases}$$

where the operator  $\mathcal{A}_\varepsilon(x, \partial_x)$  has a symbol  $\mathcal{A}_\varepsilon(x; \xi)$  of (2.7). In that case the constant  $C_s(T)$  depends on  $\varepsilon$  as well. The conservation of the energy of the solution of the equation

$$\frac{\partial^2 \psi}{\partial t^2} - \mathcal{A}(x, \partial_x)\psi + F(r) \frac{m^2 c^4}{h^2} \psi = 0$$

is known (see, e.g., [18]).

### 2.4 Equation in self-adjoint form

The semilinear Klein-Gordon equation without potential is

$$\frac{\partial^2 \psi}{\partial t^2} + \frac{3\ell}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}(x, \partial_x)\psi + t^{-2} F(r) m_c^2 \psi = c^2 F(r) \Psi(\psi), \quad (2.19)$$



where  $\mathcal{A}(x, \partial_x)$  is defined in (1.3). The operator  $\mathcal{A}(x, \partial_x) = \mathcal{A}(\vec{x}, \partial_x) = \mathcal{A}(x, y, z; D_x, D_y, D_z)$  has the symbol (2.1) in Cartesian coordinates. We apply the Liouville substitution  $\psi(x, t) = t^{k_\pm} \sqrt{F(r)} u(x, t)$  to (2.19), then the covariant Klein-Gordon equation (2.19) became

$$\frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} - t^{-2\ell} \mathcal{A}_{3/2}(x, \partial_x) u + (t^{-1}(\ell + 2i\mathcal{M}_\pm)) \frac{\partial}{\partial t} u - \frac{R_{Sch}}{r} t^{-2} \frac{m_c^2}{c^2} u = t^{-k_+} \sqrt{F(r)} \Psi \left( t^{k_+} \sqrt{F(r)} u \right),$$

where the operator  $\mathcal{A}_{3/2}(x, \partial_x)$  in the spherical coordinates is defined (see, [18, Sec.2.3]) by

$$\mathcal{A}_{3/2}(x, \partial_x) v := F(r)^{3/2} \frac{\partial^2}{\partial r^2} \sqrt{F(r)} v + \sqrt{F(r)} \frac{2}{r} \left( 1 - \frac{R_{Sch}}{2r} \right) \frac{\partial}{\partial r} \sqrt{F(r)} v + F(r) \frac{1}{r^2} \Delta_{S^2} v,$$

while the term  $-t^{-2} \frac{R_{Sch}}{r} \frac{m_c^2 c^2}{h^2}$  can be regarded as a potential  $V = V(x, y, z, t)$ .

The symbol  $\mathcal{A}_{3/2}(x, \xi)$  of operator  $\mathcal{A}_{3/2}(x, \partial_x)$  in Cartesian coordinates is

$$\mathcal{A}_{3/2}(\vec{x}, \vec{\xi}) := \left( 1 - \frac{R_{Sch}}{|\vec{x}|} \right) \left( -|\vec{\xi}|^2 + \frac{R_{Sch} (\vec{x} \cdot \vec{\xi})^2}{|\vec{x}|^3} \right) + \frac{R_{Sch}^2}{4|\vec{x}|^4}.$$

According to Lemma 2.4 [18], the operator  $\mathcal{A}_{3/2}(x, \partial_x)$  is essentially self-adjoint on  $C_0^\infty(B_{Sch}^{ext})$ , and on every closed subset in  $B_{Sch}^{ext}$  it is an elliptic operator that is non-positive on the subspace of functions with supports in  $B_{Sch}^{ext}$ .

### 3 The case of large mass. Proof of Theorem 1.1

In this case, for the partial Liouville transform we choose  $k_-$  and  $\mathcal{M}_-$  from (2.8) and (2.10), respectively. We also denote

$$M := \frac{\sqrt{4m_c^2 - (1 - 3\ell)^2}}{2(\ell - 1)} \geq 0, \quad \text{then} \quad i \frac{\mathcal{M}_-}{1 - \ell} = \frac{1}{2} + iM.$$

In this section, we drop the subindex of  $\mathcal{M}_-$ .

**Lemma 3.1** *For  $M \in [0, \infty)$  and the given above parameter  $\ell$  with some constant  $C$*

$$\begin{aligned} \left| F \left( \frac{1}{2}, \frac{1}{2} + iM, \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2} \right) \right| &\leq C \quad \text{for all } z \in [1, \infty), \\ \left| F \left( \frac{1}{2}, \frac{3}{2} + iM; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2} \right) \right| &\leq C + C(1 - \text{sign}(M)) z^{\frac{1}{2}} \quad \text{for all } z \in [1, \infty). \end{aligned}$$

**Proof.** According to [2, (1) Sec. 2.10]

$$\begin{aligned} F \left( \frac{1}{2}, \frac{1}{2} + iM, \frac{3}{2}; \zeta \right) &= \frac{\Gamma(\frac{3}{2})\Gamma(\frac{1}{2} - iM)}{\Gamma(1)\Gamma(1 - iM)} F \left( \frac{1}{2}, \frac{1}{2} + iM, \frac{1}{2} + iM; 1 - \zeta \right) \\ &\quad + (1 - \zeta)^{\frac{1}{2} - iM} \frac{\Gamma(\frac{3}{2})\Gamma(\frac{1}{2} - iM)}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2} + iM)} F \left( 1, 1 - iM, \frac{3}{2} - iM; 1 - \zeta \right), \end{aligned}$$

where  $\zeta = \frac{(z-1)^2}{(z+1)^2}$  and  $1 - \zeta = \frac{4z}{(z+1)^2} \rightarrow 0$  as  $z \rightarrow \infty$ . Consequently, the first estimate follows. Similarly, with  $M > 0$  we obtain

$$\left| F \left( \frac{1}{2}, \frac{3}{2} + iM; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2} \right) \right| \leq C \quad \text{for all } z \in [1, \infty),$$

while according to [2, (23) Sec. 2.1] with  $M = 0$  we obtain

$$\left| F \left( \frac{1}{2}, \frac{3}{2} + iM; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2} \right) \right| \leq C + C(1 - \text{sign}(M)) \left( \frac{4z}{(z+1)^2} \right)^{-\frac{1}{2}}.$$

The lemma is proved. □

**Lemma 3.2** *Let  $4m_c^2 \geq (3\ell - 1)^2$ , then*

$$\begin{aligned} & \int_0^{z-1} \left| \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} F \left( \frac{1}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right| dy \\ & \lesssim (1 + \ln(z))^{1 - \text{sign } M} \quad \text{for all } z \in [1, \infty). \end{aligned}$$

**Proof.** According to [14, (7.3)]

$$\left| F \left( \frac{1}{2} + iM, \frac{1}{2} + iM; 1; \zeta \right) \right| \leq C(1 - \ln(1 - \zeta))^{1 - \text{sign } M} \quad (3.20)$$

we obtain

$$\begin{aligned} & \int_0^{z-1} \left| \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} F \left( \frac{1}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right| dy \\ & \leq C \int_0^{z-1} \left| \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} \left( 1 - \ln \left( \frac{4z}{(z+1)^2 - y^2} \right) \right)^{1 - \text{sign } M} \right| dy. \end{aligned}$$

If  $M > 0$ , then

$$\int_0^{z-1} \left| \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} F \left( \frac{1}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right| dy \leq C \quad \text{for all } z \in [1, \infty).$$

If  $M = 0$ , then

$$\begin{aligned} & \int_0^{z-1} \left| \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} \left( 1 - \ln \left( \frac{4z}{(z+1)^2 - y^2} \right) \right) \right| dy \\ & \leq C + \int_0^{z-1} \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} \left| \ln \left( \frac{4z}{(z+1)^2 - y^2} \right) \right| dy. \end{aligned}$$

For  $z > R > 6$ , in the last integral, we split the interval of integration  $[0, z-1]$  into  $[0, \sqrt{(z+1)^2 - 8z}]$  and  $[\sqrt{(z+1)^2 - 8z}, z-1]$ . For the first interval  $y \leq \sqrt{(z+1)^2 - 8z}$  implies  $8z \leq (z+1)^2 - y^2$ . It follows

$$\begin{aligned} \int_0^{\sqrt{(z+1)^2 - 8z}} \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} \left| \ln \left( \frac{4z}{(z+1)^2 - y^2} \right) \right| dy & \leq (\ln z) \int_0^{z-1} \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} dy \\ & \leq C \ln z. \end{aligned}$$

For the second interval  $y \in [\sqrt{(z+1)^2 - 8z}, z-1]$  and we obtain

$$\begin{aligned} \int_{\sqrt{(z+1)^2 - 8z}}^{z-1} \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} \left| \ln \left( \frac{4z}{(z+1)^2 - y^2} \right) \right| dy & \leq C \int_{\sqrt{(z+1)^2 - 8z}}^{z-1} \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} dy \\ & \leq C. \end{aligned}$$

The lemma is proved. □

### 3.1 Estimates of integrals of the kernel functions. Proof of Theorem 1.1

**Lemma 3.3** *Let  $4m_c^2 \geq (3\ell - 1)^2$ , then*

$$\int_0^{\phi(t) - \phi(b)} |E(r, t; b; \mathcal{M})| dr \leq C |\phi(b)|^{\frac{1}{1-\tau}} \left( 1 + \ln \left( \frac{\phi(b)}{\phi(t)} \right) \right)^{1 - \text{sign } M}$$

for all  $t \in [1, \infty)$ ,  $1 \leq b \leq t$ .

**Proof.** Denote

$$z = \left( \frac{\phi(b)}{\phi(t)} \right) = \left( \frac{t}{b} \right)^{\ell-1}, \quad y = -\frac{r}{\phi(t)}, \quad (3.21)$$

then, from (2.11) we derive

$$\begin{aligned} & \int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \\ & \leq C_\ell \int_0^{\phi(t)-\phi(b)} \left| \phi(b)^{\frac{\ell+2i\mathcal{M}}{1-\ell}} \left( (\phi(t) + \phi(b))^2 - r^2 \right)^{-\frac{i\mathcal{M}}{1-\ell}} F \left( i \frac{\mathcal{M}}{1-\ell}, i \frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(b))^2 - r^2}{(\phi(t) + \phi(b))^2 - r^2} \right) \right| dr \\ & \lesssim |\phi(b)|^{\frac{1}{1-\ell}} \int_0^{z^{-1}} \left( (z+1)^2 - y^2 \right)^{-\frac{1}{2}} \left| F \left( \frac{1}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right| dy. \end{aligned}$$

Then we apply Lemma 3.2 and obtain

$$\int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \leq C |\phi(b)|^{\frac{1}{1-\ell}} (1 + \ln(z))^{1 - \text{sign } M}.$$

The lemma is proved.  $\square$

**Lemma 3.4** *Let  $4m_c^2 \geq (3\ell - 1)^2$ , then*

$$\int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr \leq C (1 + \ln t)^{1 - \text{sign } M} \quad \text{for all } t \in [1, \infty).$$

**Proof.** We use notations (3.21) with  $b = 1$ :

$$z := \frac{\phi(1)}{\phi(t)} = t^{\ell-1} \in [1, \infty), \quad y := -\frac{r}{\phi(t)} \quad (3.22)$$

and obtain

$$\int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr \leq C \int_0^{z^{-1}} \left| \left( (1+z)^2 - y^2 \right)^{-\frac{1}{2}} F \left( \frac{1}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right| dy.$$

The application of Lemma 3.2 completes the proof.  $\square$

**Lemma 3.5** *Let  $4m_c^2 \geq (3\ell - 1)^2$ , then*

$$\int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr \leq C t^{\frac{\ell-1}{2}} (1 + \ln(t))^{1 - \text{sign } M} \quad \text{for all } t \in [1, \infty).$$

**Proof.** We use notations (3.22) and obtain

$$\begin{aligned} K_2(r, t; \mathcal{M}; 1) &= \frac{2^{2+2iM} \left( \frac{1}{2} + iM \right) z^{2+2iM} \left( (z+1)^2 - y^2 \right)^{-\frac{1}{2} - iM}}{(z-1)^2 - y^2} \\ &\quad \times \left\{ (z-1) F \left( iM + \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right. \\ &\quad \left. + \frac{2(1-y^2-z^2)}{(z+1)^2 - y^2} F \left( iM + \frac{1}{2}, iM + \frac{3}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right\}. \end{aligned}$$

By [2, (28) Sec. 2.8],

$$F(a+1, a; 1; z) = -\frac{(1-a)F(a-1, a; 1; z) + (2a-1)F(a, a; 1; z)}{a(z-1)}, \quad (3.23)$$

we derive

$$F\left(\frac{3}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) = \frac{((z+1)^2 - r^2)}{2(1+2iM)z} \left( 2iMF\left(iM + \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - r^2}{(z+1)^2 - r^2}\right) + \left(\frac{1}{2} - iM\right) F\left(iM - \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - r^2}{(z+1)^2 - r^2}\right) \right).$$

It follows

$$\begin{aligned} K_2(r, t; \mathcal{M}; 1) &= 2^{2+2iM} \left(\frac{1}{2} + iM\right) \frac{z^{2+2iM}}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2} - iM} \\ &\times \left\{ \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z - 1 \right) F\left(iM + \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) + \left(\frac{1}{2} - iM\right) \frac{1-y^2-z^2}{(1+2iM)z} F\left(iM - \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) \right\} \end{aligned}$$

and

$$\begin{aligned} \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr &\lesssim z \int_0^{z-1} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}} \\ &\times \left| \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z - 1 \right) F\left(iM + \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) + \left(\frac{1}{2} - iM\right) \frac{1-y^2-z^2}{(1+2iM)z} F\left(iM - \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) \right| dy. \end{aligned}$$

For a small positive  $\epsilon$  we have

$$\begin{aligned} F\left(\frac{1}{2} + iM, \frac{1}{2} + iM; 1; \epsilon\right) &= 1 + \left(\frac{1}{2} + iM\right)^2 \epsilon + \frac{1}{64}(2M-i)^2(2M-3i)^2\epsilon^2 + O(\epsilon^3), \\ F\left(-\frac{1}{2} + iM, \frac{1}{2} + iM; 1, \epsilon\right) &= 1 - \frac{1}{4}(2M-i)(2M+i)\epsilon + \frac{1}{64}(2M-3i)(2M-i)^2(2M+i)\epsilon^2 + O(\epsilon^3). \end{aligned}$$

We divide the domain of integration into two zones,

$$Z_1(\epsilon, z) := \left\{ (z, y) \left| \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \leq \epsilon, 0 \leq y \leq z-1 \right. \right\}, \quad (3.24)$$

$$Z_2(\epsilon, z) := \left\{ (z, y) \left| \epsilon \leq \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}, 0 \leq y \leq z-1 \right. \right\}, \quad (3.25)$$

and split the integral into two parts,

$$\begin{aligned} &\int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr \\ &= \int_{(z,y) \in Z_1(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| |\phi(t)| dy + \int_{(z,y) \in Z_2(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| |\phi(t)| dy. \end{aligned}$$

In the first zone we have

$$\frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \leq \epsilon < 1. \quad (3.26)$$

Consequently, the integral over the first zone reads

$$\begin{aligned}
& \int_{Z_1(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| dr \\
& \lesssim z \int_{y \in Z_1(\epsilon, z)} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}} \\
& \quad \times \left| \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z-1 \right) \left\{ 1 + \left( \frac{1}{2} + iM \right)^2 \epsilon + \frac{1}{64} (2M-i)^2 (2M-3i)^2 \epsilon^2 + O(\epsilon^3) \right\} \right. \\
& \quad + \left( \frac{1}{2} - iM \right) \frac{1-y^2-z^2}{(1+2iM)z} \left\{ 1 - \frac{1}{4} (2M-i)(2M+i)\epsilon \right. \\
& \quad \left. \left. + \frac{1}{64} (2M-3i)(2M-i)^2 (2M+i)\epsilon^2 + O(\epsilon^3) \right\} \right| dy.
\end{aligned}$$

If we split the last integral into three parts, then for the first part, which is defined by

$$\begin{aligned}
A_0 & := z \int_{y \in Z_1(\epsilon, z)} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}} \\
& \quad \times \left| \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z-1 \right) + \left( \frac{1}{2} - iM \right) \frac{1-y^2-z^2}{(1+2iM)z} \right| dy,
\end{aligned}$$

we have

$$A_0 \lesssim \int_0^{z-1} \frac{1}{\sqrt{(z+1)^2 - y^2}} dy = \tan^{-1} \left( \frac{z-1}{\sqrt{z}} \right) \lesssim 1 \quad \text{for all } z \in [1, \infty).$$

Next, for  $A_1$ , which is defined by

$$\begin{aligned}
A_1 & := z \int_{y \in Z_1(\epsilon, z)} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}} \\
& \quad \times \left| \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z-1 \right) \left\{ \left( \frac{1}{2} + iM \right)^2 \epsilon \right\} \right. \\
& \quad \left. + \left( \frac{1}{2} - iM \right) \frac{1-y^2-z^2}{(1+2iM)z} \left\{ -\frac{1}{4} (2M-i)(2M+i)\epsilon \right\} \right| dy,
\end{aligned}$$

where  $\epsilon = \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}$ , we obtain

$$\begin{aligned}
A_1 & = z \int_{y \in Z_1(\epsilon, z)} ((z+1)^2 - y^2)^{-\frac{3}{2}} \\
& \quad \times \left| \frac{4M^2(y-z+1)(y+z-1) - 8iM(y^2+z-1) + y^2 + (z-1)(3z+1)}{8z} \right| dy \\
& \lesssim \int_{y \in Z_1(\epsilon, z)} ((z+1)^2 - y^2)^{-\frac{3}{2}} z^2 dy \\
& \lesssim z^{\frac{1}{2}} \quad \text{for all } z \in [1, \infty).
\end{aligned}$$

Similarly, for  $A_2$ , which is defined by

$$A_2 := z \int_{y \in Z_1(\epsilon, z)} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}}$$

$$\begin{aligned} & \times \left| \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z - 1 \right) \left\{ \frac{1}{64} (2M-i)^2 (2M-3i)^2 \epsilon^2 \right\} \right. \\ & \left. + \left( \frac{1}{2} - iM \right) \frac{1-y^2-z^2}{(1+2iM)z} \left\{ \frac{1}{64} (2M-3i)(2M-i)^2 (2M+i)\epsilon^2 \right\} \right| dy, \end{aligned}$$

we derive

$$\begin{aligned} A_2 & \lesssim z^2 \int_{y \in Z_1(\epsilon, z)} ((z+1)^2 - y^2)^{-\frac{5}{2}} ((z-1)^2 - y^2) dy \\ & \lesssim z^{\frac{1}{2}} \quad \text{for all } z \in [1, \infty). \end{aligned}$$

Finally, for  $A_3$ , which is defined by

$$\begin{aligned} A_3 & := z \int_{y \in Z_1(\epsilon, z)} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}} \\ & \times \left| \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z - 1 \right) O(\epsilon^3) + \left( \frac{1}{2} - iM \right) \frac{1-y^2-z^2}{(1+2iM)z} O(\epsilon^3) \right| dy, \end{aligned}$$

we obtain

$$\begin{aligned} A_3 & \lesssim z^2 \int_0^{z-1} ((z+1)^2 - y^2)^{-\frac{7}{2}} ((z-1)^2 - y^2)^2 dy = z^2 \frac{4(z-1)^5}{15\sqrt{z}(z+1)^6} \\ & \lesssim z^{\frac{1}{2}} \quad \text{for all } z \in [1, \infty). \end{aligned}$$

Thus,

$$\int_{Z_1(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| dr \leq Cz^{\frac{1}{2}} \quad \text{for all } z \in [1, \infty).$$

In the second zone,  $Z_2(\epsilon, z)$ , we have

$$\frac{1}{(z-1)^2 - y^2} \leq \frac{1}{\epsilon} \frac{1}{(z+1)^2 - y^2} \quad (3.27)$$

and due to (3.20)

$$\begin{aligned} \left| F \left( \frac{1}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right| & \leq C(1 + \ln(z))^{1 - \text{sign } M}, \\ \left| F \left( -\frac{1}{2} + iM, \frac{1}{2} + iM; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right| & \leq C. \end{aligned}$$

Then

$$\begin{aligned} \int_{Z_2(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| dr & \lesssim z \int_{y \in Z_1(\epsilon, z)} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}} \\ & \times \left| \left\{ \left( \frac{2M(1-y^2-z^2)}{(2M-i)z} + z - 1 \right) F \left( iM + \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right. \right. \\ & \left. \left. + \left( \frac{1}{2} - iM \right) \frac{1-y^2-z^2}{(1+2iM)z} F \left( iM - \frac{1}{2}, iM + \frac{1}{2}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right\} \right| dy \\ & \lesssim z^2 (1 + \ln(z))^{1 - \text{sign } M} \int_0^{z-1} \frac{1}{(z-1)^2 - y^2} ((z+1)^2 - y^2)^{-\frac{1}{2}} dy \\ & \lesssim (1 + \ln(z))^{1 - \text{sign } M} z^{\frac{1}{2}} \quad \text{for all } z \in [1, \infty). \end{aligned}$$

The lemma is proved.  $\square$

### 3.2 Estimates of the solution of the problem without potential and nonlinear terms. Proof of Theorem 1.1

**Proposition 3.6** *Let  $\ell > 1$  and  $4m_c^2 \geq (3\ell - 1)^2$  and  $\gamma \in \mathbb{R}$ . Then the solution  $\psi_{ID}$  of the problem (1.7) for the equation*

$$\frac{\partial^2 \psi}{\partial t^2} + \frac{3\ell}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}(x, \partial_x) \psi + t^{-2} m_c^2 \psi = 0, \quad (3.28)$$

satisfies

$$t^\gamma \|\psi_{ID}(\cdot, t)\|_{H(s)} \leq C t^{\gamma-\ell} (\|\psi_1\|_{H(s)} t^{-\frac{\ell-1}{2}} + \|\psi_0\|_{H(s)}) (1 + \ln(t))^{1- \text{sign } M} \quad \text{for all } t \in [1, \infty).$$

**Proof.** Indeed, according to [17, Theorem 4.1] if  $f = 0$ , then

$$\begin{aligned} \psi_{ID}(x, t) &= t^{k-} \int_0^{\phi(t)-\phi(1)} K_1(r, t; \mathcal{M}; 1) v_{\psi_1(x)}(x, r) dr \\ &\quad - k_- t^{k-} \int_0^{\phi(t)-\phi(1)} K_1(r, t; \mathcal{M}; 1) v_{\psi_0(x)}(x, r) dr \\ &\quad + t^{k-} \left( \frac{\phi(t)}{\phi(1)} \right)^{-i \frac{\mathcal{M}}{1-\ell}} v_{\psi_0}(x, \phi(t) - \phi(1)) + t^{k-} \frac{1}{\phi(1)} \int_0^{\phi(t)-\phi(1)} K_2(r, t; \mathcal{M}; 1) v_{\psi_0}(x, r) dr. \end{aligned}$$

It follows,

$$\begin{aligned} \|\psi(x, t)\|_{H(s)} &\leq C t^{\frac{1-3\ell}{2}} \int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| \|v_{\psi_1}(x, r)\|_{H(s)} dr \\ &\quad + C t^{\frac{1-3\ell}{2}} \int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| \|v_{\psi_0}(x, r)\|_{H(s)} dr \\ &\quad + C t^{\frac{1-3\ell}{2}} t^{-\frac{1-\ell}{2}} \|v_{\psi_0}(x, \phi(t) - \phi(1))\|_{H(s)} \\ &\quad + C t^{\frac{1-3\ell}{2}} \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| \|v_{\psi_0}(x, r)\|_{H(s)} dr. \end{aligned}$$

Next we apply (2.18) and Lemmas 3.4, 3.5 and obtain

$$\begin{aligned} \|\psi(x, t)\|_{H(s)} &\leq C \|\psi_1\|_{H(s)} t^{\frac{1-3\ell}{2}} (1 + \ln(t))^{1- \text{sign } M} + C \|\psi_0\|_{H(s)} t^{\frac{1-3\ell}{2}} (1 + \ln(t))^{1- \text{sign } M} \\ &\quad + C \|\psi_0\|_{H(s)} t^{-\ell} + C \|\psi_0\|_{H(s)} t^{-\ell} (1 + \ln(t))^{1- \text{sign } M} \\ &\leq C (\|\psi_1\|_{H(s)} t^{-\frac{\ell-1}{2}} + \|\psi_0\|_{H(s)}) t^{-\ell} (1 + \ln(t))^{1- \text{sign } M}. \end{aligned}$$

The proposition is proved. □

### 3.3 Estimates of the potential term. Proof of Theorem 1.1

**Proposition 3.7** *If  $\ell > 1$ ,  $\gamma \in \mathbb{R}$ ,  $4m_c^2 \geq (3\ell - 1)^2$ , and the potential  $V$  satisfies condition (1.6), then for  $M \geq 0$  we have*

$$t^\gamma \|G[V\Phi]\|_{H(s)} \leq C \epsilon_P \left( \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right) \begin{cases} 1 & \text{if } 3\ell - 1 > 2\gamma \\ (\ln(t))^{1- \text{sign } M} t^{\frac{1}{2}(2\gamma - 3\ell + 1)} & \text{if } 3\ell - 1 < 2\gamma \\ \ln(t)(\ln(t) + 1)^{1- \text{sign } M} & \text{if } 3\ell - 1 = 2\gamma \end{cases}$$

for all  $t \in [1, \infty)$ .

**Proof.** Indeed, from Lemma 3.3 it follows

$$\begin{aligned}
\|G[F]\|_{H(s)} &\lesssim t^{\mathfrak{R}k-} \int_1^t b^{-\mathfrak{R}k-} db \int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| \|\mathcal{E}\mathcal{E}[F](x, r; b)\|_{H(s)} dr \\
&\lesssim t^{\mathfrak{R}k-} \int_1^t b^{-\mathfrak{R}k-} \|F(x, b)\|_{H(s)} db \int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \\
&\lesssim t^{\mathfrak{R}k-} \int_1^t b^{-\mathfrak{R}k-} \|F(x, b)\|_{H(s)} |\phi(b)|^{\frac{1}{1-\ell}} \left(1 + \ln\left(\frac{\phi(b)}{\phi(t)}\right)\right)^{1-\text{sign } M} db \\
&\lesssim t^{\frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell+1}{2}} \|F(x, b)\|_{H(s)} \left(1 + \ln\left(\frac{t}{b}\right)\right)^{1-\text{sign } M} db. \tag{3.29}
\end{aligned}$$

In particular, for the potential term we obtain

$$\begin{aligned}
\|G[V\Phi]\|_{H(s)} &\lesssim t^{\frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell+1}{2}-\delta} \|\Phi(x, b)\|_{H(s)} \left(1 + \ln\left(\frac{t}{b}\right)\right)^{1-\text{sign } M} db \\
&\lesssim \left(\sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)}\right) t^{\frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell+1}{2}-\delta-\gamma} \left(1 + \ln\left(\frac{t}{b}\right)\right)^{1-\text{sign } M} db.
\end{aligned}$$

If  $\delta = 2$ , then

$$t^\gamma \|G[V\Phi]\|_{H(s)} \lesssim \left(\sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)}\right) t^{\gamma+\frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell-3}{2}-\gamma} \left(1 + \ln\left(\frac{t}{b}\right)\right)^{1-\text{sign } M} db$$

and with  $\frac{3\ell-3}{2} - \gamma \neq -1$  and  $M > 0$ , we obtain

$$t^{\gamma+\frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell-3}{2}-\gamma} db \lesssim \begin{cases} 1 & \text{if } 3\ell - 1 - 2\gamma > 0, \\ t^{\gamma+\frac{1-3\ell}{2}} & \text{if } 3\ell - 1 - 2\gamma < 0. \end{cases}$$

If  $\frac{3\ell-3}{2} - \gamma = -1$  and  $M > 0$ , then

$$t^{\gamma+\frac{1-3\ell}{2}} \int_1^t b^{-1} db = t^{\gamma+\frac{1-3\ell}{2}} \ln(t).$$

Thus, if  $M > 0$ , then

$$t^\gamma \|G[V\Phi]\|_{H(s)} \lesssim \left(\sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)}\right) \begin{cases} 1 & \text{if } 3\ell - 1 - 2\gamma > 0, \\ t^{\gamma+\frac{1-3\ell}{2}} & \text{if } 3\ell - 1 - 2\gamma < 0, \\ \ln(t) & \text{if } 3\ell - 1 - 2\gamma = 0. \end{cases}$$

If  $M = 0$  and  $2\gamma - 3\ell + 1 \neq 0$ , then we obtain

$$\begin{aligned}
&t^{\gamma+\frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell+1}{2}-2-\gamma} \left(1 + \ln\left(\frac{t}{b}\right)\right)^{1-\text{sign } M} db \\
&= \frac{1}{(2\gamma - 3\ell + 1)^2} \left( -6\ell t^{\gamma-\frac{3\ell}{2}+\frac{1}{2}} + 4\gamma t^{\gamma-\frac{3\ell}{2}+\frac{1}{2}} - 2t^{\gamma-\frac{3\ell}{2}+\frac{1}{2}} \right. \\
&\quad \left. -6\ell \ln(t) t^{\gamma-\frac{3\ell}{2}+\frac{1}{2}} + 4\gamma \ln(t) t^{\gamma-\frac{3\ell}{2}+\frac{1}{2}} + 2\ln(t) t^{\gamma-\frac{3\ell}{2}+\frac{1}{2}} + 6\ell + 2 - 4\gamma \right).
\end{aligned}$$

Then if  $M = 0$  we obtain

$$t^{\gamma+\frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell+1}{2}-2-\gamma} \left(1 + \ln\left(\frac{t}{b}\right)\right)^{1-\text{sign } M} db \lesssim \begin{cases} 1 & \text{if } 2\gamma - 3\ell + 1 < 0, \\ t^{\gamma-\frac{3\ell}{2}+\frac{1}{2}} \ln(t) & \text{if } 2\gamma - 3\ell + 1 > 0, \\ \ln(t)(\ln(t) + 2) & \text{if } 2\gamma - 3\ell + 1 = 0. \end{cases}$$



Thus, if  $M = 0$ , then

$$t^\gamma \|G[V\Phi]\|_{H(s)} \lesssim \left( \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right) \begin{cases} 1 & \text{if } 3\ell - 1 - 2\gamma > 0, \\ t^{\gamma - \frac{3\ell}{2} + \frac{1}{2}} \ln(t) & \text{if } 3\ell - 1 - 2\gamma < 0, \\ \ln(t)(\ln(t) + 1) & \text{if } 3\ell - 1 - 2\gamma = 0. \end{cases}$$

Note, if  $\gamma \leq \ell$  and  $\ell > 1$ , then  $3\ell - 1 - 2\gamma > 0$ . The lemma is proved.  $\square$

### 3.4 Estimates of the nonlinear term. Proof of Theorem 1.1

**Theorem 3.8** *Assume that  $\ell > 1$ ,  $\gamma \in \mathbb{R}$  and the semilinear term is Lipschitz continuous with exponent  $\alpha \geq 0$  in the space  $H(s)$ . Then for the operator  $G$  with  $M \geq 0$ , the following estimate holds:*

$$t^\gamma \|G[\Psi(\Phi)]\|_{H(s)} \lesssim \left( \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right)^{1+\alpha} \begin{cases} t^{\frac{1}{2}(2\gamma - 3\ell + 1)} (\log(t))^{1 - \text{sign } M} & \text{if } \gamma > \frac{3(\ell+1)}{2(\alpha+1)}, \\ t^{2-\gamma\alpha} & \text{if } \gamma < \frac{3(\ell+1)}{2(\alpha+1)}, \\ t^{2-\gamma\alpha} \ln(t)(\ln(t) + 1)^{1 - \text{sign } M} & \text{if } \gamma = \frac{3(\ell+1)}{2(\alpha+1)} \end{cases}$$

for all  $t \in [1, \infty)$  and all  $\Phi \in H(s)$ .

**Proof.** Indeed, with  $F = \Psi(\Phi)$  from (3.29) we derive

$$t^\gamma \|G[\Psi(\Phi)]\|_{H(s)} \lesssim \left( \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right)^{1+\alpha} I_{\ell, \alpha, M}(t),$$

where

$$I_{\ell, \alpha, M}(t) := t^{\gamma + \frac{1-3\ell}{2}} \int_1^t b^{\frac{3\ell+1}{2} - \gamma(1+\alpha)} \left( 1 + \ln\left(\frac{t}{b}\right) \right)^{1 - \text{sign } M} db.$$

We use the change of variable  $z = t/b$ ,  $db = -tz^{-2}dz$ , and obtain

$$I_{\ell, \alpha, M}(t) = t^{2-\gamma\alpha} \int_1^t z^{-\frac{3\ell+1}{2} + \gamma(1+\alpha) - 2} (1 + \ln z)^{1 - \text{sign } M} dz.$$

If  $M > 0$  and  $-\frac{3\ell+1}{2} + \gamma(1+\alpha) - 2 \neq -1$ , then

$$I_{\ell, \alpha, M}(t) \lesssim \begin{cases} t^{-\frac{3\ell+1}{2} + \gamma + 1} & \text{if } -\frac{3\ell+1}{2} + \gamma(1+\alpha) - 1 > 0, \\ t^{2-\gamma\alpha} & \text{if } -\frac{3\ell+1}{2} + \gamma(1+\alpha) - 1 < 0. \end{cases}$$

If  $M > 0$  and  $-\frac{3\ell+1}{2} + \gamma(1+\alpha) - 2 = -1$ , then  $I_{\ell, \alpha, M}(t) = t^{2-\gamma\alpha} \ln(t)$ . Thus, if  $M > 0$

$$t^\gamma \|G[\Psi(\Phi)]\|_{H(s)} \lesssim \left( \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right)^{1+\alpha} \begin{cases} t^{-\frac{3\ell+1}{2} + \gamma + 1} & \text{if } -\frac{3\ell+1}{2} + \gamma(1+\alpha) - 1 > 0, \\ t^{2-\gamma\alpha} & \text{if } -\frac{3\ell+1}{2} + \gamma(1+\alpha) - 1 < 0, \\ t^{2-\gamma\alpha} \ln(t) & \text{if } -\frac{3\ell+1}{2} + \gamma(1+\alpha) - 1 = 0. \end{cases}$$

If  $M = 0$  and  $-\frac{3\ell+1}{2} + \gamma(1+\alpha) - 2 \neq -1$ , then

$$\begin{aligned} I_{\ell, \alpha, M}(t) &= \frac{2}{(2(\alpha+1)\gamma - 3(\ell+1))^2} t^{2-\gamma\alpha} t^{\frac{1}{2}(-3)(\ell+1)} \\ &\quad \times \left[ \ln(t)(2(\alpha+1)\gamma - 3(\ell+1))t^{(\alpha+1)\gamma} + (2(\alpha+1)\gamma - 3\ell - 5) \left( t^{(\alpha+1)\gamma} - t^{\frac{3(\ell+1)}{2}} \right) \right] \\ &\lesssim t^{\frac{1}{2}(-2\alpha\gamma - 3\ell + 1)} \left[ \ln(t)t^{(\alpha+1)\gamma} + |2(\alpha+1)\gamma - 3\ell - 5| t^{\frac{3(\ell+1)}{2}} \right] \\ &\lesssim t^{\frac{1}{2}(-2\alpha\gamma - 3\ell + 1)} \begin{cases} t^{(\alpha+1)\gamma} \ln(t) & \text{if } (\alpha+1)\gamma \geq \frac{3(\ell+1)}{2}, \\ t^{\frac{3(\ell+1)}{2}} & \text{if } (\alpha+1)\gamma < \frac{3(\ell+1)}{2}. \end{cases} \end{aligned}$$

If  $M = 0$  and  $-\frac{3\ell+1}{2} + \gamma(1 + \alpha) - 2 = -1$ , then

$$I_{\ell,\alpha,M}(t) = t^{2-\gamma\alpha} \frac{1}{2} \ln(t)(\ln(t) + 2) \lesssim t^{2-\gamma\alpha} \ln(t)(\ln(t) + 1).$$

Thus, if  $M = 0$ , then

$$\begin{aligned} & t^\gamma \|G[\Psi(\Phi)]\|_{H(s)} \\ & \lesssim \left( \sup_{\tau \in [1,t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right)^{1+\alpha} \begin{cases} t^{\frac{1}{2}(2\gamma-3\ell+1)} \log(t) & \text{if } (\alpha+1)\gamma > \frac{3(\ell+1)}{2}, \\ t^{2-\gamma\alpha} & \text{if } (\alpha+1)\gamma < \frac{3(\ell+1)}{2}, \\ t^{2-\gamma\alpha} \ln(t)(\ln(t) + 1) & \text{if } -\frac{3\ell+1}{2} + \gamma(1 + \alpha) - 1 = 0. \end{cases} \end{aligned}$$

The lemma is proved.  $\square$

### 3.5 Proof of global existence. Proof of Theorem 1.1

We intend to apply Banach's fixed-point theorem. We use the Lipschitz condition  $(\mathcal{L})$  to estimate nonlinear term. First, we consider the following integral equation

$$\psi(x, t) = \psi_{ID}(x, t) + G[V\psi](x, t) + G[F(\cdot)\Psi(\cdot, \psi)](x, t), \quad (3.30)$$

where the function  $\psi_{ID}(x, t) \in C([1, \infty); H(s))$  is given. We set  $a_{acc} = 1$  and  $c = 1$ .

We consider the Cauchy problem in the Sobolev space  $H(s)$  with  $s > 3/2$ , which is an algebra.

Denote  $\tilde{C}^\ell([1, T]; H(s))$  the complete subspace of  $C^\ell([1, T]; H(s))$  of all functions  $f = f(x, t)$  with  $\text{supp } f \subset \{(x, t) \in \mathbb{R}^3 \times [1, \infty) \mid |x| \geq R_{ID} - (\phi(t) - \phi(1))\}$ . For every  $T > 1$ , due to Section 2, the operator  $G$  maps

$$G : \tilde{C}([1, T]; H(s)) \longrightarrow \tilde{C}^2([1, T]; H(s))$$

continuously. Consider the mapping

$$S[\Phi] = \psi_{ID} + G[V\Phi] + G[F\Psi(\Phi)], \quad (3.31)$$

where  $\psi_{ID} \in X(R, H(s), \gamma)$ . The function  $\psi_{ID}$  can be generated by initial data (1.5),(1.7). In that case it belongs to  $X(R, H(s), \gamma)$  if  $\gamma \leq \ell$  and

$$t^\gamma \|\psi_{ID}(x, t)\|_{H(s)} \leq C(\|\psi_0\|_{H(s)} + \|\psi_1\|_{H(s)}).$$

It is easily seen that  $\text{supp } S[\Phi] \subset \text{supp } \Phi$  if  $\text{supp } \Phi \subset \text{supp } \psi_{ID}$ . We claim that since  $\ell > 1$  and  $\gamma \leq \ell$ , if  $\Phi \in X(R, H(s), \gamma)$  with  $\gamma \leq \frac{3\ell+1}{\alpha}$  and if  $\text{supp } \Phi \subseteq \{(x, t) \in \mathbb{R}^3 \times [0, \infty) \mid |x| > R_{ID} - (\phi(t) - \phi(1))\}$ , then  $S[\Phi] \in X(R, H(s), \gamma)$ . Moreover,  $S$  is a contraction, provided that  $\varepsilon$ ,  $\varepsilon_P$ , and  $R$  are sufficiently small.

First, we note that due to Theorem 3.8 and Propositions 3.6, 3.7, for the values of  $\alpha$ ,  $\ell$ , and  $\gamma$  assumed in the theorem, we have  $\gamma - \ell \leq 0$  and, consequently,

$$\begin{aligned} t^\gamma \|S[\Phi]\|_{H(s)} & \leq t^\gamma \|\psi_{ID}\|_{H(s)} + t^\gamma \|G[V\Phi]\|_{H(s)} + t^\gamma \|G[F\Psi(\Phi)]\|_{H(s)} \\ & \leq C t^{\gamma-\ell} (\|\psi_1\|_{H(s)} t^{-\frac{\ell-1}{2}} + \|\psi_0\|_{H(s)}) (1 + \ln(t))^{1-\text{sign } M} \\ & \quad + C \varepsilon_P \left( \sup_{\tau \in [1,t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right) \begin{cases} 1 & \text{if } 3\ell - 1 > 2\gamma \\ (\ln(t))^{1-\text{sign } M} t^{\frac{1}{2}(2\gamma-3\ell+1)} & \text{if } 3\ell - 1 < 2\gamma \\ \ln(t)(\ln(t) + 1)^{1-\text{sign } M} & \text{if } 3\ell - 1 = 2\gamma \end{cases} \\ & \quad + C_{NT} \left( \sup_{\tau \in [1,t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right)^{1+\alpha} \begin{cases} t^{\frac{1}{2}(2\gamma-3\ell+1)} (\log(t))^{1-\text{sign } M} & \text{if } \gamma > \frac{3(\ell+1)}{2(\alpha+1)}, \\ t^{2-\gamma\alpha} & \text{if } \gamma < \frac{3(\ell+1)}{2(\alpha+1)}, \\ t^{2-\gamma\alpha} \ln(t)(\ln(t) + 1)^{1-\text{sign } M} & \text{if } \gamma = \frac{3(\ell+1)}{2(1+\alpha)}. \end{cases} \end{aligned}$$

Hence, to get a contraction we set all time-dependent factors to be bounded for all times. For  $M > 0$ : we choose  $\frac{3(\ell+1)}{2(\alpha+1)} < \gamma \leq \ell \implies \alpha > \frac{\ell+3}{2\ell}$ ;  $\frac{2}{\alpha} \leq \gamma < \frac{3(\ell+1)}{2(\alpha+1)}$ ,  $\gamma \leq \ell$ ,  $\alpha > \frac{4}{3\ell-1}$ ;  $\gamma = \frac{3(\ell+1)}{2(1+\alpha)}$ ,  $\alpha > \frac{4}{3\ell-1}$ . For  $M = 0$

we choose:  $\frac{3(\ell+1)}{2(\alpha+1)} < \gamma < \ell \implies \alpha > \frac{\ell+3}{2\ell}$ ;  $\frac{2}{\alpha} \leq \gamma < \frac{3(\ell+1)}{2(\alpha+1)}$ ,  $\gamma < \ell$ ,  $\alpha > \frac{4}{3\ell-1}$ ;  $\gamma = \frac{3(\ell+1)}{2(1+\alpha)}$ ,  $\alpha > \frac{4}{3\ell-1}$ ,  $\gamma < \ell$ .  
 Note that  $\frac{\ell+3}{2\ell} > \frac{4}{3\ell-1}$ .

Thus, we choose  $\gamma$  in accordance to the cases above, and the last inequality proves that the operator  $S$  maps  $X(R, s, \gamma)$  into itself if

$$\|\psi_0\|_{H(s)} + \|\psi_1\|_{H(s)} \leq \varepsilon,$$

$\varepsilon_P$ ,  $\varepsilon$ , and  $R$  are sufficiently small, namely, if  $\varepsilon_P C_{pt} < 1$  and

$$\frac{1}{1 - \varepsilon_P C_{pt}} C_{id} \varepsilon + \frac{1}{1 - \varepsilon_P C_{pt}} C_{nt} R^{\alpha+1} < R.$$

We derive the contraction property of  $S$  from

$$d(S[\Phi_1], S[\Phi_2]) \leq d(\Phi_1, \Phi_2)(C_{pt}\varepsilon_0 + C_{nt}R^\alpha).$$

Indeed, due to Theorem 3.8, Proposition 3.7, and conditions of theorem we have

$$\begin{aligned} & t^\gamma \|S[\Phi_1](\cdot, t) - S[\Phi_2](\cdot, t)\|_{H(s)} \\ & \leq t^\gamma \|G[V(\Phi_1 - \Phi_2)]\|_{H(s)} + t^\gamma \|G[F(\Psi(\Phi_1) - \Psi(\Phi_2))]\|_{H(s)} \\ & \leq C_{pt}\varepsilon_P \sup_{\tau \in [1, t]} \tau^\gamma \|(\Phi_1 - \Phi_2)(x, \tau)\|_{H(s)} + Ct^\gamma \|G[F(\Psi(\Phi_1) - \Psi(\Phi_2))]\|_{H(s)}, \end{aligned}$$

while according to above used arguments and due to the condition  $(\mathcal{L})$ , we obtain

$$\begin{aligned} & \|G[(\Psi(\Phi_1) - \Psi(\Phi_2))]\|_{H(s)} \\ & \leq Ct^{\mathfrak{R}k-} \int_1^t b^{-\mathfrak{R}k-} \|(\Psi(\Phi_1) - \Psi(\Phi_2))(x, b)\|_{H(s)} |\phi(b)|^{\frac{1}{1-\ell}} \left(1 + \ln \frac{\phi(b)}{\phi(t)}\right)^{1 - \text{sign } M} db \\ & \leq Ct^{\frac{1-3\ell}{2}} \int_1^t b^{\frac{1+3\ell}{2}} \|\Phi_1(x, b) - \Phi_2(x, b)\|_{H(s)} \left(\|\Phi_1(x, b)\|_{H(s)}^\alpha + \|\Phi_2(x, b)\|_{H(s)}^\alpha\right) \left(1 + \ln \frac{\phi(b)}{\phi(t)}\right)^{1 - \text{sign } M} db \\ & \leq Ct^{\frac{1-3\ell}{2}} \int_1^t b^{\frac{1+3\ell}{2} - \gamma(1+\alpha)} \left(\sup_{b \in [1, t]} b^\gamma \|\Phi_1(x, b) - \Phi_2(x, b)\|_{H(s)}\right) \\ & \quad \times \left(\left(\sup_{b \in [1, t]} b^\gamma \|\Phi_1(x, b)\|_{H(s)}\right)^\alpha + \left(\sup_{b \in [1, t]} b^\gamma \|\Phi_2(x, b)\|_{H(s)}\right)^\alpha\right) \left(1 + \ln \frac{\phi(b)}{\phi(t)}\right)^{1 - \text{sign } M} db. \end{aligned}$$

Further, if

$$d(\Phi_1, \Phi_2) := \sup_{b \in [1, \infty]} b^\gamma \|\Phi_1(x, b) - \Phi_2(x, b)\|_{H(s)}$$

and

$$R := \max \left\{ \sup_{b \in [1, t]} b^\gamma \|\Phi_1(x, b)\|_{H(s)}, \sup_{b \in [1, t]} b^\gamma \|\Phi_2(x, b)\|_{H(s)} \right\},$$

then

$$\|G[(\Psi(\Phi_1) - \Psi(\Phi_2))]\|_{H(s)} \leq Cd(\Phi_1, \Phi_2) R^\alpha t^{\frac{1-3\ell}{2}} \int_1^t b^{\frac{1+3\ell}{2} - \gamma(1+\alpha)} \left(1 + \ln \frac{\phi(b)}{\phi(t)}\right)^{1 - \text{sign } M} db$$

and, consequently, we obtain

$$t^\gamma \|G[(\Psi(\Phi_1) - \Psi(\Phi_2))]\|_{H(s)} \leq Cd(\Phi_1, \Phi_2) R^\alpha \begin{cases} t^{\frac{1}{2}(2\gamma-3\ell+1)} (\log(t))^{1 - \text{sign } M} & \text{if } \gamma > \frac{3(\ell+1)}{2(\alpha+1)}, \\ t^{2-\gamma\alpha} & \text{if } \gamma < \frac{3(\ell+1)}{2(\alpha+1)}, \\ t^{2-\gamma\alpha} \ln(t) (\ln(t) + 1)^{1 - \text{sign } M} & \text{if } \gamma = \frac{3(\ell+1)}{2(1+\alpha)}. \end{cases}$$

Thus, in all cases we have

$$t^\gamma \|S[\Phi_1](\cdot, t) - S[\Phi_2](\cdot, t)\|_{H(s)} \leq t^\gamma \|G[V(\Phi_1 - \Phi_2)]\|_{H(s)} + t^\gamma \|G[F(\Psi(\Phi_1) - \Psi(\Phi_2))]\|_{H(s)}.$$

Finally,

$$t^\gamma \|S[\Phi_1](\cdot, t) - S[\Phi_2](\cdot, t)\|_{H(s)} \leq C_{pt}\varepsilon_P d(\Phi_1, \Phi_2) + C_{nt}d(\Phi_1, \Phi_2)R^\alpha.$$

Then we choose  $\|\psi_0\|_{H(s)} + \|\psi_1\|_{H(s)} \leq \varepsilon$  with sufficiently small  $\varepsilon$  and  $R$  such that  $C_{pt}\varepsilon_P + C_{nt}R^\alpha < 1$ . Banach's fixed point theorem completes the proof.  $\square$

## 4 Case of small mass. Proof of Theorem 1.2

Since  $\ell > 1$  for the case of real small mass  $m_c \leq [0, (3\ell - 1)/2)$ . We choose  $k_+$  in the Liouville substitution:

$$\begin{aligned} \psi(x, t) &= t^{k_+} u(x, t) = t^{\frac{1}{2}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2})} u(x, t), \\ \psi_0(x) &= u_0(x), \quad \psi_1(x) = u_1(x) + k_+ u_0(x), \quad u_1(x) = \psi_1(x) - k_+ \psi_0(x), \quad g(x, t) = t^{-k_+} f(x, t). \end{aligned}$$

Hence, by Theorem 2.2,

$$\begin{aligned} \psi(x, t) &= t^{k_+} \int_1^t b^{-k_+} db \int_0^{\phi(t)-\phi(b)} E(r, t; b; \mathcal{M}) v_f(x, r; b) dr \\ &\quad + t^{k_+} \int_0^{\phi(t)-\phi(1)} K_1(r, t; \mathcal{M}; 1) v_{\psi_1(x)}(x, r) dr \\ &\quad - k_+ t^{k_+} \int_0^{\phi(t)-\phi(1)} K_1(r, t; \mathcal{M}; 1) v_{\psi_0(x)}(x, r) dr \\ &\quad + t^{k_+} \left(\frac{\phi(t)}{\phi(1)}\right)^{-i\frac{\mathcal{M}}{1-\ell}} v_{\psi_0}(x, \phi(t) - \phi(1)) + t^{k_+} \frac{1}{\phi(1)} \int_0^{\phi(t)-\phi(1)} K_2(r, t; \mathcal{M}; 1) v_{\psi_0}(x, r) dr. \end{aligned}$$

Since  $-\frac{\ell}{\ell-1} < i\frac{\mathcal{M}}{1-\ell} < \frac{1}{2}$ , by use of (2.18) we derive

$$\begin{aligned} \|u(x, t)\|_{H(s)} &\leq C_s t^{-\frac{1}{2}\left(1-\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}\right)} \|u_0\|_{H(s)} + C_s \|u_1\|_{H(s)} \int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr \\ &\quad + C_s \|u_0\|_{H(s)} \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr. \end{aligned}$$

For the real small mass, the condition  $1 - \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1} \geq 0$  is equivalent to  $4m_c^2 \geq 4\ell(2\ell - 1)$ . Thus,

$$\begin{aligned} \|u(x, t)\|_{H(s)} &\leq C_s \|u_0\|_{H(s)} \begin{cases} t^{-\frac{1}{2}\left(1-\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}\right)} & \text{if } 4m_c^2 < 4\ell(2\ell - 1), \\ 1 & \text{if } 4m_c^2 \geq 4\ell(2\ell - 1), \end{cases} \\ &\quad + C_s \|u_1\|_{H(s)} \int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr + C_s \|u_0\|_{H(s)} \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr. \end{aligned}$$

### 4.1 Estimates of the kernel functions. Proof of Theorem 1.2

**Lemma 4.1** For  $4m_c^2 < (1 - 3\ell)^2$  and the given above parameters  $\mathcal{M}$  and  $\ell$ ,

$$\begin{aligned} \left| F\left(\frac{1}{2}, i\frac{\mathcal{M}}{1-\ell}, \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2}\right) \right| &\leq C, \\ \left| F\left(\frac{1}{2}, 1 + i\frac{\mathcal{M}}{1-\ell}; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2}\right) \right| &\leq C \begin{cases} 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell - 1), \\ z^{\frac{1}{2} - \frac{1}{2(\ell-1)}\sqrt{(1-3\ell)^2-4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell - 1) \end{cases} \end{aligned}$$

for all  $z \in [1, \infty)$ .

**Proof.** The first inequality follows from [2, (14) Sec.2.1] since  $1 + i\frac{\mathcal{M}}{\ell-1} > 0$ . Similarly, for the second one we use [2, (14),(23) Sec.2.1] and derive

$$\left| F\left(\frac{1}{2}, 1 + i\frac{\mathcal{M}}{1-\ell}; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2}\right) \right| \leq C \left(1 + \left(\frac{(z+1)^2}{4z}\right)^{i\frac{\mathcal{M}}{1-\ell}}\right)$$

for all  $z \in [1, \infty)$ . The lemma is proved.  $\square$

**Lemma 4.2** *If  $\ell > 1$  and  $4m_c^2 < (1-3\ell)^2$ , then for the given above parameters  $\mathcal{M}$  and  $\ell$ ,*

$$\int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \lesssim b^{-1+\sqrt{(1-3\ell)^2-4m_c^2}} |\phi(b) + \phi(t)|^{\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}}$$

for all  $t \in [1, \infty)$ .

**Proof.** From (2.11)

$$\begin{aligned} \int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr &\lesssim \int_0^{\phi(t)-\phi(b)} \left| \phi(b)^{\frac{\ell+2i\mathcal{M}}{1-\ell}} \left( (\phi(t) + \phi(b))^2 - r^2 \right)^{-\frac{i\mathcal{M}}{1-\ell}} \right. \\ &\quad \left. \times F\left(i\frac{\mathcal{M}}{1-\ell}, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(b))^2 - r^2}{(\phi(t) + \phi(b))^2 - r^2}\right) \right| dr. \end{aligned}$$

In the notations (3.21), we write

$$\begin{aligned} &\int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \\ &\lesssim |\phi(b)^{\frac{\ell+2i\mathcal{M}}{1-\ell}} \phi(t)^{1-2\frac{i\mathcal{M}}{1-\ell}}| \int_0^{z-1} \left| \left( (z+1)^2 - y^2 \right)^{-\frac{i\mathcal{M}}{1-\ell}} F\left(i\frac{\mathcal{M}}{1-\ell}, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) \right| dy \end{aligned}$$

and then, since  $2i\frac{\mathcal{M}}{1-\ell} < 1$ ,  $\frac{\ell+2i\mathcal{M}}{1-\ell} = \frac{1}{\ell-1} - \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}$ , and  $1-2\frac{i\mathcal{M}}{1-\ell} = \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}$ , we obtain

$$\int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \lesssim \left| \phi(b)^{\frac{1}{\ell-1} - \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}} \phi(t)^{\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}} \right| \int_0^{z-1} \left( (z+1)^2 - y^2 \right)^{-\frac{i\mathcal{M}}{1-\ell}} dy.$$

On the other hand, since  $\frac{1}{2} + \frac{i\mathcal{M}}{1-\ell} = 1 - \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{2(\ell-1)} < \frac{3}{2}$ , we have

$$\begin{aligned} \int_0^{z-1} \left( (z+1)^2 - y^2 \right)^{-\frac{i\mathcal{M}}{1-\ell}} dy &= (z-1)(z+1)^{-2\frac{i\mathcal{M}}{1-\ell}} F\left(\frac{1}{2}, \frac{i\mathcal{M}}{1-\ell}; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2}\right) \\ &\lesssim (z-1)(z+1)^{-1+\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}}. \end{aligned}$$

It follows

$$\int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \lesssim |\phi(b)|^{\frac{1}{\ell-1} - \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}} |\phi(b) + \phi(t)|^{\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}}.$$

The lemma is proved.  $\square$

**Lemma 4.3** *If  $\ell > 1$  and  $4m_c^2 < (1-3\ell)^2$ , then for the given above parameters  $\mathcal{M}$  and  $\ell$*

$$\int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr \leq C \quad \text{for all } t \in [1, \infty).$$

**Proof.** According to (2.12) and  $2i\frac{\mathcal{M}}{1-\ell} < 1$  we obtain

$$\begin{aligned} & \int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr \\ & \leq C \int_0^{\phi(t)-\phi(1)} \left| \left( (\phi(t) + \phi(1))^2 - r^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} F \left( i\frac{\mathcal{M}}{1-\ell}, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(1))^2 - r^2}{(\phi(t) + \phi(1))^2 - r^2} \right) \right| dr \\ & \leq C \int_0^{\phi(t)-\phi(1)} \left( (\phi(t) + \phi(1))^2 - r^2 \right)^{i\frac{\mathcal{M}}{\ell-1}} dr. \end{aligned}$$

Here  $\phi(t) - \phi(1) \geq 0$ . In the notations (3.22), since  $2i\frac{\mathcal{M}}{1-\ell} < 1$ , we obtain

$$\int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr \lesssim |(\phi(t))^{1+2i\frac{\mathcal{M}}{\ell-1}}| \int_0^{z-1} \left( (z+1)^2 - y^2 \right)^{i\frac{\mathcal{M}}{\ell-1}} dy.$$

Then according to [2, (14) Sec. 2.1] and since  $1 + i\frac{\mathcal{M}}{\ell-1} > 0$ , we have

$$\begin{aligned} \int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr & \lesssim |(\phi(t))^{1+2i\frac{\mathcal{M}}{\ell-1}}| |(z-1)(z+1)^{2i\frac{\mathcal{M}}{\ell-1}}| F \left( \frac{1}{2}, i\frac{\mathcal{M}}{1-\ell}, \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2} \right) \\ & \lesssim 1. \end{aligned}$$

The lemma is proved. □

**Lemma 4.4** *If  $\ell > 1$  and  $4m_c^2 < (1 - 3\ell)^2$ , then we have*

$$\int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr \leq C \begin{cases} t^{\frac{\ell-1}{2} - \frac{1}{2}\sqrt{(1-3\ell)^2 - 4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1) \end{cases}$$

for all  $t \in [1, \infty)$ .

**Proof.** According to the definition of the kernel  $K_2$  we have

$$\begin{aligned} \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr & \leq \int_0^{\phi(t)-\phi(1)} \left| -i\mathcal{M}2^{1+2i\frac{\mathcal{M}}{1-\ell}} \phi^{\frac{2i\mathcal{M}}{1-\ell}}(1) \left( (\phi(t) + \phi(1))^2 - r^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \right. \\ & \quad \times \left[ \left( \frac{-\phi(t)\phi(1) + \phi^2(1)}{r^2 - (\phi(t) - \phi(1))^2} \right) F \left( i\frac{\mathcal{M}}{1-\ell}, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(1))^2 - r^2}{(\phi(t) + \phi(1))^2 - r^2} \right) \right. \\ & \quad \left. \left. - \frac{2\phi(t)\phi(1)(\phi^2(t) - \phi^2(1) - r^2)}{((\phi(t) - \phi(1))^2 - r^2)((\phi(t) + \phi(1))^2 - r^2)} \right] \right| \\ & \quad \times F \left( i\frac{\mathcal{M}}{1-\ell} + 1, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(\phi(t) - \phi(1))^2 - r^2}{(\phi(t) + \phi(1))^2 - r^2} \right) \Bigg| dr. \end{aligned}$$

In the notations (3.22), it follows

$$\begin{aligned} & \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr \\ & \leq C \left( (\phi(t))^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \int_0^{\phi(t)-\phi(1)} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \right. \\ & \quad \times \left[ \left( \frac{-z + z^2}{y^2 - (z-1)^2} \right) F \left( i\frac{\mathcal{M}}{1-\ell}, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \right. \end{aligned}$$

$$\begin{aligned}
& - \frac{2z(1-z^2-y^2)}{\left((z-1)^2-y^2\right)\left((z+1)^2-y^2\right)} F\left(i\frac{\mathcal{M}}{1-\ell}+1, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \Bigg| dr \\
\leq & C \left( (\phi(t))^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \int_0^{\phi(t)-\phi(1)} \left| \left( (z+1)^2-y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \frac{z}{(z-1)^2-y^2} \right. \\
& \times \left[ -(z-1)F\left(i\frac{\mathcal{M}}{1-\ell}, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \right. \\
& \left. \left. - \frac{2(1-z^2-y^2)}{(z+1)^2-y^2} F\left(i\frac{\mathcal{M}}{1-\ell}+1, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \right] \right| dr.
\end{aligned}$$

By (3.23) we derive

$$\begin{aligned}
& F\left(i\frac{\mathcal{M}}{1-\ell}+1, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \\
= & -\frac{1}{i\frac{\mathcal{M}}{1-\ell}\left(\frac{(z-1)^2-y^2}{(z+1)^2-y^2}-1\right)} \left\{ \left(1-i\frac{\mathcal{M}}{1-\ell}\right) F\left(i\frac{\mathcal{M}}{1-\ell}-1, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \right. \\
& \left. + \left(2i\frac{\mathcal{M}}{1-\ell}-1\right) F\left(i\frac{\mathcal{M}}{1-\ell}, i\frac{\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \right\}.
\end{aligned}$$

Then

$$\begin{aligned}
& \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr \\
\leq & Cz^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left| \left( (z+1)^2-y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \frac{1}{i2\mathcal{M}(y^2-(z-1)^2)} \right. \\
& \times \left( i(-2\mathcal{M}(y^2+z-1) + i(\ell-1)(y^2+z^2-1)) F\left(\frac{i\mathcal{M}}{1-\ell}, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \right. \\
& \left. \left. + (-\ell-i\mathcal{M}+1)(-y^2-z^2+1) F\left(\frac{i\mathcal{M}}{1-\ell}-1, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \right) \right| dy.
\end{aligned}$$

For small  $\epsilon$  we have

$$\begin{aligned}
F\left(\frac{i\mathcal{M}}{1-\ell}, \frac{i\mathcal{M}}{1-\ell}; 1; \epsilon\right) &= 1 - \frac{\mathcal{M}^2}{(\ell-1)^2}\epsilon + \frac{\mathcal{M}^2(i\ell+\mathcal{M}-i)^2}{4(\ell-1)^4}\epsilon^2 + O(\epsilon^3), \\
F\left(\frac{i\mathcal{M}}{1-\ell}-1, \frac{i\mathcal{M}}{1-\ell}; 1; \epsilon\right) &= 1 - \frac{\mathcal{M}(-i\ell+\mathcal{M}+i)}{(\ell-1)^2}\epsilon + \frac{\mathcal{M}^2(-i\ell+\mathcal{M}+i)(i\ell+\mathcal{M}-i)}{4(\ell-1)^4}\epsilon^2 + O(\epsilon^3).
\end{aligned}$$

We divide the domain of integration into two zones, according to (3.24) and (3.25) and, accordingly, split the integral into two parts,

$$\int_0^{z-1} |K_2(r, t; \mathcal{M}; 1)| dr = \int_{(z,y) \in Z_1(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| dy + \int_{(z,y) \in Z_2(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| dy.$$

In the first zone we have (3.26), then

$$\begin{aligned}
& \frac{1}{i2\mathcal{M}(y^2-(z-1)^2)} \\
& \times \left( i(-2\mathcal{M}(y^2+z-1) + i(\ell-1)(y^2+z^2-1)) F\left(\frac{i\mathcal{M}}{1-\ell}, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2-y^2}{(z+1)^2-y^2}\right) \right.
\end{aligned}$$

$$\begin{aligned}
& +(-\ell - i\mathcal{M} + 1)(-y^2 - z^2 + 1) F\left(\frac{i\mathcal{M}}{1-\ell} - 1, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) \\
= & -\frac{1}{2} + \frac{(\mathcal{M}^2(y^2 - (z-1)^2) + i(\ell-1)\mathcal{M}(y^2 + z^2 - 1) + (\ell-1)^2(y^2 + z^2 - 1))(z-1)^2 - y^2}{2(\ell-1)^2(y^2 - (z-1)^2)} \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \\
& - \frac{\mathcal{M}(\mathcal{M} + i(\ell-1))}{8((\ell-1)^4(y^2 - (z-1)^2))} \\
& \times (\mathcal{M}^2(y^2 - (z-1)^2) + i(\ell-1)\mathcal{M}(3y^2 + z^2 + 2z - 3) + 2(\ell-1)^2(y^2 + z^2 - 1)) \\
& \times \left(\frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right)^2 + O\left(\left(\frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right)^3\right).
\end{aligned}$$

Hence

$$\begin{aligned}
& \int_{Z_1(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| dr \\
\leq & C z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_{Z_1(\epsilon, z)} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \right. \\
& \times \left[ -\frac{1}{2} + \frac{(\mathcal{M}^2(y^2 - (z-1)^2) + i(\ell-1)\mathcal{M}(y^2 + z^2 - 1) + (\ell-1)^2(y^2 + z^2 - 1))(z-1)^2 - y^2}{2(\ell-1)^2(y^2 - (z-1)^2)} \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right. \\
& - \frac{\mathcal{M}(\mathcal{M} + i(\ell-1))}{8((\ell-1)^4(y^2 - (z-1)^2))} \\
& \times (\mathcal{M}^2(y^2 - (z-1)^2) + i(\ell-1)\mathcal{M}(3y^2 + z^2 + 2z - 3) + 2(\ell-1)^2(y^2 + z^2 - 1)) \\
& \left. \left. \times \left( \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right)^2 + O\left( \left( \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right)^3 \right) \right] \right| dy.
\end{aligned}$$

We split integral into three parts and consider the first one. Since  $i\frac{\mathcal{M}}{1-\ell} < 1$ , we obtain

$$\begin{aligned}
A_0 & := z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_{Z_1(\epsilon, z)} \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} dy \\
& \lesssim z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} dy \\
& = z^{-1+2i\frac{\mathcal{M}}{1-\ell}} (z-1)(z+1)^{-2i\frac{\mathcal{M}}{1-\ell}} F\left(\frac{1}{2}, i\frac{\mathcal{M}}{1-\ell}; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2}\right) \\
& \lesssim 1 + \left( \frac{4z}{(z+1)^2} \right)^{1+i\frac{\mathcal{M}}{1-\ell}}.
\end{aligned}$$

Hence  $A_0 \lesssim 1$  for all  $z \in [1, \infty)$ . Similarly,

$$\begin{aligned}
A_1 & := z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_{Z_1(\epsilon, z)} \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \right. \\
& \times \left[ \frac{(\mathcal{M}^2(y^2 - (z-1)^2) + i(\ell-1)\mathcal{M}(y^2 + z^2 - 1) + (\ell-1)^2(y^2 + z^2 - 1))}{2(\ell-1)^2(y^2 - (z-1)^2)} \right] \Big| dy \\
& \lesssim z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \frac{z^2}{(y^2 - (z-1)^2)} \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right| dy \\
& \lesssim z^{1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left( (z+1)^2 - y^2 \right)^{-1-i\frac{\mathcal{M}}{1-\ell}} dy \\
& \lesssim F\left(\frac{1}{2}, 1 + i\frac{\mathcal{M}}{1-\ell}; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2}\right).
\end{aligned}$$



According to Lemma 4.1 we have

$$A_1 \lesssim \begin{cases} z^{\frac{1}{2} - \frac{1}{2(\ell-1)} \sqrt{(1-3\ell)^2 - 4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1). \end{cases}$$

Similarly,

$$\begin{aligned} A_2 &:= z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_{Z_1(\epsilon, z)} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \right. \\ &\quad \times \left[ -\frac{\mathcal{M}(\mathcal{M} + i(\ell-1))}{8((\ell-1)^4(y^2 - (z-1)^2))} \right. \\ &\quad \times \left( \mathcal{M}^2(y^2 - (z-1)^2) + i(\ell-1)\mathcal{M}(3y^2 + z^2 + 2z - 3) + 2(\ell-1)^2(y^2 + z^2 - 1) \right) \\ &\quad \left. \left. \times \left( \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right)^2 \right] \right| dy \\ &\lesssim z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \left[ \frac{1}{(y^2 - (z-1)^2)} z^2 \left( \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right)^2 \right] \right| dy \\ &\lesssim z^{1+2i\frac{\mathcal{M}}{1-\ell}} (z-1)(z+1)^{2(-1-i\frac{\mathcal{M}}{1-\ell})} F\left(\frac{1}{2}, 1 + i\frac{\mathcal{M}}{1-\ell}; \frac{3}{2}; \frac{(z-1)^2}{(z+1)^2}\right) \\ &\lesssim \begin{cases} z^{\frac{1}{2} - \frac{1}{2(\ell-1)} \sqrt{(1-3\ell)^2 - 4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1). \end{cases} \end{aligned}$$

The last term is defined by

$$A_3 := z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_{Z_1(\epsilon, z)} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \left[ O\left(\left(\frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right)^3\right) \right] \right| dy$$

and estimated as follows:

$$\begin{aligned} A_3 &\lesssim z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} dy \\ &\lesssim \begin{cases} z^{\frac{1}{2} - \frac{1}{2(\ell-1)} \sqrt{(1-3\ell)^2 - 4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1). \end{cases} \end{aligned}$$

In  $Z_2(\epsilon, z)$  we have (3.27) and

$$\begin{aligned} \left| F\left(\frac{i\mathcal{M}}{1-\ell}, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) \right| &\leq C \quad \text{for all } z \in [1, \infty), \quad y \in [0, z-1], \\ \left| F\left(\frac{i\mathcal{M}}{1-\ell} - 1, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) \right| &\leq C \quad \text{for all } z \in [1, \infty), \quad y \in [0, z-1], \end{aligned}$$

then

$$\begin{aligned} &\int_{Z_2(\epsilon, z)} |K_2(r, t; \mathcal{M}; 1)| dr \\ &\lesssim z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_{Z_2(\epsilon, z)} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \left[ \frac{1}{i2\mathcal{M}(y^2 - (z-1)^2)} \right. \right. \\ &\quad \left. \left. \times \left( i(-2\mathcal{M}(y^2 + z - 1) + i(\ell-1)(y^2 + z^2 - 1)) F\left(\frac{i\mathcal{M}}{1-\ell}, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2}\right) \right) \right] \right| dy \end{aligned}$$

$$\begin{aligned}
& +(-\ell - i\mathcal{M} + 1)(-y^2 - z^2 + 1) F \left( \frac{i\mathcal{M}}{1-\ell} - 1, \frac{i\mathcal{M}}{1-\ell}; 1; \frac{(z-1)^2 - y^2}{(z+1)^2 - y^2} \right) \Bigg] \Bigg| dy \\
& \lesssim z^{-1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left| \left( (z+1)^2 - y^2 \right)^{-i\frac{\mathcal{M}}{1-\ell}} \left[ \frac{1}{i2\mathcal{M}(y^2 - (z-1)^2)} z^2 \right] \right| dy \\
& \lesssim \varepsilon^{-1} z^{1+2i\frac{\mathcal{M}}{1-\ell}} \int_0^{z-1} \left( (z+1)^2 - y^2 \right)^{-1-i\frac{\mathcal{M}}{1-\ell}} dy \\
& \lesssim \begin{cases} z^{\frac{1}{2} - \frac{1}{2(\ell-1)} \sqrt{(1-3\ell)^2 - 4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1), \end{cases}
\end{aligned}$$

for all  $z \in [1, \infty)$ . The lemma is proved.  $\square$

## 4.2 Problem without potential and nonlinear term. Proof of Theorem 1.2

**Proposition 4.5** *Let  $\ell > 1$  and  $\gamma \in \mathbb{R}$ . Then the solution  $\psi_{ID}(x, t)$  of the linear equation (3.28) that takes initial values (1.7) satisfies the estimate*

$$\begin{aligned}
t^\gamma \|\psi_{ID}(x, t)\|_{H(s)} & \leq Ct^{\gamma+k_+} (\|\psi_0\|_{H(s)} + \|\psi_1\|_{H(s)}) + Ct^{\gamma-\ell} \|\psi_0\|_{H(s)} \\
& \quad + Ct^{\gamma+k_+} \|\psi_0\|_{H(s)} \begin{cases} t^{\frac{\ell-1}{2} - \frac{1}{2} \sqrt{(1-3\ell)^2 - 4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1) \end{cases}
\end{aligned}$$

for all  $t \in [1, \infty)$ .

**Proof.** The solution, after the partial Liouville transformation, can be written as follows:

$$\begin{aligned}
\psi_{ID}(x, t) & = t^{k_+} \int_0^{\phi(t)-\phi(1)} K_1(r, t; \mathcal{M}; 1) v_{u_1}(x, r) dr + t^{k_+} \left( \frac{\phi(t)}{\phi(1)} \right)^{-i\frac{\mathcal{M}}{1-\ell}} v_{u_0}(x, \phi(t) - \phi(1)) \\
& \quad + t^{k_+} \frac{1}{\phi(1)} \int_0^{\phi(t)-\phi(1)} K_2(r, t; \mathcal{M}; 1) v_{u_0}(x, r) dr.
\end{aligned}$$

Then, by (2.18), we obtain

$$\begin{aligned}
\|\psi_{ID}(x, t)\|_{H(s)} & \lesssim t^{k_+} \|u_1\|_{H(s)} \int_0^{\phi(t)-\phi(1)} |K_1(r, t; \mathcal{M}; 1)| dr + t^{k_+} \left( \frac{\phi(t)}{\phi(1)} \right)^{-i\frac{\mathcal{M}}{1-\ell}} \|u_0\|_{H(s)} \\
& \quad + t^{k_+} \|u_0\|_{H(s)} \int_0^{\phi(t)-\phi(1)} |K_2(r, t; \mathcal{M}; 1)| dr \\
& \lesssim t^{k_+} \|u_1\|_{H(s)} + t^{-\ell} \|u_0\|_{H(s)} \\
& \quad + t^{k_+} \|u_0\|_{H(s)} \begin{cases} t^{\frac{\ell-1}{2} - \frac{1}{2} \sqrt{(1-3\ell)^2 - 4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1). \end{cases}
\end{aligned}$$

Here  $u_0(x, t) = \psi_0(x, t)$ ,  $u_1(x, t) = \psi_1(x, t) - k_+ \psi_0(x, t)$ . Hence, the proposition is proved.  $\square$

## 4.3 Problem with vanishing initial conditions and without potential. Estimate of nonlinear term. Proof of Theorem 1.2

In this subsection, we obtain a decay estimates of the solution of the linear equation

$$\frac{\partial^2 \psi}{\partial t^2} + \frac{3\ell}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}(x, \partial_x) \psi + t^{-2} \frac{m^2 c^4}{h^2} \psi = \Psi.$$

For  $k_+ = -\frac{1}{2}(3\ell - 1 - \sqrt{(3\ell - 1)^2 - 4m_c^2})$ , we define operator  $G$  by

$$G[F] := t^{k_+} \int_1^t b^{-k_+} db \int_0^{\phi(t)-\phi(b)} E(r, t; b; \mathcal{M}) \mathcal{E}\mathcal{E}[F](x, r; b) dr.$$

Hence, the operator  $G$  gives a solution to the problem

$$\begin{cases} \frac{\partial^2 \psi}{\partial t^2} + \frac{3\ell}{t} \frac{\partial \psi}{\partial t} - t^{-2\ell} \mathcal{A}_{3/2}(x, \partial_x) \psi + t^{-2} m_c^2 \psi = F, \\ \psi(x, 1) = 0, \quad \psi_t(x, 1) = 0. \end{cases}$$

**Theorem 4.6** *Assume that  $\ell > 1$ . Then for the operator  $G$  the following estimate holds*

$$\begin{aligned} t^\gamma \|G[\Psi(\Phi)]\|_{H(s)} &\leq C_{NT} \left( \max_{b \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right)^{1+\alpha} \\ &\times \begin{cases} t^{-\gamma\alpha} & \text{if } \gamma < \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)}, \\ t^{\gamma+\frac{1}{2}}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2}) & \text{if } \gamma > \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)}, \\ t^{\gamma+\frac{1}{2}}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2}) \ln(t) & \text{if } \gamma = \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)} \end{cases} \end{aligned}$$

for all  $t \in [1, \infty)$ .

**Proof.** Indeed, from (2.15), (2.18), and Lemma 4.2 we obtain

$$\begin{aligned} \|G[F]\|_{H(s)} &\lesssim t^{k_+} \int_1^t b^{-k_+} \|F(x, b)\|_{H(s)} db \int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \\ &\lesssim t^{k_+} \int_1^t b^{-k_+} \|F(x, b)\|_{H(s)} |\phi(b)|^{\frac{1}{\ell-1} - \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}} |\phi(b) + \phi(t)|^{\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}} db \\ &\lesssim t^{k_+} \int_1^t b^{-k_+-1} \|F(x, b)\|_{H(s)} db. \end{aligned}$$

Further,

$$\begin{aligned} t^\gamma \|G[\Psi(\Phi)]\|_{H(s)} &\lesssim t^{\gamma+k_+} \int_1^t b^{-k_+-1} (\|\Phi(x, b)\|_{H(s)})^{1+\alpha} db \\ &\lesssim \left( \max_{b \in [1, t]} b^\gamma \|\Phi(x, b)\|_{H(s)} \right)^{1+\alpha} I(t), \end{aligned}$$

where with  $-k_+ - \gamma(1+\alpha) \neq 0$  we obtain

$$\begin{aligned} I(t) &:= t^{\gamma+k_+} \int_1^t b^{-k_+-1-\gamma(1+\alpha)} db \\ &\lesssim \begin{cases} t^{-\gamma\alpha} & \text{if } \gamma < \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)}, \\ t^{\gamma+\frac{1}{2}}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2}) & \text{if } \gamma > \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)}. \end{cases} \end{aligned}$$

For  $-k_+ - \gamma(1+\alpha) = 0$  the estimate  $I(t) \lesssim t^{\gamma+k_+} \ln(t)$  is evident. The theorem is proved.  $\square$

#### 4.4 Estimate of equation with potential and with vanishing initial data. Proof of Theorem 1.2

**Theorem 4.7** *Assume that with  $\delta \geq 0$  the potential  $V$  satisfies the condition (1.6). Then*

$$\begin{aligned} t^\gamma \|G[V\Phi]\|_{H(s)} &\leq C_{EP} \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \\ &\times \begin{cases} t^{-\delta} & \text{if } \gamma < \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta, \\ t^{\gamma+\frac{1}{2}}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2}) & \text{if } \gamma > \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta, \\ t^{-\delta} \ln(t) & \text{if } \gamma = \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta \end{cases} \end{aligned}$$

for all  $t \in [1, \infty)$ .

**Proof.** Indeed, from (2.15), (2.18), and Lemma 4.2 we obtain

$$\begin{aligned}
t^\gamma \|G[V\Phi]\|_{H(s)} &\leq t^{\gamma+k_+} \int_1^t b^{-k_+} db \int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| \|\mathcal{E}\mathcal{E}[V\Phi](x, r; b)\|_{H(s)} dr \\
&\leq t^{\gamma+k_+} \int_1^t b^{-k_+} \|V\Phi(x, b)\|_{H(s)} db \int_0^{\phi(t)-\phi(b)} |E(r, t; b; \mathcal{M})| dr \\
&\leq C_{\varepsilon_P} t^{\gamma+k_+} \int_1^t b^{-k_+-\delta} \|\Phi(x, b)\|_{H(s)} |\phi(b)^{\frac{1}{\ell-1} - \frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}} (\phi(b) + \phi(t))^{\frac{\sqrt{(1-3\ell)^2-4m_c^2}}{\ell-1}}| db \\
&\leq C_{\varepsilon_P} t^{\gamma+k_+} \int_1^t b^{-k_+-1-\delta} \|\Phi(x, b)\|_{H(s)} db \\
&\leq C_{\varepsilon_P} \left( \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right) J(t),
\end{aligned}$$

where  $J(t) := t^{\gamma+k_+} \int_1^t b^{-k_+-1-\delta-\gamma} db$ . With  $-k_+ - \delta - \gamma \neq 0$  we obtain

$$J(t) \lesssim \begin{cases} t^{-\delta} & \text{if } \gamma < \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta, \\ t^{\gamma+\frac{1}{2}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2})} & \text{if } \gamma > \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta. \end{cases}$$

If  $-k_+ - \gamma - \delta = 0$ , then  $J(t) = t^{-\delta} \ln(t)$ . The theorem is proved.  $\square$

#### 4.5 Existence of solution to semilinear equation. Proof of Theorem 1.2

We solve the Cauchy problem (1.4)&(1.7) through the integral equation. We appeal to the operator (2.15),  $G := \mathcal{K} \circ \mathcal{E}\mathcal{E}$ , where  $\mathcal{E}\mathcal{E}$  stands for the evolution (wave) equation in the exterior of BH in the universe without expansion, while  $\mathcal{K}$  is introduced by (2.17) with  $k_+$ ,  $\mathcal{M}_+$ . The kernel  $E(r, t; 0; b; M)$  is given by (2.11). For the function  $f(x, t)$  we define  $v(x, t; b) := \mathcal{E}\mathcal{E}[f](x, t; b)$ , where the function  $v(x, t; b)$  is a solution to the Cauchy problem (2.16).

We intend to apply Banach's fixed-point theorem. We use the Lipschitz condition ( $\mathcal{L}$ ) to estimate nonlinear term. We start with the integral equation (3.30), where the function  $\psi_{ID}(x, t) \in C([0, \infty); H(s))$  is given.

Consider the mapping  $S[\Phi]$  (3.31), where the function  $\psi_{ID}$  is generated by initial data, that is, by (1.7). It is easily seen that  $\text{supp } S[\Phi] \subset \text{supp } \Phi$  if  $\text{supp } \Phi \subset \text{supp } \psi_{ID}$ . We claim that if  $\Phi \in X(R, H(s), \gamma)$  with  $\gamma \in [0, H]$  and if  $\text{supp } \Phi \subseteq \{(x, t) \in \mathbb{R}^3 \times [0, \infty) \mid |x| > R_{ID} - (\phi(t) - \phi(1))\}$ , then  $S[\Phi] \in X(R, H(s), \gamma)$ . Moreover,  $S$  is a contraction, provided that  $\varepsilon$ ,  $\varepsilon_P$ , and  $R$  are sufficiently small.

First, we note that due to Proposition 4.5, we have

$$\begin{aligned}
t^\gamma \|\psi_{ID}(x, t)\|_{H(s)} &\leq C_{ID} t^{\gamma+k_+} (\|\psi_0\|_{H(s)} + \|\psi_1\|_{H(s)}) + C_{ID} t^{\gamma-\ell} \|\psi_0\|_{H(s)} \\
&\quad + C_{ID} t^{\gamma+k_+} \|\psi_0\|_{H(s)} \begin{cases} t^{\frac{\ell-1}{2} - \frac{1}{2}\sqrt{(1-3\ell)^2-4m_c^2}} & \text{if } 4m_c^2 > 4\ell(2\ell-1), \\ 1 & \text{if } 4m_c^2 \leq 4\ell(2\ell-1). \end{cases}
\end{aligned}$$

According to Theorem 4.7

$$\begin{aligned}
t^\gamma \|G[V\Phi]\|_{H(s)} &\leq C_{\varepsilon_P} \sup_{\tau \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \\
&\quad \times \begin{cases} t^{-\delta} & \text{if } \gamma < \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta, \\ t^{\gamma+\frac{1}{2}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2})} & \text{if } \gamma > \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta, \\ t^{-\delta} \ln(t) & \text{if } \gamma = \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2} - \delta, \end{cases}
\end{aligned}$$

while due to Theorem 4.6

$$t^\gamma \|G[\Psi(\Phi)]\|_{H(s)} \leq C_{NT} \left( \max_{b \in [1, t]} \tau^\gamma \|\Phi(x, \tau)\|_{H(s)} \right)^{1+\alpha}$$

$$\times \begin{cases} t^{-\gamma\alpha} & \text{if } \gamma < \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)}, \\ t^{\gamma+\frac{1}{2}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2})} & \text{if } \gamma > \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)}, \\ t^{\gamma+\frac{1}{2}(1-3\ell+\sqrt{(1-3\ell)^2-4m_c^2})} \ln(t) & \text{if } \gamma = \frac{3\ell-1-\sqrt{(1-3\ell)^2-4m_c^2}}{2(1+\alpha)}. \end{cases}$$

All time-dependent factors are bounded at all times due to the conditions of the theorem for  $\gamma$ . Thus, the last inequalities prove that the operator  $S$  maps  $X(R, s, \gamma)$  into itself if  $\|\psi_0\|_{H(s)} + \|\psi_1\|_{H(s)} \leq \varepsilon$ , while  $\varepsilon_P$ ,  $\varepsilon$ , and  $R$  are sufficiently small, namely, provided that  $\frac{1}{1-\varepsilon_P C_P} C_{ID} \varepsilon + \frac{1}{1-\varepsilon_P C_P} C_{NT} R^{\alpha+1} < R$ . To prove that  $S$  is a contraction mapping, we derive the contraction property from

$$\sup_{t \in [0, \infty)} t^\gamma \|S[\Phi_1](\cdot, t) - S[\Phi_2](\cdot, t)\|_{H(s)(\mathbb{R}^n)} \leq CR^\alpha d(\Phi_1, \Phi_2).$$

The theorem is proved. □

## Acknowledgments

The authors are indebted to the referees for the valuable remarks and suggestions, which improved the text.

## References

- [1] H. S. Aslan, M. R. Ebert and M. Reissig, *Scale-invariant semilinear damped wave models with mass term and integrable in time speed of propagation*, Differential and Integral Equations, 36 (2023), 453–490.
- [2] H. Bateman and A. Erdelyi, “Higher Transcendental Functions”, Vol. 1, 2, McGraw-Hill, 1954.
- [3] S. Chen, G. W. Gibbons, Y. Li and Y. Yang, *Friedmann’s equations in all dimensions and Chebyshev’s theorem*, J. Cosmol. Astropart. Phys., 12 (2014), 035.
- [4] J. L. Costa, A. T. Franzen and J. Oliver, *Semilinear wave equations on accelerated expanding FLRW spacetimes*, Ann. Henri Poincaré, 24 (2023), 3185-3207.
- [5] D. Farrah et al., *Observational evidence for cosmological coupling of black holes and its implications for an astrophysical source of dark energy*, The Astrophysical Journal Letters, 944 (2023), L31.
- [6] A. Galstian and K. Yagdjian, *Global solutions for semilinear Klein-Gordon equations in FLRW spacetimes*, Nonlinear Anal., 113 (2015), 339-356.
- [7] M. Nakamura, *The Cauchy problem for semi-linear Klein-Gordon equations in de Sitter spacetime*, J. Math. Anal. Appl., 410 (2014), 445-454.
- [8] J. Natário and F. Rossetti, *Explicit formulas and decay rates for the solution of the wave equation in cosmological spacetimes*, J. Math. Phys., 64 (2023), 032504.
- [9] J. Natário and A. Sasane, *Decay of solutions to the Klein-Gordon equation on some expanding cosmological spacetimes*, Ann. Henri Poincaré, 23 (2022), 2345–2389.
- [10] A. Palmieri, *Integral representation formulae for the solution of a wave equation with time-dependent damping and mass in the scale-invariant case*, Math. Methods Appl. Sci., 44 (2021), 13008-13039.
- [11] V. Perlick, O. Y. Tsupko and G. S. Bisnovatyi-Kogan, *Black hole shadow in an expanding universe with a cosmological constant*, Phys. Rev. D, 97 (2018), 104062.
- [12] M. Taylor, “Partial Differential Equations III. Nonlinear Equations”, 2nd edition, Applied Mathematical Sciences 117, Springer, New York, 2011.
- [13] K. Yagdjian, “The Cauchy Problem for Hyperbolic Operators. Multiple Characteristics. Micro-Local Approach”, Mathematical Topics, 12, Akademie Verlag, Berlin, 1997.

- [14] K. Yagdjian and A. Galstian, *Fundamental solutions for the Klein-Gordon equation in de Sitter spacetime*, Comm. Math. Phys., 285 (2009), 293–344.
- [15] K. Yagdjian, *Integral transform approach to time-dependent partial differential equations*, Mathematical analysis, probability and applications—plenary lectures, Springer Proc. Math. Stat., Vol. 177, Springer, (2016), 281–336.
- [16] K. Yagdjian, *Global existence of the self-interacting scalar field in the de Sitter universe*, J. Math. Phys., 60 (2019), 051503.
- [17] K. Yagdjian and A. Galstian, *Fundamental solutions for the Dirac equation in curved spacetime and generalized Euler–Poisson–Darboux equation*, J. Differential Equations, 300 (2021), 80–117.
- [18] K. Yagdjian, *Waves in cosmological background with static Schwarzschild radius in the expanding universe*, J. Math. Phys., 65, (2024), 083503.