

Article

# Look beyond additivity and extensivity of entropy for black hole and cosmological horizons.

Mariusz P. Dąbrowski <sup>1,2,3</sup>

<sup>1</sup> Institute of Physics, University of Szczecin, Wielkopolska 15, 70-451 Szczecin, Poland; Mariusz.Dabrowski@usz.edu.pl

<sup>2</sup> National Centre for Nuclear Research, Andrzeja Sołtana 7, 05-400 Otwock, Poland;

<sup>3</sup> Copernicus Center for Interdisciplinary Studies, Szczepańska 1/5, 31-011 Kraków, Poland

Academic Editors: Stephen Xsu, Xavier Calmet, Roberto Casadio

Version September 30, 2024 submitted to Journal Not Specified

**Abstract:** We present a comparative analysis of the plethora of nonextensive and/or nonadditive entropies which go beyond the standard Boltzmann-Gibbs formulation. After defining the basic notions of additivity, extensivity, and composability, we discuss the properties of these entropies and their mutual relations, if they exist. The results are presented in two informative tables supposedly of strong interest to gravity and cosmology community in the context of intensively explored recent days the horizon entropies for black hole and cosmological models. This is since gravitational systems admit long-range interactions which usually lead to a break of the standard additivity rule for thermodynamical systems composed of subsystems in Boltzmann-Gibbs thermodynamics. The features of additivity, extensivity, and the composability are listed systematically. Some brief discussion on the validity of the notion of equilibrium temperature for nonextensive systems is also presented.

**Keywords:** thermodynamics, entropy, additivity, extensivity, long-range interactions, horizons.

## 1. Introduction

It is widely known that the Boltzmann-Gibbs thermodynamics (from now on BG) and statistical mechanics are additive and extensive [1]. The core physical quantity which refers to these theories is the entropy, which is assumed to be extensive, since it relates to the negligence of the long-range forces between thermodynamic subsystems. This assumption is justified only, when the size of the system exceeds the range of the interaction between its components. As a result, the total entropy of a composite system is equal to the sum of the entropies of the individual subsystems (additivity) and the entropy grows with the size of the system or its configuration space (extensivity).

However, contemporary physics exhibits a number of systems for which the long-range forces are important. The very examples of such systems are gravitational systems since gravity is long-range interactive, and besides, it is strongly non-linear when its extreme regimes are taken into account. Strong gravity characterises all the compact astrophysical objects in the Universe like white dwarfs, neutron and boson stars, quark stars etc., with the most extreme and most intriguing - the black holes. The latter are surrounded by the horizons which areas, according to Bekenstein and Hawking [2,3], can be interpreted as the entropy, and so it is possible to formulate the appropriate laws of thermodynamics. Since for the black holes the Bekenstein entropy *scales with the area* and not with the volume (size), it is consequently a nonextensive quantity [4–9]. In addition to that, because of a long-range interaction nature of gravity, the Bekenstein entropy is also nonadditive.

In fact, a number of nonadditive and/or nonextensive entropies have been proposed in the literature [10–16]. Most of them have been applied to gravitational systems both in astrophysics and

in cosmology and there is a debate, if they can serve dark energy [17–34], which is specifically called the *holographic dark energy* [35]. Amazingly, a number of these explorations do not acknowledge the fact of nonextensivity of the systems, which makes some issues related to their firm thermodynamical background [36]. All this motivates us to investigate the problem of nonextensive entropies applications in some gravitational framework.

In this paper, we explore the topic of additivity and extensivity of entropies which goes beyond the standard BG thermodynamics, being strongly motivated by gravitational interaction. Our focus is put on non-standard, but better fitting to gravitational systems entropies, such as: Bekenstein entropy [2,3], Tsallis  $q$ -entropy [10,37], Tsallis-Cirto  $\delta$ -entropy [4], Barrow  $\Delta$ -entropy [16], Tsallis  $q, \delta$  entropy, Tsallis-Jensen  $q, \gamma$ -entropy [38], Rényi entropy [11], Landsberg  $U$ -entropy [39], Sharma-Mittal entropy [12,13], and Kaniadakis entropy [14,15].

The following is the outline of the paper. In Sec. 2, we define additivity and extensivity in thermodynamical systems and try to establish some generalities about possible composition rules for the entropy. In Sec. 3 we go beyond the definitions of additivity and extensivity. In Sec. 4 we constructively review and compare the plethora of nonadditive and/or nonextensive entropies together with accompanying nonadditive and/or nonextensive thermodynamical quantities. Then, we make the classification of the entropies under study with respect to additivity and extensivity properties as well as with the application of the appropriate composition rules. Finally, in Sec. 5, we summarize the paper.

## 2. Boltzmann-Gibbs thermodynamics and statistical mechanics

Boltzmann-Gibbs thermodynamics and statistical mechanics are based on two key hypotheses which are that the entropy is extensive and that the internal energy and entropy follow the additive composition rule for the system made of some subsystems. All physical relations in BG statistical mechanics are defined in light of these conditions which, in fact, rely on ignoring long-range forces between thermodynamical subsystems.

The Boltzmann-Gibbs (BG) entropy is defined as [1]

$$S_{BG} = -k_B \sum_{i=1}^n p_i \ln p_i = k_B \sum_{i=1}^n p_i \ln \frac{1}{p_i}, \quad (1)$$

where  $p_i$  is the probability distribution defined on a configuration space  $\Omega$  with the number of degrees of freedom (states)  $n$ ,  $k_B$  is the Boltzmann constant, and the condition that the total probability must be equal to one  $\sum p_i = 1$  is fulfilled. For the case of all probabilities equal, i.e. for  $p_i = \text{const.} = p$ , we get

$$\sum_{i=1}^n p_i = 1 = np \Rightarrow p = 1/n. \quad (2)$$

After applying (2) to (1), one obtains that

$$S_{BG} = k_B \ln n, \quad (3)$$

which means that the entropy is proportional to the number of states  $n$  in the configuration space  $\Omega$ .

In view of the key properties of BG thermodynamics, and in the context of our investigations beyond these properties, we can define additivity and extensivity in quite a general way following some literature [39–41] as below.

### 2.1. Additivity

Additivity means that for a given physical or thermodynamical quantity  $f$ , the following composition rule is fulfilled:

$$f(A + B) = f(p_{A \cup B}) = f(p_A p_B) = f(p_A) + f(p_B) = f(A) + f(B), \quad (4)$$

where  $A, B$  are independent subsystems, equipped with the sets of configuration space degrees of freedom  $\Omega_A$  and  $\Omega_B$ , and corresponding probabilities  $p_A$  and  $p_B$ . The composite system  $A \cup B$  allows the probability  $p_{A \cup B}$  and it is equipped with the set of configuration space degrees of freedom  $\Omega_{A \cup B}$ . If the subsystems  $A$  and  $B$  are assumed to be independent, then it happens that the probabilities are related by  $p_{A \cup B} = p_A p_B$ , which allows the transition leading to the additivity rule (4) [42].

If a particular case of the entropy  $S$  is taken into account, then (4) reads

$$S(A + B) = S(A) + S(B). \quad (5)$$

### 2.2. Extensivity

Let us assume that there is a set of physical quantities  $(X_0, X_1, X_2, \dots, X_k)$  such that  $X_0 = f(X_1, X_2, \dots, X_k)$ . Extensivity of a selected physical quantity means that the function  $f$  which describes this quantity is *homogeneous degree one* [1,39,40] i.e. that

$$f(aX_1, aX_2, \dots, aX_k) = af(X_1, X_2, \dots, X_k) \quad (6)$$

for every positive real number  $a > 0$ , for all  $X_1, X_2, \dots, X_k$ . Taking  $k = 3$ , so that we have only four quantities  $X_0, X_1, X_2, X_3$ , and assuming that they are the entropy  $S$ , the energy  $E$ , the volume  $V$ , and the mole number  $N$  accordingly, we can obtain the standard Boltzmann-Gibbs thermodynamical extensivity relation for the entropy [39]

$$S(aE, aV, aN) = aS(E, V, N). \quad (7)$$

In fact, the property (6) is called 'homogeneity', and is considered the most general definition of extensivity (cf. [39]).

In standard textbooks of thermodynamics, one commonly uses less general definition of an extensive quantity, which says that if a system's total number of states in the configuration space  $\Omega$  is proportional to its number of degrees of freedom, then this quantity (such as the entropy, for example) is extensive. For BG entropy, as we have shown in (3), one has that  $S_{BG}(n) = k_B \ln(n) \propto n$ , where  $n$  is the total number of states in the system.

The advantage of definition (6) is that one does not refer to any kind of geometrical or bulk properties of a system such as the 'size', though the geometrical size of a system seems intuitively to be related to the number of states or degrees of freedom.

### 2.3. Concavity

Concavity is the feature of the functions which read as [4,39]

$$f(ax + (1 - a)y) \geq af(x) + (1 - a)f(y) \quad (a > 0). \quad (8)$$

In the context of thermodynamics, concavity of entropy guarantees that the system in thermodynamic equilibrium is *stable*.

### 3. Beyond Boltzmann-Gibbs thermodynamics

#### 3.1. Composability

Let us consider two independent systems  $A$  and  $B$ , combined as a single Cartesian product  $A \times B$  of the states of  $A$  and  $B$  with the requirement that [38]:

$$S(A \times B, Y) = k_B g \left( \frac{S(A)}{k_B}, \frac{S(B)}{k_B} \right), \quad (9)$$

where  $g$  is a smooth function of  $S(A)$  and  $S(B)$  and  $Y$  is a parameter, which in the limit  $Y \rightarrow 0$  give an additive composition rule (5). If the systems  $A$  and  $B$  fulfil the condition (9), then their combined system  $A + B$  is called *composable*. Of course, the BG system is composable.

#### 3.2. Beyond additivity

Additivity is violated, if the rule (5) does not hold. In such a case, one can have two options [39]. The first one is when

$$S(A + B) \geq S(A) + S(B), \quad (10)$$

which is called *superadditivity*, and it leads to the tendency of the system to clump its pieces/subsystems. The second one is when

$$S(A + B) < S(A) + S(B), \quad (11)$$

which is *subadditivity*, and it tends to fragment the system into pieces, rather than clump. A cosmological similarity of such a system is phantom [43,44], since it splits spontaneously into pieces under (anti)gravity beginning with the largest size pieces and terminating at the smallest [45].

In the literature, there are a number composition rules for nonadditive systems, which we introduce in Section 4. One of them, which generalizes the additive composition rule (5) into a nonadditive case, is the Abé rule [46–48]. It fulfils the composability requirement given by (9). If applied to entropy, it reads as follows

$$S(A + B) = S(A) + S(B) + \frac{Y}{k_B} S(A)S(B), \quad (12)$$

where  $Y$  takes numerical values according to the statistical definition of a specific entropy type. For BG entropy, one just has  $Y = 0$ . With the assumption that all the entropies in (12) are positive, one deals with superadditivity for  $Y \geq 0$ , and with subadditivity for  $Y < 0$ . In fact, the physical interpretation of  $Y$  is the result of the long-range interactions between subsystems, which leads to nonadditivity.

#### 3.3. Beyond extensivity

In BG thermodynamics the additivity and the extensivity are closely related - additivity implies extensivity and extensivity implies additivity [1]. This is not the case in general, and so the extensivity and the additivity may not be related, i.e. the extensivity may not imply the additivity and vice versa. An example of such a kind of a quantity, which is based on the definition (6) is given by the function  $f(X_1, X_2) = x_1^2 / \sqrt{X_1^2 + X_2^2}$ . It obeys extensivity, but not additivity [39].

Generally, the entropy  $S$  is nonextensive if

$$S(aX) \neq aS(X), \quad (13)$$

where  $X$  is a thermodynamical quantity and  $a > 0$ , i.e. when the relations (6) and (7) are violated.

## 4. A comparable analysis of nonextensive entropies plethora

### 4.1. Bekenstein entropy

The Bekenstein entropy is not motivated by anything like statistical mechanics, but it is a well established notion in gravity theory [2]. For a Schwarzschild black hole it reads

$$S_{Bek} = 4\pi k_B \left( \frac{M}{m_p} \right)^2 = \frac{4\pi k_B G M^2}{\hbar c}, \quad (14)$$

and it is usually presented with its accompanying Hawking temperature, which reads

$$T_H = \frac{\hbar c^3}{8\pi G k_B M}, \quad (15)$$

where  $M$  is the mass of a black hole,  $c$  is the speed of light,  $G$  is the gravitational constant,  $\hbar$  is the reduced Planck constant, and  $m_p$  is the Planck mass. In fact, the temperature (15) can be calculated from the entropy (14), by applying the Clausius formula

$$\frac{k_B}{T} = \frac{\partial S}{\partial E}, \quad (16)$$

and using the Einstein mass-energy equivalence formula  $E = Mc^2$ .

It is not always understood in the literature that because of the area rather than volume scaling, the Bekenstein entropy is nonextensive, and that it obeys the following nonadditive composition rule (see e.g. [49])

$$S_{A+B} = S_A + S_B + 2\sqrt{S_A}\sqrt{S_B}, \quad (17)$$

which we will call the *square root rule* from now on. This rule comes directly as a consequence of (14), according to which the entropy  $S_{Bek} \propto M^2$ , so that  $S_A \propto M_A^2$ , and  $S_B \propto M_B^2$ . If black holes merge in an adiabatic way, then their mass after merging is the sum  $M_{A+B} = M_A + M_B$ , but the entropy  $S_{A+B} \propto M_{A+B}^2$ , giving an extra term  $1/2M_A M_B$ , which is an extra nonadditive term in (17).

Curiously, after making a redefinition of the Bekenstein entropies as  $\tilde{S}_{A+B} \equiv \sqrt{S_{A+B}} = M_{A+B} = M_A + M_B$ ,  $\sqrt{S_A} = M_A \equiv \tilde{S}_A$ , and  $\sqrt{S_B} = M_B \equiv \tilde{S}_B$ , one can rewrite the composition rule (17) in an additive way

$$\tilde{S}_{A+B} = \tilde{S}_A + \tilde{S}_B, \quad (18)$$

but this is not of any physical meaning.

In conclusion, Bekenstein entropy addition formula (17) does not fulfil Abé rule (12), though it looks quite similar. We comment on this point in relation to other entropies in subsection 4.3.

### 4.2. Tsallis $q$ , Tsallis-Cirto $\delta$ , Tsallis $q, \delta$ , and Tsallis-Jensen $q, \gamma$ entropies

#### 4.2.1. Tsallis $q$ -entropy

The Tsallis  $q$ -entropy [10,50] is one of the earliest proposals for generalization of BG entropy. It encompasses an issue of the long-range interaction between thermodynamical subsystems by introducing a nonextensivity parameter  $q$  ( $q \in \mathbb{R}$ ) into the BG entropy definition (1) keeping the standard BG condition that the sum of all the probabilities is equal to one, i.e. that  $\sum p_i = 1$ . It reads as follows

$$\mathcal{S}_q = k_B \sum_{i=1}^n p_i \ln_q \frac{1}{p_i} = -k_B \sum_{i=1}^n (p_i)^q \ln_q p_i = -k_B \sum_{i=1}^n \ln_{2-q} p_i, \quad (19)$$

where a newly defined  $q$ -logarithmic function  $\ln_q p$ , is introduced:

$$\ln_q p \equiv \frac{p^{1-q} - 1}{1 - q}. \quad (20)$$

In the limit  $q \rightarrow 1$ , one has the standard logarithm  $\ln_1 p = \ln p$ . It is important that the  $q$ -logarithm does not fulfil the standard logarithm addition rule  $\ln ab = \ln a + \ln b$ , where  $a, b$  are some arbitrary numbers. Instead, it fulfils a nonadditive composition rule given by

$$\ln_q ab = \ln_q a + \ln_q b + (1 - q) \ln_q a \ln_q b, \quad (21)$$

which, in fact, is the origin of the Abé rule (12). Interestingly, by the introduction of some specific  $q$ -product defined as [51]

$$(x \otimes y)_q \equiv \left[ x^{1-q} + y^{1-q} - 1 \right]^{\frac{1}{1-q}}, \quad (x \geq 0, y \geq 0), \quad (22)$$

one can make the rule (21) additive, i.e.

$$\ln_q [(x \otimes y)_q] = \ln_q x + \ln_q y. \quad (23)$$

It is also possible to define the  $q$ -exponential function

$$e_q^p \equiv [1 + (1 - q)p]^{\frac{1}{1-q}}, \quad (24)$$

which does not fulfil the standard exponent addition rule  $e^{a+b} = e^a e^b$ , though in the limit  $q \rightarrow 1$  it does, since  $e_1^p = e^p$ .

Tsallis  $q$ -entropy definition (19) is usually presented in three equivalent forms. However, using the definition of  $q$ -logarithm (20), all of them can be brought into the same form (cf. Appendix A):

$$\mathcal{S}_q = k_B \frac{1 - \sum_{i=1}^n (p_i)^q}{q - 1}. \quad (25)$$

It is important to mention that in order to fulfil the requirements of concavity for  $S_q$  given by (8), the nonextensivity parameter should be positive  $q > 0$  [4].

In the limit  $q \rightarrow 1$ , Tsallis entropy  $S_q$  given by (19) or (25) reduces to BG entropy (1). After some check, it is possible to find that Tsallis  $q$ -entropy (19) or (25) satisfies the nonadditive composition Abé rule (12), if one defines a nonextensivity parameter as  $Y = 1 - q$  (cf. Appendix B). For equal probability states (2), the formula (25) gives the Tsallis  $q$ -entropy as

$$\mathcal{S}_q = k_B \ln_q n = k_B \frac{n^{q-1} - 1}{1 - q}, \quad (26)$$

which nicely shows, how it generalizes BG entropy (3) via a new parameter  $q$ .

#### 4.2.2. Rényi entropy

The Rényi entropy [11], which is in fact a measure of entanglement in quantum information theory, is additive and preserves event independence. It is another important generalization of BG entropy, which is defined by

$$S_R = k_B \frac{\ln \sum_{i=1}^n (p_i)^q}{1 - q}. \quad (27)$$

By assuming that all the states are equally probable, as in (2), it follows from (27) that

$$S_R = k_B \ln n, \quad (28)$$

which is the same as BG entropy (1).

In fact, the Rényi entropy (27) can be written in terms of the Tsallis  $q$ -entropy by using the formal logarithm approach [52] as follows

$$S_R = \frac{k_B}{1-q} \ln \left[ 1 + \frac{1-q}{k_B} S_q \right]. \quad (29)$$

Quite a unique feature of Rényi entropy is that *it is additive*, which results from some more general Abé composition rule given by [49]

$$H(S_{A+B}) = H(S_A) + H(S_B) + \frac{Y}{k_B} H(S_A) H(S_B), \quad (30)$$

together with redefinition using the logarithm in the form

$$L(S) = \frac{k_B}{Y} \ln \left( 1 + \frac{Y}{k_B} H(S) \right), \quad (31)$$

which applied to (29) gives an additive formula

$$L(S_{A+B}) = L(S_A) + L(S_B), \quad (32)$$

where  $L(S)$  corresponds to the Rényi entropy, and  $H(S)$  corresponds to the Tsallis  $q$ -entropy. In such a formulation, one can write that the Rényi entropy fulfils the Abé rule (12), with the parameter  $Y = 0$ .

#### 4.2.3. Tsallis-Cirto $\delta$ -entropy

The Tsallis-Cirto  $\delta$ -entropy [4,9], sometimes also known in the literature as just the Tsallis entropy, is yet another generalization of BG entropy (1) by the introduction of another nonextensivity parameter  $\delta$  as follows

$$S_\delta = k_B \sum_{i=1}^n p_i (\ln p_i)^\delta \quad (\delta > 0, \delta \in \mathbb{R}), \quad (33)$$

and this difference is easily recognized, when one compares it with the Tsallis  $q$ -entropy (19), and with BG entropy (1).

Under the assumption that all the states are equally probable, as in (2), one gets from (33) that

$$S_\delta = k_B (\ln n)^\delta \equiv k_B \ln^\delta n. \quad (34)$$

Making another assumption that we deal with two independent systems  $A$  and  $B$ , fulfilling the condition  $n^{A+B} = n^A \cdot n^B$ , one realizes that the composition rule for the Tsallis-Cirto entropy (34) reads

$$\left( \frac{S_{\delta, A+B}}{k_B} \right)^{1/\delta} = \left( \frac{S_{\delta, A}}{k_B} \right)^{1/\delta} + \left( \frac{S_{\delta, B}}{k_B} \right)^{1/\delta}, \quad (35)$$

which is another example of a composition rule, which is *different* from the Abé rule (12). We start calling it  $\delta$ -addition rule from now on. In fact, Tsallis and Cirto suggest that [4,9]

$$S_\delta = k_B \left( \frac{S_{Bek}}{k_B} \right)^\delta, \quad (36)$$

where  $S_{Bek}$  is the Bekenstein entropy (14). According to a new composition rule (35), one realizes that the Bekenstein entropy, as given by  $S_{Bek} \propto (S_\delta)^{1/\delta}$ , can be additive, while the Tsallis-Cirto entropy  $S_\delta$  itself is nonadditive. Besides, bearing in mind the definition of Bekenstein entropy for a Schwarzschild black hole (14), one can easily notice that for  $\delta = 3/2$  the Tsallis-Cirto entropy (36) is proportional to the volume  $S_\delta \propto M^3$ , and so it is an extensive quantity in view of the standard definition of extensivity. We come back to this issue in subsection 4.3.

#### 4.2.4. Tsallis $q, \delta$ -entropy

The Tsallis  $q, \delta$ -entropy generalizes both the Tsallis  $q$ -entropy (19) and the Tsallis-Cirto  $\delta$ -entropy (33), combining them as follows [4,9]

$$S_{q,\delta} = k_B \sum_{i=1}^n p_i (\ln_q p_i)^\delta \quad (\delta > 0, q \in \mathbb{R}, \delta \in \mathbb{R}). \quad (37)$$

Now both  $q$  and  $\delta$  play the role of two independent nonextensivity parameters. By assuming that all the states are equally probable as in (2), one gets from (37) that

$$S_{q,\delta} = k_B (\ln_q n)^\delta \equiv k_B \ln_q^\delta n. \quad (38)$$

The Tsallis  $q, \delta$ -entropy fulfils neither Abé addition rule nor  $\delta$ -addition rule, though it does the former in the limit  $\delta \rightarrow 0$ , and the latter in the limit  $q \rightarrow 0$ .

#### 4.2.5. Tsallis-Jensen $q, \gamma$ -entropy

Recently, Tsallis and Jensen [38] proposed another generalization of BG entropy which reads

$$S_{q,\gamma} = k_B \left[ \frac{\ln \sum_{i=1}^n p_i^q}{1-q} \right]^{\frac{1}{\gamma}} = k_B \left( \frac{S_R}{k_B} \right)^{\frac{1}{\gamma}}, \quad (39)$$

where  $S_R$  is the Rényi entropy and  $\gamma$  is a new parameter somewhat analogous to the parameter  $\delta$  in the Tsallis-Cirto entropy (33).

Since the Rényi entropy possesses BG limit for  $q \rightarrow 1$ , then we can write [38]

$$S_{1,\gamma} = k_B \left( \frac{S_{BG}}{k_B} \right)^{\frac{1}{\gamma}}, \quad (40)$$

and analogously, if we take Bekenstein entropy (14) instead of the BG in (40), in the same limit  $q \rightarrow 1$ , we get

$$S_{1,\gamma}^{Bek} = k_B \left( \frac{S_{Bek}}{k_B} \right)^{\frac{1}{\gamma}}. \quad (41)$$

Bearing in mind the additive composition rule (5), for the BG entropy, and using (40), one can write the additivity rule for  $S_{1,\gamma}$  in such a case as

$$[S_{1,\gamma}(A+B)]^\gamma = [S_{1,\gamma}(A)]^\gamma + [S_{1,\gamma}(B)]^\gamma. \quad (42)$$

Similarly, taking into account the square root additivity rule (17) for the Bekenstein-like entropy (41), one can write the composition rule as

$$[S_{1,\gamma}^{Bek}(A+B)]^\gamma = [S_{1,\gamma}^{Bek}(A)]^\gamma + [S_{1,\gamma}^{Bek}(B)]^\gamma + 2 [S_{1,\gamma}^{Bek}(A)S_{1,\gamma}^{Bek}(B)]^{\frac{\gamma}{2}}. \quad (43)$$



**Table 1.** Tsallis entropies

Entropy Type	Extensivity	Additivity	Abé addition rule	$\delta$ -addition rule
Boltzmann-Gibbs $S_{BG}$	yes	yes	yes, $Y = 0$	yes, $\delta = 1$
Tsallis $S_{q,1} = S_q$	no	no	yes, $Y = 1 - q$	no
Tsallis-Cirto $S_{1,\delta} = S_\delta$	no	no	no	yes
General Tsallis $S_{q,\delta}$	no	no	no	no
Tsallis-Jensen $S_{q,\gamma}$	no	no	no	yes, $\delta = 1/\gamma$

Finally, since the Rényi entropy  $S_R$  in (39) is in general additive according to the composition rule (32), then we can write a generic composition rule for the Tsallis-Jensen entropy (39) as

$$[S_{q,\gamma}(A+B)]^\gamma = [S_{q,\gamma}(A)]^\gamma + [S_{q,\gamma}(B)]^\gamma, \quad (44)$$

and this is exactly the  $\delta$  composition rule (35) with  $\gamma = 1/\delta$ .

Table 1 gives the summary of four different Tsallis invented entropies.

#### 4.3. Barrow fractal horizon $\Delta$ -entropy and its relation to Bekenstein and Tsallis-Cirto $\delta$ -entropy

Barrow entropy [16] has no statistical roots at all. It is closely tied to black hole horizon geometry influenced by quantum fluctuations, which make initially smooth black hole horizon a fractal composed of spheres, forming the so-called sphereflake. This structure is characterised by the fractal dimension  $d_f$  which in the extreme cases are either the surface or the volume i.e.  $2 \leq d_f \leq 3$ , and results in an effective horizon area of  $r^{d_f}$ , where  $r$  is the black hole horizon radius. After quantum-motivated modification of the area, the entropy reads

$$S_{Bar} = k_B \left( \frac{A}{A_p} \right)^{1+\frac{\Delta}{2}} = k_B \left( \frac{S_{Bek}}{k_B} \right)^{1+\frac{\Delta}{2}}, \quad (45)$$

where  $S_{Bek}$  is Bekenstein entropy,  $A$  - the horizon area,  $A_p$  - the Planck area,  $A_p \propto l_p^2$  with  $l_p$  - Planck length, and  $\Delta$  is the parameter related to the fractal dimension by the relation  $\Delta = d_f - 2$ . In fact,  $0 \leq \Delta \leq 1$  with  $\Delta \rightarrow 1$  limit yielding maximally fractal structure, where the horizon area behaves effectively like a 3-dimensional volume, and with  $\Delta \rightarrow 0$  limit yielding the Bekenstein area law, where no fractalization occurs. Although Barrow entropy has geometrical roots, and is not motivated by thermodynamics, it has the same form as Tsallis-Cirto  $\delta$  entropy (36) [53] being also related to Bekenstein entropy  $S_{Bek}$ , as in (14), provided that

$$\delta = 1 + \frac{\Delta}{2}. \quad (46)$$

However, the ranges of parameters  $\delta$  and  $\Delta$  are different -  $\delta$  has only the bound  $\delta > 0$ , while  $0 \leq \Delta \leq 1$  is equivalent to  $1 \leq \delta \leq 3/2$ . Thus, qualitatively, both entropic forms yield the same temperatures as a function of a black hole mass. Both Tsallis-Cirto entropy limit  $\delta \rightarrow 3/2$  and Barrow limit  $\Delta \rightarrow 1$  yield an extensive, but still nonadditive entropy for black holes. In fact, the cosmological studies of Barrow entropy as holographic dark energy have been performed intensively [17–34], pointing towards this extensive case [18,38,54].

#### 4.4. Landsberg $U$ -entropy

The Landsberg  $U$ -entropy is defined in relation to Tsallis  $q$ -entropy (25) as [39]

$$S_U = \frac{k_B}{1-q} \left( 1 - \frac{1}{\sum_{i=1}^n (p_i)^q} \right) = k_B \frac{1 - \sum_{i=1}^n (p_i)^q}{q-1} \frac{1}{(p_i)^q} = \frac{S_q}{\sum_{i=1}^n (p_i)^q}, \quad (47)$$

and it fulfils the Abé rule (12) for  $Y = q - 1$  (cf. Appendix B). By assuming that all the states are equally probable as in (2), it simplifies (47) to the form

$$S_U = n^{q-1} S_q, \quad (48)$$

so it simply relates to Tsallis  $q$ -entropy.

#### 4.5. Sharma-Mittal entropy

The Sharma-Mittal (SM) entropy [12,55] combines the Rényi entropy with the Tsallis  $q$ -entropy, and is defined as

$$S_{SM} = \frac{k_B}{R} \left[ \left( \sum_{i=1}^n (p_i)^q \right)^{\frac{R}{1-q}} - 1 \right], \quad (49)$$

where  $R$  is another dimensionless parameter apart from  $q$ . For equally probable states in (2), one gets from (49) that [56]

$$S_{SM} = \frac{k_B}{R} \left\{ \left[ 1 + \frac{1-q}{k_B} S_q \right]^{\frac{R}{1-q}} - 1 \right\}, \quad (50)$$

where  $R \rightarrow 1 - q$  limit yields the Tsallis entropy, and  $R \rightarrow 0$  limit yields the Rényi entropy. It is interesting to notice that the SM entropy obeys the composition rule of Abé (12) for  $Y = 1$  (cf. Appendix C).

#### 4.6. Kaniadakis entropy

The Kaniadakis entropy [14,15,24] results from taking into account Lorentz transformations of special relativity. It is a single  $K$ -parameter ( $-1 < K < 1$ ) deformation of BG entropy (1), with  $K$  parameter related to the dimensionless rest energy of the various parts of a multibody relativistic system. The basic definition of Kaniadakis entropy, which directly generalizes BG entropy reads

$$S_K = -k_B \sum_{i=1}^n p_i \ln_K p_i, \quad (51)$$

where  $p_i$  is the probability distribution and  $n$  is the total number of states, as mentioned in Section 2. The formula (51) introduces the  $K$ -logarithm

$$\ln_K x \equiv \frac{x^K - x^{-K}}{2K} = \frac{1}{K} \sinh(K \ln x) \quad (52)$$

with some basic properties like  $\ln_K x^{-1} = -\ln_K x$  and  $\ln_{-K} x = \ln_K x$ , giving the standard logarithm  $\ln x$  in the limit  $K \rightarrow 0$ . An equivalent definition of Kaniadakis entropy, which can be obtained after the application of  $K$ -logarithm (52) reads

$$S_K = -k_B \sum_{i=1}^n \frac{(p_i)^{1+K} - (p_i)^{1-K}}{2K}. \quad (53)$$

The  $K$ -deformed logarithm is associated with the  $K$ -exponential as follows

$$\exp_K x = \exp \left[ \frac{1}{K} \operatorname{arcsinh}(Kx) \right] = \left( \sqrt{1 + K^2 x^2} + Kx \right)^{1/K}, \quad (54)$$

and it fulfils some basic relations like  $\exp_K(x)\exp_K(-x) = 1$ , and  $\exp_K(x)\exp_{-K}(x)$ , giving the standard exponential function  $\exp(x)$  in the limit  $K \rightarrow 0$ . In fact,  $K$ -logarithm and  $K$ -exponential are the inverse functions, which means that they fulfil the relation

$$\ln_K(\exp_K x) = \exp_K(\ln_K x) = x. \quad (55)$$

The  $K$ -logarithm fulfils a generalized composition rule which reads

$$\ln_K(xy) = \ln_K x \sqrt{1 + K^2(\ln_K y)^2} + \ln_K y \sqrt{1 + K^2(\ln_K x)^2}, \quad (56)$$

and it admits the standard logarithm rule  $\ln(xy) = \ln x + \ln y$ , in the limit  $K \rightarrow 0$ . The rule (56) comes from the definition of  $K$ -sum

$$(x \oplus y)_K = x \sqrt{1 + K^2 y^2} + y \sqrt{1 + K^2 x^2}, \quad (57)$$

where one replaced  $x \rightarrow \ln x$  and  $y \rightarrow \ln y$ , and gives standard additivity rule  $(x \oplus y)_K = x + y$  in the limit  $K \rightarrow 0$ . Using the definition of Kaniadakis entropy (51), as well as the  $K$ -logarithm addition rule, we can write down the Kaniadakis entropy composition rule as follows

$$S_K(A + B) = S_K(A) \sqrt{1 + \frac{K^2}{k_B^2} S_K(B)} + S_K(B) \sqrt{1 + \frac{K^2}{k_B^2} S_K(A)}, \quad (58)$$

which we start calling  $K$ -addition rule from now on. It is interesting to note that by the application of the  $K$ -sum [14] defined as

$$(x \otimes y)_K = \frac{1}{K} \sinh \left[ \frac{1}{K} \operatorname{arcsinh}(Kx) \operatorname{arcsinh}(Ky) \right], \quad (59)$$

one has for the  $K$ -logarithm

$$\ln_K [(x \otimes y)_K] = \ln_K x + \ln_K y, \quad (60)$$

so that applying it to (51), the Kaniadakis entropy (in full analogy to the  $q$ -product of Tsallis given by (22)), can take a completely *additive form* as below (cf. the definition of additivity (4) for statistically independent systems)

$$S_K(A + B)_K = S_K(p_A p_B) = S_K(p_A) + S_K(p_B) = S_K(A) + S_K(B). \quad (61)$$

Using the  $K$ -product, the Kaniadakis entropy can also be *extensive*

$$S_K(x^{\otimes r}) = r S_K(x), \quad (62)$$

where  $r = \text{const.}$ , and it comes as a result of the identity

$$\ln_K(x^{\otimes r}) = r \ln_{rK}(x). \quad (63)$$

Finally, in analogy to the previous considerations, and under the assumption that all the states are equally probable, as in (2), one gets from (51) that

$$\ln_K p_i = -\frac{1}{K} \frac{e^{K \ln n} - e^{-K \ln n}}{2}, \quad (64)$$

which can further be transformed into

$$S_K = \frac{k_B}{K} \sinh\left(\frac{K}{k_B} S\right), \quad (65)$$

where  $S = k_B \ln n$  is the BG entropy (3).

#### 4.7. Thermal equilibrium temperature vs equilibrium temperature for nonextensive entropies

One of the important issues related to the nonextensive systems is the formulation of a proper definition of the thermal equilibrium temperature which is necessary according to the zeroth law of thermodynamics [36,52,57]. This problem can be solved by using the notion of the *effective equilibrium temperature* based on the equilibrium condition [46], recently discussed in more detail in Ref. [42]. In defining this temperature, one uses an analogy to some strongly coupled quantum systems which can be in equilibrium at the effective temperature, but not in the thermal equilibrium, as for the extensive systems. The equilibrium temperature is obtained by maximising the composition rule (9) with the fixed total energy of the system  $U_{AB} = U_A + U_B$  which leads to the conditions  $\delta S_{AB} = 0$ , with  $S_{AB} = g(S_A, S_B)$ , and  $\delta U(A+B) = 0$ . The equilibrium temperature  $T_{eq}$  then reads

$$T_{eq} \equiv \frac{1}{k_B \beta^*} = \frac{\frac{\partial g}{\partial S_B}}{k_B \beta_A} = \frac{\frac{\partial g}{\partial S_A}}{k_B \beta_B}, \quad (66)$$

where we have defined  $\beta$  for each system i.e.

$$k_B \beta_A = \frac{\partial S_A}{\partial U_A} \quad \text{and} \quad k_B \beta_B = \frac{\partial S_B}{\partial U_B}. \quad (67)$$

The application of the above procedure to the specific system described by the Tsallis  $q$ -entropy (19), which fulfils the Abé composition rule (12) as an example of the general composition rule (9), gives the equilibrium temperature [42]

$$T_{eq} = T_R = \left(1 + \frac{1-q}{k_B} S_q\right) \frac{1}{k_B \beta'}, \quad (68)$$

which is the Rényi temperature  $T_R$  [58] corresponding to Rényi entropy  $S_R$  given by (29) according to the Clausius formula (16).

It is interesting that the Rényi entropy and the Rényi effective equilibrium temperature, can be defined on a horizon of a Schwarzschild black hole [5–8,59], by assuming that the Bekenstein entropy  $S_{Bek}$  given by (14) is the Tsallis entropy  $S_q$  in (29) and (68).

Similar procedure of introducing the equilibrium temperature can be performed for the Tsallis-Cirto  $\delta$ -entropy by calculating the corresponding temperature from the Clausius relation (16) as follows [42,58]

$$T_\delta = \frac{T_H}{\delta} \left(\frac{S_{Bek}}{k_B}\right)^{1-\delta}, \quad (69)$$

which scales with  $1/M^2$  for  $\delta = 3/2$ , i.e.,  $T_\delta \propto 1/M^2$  for a Schwarzschild black hole.

#### 4.8. Classification of entropies

Bearing in mind all the considerations of the whole Section 4, we present the summary of the additivity and extensivity properties of entropies in the Table 2.

There is the whole group of Tsallis invented thermodynamical entropies, which generalize BG entropy in some different ways (cf. also Table 1). They obey either the Abé composition rule or  $\delta$ -addition rule. The Tsallis  $q$ -entropy relates to both the Rényi and the Landsberg  $\mathcal{U}$  entropies, while it is generalized by the Sharma-Mittal entropy. On the other hand, the Tsallis-Cirto  $\delta$ -entropy is related to the Barrow entropy and the Tsallis-Jensen  $q, \gamma$  entropy and, interestingly, it points towards extensivity,

**Table 2.** The additivity, extensivity, and composability properties of entropies.

Entropy Type	Extensivity	Additivity	Abé rule	$\delta$ -rule	$K$ -rule
Boltzmann-Gibbs $S_{BG}$	yes	yes	yes, $Y = 0$	yes, $\delta = 1$	yes, $K = 0$
Bekenstein $S_{Bek}$	no	no*	no	no	no
Tsallis $q$ -entropy $S_q$	no	no	yes, $Y = 1 - q$	no	no
Tsallis-Cirto $S_\delta$ ( $\delta \neq \frac{3}{2}$ )	no	no	no	yes	no
Tsallis-Cirto $S_\delta$ ( $\delta = \frac{3}{2}$ )	yes	no	no	yes, $\delta = \frac{3}{2}$	no
Barrow $S_{Bar} = S_{Bek}$ ( $\Delta = 0$ )	no	no*	no	no	no
Barrow $S_{Bar}$ ( $0 < \Delta < 1$ )	no	no	no	yes	no
Barrow $S_{Bar}$ ( $\Delta = 1$ )	yes	no	no	yes, $\delta = \frac{3}{2}$	no
Rényi $S_R$	no	yes	yes, $Y = 0$	no	no
Landsberg $U$ -entropy $S_U$	no	no	yes, $Y = q - 1$	no	no
Kaniadakis $S_K$	no	no	no	no	yes
Sharma-Mittal $S_{SM}(q, R)$	no	no	yes, $Y = R$	no	no
Tsallis $q, \delta$ -entropy $S_{q,\delta}$	no	no	no	no	no
Tsallis-Jensen $S_{q,\gamma}$	no	no	no	yes, $\delta = 1/\gamma$	no
Tsallis-Jensen $S_{1,\gamma}$	no	no	no	yes, $\delta = 1/\gamma$	no
Tsallis-Jensen $S_{q,1} = S_R$	no	yes	yes, $Y = 0$	no	no

\* obeys square root composition rule (17)

when observations are taken into account [18,38,54]. The Kaniadakis entropy seem to form a separate group of nonextensive entropies because of its hyperbolic formulation as a consequence of relativity theory being taken into account, but it stil has a BG limit. In fact, all the entropies in our study have BG limit, except Bekenstein entropy, which is not formulated as proper statistics. However, it is composable though its composition rule is unique among any other rules, which are often shared with themselves. Besides, in the microscopic counting of states in the string theory [60,61], the nonextensive Bekenstein formula is recovered, though it was also found that some higher-dimensional extremal black holes allow Bekenstein entropy, which is consistent with the Boltzmann-Gibbs extensive limit.

#### 4.9. Generalized four- and five-parameter entropic forms

There exists a four-parameter entropic formula [62], which reads

$$S_g(\alpha_{\pm}, \beta, \sigma) = \frac{k_B}{\sigma} \left[ \left( 1 + \frac{\alpha_+ S}{\beta k_B} \right)^\beta - \left( 1 + \frac{\alpha_- S}{\beta k_B} \right)^{-\beta} \right], \quad (70)$$

as well as the five-parameter formula [63] of the form

$$S_g(\alpha_{\pm}, \beta, \sigma, \epsilon) = \frac{k_B}{\sigma} \left\{ \left[ 1 + \frac{1}{\epsilon} \tanh \left( \frac{\epsilon \alpha_+ S}{\beta k_B} \right) \right]^\beta - \left[ 1 + \frac{1}{\epsilon} \tanh \left( \frac{\epsilon \alpha_- S}{\beta k_B} \right) \right]^{-\beta} \right\}, \quad (71)$$

Both these formulas generalize some of the entropies which are contained in the Table 2 and have the following limits:

1. if  $\epsilon \rightarrow 0$ , then one recovers Tsallis-Cirto (33) and Barrow (45) entropies;
2. if  $\epsilon \rightarrow 0$ ,  $\alpha_- \rightarrow 0$ ,  $\beta \rightarrow 0$ , and  $\alpha_+/\beta$  finite, then one recovers Rényi entropy (29);
3. if  $\epsilon \rightarrow 0$ ,  $\alpha_- \rightarrow 0$ ,  $\sigma = \alpha_+ = R$ , and  $\beta = R/\delta$ , then one recovers Sharma-Mittal entropy formula (50), though only when one replaces Tsallis  $q$ -entropy  $S_q$  with Tsallis-Cirto  $\delta$ -entropy  $S_\delta$ ;
4. if  $\epsilon \rightarrow 0$ ,  $\beta \rightarrow \infty$ ,  $\alpha_+ = \alpha_- = \sigma/2 = K$ , then one recovers Kaniadakis entropy (65).

These entropies have an important advantage for cosmology. Namely, they are singular-free at the cosmological bounce, which is characterized by vanishing of the Hubble parameter  $H = 0$  in a bouncing scenarios [64]. Besides, they allow microscopic interpretation [65,66].

Since all of them are in general nonadditive and nonextensive, then it is not necessary to discuss them in more detail, though they may recover BG behaviour in some special cases. However, we will not consider these cases in detail here.

## 5. Summary and discussion

Beginning with the underlying properties of the Boltzmann-Gibbs classical entropy, we have investigated the problem of nonextensivity and nonadditivity in thermodynamics aiming towards gravitational systems which admit long-range interactions. The focus has been on such extensions of Boltzmann-Gibbs entropic form, which allow various deformations of it, via some new parameters modifying the space of microstates  $\Omega$ . These new parameters are given some interpretations according to a deformation and can be enumerated as: Tsallis nonextensivity  $q$ -parameter, Tsallis-Cirto nonextensivity  $\delta$ -parameter which is equivalent to Barrow fractality  $\Delta$ -parameter, Tsallis-Jensen nonextensivity parameter  $\gamma$ , Sharma-Mittal  $R$ -parameter, and Kaniadakis relativistic  $K$ -deformation parameter.

The entropies under study may fulfil some composition rules which are the Abé rule, and  $\delta$ -addition rule, both of which are nonadditive. The Bekenstein entropy obeys some other nonadditive rule, called the square root rule, which is somewhat similar, but not the type of the Abé rule. The Kaniadakis entropy can be made additive within a special  $K$ -deformed algebra, and it reaches the standard Boltzmann-Gibbs additive rule in the limit  $K \rightarrow 0$ . The same may refer to other entropies which are composable, such as the Tsallis  $q$ -entropy and the Bekenstein entropy. A separate matter is whether these "made-additive" entropies have a proper physical interpretation. Another point that we have discussed was the definition of the equilibrium temperature for nonextensive systems in contrast to the thermal equilibrium temperature, which is ambiguous for these systems.

We have presented two important comparative tables with the additivity, extensivity, and composability properties of the entropies under study, and the relations between them have been shown explicitly. Up to our knowledge, such a collective comparison of entropic forms have not yet been presented in the literature. In view of recent interest of both astrophysicists and cosmologists in the application of the plethora of alternative to Boltzmann-Gibbs entropies, this paper may then serve a useful guide to these applications. It is worth noting that a number of these entropies are considered suitable as the candidates for the holographic dark energy and so the presented summary gives some structure to their general properties in a systematic way, which can help astrophysicists and cosmologists sort them out into categories which can then be systematically verified by the data. Nowadays, it seems that most of the tests are being done mainly for some particular types of entropies without much regard to the relation to others.

**Funding:** This research was partially supported by the Polish National Science Centre grant No. DEC-2020/39/O/ST2/02323.

**Acknowledgments:** The discussions with Ilim Çimdiker and Hussain Gohar are acknowledged.

**Conflicts of Interest:** The author declares no conflict of interest.

## Appendix A Equivalent forms of Tsallis $q$ -entropy

There are three equivalent forms of Tsallis  $q$ -entropy given by (19), which we can label according to their appearance in (19):  $S_{qI}$ ,  $S_{qII}$ ,  $S_{qIII}$ . We show their equivalence, by reducing each of them to the form (25) by the application the probability sum rule  $\sum p_i = 1$  and the formula (20). For  $S_{qI}$  one has

$$S_{qI} \equiv k_B \sum_{i=1}^n p_i \ln_q \frac{1}{p_i} = k_B \sum_{i=1}^n p_i \frac{\left(\frac{1}{p_i}\right)^{1-q} - 1}{1-q} = k_B \sum_{i=1}^n \frac{(p_i)^q - 1}{1-q} = k_B \frac{1 - \sum_{i=1}^n (p_i)^q}{q-1}, \quad (A1)$$

which is equivalent to (25). For  $S_{qII}$ , one has

$$S_{qII} = -k_B \sum_{i=1}^n (p_i)^q \ln_q p_i = -k_B \sum_{i=1}^n (p_i)^q \frac{(p_i)^{1-q} - 1}{1-q} = -k_B \frac{\sum_{i=1}^n [(p_i) - (p_i)^q]}{1-q} = k_B \frac{1 - \sum_{i=1}^n (p_i)^q}{q-1}, \quad (\text{A2})$$

which is equivalent to (25). For  $S_{qIII}$  one has

$$S_{qIII} = -k_B \sum_{i=1}^n \ln_{2-q} p_i = -k_B \sum_{i=1}^n (p_i) \frac{(p_i)^{q-1} - 1}{q-1} = -k_B \sum_{i=1}^n \frac{(p_i)^q - (p_i)}{q-1} = k_B \frac{1 - \sum_{i=1}^n (p_i)^q}{q-1}, \quad (\text{A3})$$

where we have applied a redefined  $q$ -logarithm (20) with  $q' = 2 - q$ . This finishes the prove.

## Appendix B Validity of Abé composition rule for Tsallis $q$ -entropy and Landsberg $U$ -entropy

We first assume two probabilistically independent systems  $A$  and  $B$  fulfilling

$$\sum_{i=1}^n \sum_{j=1}^m p_{ij}^{A+B} = \sum_{i=1}^n \sum_{j=1}^m p_i^A p_j^B \quad (\text{A4})$$

for every  $i, j$  which gives

$$\sum_{i=1}^n \sum_{j=1}^m (p_{ij}^{A+B})^q = \sum_{i=1}^n \sum_{j=1}^m (p_i^A p_j^B)^q = \sum_{i=1}^n \sum_{j=1}^m (p_i^A)^q (p_j^B)^q. \quad (\text{A5})$$

Following the definition of Tsallis  $q$ -entropy (25), one can write using (A5):

$$S_q(A+B) = k_B \frac{1 - \sum_{i=1}^n \sum_{j=1}^m (p_{ij}^{A+B})^q}{q-1} = k_B \frac{1 - \sum_{i=1}^n \sum_{j=1}^m (p_i^A)^q (p_j^B)^q}{q-1}, \quad (\text{A6})$$

as well as

$$S_q(A) = k_B \frac{1 - \sum_{i=1}^n (p_i^A)^q}{q-1}, \quad (\text{A7})$$

and analogously for  $S_q(B)$ .

Following Abé rule (12) we can then write the right-hand side of it as

$$\begin{aligned} RHS &= S_q(A) + S_q(B) + \frac{Y}{k_B} S_q(A) S_q(B) = k_B \frac{1 - \sum_{i=1}^n (p_i^A)^q}{q-1} + k_B \frac{1 - \sum_{j=1}^m (p_j^B)^q}{q-1} \\ &+ Y k_B \left( \frac{1 - \sum_{i=1}^n (p_i^A)^q}{q-1} \right) \left( \frac{1 - \sum_{j=1}^m (p_j^B)^q}{q-1} \right) = \frac{k_B}{q-1} \left[ 1 - \sum_{i=1}^n (p_i^A)^q + 1 - \sum_{j=1}^m (p_j^B)^q \right. \\ &\left. + \frac{Y}{q-1} - \frac{Y}{q-1} \sum_{i=1}^n (p_i^A)^q - \frac{Y}{q-1} \sum_{j=1}^m (p_j^B)^q + \frac{Y}{q-1} \sum_{i=1}^n (p_i^A)^q \sum_{j=1}^m (p_j^B)^q \right], \quad (\text{A8}) \end{aligned}$$

which after selecting  $Y = 1 - q$ , cancels six out of eight terms giving on the base of (A5) that

$$RHS = k_B \frac{1 - \sum_{i=1}^n \sum_{j=1}^m (p_i^A)^q (p_j^B)^q}{q-1} = k_B \frac{1 - \sum_{i=1}^n \sum_{j=1}^m (p_{ij}^{A+B})^q}{q-1} = S_q(A+B). \quad (\text{A9})$$

The proof of applicability of the Abé rule for Landsberg  $U$ -entropy (47), proceeds analogously, provided that  $Y = q - 1$ , instead.

### Appendix C Validity of Abé composition rule for Sharma-Mittal entropy

Let us write the Abé rule for the Sharma-Mittal entropy as follows

$$S_{SM}(A+B) = S_{SM}(A) + S_{SM}(B) + \frac{Y}{k_B} S_{SM}(A) S_{SM}(B) \quad (\text{A10})$$

with

$$S_{SM}(A+B) = \frac{k_B}{R} \left[ \left( \sum_{i=1}^n \sum_{j=1}^m (p_{ij}^{A+B})^q \right)^{\frac{R}{1-q}} - 1 \right] \quad (\text{A11})$$

and

$$S_{SM}(A) = \frac{k_B}{R} \left[ \left( \sum_{i=1}^n (p_i^A)^q \right)^{\frac{R}{1-q}} - 1 \right]. \quad (\text{A12})$$

Calculation of the RHS of (A10) gives

$$\begin{aligned} RHS &= \frac{k_B}{R} \left[ \left( \sum_{i=1}^n (p_i^A)^q \right)^{\frac{R}{1-q}} - 1 \right] + \frac{k_B}{R} \left[ \left( \sum_{j=1}^m (p_j^B)^q \right)^{\frac{R}{1-q}} - 1 \right] + \frac{Y}{k_B} \frac{k_B}{R} \left[ \left( \sum_{i=1}^n (p_i^A)^q \right)^{\frac{R}{1-q}} - 1 \right] \times \\ &\frac{k_B}{R} \left[ \left( \sum_{j=1}^m (p_j^B)^q \right)^{\frac{R}{1-q}} - 1 \right] = \frac{k_B}{R} \left[ \left( \sum_{i=1}^n (p_i^A)^q \right)^{\frac{R}{1-q}} - 1 + \left( \sum_{j=1}^m (p_j^B)^q \right)^{\frac{R}{1-q}} - 1 \right. \\ &\left. + \frac{Y}{R} \left( \sum_{i=1}^n (p_i^A)^q \right)^{\frac{R}{1-q}} \left( \sum_{j=1}^m (p_j^B)^q \right)^{\frac{R}{1-q}} - \frac{Y}{R} \left( \sum_{i=1}^n (p_i^A)^q \right)^{\frac{R}{1-q}} - \frac{Y}{R} \left( \sum_{j=1}^m (p_j^B)^q \right)^{\frac{R}{1-q}} + \frac{Y}{R} \right] \\ &= \frac{k_B}{R} \left[ \left( \sum_{i=1}^n \sum_{j=1}^m (p_i^A)^q (p_j^B)^q \right)^{\frac{R}{1-q}} - 1 \right] = \frac{k_B}{R} \left[ \left( \sum_{i=1}^n \sum_{j=1}^m (p_{ij}^{A+B})^q \right)^{\frac{R}{1-q}} - 1 \right] = LHS, \end{aligned} \quad (\text{A13})$$

where we have taken  $Y = R$ , and applied (A5).

### References

1. Swendsen, R.H. *An Introduction to Statistical Mechanics and Thermodynamics*; Oxford University Press, Oxford - New York, 2012.
2. Bekenstein, J.D. Black holes and entropy. *Phys. Rev. D* **1973**, *7*, 2333–2346. doi:10.1103/PhysRevD.7.2333.
3. Hawking, S.W. Particle Creation by Black Holes. *Commun. Math. Phys.* **1975**, *43*, 199–220. [Erratum: *Commun.Math.Phys.* *46*, 206 (1976)], doi:10.1007/BF02345020.
4. Tsallis, C.; Cirto, L.J.L. Black hole thermodynamical entropy. *Eur. Phys. J. C* **2013**, *73*, 2487, [arXiv:cond-mat.stat-mech/1202.2154]. doi:10.1140/epjc/s10052-013-2487-6.
5. Biró, T.S.; Czinner, V.G. A  $q$ -parameter bound for particle spectra based on black hole thermodynamics with Rényi entropy. *Phys. Lett. B* **2013**, *726*, 861–865, [arXiv:gr-qc/1309.4261]. doi:10.1016/j.physletb.2013.09.032.
6. Czinner, V.G.; Iguchi, H. A Zeroth Law Compatible Model to Kerr Black Hole Thermodynamics. *Universe* **2017**, *3*, 14. doi:10.3390/universe3010014.
7. Czinner, V.G.; Iguchi, H. Rényi Entropy and the Thermodynamic Stability of Black Holes. *Phys. Lett. B* **2016**, *752*, 306–310, [arXiv:gr-qc/1511.06963]. doi:10.1016/j.physletb.2015.11.061.
8. Czinner, V.G.; Iguchi, H. Thermodynamics, stability and Hawking–Page transition of Kerr black holes from Rényi statistics. *Eur. Phys. J. C* **2017**, *77*, 892, [arXiv:gr-qc/1702.05341]. doi:10.1140/epjc/s10052-017-5453-x.
9. Tsallis, C. Black Hole Entropy: A Closer Look. *Entropy* **2019**, *22*, 17. doi:10.3390/e22010017.
10. Tsallis, C. Possible Generalization of Boltzmann-Gibbs Statistics. *J. Statist. Phys.* **1988**, *52*, 479–487. doi:10.1007/BF01016429.
11. Rényi, A. On the dimension and entropy of probability distributions. *Acta Mathematica Academiae Scientiarum Hungaricae* **1959**, *10*, 193–215. doi:https://doi.org/10.1007/BF02063299.



12. Sharma, B.D.; Mittal, D.P. New non-additive measures of relative information. *J. Comb. Inf. Syst. Sci.* **1977**, *2*, 122–133.
13. Sharma, B.D.; Mittal, D.P. New non-additive measures of entropy for discrete probability distributions. *J. Math. Sci.* **10**, 28–40.
14. Kaniadakis, G. Statistical mechanics in the context of special relativity. *Phys. Rev. E* **2002**, *66*, 056125, [[cond-mat/0210467](#)]. doi:10.1103/PhysRevE.66.056125.
15. Kaniadakis, G. Statistical mechanics in the context of special relativity. II. *Phys. Rev. E* **2005**, *72*, 036108, [[cond-mat/0507311](#)]. doi:10.1103/PhysRevE.72.036108.
16. Barrow, J.D. The area of a rough black hole. *Physics Letters B* **2020**, *808*, 135643. doi:<https://doi.org/10.1016/j.physletb.2020.135643>.
17. Saridakis, E.N. Barrow holographic dark energy. *Phys. Rev. D* **2020**, *102*, 123525, [[arXiv:gr-qc/2005.04115](#)]. doi:10.1103/PhysRevD.102.123525.
18. Dąbrowski, M.P.; Salzano, V. Geometrical observational bounds on a fractal horizon holographic dark energy. *Phys. Rev. D* **2020**, *102*, 064047. doi:10.1103/PhysRevD.102.064047.
19. Saridakis, E.N. Modified cosmology through spacetime thermodynamics and Barrow horizon entropy. *Journal of Cosmology and Astroparticle Physics* **2020**, *2020*, 031. doi:10.1088/1475-7516/2020/07/031.
20. Al Mamon, A. Study of Tsallis holographic dark energy model in the framework of Fractal cosmology. *Mod. Phys. Lett. A* **2020**, *35*, 2050251, [[arXiv:gr-qc/2007.01591](#)]. doi:10.1142/S021773232050251X.
21. Asghari, M.; Sheykhi, A. Observational constraints on Tsallis modified gravity. *Mon. Not. Roy. Astron. Soc.* **2021**, *508*, 2855–2861, [[arXiv:gr-qc/2106.15551](#)]. doi:10.1093/mnras/stab2671.
22. Adhikary, P.; Das, S.; Basilakos, S.; Saridakis, E.N. Barrow holographic dark energy in a nonflat universe. *Phys. Rev. D* **2021**, *104*, 123519, [[arXiv:gr-qc/2104.13118](#)]. doi:10.1103/PhysRevD.104.123519.
23. Nojiri, S.; Odintsov, S.D.; Faraoni, V. Area-law versus Rényi and Tsallis black hole entropies. *Phys. Rev. D* **2021**, *104*, 084030, [[arXiv:gr-qc/2109.05315](#)]. doi:10.1103/PhysRevD.104.084030.
24. Drepanou, N.; Lymperis, A.; Saridakis, E.N.; Yesmakhanova, K. Kaniadakis holographic dark energy and cosmology. *Eur. Phys. J. C* **2022**, *82*, 449, [[arXiv:gr-qc/2109.09181](#)]. doi:10.1140/epjc/s10052-022-10415-9.
25. Di Gennaro, S.; Ong, Y.C. Sign Switching Dark Energy from a Running Barrow Entropy. *Universe* **2022**, *8*, 541, [[arXiv:gr-qc/2205.09311](#)]. doi:10.3390/universe8100541.
26. Di Gennaro, S.; Xu, H.; Ong, Y.C. How barrow entropy modifies gravity: with comments on Tsallis entropy. *Eur. Phys. J. C* **2022**, *82*, 1066, [[arXiv:gr-qc/2207.09271](#)]. doi:10.1140/epjc/s10052-022-11040-2.
27. Nojiri, S.; Odintsov, S.D.; Paul, T. Barrow entropic dark energy: A member of generalized holographic dark energy family. *Phys. Lett. B* **2022**, *825*, 136844, [[arXiv:gr-qc/2112.10159](#)]. doi:10.1016/j.physletb.2021.136844.
28. Anagnostopoulos, F.K.; Basilakos, S.; Saridakis, E.N. Observational constraints on Barrow holographic dark energy **2020**. [[arXiv:gr-qc/2005.10302](#)].
29. Sheykhi, A. Modified Friedmann Equations from Tsallis Entropy. *Phys. Lett. B* **2018**, *785*, 118–126, [[arXiv:gr-qc/1806.03996](#)]. doi:10.1016/j.physletb.2018.08.036.
30. Feizi Mangoudehi, Z. Observational constraints on Tsallis holographic dark energy with Ricci horizon cutoff. *Astrophys. Space Sci.* **2022**, *367*, 115, [[arXiv:gr-qc/2211.17212](#)]. doi:10.1007/s10509-022-04154-2.
31. Ghoshal, A.; Lambiase, G. Constraints on Tsallis Cosmology from Big Bang Nucleosynthesis and Dark Matter Freeze-out **2021**. [[arXiv:astro-ph.CO/2104.11296](#)].
32. Luciano, G.G.; Gine, J. Baryogenesis in non-extensive Tsallis Cosmology. *Phys. Lett. B* **2022**, *833*, 137352, [[arXiv:gr-qc/2204.02723](#)]. doi:10.1016/j.physletb.2022.137352.
33. Leon, G.; Magaña, J.; Hernández-Almada, A.; García-Aspeitia, M.A.; Verdugo, T.; Motta, V. Barrow Entropy Cosmology: an observational approach with a hint of stability analysis. *JCAP* **2021**, *12*, 032, [[arXiv:astro-ph.CO/2108.10998](#)]. doi:10.1088/1475-7516/2021/12/032.
34. Hernández-Almada, A.; Leon, G.; Magaña, J.; García-Aspeitia, M.A.; Motta, V.; Saridakis, E.N.; Yesmakhanova, K. Kaniadakis-holographic dark energy: observational constraints and global dynamics. *Monthly Notices of the Royal Astronomical Society* **2022**, *511*, 4147–4158, [<https://academic.oup.com/mnras/article-pdf/511/3/4147/42594394/stac255.pdf>]. doi:10.1093/mnras/stac255.
35. Wang, S.; Wang, Y.; Li, M. Holographic Dark Energy. *Phys. Rept.* **2017**, *696*, 1–57, [[arXiv:astro-ph.CO/1612.00345](#)]. doi:10.1016/j.physrep.2017.06.003.

36. Nauenberg, M. Critique of  $q$ -entropy for thermal statistics. *Phys. Rev. E* **2003**, *67*, 036114. doi:10.1103/PhysRevE.67.036114.
37. Tsallis, C.; Mendes, R.; Plastino, A. The role of constraints within generalized nonextensive statistics. *Physica A: Statistical Mechanics and its Applications* **1998**, *261*, 534–554. doi:https://doi.org/10.1016/S0378-4371(98)00437-3.
38. Tsallis, C.; Jensen, H.J. Extensive Composable Entropy for the Analysis of Cosmological Data, 2024, [arXiv:cond-mat.stat-mech/2408.08820].
39. Landsberg, P. Entropies galore! *Brazilian Journal of Physics* **1999**, *29*, 46–49.
40. Swendsen, R.H. How physicists disagree on the meaning of entropy. *American Journal of Physics* **2011**, *79*, 342–348, [https://doi.org/10.1119/1.3536633]. doi:10.1119/1.3536633.
41. Mannaerts, S.H. Extensive quantities in thermodynamics. *European Journal of Physics* **2014**, *35*, 035017. doi:10.1088/0143-0807/35/3/035017.
42. Çimdiker, I.; Dąbrowski, M.P.; Gohar, H. Equilibrium Temperature for Black Holes with Nonextensive Entropy. *European Physical Journal C: Particles and Fields* **2023**, *83*. doi:https://doi.org/10.1140/epjc/s10052-023-11317-0.
43. Caldwell, R. A phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state. *Physics Letters B* **2002**, *545*, 23–29. doi:https://doi.org/10.1016/S0370-2693(02)02589-3.
44. Caldwell, R.R.; Kamionkowski, M.; Weinberg, N.N. Phantom Energy: Dark Energy with  $w < -1$  Causes a Cosmic Doomsday. *Phys. Rev. Lett.* **2003**, *91*, 071301. doi:10.1103/PhysRevLett.91.071301.
45. Dąbrowski, M.P. Puzzles of dark energy in the Universe - phantom. *European Journal of Physics* **2015**, *36*, 065017. doi:10.1088/0143-0807/36/6/065017.
46. Abé, S.; Martinez, S.; Pennini, F.; Plastino, A. Nonextensive thermodynamic relations. *Physics Letters A* **2001**, *281*, 126–130. doi:10.1016/s0375-9601(01)00127-x.
47. Abé, S.; Rajagopal, A.K. Macroscopic thermodynamics of equilibrium characterized by power law canonical distributions. *Europhysics Letters (EPL)* **2001**, *55*, 6–11. doi:10.1209/epl/i2001-00373-4.
48. Abé, S. General pseudoadditivity of composable entropy prescribed by the existence of equilibrium. *Physical Review E* **2001**, *63*. doi:10.1103/physreve.63.061105.
49. Alonso-Serrano, A.; Dabrowski, M.P.; Gohar, H. Nonextensive Black Hole Entropy and Quantum Gravity Effects at the Last Stages of Evaporation. *Phys. Rev. D* **2021**, *103*, 026021, [arXiv:gr-qc/2009.02129]. doi:10.1103/PhysRevD.103.026021.
50. Tsallis, C. *Introduction to Nonextensive Statistical Mechanics: Approaching a Complex World*; Springer New York, NY, 2009. doi:10.1007/978-0-387-85359-8.
51. Tsallis, C. Nonadditive entropy and nonextensive statistical mechanics -an overview after 20 years. *Brazilian Journal of Physics* **2009**, *39*, 337–356. doi:https://doi.org/10.1590/S0103-97332009000400002.
52. Biró, T.S.; Ván, P. Zeroth law compatibility of nonadditive thermodynamics. *Phys. Rev. E* **2011**, *83*, 061147. doi:10.1103/PhysRevE.83.061147.
53. Abreu, E.M.; Ananias Neto, J. Barrow black hole corrected-entropy model and Tsallis nonextensivity. *Physics Letters B* **2020**, *810*, 135805. doi:https://doi.org/10.1016/j.physletb.2020.135805.
54. Denkiewicz, T.; Salzano, V.; Dąbrowski, M.P. Barrow nearly-extensive Gibbs-like entropy favored by dynamical and geometrical data sets in cosmology. *Phys. Rev. D* **2023**, *108*, 103533. doi:10.1103/PhysRevD.108.103533.
55. Masi, M. A step beyond Tsallis and Rényi entropies. *Physics Letters A* **2005**, *338*, 217–224. doi:https://doi.org/10.1016/j.physleta.2005.01.094.
56. Sayahian Jahromi, A.; Moosavi, S.A.; Moradpour, H.; Morais Graça, J.P.; Lobo, I.P.; Salako, I.G.; Jawad, A. Generalized entropy formalism and a new holographic dark energy model. *Phys. Lett. B* **2018**, *780*, 21–24, [arXiv:gr-qc/1802.07722]. doi:10.1016/j.physletb.2018.02.052.
57. Ou, C.; Chen, J. Two long-standing problems in Tsallis' statistics. *Physica A: Statistical Mechanics and its Applications* **2006**, *370*, 525–529. doi:https://doi.org/10.1016/j.physa.2006.03.065.
58. Çimdiker, I.; Dąbrowski, M.P.; Gohar, H. Generalized uncertainty principle impact on nonextensive black hole thermodynamics. *Classical and Quantum Gravity* **2023**, *40*, 145001. doi:10.1088/1361-6382/acdb40.
59. Czinner, V.G. Black hole entropy and the zeroth law of thermodynamics. *Int. J. Mod. Phys. D* **2015**, *24*, 1542015. doi:10.1142/S0218271815420158.

60. Strominger, A.; Vafa, C. Microscopic origin of the Bekenstein-Hawking entropy. *Physics Letters B* **1996**, *379*, 99–104. doi:[https://doi.org/10.1016/0370-2693\(96\)00345-0](https://doi.org/10.1016/0370-2693(96)00345-0).
61. De Haro, S.; van Dongen, J.; Visser, M.; Butterfield, J. Conceptual analysis of black hole entropy in string theory. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* **2020**, *69*, 82–111. doi:<https://doi.org/10.1016/j.shpsb.2019.11.001>.
62. Nojiri, S.; Odintsov, S.D.; Paul, T. Early and late universe holographic cosmology from a new generalized entropy. *Physics Letters B* **2022**, *831*, 137189. doi:<https://doi.org/10.1016/j.physletb.2022.137189>.
63. Odintsov, S.D.; Paul, T. A non-singular generalized entropy and its implications on bounce cosmology. *Physics of the Dark Universe* **2023**, *39*, 101159. doi:<https://doi.org/10.1016/j.dark.2022.101159>.
64. Nojiri, S.; Odintsov, S.D.; Faraoni, V. From nonextensive statistics and black hole entropy to the holographic dark universe. *Phys. Rev. D* **2022**, *105*, 044042, [[arXiv:gr-qc/2201.02424](https://arxiv.org/abs/2201.02424)]. doi:10.1103/PhysRevD.105.044042.
65. Nojiri, S.; Odintsov, S.D.; Paul, T. Microscopic interpretation of generalized entropy. *Physics Letters B* **2023**, *847*, 138321. doi:<https://doi.org/10.1016/j.physletb.2023.138321>.
66. Nojiri, S.; Odintsov, S.D. Micro-canonical and canonical description for generalised entropy. *Physics Letters B* **2023**, *845*, 138130. doi:<https://doi.org/10.1016/j.physletb.2023.138130>.

© 2024 by the authors. Submitted to *Journal Not Specified* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).