

## LIEB-THIRRING INEQUALITIES FOR THE SHIFTED COULOMB HAMILTONIAN

THIAGO CARVALHO CORSO, TIMO WEIDL, AND ZHUOYAO ZENG

ABSTRACT. In this paper we prove sharp Lieb-Thirring (LT) inequalities for the family of shifted Coulomb Hamiltonians. More precisely, we prove the classical LT inequalities with the semi-classical constant for this family of operators in any dimension  $d \geq 3$  and any  $\gamma \geq 1$ . We also prove that the semi-classical constant is never optimal for the Cwikel-Lieb-Rozenblum (CLR) inequalities for this family of operators in any dimension. In this case, we characterize the optimal constant as the minimum of a finite set and provide an asymptotic expansion as the dimension grows. Using the same method to prove the CLR inequalities for Coulomb, we obtain more information about the conjectured optimal constant in the CLR inequality for arbitrary potentials.

## CONTENTS

1. Introduction and statement of the main results	1
1.1. Main results	2
1.2. Outline of the paper	5
2. On the sharp LT inequalities for the shifted Coulomb Hamiltonian	5
3. On the sharp CLR inequalities for the shifted Coulomb Hamiltonian	9
4. On the CLR conjecture	12
Appendix A. Proof of (3.5)	13
References	14

## 1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Let  $V = V_+ - V_- : \mathbb{R}^d \rightarrow \mathbb{R}$  be a potential function, such that the resulting Schrödinger operator  $-\Delta - V$  is a lower semi-bounded operator with compact negative part. We denote the negative eigenvalues of  $-\Delta - V$  in increasing order by

$$\lambda_0 < \lambda_1 < \lambda_2 < \dots < 0$$

and the multiplicity of each  $\lambda_j$  by  $\mu_j \in \mathbb{N}$ . In the study of spectral behavior of Schrödinger operators, the Lieb-Thirring inequalities play an important role.

**Theorem 1.1** (Lieb-Thirring family of inequalities). *Let  $d \in \mathbb{N}$  and*

$$\gamma \in \begin{cases} [1/2, \infty), & \text{if } d = 1, \\ (0, \infty), & \text{if } d = 2, \\ [0, \infty), & \text{if } d \geq 3. \end{cases}$$

*Date:* September 17, 2024.

*2020 Mathematics Subject Classification.* Primary 35J10 ; Secondary 35P15, 47A75, 81Q10.

*Key words and phrases.* Cwikel-Lieb-Rozenblum inequalities, Lieb-Thirring inequalities, Schrödinger operators, Coulomb Hamiltonian, spectral estimations .

*Funding information:* DFG – Project-ID 442047500 – SFB 1481.

©2024 by the authors. Faithful reproduction of this article, in its entirety, by any means is permitted for noncommercial purposes.

Then, there exists a constant  $R \in \mathbb{R}_+$  such that for all  $V \in L^1_{\text{loc}}(\mathbb{R}^d)$  with  $V_+ \in L^{\gamma+d/2}(\mathbb{R}^d)$  it holds

$$\text{Tr}(-\Delta - V)_-^\gamma \leq R L_{\gamma,d}^{\text{cl}} \int_{\mathbb{R}^d} V_+^{\gamma+d/2} dx, \quad (1.1)$$

where  $\text{Tr}(-\Delta - V)_-^\gamma := \sum_j \mu_j |\lambda_j|^\gamma$  is the Riesz means and  $L_{\gamma,d}^{\text{cl}}$  is the semi-classical constant

$$L_{\gamma,d}^{\text{cl}} := \frac{\Gamma(\gamma + 1)}{(4\pi)^{d/2} \Gamma(\gamma + 1 + d/2)} \quad (1.2)$$

with  $\Gamma$  denoting the standard gamma function.

The case  $\gamma = 0$  is more commonly known as Cwikel-Lieb-Rozenblum (CLR) inequality. Historically, the CLR-inequality was proven independently by Cwikel [Cwi77], Lieb [Lie80] and Rozenblum [Roz76]. Besides, Lieb and Thirring [LT76] proved the cases  $\gamma > 1/2$  in  $d = 1$  and  $\gamma > 0$  in  $d \geq 2$ . The "limit" case  $\gamma = 1/2$  in  $d = 1$  was settled by Weidl [Wei96]. Since the validity of LT inequalities has been completely settled, the focus nowadays lies in the optimal constants. For the best bounds on the optimal constant up to date we refer the reader to the book [FLW23] and to the recent works [CCR24, Cor24].

In this paper, we are interested in the LT family of inequalities for a special kind of Schrödinger operators, namely the shifted Coulomb Hamiltonian in  $L^2(\mathbb{R}^d)$ , defined as

$$-\Delta - \frac{\kappa}{|x|} + \Lambda \quad \text{for } \kappa, \Lambda > 0.$$

Among many possible Schrödinger operators, the Coulomb Hamiltonian is special for two reasons. For one thing, its spectrum can be explicitly resolved, which greatly simplifies its analysis. For another, it is one of the most physically relevant Schrödinger operators, as it serves as a basic quantum model for non-interacting electrons bound to a point nuclei with charge  $\kappa > 0$ . Despite these facts, there are (to the best of our knowledge) only few works attempting to explicitly compute optimal constants in LT inequalities for this family of operators, namely [FLW23] and [Sel24].

The goal of this work is in some sense to fill in this gap. More precisely, the main contributions presented here are the following:

- (i) We prove that the semi-classical constant is optimal for the family of shifted Coulomb Hamiltonians in the case  $\gamma \in [1, d/2)$  and for all dimensions  $d \geq 3$ .
- (ii) We prove that the semi-classical constant is never optimal for the CLR inequality within the family of shifted Coulomb Hamiltonians. Moreover, we characterize the optimal constant in the CLR case and give an asymptotic expansion as the dimension increases.
- (iii) We prove that the conjectured value of the optimal CLR constant for arbitrary potentials [GGM78], which is given by the minimization of a specific function over integers, can be reduced to a minimization on an interval of length  $d$  around the point  $d^2/6$ . As a by-product of this analysis, we obtain an explicit asymptotic expansion of the conjectured optimal value as the dimension grows.

**1.1. Main results.** We now present the precise statements of our main results. Our first result shows that the semi-classical constant is optimal for the Lieb-Thirring inequalities for the family of shifted Coulomb Hamiltonians with  $\gamma \geq 1$ . This result extends a result by Frank, Laptev, and Weidl [FLW23, Section 5.2.2] for the case  $d = 3$  to all dimensions  $d \geq 3$ .

**Theorem 1.2** (Optimal LT inequalities for the shifted Coulomb Hamiltonian). *Let  $d \geq 3$  and  $\gamma \in [1, d/2)$ . Then for any  $\kappa, \Lambda > 0$  we have*

$$\text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_-^\gamma < L_{\gamma,d}^{\text{cl}} \int_{\mathbb{R}^d} \left( \frac{\kappa}{|x|} - \Lambda \right)_+^{\gamma+d/2} dx, \quad (1.3)$$

where  $L_{\gamma,d}^{\text{cl}}$  is the semi-classical constant defined in (1.2).

We refer to (2.4) and (2.6) for the concrete formulae of both sides in (1.3). Since this bound for general  $\gamma \in [1, d/2)$  follows by the Aizenman-Lieb argument [AL78] from (1.3) with  $\gamma = 1$ , this case is of particular interest. Here one has

$$L_{1,d}^{\text{cl}} \int_{\mathbb{R}^d} \left( \frac{\kappa}{|x|} - \Lambda \right)_+^{1+d/2} dx = \frac{2^{2-d} \Lambda^{1-d/2} \kappa^d}{d!(d-2)}.$$

**Remark 1.3** (Improved estimate). The bound (1.3) turns out to be strict. Again, by Aizenman-Lieb, it suffices to consider the case  $\gamma = 1$ . In fact, for  $d \geq 4$  we shall prove the somewhat stronger inequality

$$\text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_- \leq \left( \frac{2^{2-d} \Lambda^{1-d/2} \kappa^d}{d!(d-2)} - \frac{\kappa^2}{4(d-1)(d-2)^2} \right)_+ \quad (1.4)$$

for any  $\kappa, \Lambda > 0$ , cf. (2.16).

For  $d = 3$  the bound (1.4) itself does not hold. However, straightforward computations yield the following modified elementary upper and lower bounds.

**Remark 1.4** (Sharp corrections to Lieb-Thirring for  $d = 3$ ). For all  $\kappa, \Lambda > 0$  we have

$$\text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_- \leq \left( \frac{\kappa^3}{12\sqrt{\Lambda}} - \frac{\kappa^2}{8} + \frac{\sqrt{\Lambda}\kappa}{24} \right)_+ \quad (1.5)$$

$$\text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_- \geq \left( \frac{\kappa^3}{12\sqrt{\Lambda}} - \frac{\kappa^2}{8} - \frac{\sqrt{\Lambda}\kappa}{12} \right)_+ \quad (1.6)$$

These inequalities are sharp as equality is achieved in (1.5) whenever  $\kappa/\sqrt{\Lambda}$  is an odd natural number and in (1.6) whenever  $\kappa/\sqrt{\Lambda}$  is an even natural number. Since  $\kappa^2/8 - \sqrt{\Lambda}\kappa/24 > 0$  whenever the LHS of (1.5) is positive, that is for  $\kappa/\sqrt{\Lambda} > 2$ , the bound (1.3) is strict for  $d = 3$ , too.

The term  $-\kappa^2/8$  in (1.5) and (1.6) is related to the so-called Scott correction [SW87]. The bounds in Remark 1.4 are instructive, since the asymptotic envelopes of the eigenvalue sum, including the semiclassical term, the Scott correction and the oscillatory third term, serve also as sharp universal upper and lower bounds. For an illustration see Figure 1.




FIGURE 1. Behavior of the bound (1.4) for  $d = 3, \Lambda = 1$  and  $\kappa \in (2, 20]$ . We see that (1.4) for  $d = 3$  oscillates between the further correction terms from Remark 1.4.

Finally, let us point out that for  $d \geq 4$  the correction term in (1.4) is *not* related to a Scott type term, since the asymptotics of the eigenvalue sums for the Coulomb Hamiltonian show a different behavior in higher dimensions [Sol96]. The sole purpose of this term is to show strictness of (1.3).

Our next result concerns the CLR inequality for the Coulomb Hamiltonian. In [Sel24], it is claimed that the optimal constant of the CLR inequality for the shifted Coulomb Hamiltonian is also given by the semi-classical constant in case  $d \geq 6$ . However, this turns out to be incorrect<sup>1</sup>. To study this optimal constant, let us define for  $d \geq 3$  the function

$$Q_d : \mathbb{R} \setminus \left\{-\frac{d-1}{2}\right\} \rightarrow \mathbb{R}, \quad t \mapsto \frac{t + \frac{d}{2}}{\left(t + \frac{d-1}{2}\right)^d} \prod_{j=1}^{d-1} (t + j) \quad (1.7)$$

and set  $Q_3^* := Q_3(0)$  as well as

$$Q_d^* := \max \left\{ Q_d(\ell) : \ell \in \mathbb{N}_0 \wedge \frac{d^2}{6} - \frac{3d}{2} + \frac{7}{3} \leq \ell \leq \frac{d^2}{6} - \frac{d}{2} - \frac{2}{3} \right\}$$

for  $d \geq 4$ . It turns out that  $Q_d^* > 1$  is optimal for the CLR inequality for the shifted Coulomb Hamiltonian for all  $d \geq 3$ . In particular, the optimal constant in the CLR inequality for the family of shifted Coulomb Hamiltonians is in fact strictly larger than the semi-classical constant *in all dimensions*  $d \geq 3$ .

**Theorem 1.5** (Optimal CLR inequalities for the shifted Coulomb Hamiltonian). *For all  $d \geq 3$ , it holds that*

$$\inf \left\{ R \in \mathbb{R}_+ : \text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_-^0 \leq R L_{0,d}^{\text{cl}} \int_{\mathbb{R}^d} \left( \frac{\kappa}{|x|} - \Lambda \right)_+^{d/2} dx \text{ for all } \kappa, \Lambda > 0 \right\} = Q_d^* > 1.$$

We can also derive an asymptotic expansion for  $Q_d^*$ .

**Corollary 1.6** (Asymptotic expansion in high dimensions). *It holds for  $d \rightarrow +\infty$  that*

$$Q_d^* = 1 + \frac{3}{2d} + \frac{45}{8d^2} + O(d^{-3}).$$

**Remark 1.7.** Our analysis shows that  $Q_d(t)$  has a unique global maximum for  $t \in [0, +\infty)$ , and that  $\lim_{t \rightarrow +\infty} Q_d(t) = 1$ . For any  $d \geq 3$ , one can find (uncountably many)  $\kappa, \Lambda > 0$ , s.t.

$$\text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_-^0 > L_{0,d}^{\text{cl}} \int_{\mathbb{R}^d} \left( \frac{\kappa}{|x|} - \Lambda \right)_+^{d/2} dx.$$

**Remark 1.8.** From Theorem 1.5, the supremum of  $Q_d$  over  $\mathbb{N}_0$ , which gives a tight upper bound to the CLR excess factor for  $-\Delta - \kappa/|x|^{-1} + \Lambda$ , is attained for  $\ell \in \mathbb{N}_0$  in an interval of length  $d - 3$  around the point  $d^2/6 - d + 5/6$ . This value is strictly larger than 1 for  $d \geq 8$ . In particular, as  $\ell$  represents the level of eigenvalues that are counted after the shift (see Subsection 2.1), Theorem 1.5 implies that the maximal excess factor in the CLR inequality for the Coulomb Hamiltonian in dimension  $d \geq 3$  is achieved when we consider about  $d^2/6 + O(d)$  distinct eigenvalues. Hence, Theorem 1.5 provides – besides the one in [GGM78] – another class of potentials for which the optimal constant in the CLR inequality is larger than the semi-classical constant (and the one-particle constant) in high dimensions.

Our last result concerns the conjectured value of the optimal CLR constant *for arbitrary potentials* proposed by Glaser, Gloser and Martin [GGM78]. We now briefly recall their conjecture. For  $d \geq 3$ , we denote the optimal excess factor in the  $d$ -dimensional CLR inequality by  $R_{0,d}$ , i.e.

$$R_{0,d} := \inf \left\{ R \in \mathbb{R}_+ : \text{Tr}(-\Delta - V)_-^0 \leq R L_{0,d}^{\text{cl}} \int_{\mathbb{R}^d} V_+^{d/2} dx \text{ for all } V \in L_{\text{loc}}^1(\mathbb{R}^d) \text{ with } V_+ \in L^{d/2}(\mathbb{R}^d) \right\}.$$

<sup>1</sup>For instance, in case  $d = 6$  and  $\kappa/\sqrt{\Lambda} = 11.1$ , we have  $\text{Tr}(-\Delta - \kappa/|x| + \Lambda)_-^0 = 121$ , but  $L_{0,d}^{\text{cl}} \int (\kappa/|x| - \Lambda)_+^{d/2} dx \approx 81.81$ . In fact, there are uncountably many such counter examples in every dimension, see Remark 1.7 and Figure 2.

Besides, for  $d \geq 3$ , we define the function

$$a_d : \mathbb{R} \setminus \left\{-\frac{d}{2}, -\frac{d}{2} + 1\right\} \rightarrow \mathbb{R}, t \mapsto \frac{t + \frac{d}{2}}{\left(t + \frac{d}{2}\right)^{\frac{d}{2}} \left(t + \frac{d}{2} - 1\right)^{\frac{d}{2}}} \prod_{k=1}^{d-1} (t+k)$$

and set  $a_d^* := \sup \{a_d(\ell) : \ell \in \mathbb{N}_0\}$ .

**Conjecture 1.9.** *It holds that*

$$R_{0,d} = a_d^*.$$

The function  $a_d$  and the constant  $a_d^*$  for the aforementioned conjecture are quite similar to  $Q_d$  and  $Q_d^*$  for Theorem 1.5. In fact, it is not hard to see that  $Q_d \leq a_d$ , and therefore, our results in Theorem 1.5 do not contradict this conjecture. Furthermore, by adapting our analysis of  $Q_d$  to the function  $a_d$ , we can show that these functions have essentially the same supremum as the dimension increases. The following theorem is a precise formulation of our results.

**Theorem 1.10** (On the conjectured optimal CLR excess factor). *For  $d = 3, 4$  we have  $a_d^* = a_d(0)$  and for  $d \geq 5$  we have*

$$a_d^* = \max \left\{ a_d(\ell) : \ell \in \mathbb{N}_0 \wedge \frac{d^2}{6} - \frac{3d}{2} + \frac{7}{3} \leq \ell \leq \frac{d^2}{6} - \frac{d}{2} - \frac{5}{3} \right\}.$$

Moreover, we have for  $d \rightarrow +\infty$  the asymptotic expansion

$$a_d^* = Q_d^* + O(d^{-5}) = 1 + \frac{3}{2d} + \frac{45}{8d^2} + O(d^{-3}). \quad (1.8)$$

**Remark 1.11.** Theorem 1.10 shows that, firstly, the conjectured value of the optimal constant in the CLR inequality approaches the semi-classical constant (with convergence rate  $1/d$ ) as the dimension increases, and secondly, that this value is essentially achieved by the shifted Coulomb Hamiltonian, i.e., up to an error of order  $1/d^5$ . To the best of our knowledge, both results are new.

**1.2. Outline of the paper.** In Section 2, we firstly recall some basic facts about the spectrum of the shifted Coulomb Hamiltonian, then we study the case  $\gamma \geq 1$  and prove Theorem 1.2. In Section 3, we study the case  $\gamma = 0$  and prove Theorem 1.5 as well as Corollary 1.6. In Section 4, we study the conjectured optimal excess factor in CLR inequalities and prove Theorem 1.10. Auxiliary properties and elementary calculations will be presented in the appendix.

## 2. ON THE SHARP LT INEQUALITIES FOR THE SHIFTED COULOMB HAMILTONIAN

**2.1. Spectrum of the shifted Coulomb Hamiltonian.** Let us recall the explicit formulae for the eigenvalues and multiplicities of the shifted Coulomb Hamiltonian. For a detailed derivation of these formulae, see [FLW23, Section 4.2.3].

Let  $d \geq 3$  and  $\kappa, \Lambda > 0$ . We set

$$\eta := \kappa/\sqrt{\Lambda} \quad \text{and} \quad \ell := \lceil (\eta - d + 1)/2 \rceil - 1, \quad (2.1)$$

where  $\lceil \bullet \rceil$  is the ceiling function. If  $\ell \leq 0$ , i.e.  $(\eta - d + 1)/2 \leq 0$ , then  $-\Delta - \kappa|x|^{-1} + \Lambda$  has no negative spectrum. Thus, it is sufficient to consider the case  $\ell = \lceil (\eta - d + 1)/2 \rceil - 1 \geq 0$ , i.e.  $\eta > d - 1$ . In this case, the negative eigenvalues of  $-\Delta - \kappa|x|^{-1} + \Lambda$  are given by

$$\lambda_j := -\frac{\kappa^2}{(2j + d - 1)^2} + \Lambda \quad \text{for } j = 0, 1, 2, \dots, \ell, \quad (2.2)$$

and each  $\lambda_j$  has the multiplicity

$$\mu_j := \frac{\Gamma(d - 1 + j) \Gamma(d - 1 + 2j)}{\Gamma(d) \Gamma(j + 1)}. \quad (2.3)$$

By (2.2) and (2.3), the Riesz means of the shifted Coulomb Hamiltonian for  $\gamma \geq 0$  reads as

$$\mathrm{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_-^\gamma = \sum_{j=0}^{\ell} \frac{\Gamma(d-1+j) \Gamma(d-1+2j)}{\Gamma(d) \Gamma(j+1)} \left( \frac{\kappa^2}{(2j+d-1)^2} - \Lambda \right)^\gamma. \quad (2.4)$$

Regarding the phase space integral, which can be explicitly calculated using variable transformation and properties of the beta function, we see that

$$\int_{\mathbb{R}^d} \left( \frac{\kappa}{|x|} - \Lambda \right)_+^{\gamma+d/2} dx = (4\pi)^{\frac{d}{2}} \frac{\Lambda^\gamma \eta^d \Gamma(\gamma+1+d/2) \Gamma(d/2-\gamma)}{2^{d-1} \Gamma(d+1) \Gamma(d/2)} < \infty \quad \text{for } \gamma < \frac{d}{2}, \quad (2.5)$$

and so we only consider  $\gamma < d/2$ . With (1.2) and (2.5), the RHS of (1.3) reads as

$$L_{\gamma,d}^{\mathrm{cl}} \int_{\mathbb{R}^d} \left( \frac{\kappa}{|x|} - \Lambda \right)_+^{\gamma+d/2} dx = \frac{\Lambda^\gamma \eta^d \Gamma(\gamma+1) \Gamma(d/2-\gamma)}{2^{d-1} \Gamma(d+1) \Gamma(d/2)}. \quad (2.6)$$

We also observe that each  $\mu_j$  can be rewritten as

$$\mu_j = \binom{d-1+j}{d-1} + \binom{d-2+j}{d-1}.$$

Therefore, we can use elementary properties of binomial coefficients and the *Hockey-Stick identity*, i.e.  $\sum_{i=r}^n \binom{i}{r} = \binom{n+1}{r+1}$  for  $n, r \in \mathbb{N}$  and  $n \geq r$ , to obtain

$$N_j := \sum_{k=0}^j \mu_k = \frac{(d+2j)\Gamma(d+j)}{\Gamma(d+1)\Gamma(j+1)} \quad \text{for } j = 0, 1, 2, \dots, \ell. \quad (2.7)$$

In particular, we see that  $\mathrm{Tr}(-\Delta - \kappa/|x| + \Lambda)_-^0 = N_\ell$ .

## 2.2. Proving the optimal LT inequality for the shifted Coulomb Hamiltonian.

*Proof of Remark 1.4.* For  $d = 3$ , the Riesz means (2.4) can be explicitly summed, and we can use the observation  $\ell = \lceil (\eta - 2)/2 \rceil - 1 = \lceil \eta/2 \rceil - 2$  with  $\eta = \kappa/\sqrt{\Lambda}$  to rewrite it as

$$\mathrm{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_- = \Lambda \left( \frac{\eta^2(\ell+1)}{4} - \frac{(\ell+1)(\ell+2)(2\ell+3)}{6} \right) = \Lambda \left( \frac{\eta^3}{12} - \frac{\eta^2}{8} + \varphi(\eta) \right) \quad (2.8)$$

with

$$\varphi : (2, +\infty) \rightarrow \mathbb{R}, \quad t \mapsto t \frac{6\delta_t(1-\delta_t) - 1}{12} - \frac{\delta_t(1-\delta_t)(1-2\delta_t)}{6}$$

and  $\delta_t := \lceil t/2 \rceil - t/2$ . Notice that  $\ell = \eta/2 + \delta_\eta - 2$ . Since for any  $m \in \mathbb{N}$ , the function

$$(0, 1] \rightarrow \mathbb{R}, \quad \varepsilon \mapsto \varphi(2m+2\varepsilon) = -\frac{2}{3}\varepsilon^3 + \frac{1-2m}{2}\varepsilon^2 + m\varepsilon^3 - \frac{m}{6}$$

takes its maximum  $(2m+1)/24$  at  $\varepsilon = 1/2$  and its minimum  $-(2m+2)/12$  at  $\varepsilon = 1$ , we deduce that

$$\Lambda \left( \frac{\eta^3}{12} - \frac{\eta^2}{8} - \frac{\eta}{12} \right) \leq \mathrm{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_- \leq \Lambda \left( \frac{\eta^3}{12} - \frac{\eta^2}{8} + \frac{\eta}{24} \right) \quad \text{for } \eta > 2. \quad (2.9)$$

As the middle term is zero for  $\eta \leq 2$  and  $\Lambda\eta^3/12$  is the semi-classical term in (2.6), the proof of Remark 1.4 is completed.  $\blacksquare$

*Proof of Theorem 1.2.* By Remark 1.3 and Remark 1.4, it remains to prove (1.4) for  $d \geq 4$ .

To this end, we begin with using (2.7) to rewrite (2.4) as

$$\mathrm{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_- = \frac{\Lambda \eta^d}{d!(d-2)} \left( d\eta^{2-d}(d-2) \sum_{k=0}^{\ell} \frac{(d-1)! \mu_k}{(d+2k-1)^2} - \eta^{-d}(d-2)d!N_\ell \right), \quad (2.10)$$

where  $\eta$  and  $\ell$  are defined as in (2.1). Recall that we only consider  $\eta > d-1$ , i.e.  $\ell \geq 0$ , as otherwise the Riesz means is zero and inequality (1.3) trivially holds. In particular, the RHS of (2.10) is non-negative.

We proceed to rewrite and estimate the sum in the RHS of (2.10). Let us define  $p_0(t) := 1$  and

$$p_m(t) := \prod_{k=1}^m (t+k) \quad \text{for } m \in \mathbb{N}.$$

A key property of  $p_m$ , which can be directly verified from its definition, is the recursive relation

$$p_m(t) = \frac{p_{m+1}(t) - p_{m+1}(t-1)}{m+1} \quad \text{for all } m \in \mathbb{N}_0, t \in \mathbb{R}. \quad (2.11)$$

This relation allows us to explicitly evaluate the sum  $\sum_{k=0}^{\ell} p_m(k)$  via the formula

$$\sum_{k=0}^{\ell} p_m(k) = \sum_{k=0}^{\ell} \frac{p_{m+1}(k) - p_{m+1}(k-1)}{m+1} = \frac{p_{m+1}(\ell)}{m+1}. \quad (2.12)$$

Therefore, we find that

$$\begin{aligned} (d-2) \sum_{k=0}^{\ell} \frac{(d-1)! \mu_k}{(2k+d-1)^2} &= \frac{d-2}{2} \left( \sum_{k=0}^{\ell} p_{d-3}(k) \right) + \frac{d-2}{2} \left( \sum_{k=0}^{\ell} \frac{d-3}{2} \frac{p_{d-3}(k)}{k + \frac{d-1}{2}} \right) \\ &= \frac{p_{d-2}(\ell)}{2} + \frac{(d-3)}{4} \sum_{k=0}^{\ell} (p_{d-2}(k) - p_{d-2}(k-1)) \frac{1}{k + \frac{d-1}{2}} \\ &= \frac{p_{d-2}(\ell)}{2} + \frac{(d-3)}{4} \frac{p_{d-2}(\ell)}{\ell + \frac{d-1}{2}} + \frac{d-3}{4} \sum_{k=0}^{\ell-1} \frac{p_{d-2}(k)}{\left(k + \frac{d+1}{2}\right) \left(k + \frac{d-1}{2}\right)}, \end{aligned} \quad (2.13)$$

where the first equality results from the identity

$$\frac{(d-1)! \mu_k}{(2k+d-1)^2} = \frac{p_{d-2}(k)}{2k+d-1} = \frac{1}{2} \left( p_{d-3}(k) + \frac{d-3}{2} \frac{p_{d-3}(k)}{k + \frac{d-1}{2}} \right),$$

and the second equality is due to (2.11) and (2.12) respectively applied to each sum. The last equality in (2.13) is derived using the formula of summation by parts, i.e.  $\sum_{k=0}^{\ell} A_k (B_k - B_{k-1}) = A_{\ell} B_{\ell} + \sum_{k=0}^{\ell-1} (A_k - A_{k+1}) B_k$ , applied to  $A_k := (k + (d-1)/2)^{-1}$  and  $B_k := p_{d-2}(k)$  (with the convention  $0 = \sum_{k=0}^{-1} a_k$  for any  $a_k$  in case  $\ell = 0$ ).

Regarding the last sum in (2.13), we see that

$$\begin{aligned} \frac{d-3}{4} \sum_{k=0}^{\ell-1} \frac{p_{d-2}(k)}{\left(k + \frac{d+1}{2}\right) \left(k + \frac{d-1}{2}\right)} &\leq \frac{d-3}{4} \sum_{k=0}^{\ell-1} p_{d-4}(k+1) \\ &= \frac{p_{d-3}(\ell) - p_{d-3}(0)}{4} = \frac{p_{d-2}(\ell)}{4(\ell+d-2)} - \frac{(d-3)!}{4}, \end{aligned} \quad (2.14)$$

where the inequality follows from the elementary estimate

$$\frac{(k+1)(k+d-2)}{\left(k + \frac{d+1}{2}\right) \left(k + \frac{d-1}{2}\right)} \leq 1 \quad \text{for all } d, k \in \mathbb{N}$$

together with an index and variable shift. The first equality in (2.14) follows from (2.12), and the second equality results from the fact that  $p_{d-3}(\ell) = p_{d-2}(\ell)/(\ell+d-2)$  and  $p_{d-3}(0) = (d-3)!$ . We also note that for  $\ell = 0$ , (2.14) still holds because the very left sum is zero due to the convention  $\sum_{k=0}^{-1} a_k = 0$  and the very right difference is clearly zero.

All in all, by estimating the sum in (2.13) with (2.14), plugging the resulting expression back in (2.10), and using the identity  $d! N_{\ell} = (d+2\ell)(\ell+d-1)p_{d-2}(\ell)$ , we find

$$\text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_{-} \leq \frac{\Lambda \eta^d}{d!(d-2)} p_{d-2}(\ell) \left( d\eta^{2-d} \alpha - \eta^{-d} (d-2) \beta \right) - \frac{\kappa^2}{4(d-1)(d-2)^2} \quad (2.15)$$

with

$$\alpha := \frac{1}{2} \left( 1 + \frac{d-3}{2(\ell + \frac{d-1}{2})} + \frac{1}{2(\ell + d - 2)} \right) \quad \text{and} \quad \beta := 2(\ell + \frac{d}{2})(\ell + d - 1).$$

We can now use the fact that

$$\sup_{\eta \geq 0} d\eta^{2-d}\alpha - \eta^{-d}(d-2)\beta = 2\alpha^{\frac{d}{2}}\beta^{1-\frac{d}{2}},$$

to further estimate (2.15) from above and obtain

$$\text{Tr} \left( -\Delta - \frac{\kappa}{|x|} + \Lambda \right)_- \leq \frac{2^{2-d}\Lambda\eta^d}{d!(d-2)} G(\ell) - \frac{\kappa^2}{4(d-1)(d-2)^2} \quad (2.16)$$

with  $\ell = \lceil (\eta + 1 - d)/2 \rceil - 1 \in \mathbb{N}_0$  as before and  $G$  the function defined as

$$G : [0, +\infty) \rightarrow \mathbb{R}, \quad t \mapsto \frac{p_{d-2}(t)}{\left( (t + \frac{d}{2})(t + d - 1) \right)^{\frac{d-2}{2}}} \left( 1 + \frac{d-3}{2(t + \frac{d-1}{2})} + \frac{1}{2(t + d - 2)} \right)^{\frac{d}{2}}. \quad (2.17)$$

The upcoming Lemma 2.1 shows that  $G(\ell) \leq 1$  for all  $\ell \in \mathbb{N}_0$ . Consequently, (1.4) is proven.  $\blacksquare$

**Lemma 2.1.** *For  $d \geq 4$ , the function  $G$  defined in (2.17) is strictly increasing for  $t \geq 0$  and satisfies  $\lim_{t \rightarrow +\infty} G(t) = 1$ .*

*Proof.* To simplify some calculations, it is more convenient to work with the translated function

$$g(t) := G \left( t - \frac{d-1}{2} \right) = \left( 1 + \frac{d-3}{2t} + \frac{1}{2(t + \frac{d-3}{2})} \right)^{\frac{d}{2}} \left( \left( t + \frac{1}{2} \right) \left( t + \frac{d-1}{2} \right) \right)^{\frac{2-d}{2}} \prod_{k=0}^{d-3} \left( t - \frac{d-3}{2} + k \right).$$

Our goal is then to show that  $g(t) \leq 1$  for  $t \geq (d-1)/2$  and  $\lim_{t \rightarrow +\infty} g(t) = 1$ .

The limit is immediate to compute. Hence, to prove the inequality, it suffices to show that the derivative of  $\log g(t)$  is non-negative for  $t \geq (d-1)/2$ . To this end, we first note that

$$\begin{aligned} (\log g)'(t) &= \frac{1}{t - \frac{d-3}{2}} + \sum_{k=1}^{d-3} \left( \frac{1}{t - \frac{d-3}{2} + k} \right) - \frac{d-2}{2} \left( \frac{1}{t + \frac{1}{2}} + \frac{1}{t + \frac{d-1}{2}} \right) \\ &\quad + \frac{d}{2} \left( \frac{4t + 2d - 5}{2t^2 + (2d-5)t + \frac{(d-3)^2}{2}} - \frac{1}{t} - \frac{1}{t + \frac{d-3}{2}} \right). \end{aligned} \quad (2.18)$$

Next, we use two elementary inequalities to find a simpler lower bound for  $(\log g)'(t)$ . The first inequality is immediate to verify and reads

$$\frac{4t + 2d - 5}{2t^2 + (2d-5)t + \frac{(d-3)^2}{2}} \geq \frac{4t + 2d - 5}{2t^2 + (2d-5)t + \frac{(d-2)(d-3)}{2}} = \frac{1}{t + \frac{d-2}{2}} + \frac{1}{t + \frac{d-3}{2}} \quad \text{for } t \geq 0. \quad (2.19)$$

The second inequality is a consequence of the convexity of the function  $r \in \mathbb{R}_+ \mapsto 1/r$  and can be stated as

$$\sum_{k=1}^{d-3} \frac{1}{t - \frac{d-3}{2} + k} = \sum_{k=0}^{d-4} \frac{1}{(t + \frac{1}{2}) - \frac{d-4}{2} + k} \geq \frac{d-3}{t + \frac{1}{2}} \quad \text{for } t \geq \frac{d-3}{2}. \quad (2.20)$$

Hence, applying (2.19) and (2.20) to (2.18), we find that  $(\log g)'(t) \geq h(t)$  for  $t \geq (d-1)/2$  with the function  $h : [(d-1)/2, \infty) \rightarrow \mathbb{R}$  defined as

$$h(t) := \frac{1}{t - \frac{d-3}{2}} - \frac{d}{2} + \frac{d-4}{2} + \frac{d}{2} - \frac{d-2}{2}.$$



A somewhat cumbersome but straightforward calculation then shows that

$$h(t) = \frac{h_1(t)}{(t - \frac{d-3}{2})t(t + \frac{1}{2})(t + \frac{d-2}{2})(t + \frac{d-1}{2})}$$

with

$$h_1(t) = \frac{1}{4}(d-2)(d-4)t^2 + \frac{1}{16}(5d^2 + 5d + 32)(d-1)t + \frac{1}{32}d(d-1)(d-2)(d-3).$$

One can easily see that all coefficients of  $h_1(t)$  are non-negative for  $d \geq 4$ . Therefore, we obtain  $(\log g)'(t) \geq h(t) \geq 0$  for  $t \geq (d-1)/2$ , which completes the proof. ■

### 3. ON THE SHARP CLR INEQUALITIES FOR THE SHIFTED COULOMB HAMILTONIAN

*Proof of Theorem 1.5.* By (2.7) and (2.5), we need to study the behaviour of the quotient

$$R_d(\eta) := \frac{\text{Tr} \left( -\Delta - \frac{\kappa}{|\cdot|} + \Lambda \right)_-^0}{L_{0,d}^{\text{cl}} \int_{\mathbb{R}^d} \left( \frac{\kappa}{|\cdot|} - \Lambda \right)_+^{d/2} dx} = \frac{d+2\ell}{2^{1-d}\eta^d} \prod_{j=1}^{d-1} (\ell+j), \quad (3.1)$$

where again  $\eta = \kappa/\sqrt{\Lambda}$  and  $\ell = \lceil (\eta + 1 - d)/2 \rceil - 1$ . For this, let us set  $\tau := (\eta + 1 - d)/2$  and observe that, since  $\tau \geq \ell$ , we have

$$R_d(\eta) = \frac{d+2\ell}{2^{1-d}\eta^d} \prod_{j=1}^{d-1} (\ell+j) \leq \frac{\tau + \frac{d}{2}}{\left(\tau + \frac{d-1}{2}\right)^d} \prod_{j=1}^{d-1} (\tau+j) = Q_d(\tau) \quad (3.2)$$

with  $Q_d$  defined in (1.7). Moreover, we note that, even though  $Q_d(\tau) \neq R_d(2\tau + d - 1)$  for any  $\tau > 0$ , we have the equality

$$Q_d(\tau_0) = \lim_{\tau \downarrow \tau_0} R_d(2\tau + d - 1) \quad \text{for any } \tau_0 \in \mathbb{N}_0.$$

As  $R_d(2\tau + d - 1)$  is strictly decreasing in any interval  $(\tau_0, \tau_0 + 1]$  with  $\tau_0 \in \mathbb{N}_0$ , the identity above implies that the supremum of  $Q_d$  over  $\mathbb{N}_0$  gives the optimal excess factor in the CLR inequality. For a visual illustration of  $Q_d$  and  $R_d$  see Figure 1. The proof is then completed with the upcoming Lemma 3.1. ■




FIGURE 2. Comparison of  $R_d(2\tau + d - 1)$ ,  $Q_d(\tau)$  for  $d = 5, 6$ . The bold marked dots emphasize where  $R_d$  takes values. As stated in Lemma 3.1,  $Q_d$  is monotonically decreasing for  $d \leq 5$  and firstly increasing then decreasing for  $d \geq 6$ . The monotonic decrease of  $R_d(2\tau + d - 1)$  for  $\tau \in (\ell, \ell + 1]$  and  $\ell \in \mathbb{N}_0$  is due to the ceiling function in  $R_d$ , see (3.1). Also observe that  $R_d(2\tau + d - 1) < 1$  for  $\tau \in \mathbb{N}_0$ .

**Lemma 3.1.** For  $d = 3$ , the function  $Q_d$  defined in (1.7) is strictly decreasing in  $(-1, +\infty)$ . For  $d \geq 4$ ,  $Q_d$  has a unique maximum at  $t_d^* \in (-1, +\infty)$  satisfying

$$\frac{d^2}{6} - \frac{3d}{2} + \frac{7}{3} < t_d^* < \frac{d^2}{6} - \frac{d}{2} - \frac{2}{3}. \quad (3.3)$$

*Proof of Lemma 3.1.* Let us define an auxiliary set

$$P := \left\{-\frac{d}{2}, -\frac{d-1}{2}\right\} \cup \{-k\}_{k=1}^{d-1} = \left\{\left[\frac{d}{2}\right] - \frac{1}{2}\right\} \cup \{-k\}_{k=1}^{d-1},$$

where the equality holds true because either  $d/2$  or  $(d-1)/2$  is an integer. Then, an explicit calculation yields for  $t \in \mathbb{R} \setminus P$  the representation

$$Q'_d(t) = Q_d(t) f_d(t)$$

with

$$f_d : \mathbb{R} \setminus P \rightarrow \mathbb{R}, \quad t \mapsto \frac{1}{t + \frac{d}{2}} - \frac{d}{t + \frac{d-1}{2}} + \sum_{k=1}^{d-1} \frac{1}{t+k}. \quad (3.4)$$

For  $d = 3$ , one can immediately verify that  $f_3(t) < 0$  for all  $t > -1$ . Since  $Q_d(t) > 0$  for  $t > -1$ , the case  $d = 3$  is proven.

Consider now  $d \geq 4$ . Since  $Q_d(t) > 0$  for  $t > -1$ , it suffices to prove that  $f_d$  has a unique zero in  $(-1, +\infty)$  located in the interval (3.3). To this end, we rewrite  $f_d$  as a quotient of co-prime polynomials, i.e.,

$$f_d(t) = \frac{p(t)}{q(t)},$$

where

$$q(t) := \left(t + \left[\frac{d}{2}\right] - \frac{1}{2}\right) \prod_{k=1}^{d-1} (t+k),$$

and  $p$  is a polynomial of order at most  $d-1$ . In fact, a long calculation (see appendix A) shows that  $p$  actually has degree  $d-2$  and

$$p(t) = -\frac{d}{2} \left( t^{d-2} + \left( \frac{d^2}{3} - \left[\frac{d}{2}\right] - \frac{1}{3} \right) t^{d-3} + r(t) \right), \quad (3.5)$$

where  $r$  is a polynomial of degree at most  $d-4$ . Since  $p$  and  $q$  are co-prime, every zero of  $f_d$  must be a zero of  $p$ . This implies in particular that the function  $f_d$  has at most  $d-2$  zeros.

We proceed with inferring the location of the zeros of  $f_d$  by analysing the sign changes of  $f_d$  in the intervals between two consecutive poles. More precisely, we note from (3.4) that

$$\begin{aligned} \lim_{t \downarrow -k} f_d(t) &= +\infty = - \lim_{t \uparrow -k} f_d(t) = -\infty && \text{for } k \in \{1, \dots, d\} \setminus \left\{\left[\frac{d}{2}\right]\right\}, \\ \lim_{t \downarrow -\frac{d-1}{2}} f_d(t) &= -\infty = - \lim_{t \uparrow -\frac{d-1}{2}} f_d(t), \\ \lim_{t \downarrow -\frac{d}{2}} f_d(t) &= +\infty = - \lim_{t \uparrow -\frac{d}{2}} f_d(t). \end{aligned}$$

So, by defining the open intervals  $\{I_j\}_{j=1}^{d-2}$  as

$$I_j := \begin{cases} \left(-\frac{d}{2}, -\left[\frac{d}{2}\right] + 1\right), & \text{for } j = \left[\frac{d}{2}\right] - 1, \\ \left(-\left[\frac{d}{2}\right] - 1, -\frac{d}{2}\right), & \text{for } j = \left[\frac{d}{2}\right], \\ (-j-1, -j), & \text{otherwise,} \end{cases}$$

we deduce with the intermediate value theorem that  $f_d(t)$  has at least one zero in each interval  $I_j$  for  $j \in \{1, \dots, d-2\} \setminus \{\lfloor d/2 \rfloor - 1\}$ . In particular,  $f_d$  has at least  $d-3$  zeros inside the interval  $(-d+1, -1)$ . Moreover, since the leading coefficient of  $p$  is negative, we have

$$\lim_{t \rightarrow +\infty} p(t) = -\infty \quad \text{while} \quad q(t) > 0 \quad \text{for } t \geq -1.$$

Consequently,  $f_d(t)$  is negative for  $t$  sufficiently large. From this observation and the fact that  $\lim_{t \downarrow -1} f_d(t) = +\infty$ , we deduce again with the intermediate value theorem that  $f_d$  has at least one zero in the interval  $(-1, +\infty)$ . Since  $f_d$  has at most  $d-2$  zeros, we conclude that  $f_d$  must have exactly one zero in each of the intervals  $I_j$  for  $j \in \{1, \dots, d-2\} \setminus \{\lfloor d/2 \rfloor - 1\}$  and one zero in the interval  $(-1, +\infty)$  (see Figure 3 for an example of  $f_d$ ). Moreover, the rightmost zero corresponds to a local maximum of  $Q_d$ . This implies that  $Q_d$  has exactly one maximum in the interval  $(-1, +\infty)$ .




FIGURE 3. The function  $f_d$  for  $d = 6$  and its four zeros: 3 negative ones and 1 positive one, as stated in the proof of Lemma 3.1.

We now proceed to prove the bound in (3.3). Note that the previous arguments not only give us a good qualitative picture of the function  $f_d$ , but also a quantitative estimate on the negative zeros of the polynomial  $p$ : From the previous discussion we know that  $p$  has exactly one zero in each of the intervals  $I_j$  for  $j \neq \lceil (d-1)/2 \rceil - 1$ . Therefore, we can write  $p$  in the factorized form

$$p(t) = -\frac{d}{2}(t - t_d^*) \prod_{k=1, k \neq \lfloor \frac{d}{2} \rfloor - 1}^{d-2} (t + k + \varepsilon_k), \quad (3.6)$$

where  $\varepsilon_k \in (0, 1)$  for any  $k$ , and  $t_d^*$  is the root that we would like to estimate. Hence, by comparing (3.6) with (3.5), we find with Vieta's formula applied to the term of order  $t^{d-3}$  that

$$-t_d^* + \sum_{k=1, k \neq \lfloor \frac{d}{2} \rfloor - 1}^{d-2} (k + \varepsilon_k) = \frac{d^2}{3} - \left\lfloor \frac{d}{2} \right\rfloor - \frac{1}{3},$$

from which (3.3) follows by using the estimates  $0 \leq \varepsilon_k \leq 1$ . ■

Let us now turn to the proof of Corollary 1.6.

*Proof of Corollary 1.6.* Note that the derivative of  $f_d(t)$  is given by

$$f_d'(t) = -\frac{1}{(t + \frac{d}{2})^2} + \frac{d}{(t + \frac{d-1}{2})^2} - \sum_{k=1}^{d-1} \frac{1}{(t+k)^2} = \frac{t - \frac{1}{4} + \frac{d}{2}}{(t + \frac{d}{2})^2 (t + \frac{d-1}{2})^2} + \frac{d-1}{(t + \frac{d-1}{2})^2} - \sum_{k=1}^{d-1} \frac{1}{(t+k)^2}.$$

Since  $1/(t+k)^2$  is monotonically decreasing in  $k$  for  $t \geq 0$ , the sum  $\sum_{k=1}^{d-1} (t+k)^{-2}$  can be bounded from above and below by

$$\frac{d-1}{(t+1)(t+d)} = \int_1^d \frac{1}{(t+k)^2} dk \leq \sum_{k=1}^{d-1} \frac{1}{(t+k)^2} \leq \int_0^{d-1} \frac{1}{(t+k)^2} dk = \frac{d-1}{t(t+d-1)}.$$

Then, the foregoing lower bound on the sum leads to

$$\begin{aligned} f'_d(t) &\leq \frac{t - \frac{1}{4} + \frac{d}{2}}{(t + \frac{d}{2})^2 (t + \frac{d-1}{2})^2} + (d-1) \left( \frac{1}{(t + \frac{d-1}{2})^2} - \frac{1}{(t+1)(t+d)} \right) \\ &= \frac{t - \frac{1}{4} + \frac{d}{2}}{(t + \frac{d}{2})^2 (t + \frac{d-1}{2})^2} + \frac{(d-1) \left( 2t + d - (\frac{d-1}{2})^2 \right)}{(t + \frac{d-1}{2})^2 (t+1)(t+d)}. \end{aligned}$$

Hence, for  $t = cd^2 + O(d)$  with  $c > 0$ , we see that  $f'_d(t) = O(d^{-5})$  asymptotically in the limit  $d \rightarrow +\infty$ . A similar calculation using the upper bound on  $\sum_{k=1}^{d-1} (t+k)^{-2}$  shows that  $f'_d(t)$  is also bounded from below by terms of order  $1/d^5$  for  $d \rightarrow +\infty$  and  $t = cd^2 + O(d)$  with any  $c > 0$ . In particular, we find that

$$f'_d \left( \frac{d^2}{6} + O(d) \right) = O(d^{-5}), \quad (3.7)$$

as long as  $d$  is large enough. To conclude, we observe that  $f_d(t)$  is the derivative of  $\log Q_d(t)$  and the maximising point of  $\log Q_d(t)$  satisfies  $t_d^* = d^2/6 + O(d)$  due to Lemma 3.1. Thus, by (3.7) and a Taylor expansion of  $\log Q_d$  around  $t_d^*$ , we find that

$$|\log Q_d(t) - \log Q_d(t_d^*)| \leq \sup_{s \in [t, t_d^*]} |f'_d(s)| \cdot |t - t_d^*|^2 = O(d^{-3})$$

for any  $|t - t_d^*|$  of order  $O(d)$ . Hence, we have

$$Q_d \left( \frac{d^2}{6} + O(d) \right) = Q_d(t_d^*) + O(d^{-3}).$$

By plugging in  $t := d^2/6 - 3d/2 + 7/3$  into  $Q_d$ , we obtain after some calculations the desired formula (1.8).  $\blacksquare$

#### 4. ON THE CLR CONJECTURE

*Proof of Theorem 1.10.* The proof here follows essentially the same steps as in the proof of Theorem 1.5. Therefore, we shall only sketch the main steps below. Also, for simplicity, we shall consider only the case  $d$  even. For  $d$  odd, the proof follows from similar arguments.

As before, we start with computing the derivative of  $a_d$ . For  $t \neq -1, -2, \dots, -d+1$ , we obtain after some straightforward calculations the representation  $a'_d(t) = a_d(t)g_d(t)$  with

$$g_d : \mathbb{R} \setminus \{-1, -2, \dots, -d+1\} \rightarrow \mathbb{R}, \quad t \mapsto \frac{1 - \frac{d}{2}}{t + \frac{d}{2}} - \frac{\frac{d}{2}}{t + \frac{d}{2} - 1} + \sum_{k=1}^{d-1} \frac{1}{t+k}.$$

Note that  $-d/2, -d/2 + 1 \in \{-1, -2, \dots, -d+1\}$  due to the assumption of  $d$  being even. As  $a_d$  is positive for  $t \geq 0$ , it suffices to study the behaviour of  $g_d$ . For  $d = 4$ , it is immediate to see that  $g_d \leq 0$  and therefore  $a_d$  is decreasing, which completes the proof in this case. For  $d \geq 6$ , we rewrite  $g_d$  as a quotient of co-prime polynomials, i.e.

$$g_d(t) = \frac{p(t)}{q(t)},$$

where  $q(t) := \prod_{k=1}^{d-1} (t+k)$  and  $p(t)$  is a polynomial of order  $d-2$  satisfying

$$p(t) = -\frac{d}{2} \left( t^{d-3} + \left( \frac{d^2}{3} - d + \frac{2}{3} \right) t^{d-4} + O(t^{d-5}) \right). \quad (4.1)$$

Thus,  $g_d$  has at most  $d-3$  zeros. By analysing the poles of  $g_d$  and using the intermediate value theorem, as we did in the proof of Theorem 1.5, we conclude that  $g_d$  has at least one zero along each interval of the form  $(-k-1, -k)$  for  $k \in \{1, \dots, d-2\} \setminus \{d/2, d/2-2\}$  and at least one zero in the interval  $(-1, +\infty)$ . Since  $g_d$  has at most  $d-3$  zeros, each of these intervals has exactly one zero. Hence,  $p(t)$  can be written in factorized form as

$$p(t) = -\frac{d}{2} (t - t_d^*) \prod_{k=1, k \notin \{\frac{d}{2}, \frac{d}{2}-2\}}^{d-2} (t + k + \varepsilon_j), \quad (4.2)$$

where  $t_d^*$  is the unique zero in the interval  $(-1, +\infty)$ . Expanding the RHS of (4.2) then yields

$$p(t) = -\frac{d}{2} t^{d-3} + \frac{d}{2} \left( t_d^* - \sum_{k=1, k \notin \{\frac{d}{2}, \frac{d}{2}-2\}}^{d-2} (k + \varepsilon_k) \right) t^{d-4} + O(t^{d-5}). \quad (4.3)$$

The estimate on  $t_d^*$  now follows by equating the second coefficient in (4.3) with the one in (4.1) and using the estimates  $0 \leq \varepsilon_k \leq 1$ .

To prove the asymptotic expansion in (1.8) we note that

$$a_d(t) = Q_d(t) \frac{(t + \frac{d-1}{2})^d}{(t + \frac{d}{2})^{\frac{d}{2}} (t + \frac{d}{2} - 1)^{\frac{d}{2}}}.$$

Hence, for  $t = cd^2 + O(d)$  with any  $c > 0$ , we see that

$$\log a_d(t) - \log Q_d(t) = \frac{d}{t + \frac{d-1}{2}} - \frac{d/2}{t + \frac{d}{2}} - \frac{d/2}{t + \frac{d}{2} - 1} = \frac{d}{(t + \frac{d-1}{2})(t + \frac{d}{2})(t + \frac{d}{2} - 1)} = O(d^{-5}).$$

As the maximising points of both  $Q_d$  and  $a_d$  lies within the interval  $[d^2/6 - 3d/2, d^2/6 - d/2]$ , we obtain the first equality in (1.8). The second equality in (1.8) follows then with the expansion from Corollary 1.6.  $\blacksquare$

#### APPENDIX A. PROOF OF (3.5)

*Proof of equation (3.5).* Since  $p(t) = q(t)f_d(t)$ , (3.4) yields

$$\begin{aligned} p(t) &= q(t) \left( \left( \frac{1}{t + \frac{d}{2}} - \frac{1}{t + \frac{d-1}{2}} \right) + \sum_{k=1}^{d-1} \left( \frac{1}{t+k} - \frac{1}{t + \frac{d-1}{2}} \right) \right) \\ &= -\frac{1}{2} \prod_{k=1, k \neq \lfloor \frac{d}{2} \rfloor}^{d-1} (t+k) + \sum_{k=1}^{d-1} \left( \frac{d-1}{2} - k \right) \frac{t + \frac{d}{2}}{t + \lfloor \frac{d}{2} \rfloor} \prod_{j=1, j \neq k}^{d-1} (t+j) = I(t) + J(t) \end{aligned}$$

where

$$\begin{aligned} I(t) &:= \left( \frac{d-1}{2} - \left\lfloor \frac{d}{2} \right\rfloor - \frac{1}{2} \right) \prod_{k=1, k \neq \lfloor \frac{d}{2} \rfloor}^{d-1} (t+k), \\ J(t) &:= \sum_{k=1, k \neq \lfloor \frac{d}{2} \rfloor}^{d-1} \left( \frac{d-1}{2} - k \right) \left( t + \frac{d}{2} \right) \prod_{j=1, j \neq \{k, \lfloor \frac{d}{2} \rfloor\}}^{d-1} (t+j). \end{aligned}$$

We can further expand  $I(t)$  as

$$I(t) = \left( \frac{d}{2} - \left\lfloor \frac{d}{2} \right\rfloor - 1 \right) \left( t^{d-2} + S(d)t^{d-3} + O(t^{d-4}) \right)$$

with  $S(d) := \sum_{k=1, k \neq \lfloor \frac{d}{2} \rfloor}^{d-1} k$ . Regarding  $J(t)$ , we find

$$\begin{aligned} J(t) &= \sum_{k=1, k \neq \lfloor \frac{d}{2} \rfloor}^{d-1} \left( \frac{d-1}{2} - k \right) \left( t^{d-2} + \left( \frac{d}{2} + \sum_{j=1, j \notin \{k, \lfloor \frac{d}{2} \rfloor\}}^{d-1} j \right) t^{d-3} + O(t^{d-4}) \right) \\ &\approx \left( 1 - \left\lfloor \frac{d}{2} \right\rfloor \right) t^{d-2} + \left( \sum_{k=1, k \neq \lfloor \frac{d}{2} \rfloor}^{d-1} \left( \frac{d-1}{2} - k \right) \left( S(d) + \frac{d}{2} - k \right) \right) t^{d-3} \\ &= \left( 1 - \left\lfloor \frac{d}{2} \right\rfloor \right) t^{d-2} + \left( \left( \frac{3}{2} - \left\lfloor \frac{d}{2} \right\rfloor - d \right) S(d) + \frac{(d-2)(d-1)d}{4} + \sum_{k=1, k \neq \lfloor \frac{d}{2} \rfloor}^{d-1} k^2 \right) t^{d-3} \end{aligned}$$

where  $\approx$  means equality up to a polynomial (in  $t$ ) of order  $d-4$ . To finish the proof, we can now add the coefficients of  $I(t)$  and  $J(t)$ , and use the identities

$$S(d) = \frac{d-2}{2}d + \frac{d}{2} - \left\lfloor \frac{d}{2} \right\rfloor \quad \text{and} \quad \sum_{j=1}^{d-1} j^2 = \frac{d^3}{3} - \frac{d^2}{2} + \frac{d}{6}$$

to obtain (3.5). ■

#### ACKNOWLEDGEMENTS

The authors thank Professor Heinz Siedentop for calling attention to the relation between the second coefficient in Remark 1.3 and the Scott correction term in  $d=3$ , which lead us to Theorem 1.4.

T.C. Corso acknowledges funding by the *Deutsche Forschungsgemeinschaft* (DFG, German Research Foundation) - Project number 442047500 through the Collaborative Research Center "Sparsity and Singular Structures" (SFB 1481).

#### REFERENCES

- [AL78] M. AIZENMAN and E.H. LIEB, On semi-classical bounds for eigenvalues of Schrödinger operators. *Physics Letters A* **66**, (1978), no.6, 427–429. MR 0598768.
- [CCR24] T.C. CORSO and T. RIED, *On a variational problem related to the Cwikel-Lieb-Rozenblum and Lieb-Thirring inequalities*. arxiv 2403.04347
- [Cor24] T.C. CORSO, *A generalized three lines lemma in Hardy-like spaces*. arxiv 2407.10117
- [Cwi77] M. CWIKEL, Weak Type Estimates for Singular Values and the Number of Bound States of Schrodinger Operators. *Annals of Mathematics* **106**, (1977), no.1, 93–100. MR 0473576. Zbl 0362.47006.
- [FLW23] R.L. FRANK, A. LAPTEV, and T. WEIDL, *Schrödinger operators: eigenvalues and Lieb-Thirring inequalities*. Cambridge Studies in Advanced Mathematics Vol. 200, Cambridge University Press, Cambridge, 2023. MR 4496335. Zbl 07595814.
- [Fra21] R.L. FRANK, *The Lieb–Thirring inequalities: recent results and open problems*. In *Nine mathematical challenges—an elucidation* (eds. A. Kechris, N. Makarov, D. Ramakrishnan, X. Zhu), Proceedings of Symposia in Pure Mathematics **104**, American Mathematical Society, Providence, RI, 2021, pp. 210–235. MR 4337417. Zbl 1518.35002.
- [GGM78] V. GLASER, H. GROSSE, and A. MARTIN, Bounds on the number of eigenvalues of the Schrödinger operator. *Communications in Mathematical Physics* **59** (1978), 197–212 MR 491613. Zbl 0373.35050.
- [Lie80] E.H. LIEB, The number of bound states of one-body Schroedinger operators and the Weyl problem. In *Geometry of the Laplace operator*, Proceedings of Symposia in Pure Mathematics Vol. 36, American Mathematical Society, Providence, RI, 1980, 241–252. MR 573436, Zbl 0445.58029
- [LT76] E.H. LIEB and W. THIRRING, Bounds on the eigenvalues of the Laplace and Schrödinger operators. *Studies in Mathematical Physics: Essays in Honor of Valentine Bargmann (Princeton Series in Physics)* (1976), 269–303. Zbl 0342.35044.
- [Roz76] G.V. ROZENBLUM, Distribution of the discrete spectrum of singular differential operators (Russian). *Izv. Vysš. Učebn. Zaved. Matematika*, (1976), no.1, 75–86. MR 0430557. Zbl 0249.35069.
- [Sel24] A.D. SELVI, Cwikel–Lieb–Rozenblum inequalities for the Coulomb Hamiltonian. *Analysis and Mathematical Physics* **14**, (2024). MR 4701844. Zbl 1535.35046.

- [Sol96] A.V. SOBOLEV, Discrete spectrum asymptotics for the Schrödinger operator in a moderate magnetic field. *Reviews in Mathematical Physics* **8**, (1996), no.6, 861–903. [MR 1365349](#). [Zbl 0843.35093](#).
- [SW87] H. SIEDENTOP and R. WEIKARD, On the Leading Energy Correction for the Statistical Model of the Atom: Interacting Case. *Communications in Mathematical Physics* **112**, (1987), no.3, 471–491. [MR 0908549](#). [Zbl 0920.35120](#).
- [Wei96] T. WEIDL, On the Lieb-Thirring constants  $L_{\gamma, 1}$  for  $\gamma \geq 1/2$ . *Communications in Mathematical Physics* **178** No.1 (1996), 135–146. [MR 2714989](#). [Zbl 0858.34075](#).

(T. Carvalho Corso) INSTITUTE OF APPLIED ANALYSIS AND NUMERICAL SIMULATION, UNIVERSITY OF STUTTGART, PFAFFENWALDRING 57, 70569 STUTTGART, GERMANY  
*Email address:* thiago.carvalho-corso@mathematik.uni-stuttgart.de

(T. Weidl) INSTITUTE OF ANALYSIS, DYNAMICS AND MODELING, UNIVERSITY OF STUTTGART, PFAFFENWALDRING 57, 70569 STUTTGART, GERMANY  
*Email address:* timo.weidl@mathematik.uni-stuttgart.de

(Z. Zeng) INSTITUTE OF APPLIED ANALYSIS AND NUMERICAL SIMULATION, UNIVERSITY OF STUTTGART, PFAFFENWALDRING 57, 70569 STUTTGART, GERMANY  
*Email address:* zhuoyao.zeng@mathematik.uni-stuttgart.de