

Confidence regions for the multidimensional density in the uniform norm based on the recursive Wolverton-Wagner estimation

Maria Rosaria Formica ¹, Eugeny Ostrovsky ² and Leonid Sirota ³

¹ Università degli Studi di Napoli “Parthenope”, via Generale Parisi 13,
Palazzo Pacanowsky, 80132, Napoli, Italy.

e-mail: mara.formica@uniparthenope.it

^{2,3} Bar-Ilan University, Department of Mathematics and Statistics,
52900, Ramat Gan, Israel.

e-mail: eugostrovsky@list.ru

e-mail: sirota3@bezeqint.net

Abstract

We construct an optimal *exponential tail decreasing* confidence region for an unknown density of distribution in the Lebesgue-Riesz as well as in the *uniform* norm, built on the sample of the random vectors based of the famous *recursive* Wolverton-Wagner density estimation.

Keywords:

Probability space, random variables and processes (fields), uniform and Lebesgue - Riesz norms and spaces, density, bias, Wolverton - Wagner and Parzen - Rosenblatt estimations, Rosenthal’s inequality, sample, independence, ordinary Euclidean space, loss function, Chernoff’s exponential tail estimate, Lebesgue - Riesz norm and spaces, smooth functions, distribution and tail of distribution, confidence regions, Young - Fenchel (Legendre) transform, conditions of orthogonality, bias and variation, convex domain, Lipschitz condition, moments, kernel, windows or bandwidth, multivariate Banach spaces of random variables, exponential estimates, Grand Lebesgue Spaces.

1 Introduction

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a non-trivial probability space with an expectation \mathbf{E} and a variance \mathbf{Var} . Let also $\xi_1, \xi_2, \dots, \xi_k, \dots, \xi_n$; $k \in \mathbb{N} = 1, 2, \dots, n$; $n \geq 2$ be a *sample*, i.e. finite sequence of independent, identical distributed, (i, i.d.) random vectors (r.v.) taking the values in the ordinary Euclidean space $\mathbb{R}^d, d \geq 1$, equipped with usually distance function $\|x - y\| = (x - y, x - y)^{1/2}$, $x, y \in \mathbb{R}^d$, and having an unknown distribution *density* function $f = f(x)$, $x \in \mathbb{R}^d$. C. Wolverton and T.J. Wagner in [34] introduced the famous statistical estimation $f_n^{WW}(x) = f_n(x)$ for the function $f(\cdot)$ as follows.

Let $\{h_k\}$, $k \in \mathbb{N}$, be a positive deterministic sequence of real numbers such that $\lim_{k \rightarrow \infty} h_k = 0$, $h_1 = 1$, (*windows, or bandwidth*). Let also $K = K(x)$, $x \in \mathbb{R}^d$, be certain *kernel*, i.e. a measurable numerical valued fixed "*sufficiently smooth*," see an exact definition further, even normalized function, i.e. for which

$$\int_{\mathbb{R}^d} K(x) dx = 1.$$

Recall that following definitions.

Definition of the Wolverton - Wagner recursive estimate, see [34]; as well as [20].

$$f_n^{WW}(x) = f_n(x) \stackrel{def}{=} \frac{1}{n} \sum_{k=1}^n \frac{1}{h_k^d} K\left(\frac{x - \xi_k}{h_k}\right). \quad (1)$$

Definition of the Parzen - Rosenblatt (or Kernel) estimate (see, e.g., [25, 29]).

$$g_n^{PR}(x) = g_n(x) \stackrel{def}{=} \frac{1}{n} \frac{1}{h_n^d} \sum_{k=1}^n K\left(\frac{x - \xi_k}{h_n}\right). \quad (2)$$

Note that the Wolverton - Wagner estimate obeys a very important recursion property:

$$f_{n+1}^{WW}(x) = \frac{n}{n+1} f_n^{WW}(x) + \frac{h_{n+1}^{-d}}{n+1} \cdot K\left(\frac{x - \xi_{n+1}}{h_{n+1}}\right).$$

The recurrent definition of probability density estimates $f_n^{WW}(x)$ has two obvious advantages: 1) there is no need to memorize data, i.e. if the estimate $f_n^{WW}(x)$ is known, then the following one $f_{n+1}^{WW}(x)$ can be calculated by means of the last observation f_n^{WW} only, without using the sampling $\xi_1, \xi_2, \dots, \xi_n$; 2) the asymptotic variation as well as *bias* of the estimate $f_n^{WW}(x)$ does not exceed the asymptotic variation and bias of the estimate $g_n^{PR}(x)$, see e.g. [4, 5, 9]. This proposition holds true still when the error of estimation is understood in the classical Lebesgue - Riesz norm sense

$$R_p[f, f_n] \stackrel{def}{=} \mathbf{E} \left\{ \int_{\mathbb{R}^d} |f_n^{WW}(x) - f(x)|^p dx \right\}^{1/p}, \quad p \in [1, \infty).$$

The *loss function* is understood in this report in an *uniform norm* deviation

$$R_{\Psi, B_n, D}[f, f_n] \stackrel{def}{=} \mathbf{E} \Psi \left[B_n \sup_{x \in D} |f_n^{WW}(x) - f(x)| \right], \quad (3)$$

where $\Psi(\cdot)$ is certain *weight* function; indeed some non - negative Young function.

This means by definition that this function is such that $\Psi : [0, \infty) \rightarrow [0, \infty)$, is *strictly* increasing, continuous, convex, and

$$\Psi(0) = 0; \quad \lim_{x \rightarrow \infty} \Psi(x) = \infty.$$

HERE AND HEREAFTER D IS FIXED NON - EMPTY CLOSED CONVEX BOUNDED SUBDOMAIN (COMPACT) SUBSET OF THE WHOLE SPACE R^d .

B_n is some deterministic normed positive numerical sequence such that $\lim_{n \rightarrow \infty} B_n = \infty$. For instance, $\Psi(y) = |y|^m$, $m = \text{const} > 1$ or $\Psi(y) = \exp(|y|^m) - 1$, $m = \text{const} > 0$.

Our aim in this report is to deduce the exact exponential decreasing estimate for the tail of deviation probability $S^{WW}(u) = S_{D, n, d}^{WW}(u) \stackrel{def}{=}$

$$\mathbf{P}(B_n \sup_{x \in D} |f_n^{WW}(x) - f(x)| > u); \quad u > u_0 = \text{const} > 0, \quad (4)$$

of course under appropriate restrictions and for exact *optimal* deterministic positive normed numerical sequence B_n , tending to infinity.

To be more precise, one can suppose that the function $f(\cdot)$ belongs to one or another smoothness class of functions $U = \{f\}$. Of course, the *optimal* norming sequence B_n depends on U : $B_n = B_n(U)$.

For the classical Parzen - Rosenblatt estimate $f_n^{PR}(x)$ analogous results were obtained, e.g., in [8, 9], [22, Chapter 5, section 5.2].

2 Auxiliary considerations.

We must introduce several notations and definitions. Let β be a fixed positive constant: $\beta = \text{const} \in (0, \infty)$. Denote by $[\beta]$ its integer part and correspondingly by $\{\beta\}$ its fraction part $\{\beta\} = \beta - [\beta]$. Define as ordinary the following (very popular) functional class $\Sigma(\beta, L)$ consisting on all the continuous bounded functions such that all its partial derivatives of (integer vector) order $\vec{\alpha} = \{\alpha_1, \alpha_2, \dots, \alpha_d\}$, where

$$|\alpha| \stackrel{\text{def}}{=} \sum_{j=1}^d \alpha_j \leq [\beta], \quad (5)$$

are bounded and satisfy the Hölder condition with degree $\{\beta\}$: $\exists L = \text{const} < \infty$ such that, for all the values α from the set (5),

$$\left| \frac{D^{|\alpha|} f(x)}{\prod_{j=1}^d \partial^{\alpha_j} x_j} - \frac{D^{|\alpha|} f(y)}{\prod_{j=1}^d \partial^{\alpha_j} y_j} \right| \leq L \|x - y\|^{\{\beta\}}. \quad (6)$$

In the case when β is integer: $\{\beta\} = 0$, in the right hand of (6) must stay $L \|x - y\|$; the so - called Lipschitz condition.

Set also $\Sigma(\beta) := \cup_{L>0} \Sigma(\beta, L)$. Obviously, the set $\Sigma(\beta)$ forms a Banach functional space.

WE SUPPOSE FURTHER THAT THE UNKNOWN DENSITY FUNCTION BELONGS TO THE SET $\Sigma(\beta)$, or equally

$$\exists \beta \geq 0, \exists L \in (0, \infty) : f(\cdot) \in \Sigma(\beta, L). \quad (7)$$

See the following works devoted to the consistent measurement (estimation) of the parameters (β, L) [10, 14, 1, 2, 4] etc.

Let us now impose also several conditions on the (measurable) kernel - function $K(\cdot)$.

$$K(x) = K(-x); \int_{\mathbb{R}^d} K(x) dx = 1; \int_{\mathbb{R}^d} K^2(x) dx < \infty, K(\cdot) \in L_1(\mathbb{R}^d) \cap C(\mathbb{R}^d);$$

$$\exists \delta = \text{const} \in (0, 1], \exists c = \text{const}(0, \infty) : |K(x) - K(y)| \leq c|x - y|^\delta.$$

The following imposed by us important conditions (of orthogonality) are needed only by the bias estimation: for arbitrary non negative integer vector $l = \vec{l} \in \mathbb{R}^d$ such that $|l| \leq [\beta]$ we suppose that

$$\int_{\mathbb{R}^d} x^l K(x) dx = 0; \quad x^l \stackrel{\text{def}}{=} \prod_{j=1}^d x_j^{l_j}. \quad (8)$$

$$|x| := \sqrt{\sum_{j=1}^d x_j^2}.$$

The optimal chooses (in different senses) of the kernel $K(\cdot)$ is investigated in the report [9].

So, we allow consider in general case as a capacity of this kernels alternating ones.

Let us discuss about the choice of the windows (bandwidth) $\{h_k\}$. For the classical Parzen - Rosenblatt estimate h_k depends only on n :

$$h_n \stackrel{\text{def}}{=} c \left\{ \frac{\ln n}{n} \right\}^{\beta/(2\beta+d)}, \quad k = 2, 3, \dots, n,$$

the asymptotical optimal choice.

Let us return to the Wolverton - Wagner estimate.

Pick then for the definiteness $h_1 = 1$. The asymptotically as $n \rightarrow \infty$ optimal in the uniform norm $\sup_x |f_n(x) - f(x)| = \|f_n - f\|_C$ the Wolverton - Wagner estimates $f_n(\cdot) = f_n^{WW}(\cdot)$ of the density $f(\cdot)$ values of the windows (bandwidth) $\{h_k\}$ under formulated above restrictions have the following form.

$$h_k \stackrel{\text{def}}{=} c_1 \left\{ \frac{\ln k}{k} \right\}^{\beta/(2\beta+d)}, \quad k = 2, 3, \dots, n; \quad (9)$$

see e.g. [1], [2], [10] etc.

So, we choose furthermore the values $\{h_k\}$ in accordance with the relation (9) for the Wolverton - Wagner estimate.

Introduce the following variables $B_1 = 1$,

$$B_n = B_n(\beta, d) \stackrel{\text{def}}{=} \left[\frac{n}{\ln n} \right]^{\beta/(2\beta+d)} \asymp 1/h_n, \quad n \geq 2;$$

$$Q_n(u) = Q_n(d; \beta, L; u) \stackrel{\text{def}}{=} \mathbf{P} \left(B_n \cdot \sup_{x \in D} |f_n^{WW}(x) - f(x)| > u \right), \quad u \geq e.$$

Notice that under formulated above notations and conditions

$$\sup_n \sup_{x \in D} B_n(\beta, d) | \mathbf{E} f_n^{WW}(x) - f(x) | < \infty,$$

therefore it is sufficient to investigate only the variable

$$Q_n^o(u) = Q_n^o(d; \beta, L; u) \stackrel{\text{def}}{=} \mathbf{P} \left(B_n \cdot \sup_{x \in D} |f_n^{WW}(x) - \mathbf{E} f_n^{WW}(x)| > u \right), \quad u \geq e.$$

3 Main result.

Let (Z, \mathcal{Z}, ν) be arbitrary measurable space equipped with a sigma - finite measure ν . Denote as ordinary the classical Lebesgue - Riesz space

$L_{p,Z,\nu} = L_p(Z)$, $p \in [1, \infty)$ as a set of all the measurable numerical valued functions $h : Z \rightarrow \mathbb{R}$ having a finite norm

$$\|h\|_{p;Z,\nu} \stackrel{def}{=} \left[\int_Z |h(z)|^p \nu(dz) \right]^{1/p}.$$

In particular, let $\zeta : \Omega \rightarrow \mathbb{R}$ be an arbitrary measurable function (random variable), denote as usual the corresponding Lebesgue - Riesz $L_p = L_p(\Omega)$ norm

$$\|\zeta\|_p \stackrel{def}{=} [\mathbf{E}|\zeta|^p]^{1/p}, \quad p \in [2, \infty).$$

We conclude under our notations and restrictions

$$\sup_{x \in D} \| |f_n^{WW}(x) - f(x)| \|_p \leq C_1(\beta, L, d, D) \times \frac{p}{\ln p} \times \quad (10)$$

$$\left(\frac{\ln n}{n} \right)^{\beta/(2\beta+d)}, \quad n \geq 2; \quad C_1 \stackrel{def}{=} C_1(\beta, L, d) \in (0, \infty).$$

Proof.

It follows immediately, quite similarly as in [8], from the classical Rosenthal inequality for the moment for sums of independent centered r.v. ([30]); see also [11, 12, 13].

The exact value of the correspondent constant is derived in [21].

□

Remark 3.1. The relation (10) may be rewritten on the language of the so - called Grand Lebesgue Spaces (GLS), see e.g. [7, 16, 17, 19, 22, 23, 24].

Namely, let $Y(\cdot)$ be an arbitrary measurable numerical valued function, for instance a random variable and, for $1 < a < b \leq \infty$, let $\psi(p)$, $p \in (a, b)$, be a strictly positive numerical valued continuous function. The norm in the Grand Lebesgue spaces $G\psi(a, b)$ is defined by

$$\|Y\|_{G\psi(a,b)} \stackrel{def}{=} \sup_{p \in (a,b)} \left\{ \frac{\|Y\|_p}{\psi(p)} \right\}. \quad (11)$$

Let for instance ξ be a r.v. such that $\|\xi\|_{G\psi} \leq \kappa$, $0 < \kappa < \infty$. Define the auxiliary function $h(p) = h[\psi](p) := p \ln \psi(p)$. and introduce its Young - Fenchel, or Legendre transform

$$h^*(t) = h^*[\psi](t) \stackrel{def}{=} \sup_{p \geq 2} (pt - h[\psi](p)), \quad t \geq 1.$$

Another name: conjugate function.

It is well known the following Chentzov's inequality

$$T[\xi](t) \leq \exp(-h^*(t/\kappa)), \quad t \geq \kappa.$$

If for example $\psi(p) = \psi_l(p) \stackrel{def}{=} \frac{p}{\ln p}$, $p \geq 2$, then

$$h^*[\psi_l](t) \sim t + \ln t \cdot \ln \ln t, \quad t \geq e^e. \quad (12)$$

Define the function

$$\psi_l(p) \stackrel{def}{=} \frac{p}{\ln p}, \quad p \in (2, \infty);$$

then

$$B_n(\beta, d) \sup_{x \in D} \|f_n^{WW}(x) - f(x)\|_{G\psi_l(2, \infty)} \leq C_1(\beta, L, d, D). \quad (13)$$

This proposition follows from the last relation (13) a corresponding tail estimation, see e.g. [7, 22]. Define, as ordinarily, for an arbitrary numerical valued random variable (r.v.) η , its tail function

$$T_\eta(t) \stackrel{def}{=} \mathbf{P}(|\eta| \geq t), \quad t \geq t_0 = \text{const} > 0.$$

Introduce also the following family of random fields

$$\zeta_n(x) := B_n(\beta, d) \cdot \{ f_n^{WW}(x) - f(x) \}; \quad z := t/e, \quad z \geq 1;$$

$$\nu(z) \stackrel{def}{=} \exp(-z) \cdot \exp(-\ln z \cdot \ln \ln z);$$

then

$$\sup_n \sup_x T_{\zeta_n, x}(t) \leq C_1 \nu(z) = C_1 \nu(t/e). \quad (14)$$

Further, we deduce analogously as in (13),

$$\begin{aligned} & B_n(\beta, d) \cdot \|(f_n^{WW}(x) - f(x)) - (f_n^{WW}(y) - f(y))\|_{G\psi_l(2, \infty)} \\ & \leq C_2(\beta, L, d) \times \|x - y\|/h_n, \quad x, y \in \mathbb{R}^d. \end{aligned} \quad (15)$$

It follows from the well - known theory of Grand Lebesgue Spaces (GLS), see e.g. [7], [22], chapter 5, section 5.4 - 5.5, pages 253 - 264, the following result.

Theorem 3.1.

Let D be a non - empty compact subset on the whole domain \mathbb{R}^d . We propose under all the formulated above conditions and notations

$$\mathbf{P} \left(B_n(\beta, d) \cdot \sup_{x \in D} |f_n^{WW}(x) - f(x)| > t \right) \leq C_{12}(D, \beta, d, L) \nu(t/e), \quad t \geq e, \quad (16)$$

where as ordinary $C_{12}(D, \beta, d, L) < \infty$.

The **proof** is completely alike to the one for the classical Parzen - Rosenblatt estimate, see [22], chapter 5, section 5.4. It used a famous method belonging to R.Reiss [28] of partition of the set D on the subsets of the variable small volumes.

In detail, introduce an auxiliary numerical valued *centered* random process (field), (r.f.), more precisely, a family ones

$$\zeta_n(x) \stackrel{def}{=} B_n(\beta, d) \times [f_n^{WW}(x) - \mathbf{E}f_n^{WW}(x)], \quad n = 1, 2, \dots; \quad x \in D.$$

We must estimate the tail of maximum distribution for this r.f.

$$\mathbf{P}_z(u) \stackrel{def}{=} \mathbf{P} \left(\sup_{x \in D} |\zeta_n(x)| > u \right),$$

of course, for all sufficiently greatest values u , say, for $u \geq e$.

One can assume without loss of generality that the set D is unit "cube" $D = [0, 1]^d$. Put now $q = (2\beta + d)/(\beta + d)$. Introduce also the following system of cubes $\vec{k} = \{k_1, k_2, \dots, k_d\}$;

$$L(\vec{k}) := \otimes_{j=1}^d \left\{ \left[\frac{k_j}{n}, \frac{k_j + 1}{n} \right) \right\},$$

where $j = 1, 2, \dots, d$; $k_j = 0, 1, 2, \dots, n - 1$; which covers the whole set D .

Define also the functions

$$\mathbf{P}_a(u) \stackrel{\text{def}}{=} \max_{\vec{k}} \mathbf{P} \left[\sup_{x \in L(\vec{k})} B_n |f_n(x) - f(x)| > u \right].$$

Further, we deduce quite analogously as in (13)

$$B_n(\beta, d) \cdot \| [f_n^{WW}(x) - \mathbf{E}f_n^{WW}(x)] - [f_n^{WW}(y) - \mathbf{E}f_n^{WW}(y)] \|_{G\psi_1(2, \infty)} \leq$$

$$C_4(\beta, L, d) \times \|x - y\|/h_n, \quad x, y \in \mathbb{R}^d. \quad (17)$$

Notice that the metric entropy $H(D, \rho, \epsilon)$ of the set D relative the distance function $\rho(x, y) := \|x - y\|/h_n$ may be estimated as follows

$$H(D, \rho, \epsilon) \leq C_1 + C_2(d) \ln n + C_3 |\ln \epsilon|, \quad \epsilon \in (0, 1).$$

Note also that it is sufficient for this purpose to ground the estimate of the form (16) function for the alike but for the following *centered* correspondent random variables

Consider for definiteness the first cube $Q \stackrel{\text{def}}{=} [0, 1/n]^d$, i.e. when $k_j = 0, 1; \dots, j = 1, 2, \dots, d$. It follows immediately from the Rosenthal's moments estimates for the sums of the independent centered r.v. - s, see e.g. [12], [21], [30], that for $x \in Q$ and $u \geq e$

$$\mathbf{P}_o(u; x) \stackrel{\text{def}}{=} \mathbf{P} [B_n |f_n(x) - \mathbf{E}f_n(x)| > u] \leq \exp [-C_7 \nu(u/V(n))]; \quad (18)$$

$$\mathbf{P}_a^o(u) \stackrel{def}{=} \max_{\vec{k}} \mathbf{P} \left[\sup_{x \in L(\vec{k})} B_n(f_n(x) - \mathbf{E}f_n(x)) > u \right]. \quad (19)$$

Notice that

$$\max_{x \in Q} \|\zeta_n(x)\|_{G\psi_l(2, \infty)} \leq \frac{C_6}{n h_n}, \quad (20)$$

$$\|\zeta_n(x) - \zeta_n(y)\|_{G\psi_l(2, \infty)} \leq \frac{C_8 \|x - y\|}{h_n}, \quad (21)$$

and that when $n \geq 2$

$$V(n) \stackrel{def}{=} n h_n \sim C_7 n^{(\beta+d)/(2\beta+d)} (\ln n)^{\beta/(2\beta+d)} \rightarrow \infty, \quad n \rightarrow \infty.$$

We propose attracting the main result of the section 3.4 of the monograph [22] that

$$\mathbf{P}_a^o(u) \leq \exp (C_4 \ln n - \nu(uV(n))), \quad u \geq u_0 = \text{const} \geq e. \quad (22)$$

Following we deduce after mild calculations, when $u > e$

$$\begin{aligned} \mathbf{P}_z(u) &\leq \sum_{\vec{k}} \exp (C_5(\beta, d, L) \ln n - \nu(u V(n))) \leq \\ &n^d \exp (C_6(\beta, d, L) \ln n - \nu(u V(n))) \leq \\ &\sup_n \{ n^d \exp (C_6(\beta, d, L) \ln n - \nu(u V(n))) \} \leq \end{aligned}$$

$$C_9(D, \beta, d, L) \exp (- \nu(u/C_{10}(D, \beta, d, L))), \quad C_{9,10}(D, \beta, d, L) \in (0, \infty).$$

Ultimately, the proposition (16) is proved.

Acknowledgements. M.R. Formica is member of Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di

Alta Matematica (INdAM) and member of the UMI group “Teoria dell’Approssimazione e Applicazioni (T.A.A.)” and is partially supported by the INdAM-GNAMPA project, *Risultati di regolarità per PDEs in spazi di funzione non-standard*, codice CUP_E53C22001930001.

References

- [1] **K. Bertin**, *Asymptotically exact minimax estimation in sup-norm for anisotropic Hölder classes*. Bernoulli, **10**, (2004), 873–888.
- [2] **K. Bertin**, *Estimation asymptotiquement exacte en norme sup de fonctions multidimensionnelles*. Ph.D. thesis, University Paris 6, (2004).
- [3] **V.V. Buldygin and Yu.V. Kozachenko**, *Metric Characterization of Random Variables and Random Processes*. 1998, Translations of Mathematics Monograph, AMS, v.188.
- [4] **L. Devroye**, *On the pointwise and integral convergence of recursive kernel estimates of probability densities*. Util. Math., **15**, (1979), 113–128.
- [5] **L. Devroye, L. Györfi**, *Nonparametric density estimation: The L_1 View*. Wiley, New York, 1975.
- [6] **S.W. Dharmadhikari, K. Jogdeo**, *Bounds on the moment of certain random variables*. Ann. Math. Statist., **40**, no. 4 (1969), 1506–1508.
- [7] **S. V. Ermakov, and E. I. Ostrovsky**. *Continuity Conditions, Exponential Estimates, and the Central Limit Theorem for Random Fields*. Moscow, VINITI, 1986. (in Russian).
- [8] **M.R. Formica, E. Ostrovsky and L. Sirota**, *Exponential confidence interval based on the recursive Wolverton - Wagner density estimation*. arXiv:2102.07867 [math.ST] 15 Feb 2021
- [9] **M.R. Formica, E. Ostrovsky and L. Sirota**, *Signed variable optimal kernel for non - parametric density estimation*. arXiv:2102.07858v1 [math.ST] 15 Feb 2021
- [10] **A. Goldenshluger, O. Lepski**, *Bandwidth selection in kernel density estimation: oracle inequalities and adaptive minimax optimality*. Ann Stat., **39** (3) (2011), 1608–1632.
- [11] **I.A. Ibragimov and R.Z. Has’minskii**, *Statistical estimation*, Vol. 16, Applications of Mathematics. Springer-Verlag, New York, 1981. (Asymptotic theory, Translated from the Russian by Samuel Kotz).
- [12] **R. Ibragimov, Sh. Sharachmedov**, *On the exact constant in the Rosenthal Inequality*. Theory Probab. Appl., **42** (1997), 294–302.
- [13] **R. Ibragimov, Sh. Sharachmedov**, *The exact constant in the Rosenthal Inequality for random variables with mean zero*. Probab. Theory Appl., **46**, no. 1, (2001), 127–132.

- [14] **S. Khardani, Y. Slaoui**, *Recursive Kernel Density Estimation and Optimal Bandwidth Selection Under α : Mixing Data*. J. Stat. Theory Pract., **13**, 36 (2019).
<https://doi.org/10.1007/s42519-018-0031-6>
- [15] **A. Korostelev, M. Nussbaum**, *The asymptotic minimax constant for sup-norm loss in nonparametric density estimation*. Bernoulli, **5** (6) (1999), 1099–1118.
- [16] **Yu. V. Kozachenko and E. I. Ostrovsky**, *The Banach Spaces of random variables of sub-Gaussian type*. Probability and Mathematical Statistics, **32**, (1985), (in Russian). Kiev, KSU, 43–57.
- [17] **Yu.V. Kozachenko, Yu.Yu. Mlavets and N.V. Yurchenko**, *Weak convergence of random processes from spaces $F_\psi(\Omega)$* . Stat. Optim. Inf. Comput. **6** (2018), no. 2, 266–277.
- [18] **R. Latala**, *Estimation of moments of sums of independent real random variables*. Ann. Probab., **25** (3) (1997), 1502–1513.
- [19] **E. Lifyand, E. Ostrovsky and L. Sirota**, *Structural properties of bilateral grand Lebesgue spaces*. Turkish J. Math. **34** (2010), no. 2, 207–219.
- [20] **E. Nadaraya, P. Babilua**, *On the Wolverton-Wagner Estimate of a Distribution Density*. Bull. Georgian Natl. Acad. Sci., **175** (2007), 40–44.
- [21] **B. Naimark, E. Ostrovsky**, *Exact Constants in the Rosenthal Moment Inequalities for Sums of independent centered Random Variables*. arXiv:math/0411614 [math.PR] 27 Nov 2004
- [22] **E. Ostrovsky**, *Exponential estimates for random fields and its applications*. 1999, OINPE, Moscow - Obninsk. (In Russian).
- [23] **E. Ostrovsky and L. Sirota**, *Moment Banach spaces: theory and applications*. HIAT Journal of Science and Engineering, C, 2011, Volume 4, Issues 1 - 2, pp. 233–262, (2007).
- [24] **E. Ostrovsky and L. Sirota**, *Vector rearrangement invariant Banach spaces of random variables with exponential decreasing tails of distributions*. arXiv:1510.04182v1 [math.PR] 14 Oct 2015
- [25] **E. Parzen**, *On estimation of a probability density and mode*. Ann. Math. Stat., **33** (1962), 1065–1076.
- [26] **I.S. Pinelis and S.A. Utev**, *Estimates of moments of sums of independent random variables*, Theory Probab. App., **29** (3) (1985), 574–578.
- [27] **G. Pisier**, *Conditions d'entropie assurant la continuité de certains processus et applications à l'analyse harmonique*. (French) Seminaire d'analyse fonctionnelle (1980), Exp. No. 13-14, pp. 43–46.
- [28] **R. Reiss**, *On the deviation empirical density function*. Metrica, 1981, B.22, 107–120.
- [29] **M. Rosenblatt**, *Remarks on some nonparametric estimates of a density function*. Ann. Math. Stat., **27** (1956), 832–837.

- [30] **H.P. Rosenthal**, *On the subspaces of L^p ($p > 2$) spanned by sequences of independent variables*. Israel J. Math., **8** (1970), 273–303.
- [31] **A.B. Tsybakov**, *Recurrent estimation of the mode of a multidimensional distribution*. Probl. Inf. Transm., **8** (1990), 119–126.
- [32] **A.B. Tsybakov**, *Introduction to Nonparametric Estimation*. Springer, 2009.
- [33] **E.J. Wegman and H.I. Davies**, *Remarks on Some Recursive Estimators of a Probability Density*. The Annals of Statistics, **7** (1979), 316–327.
- [34] **C. Wolverton, T.J. Wagner**, *Asymptotically optimal discriminant functions for pattern classification*. IEEE Trans Inform Theory, **15** (1969), 258–265.