

# Configurational entropy and stability conditions of fermion and boson stars

P.S. Koliogiannis<sup>1,2,\*</sup>, M. Vikiaris<sup>2,†</sup>, C. Panos<sup>2,‡</sup>, V. Petousis<sup>3,§</sup>, M. Veselský<sup>3,¶</sup> and Ch.C. Moustakidis<sup>2,\*\*</sup>

<sup>1</sup>*Department of Physics, Faculty of Science, University of Zagreb, Bijenička cesta 32, 10000 Zagreb, Croatia*

<sup>2</sup>*Department of Theoretical Physics, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece*

<sup>3</sup>*Institute of Experimental and Applied Physics, Czech Technical University, Prague, 110 00, Czechia*

In a remarkable study by M. Gleiser and N. Jiang (Phys. Rev. D **92**, 044046, 2015), the authors demonstrated that the stability regions of neutron stars, within the framework of the simple Fermi gas model, and self-gravitating configurations of complex scalar field (boson stars) with various self couplings, obtained through traditional perturbation methods, correlates with critical points of the configurational entropy with an accuracy of a few percent. Recently, P. Koliogiannis *et al.* (Phys. Rev. D **107**, 044069 2023) found that while the minimization of the configurational entropy generally anticipates qualitatively the stability point for neutron stars and quark stars, this approach lacks universal validity. In this work, we aim to further elucidate this issue by seeking to reconcile these seemingly contradictory findings. Specifically, we calculate the configurational entropy of bosonic and fermionic systems, described by interacting Fermi and Boson gases, respectively, that form compact objects stabilized by gravity. We investigate whether the minimization of configurational entropy coincides with the stability point of the corresponding compact objects. Our results indicate a strong correlation between the stability points predicted by configurational entropy and those obtained through traditional methods, with the accuracy of this correlation showing a slight dependence on the interaction strength. Consequently, the stability of compact objects, composed of components obeying Fermi or Boson statistics, can alternatively be assessed using the concept of configurational entropy.

PACS numbers: 03.67.-a, 04.40.Dg, 97.60.Jd, 05.30.-d, 02.30.Nw

Keywords: Configurational entropy; stability condition; compact objects; equation of state

## I. INTRODUCTION

In recent years there has been an extensive interest in the study of astrophysical objects with the help of the concept of information entropy and related quantities. In particular, Sañudo and Pacheco [1] studied the relation between the complexity and the structure of white dwarfs. Later on, the aforementioned study has been applied in neutron star's structure [2] and it was found that the interplay between gravity, the short-range nuclear force, and the very short-range weak interaction shows that neutron stars, under the current theoretical framework, are ordered systems. Similar studies took place in the following years in a series of papers [3–7] and Herrera *et al.* [8–11] elaborated the definition of the complexity factors in self-gravitating systems, approaching the problem in a different way. Some additional applications of the concept of the information measures may be found in Refs. [12–20].

A more specific application of the information measure is the configurational entropy (CE). The concept of the CE has been introduced by Gleiser and Stamatopoulos [21] in order to study possible relation between the dynamical and information content of physics models with localized energy configurations. In the next years, the CE has been applied in several similar studies [22–32]. In a notable study, Gleiser and Jiang [23] investigated the connection between the stability of compact objects (white dwarfs, neutron stars and boson stars) and the

corresponding information-entropic measure. According to their notable finding, the minimization of the CE offers an alternative way to predict the stability condition through the maximum mass configuration, for a variety of stellar objects. It is worth noting that there is no theoretical argument (or proof) to relate the stability point to the minimum of the CE. However, it is intuitive to expect that since the maximum mass corresponds to the most compact configuration (maximum mass and minimum radius of a stable configuration), the corresponding CE will exhibit an extreme value (in this case a total minimum).

Recently in Ref. [24] we extended the aforementioned study in various compact objects including neutron stars, quark stars, and hybrid stars. Employing a large set of realistic equations of state (EoS), in each case, we found that the suggested prediction of the stability by the minimization of the CE, concerning neutron stars and quark stars, does not have, at least quantitatively, universal validity.

It is worth mentioning that the study of the longstanding problem of the stability of relativistic stars [33–39] is mainly carried out by the following three methods: (a) the method of locating the point that corresponds to the minimization of the binding energy defined as  $E_B = (M - m_b N)c^2$  (where  $m_b$  is the mass of a single nucleon,  $M$  stands for the gravitational mass, and  $N$  is the total number of nucleons) [33–39], (b) the variational method developed by Chandrasekhar [40, 41], and (c) the method based on the dependence of the gravitational mass  $M$  and the radius  $R$  on the central energy density  $\mathcal{E}_c$  (hereafter traditional method TM). The stability condition demands that the mass increases with increasing central energy density  $dM/d\mathcal{E}_c > 0$ . The extrema in the mass indicates a change in the stability of the compact star configuration [33–39].

In the present work we employ the third method in order to investigate a possible relation between the stability of a relativistic star and the corresponding CE. In this case, the ques-

\* pkoliogi@auth.gr

† mvikiari@auth.gr

‡ chpanos@math.auth.gr

§ Vlasios.Petousis@cvut.cz

¶ Martin.Veselsky@cvut.cz

\*\* moustaki@auth.gr

tions that arise and must be answered (or at least investigated) are the following: Is there any one-to-one correspondence between the minimum of the CE and the stability point for each realistic EoS? Is this rule universal or it depends on the specific character of each EoS? Is it possible, even in some special cases, to associate stability with minimization of CE, and if so, which is the underlying reason? Obviously, one can appreciate the importance of discovering new ways, beyond the classical ones, to find stability conditions for compact objects.

In view of the above questions, the main motivation of the present work is to provide an extended examination of the statements of Refs. [23] and [24]. In Ref. [23] the authors found that the stability regions of neutron stars (in the framework of the simple Fermi gas model) as well as of self-gravitating configurations of complex scalar field (boson stars) with various self couplings (detailed reviews in Refs. [42–44]), obtained from traditional perturbation methods, correlating the critical points of the CE with an accuracy of a few percent. In Ref. [24] the authors found that the suggested prediction of the stability by the minimization of the CE, concerning neutron stars and quark stars, does not have, at least quantitatively, universal validity although in several cases it qualitatively predicts the existence of the stability point. In this work, we aim to further elucidate this issue by seeking to reconcile these seemingly contradictory findings. Specifically, we calculate the configurational entropy of bosonic and fermionic systems, described by interacting Fermi and Boson gases, respectively, that form compact objects stabilized by gravity. We investigate whether the minimization of configurational entropy coincides with the stability point of the corresponding compact objects. Our results indicate a strong correlation between the stability points predicted by configurational entropy and those obtained through traditional methods, with the accuracy of this correlation showing a slight dependence on the interaction strength.

The paper is organized as follows. In Sec. II we present the basic formalism of the hydrodynamic equilibrium and the role of analytical solutions while in Sec. III, we review the definition of the configurational entropy. The parametrization of the equations of state is provided in Sec. IV and in Sec. V the results of the present study are laid out and discussed. Finally, Sec. VI contains the concluding remarks.

## II. HYDRODYNAMIC EQUILIBRIUM AND ANALYTICAL SOLUTIONS

To construct the related configuration in each compact object, which is the key property to calculate the CE, we employ the Einstein's field equations of a spherical fluid. In this case the mechanical equilibrium of the star matter is determined by the well known Tolman-Oppenheimer-Volkoff (TOV) equations [33–36]

$$\begin{aligned} \frac{dP(r)}{dr} &= -\frac{G\mathcal{E}(r)M(r)}{c^2 r^2} \left(1 + \frac{P(r)}{\mathcal{E}(r)}\right) \\ &\times \left(1 + \frac{4\pi P(r)r^3}{M(r)c^2}\right) \left(1 - \frac{2GM(r)}{c^2 r}\right)^{-1}, \quad (1) \end{aligned}$$

$$\frac{dM(r)}{dr} = \frac{4\pi r^2}{c^2} \mathcal{E}(r). \quad (2)$$

In general, to obtain realistic solutions, it is most natural to numerically solve the TOV equations by incorporating an EoS that describes the relationship between pressure and density within the fluid interior. Alternatively, one can seek analytical solutions to the TOV equations, though these solutions may lack physical relevance. While there are hundreds of analytical solutions to the TOV equations [45, 46], only a few are of significant physical interest. In this work, we employ two of these noteworthy solutions: the Schwarzschild (constant-density interior solution) and the Tolman VII solution [45, 46]. It is important to note that analytical solutions are highly valuable, as they often provide explicit expressions for the quantities of interest and are instrumental in verifying the accuracy of numerical calculations. Below, we briefly describe these two fundamental analytical solutions.

- **Schwarzschild solution:** In the case of the Schwarzschild interior solution, the density is constant throughout the star [37, 38]. The energy density and the pressure read as

$$\mathcal{E} = \mathcal{E}_c = \frac{3M}{4\pi R^3}, \quad (3)$$

$$\frac{P(x)}{\mathcal{E}_c} = \frac{\sqrt{1-2\beta} - \sqrt{1-2\beta x^2}}{\sqrt{1-2\beta x^2} - 3\sqrt{1-2\beta}}, \quad (4)$$

where  $x = r/R$ ,  $\beta = GM/Rc^2$  is the compactness of the star and  $\mathcal{E}_c = \rho_c c^2$  is the central energy density.

- **Tolman VII solution:** The Tolman VII solution has been extensively employed in neutron star studies while its physical realization has been examined, very recently, in detail [47, 48]. The stability of this solution has been examined by Negi *et al.* [49, 50] and also confirmed in Ref. [51]. The energy density and the pressure read as [51]

$$\frac{\mathcal{E}(x)}{\mathcal{E}_c} = (1 - x^2), \quad \mathcal{E}_c = \frac{15Mc^2}{8\pi R^3}, \quad (5)$$

$$\frac{P(x)}{\mathcal{E}_c} = \frac{2}{15} \sqrt{\frac{3e^{-\lambda}}{\beta}} \tan \phi - \frac{1}{3} + \frac{x^2}{5}. \quad (6)$$

It is worth mentioning that these solutions are applicable to any kind of compact object independently of the values of mass and radius. Consequently, they are suitable in studying any kind of massive or supramassive object which the hydrodynamic stability obeys to TOV equations. In any case, useful insight can be gained by the use of analytical solutions concerning both the qualitative and quantitative behavior of CE as a function of central density  $\rho_c$ .

## III. CONFIGURATIONAL ENTROPY IN MOMENTUM SPACE

The key quantity to calculate the CE in momentum space is the Fourier transform  $F(\mathbf{k})$  of the density  $\rho(r) = \mathcal{E}(r)/c^2$ ,

originating from the solution of the TOV equations, that is

$$\begin{aligned} F(\mathbf{k}) &= \int \int \int \rho(r) e^{-i\mathbf{k}\cdot\mathbf{r}} d^3\mathbf{r} \\ &= \frac{4\pi}{k} \int_0^R \rho(r) r \sin(kr) dr. \end{aligned} \quad (7)$$

It is notable that the function  $F(k)$  in the case of zero momentum, coincides with the gravitational mass of the compact object, that is  $F(0) \equiv M$ , since by definition (see Eq. (2))

$$M = 4\pi \int_0^R \rho(r) r^2 dr, \quad (8)$$

where the density  $\rho(r)$  derives from the solution of the TOV equations. Moreover, we define the modal fraction  $f(\mathbf{k})$  [23]

$$f(\mathbf{k}) = \frac{|F(\mathbf{k})|^2}{\int |F(\mathbf{k})|^2 d^3\mathbf{k}}, \quad (9)$$

and also the function  $\tilde{f}(\mathbf{k}) = f(\mathbf{k})/f(\mathbf{k})_{\max}$ , where  $f(\mathbf{k})_{\max}$  is the maximum fraction, which is given in many cases by the zero mode  $k = 0$ , or by the system's longest physics mode,  $|k_{\min}| = \pi/R$ . The above normalization guarantees that  $\tilde{f}(\mathbf{k}) \leq 1$  for all values of  $\mathbf{k}$ .

Finally, the CE,  $S_C$ , as a functional of  $\tilde{f}(\mathbf{k})$ , is given by

$$S_C[\tilde{f}] = - \int \tilde{f}(\mathbf{k}) \ln[\tilde{f}(\mathbf{k})] d^3\mathbf{k}. \quad (10)$$

Summarizing, for each EoS, an infinite number of configurations can appear, leading to the construction of  $M - \rho_c$  and  $S_C - \rho_c$  dependence. The latter facilitates the investigation of any possible correlation between the minimum of  $S_C$  and the stability point of compact objects under examination.

#### IV. EQUATION OF STATE

The present study focuses on the role of the CE as a stability condition of compact objects obeying Fermi or Boson statistics including a parameterized self-interaction. The compact objects under consideration are neutron stars (introduced by a simplified EoS), boson stars and other type of astrophysical objects composed of fermions or bosons, such as dark matter stars. In following, the equations of interacting Fermi and boson gases are introduced.

##### A. Interacting Fermi Gas (FG)

In the case of compact objects consisting of solely interacting Fermi gas (FG), we considered the simplest extension of the free fermion gas in which an extra term that introduces the repulsive interaction between fermions is added. Therefore, the energy density and pressure of the fermions are described

as (for an extensive analysis see Ref. [52])

$$\begin{aligned} \mathcal{E}(n_\chi) &= \frac{(m_\chi c^2)^4}{(\hbar c)^3 8\pi^2} \left[ x \sqrt{1+x^2} (1+2x^2) \right. \\ &\quad \left. - \ln(x + \sqrt{1+x^2}) \right] + \frac{y^2}{2} (\hbar c)^3 n_\chi^2, \end{aligned} \quad (11)$$

$$\begin{aligned} P(n_\chi) &= \frac{(m_\chi c^2)^4}{(\hbar c)^3 8\pi^2} \left[ x \sqrt{1+x^2} (2x^2/3 - 1) \right. \\ &\quad \left. + \ln(x + \sqrt{1+x^2}) \right] + \frac{y^2}{2} (\hbar c)^3 n_\chi^2, \end{aligned} \quad (12)$$

where  $m_\chi$  is the particle mass, which considered equal to  $m_\chi = 939 \text{ MeV}/c^2$  for reasons of simplicity (this holds throughout the study),  $n_\chi$  is the number density and

$$x = \frac{(\hbar c)(3\pi^2 n_\chi)^{1/3}}{m_\chi c^2}.$$

The last parameter,  $y$  (in units of  $\text{MeV}^{-1}$ ), is the one that introduces the repulsive interaction. In this study we have considered the values  $y = [0, 0.001, 0.005, 0.01, 0.05, 0.1, 0.3, 0.5]$  ( $\text{MeV}^{-1}$ ) where increasing  $y$  increases the strength of the interaction and vice-versa.

##### B. Interacting Boson Gas (BG)

To enrich the study, we included the case of compact stars composed of bosonic matter and specifically that of an interacting Boson gas. As the construction of the EoS for the aforementioned gas is not unambiguously defined, we introduce three cases based on different assumptions. It needs to be noted that since the scalar field only vanishes at spatial infinity, boson stars do not have a specific radius where the energy density and pressure vanish. Thus, we do not use a momentum cut-off scheme,  $0 \leq |\mathbf{k}| \leq \infty$ .

1. **BG - C1:** The first derivation of the EoS of boson stars with a repulsive interaction was given in Ref. [53] and since then it has been used extensively in the corresponding calculations. In particular, the energy density is given as

$$\mathcal{E}(P) = \frac{4}{3w} \left[ \left( \sqrt{\frac{9w}{4} P + 1} \right)^2 - 1 \right], \quad (13)$$

where  $w = 4\lambda(\hbar c)^3/(m_\chi c^2)^4$  (in units of  $\text{MeV}^{-1} \text{ fm}^3$ ). In fact, the parameter  $\lambda$  is the one that is related with the strength of the interaction. However, it is usual to employ the combination of  $m_\chi$  and  $\lambda$  defined as  $w$ . In this study we have considered the values  $w = [0.001, 0.005, 0.01, 0.05, 0.1, 0.3, 0.5]$  ( $\text{MeV}^{-1} \text{ fm}^3$ ) where increasing  $w$  increases the strength of the interaction and vice-versa.

2. **BG - C2:** The second way to describe the interior of a boson star is through the EoS provided in Ref. [54] and

used recently in Ref. [55], where the energy density is given by

$$\mathcal{E}(P) = P + \sqrt{\frac{2P}{z}}, \quad (14)$$

where the interaction parameter  $z$  is given by  $z = u^2(\hbar c)^3/(m_\chi c^2)^2$  (in units of  $\text{MeV}^{-1} \text{fm}^3$ ) and the quantity  $u = g_\chi/m_\phi c^2$  defines the strength of the interaction in analogy to the case of fermions. In this study we have considered the values  $z = [0.001, 0.005, 0.01, 0.05, 0.1, 0.3, 0.5]$  ( $\text{MeV}^{-1} \text{fm}^3$ ).

3. **BG - C3:** Recently, in Ref. [56] the authors studied the properties of self-interacting boson stars with different scalar potentials. They concluded that the resulting properties of the boson star configurations differ considerably from previous calculations. Therefore, to enhance the connection between the stability criterion and the CE, we employed two cases of the EoSs introduced in Ref. [56]: (a) one with a mass term (MT) and (b) one with a vacuum term (VT) without a mass term. The scaling EoSs read as

$$\mathcal{E}(P) = \begin{cases} P^{2/n} + (n+2)P/(n-2), & \text{MT,} \\ 1 + (n+2)P/(n-2), & \text{VT,} \end{cases} \quad (15)$$

where the index  $n$  is restricted to  $n > 2$ . In this study we have considered the values  $n = [4, 5]$  for MT (hereafter case (a)) and  $n = [3, 4, 5]$  for VT (hereafter case (b)). An additional reason for using the mentioned cases is because they lead to different mass-radius diagrams (depending on the index values) and thus, cover a large range of cases that may correspond to boson stars.

We consider that pluralism in the use of EoS will greatly help to test the plausibility of the stability criterion through the CE in the case of boson stars.

## V. RESULTS AND DISCUSSION

As a first step, we employed two analytical solutions, namely the Schwarzschild's and Tolman's VII solutions, in order to calculate the CE. It is worth pointing out that although it is more natural to use a realistic EoS for the fluid interior in order to solve the Einstein's field equations, the use of analytical solutions has the advantage that by having an explicit form, the examination of the implied physics becomes simpler. We should remark that the analytical solutions are a source of infinite number of EoSs (plausible or not). Consequently, they can be used extensively, to introduce and establish some universal approximations (for more details and discussion see also Ref. [51]). In both cases, we formulate the dependence in the form

$$\frac{S_C \rho_c^{-1}}{4\pi b^{-1}} = \mathcal{C} \times 10^5 \times \left(\frac{\text{km}}{R}\right)^3 \frac{1}{\rho_c/ba^{-3}}, \quad (16)$$

where

$$a = \frac{1}{\pi} \left(\frac{\hbar}{mc}\right)^{3/2} \frac{c}{(mG)^{1/2}}, \quad b = \frac{c^2}{G} a,$$

and  $\mathcal{C}$  taking the values 1.728 and 0.145 for the Schwarzschild and Tolman-VII solutions, respectively.

Although the above expression does not ensure the location of the stability point, it is very useful for two reasons: (a) comparison with the results produced by using realistic EoSs, for a fixed value of the radius  $R$  (see Fig. 1(c)), and (b) check and ensure the accuracy of our numerical calculations.

In Fig. 1 we display in order the four cases corresponding to: (first) Fermi gas, (second) Boson gas - C1, (third) Boson gas - C2, and (fourth) Boson gas - C3. In particular, Fig. 1(a) manifests the dependence of the gravitational mass on the radius, Fig. 1(b) presents the dependence of the gravitational mass on the central density, and Fig. 1(c) indicates the CE as a function of the central density for various values of the interaction parameters. In addition, diamonds demonstrate the stability points due to the TM, while open circles mark the minimum of the CE.

In the case of the FG, the first panel of Fig. 1 displays that the points due to TM and CE are located in close proximity, validating the CE method for the location of the stability point. However, a detailed presentation of the percentage error on some interesting quantities, namely the gravitational mass, the radius, the central energy density and the compactness, as shown in Table I, signal a different behavior. While the error in the gravitational mass is lower than 5%, the error in the central energy density can reach up to almost 99%, depending each time to the value of the interaction. The latter have its origin in the creation of a plateau immediately after the rapid decrease of the CE. The existence of a plateau maintains the CE in a narrow region, while the central energy density is spanning in a wide region. In addition, as the data indicate, there is no simple relation between the interaction and the corresponding error to establish a pattern. Thus, in the FG case, while the CE can potentially establish the maximum gravitational mass with good accuracy, the proper description of the central energy density is almost impossible. It needs to be noted that for some specific values of the interaction, the location of the total minimum in CE was not successful, even at high values of densities beyond the maximum mass configuration (unstable region). In that cases, we located a local minimum near the density that corresponds to the maximum mass configuration. The aforementioned statement holds for all cases under consideration in the present study.

In the case of the BG, a similar behavior with the FG is observed. Once more, Fig. 1 in the second, third and fourth panel, displays a visually small difference between the TM and CE points. Nevertheless, Table I illustrates that the error in the underlying quantities aligns with the trend observed in the FG. Specifically, the error in gravitational mass is less than 5%, while the error in central energy density can extend up to 100%. In this instance as well, the CE can be employed to determine the maximum mass, but not the associated central energy density. The observed behavior in both the FG and BG




FIG. 1. (a) Gravitational mass as a function of the radius. (b) The corresponding dependence of the gravitational mass as a function of the central density. (c) The corresponding CE as a function of the central density and two analytical solutions of TOV equations (a and b are constants [23]). The inset figure indicates the location of the minimization of the CE. The black diamonds indicate the stability points due to the TM while the open circles correspond to the minimum of the CE. The panels in order correspond to: EoSs of Sec. IV A, EoSs of Sec. IV B: case 1, EoSs of Sec. IV B: case 2, EoSs of Sec. IV B: case 3.

TABLE I. Percentage (%) error in gravitational mass, radius, central energy density and compactness of the CE method with respect to the TM method.

	(y, w, z, n)	M	R	$\rho_c$	$\beta$	
FG (y)	0.000	0.030	1.648	7.034	1.645	
	0.001	0.022	1.442	6.014	1.440	
	0.005	0.176	3.580	15.320	3.530	
	0.010	0.655	6.158	29.650	5.863	
	0.050	1.177	7.276	38.295	6.577	
	0.100	0.984	6.743	35.102	6.176	
	0.300	4.433	14.491	98.601	11.762	
	0.500	0.093	1.690	7.006	1.754	
BG - C1 (w)	0.001	1.776	9.012	58.881	7.953	
	0.005	1.789	9.045	59.167	7.978	
	0.010	1.789	9.120	59.813	8.067	
	0.050	1.753	8.591	55.327	7.480	
	0.100	1.916	9.001	58.827	7.785	
	0.300	2.252	10.222	69.804	8.877	
	0.500	1.750	8.585	55.422	7.477	
	BG - C2 (z)	0.001	3.396	12.756	81.381	10.728
0.005		3.436	12.709	80.985	10.622	
0.010		3.369	12.891	82.586	10.931	
0.050		3.237	12.633	80.190	10.755	
0.100		3.238	12.627	80.190	10.746	
0.300		4.161	14.080	94.215	11.544	
0.500		3.904	13.557	89.078	11.167	
BG - C3 (n)		case (a)	4	0.418	3.195	14.332
	5		4.264	11.592	98.208	8.289
	case (b)	3	4.318	7.901	100.000	3.890
		4	4.822	7.957	97.959	3.406
		5	4.664	7.899	98.438	3.513

cases strengthens the argument that the location of the stability point is an intrinsic property of the EoS.

As one might easily suspect the CE is related to the star's compactness, Fig. 2 displays the dependence of the CE on the compactness for (a) FG, (b) BG - C1, (c) BG - C2, and (d) BG - C3. In the compactness plane the CE creates also a plateau, similar to the central energy density plane in Fig. 1(c), but in the majority of the cases it is not so extended, depending each time on the corresponding mass-radius diagram of Fig. 1(a). In the FG case, the error can reach values up to 12%, where in the majority of the cases the error is lower than 6%. This result is in accordance with the error in the gravitational mass along with the corresponding error values in the radius. As the radius is more sensitive to the structure of the star, this sensitivity is also presented in the error, reaching values close to 15%. Concerning the BG case, the error in the compactness is established in general under 12% and in the corresponding radius under 14%.

For a visual presentation of Table I for the predictions of the two methods, Fig. 3 displays the percentage error on the predictions for various fermionic and bosonic stars properties (gravitational mass, radius, central density and compactness) as a function of the relative parameterization of the interaction strength  $(y, w, z, n)$  in each particular case. As a general comment, the convergence of the two methods is clearly better in the case of fermion stars compared to that of boson stars (at

least for the selected range of parameterizations). Furthermore, for the case of a boson star, the choice of the EoS is decisive for the accuracy of the convergence of the two methods.

## VI. CONCLUDING REMARKS

Compact objects composed of either fermionic or bosonic matter have been employed for studying the configurational entropy as a mean of stability. The aforementioned quantity should be in alignment with the stability criterion of TM method. It is important to note that the existence of a stable configuration is a property of gravity and independent of the EoS. However, the specific location of the stability point is influenced by the underlying EoS [57]. Considering the above points, one might expect, as suggested in Ref. [23], that the minimization of the CE is a consequence of gravity within the framework of general relativity. If this is the case, the relevant minimum (which should be a total minimum) should be independent of the applied EoS. However, our findings indicate that this is not universally true for compact objects.

The CE was studied in light of the gravitational mass, radius, central energy density and compactness. The TM and the CE method for the location of the stability point converge, with a good or moderate accuracy, for the three out of four quantities under consideration. In fact, the most accurate prediction lies with the maximum mass, where the difference reaches values lower than 5%. In addition, a quantity with good accuracy is also the corresponding radius with errors up to 15%, while in the majority of the cases, the error is lower than 10%. As a result, the combination of the aforementioned macroscopic quantities, which is the compactness, is also a quantity with accuracy lying on values lower than 12%. These three quantities are decisive indicators of the macroscopic quantities of compact stars and leading to the ultimate result that in a macroscopic scale, the two methods for locating the stability point, are in agreement.

The last quantity under consideration, which is the central energy density, leads to enormous amounts of error that can reach values up to 100%. From this point of view, the two methods contradict each other rendering the central energy density unreliably calculated through the CE method.

The above result does not agree with what was recently found that the stability by the minimization of the CE, concerning neutron stars and quark stars, does not have, at least quantitatively, universal validity. A possible explanation, at least for neutron stars, is that the existence of the crust, which has a special constitutive explanation, has a dramatic effect on locating the stability point by the CE minimization method. On the other hand, in the case of quark stars, where there is no crust, the failure of the method does not currently have a solid explanation. Thus, the accurate prediction of the stability point is not only related to the uniformity of the EoS, such as in the case of interacting Fermi and Boson gas, where no crust is added, but also to its specific form.

In conclusion, the CE method can be used as a qualitative, rather than a quantitative, tool to macroscopically locate the instability region of certain configurations of compact objects.




FIG. 2. Configurational entropy as a function of the compactness parameter. The black diamonds indicate the stability points due to the TM while the open circles correspond to the minimum of the CE. The panels in order correspond to: (a) EoSs of Sec. IV A, (b) EoSs of Sec. IV B: case 1, (c) EoSs of Sec. IV B: case 2, (d) EoSs of Sec. IV B: case 3.




FIG. 3. Percentage error in gravitational mass (diamonds), radius (squares), central energy density (plus signs) and compactness (triangles) as a function of the interaction strength parameter of the CE method with respect to the TM method. The panels in order correspond to: (a) EoSs of Sec. IV A, (b) EoSs of Sec. IV B: case 1, (c) EoSs of Sec. IV B: case 2, (d) EoSs of Sec. IV B: case 3.

Finally, although the CE method is an alternative approach for exploring the instability regions of compact objects, the dependence on the specific EoS and the internal structure of the compact star are factors with a decisive role in the validity of the approach.

## ACKNOWLEDGMENTS

The authors would like to thank Dr. Nan Jiang for correspondence and useful comments. All numerical calculations were performed on a workstation equipped with 2 Intel Xeon Gold 6140 Processors (72 cpu cores in total) provided by the MSc program “Computational Physics” of the Physics Department, Aristotle University of Thessaloniki. This work was supported by the Croatian Science Foundation under the project number HRZZ- MOBDOL-12-2023-6026, by the Croatian Science Foundation under the project number IP-2022-10-7773 and by the Czech Science Foundation (GACR Contract No. 21-24281S).

- 
- [1] J. Sañudo and A. Pacheco, Complexity and white-dwarf structure, *Physics Letters A* **373**, 807 (2009).
- [2] K. Chatzisavvas, V. Psonis, C. Panos, and C. Moustakidis, Complexity and neutron star structure, *Physics Letters A* **373**, 3901 (2009).
- [3] M. de Avellar and J. Horvath, Entropy, complexity and disequilibrium in compact stars, *Physics Letters A* **376**, 1085 (2012).
- [4] M. de Avellar, R. de Souza, J. Horvath, and D. Paret, Information theoretical methods as discerning quantifiers of the equations of state of neutron stars, *Physics Letters A* **378**, 3481 (2014).
- [5] H. Adhitya and A. Sulaksono, *Journal of Physics: Conference Series* **572**, 012012 (2020).
- [6] E. Contreras and E. Fuenmayor, Gravitational cracking and complexity in the framework of gravitational decoupling, *Phys. Rev. D* **103**, 124065 (2021).
- [7] C. Posada, J. Hladík, and Z. c. v. Stuchlík, Dynamical stability of the modified tolmán vii solution, *Phys. Rev. D* **103**, 104067 (2021).
- [8] L. Herrera, New definition of complexity for self-gravitating fluid distributions: The spherically symmetric, static case, *Phys. Rev. D* **97**, 044010 (2018).
- [9] L. Herrera, A. Di Prisco, and J. Ospino, Definition of complexity for dynamical spherically symmetric dissipative self-gravitating fluid distributions, *Phys. Rev. D* **98**, 104059 (2018).
- [10] L. Herrera, A. Di Prisco, and J. Ospino, Complexity factors for axially symmetric static sources, *Phys. Rev. D* **99**, 044049 (2019).
- [11] L. Herrera, A. Di Prisco, and J. Carot, Complexity of the bondi metric, *Phys. Rev. D* **99**, 124028 (2019).
- [12] Sharif, M. and Butt, Iqra Ijaz, Complexity factor for charged spherical system, *Eur. Phys. J. C* **78**, 688 (2018).
- [13] Sharif, M. and Butt, Iqra Ijaz, Complexity factor for static cylindrical system, *Eur. Phys. J. C* **78**, 850 (2018).
- [14] M. Sharif, A. Majid, and M. M. M. Nasir, Complexity factor for self-gravitating system in modified gauss–bonnet gravity, *International Journal of Modern Physics A* **34**, 1950210 (2019), <https://doi.org/10.1142/S0217751X19502105>.
- [15] M. Sharif and K. Hassan, Complexity of dynamical cylindrical system in  $f(g, t)$  gravity, *Modern Physics Letters A* **37**, 2250027 (2022), <https://doi.org/10.1142/S0217732322500274>.
- [16] Yousaf, Z., Bhatti, M. Z., and Naseer, T., Study of static charged spherical structure in  $f(t, r, q)$  gravity, *Eur. Phys. J. Plus* **135**, 323 (2020).
- [17] Z. Yousaf, M. Y. Khlopov, M. Z. Bhatti, and T. Naseer, Influence of modification of gravity on the complexity factor of static spherical structures, *Monthly Notices of the Royal Astronomical Society* **495**, 4334 (2020), <https://academic.oup.com/mnras/article-pdf/495/4/4334/33372154/staa1470.pdf>.
- [18] Z. Yousaf, M. Z. Bhatti, and T. Naseer, Measure of complexity for dynamical self-gravitating structures, *International Journal of Modern Physics D* **29**, 2050061 (2020), <https://doi.org/10.1142/S0218271820500613>.
- [19] Z. Yousaf, K. Bamba, M. Bhatti, and K. Hassan, Measure of complexity in self-gravitating systems using structure scalars, *New Astronomy* **84**, 101541 (2021).
- [20] Z. Yousaf, M. Bhatti, and M. Nasir, On the study of complexity for charged self-gravitating systems, *Chinese Journal of Physics* **77**, 2078 (2022).
- [21] M. Gleiser and N. Stamatopoulos, Entropic measure for localized energy configurations: Kinks, bounces, and bubbles, *Physics Letters B* **713**, 304 (2012).
- [22] M. Gleiser and D. Sowinski, Information-entropic stability bound for compact objects: Application to q-balls and the chandrasekhar limit of polytropes, *Physics Letters B* **727**, 272 (2013).
- [23] M. Gleiser and N. Jiang, Stability bounds on compact astrophysical objects from information-entropic measure, *Phys. Rev. D* **92**, 044046 (2015).
- [24] P. S. Koliogiannis, G. A. Tsalis, C. P. Panos, and C. C. Moustakidis, Configurational entropy as a probe of the stability condition of compact objects, *Phys. Rev. D* **107**, 044069 (2023).
- [25] M. Gleiser and D. Sowinski, Information-entropic signature of the critical point, *Physics Letters B* **747**, 125 (2015).
- [26] N. R. Braga, Information versus stability in an anti-de sitter black hole, *Physics Letters B* **797**, 134919 (2019).
- [27] N. R. F. Braga and R. da Mata, Configuration entropy for quarkonium in a finite density plasma, *Phys. Rev. D* **101**, 105016 (2020).
- [28] S. H. Alexander, K. Yagi, and N. Yunes, An entropy-area law for neutron stars near the black hole threshold, *Classical and Quantum Gravity* **36**, 015010 (2018).
- [29] R. da Rocha, Ads graviton stars and differential configurational entropy, *Physics Letters B* **823**, 136729 (2021).
- [30] G. Karapetyan, The nuclear configurational entropy approach to dynamical qcd effects, *Physics Letters B* **786**, 418 (2018).
- [31] R. A. C. Correa, P. H. R. S. Moraes, A. de Souza Dutra, W. de Paula, and T. Frederico, Configurational entropy as a constraint for gauss-bonnet braneworld models, *Phys. Rev. D* **94**, 083509 (2016).



- [32] W. Barreto, A. Herrera–Aguilar, and R. da Rocha, Configurational entropy of generalized sine–gordon-type models, *Annals of Physics* **447**, 169142 (2022).
- [33] S. L. Shapiro and S. A. Teukolsky, *Black Holes, White Dwarfs, and Neutron Stars* (John Wiley & Sons, New York, 1983).
- [34] N. Glendenning, *Compact Stars: Nuclear Physics, Particle Physics, and General Relativity* (Springer, Berlin, 2000).
- [35] P. Haensel, A. Potekhin, and D. Yakovlev, *Neutron Stars I: Equation of State and Structure* (Springer-Verlag, New York, 2007).
- [36] Y. Zeldovich and I. Novikov, *Stars and Relativity* (Dover Publications, INC, Mineapolis New York, 1971).
- [37] S. Weinberg, *Gravitational and Cosmology: Principles and Applications of the General Theory of Relativity* (Wiley, New York, 1972).
- [38] B. Schutz, *A First Course in General Relativity* (Cambridge University Press, Cambridge, 1985).
- [39] J. Schaffner-Bielich, *Compact Star Physics* (Cambridge University Press, Cambridge, 2020).
- [40] S. Chandrasekhar, The dynamical instability of gaseous masses approaching the schwarzschild limit in general relativity., *The Astrophysical Journal* **140**, 417 (1964).
- [41] S. Chandrasekhar, Dynamical instability of gaseous masses approaching the schwarzschild limit in general relativity, *Physical Review Letters* **12**, 114 (1964).
- [42] A. R. Liddle and M. S. Madsen, The structure and formation of boson stars, *International Journal of Modern Physics D* **01**, 101 (1992), <https://doi.org/10.1142/S0218271892000057>.
- [43] S. L. Liebling and C. Palenzuela, Dynamical boson stars, *Living Reviews in Relativity* **20**, 5 (2017).
- [44] F. E. Schunck and E. W. Mielke, General relativistic boson stars, *Classical and Quantum Gravity* **20**, R301 (2003).
- [45] D. Kramer, H. Stephani, M. A. MacCallum, and E. Hert, *Exact Solutions of Einstein’s Field Equations* (Deutsche Verlag der Wissenschaften, Berlin/Cambridge University Press, Cambridge, 1980).
- [46] M. Delgaty and K. Lake, Physical acceptability of isolated, static, spherically symmetric, perfect fluid solutions of einstein’s equations, *Computer Physics Communications* **115**, 395 (1998), computer Algebra in Physics Research.
- [47] J. R. Oppenheimer and G. M. Volkoff, On massive neutron cores, *Physical Review* **55**, 374 (1939).
- [48] A. M. Raghoonundun and D. W. Hobill, Possible physical realizations of the tolman vii solution, *Physical Review D* **92**, 124005 (2015).
- [49] P. S. Negi and M. C. Durgapal, Stable ultracompact objects, *General Relativity and Gravitation* **31**, 13 (1999).
- [50] P. Negi and M. Durgapal, Relativistic supermassive stars, *Astrophysics and Space Science* **275**, 185 (2001).
- [51] C. Moustakidis, The stability of relativistic stars and the role of the adiabatic index, *General Relativity and Gravitation* **49**, 68 (2017).
- [52] G. Narain, J. Schaffner-Bielich, and I. N. Mishustin, Compact stars made of fermionic dark matter, *Phys. Rev. D* **74**, 063003 (2006).
- [53] M. Colpi, S. L. Shapiro, and I. Wasserman, Boson stars: Gravitational equilibria of self-interacting scalar fields, *Phys. Rev. Lett.* **57**, 2485 (1986).
- [54] P. Agnihotri, J. Schaffner-Bielich, and I. N. Mishustin, Boson stars with repulsive self-interactions, *Phys. Rev. D* **79**, 084033 (2009).
- [55] N. Rutherford, G. Raaijmakers, C. Prescod-Weinstein, and A. Watts, Constraining bosonic asymmetric dark matter with neutron star mass-radius measurements, *Phys. Rev. D* **107**, 103051 (2023).
- [56] S. L. Pitz and J. Schaffner-Bielich, Generating ultracompact boson stars with modified scalar potentials, *Phys. Rev. D* **108**, 103043 (2023).
- [57] N. K. Glendenning and C. Kettner, Possible third family of compact stars more dense than neutron stars, *Astronomy & Astrophysics* **353**, L9 (2000), [arXiv:astro-ph/9807155](https://arxiv.org/abs/astro-ph/9807155) [astro-ph].